Sex-dependent effects of infection on guppy reproductive fitness and offspring parasite

resistance

Isabella L. G. Weiler*, Rachael D. Kramp*, Faith Rovenolt, Jessica F. Stephenson[§]

Department of Biological Sciences, University of Pittsburgh, Pittsburgh, PA, USA

*these authors contributed equally to this work.

[§]Author for correspondence: jfrstephenson@gmail.com

Acknowledgements: We thank D. Clark, L. Colgan, D. Henry, M. Janecka, R. Patrick, J.

Walsman, C. Wynne for technical assistance. This work was approved by the University of

Pittsburgh's Institutional Animal Care and Use Committee (IACUC), protocol 21069471, and

supported by the University of Pittsburgh, NSF DGE: 1747452 (to FR), and an HHMI Gilliam

Fellowship (to RDK).

Data availability: The data are available in the supplement and, upon manuscript acceptance,

will be deposited in Dryad and the doi included here. The code and output we used to analyse

the data is included in the supplement.

Conflict of interest: We have no conflicting interests.

CRediT authorship statement: Conceptualization = JFS; Data curation = ILGW, RDK, FR;

Formal analysis = JFS; Funding acquisition = FR, JFS; Methodology = ILGW, RDK, FR; Project

administration = JFS; Resources = JFS; Visualization = JFS; Writing – original draft = ILGW,

JFS; Writing – review & editing = ILGW, RDK, FR, JFS. All authors approved gave approval for

publication and agreed to be accountable.

Abstract:

1. Infection imposes energetic costs on hosts. Hosts typically respond by shifting resources, potentially affecting the quantity and quality of offspring they produce. As the sexes differ in their optimal reproductive strategies, infection of mothers versus fathers may affect offspring quantity and quality in different ways.

2. Here, we test how experimental infection of guppies *Poecilia reticulata* with the ectoparasite *Gyrodactylus turnbulli* affects parental reproductive fitness and offspring parasite resistance. We compared breeding pairs in which one or neither parent had previously been infected.

3. In terms of reproductive fitness, parental infection experience did not affect the size, body condition, or number of offspring produced, but fathers who experienced the heaviest infections produced offspring ~55 days sooner than average. This result may represent terminal investment by the males most affected by infection, or may indicate that these males have a faster pace of life, investing in reproduction at the expense of parasite defence.

4. We found that offspring age, parental infection experience, and parental infection severity together strongly predicted offspring parasite resistance. Only among pairs in which one parent had been infected, older offspring, which were those born soonest after the parent's infection, tended to experience heavier infections. This result may therefore reflect temporary infection-induced reductions in parental investment in offspring quality. Beyond this effect of offspring age, offspring of infected mothers experienced 105 fewer worm days than those of infected fathers: fathers, but not mothers, that experienced heavy infections themselves produced offspring that also experienced heavy infections. The parent-offspring regression for infected fathers is therefore consistent with previous evidence that parasite resistance is heritable in this

system, and yields a narrow sense heritability estimate of 0.69±0.13. By contrast, the

mother-offspring regression (slope: -0.13±0.17) provides novel insight that mothers may

engage in transgenerational immune priming.

5. Overall, our results suggest that the sexes strike a different balance between offspring

quantity and quality when faced with infection, with potentially broad implications for

disease and host-parasite coevolutionary dynamics in nature.

Keywords: *Gyrodactylus*, heritable parasite resistance, life history, parental investment theory,

parent-offspring regression, transgenerational immune priming, sex-specific response to

infection

Introduction: Infected parents may change the quantity and quality of the offspring they

produce, both because of the energetic constraints (Tizard, 2008) or pathology (Armour et al.,

2020; Billi et al., 2019; Khan et al., 2017) associated with infection, and because the optimal

investment in defence (Boots & Bowers, 2004; Graham, 2013; Houston et al., 2007; Tschirren &

Richner, 2006; Viney et al., 2005) and reproductive strategy may differ under the threat of

infection (Schmid-Hempel, 2003; Sheldon & Verhulst, 1996). Intuitively, parental infection could

induce the production of fewer, lower quality offspring due to decreased parental resources

(Hagmayer et al., 2020; Vantaux et al., 2014). However, infection can also induce the

production of more offspring as infected parents 'terminally invest': hosts increase reproductive

output at the expense of their own defence against infection in an effort to produce offspring

before their own death (Duffield et al., 2017). Infection may also induce the production of higher

quality offspring, particularly if they are likely to be exposed to the same parasite, through

parental investment in the immunological defences of their offspring, known as

"transgenerational immune priming", or TGIP (Roth et al., 2018). TGIP can last from shortly after

birth, (in the case of e.g. maternal antibodies (Breuil et al., 1997; Brown et al., 1997)) to

subsequent generations (in the case of epigenetic changes (Beemelmanns, 2019)), depending

on the host species and the disease to which they are responding (Roth et al., 2018). TGIP is

costly to both parents and offspring, however: manufacturing parentally donated factors such as

proteins or antibodies requires resources and energy, and any form of priming may require

offspring to mount an immune response, even in the absence of the disease (Roth et al., 2018).

Mothers and fathers typically take different strategies to maximise fitness, even in the absence

of infection. At the ultimate level, parental investment theory (Trivers, 1972) posits that as

females (in species with typical sex roles) invest more in individual gametes or zygotes, they are

limited in the rate at which they can produce offspring. Females therefore maximise fitness by

investing in the probability of survival and reproduction of the offspring they produce. Males, by

contrast, produce many more gametes at a much higher rate and, all else being equal,

maximise fitness by mating with as many females as possible. Therefore, in the absence of

infection, males should optimally invest in offspring quantity, and females, quality.

These life history differences between the sexes have implications for their investment in their

own immunity (Fischer et al., 2011; Rolff, 2002; Stoehr & Kokko, 2006), and potentially how they

shift their parental investment post-infection. Females typically invest more in their immune and

self-repair systems to maximise longevity, and thus lifetime reproductive success, whereas

males do not experience this same selection on longevity (Rolff, 2002; Stoehr & Kokko, 2006).

As well as ultimate drivers, the sexes may differ in the proximate mechanisms through which

they can affect offspring quantity or quality. For example, in many systems, mothers have more

opportunity to affect the phenotypes of their offspring (Wassarman et al., 2001; Wolf & Wade,

2009), and as a result offspring phenotypes are often strongly correlated with maternal

phenotypes, for example in live-bearing fish (Hagmayer et al., 2018) and humans (Zhang et al.,

2018). This may hold true for TGIP: although the underlying mechanisms have not been fully

elucidated, promising candidates include parentally donated proteins (Adrian-Kalchhauser et al.,

2020) that are likely too large to be transferred by sperm (Arnqvist & Rowe, 2005; Wassarman

et al., 2001). Indeed, no relationships among antibody titres were detected in vertebrates when

fathers but not mothers were primed (Gasparini et al., 2002; Reid et al., 2006). However, other

mechanisms, such as epigenetic modifications (i.e., altered immune gene expression in

offspring (Vilcinskas, 2021)) or nonnuclear effects (Simms & Triplett, 1996), could be mediated

by either parent. Indeed, a recent meta-analysis of TGIP in invertebrates found that the sex of

the parent primed did not affect the strength of TGIP (Rutkowski et al., 2023). Such

mechanisms likely function among vertebrates, too: among pipefish, both offspring and grand-

offspring of primed parents show altered gene expression (Beemelmanns & Roth, 2017).

However, more broadly, work in vertebrates has historically been heavily biased towards

investigating maternal effects (Roth et al., 2012, 2018). Indeed, evidence for paternally-

mediated TGIP in vertebrates is limited to sex-role reversed species (seahorses and pipefish

biparental TGIP influences offspring lymphocyte proliferation and immune gene expression

(Beemelmanns & Roth, 2016, 2017; Roth et al., 2012)), or those with some paternal care

(Pizzolon et al., 2010).

Here, we test how infection with *Gyrodactylus turnbulli* affects the reproductive fitness of male

and female guppies, and the quality of the offspring produced. This highly tractable and

manipulable system has been used extensively (Harris & Lyles, 1992; Madhavi & Anderson,

1985; Stephenson et al., 2018; Walsman et al., 2022), and has characteristics that make it

particularly appropriate for understanding transgenerational effects. Indeed, guppy exposure to

predators has transgenerational effects mediated by both parents (Stein & Hoke, 2022). For

infection, resistance to *G. turnbulli* is heritable (Dargent et al., 2016; Madhavi & Anderson,

1985), enabling us to partially control for host genotype when investigating other potential

sources of resistance. The parasite is associated with lower survival (Stephenson et al., 2016;

van Oosterhout et al., 2007) and body condition (Stephenson, van Oosterhout, et al., 2015) of

wild guppies, indicating it depletes host resources. Furthermore, guppy immune gene

expression is affected by infection with *Gyrodactylus* spp. (Konczal et al., 2020). The key

conditions for TGIP are present in this system (Roth et al., 2018): *G. turnbulli* can be directly

transmitted from mother to offspring during birth, so offspring of infected mothers are likely to be

exposed to the same infection; and *G. turnbulli* primarily reproduce asexually (Schelkle et al.,

2012), indicating infections in natural populations have some genotypic stability.

The biology of our system also suggests that the sexes may differ in how infection with *G.*

turnbulli affects parental investment. First, the parasite interacts with the host sexes differently:

males are less likely to be found infected in nature (Stephenson, Van Oosterhout, et al., 2015),

and experience lower infection loads both in the wild (Stephenson, Van Oosterhout, et al., 2015;

Walsman et al., 2022) and during experimental infections (Dargent et al., 2016) than females.

Despite this, females may be more likely than males to survive their infections in nature (Clark

et al., 2023; Stephenson, van Oosterhout, et al., 2015; van Oosterhout et al., 2007), and appear

better able to maintain the same behavioural phenotypes as uninfected fish (Jog et al., 2022;

Stephenson, 2019). Female guppies may be well-positioned to prime their offspring: they give

birth to live offspring whose eggs are provisioned only at the time of fertilization, and female

phenotype affects offspring phenotype (Shikano & Taniguchi, 2005). However, in other fish,

maternally-provisioned antibodies often disappear after offspring pass certain developmental

stages such as absorbing the yolk sac (Breuil et al., 1997; Brown et al., 1997) which, in

livebearers such as the guppy, usually happens before they are born (Norazmi-Lokman et al.,

2016). Male guppies may also be able to directly affect offspring phenotype through sperm

quality: in an artificial insemination experiment, sperm from males fed a high quality diet was of

a higher quality itself, and produced offspring that were heavier at birth (Evans et al., 2017).

Sperm is the only potential route for paternally-derived TGIP in this system: infection may affect

a father's investment in his sperm, which in turn could affect the quality of his offspring, as

potentially reflected in their parasite resistance. We also know that infection with *G. turnbulli*

affects DNA methylation in guppies (Hu et al., 2018a), raising the possibility of epigenetically-

mediated (and thus potentially bi-parental) TGIP. Such epigenetic changes can last generations

after an initial immune challenge (Beemelmanns, 2019).

We address two main questions: 1) Does *G. turnbulli* infection affect guppy reproductive

fitness? 2) Does parent infection affect offspring quality and response to *G. turnbulli* infection?

Based on parental investment theory, we predicted that males should prioritize offspring

number, whereas females should prioritise offspring quality, after exposure to a parasite. We

quantified offspring number as the total number of offspring a pair produced during our

observation period, and latency to birth: together these predict how many offspring a female

(and her mate in monogamous pairs, such as in our experiment) may produce in her lifetime,

assuming all else is constant. We quantified offspring quality as their size, body condition, and

how heavily infected they became during experimental infection.

Materials and Methods:

(a) Fish Origin and Maintenance

We used laboratory-bred descendants of guppies from the Caura River in Trinidad. This

laboratory population was founded by fish collected in 2012 from a high-predation population

located at an undeveloped, rural site on the Caura River, Trinidad (UTM: 20 P 679527.7 m E,

1180376.4 m N based on WGS84 Datum; elevation 112m). Since 2012, the population has

been maintained in large numbers, and we continue to see substantial variation in disease

resistance, as is clear from our results. Fish were maintained and bred in 20 L tanks of mixed

sexes, under standard conditions, as follows. Tanks were on a recirculating system

(Aquaneering) with approximately 20% daily water changes, a lighting schedule of 12L:12D, and

an average temperature of 25±1°C. In these systems, wastewater from tanks passed through

foam, sand, and ultraviolet filters before re-entering other tanks, preventing unintentional

parasite transmission. Fish were fed flake and Artemia daily.

(b) Parasite origin and maintenance

Our parasite line was established by transferring one worm from a commercially obtained guppy

to an uninfected host from a mixed laboratory-bred stock, a 'culture fish'. This parasite line was

maintained on groups of three to six culture fish per 1.8 L tank under standard conditions. Twice

per week, all culture fish were anesthetized (tricaine methanesulfonate 'MS222'; 4g per L) and

parasites were counted using a dissecting microscope. Uninfected fish were added to replace

recovered or dead fish.

(c) Parental infection and breeding design

Parental fish were anaesthetized (using MS222 as above) and placed in close proximity to a

dead, highly infected culture fish until at least two worms transmitted (mean±SEM=2.8±0.27), as

observed under a dissecting microscope. Once infected, parental fish were isolated in 1.8 L

tanks maintained under standard conditions for the duration of their infection. We counted

worms every three days for 15 days to record individual infection trajectory over time. After 15

days, we treated fish with levamisole (0.002g per L) to clear infection. We verified that this

treatment had cleared infection by screening the fish for parasites, under anaesthetic, three

times with a minimum of four days between screens. We recorded the length and weight of the

parents before and after infection. Parasite naïve parents were exposed to the same conditions

and manipulations, with the exception of parasites.

We set up 38 total breeding pairs of one male and one female using the recovered fish and

parasite naïve fish in three different treatments: recovered mother, naïve father (13 pairs);

recovered father, naïve mother (11 pairs); and both parents naïve (14 pairs). We did not set up

pairs in which both parents were recovered because previous experience suggested this

treatment may not have yielded sufficient offspring, and our primary objective was to examine

the independent effects of maternal and paternal infection. These breeding pairs were housed in

1.8 L tanks under standard conditions, including fry mesh structures and gravel to reduce

possible fry cannibalism by the parents. Between May and December 2021, these breeding

tanks were checked daily for the presence of offspring, which were moved to a separate 1.8 L

tank. Each brood produced from a pair was moved to a separate 1.8 L tank.

(d) Offspring infection

Once they reached adulthood, offspring were separated into individual 1.8 L tanks. A total of

108 offspring were collected: 32 from recovered fathers and naïve mothers; 38 from recovered

mothers and naïve fathers; and 38 from naïve parents. We used two batches of fish, one in

August 2021 (52 offspring from 33 breeding pairs, with an average age at infection of

approximately 4.4 months) and one in May 2022 (56 offspring from 32 breeding pairs, with an

average age at infection of approximately 7.8 months). According to the needs of other

concurrent experiments, 94 fish were infected manually, as described under parental infection,

and 14 were infected via "co-housing": they were housed in 1.8 L tanks with an infected

conspecific and screened for infection every 24 hours. Once infected, these fish were isolated in

a 1.8 L tank and treated the same as the manually infected offspring and parents from that point

forwards. As noted below, we included infection type as a fixed effect in the statistical models

and found that it did not explain significant variation in any of our response variables. On all

infected offspring, we counted the worms every three days for 12 days and then sacrificed the

host.

(e) Data Analysis

We used R statistical software (R Core Team, 2020) to analyse the data, and provide the code

and output in the supplement. We used the data from this experiment to address two main

questions: 1) Does infection affect reproductive fitness? 2) Does parent infection affect offspring

response to infection? We addressed each question using a) data from all pairs, whether or not

one parent had been infected; and b) only data from pairs in which one parent had been

infected. This enabled us to evaluate the extent to which parental exposure to infection affected

reproductive fitness and offspring infection (a); and the extent to which the intensity of the

parent's infection explained variation in these two outcomes (b). As our measure of the parent's

infection intensity, and offspring response to infection, we used the area under the curve of

infection load over time ('infection integral').

As our metrics of reproductive fitness, we used: the latency between the parent's infection and

the birth of the offspring; the total number of offspring a pair produced during the observation

period; and the body length and body condition of those offspring at adulthood. We used each

of these four metrics as the response variable in separate linear (mixed) models (Gaussian error

family, identity link function) in glmmTMB (Brooks et al., 2017) using first the data from all pairs,

and then data from only pairs in which one parent had been infected (i.e., a total of eight

models). Each of these eight models included as fixed effects the parental infection treatment

(mother, father, or neither parent infected), mother's body condition, mother's body length, and

father's body condition. For response variables relevant to the level of the individual offspring

(birth latency, body length, body condition), the models included mother identity as a random

factor as many pairs produced multiple offspring. Because male and female guppies typically

differ in body condition metrics and length, we included sex of the offspring as a fixed effect in

those models. For the total number of offspring models, we used only one observation per pair.

The four models using only data from pairs in which one parent had been infected additionally

included the (log transformed) infected parent's infection integral and the interaction with

parental infection treatment as fixed effects.

We used two separate linear mixed models to test how a) parental exposure to infection, and b)

the severity of that infection, affects offspring response to infection (Gaussian error family,

identity link function in glmmTMB (Brooks et al., 2017)). Both models included as fixed effects

the sex, body condition, body length (residuals on sex), age at infection, the method of

experimental infection (manual vs. cohoused), and the number of parasites establishing an

infection on the offspring. Both models also included the parental infection treatment (mother,

father, neither), and the interaction between parental infection treatment and offspring age at

infection. As random effects, both models included the mother's identity to control for the non-

independence of data from multiple offspring from the same pair. In addition, the model

including parental infection severity included parental infection integral (log transformed), and its

interaction with parental infection treatment, plus a three-way interaction with these two

variables and offspring sex. We checked the fit of all models by visually assessing the plots and

tests produced by the DHARMa package ((Hartig, 2022) provided in the supplement). DHARMa

simulates standardized residuals to thoroughly validate complex mixed models such as those

we use here. The figures were plotted using visreg (Breheny & Burchett, 2017) and ggplot2

(Wickham, 2016).

We estimated narrow-sense heritability from the slope of the parent-offspring regression

(Åkesson et al., 2007, 2008; Falconer & Mckay, 1997). We rescaled parent and offspring

infection integrals and included offspring age at infection (which in our main analysis explained

a highly significant portion of the variation in offspring infection integral) as a fixed effect in our

models, and mother ID as a random effect to account for the fact that pairs contributed more

than one offspring. We used glmmTMB (Brooks et al., 2017) to estimate slope, standard error,

and probability the slope differed from 0.

Results: We did not find concerning issues with the fits of our models and present the plots and

tests of assumptions in the supplement.

(a) Does infection affect reproductive fitness?

We found that previously infected and parasite naïve guppies produced the same number of

offspring, at the same rate, and these did not differ in either size or body condition at adulthood

(all $P > 0.25$, see supplement).

Similarly, we found that the intensity of parental infection did not significantly affect the number

of offspring a pair produced, or the size or body condition of those offspring at adulthood ($P =$

0.06-0.97, see supplement). One of these results was marginally non-significant: parents that

experienced heavier infection loads produced offspring with higher body condition at adulthood

(Chisq = 3.51, df = 1, $P = 0.061$).

We found that the number of days between parental infection and offspring birth depended on

the intensity of that infection, and this relationship differed between infected mothers and fathers

(Fig. 1; interaction between parental infection treatment and integral: Chisq = 5.13, df = 1, $P =$

0.024). We reran this analysis on the data split by sex and found that the correlation among

fathers differed significantly from 0 (effect of infection integral: Chisq = 5.49, df = 1, $P = 0.019$),

whereas that among mothers did not (Chisq = 0.87, df = 1, $P = 0.35$).

(b) Does parent infection affect offspring response to infection?

We found that the effect of parental infection treatment on offspring response to infection

depended on the age of the offspring at infection, and the parental infection treatment: among

offspring with a previously infected parent, younger offspring experienced lower infection

intensities than older offspring (Fig. 2; interaction between parental infection treatment and

offspring age at infection: Chisq = 7.39, df = 2, P = 0.025). We also found that the more

parasites initially established on the fish, the higher their infection integral (Chisq = 9.69, df = 1,

P = 0.002). No other variables, including offspring sex, explained any of the variation in offspring

infection integral (see supplement for full model output). Our *post-hoc* tests of the correlation in

Fig. 2 show that it is significantly different from 0 for offspring of infected fathers (Chisq = 16.57,

df = 1, P < 0.000) and mothers (Chisq = 4.70, df = 1, P = 0.03), but not those of parasite naïve

parents (Chisq = 2.39, df = 1, P = 0.122; see supplement).

Among offspring with one infected parent, we found their infection integral was significantly

predicted by whether the mother or the father had been infected, but this effect depended both

on offspring age at infection, and the intensity of the parent's infection. As in the model including

all offspring, younger offspring experienced lower infection intensities than older offspring (Chisq

= 19.74, df = 1, P < 0.000). We also found a significant correlation between the father's infection

integral and that of the offspring, but no significant correlation between mother and offspring

infection integral (Fig 4.; interaction between parental infection integral and parental infection

treatment: Chisq = 11.89, df = 1, P = 0.0006). No other variable explained a significant portion

of the variation (see supplement), including the interaction between parental infection treatment

and offspring age: the correlations in Fig. 2 do not therefore differ significantly between offspring

with infected mothers and fathers. Our *post-hoc* tests confirmed that the correlation between

father and offspring infection integrals differs from 0 (Chisq = 13.07, df = 1, P = 0.0003),

whereas that between mothers and offspring does not (Chisq = 0.63, df = 1, P = 0.43).

(c) What is the heritability of parasite resistance?

The slope of the father-offspring regression is 0.69±0.13 (estimate±standard error; $P <$

0.00001), and 0.53±0.18 ($P = 0.003$) after the removal of one outlier. The slope of the mother-

offspring regression is -0.13±0.17 ($P = 0.45$).

Discussion: Our results show that parental infection does affect the quantity and quality of

offspring a pair produces, and that these effects depend on which parent was infected. We

found that males that experienced the heaviest infections produced offspring ~55 days quicker

than average, which may result in these males producing more offspring over their lifetimes. We

also found that older offspring experienced heavier infections, but only among those produced

by parents with experience of infection. Finally, we found that offspring of fathers who

experienced heavy infections also experienced heavy infections, whereas offspring of mothers

showed no such pattern. On average, offspring of infected mothers experienced 105 fewer

worm days than offspring of infected fathers. Overall, our results indicate how parasites may

change reproductive investment strategies, and how this affects the next generation of

hosts. Here we discuss each of our results in turn.

Our observation that fathers that become heavily infected produced offspring more quickly (Fig.

1) is consistent with other observations from this system. First, these males may mate more

quickly than those that experienced lighter infections and thus their broods are produced earlier.

Previous work has shown that *Gyrodactylus* spp. infection (Kennedy et al., 1987; López, 1998,

1999) and immune challenges (Encel et al., 2023) affect guppy reproductive behaviour and,

consistent with this explanation, the heaviest infected males appear more interested in contact

with females (Stephenson, 2019). This behavioural explanation may be a component of a

broader explanation that fathers who become heavily infected may generally have a faster pace

of life. Such males would bias investment towards reproduction, including reproductive

behaviours, and away from immune investment, resulting in their heavier infection loads. There

is evidence from natural guppy populations that pace of life does predict resistance to

Gyrodactylus spp. (Stephenson, van Oosterhout, et al., 2015). Alternatively, rather than being

indicative of a pace of life syndrome, this result could indicate that the males that experienced

the heaviest infections are terminally investing in reproduction (Duffield et al., 2017): while

suggestive results have been reported (López, 1998; Stephenson, 2019) this phenomenon

remains to be conclusively demonstrated in guppies.

We found that older offspring experienced heavier infections, but only among those produced by

parents with experience of infection (Fig. 2). It is well known that many immune components

wane with age across animal taxa (Metcalf et al., 2020; Peters et al., 2019), but the offspring in

our experiment were young adults, and the effect of age was absent from those born to parasite

naïve parents. We therefore suggest that this result is evidence that parental infection affects

investment in offspring quality. Indeed, investment in offspring can respond to the environment

among guppy mothers (Cunningham & Russell, 2000; Reynolds & Gross, 1992) and fathers

(Evans et al., 2017). The oldest offspring in our experiment are those produced the soonest

after parental infection, so could therefore represent those that received least resources from

parents depleted by that infection. However, the mechanism underlying this potential

explanation is unclear. As maternal antibodies, mRNAs, and proteins are typically absent from

guppy offspring within a few days post-birth (Breuil et al., 1997; Brown et al., 1997; Norazmi-

Lokman et al., 2016), and such proteins are too large to be transferred by sperm (Arnqvist &

Rowe, 2005; Wassarman et al., 2001), it seems unlikely this result is due to waning TGIP.

We found that offspring infection intensity was strongly correlated with parental infection

intensity among offspring of infected fathers, but not among those of infected mothers, even

after controlling for the effect of offspring age (Fig. 3). Among offspring of infected fathers, the

positive parent-offspring regression we observe is consistent with previous findings that

resistance to *Gyrodactylus* spp. is heritable in guppies (Dargent et al., 2016; Madhavi &

Anderson, 1985; Phillips et al., 2018), and other fish (Gilbey et al., 2006). Our estimates of

narrow-sense heritability from these regressions (0.69 or 0.53 after outlier removal) are high, but

comparable to heritability estimates for male guppy ornaments (Morris et al., 2020; van der Bijl

et al., 2023), which are correlated with *G. turnbulli* resistance in our focal population

(Stephenson et al., 2020).

Despite the heritability of resistance to *Gyrodactylus* spp., we did not observe a significant

mother-offspring regression. Potential mechanisms as to why offspring resembled their fathers

but not their mothers in this trait include sex-linked inheritance, autosomal dominant traits, and

maternally derived TGIP. We believe maternally derived TGIP provides the most parsimonious

explanation. For example, if resistance to *Gyrodactylus* spp. is Y-linked, and thus passed from

fathers to sons, the sex of the offspring from infected fathers should explain variation in offspring

parasite resistance but it does not. However, there is some evidence that parasite resistance

evolves along sex-specific trajectories in natural conditions (Dargent et al., 2016). Autosomal

dominant traits also seem an unlikely explanation as males were randomly assigned across our

38 pairs. Furthermore, the Major Histocompatibility Complex (MHC) is important in guppy

resistance to these parasites (Fraser et al., 2010; Fraser & Neff, 2009; Phillips et al., 2018), and

many MHC genes are codominantly expressed (Herdegen-Radwan et al., 2020). We therefore

suggest that maternally derived TGIP most likely underlies our result, though it is clear more

work is needed to fully elucidate the mechanisms underlying the difference we observed

between father-offspring and mother-offspring regressions.

Our data are therefore consistent with maternally mediated TGIP and provide some insight into

potential mechanisms. Offspring of mothers that experienced severe infections had light

infections, comparable to those of offspring of mothers who experienced less severe infections.

Thus, any parasite load may be sufficient to induce TGIP to the same protective level, or

perhaps TGIP is compensatory to load and thus allows all offspring of infected mothers to

benefit from a similar level of resistance. The offspring were all adults, several months old,

before their infections, which could rule out mechanisms observed in other systems such as

endocrine signalling (Kvist et al., 2020) or protein transfer (Adrian-Kalchhauser et al., 2020)

from the mother to the offspring. Epigenetic mechanisms would persist in the offspring until

adulthood and beyond, and maternal stress can induce epigenetic changes, such as histone

modifications and DNA methylation, during gametogenesis in mammals (Gapp et al., 2014).

Indeed, epigenetic changes in response to infection have been observed in female guppies (Hu

et al., 2018b) and these changes may be conserved from generation to generation (Heard &

Martienssen, 2014). These changes persist after fertilization, could permanently alter offspring

immune function, and provide an evolutionary advantage as females can prepare their offspring

for the parasites they are likely to encounter in their environment (Willis et al., 2021). Epigenetic

changes could also lead to non-immune changes like increased epidermal thickness, which has

been implicated in guppy response to stress and interactions with *G. turnbulli* (Gheorghiu et al.,

2006, 2009, 2012).

Our results suggest more complexity in the interactions between host and parasite in natural

communities. In our system, Trinidadian guppy populations differ in their prevalence of

Gyrodactylus spp. and the mean intensity of infection, and these are likely not stable through

time (Clark et al., 2023; Stephenson, Van Oosterhout, et al., 2015; Walsman et al., 2022). In

populations and at times when prevalence is high, our findings suggest that more mothers may

prime their offspring, who would develop overall lower infection loads, potentially reducing

transmission (Stephenson et al., 2017), and thus affecting population-level prevalence. If

maternally derived TGIP has a measurable impact on disease dynamics in nature, it may

therefore slow the evolution of host resistance by reducing the strength of parasite-mediated

selection. However, host defences do appear to evolve rapidly in response to these parasites

(Dargent et al., 2016; Phillips et al., 2018). Heavily infected guppies may not successfully breed

in nature, which would erode the effects of TGIP on resistance evolution. Indeed, observed

natural infection loads tend to be low (Stephenson, Van Oosterhout, et al., 2015), and patterns

of aggregation show that heavily infected hosts are missing from our surveys, presumed dead

(Clark et al., 2023). Similarly, the impact of TGIP may be overwhelmed by the fact that males

are able to sire offspring (that inherit their parasite resistance) up to 10 months posthumously

(Lopez-Sepulcre et al., 2013), and engage in coercive mating so will likely mate (Magurran,

2005), even though females discriminate against males who are infected (Kennedy et al., 1987),

or those who may lack resistance (Stephenson et al., 2020). How the TGIP we observe here

affects dynamics in nature remains unclear, therefore, but the magnitude of the effect we

observe should mean it warrants future investigation.

More broadly, our comparison of the maternally and paternally derived transgenerational effects

of infection contributes novel insight valuable across taxa. In accordance with parental

investment theory, we found that males appear to prioritise offspring quantity, and females

quality, in the face of infection. We also show that maternal and paternal contributions to

offspring immunity can differ significantly. Thus, our observed transgenerational effects of

infection potentially influence disease dynamics and host-parasite co-evolutionary dynamics in

natural communities (McIntire et al., 2023).

References:

Adrian-Kalchhauser, I., Sultan, S. E., Shama, L. N. S., Spence-Jones, H., Tiso, S., Valsecchi,

C. I. K., & Weissing, F. J. (2020). Understanding'non-genetic'inheritance: insights from

molecular-evolutionary crosstalk. *Trends in Ecology & Evolution*, *35*(12), 1078–1089.

Åkesson, M., Bensch, S., & Hasselquist, D. (2007). Genetic and phenotypic associations in

morphological traits: a long term study of great reed warblers*Acrocephalus*

arundinaceus. *Journal of Avian Biology*, *38*(1), 58–72.

Åkesson, M., Hasselquist, D., Tarka, M., & Hansson, B. (2008). Estimating Heritabilities and

Genetic Correlations: Comparing the 'Animal Model' with Parent-Offspring Regression

Using Data from a Natural Population. *PloS One*, *3*(3), e1739.

Armour, E. M., Bruner, T. L., Hines, J. K., & Butler, M. W. (2020). Low-dose immune challenges

result in detectable levels of oxidative damage. *The Journal of Experimental Biology*,

223(Pt 6). https://doi.org/10.1242/jeb.220095

Arnqvist, G., & Rowe, L. (2005). *Sexual conflict*. Princeton University Press.

Beemelmanns, A. (2019). *Biparental Immune Priming in the Broad-nosed Pipefish (Syngnathus*

Typhle). Universitätsbibliothek Kiel.

Beemelmanns, A., & Roth, O. (2016). Bacteria-type-specific biparental immune priming in the

pipefish Syngnathus typhle. *Ecology and Evolution*, *6*(18), 6735–6757.

Beemelmanns, A., & Roth, O. (2017). Grandparental immune priming in the pipefish

Syngnathus typhle. *BMC Evolutionary Biology*, *17*(1), 44.

Billi, A. C., Kahlenberg, J. M., & Gudjonsson, J. E. (2019). Sex bias in autoimmunity. *Current*

Opinion in Rheumatology, *31*(1), 53–61.

Boots, M., & Bowers, R. G. (2004). The evolution of resistance through costly acquired

immunity. *Proceedings. Biological Sciences / The Royal Society*, *271*(1540), 715–723.

Breheny, P., & Burchett, W. (2017). Visualization of Regression Models Using visreg. *The R*

Journal, *9*(2), 56.

Breuil, G., Vassiloglou, B., Pepin, J. F., & Romestand, B. (1997). Ontogeny of IgM-bearing cells

and changes in the immunoglobulin M-like protein level (IgM) during larval stages in sea

bass (Dicentrarchus labrax). *Fish & Shellfish Immunology*, *7*(1), 29–43.

Brooks, M., Kristensen, K., Benthem, K. van, Magnusson, A., Berg, C., Nielsen, A., Skaug, H.,

Mächler, M., & Bolker, B. (2017). GlmmTMB balances speed and flexibility among

packages for zero-inflated generalized linear mixed modeling. *The R Journal*, *9*(2), 378.

Brown, L. L., Evelyn, T. P. T., & Iwama, G. K. (1997). Specific protective activity demonstrated

in eggs of broodstock salmon injected with rabbit antibodies raised against a fish

pathogen. *Diseases of Aquatic Organisms*, *31*, 95–101.

Clark, D. R., Young, K. A., J., K., Moore, P. J., Evans, A., & Stephenson, J. F. (2023).

Aggregation of symbionts on hosts depends on interaction type and host traits.

Ecography.

Cunningham, E. J., & Russell, A. F. (2000). Egg investment is influenced by male attractiveness

in the mallard. *Nature*, *404*(6773), 74–77.

Dargent, F., Rolshausen, G., Hendry, A. P., Scott, M. E., & Fussmann, G. F. (2016). Parting

ways: parasite release in nature leads to sex-specific evolution of defence. *Journal of*

Evolutionary Biology, *29*(1), 23–34.

Duffield, K. R., Bowers, E. K., Sakaluk, S. K., & Sadd, B. M. (2017). A dynmaic threshold model

for terminal investment. *Behavioral Ecology and Sociobiology*, *71*(12), 185.

Encel, S. A., Simpson, E. K., Schaerf, T. M., & Ward, A. J. W. (2023). Immune challenge affects

reproductive behaviour in the guppy (*Poecilia reticulata*). *Royal Society Open Science*,

10(8). https://doi.org/10.1098/rsos.230579

Evans, J. P., Lymbery, R. A., Wiid, K. S., Rahman, M. M., & Gasparini, C. (2017). Sperm as

moderators of environmentally induced paternal effects in a livebearing fish. *Biology*

Letters, *13*(4). https://doi.org/10.1098/rsbl.2017.0087

Falconer, D. S., & Mckay, T. F. C. (1997). *Introduction to Quantitative Genetics (Vol. 4).*

Longman Higher Education.

Fischer, B., Taborsky, B., & Kokko, H. (2011). How to balance the offspring quality-quantity

tradeoff when environmental cues are unreliable. *Oikos* , *120*(2), 258–270.

Fraser, B. A., & Neff, B. D. (2009). MHC class IIB additive and non-additive effects on fitness

measures in the guppy Poecilia reticulata. *Journal of Fish Biology*, *75*(9), 2299–2312.

Fraser, B. A., Ramnarine, I. W., & Neff, B. D. (2010). Selection at the MHC class IIB locus

across guppy (Poecilia reticulata) populations. *Heredity*, *104*(2), 155–167.

Gapp, K., von Ziegler, L., Tweedie-Cullen, R. Y., & Mansuy, I. M. (2014). Early life epigenetic

programming and transmission of stress-induced traits in mammals: how and when can

environmental factors influence traits and their transgenerational inheritance?

BioEssays: News and Reviews in Molecular, Cellular and Developmental Biology, *36*(5),

491–502.

Gasparini, J., McCoy, K. D., Tveraa, T., & Boulinier, T. (2002). Related concentrations of

specific immunoglobulins against the Lyme disease agent *Borrelia burgdorferi sensu lato*

in eggs, young and adults of the kittiwake (*Rissa tridactyla*). *Ecology Letters*, *5*(4), 519–

524.

Gheorghiu, C., Marcogliese, D. J., & Scott, M. (2006). Concentration-dependent effects of

waterborne zinc on population dynamics of Gyrodactylus turnbulli (Monogenea) on

isolated guppies (Poecilia reticulata). *Parasitology*, *132*(Pt 2), 225–232.

Gheorghiu, C., Marcogliese, D. J., & Scott, M. E. (2009). Temporal dynamics of epidermal

responses of guppies Poecilia reticulata to a sublethal range of waterborne zinc

concentrations. *Journal of Fish Biology*, *75*(10), 2642–2656.

Gheorghiu, C., Marcogliese, D. J., & Scott, M. E. (2012). Waterborne zinc alters temporal

dynamics of guppy *Poecilia reticulata* epidermal response to *Gyrodactylus turnbulli*

(Monogenea). *Diseases of Aquatic Organisms*, *98*, 143–153.

Gilbey, J., Verspoor, E., Mo, T. A., Sterud, E., Olstad, K., Hytterød, S., Jones, C., & Noble, L.

(2006). Identification of genetic markers associated with Gyrodactylus salaris resistance

in Atlantic salmon Salmo salar. *Diseases of Aquatic Organisms*, *71*(2), 119–129.

Graham, A. L. (2013). Optimal immunity meets natural variation: the evolutionary biology of host

defence. *Parasite Immunology*, *35*(11), 315–317.

Hagmayer, A., Furness, A. I., & Pollux, B. J. A. (2020). Parasite infestation influences life history

but not boldness behavior in placental live-bearing fish. *Oecologia*, *194*(4), 635–648.

Hagmayer, A., Furness, A. I., Reznick, D. N., & Pollux, B. J. A. (2018). Maternal size and body

condition predict the amount of post-fertilization maternal provisioning in matrotrophic

fish. *Ecology and Evolution*, *8*(24), 12386–12396.

Harris, P. D., & Lyles, A. M. (1992). Infections of Gyrodactylus bullatarudis and Gyrodactylus

turnbulli on guppies (Poecilia reticulata) in Trinidad. *The Journal of Parasitology*, *78*(5),

912–914.

Hartig, F. (2022). *DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)*

Regression Models. https://CRAN.R-project.org/package=DHARMa

Heard, E., & Martienssen, R. A. (2014). Transgenerational epigenetic inheritance: myths and

mechanisms. *Cell*, *157*(1), 95–109.

Herdegen-Radwan, M., Phillips, K. P., Babik, W., Mohammed, R. S., & Radwan, J. (2020).

Balancing selection versus allele and supertype turnover in MHC class II genes in

guppies. *Heredity*. https://doi.org/10.1038/s41437-020-00369-7

Houston, A. I., McNamara, J. M., Barta, Z., & Klasing, K. C. (2007). The effect of energy

reserves and food availability on optimal immune defence. *Proceedings. Biological*

Sciences / The Royal Society, *274*(1627), 2835–2842.

Hu, J., Pérez-Jvostov, F., Blondel, L., & Barrett, R. D. H. (2018a). Genome-wide DNA

methylation signatures of infection status in Trinidadian guppies (Poecilia reticulata).

Molecular Ecology, *27*(15), 3087–3102.

Hu, J., Pérez-Jvostov, F., Blondel, L., & Barrett, R. D. H. (2018b). Genome-wide DNA

methylation signatures of infection status in Trinidadian guppies (Poecilia reticulata).

Molecular Ecology, *27*(15), 3087–3102.

Jog, M. G., Sackett, M. E., Kisty, S. D., Hansen, J. A., & Stephenson, J. F. (2022). The

behaviour of infected guppies depends on social context, parasite tolerance and host

sex. *Animal Behaviour*, *187*, 97–104.

Kennedy, C. E. J., Endler, J. A., Poynton, S. L., & McMinn, H. (1987). Parasite load predicts

mate choice in guppies. *Behavioral Ecology and Sociobiology*, *21*(5), 291–295.

Khan, I., Agashe, D., & Rolff, J. (2017). Early-life inflammation, immune response and ageing.

Proceedings. Biological Sciences / The Royal Society, *284*(1850).

https://doi.org/10.1098/rspb.2017.0125

Konczal, M., Ellison, A., Phillips, K. P., Radwan, J., Mohammed, R. S., Cable, J., & Chadzinska,

M. (2020). RNA-Seq analysis of the guppy immune response against Gyrodactylus

bullatarudis infection. *Parasite Immunology*, e12782.

Kvist, J., Athanàsio, C. G., Pfrender, M. E., Brown, J. B., Colbourne, J. K., & Mirbahai, L. (2020).

A comprehensive epigenomic analysis of phenotypically distinguishable, genetically

identical female and male Daphnia pulex. *BMC Genomics*, *21*(1), 17.

López, S. (1998). Acquired resistance affects male sexual display and female choice in guppies.

Proceedings. Biological Sciences / The Royal Society, *265*(1397), 717–723.

López, S. (1999). Parasitized female guppies do not prefer showy males. *Animal Behaviour*,

57(5), 1129–1134.

Lopez-Sepulcre, A., Gordon, S. P., Paterson, I. G., Bentzen, P., & Reznick, D. N. (2013).

Beyond lifetime reproductive success: the posthumous reproductive dynamics of male

Trinidadian guppies. *Proceedings of the Royal Society of London, Series B: Biological*

Sciences, *280*, 20131116.

Madhavi, R., & Anderson, R. M. (1985). Variability in the susceptibility of the fish host, Poecilia reticulata, to infection with Gyrodactylus bullatarudis (Monogenea). *Parasitology*, *91*(3), 531–544.

Magurran, A. E. (2005). *Evolutionary ecology: the Trinidadian guppy* (pp. 1–224). Oxford Univeristy Press.

McIntire, K. M., Dziuba, M. K., Haywood, E. B., Robertson, M. L., Vaandrager, M., Baird, E., Corcoran, F., Cortez, M. H., & Duffy, M. A. (2023). Transgenerational virulence: Maternal pathogen exposure reduces offspring fitness. In *bioRxiv* (p. 2023.03.14.532659). https://doi.org/10.1101/2023.03.14.532659

Metcalf, C. J. E., Roth, O., & Graham, A. L. (2020). Why leveraging sex differences in immune trade-offs may illuminate the evolution of senescence. *Functional Ecology*, *34*(1), 129–140.

Morris, J., Darolti, I., van der Bijl, W., & Mank, J. E. (2020). High-resolution characterization of male ornamentation and re-evaluation of sex linkage in guppies. *Proceedings. Biological Sciences*, *287*(1937), 20201677.

Norazmi-Lokman, N. H., Purser, G. J., & Patil, J. G. (2016). Gravid Spot Predicts Developmental Progress and Reproductive Output in a Livebearing Fish, Gambusia holbrooki. *PloS One*, *11*(1), e0147711.

Peters, A., Delhey, K., Nakagawa, S., Aulsebrook, A., & Verhulst, S. (2019). Immunosenescence in wild animals: meta-analysis and outlook. *Ecology Letters*, *22*(10), 1709–1722.

Phillips, K. P., Cable, J., Mohammed, R. S., Herdegen-Radwan, M., Raubic, J., Przesmycka, K. J., van Oosterhout, C., & Radwan, J. (2018). Immunogenetic novelty confers a selective advantage in host-pathogen coevolution. *Proceedings of the National Academy of Sciences of the United States of America*, *115*(7), 1552–1557.

Pizzolon, M., Giacomello, E., Marri, L., Marchini, D., Pascoli, F., Mazzoldi, C., & Rasotto, M. B.

(2010). When fathers make the difference: efficacy of male sexually selected

antimicrobial glands in enhancing fish hatching success. *Functional Ecology*, *24*(1), 141–

148.

R Core Team. (2020). *R: A language and environment for statistical computing. R Foundation*

for Statistical Computing.

Reid, J. M., Arcese, P., Keller, L. F., & Hasselquist, D. (2006). Long-term maternal effect on

offspring immune response in song sparrows Melospiza melodia. *Biology Letters*, *2*(4),

573–576.

Reynolds, J. D., & Gross, M. R. (1992). Female mate preference enhances offspring growth and

reproduction in a fish,Poecilia reticulata. *Proceedings. Biological Sciences / The Royal*

Society, *250*(1327), 57–62.

Rolff, J. (2002). Bateman's principle and immunity. *Proceedings. Biological Sciences / The*

Royal Society, *269*(1493), 867–872.

Roth, O., Beemelmanns, A., Barribeau, S. M., & Sadd, B. M. (2018). Recent advances in

vertebrate and invertebrate transgenerational immunity in the light of ecology and

evolution. *Heredity*, *121*(3), 225–238.

Roth, O., Klein, V., Beemelmanns, A., Scharsack, J. P., & Reusch, T. B. H. (2012). Male

pregnancy and biparental immune priming. *The American Naturalist*, *180*(6), 802–814.

Rutkowski, N.-A. J., McNamara, K. B., Jones, T. M., & Foo, Y. Z. (2023). Trans-generational

immune priming is not mediated by the sex of the parent primed: a meta-analysis of

invertebrate data. *Biological Reviews of the Cambridge Philosophical Society*, *98*(4),

1100–1117.

Schelkle, B., Faria, P. J., Johnson, M. B., van Oosterhout, C., & Cable, J. (2012). Mixed

Infections and Hybridisation in Monogenean Parasites. *PloS One*, *7*(7), e39506.

Schmid-Hempel, P. (2003). Variation in immune defence as a question of evolutionary ecology. *Proceedings. Biological Sciences / The Royal Society*, *270*(1513), 357–366.

Sheldon, B. C., & Verhulst, S. (1996). Ecological immunology: costly parasite defences and trade-offs in evolutionary ecology. *Trends in Ecology & Evolution*, *11*(8), 317–321.

Shikano, T., & Taniguchi, N. (2005). Relationships between brood size and offspring body size in an ovoviviparous fish: maternal effects and genetic trade-off. *Journal of Experimental Zoology. Part A, Comparative Experimental Biology*, *303*(8), 635–642.

Simms, E. L., & Triplett, J. K. (1996). PATERNAL EFFECTS IN INHERITANCE OF A PATHOGEN RESISTANCE TRAIT IN IPOMOEA PURPUREA. *Evolution; International Journal of Organic Evolution*, *50*(6), 2178–2186.

Stein, L. R., & Hoke, K. (2022). Parental and individual experience with predation risk interact in shaping phenotypes in a sex-specific manner. *Animal Behaviour*, *191*, 75–89.

Stephenson, J. F. (2019). Parasite-induced plasticity in host social behaviour depends on sex and susceptibility. *Biology Letters*, *15*(11), 20190557.

Stephenson, J. F., Kinsella, C., & Cable, J. (2016). A further cost for the sicker sex? Evidence for male-biased parasite-induced vulnerability to predation. *Ecology*. https://onlinelibrary.wiley.com/doi/abs/10.1002/ece3.2049

Stephenson, J. F., Perkins, S. E., & Cable, J. (2018). Transmission risk predicts avoidance of infected conspecifics in Trinidadian guppies. *The Journal of Animal Ecology*, *87*(6), 1525–1533.

Stephenson, J. F., Stevens, M., Troscianko, J., & Jokela, J. (2020). The size, symmetry, and color saturation of a male guppy's ornaments forecast his resistance to parasites. *The American Naturalist*, *196*(5), 597–608.

Stephenson, J. F., van Oosterhout, C., & Cable, J. (2015). Pace of life, predators and parasites: predator-induced life-history evolution in Trinidadian guppies predicts decrease in parasite tolerance. *Biology Letters*, *11*(11). https://doi.org/10.1098/rsbl.2015.0806

Stephenson, J. F., Van Oosterhout, C., & Mohammed, R. S. (2015). Parasites of Trinidadian

guppies: evidence for sex-and age-specific trait-mediated indirect effects of predators.

Ecology. https://esajournals.onlinelibrary.wiley.com/doi/abs/10.1890/14-0495.1

Stephenson, J. F., Young, K. A., Fox, J., Jokela, J., Cable, J., & Perkins, S. E. (2017). Host

heterogeneity affects both parasite transmission to and fitness on subsequent hosts.

Philosophical Transactions of the Royal Society of London. Series B, Biological

Sciences, *372*(1719). https://doi.org/10.1098/rstb.2016.0093

Stoehr, A. M., & Kokko, H. (2006). Sexual dimorphism in immunocompetence: what does life-

history theory predict? *Behavioral Ecology: Official Journal of the International Society*

for Behavioral Ecology, *17*(5), 751–756.

Tizard, I. (2008). Sickness behavior, its mechanisms and significance. *Animal Health Research*

Reviews / Conference of Research Workers in Animal Diseases, *9*(1), 87–99.

Trivers, R. L. (1972). Parental investment and sexual selection. In B. Campbell (Ed.), *Sexual*

selection and the descent of man. Aldine Publishing Company.

Tschirren, B., & Richner, H. (2006). Parasites shape the optimal investment in immunity.

Proceedings. Biological Sciences / The Royal Society, *273*(1595), 1773–1777.

van der Bijl, W., Shu, J. J., Goberdhan, V. S., Sherin, L. M., Cortázar-Chinarro, M., Corral-

López, A., & Mank, J. E. (2023). Deep learning reveals the role of copy number variation

in the genetic architecture of a highly polymorphic sexual trait. In *bioRxiv* (p. 2023.09.

29.560175). https://doi.org/10.1101/2023.09.29.560175

van Oosterhout, C., Mohammed, R. S., Hansen, H., Archard, G. A., McMullan, M., Weese, D. J.,

& Cable, J. (2007). Selection by parasites in spate conditions in wild Trinidadian guppies

(*Poecilia reticulata*). *International Journal for Parasitology*, *37*(7), 805–812.

Vantaux, A., Dabiré, K. R., Cohuet, A., & Lefèvre, T. (2014). A heavy legacy: offspring of

malaria-infected mosquitoes show reduced disease resistance. *Malaria Journal*, *13*, 442.

Vilcinskas, A. (2021). Mechanisms of transgenerational immune priming in insects.

Developmental and Comparative Immunology, *124*, 104205.

Viney, M. E., Riley, E. M., & Buchanan, K. L. (2005). Optimal immune responses:

immunocompetence revisited. *Trends in Ecology & Evolution*, *20*(12), 665–669.

Walsman, J. C., Janecka, M. J., Clark, D. R., Kramp, R. D., Rovenolt, F., Patrick, R.,

Mohammed, R. S., Konczal, M., Cressler, C. E., & Stephenson, J. F. (2022). Shoaling

guppies evade predation but have deadlier parasites. *Nature Ecology & Evolution*, *6*(7),

945–954.

Wassarman, P. M., Jovine, L., & Litscher, E. S. (2001). A profile of fertilization in mammals.

Nature Cell Biology, *3*(2), E59-64.

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis* (Vol. 174, pp. 245–246).

Springer-Verlag.

Willis, A. R., Sukhdeo, R., & Reinke, A. W. (2021). Remembering your enemies: mechanisms of

within-generation and multigenerational immune priming in Caenorhabditis elegans. *The*

FEBS Journal, *288*(6), 1759–1770.

Wolf, J. B., & Wade, M. J. (2009). What are maternal effects (and what are they not)?

Philosophical Transactions of the Royal Society of London. Series B, Biological

Sciences, *364*(1520), 1107–1115.

Zhang, C., Hediger, M. L., Albert, P. S., Grewal, J., Sciscione, A., Grobman, W. A., Wing, D. A.,

Newman, R. B., Wapner, R., D'alton, M. E., Skupski, D., Nageotte, M. P., Ranzini, A. C.,

Owen, J., Chien, E. K., Craigo, S., Sungduk, K. I. M., Grantz, K. L., & Buck Louis, G. M.

(2018). Maternal obesity and longitudinal ultrasound measures of fetal growth: Findings

from the NICHD Fetal Growth Studies-Singletons. *JAMA Pediatrics*, *172*(1), 24.

Figures:

Fig. 1: Pairs in which fathers (darker blue) had more intense infections produced broods

quicker, whereas the intensity of the mother's (lighter yellow) infection did not affect

birth latency. Points represent the partial residuals from the model described in the main text,

lines give the model fits, and shading the standard error. We quantified infection intensity using

'infection integral', or the area under the curve of infection load over time.

Fig. 2: Among offspring with an infected parent, older fish tended to experience more intense infections, but no such relationship exists among offspring of two parasite-naïve parents. Points represent the partial residuals from the model described in the main text, lines give the model fits, and shading the standard error. We quantified infection intensity using 'infection integral', or the area under the curve of infection load over time.

Fig. 3: A father's infection intensity significantly predicts the infection intensity that his offspring experience, but no such relationship exists between the infection intensity experienced by a mother and that of her offspring. Points represent the partial residuals from the model described in the main text, lines give the model fits, and shading the standard error. We quantified infection intensity using 'infection integral', or the area under the curve of infection load over time.