Within-species floral evolution reveals convergence in adaptive walks during incipient pollinator shift

Katherine E. Wenzell† [1,2,3], Mikhaela Neequaye†[1], Pirita Paajanen[1], Lionel Hill[1], Paul Brett[1], Kelsey J.R.P. Byers[1]*

[1] John Innes Centre, Norwich, UK, [2] California Academy of Sciences, San Francisco, California, USA, [3] University of Maryland, College Park, Maryland, USA
† Co-first authors
*Author for correspondence (Kelsey.Byers@jic.ac.uk)

ABSTRACT

Understanding how evolution proceeds from molecules to organisms to interactions is critical for evolutionary biology and requires integrative studies that span biological levels. By linking phenotypic changes with associated genes and fitness implications, we gain insight on the repeatability of evolution and how adaptive walks move organisms between fitness peaks, which remains poorly understood at intermediate stages, *e.g.* potential fitness valleys. Floral trait evolution can proceed rapidly, confer reproductive isolation, and provides numerous examples of convergent evolution associated with pollinator guilds. Within the monkeyflowers (*Mimulus* sect. *Erythranthe*), intraspecific shifts to yellow flowers within red hummingbird-pollinated species have arisen at least twice and may represent incipient shifts between optima of pollinator adaptation. Here, we compare two natural yellow-flowered forms of *M. cardinalis* and *M. verbenaceus* to their red conspecifics in floral phenotypes, biochemistry, transcriptomic and genomic variation, and interactions with pollinators. We find convergent evolution in the adaptive walks of yellow morphs in both species, with consistent changes in traits of large effect (floral pigments and associated gene expression), which result in strong preference for yellow flowers by bumblebees. Shifts in additional traits (scent emissions and floral opening size) are also consistent with bee pollination, which may represent smaller-effect steps in an ongoing walk from hummingbird to bee pollination. By examining intraspecific variation representing incipient pollinator shifts in two related species, this work elucidates how adaptive walks proceed at early stages by highlighting how convergent large effect mutations (floral color) may drive pollinator attraction, preceding smaller effect changes controlling pollinator fit and reward access. This work integrates phenotypic, biochemical, genomic, and behavioral evidence to shed light on how repeated steps in adaptive walks to new fitness optima may impact reproductive isolation and incipient speciation via convergent evolution.

Key words: Convergent evolution, pollination, floral pigments, floral scent, behaviour, monkeyflowers, *Mimulus,* fitness landscapes

INTRODUCTION

Understanding how adaptations in complex traits arise remains a central question in evolutionary biology (Fisher, 1934; Orr, 1998). Theory predicts that adaptive walks to new fitness optima involve a few mutations of large effect and many "steps" of smaller-effect changes (Orr, 2005), but uncertainty remains regarding how adaptive walks proceed at intermediate stages and when moving between peaks in a fitness landscape (Arnold et al., 2001). Linking phenotype to genotype to fitness within multilevel systems in an integrative way (Byers et al., 2017) allows us to assess the repeatability and convergence of adaptation. One such multilevel system is floral trait evolution, which can display phenotypic convergence via selection by specific pollinator guilds, which may also confer reproductive isolation (Fenster et al., 2004; Dellinger 2020). Pollinators are important for flowering plant diversification (Fenster et al., 2004; Lowry et al., 2008; van der Niet and Johnson, 2012; van der Niet et al., 2014), but questions remain about how pollinator-mediated speciation arises at early stages from intraspecific floral variation (Herrera et al., 2006; Kay and Sargent, 2009; Wenzell et al., 2023). Combining knowledge of key genes underlying floral shifts and their impact on pollinator behavior thus provides a powerful assessment of how evolution drives species diversity.

Previous work suggests that genes of large effect can alter floral traits and therefore reproductive isolation via pollinator shifts (Bradshaw and Schemske, 2003; Hoballah et al., 2007; Schlüter et al., 2011), which often involves floral color transitions (Hopkins and Rausher, 2012; Stankowski et al., 2017; Liang et al., 2023, Wessinger et al., 2023). Understanding the underlying genetic changes can reveal evolutionary mechanisms driving phenotypic changes, e.g. drift or selection (Hopkins & Rausher 2011; Kellenberger et al., 2019), which may result in phenotypic convergence via independent mutations (Quattrocchio et al., 1999; Cooley et al., 2011; Smith & Rausher 2011; Sheehan et al., 2016), sorting of standing ancestral variation (Berardi et al., 2022; Sánchez-Cabrera et al., 2023), or adaptive introgression via gene flow (Stankowski & Streisfeld 2015, Short and Streisfeld, 2023). Nonetheless, few if any studies have assessed genetic changes associated with repeated natural floral color transitions in multiple related species and experimentally quantified their impact on pollinator behavior, an approach that would shed light on the repeatability and convergence of pollinator-mediated evolution.

Whether recent floral color transitions are accompanied by changes in other floral traits impacting pollinator behavior remains uncertain (Sutherland and Vickery, 1993; Schiestl and Johnson, 2013) and provides an opportunity to explore how adaptive walks may proceed between peaks in a fitness landscape. Because suites of floral traits adapted to different pollinator guilds represent a key example of complex trait adaptation (Wessinger et al., 2023), exploring variation in additional floral traits in intraspecific floral color transitions (hypothesized to represent pollinator shifts (Vickery, 1995)), provides insight into the nature of early stages of adaptive evolution. Within the genus *Mimulus* (monkeyflowers, Phrymaceae, *sensu* (Lowry et al. 2019)), section *Erythranthe* offers ideal case studies to investigate floral trait evolution via pollinator-mediated selection (Bradshaw and Schemske, 2003; Byers et al., 2014a,b) and has

become an emerging model system for floral trait evolution (Wu et al., 2008; Yuan et al., 2013;
Yuan, 2019). Section *Erythranthe* includes two independent shifts from insect to hummingbird
pollination (Beardsley et al., 2003, Nelson et al. 2021). These shifts include transitions to red
flowers, which result from high concentrations of anthocyanins and carotenoids (Yuan et al.,
2014; LaFountain et al., 2015), in addition to changes in floral morphology, scent, and reward
(Bradshaw et al., 1995; Byers et al., 2014a).
Within the two geographically widespread hummingbird-pollinated members of section
Erythranthe, *M. cardinalis* and *M. verbenaceus*, naturally-occurring yellow-flowered morphs
have arisen and have become well-established in several populations at range edges (Vickery,
1995). Previous work on these yellow morphs found evidence that they were preferred by
bumblebee pollinators compared to the red-flowered wild type in both species (Vickery and
Vickery, 1992), as well as evidence of assortative mating within the yellow morph due to
pollinator fidelity (Vickery, 1995). Yellow floral color was hypothesized to result from a loss of
anthocyanins via a single recessive locus (Vickery, 1995). In *M. cardinalis* this is due to
expression changes or deletion of the gene *PELAN*, which is responsible for the production of
anthocyanins in petal lobes (Yuan et al., 2014). Despite previous work on these color
polymorphisms, it remains unknown whether these stable floral morphs also differ in other
pollinator-relevant floral traits (including floral morphology, nectar, pigments, and floral scent),
which may represent additional steps along an adaptive walk between hummingbird and bee
pollination syndromes by further adapting these morphs to bee pollination via modifications to
floral attraction and mechanical fit traits. Does the evolution of novel color morphs with
differential pollinator attraction involve only color shifts (i.e., "first step" mutations of large
effect), or are they also altering in more complex ways (via many traits of small effect) to adapt
to a novel pollinator?
In order to identify whether these repeated floral color transitions may represent early
stages of pollinator-mediated evolution, we investigated whether these independent shifts
showed evidence of convergent evolution between the two species (*sensu* Arendt and Reznick,
2008) at the phenotypic and genetic levels. We address the following: 1) What are the
phenotypic, biochemical, and genetic changes underlying the red-yellow shifts? 2) Is there
evidence for convergence between the yellow morphs of both species in floral color and other
traits? 3) Are shifts across floral traits of yellow morphs consistent with an adaptive walk toward
bee pollination? This approach assesses if phenotypic and genetic changes are best explained by
shared (i.e., adaptive introgression of alleles or a shared ancestral polymorphism) or independent
(i.e., convergence due to consistent selection pressure by pollinators) sources of genetic
variation. By expanding our understanding of the direction, order, and integration of the
evolution of floral traits in response to an incipient pollinator shift, this work sheds light on how
adaptive walks may proceed at intermediate stages, as complex phenotypes move between peaks
in a fitness landscape. Integrating studies of floral phenotypic, biochemical, and genetic variation
with their impact on pollinator behavior within multiple species provides a comprehensive
investigation of the repeatability of early stages of adaptive evolution.

RESULTS

   Abbreviations and summary-- We used two existing inbred lines of *M. verbenaceus* (red: MVBL, hereafter MvR; yellow: MVYL, hereafter MvY, see Methods) and two existing inbred lines of *M. cardinalis* (red: CE10, hereafter McR; yellow: SM, hereafter McY).

   To characterize these repeated floral transitions from red to yellow flowers, we compared MvY and McY to their red conspecific forms in terms of floral phenotypes, biochemistry, transcriptomic and genomic variation, and interactions with bumblebee pollinators. Despite contrasting differences in some aspects of floral phenotype (nectar guides, corolla tube length, and scent composition), we find evidence of convergent evolution in pigments, gene expression, and pollinator responses in both yellow forms (Table 1). Yellow morphs share less restrictive floral tubes, increased scent emission, and patterns of carotenoid production and expression of key carotenoid biosynthetic genes. Further, we find some evidence that homologues of R2R3-MYB transcription factor (*PELAN*) are likely involved in the lack of anthocyanin expression in the petal lobes of both yellow forms, which attract novel bumblebee pollinators at a rate of 2:1 over red flowers. Finally, analysis of whole genome sequence variation reveals independent paths of molecular evolution, even when the same genes are involved, consistent with convergent evolution of yellow flowers with increased attractiveness to bee pollinators.

Floral color: steps of large effect in an adaptive walk

   Floral reflectance—We compared spectral reflectance of petal lobes of all morphs by plotting reflectance curves and performing principal component analysis of reflectance values at wavelengths from 300-700nm. We found that red flowers of both species (MvR and McR) appeared similar in both human vision and reflectance curves, while yellow morphs (MvY and McY) differed in both human vision and reflectance (Fig. 1A; Fig. 1B; Figure S2). While petal lobes of both red morphs reflect almost no UV, MvY and particularly McY are modestly reflective in the UV range, with a subtle "bullseye" pattern in both (Figure S3).

   Floral pigments–Total anthocyanins extracted from corolla tissues did not vary among species ($F_{1, 16}$= 0.89, p = 0.36; Figure 1C) but varied between color morphs within species ($F_{2,16}$= 207.8, p < 0.0001), with higher levels in red morphs than their yellow conspecifics (Table S2). Total carotenoids varied significantly among species ($F_{1, 16}$= 15.2, p = 0.001; Figure 1D) and between color morphs within species ($F_{2,16}$ = 14.04, p = 0.0003). Both yellow-flowered morphs had significantly higher total carotenoids than their red conspecifics. This suggests that both species' shifts from red to yellow flowers were mediated by a simultaneous loss of anthocyanins and an increase in carotenoids.

Individual anthocyanins—We characterized the individual anthocyanins produced by
each morph and found that anthocyanin accumulation did not show any consistent patterns
across morphs (Figure 2A). Two core anthocyanidins (cyanidin and pelargonidin) and eight
anthocyanins were found in the two species. Three anthocyanins were found in more than one
morph: cyanidin-glucoside, cyanidin-rhamnose-glucoside and pelargonidin-rhamnose-glucoside.
The lowest total anthocyanin accumulation was found in the wholly yellow MvY, which
produced cyanidin-rhamnose-glucoside and pelargonidin-rhamnose-glucoside at almost
undetectable levels. In contrast, McY, which has red nectar guides (Figure 1Aiii), produces both
anthocyanins found in MvY as well as cyanidin-glucoside. Of these three yellow morph
anthocyanins, cyanidin-rhamnose-glucoside was found at the highest levels in McY, with the
others at barely detectable levels. In contrast to McY, McR accumulates 2.4-fold more cyanidin-
rhamnose-glucoside (Welch's t-test, t=8.899, df=4, p =0.0009) in addition to 15.6-fold more
pelargonidin-rhamnose-glucoside (Welch's t-test, t=9.763, df=4, p = 0.0006) . In addition to the
three anthocyanins shared with McY, McR also produces an additional isomer of pelargonidin-
rhamnose-glucoside. In contrast with the other morphs, MvR shows the highest diversity of
anthocyanins (five, three unique to MvR). Interestingly, MvR does not share a single
anthocyanin with MvY. This is in contrast with McY, which shares all three of its anthocyanins
with McR. Moreover, MvR is the only line to accumulate anthocyanins with pentose and acetate
decorations. In addition, pelargonidin-glucoside is only found in MvR, whereas pelargonidin-
rhamnose-glucoside is found in the other three morphs. This suggests a potential loss of ability to
add rhamnose in addition to the gain of ability to add pentose and acetate decorations in MvR.
MvR also contains significantly higher levels of cyanidin-glucoside when compared to the two
M. cardinalis lines (Welch's t-test, t=13.14 df=4, p=0.0002 for McR, t=13.99 df=4, p=0.0002
for McY).

Anthocyanin biosynthesis and regulatory genes—To better characterize the loss of red
pigmentation in the yellow morphs, we analyzed expression of anthocyanin biosynthesis
pathway genes in corolla tissue. No significant difference was found between core anthocyanin
biosynthesis genes across the four morphs, with the exception of *chalcone synthase* (*CHS*) and
flavanone-3-hydroxylase (*F3H*, only significantly different in *M. verbenaceus*) (Figure 2B). *CHS*
is significantly upregulated in both yellow morphs compared to their red counterparts, (Welch's
t-test, t=13.02 df=2, p=0.0058 for McY, t=8.880 df=2, p=0.0124 for MvY). These enzymatic
reactions occur early in anthocyanin biosynthesis and lead not only to the production of
anthocyanins but also other flavonoids (Table S6) (Holton & Cornish 1995, Wang 2020).

By contrast, significant differences were found in previously characterized *Mimulus*
anthocyanin regulatory genes (Yuan et al., 2014). *NEGAN* is only expressed in McY (Figure 2C),
consistent with published work demonstrating its role in McY's red-spotted corolla throat (Yuan
et al., 2014). MvY does not have any red spots or associated *NEGAN* expression. The published
PELAN is highly upregulated in MvY despite accumulating the lowest total anthocyanins of the
four morphs(Welch's t-test, t=6.117 df=2, p=0.0257) . Closer inspection of the whole genomic

sequence in order to confirm the identity of this *PELAN* homolog showed it is deleted in McY,
again in agreement with (Yuan et al., 2014) – whereas it is both present and highly expressed in
MvY. Six *PELAN* homologs were found in the two species' genomes, all clustered on
chromosome 4. These were named *PELAN*(1-6) with *PELAN*(1) being the published *PELAN*.
PELAN(3) and *PELAN*(4) were not expressed in corolla tissue of any line. *PELAN*(6) was only
expressed in MvR and *PELAN*(2) and *PELAN*(5) were only expressed in McR (Figure 2C,Table
S7). This suggests a historical set of duplications resulting in multiple diverged copies of *PELAN*
with various functions or expression domains, which likely explains the lack of consistency
found in the different anthocyanins in these morphs.

Flavonoids—Due to the increased expression of *CHS* (the enzyme responsible for the
first committed step in flavonoid biosynthesis, Holton & Cornish 1995) in yellow morphs, we
also briefly investigated the accumulation of additional flavonoids. Preliminary analyses could
not find a relationship between anthocyanins, their core biosynthesis genes, and flavonoids
detectable at a wavelength of 350nm (Figure S12, Table S8).

Individual carotenoids—Yellow morphs had higher levels of total carotenoids compared
to their red conspecifics. In order to better characterize this shift, we quantified individual
carotenoids. While none of the eight anthocyanins characterized could be found across all
morphs, several carotenoids – first characterized in *Mimulus* by LaFountain et al., 2015 – were
detected across all four lines (Figure 3A Table S9). Zeaxanthin (first in the biosynthesis
pathway) is found in significantly higher levels in McR than all other lines, including its yellow
counterpart (Welch's t-test, t=4.208 df=4, p=0.0136). Both isomers of antheraxanthin had a
significant increase in MvY (Welch's t-test, t=7.240 df=7, p=0.0002 for antheraxanthin1,
t=9.391 df=7, p= 0.0001 for antheraxanthin2). In the last three steps of the pathway, a consistent
trend is observed in which both yellow-flowered morphs accumulate significantly higher levels
of violaxanthin (Welch's t-test, t=10.87 df=4, p=0.0004 for MvY, t=6.299 df=6, p=0.0007 for
McY), neoxanthin (Welch's t-test, t=3.563 df=7, p=0.0092 for MvY, t=4.108 df=7, p=0.0045
for McY) and mimulaxanthin (Welch's t-test, t=8.517 df=7, p= 0.0001 for MvY, t=10.53 df=5,
p=0.0001 for McY) when compared to their red conspecifics, suggesting convergence in the
final steps of biosynthesis and accumulation of carotenoids in both yellow morphs.

Carotenoid biosynthesis genes—Following the evidence of convergence in carotenoid
production, we next characterized the expression of genes involved in carotenoid biosynthesis.
No significant differences were found in the initial steps of carotenoid biosynthesis in *M.
verbenaceus*, whereas in *M. cardinalis* there is a significant difference in expression of *phytoene
synthase-1 (PSY1)*, (Welch's t-test, t=8.161 df=3, p=0.0038) and *phytoene synthase
(PDS)*(Welch's t-test, t=6.032 df=3, p=0.0091) between McR and McY. In the later steps of
biosynthesis, both species show a significant increase in one homolog each of *β-carotene
hydroxylase (BCH)* (Welch's t-test, t=4.520 df=3, p=0.0202 for MvY, t=5.184 df=2, p=0.0353
for McY) and *zeaxanthin epoxidase (ZEP)* (Welch's t-test, t=7.287 df=2, p=0.01 for MvY,

t=8.692 df=2, p=0.0130 for McY) (Sun et al., 1996; Hirschberg, 2001) which reflects the
convergent increase in accumulation of violaxanthin, neoxanthin and mimulaxanthin in both
yellow morphs (Figure 3B&C, Table S10). Overall, these results show that the two yellow
morphs are convergently increasing the production of the same carotenoid pigments by
upregulating the same biosynthesis genes across both species.

Transcriptome and whole genome variation—In order to better characterize whether
both novel yellow morphs experienced similar genomic patterns of mutation accumulation, we
characterized the number of within-species variants on each chromosome. All statistical analyses
for the variant counts by chromosome window show the trends in variant accumulation are
significantly different between species (Figure 4, Table S11). This suggests a lack of overall
convergence in variant accumulation by chromosome region in the two red-yellow transitions.

Following from the work of (Beardsley et al., 2003), we extracted ITS sequences for *M.*
cardinalis and *M. verbenaceus* from NCBI in order to visualize the phylogenetic relationship
between these morphs using *Mimulus guttatus* as an outgroup (Song et al., 2012). The resulting
tree demonstrates that the yellow morphs are indeed sister to their red counterparts for each of
the three ITS sequences (Figure S1B). Moreover, it demonstrates how recent the emergence of
these morphs is compared to the section *Erythranthe-Simiolus* (*M. guttatus*) split. These analyses
confirm that yellow morphs have arisen independently and very recently from within separate
ancestrally red-flowered species. When viewing whole transcriptome expression patterns by
NMDS (Figure S13), the morphs show clear species separation through MDS1 with distinction
of colour morphs apparent through MDS2. Coupled with different patterns of genomic variation
(Fig. 4), the color shifts are unlikely to be the product of introgression from one yellow morph
into the other, in contrast to historical color locus introgression between multiple red species and
morphs in *Mimulus* section *Diplacus* (Stankowski & Streisfeld 2015, Short & Streisfeld 2023).

Additional floral traits– To understand if recent shifts in floral color were accompanied by
changes in additional floral traits (which could represent steps of smaller effect in an adaptive
walk toward adaptation to a novel pollinator), we characterized floral scent, morphology, and
nectar rewards for all four lines.

Floral scent—*Mimulus verbenaceus* emitted considerable amounts of floral scent, while
M. cardinalis produced very little scent (Figure 5A). Scent composition was strongly different
between morphs within species (Figure 5B, 5C). We identified 39 compounds in both morphs of
M. verbenaceus, compared to ten compounds across both morphs of *M. cardinalis* (Table 2).
Mimulus verbenaceus produced a high diversity of terpenoids (26 of 39 total compounds) and
volatile composition was overall similar across color morphs (Figure 5D), though terpenoid
alcohols were mostly absent from red *M. verbenaceus*. Additionally, MvY had significantly
higher total emission rates (2.32-fold) compared to MvR (Welch's t-test, t = 8.064, df = 22.214,
p < 0.0001). In contrast, McY and McR produced markedly different bouquets of floral VOCs

(Figure 5E), which they emitted overall at different, relatively low levels, with yellow morph

total emissions approximately 1.68-fold higher than red morph emissions (Welch's t-test, t =

3.738, df = 20.904, p = 0.002). None of the compounds (including unknowns) were found in

common between the two species.

Floral scent biosynthesis genes – Monoterpenoid synthase genes responsible for the

biosynthesis of (*E*)-β-ocimene, limonene, and β-myrcene have been identified in section

Erythranthe (Byers et al. 2014b, Peng et al. 2017). One already characterized *OCIMENE*

SYNTHASE (OS) and five putative bifunctional *LIMONENE-MYRCENE SYNTHASEs (LMS)*

were identified in the genome of MvR based on homology to published *M. lewisii* sequences

(Byers et al. 2014b). *OS* was expressed only in *M. cardinalis,* where it is known to be non-

functional in McR (=CE10) (Figure S14), and no morph produces (*E*)-β-ocimene. The five

different *LMS* candidates were expressed to different degrees (Figure S14), despite limonene's

absence from all four. Although both morphs of *M. verbenaceus* produce β-myrcene, the lack of

concurrent limonene production indicates that either a different gene is responsible for β-

myrcene production or an *LMS* mutation has removed limonene production in *M. verbenaceus*.

Neither product of *LMS* is produced in *M. cardinalis*, although all five *LMS* candidates are

expressed. These putative *LMS* genes may be responsible for the production of other terpenes, or

may have become pseudogenes, as previously found in McR (=CE10) *LMS* (Byers et al. 2014b).

Floral morphology and nectar rewards—We measured floral morphology traits

involved in pollinator attraction (floral display height and width), mechanical fit (corolla tube

length, corolla opening height and width, and herkogamy) and nectar reward quality (volume and

percent sugar). Morphological traits varied significantly among species (MANOVA:

approximate $F_{1,97}= 81.628$, p < 0.0001) and between lines nested within species (MANOVA:

approximate $F_{2,97}= 18.613$, p < 0.0001). NMDS ordination of morphological traits revealed

overall separation between the two species, but little separation between MvR and MvY, in

contrast to clear separation between McR and McY (Fig. 6A).

All individual morphological traits varied significantly among species (p =< 0.001; Table

S3), with the exception of herkogamy (p = 0.1) (Figure S4). Corolla length varied among morphs

nested within species ($\chi^2_{2,\,117}= 53.99$, p < 0.0001), with MvY having significantly longer

corollas than MvR, while McY had significantly shorter corolla tubes than McR (Fig. 6B).

Opening height and width both varied significantly between morphs within species (height: $\chi^2_{2,}$

$_{98}= 65.635$, p < 0.0001; width: $\chi^2_{2,\,98}= 57.03$, p < 0.0001). For both species, yellow-flowered

lines had significantly larger openings than their conspecific red-flowered lines in both opening

height (Fig. 6C) and width (Figure S4). Display height varied significantly among lines within

species ($\chi^2_{2,\,99}= 58.16$, p < 0.0001; Figure S4), with MvY significantly greater then MvR, the

only significant pairwise species comparison. Display width varied significantly among lines

within species ($\chi^2_{2,\,99} = 11.29$, p = 0.004; Figure S4); however, no pairwise comparisons were

significantly different. Herkogamy (within-flower stigma-anther separation) did not vary

between species but did vary significantly among lines nested within species ($\chi^2_{2, 63}$= 116.34, p < 0.0001), with both yellow flowered lines displaying greater herkogamy than their red-flowered conspecifics (Fig. 6D).

Nectar volume per flower varied between species ($\chi^2_{1, 93}$= 48.94, p < 0.0001) but not between lines within species ($\chi^2_{2, 93}$= 2.28, p = 0.32; Figure S4). Percent sugar varied between species ($\chi^2_{1, 93}$= 74.25, p < 0.0001) and among lines ($\chi^2_{2, 93}$= 17.38, p = 0.0002), with the only significant pairwise comparison being in *M. verbenaceus*, where MvY had a higher sugar concentration than MvR (Figure S4).

Floral trait category correlation analysis – Floral traits may be correlated within trait categories, suggesting category-specific floral integration. Distributions of the absolute value of intra-category Pearson correlation coefficients (PCCs) were higher in *M. verbenaceus* than *M. cardinalis* for volatile (F_3 = 7.330, p = 7x10^{-5}) and nectar trait categories (Figure S5). No differences were seen between species for color (F_3 = 1.727, p = 0.16) or morphology (F_3 = 0.866, p = 0.46) trait categories, indicating that these are equally integrated between the two species. As terpenoid volatiles share the same biosynthetic precursors as carotenoid pigments (McGarvey and Croteau 1995), we also calculated correlations between total terpenoid emission and total carotenoids, with species grouped by color morph. These were significantly negatively correlated in the two grouped red lines (R = -0.866, p = 0.0012), suggesting a biochemical tradeoff, but were not significantly correlated in yellow lines (R = 0.111, p = 0.76), despite the higher production of both in yellow morphs.

Individual floral trait correlation analysis - Floral traits may also be individually correlated, even across trait categories. Traits were generally more strongly correlated in M. verbenaceus (Figures S6-S7) than in M. cardinalis (Figures S8-S9), which demonstrated very little inter-trait correlation, even within trait categories. There were 34 traits in McY yielding 544 pairwise comparisons, of which 34 (6.25%) were significantly correlated. McR had 32 traits and 496 comparisons, of which 19 (3.83%) were significant. MvY had 63 traits and 1953 comparisons, of which 288 (14.75%) were significant. MvR had 64 traits and 2016 comparisons, of which 317 (15.72%) were significant.

Pollinator perception and preference

Perception of color and scent—We visualized how bees would perceive the four lines' corolla colors by plotting central petal lobe reflectance in a trichromatic model of honeybee visual space (Figure 7A, Chittka 1992), which revealed that both red-flowered lines overlapped with each other and showed low contrast from a green background. MvY and McY appear distinct from both from each other and their red conspecifics, and also fall further outward on the axes, indicating greater contrast with a green background compared to the red-flowered morphs.

To test scent perception, we performed electroantennography (EAG) to assess the physiological responses of *Bombus terrestris* ssp. *audax* (hereafter *B. terrestris*), a novel

bumblebee pollinator with broad floral preferences (Rasmont et al. 2008) and similar size to native bumblebees observed on these species (Vickery and Vickery, 1992; Vickery, 1995; Cueva del Castillo and Fairbairn, 2012), to floral scent of all morphs and several controls. Bumblebees' electroantennographic responses varied significantly among stimuli ($\chi^2 = 78.28$, df = 6, p < 0.0001), and all four floral scent extracts elicited a significantly greater response than the air-only negative control (Figure 7B, Table S4), suggesting these floral scents can be perceived by bumblebee pollinators. However, antennal responses to floral scent extracts were only significantly greater than to extraction solvent alone for the two *M. cardinalis* morphs (McR t=3.32 df=48.4 p = 0.03; McY t=5.13 df=48.4 p = 0.0001), were marginally greater for MvY (t=3.03 df=48.5 p = 0.06), and were not significantly different for MvR (t=1.68 df=48.4 p= 0.63). Additionally, only one pairwise comparison among the different floral scents was significantly different (McY response was significantly higher than MvR, t=-3.45 df=48.4 p = 0.02). Thus, while bumblebees could perceive the floral scents of all floral morphs, we did not find evidence that morphs were likely to vary in conspicuousness to bees based on their floral scent profiles alone. This is despite the overall much higher total volatile emission in *M. verbenaceus*, suggesting that total volatile emission may only need to reach a certain level for antennal perception. To explore the response of other potential novel pollinators to floral scent, we performed equivalent EAG experiments with *Manduca sexta* hawkmoths. These revealed similar patterns to those of bumblebees, with moths displaying significantly higher antennal responses to scents of all floral morphs compared to controls, but no significant variation in response among floral morphs (Figure S10, Table S5).

Pollinator preference: bumblebee choice— Finally, we tested behavioral responses of novel bumblebee pollinators to yellow and red morphs in controlled within-species comparisons. When ten naïve *Bombus terrestris* workers were offered equal numbers of red and yellow *M. verbenaceus* flowers *M. verbenaceus* in a single-bee experimental array, they made a total of 89 floral visits (Figure S6). Fifty-nine (66.3%) of these visits were to MvY and 30 (33.7%) were to MvR, revealing a significant preference for the yellow-flowered morph (χ^2_1= 9.45, p = 0.002). For *M. cardinalis*, bees made a total of 150 visits over ten trials, visiting McY significantly more frequently, with 99 visits (66%) to McY and 51 visits (34%) to McR (χ^2_1= 15.36, p < 0.0001). Despite this clear preference for yellow in total number of choices across both species, bees were not more likely to choose a yellow flower for their first (naïve) choice (in both species, six first choices were to yellow and four to red; χ^2_1= 0.4, p = 0.53).

Bumblebees preferred yellow flowers of *M. verbenaceus* over red in metrics of total visit number (χ^2_1= 9.44, p = 0.002; Fig. 7C) and total handling time (χ^2_1= 11.74, p = 0.0006), but only marginal evidence for number of flowers probed (χ^2_1= 3.65, p = 0.056; Fig. 7D). For *M. cardinalis*, bees significantly preferred yellow flowers in metrics of total visit number (χ^2_1= 15.125, p = 0.0001), number of flowers probed (χ^2_1= 10.07, p = 0.002), and total handling time (χ^2_1= 74.07, p < 0.0001). Worker bees did not always closely contact the anthers and stigma of flowers while nectaring (Fig. 7E) and were observed to have difficulty handling flowers and

391 accessing floral rewards, particularly in the longer-tubed MvY, suggesting incomplete
392 morphological adaptation to bumblebee pollination. Bees commonly crawled around the back of
393 corollas in an attempt to nectar rob or check for existing robbing holes. When bees made
394 persistent efforts to enter corollas to reach nectar at the base of floral tubes this often resulted in
395 considerable damage to flowers, including bruising and ripping of corollas (Fig. 7E).
396
397 **DISCUSSION**
398
We characterized two yellow-flowered morphs, established as stable populations in two
otherwise red-flowered hummingbird-pollinated species, *Mimulus verbenaceus* and *M.*
cardinalis, across a suite of floral traits important to pollinator attraction and fit and examined
genetic differences underlying these repeated color transitions. We confirmed the previously
hypothesized loss of anthocyanins in petal lobes (Vickery, 1995) and involvement of the gene
PELAN in these shifts (Yuan et al., 2014). Surprisingly, both yellow morphs *also* increase their
production of the same carotenoid pigments using convergent upregulation of the same
carotenoid biosynthesis genes via independent genetic changes. These recent, intraspecific
transitions in color were accompanied by changes in additional traits related to pollinator
attraction and fit. Both yellow forms increased overall floral volatile emissions and had wider,
less restrictive corolla openings and greater herkogamy. Nectar rewards were largely similar
across color morphs within species. Bees were more likely to visually perceive the yellow
flowers of both species and could detect the floral scent of all floral morphs. Collectively, these
floral trait differences resulted in a clear preference of naïve bumblebees for yellow morphs in
both species, despite difficulty handling flowers, which likely reflects the incipient nature of this
intraspecific pollinator shift. Overall, this study reveals that repeated transitions in floral color
within two closely related hummingbird-pollinated species may follow largely convergent
adaptive walks across a suite of floral traits in an incipient shift to bumblebee pollination, which
strongly impacts the potential for reproductive isolation and adaptive evolution.
Convergence in floral pigments and their genetic basis– We found that shades of the
two yellow morphs result not only from a near lack of anthocyanins as expected (Vickery and
Vickery, 1992; Yuan et al., 2014), but also from a simultaneous increase in carotenoid pigments.
The yellow morphs in both species have independently increased production of the same three
carotenoid pigments (violaxanthin, neoxanthin, and mimulaxanthin) via upregulation of the same
two carotenoid biosynthesis genes (*BCH* and *ZEP*), suggesting convergence in these independent
color shifts. Variation in anthocyanin accumulation was reflected in the expression patterns of
known anthocyanin regulators. *PELAN* is known to be responsible for anthocyanin
accumulation in petal lobes of McR (CE10) and is deleted from the genome of McY (SM) (Yuan
et al., 2014). The high expression of this locus in MvY is surprising, given the near absence of
anthocyanins in this line. However, studies in *Lotus* (Yoshida et al., 2010), *Fagopyrum* (Katsu et
al., 2017) and most recently *Lysimachia* (Sánchez-Cabrera et al., 2023) have found that
differential regulation of gene paralogs by the same MYB transcription factors mediated via

promoter sequence polymorphisms can underlie floral color polymorphisms. We propose a
similar case here, where *PELAN* expression in MvY may play a suppressive role in anthocyanin
accumulation, though further study is required.
434   Floral color transitions in multiple related species that involve the same genes have
previously been attributed to adaptive introgression of alleles mediated by gene flow (Short and
Streisfeld, 2023) or to ancestral polymorphism predating speciation (Sánchez-Cabrera et al.,
2023). If either of these processes were responsible here, we would expect to find shared patterns
of mutations in involved loci, which was not the case. In contrast, we find evidence that
differential expression of the same loci results from independent mutations, consistent with
distinct patterns of sequence variation among color morphs throughout their genomes (Figure 4).
Analysis of additional variation found between focal morphs at genomic and transcriptomic
levels is outside  the scope of this study but opens opportunities for further exploration in this
system.
444   ***Bird-to-bee pollinator shifts***—Red floral pigmentation is often associated with
hummingbird pollination (Grant, 1966; Fægri and van der Pijl, 1979), though recent work
suggests red flowers may function less to attract hummingbirds (Waser et al., 2018) than to deter
bumblebees (Castellaños et al., 2004). We also note that the UV "bullseye" pattern seen in
yellow morphs may make them more visible to bees, whose eyes have receptors sensitive to UV
wavelengths (Briscoe and Chittka, 2001). Shifts from blue-purple bee-pollinated flowers to red
hummingbird-pollinated flowers are considered strongly directional (Whittall and Hodges, 2007;
Thomson and Wilson, 2008), and reversals from hummingbird to bee pollination are uncommon
(Kay and Grossenbacher, 2022). However, given multiple examples of transitions between red
and yellow flowers (Tripp and Manos, 2008; Nesom and Egger, 2014; Ho and Smith, 2016;
Short and Streisfeld, 2023) and reported associations of bumblebees with yellow over red
flowers (Vickery, 1992; Simonds and Plowright, 2004; Wenzell et al., 2023), perhaps shifts from
red to yellow flowers as here may represent an avenue to re-gain ancestral attraction to
bumblebees or other insects (Whittall and Hodges, 2007, Tripp and Manos, 2008).
458   The potential for these repeated transitions to yellow flowers in *M. verbenaceus* and *M.
cardinalis* to represent shifts to bumblebee pollination is consistent with changes in other floral
traits. Both yellow morphs had increased scent emission, which could reflect selection from
pollinators such as bees that use scent to forage, unlike hummingbirds (Goldsmith and
Goldsmith, 1982; Núñez et al., 2021), although this might require potentially difficult (but
possible: Esfeld et al. 2018) gain-of-function mutations. Both yellow floral forms were
characterized by wider, less restrictive tube openings, which might function to allow easier
access to nectar by bees. This increased access may be further aided in McY by shorter corolla
tubes compared to McR (though the opposite was observed in *M. verbenaceus*, discussed below),
as well as the presence of bee-orienting nectar guides (Free 1970, Leonard & Papaj 2011). This
combination of traits particular to McY could contribute to the significant increase in probed
yellow flowers in this species. Nectar volume did not vary among color morphs in either species
(though in *M. verbenaceus*, yellow flowers had higher sucrose content than red), which is

noteworthy as rewards are important in determining pollinator preference (Kaczorowski et al.,
2005; Cronk and Ojeda, 2008).

In addition to color, floral scent is a key trait involved in pollinator attraction (Raguso,
2008; Koski, 2020; Byers, 2021). Based on our EAG analysis, workers of *Bombus terrestris* are
capable of detecting the floral scents of all morphs, though we did not see a greater response
from bee antennae to either yellow morph. It is possible that the presented floral volatile stimuli
are equivalently ecologically relevant to bumblebees, resulting in no response difference despite
significant volatile composition differences, as seen in other studies (Morawo et al., 2017). It is
also possible that our EAG analysis was complicated by the inadvertently low concentration of
the positive control stimulus. We note that our scent profile of McR (= CE10) differs from that
described in Byers et al. (2014a), which is likely a result of our more stringent filtering of
compounds and a more sensitive GCMS system.

Overall, the most compelling evidence for a potential shift to bee pollination in this
system is the strong preference of the bees themselves for yellow morphs. We found significant
preference by bumblebees for yellow in both species, with a notably consistent 2:1 ratio of all
visits to yellow versus red flowers across species (Figure S11). Nonetheless, during the
experiment, bees were observed to have difficulty handling flowers. In particular, bumblebees
entered the corolla in multiple orientations, often poorly contacting floral organs (Figure 7E),
which would hinder pollen transfer efficiency. Bees were also observed attempting to nectar rob,
and often struggled to fit inside corolla tubes to access nectar, occasionally resulting in severe
corolla damage when bees persisted in trying to enter (Figure 7E, particularly in the longer-tubed
MvY).

Given that these floral morphs represent recent transitions below the species level, we
suggest they represent incipient pollinator shifts which may occupy a fitness valley between
peaks of floral phenotypes optimally adapted to either hummingbird or bee pollination. Based on
the strong increase in bumblebee attraction observed in both yellow morphs, we hypothesize that
convergent shifts in floral color (loss of anthocyanins and increased carotenoids) represent
mutations of large effect at the beginning of an ongoing adaptive walk with strong effects on
attraction of a novel bumblebee pollinator, while additional traits likely reflect smaller-effect
steps contributing further to attraction (increased scent emission), fit (wider corolla openings),
and positioning (nectar guides in McY) of bee pollinators. We speculate that additional changes
in floral traits characteristic of bumblebee-pollinated flowers, such as landing platforms and less
tubular corollas (characteristic of *M. lewisii*, the bee-pollinated member of section *Erythranthe*
sister to *M. cardinalis*) would represent further "steps" needed to complete the adaptive walk to
an optimal bee-pollinated phenotype. Future research into this putative pollinator shift should
extend to natural populations by assessing allelic variation and potential signatures of selection
of identified loci, along with characterizing variation in floral traits beyond the inbred reference
lines used here.

With this in mind, we acknowledge that pollinators besides hummingbirds and
bumblebees may pollinate and exert selection on these flowers in natural habitats. In particular,

we hypothesize that the yellow morph of *M. verbenaceus* could reflect an incipient shift to hawkmoth pollination, given its pale coloration, long corolla tubes, increased floral scent emission, and lack of nectar guides (Riffell and Alarcón, 2013). While speculative, the potential for hawkmoth pollination is consistent with our EAG results for the hawkmoth *Manduca sexta*, as hawkmoths are capable of detecting scent profiles of all morphs (Figure S6), and *Manduca sexta* preferred yellow flowers over red in controlled experiments with *Mimulus* (Byers and Bradshaw, 2021). Shifts from hummingbird to moth pollination have occurred repeatedly in other systems (e.g. *Aquilegia*, Whittall and Hodges, 2007), and may reflect that many hawkmoths, including the regionally common *Hyles lineata*, are generalists (Alarcón et al., 2008; Wenzell et al., 2023), and are often underreported in diurnal-only pollinator observations (e.g. Vickery 1992). Alternatively, observed floral trait variation could reflect selection from other biotic or abiotic drivers (Strauss and Whittall, 2006) or drift, which is expected to be stronger at range edges (Nadeau and Urban, 2019) where both yellow morphs occur. While other explanations of floral trait variation warrant further study, our reported twofold preference among bumblebees for yellow morphs in both species provide convincing evidence that these floral traits are likely to be subject to pollinator-mediated selection. Future research into the native pollinators and other environmental conditions of yellow morphs of both species *in situ*, along with nearby red-flowered populations, is needed to more fully investigate the eco-evolutionary conditions which may favor a pollinator shift within these species.

CONCLUSIONS

In this study, we document that two intraspecific floral color transitions result from convergent patterns of carotenoid accumulation, in addition to loss of anthocyanins, both of which involve independent mutations in a shared set of pigment-related genes. These changes in floral color are accompanied by variation in a suite of additional floral traits, with implications for the attraction, mechanical fit, and reward access of a novel pollinator. The novel yellow morphs in otherwise red-flowered, hummingbird-pollinated species were strongly preferred by naïve bumblebees in experimental arrays, and our results suggest this preference reflects multiple floral traits acting in concert to influence pollinator foraging decisions. We found that while increased preference by bumblebees for yellow morphs may provide the potential for a shift in pollinators, this shift appears to be incomplete, as even preferred phenotypes appear ill-suited to these pollinators. This could suggest that current phenotypes may represent an ongoing adaptive walk between peaks in a fitness landscape. Ultimately, this study reveals that suites of floral traits act in concert to shape the attraction, fit, and preference of pollinators, and while changes in attraction traits of large effect like color may initiate the process of a pollinator shift, additional changes to smaller-effect fit traits must follow to ensure efficient pollen transfer and reproductive isolation. Thus, this study sheds light on the order in which floral traits may evolve in an incipient pollinator shift, their genetic basis, and their impact on the behavior of a novel pollinator, with implications for early stages of pollinator-mediated speciation and our understanding of adaptation in complex traits.

551
METHODS
  Plant material—The yellow-flowered color morph of *M. verbenaceus* (MVYL, hereafter MvY) was originally collected from a population located at Vassey's Paradise in the Grand Canyon, Arizona, USA (exact coordinates unknown), near the northwestern extent of the species range (Vickery and Vickery, 1992), about 100 miles north of the source population of the red-flowered *M. verbenaceus* (MVBL, hereafter MvR; Figure S1A). MvR was collected along the West Fork Trail in Sedona, AZ, USA ($34.9883°N$, $111.7485°W$). Yellow-flowered morphs of *M. cardinalis* have been characterized at two populations: one at the northern extent of the species range in the Siskiyou Mountains of Jackson County, Oregon, USA (known as SM, hereafter McY), and one from the southern range extent in Cedros Island, Baja California. For this study, we characterize the yellow morph from the northern population (SM, exact coordinates unknown), though we note that previous work with pollinators include the southern Cedros Island population (Vickery and Vickery, 1992; Vickery, 1995). The red-flowered reference line of *M. cardinalis* (CE10, hereafter McR) was collected near South Fork Tuolumne River, Tuolumne County, California, USA ($37.817°N$, $119.867°W$). Plants used in experiments represent reference lines inbred for at least ten generations and were grown from seed at the John Innes Centre, Norwich, UK.

  Prior to germinating, seeds were sterilized in dilute (1.2%) sodium hypochlorite solution for 5-10 minutes, then rinsed three times with sterilized distilled water. Seeds were then sown on MS growth media in Petri dishes and placed in a controlled environment room (CER) under 14-hour day lengths at $20°C$ day/$19°C$ night. Following germination, seedlings were transferred to cereal mix to establish in CERs before being moved to the glasshouses at John Innes Centre.

  All floral phenotypic traits were measured on freshly opened flowers (i.e. opened that day), including tissue collected for pigment and RNAseq analyses.

  Floral color: reflectance and pigments—Floral color was quantified by measuring reflectance of fresh corolla tissues (collected immediately prior from plants growing in the glasshouse) using a reflectance spectrometer (FLAME-S-UV-VIS-ES Assembly, 200-850nm, Ocean Insight), with light source from a pulsed xenon lamp (220 Hz, 220-750 nm, Ocean Insight) on four areas of the corolla: the upper petal lobe, the central (bottom) petal lobe, the lower side petal lobe, and the throat of the corolla tube/nectar guide. The reflectance spectrometer collected readings every 5ms and averaged 25 scans per reading (Ocean View spectroscopy software, Ocean Insight). Readings were standardized with an absolute white color reference (Certified Reflectance Standard, Labsphere) and electric dark was used as the black standard. Three readings were taken from each tissue of each flower measured, and three flowers were measured per individual plant, with at least 5-15 plants characterized per line. Reflectance curves at wavelengths from 300-700nm were visualized for all four tissues. All three petal lobe tissues showed similar reflectance curves (Figure S2), so we present and analyze only the central petal lobe tissue hereafter, as this petal lobe is forward facing (not strongly reflexed) and thus visible to pollinators approaching the front of the flowers.

Floral color (reflectance) of the central petal lobe was compared among lines using a principal component analysis (PCA) using R v.4.2.2 (R Core Team 2022), package psych version - 2.2.5(Revelle, 2023) to allow for the varimax rotation, which prioritizes loading each trait (i.e., wavelength) onto only one PC axis), based on the intensity of reflectance at a given wavelength (based on intervals of every ten nanometers from 300-700nm, which captures wavelengths relevant to the vision of most pollinators). First, a PCA was run based on intensity values at all 41 wavelengths; then, only factors with an eigenvalue > 1 were used in the PCA, which resulted in a final PCA of 3 factors. To visualize whether these flowers reflect in the ultraviolet (UV) range (including potential patterning), which is important for bee vision, we photographed flowers of 1-3 individuals of all four color morphs under UV light using a UV-sensitive camera (Nikon D610, converted to a Full Spectrum camera by removing the Kolari Vision UV bandpass filter, with a Micro-Nikkor 105mm lens) in a dark room illuminated by a UV black light.

Characterizing floral pigments— To characterize the biochemical basis of floral color differences, we analyzed the anthocyanin and carotenoid pigments present in fully opened corolla tissues of five individuals (1 flower per individual plant for pigment group) for all four lines. Total anthocyanins were quantified by measuring absorbance at 525 nm and total carotenoids at 450 nm, using a spectrophotometer (DS-11 FX UV-vis-spectrophotometer, Cambridge Bioscience, UK). When initial absorbance readings were above the saturation point of the spectrophotometer, samples were diluted with the addition of extraction buffer (1:1 of 1% solution of hydrochloric acid in methanol for anthocyanins and 4:1 ethyl acetate for carotenoids). Because the size of corollas varied among floral morphs, and thus likely contributed different amounts of tissue, we standardized the absorbance values relative to corolla size by dividing absorbance by the average mass (g) of a fresh corolla for each line (based on an average mass from one corolla from five individual plants), and used these values for analysis (though we note that overall patterns were similar to those using un-standardized absorbance values). Differences among lines (nested in species) in total anthocyanins and total carotenoids were assessed by performing separate ANOVAs (aov() function in R) for each pigment type, followed by post doc Tukey tests (TukeyHSD() function with 95% CI).

Anthocyanins— Modified from (Butelli et al.,2008). Roughly, fully opened whole corollas were snap-frozen in liquid nitrogen and ground to a fine powder, anthocyanins were extracted using a solution of 80%methanol and HCL with samples left to shake overnight. The resulting supernatant was stored in amber vials prior to analysis. The samples were run on an Agilent 1290 Infinity II UHPLC equipped with PDA and 6546 Q-ToF. Separation was on a 100×2.1mm 2.6µ Kinetex EVO column using the following gradient of acetonitrile (solvent B) versus 1% formic acid in water (solvent A), run at 600µL.min-1 and 40°C: 0 min, 2% B; 7.5 min, 30% B; 10 min, 90% B; 10.7 min, 90% B; 10.9 min, 2% B; 15 min, 2% B.   The diode array detector collected individual channels at 350nm (bw 4nm) and 525nm (bw 50nm), and also full spectra from 220-640nm, at 10 Hz.The Q-ToF collected positive electrospray MS

using the JetStream source, together with data-dependent MS2. The MS was calibrated before

use, but also had two lock-masses infused during the runs, at 121.05087 and 922.009798;

components of Agilent's normal ESI calibration mix). Spray chamber conditions were 325°C

gas, $10L.min^{-1}$ drying gas, 20psi nebulizer pressure, 400°C sheath gas, $12L.min^{-1}$ sheath gas,

4000V Vcap (spray voltage) with a nozzle voltage of 1000V. The fragmentor voltage was 180V.

Carotenoids— Modified from (Serino et al., 2009). Roughly, fully opened whole corollas

were snap frozen in liquid nitrogen and ground to a fine powder then carotenoids were extracted

through sequential agitation and centrifugation with the following: NaCl saturated aqueous

solution, hexane, dichloromethane and ethyl acetate. Final extraction product was stored in

amber vials and run on LC/MS. To assist in developing a suitable LC/MS method the following

paper was used (Rivera et al., 2011). The samples were run on an Agilent 1290 Infinity II

UHPLC equipped with PDA and 6546 Q-ToF using APCI.   The column used was an Agilent

Zorbax Eclipse plus C18, 2.1mm X 50mm. 1.8 micron, Rapid Resolution HD with an

Acetonitrile:Methanol mobile phase at a ratio of 70:30. Flow rate was 0.4ml/min for 15 minutes

with a column temperature of 32°C.   Violaxanthin, neoxanthin, zeaxanthin, and

antheraxanthin were confirmed with standards. For mimulaxanthin, quantification was

performed using the response of antheraxanthin.

Transcriptome analysis and comparative gene expression– RNA was extracted from

fully opened whole corollas – including anthers – as was performed for the pigment analyses, in

order for these datasets to be comparable. RNA extraction was performed using

SPECTRUM(TM) PLANT TOTAL RNA KIT from Merck Catalogue #STRN50-1KT and eluted

in water. Three replicates per genotype were sent for sequencing. Library preparation and

sequencing was performed by Novogene using Illumina sequencing technologies. Expression

was quantified using kallisto version 0.46.1 (Liao et al., 2014), with the publicly available cDNA

dataset from *Mimulus verbenaceus* (MvR=MVBL) reference (http://mimubase.org/). Differential

expression data were visualised in Degust (Powell, 2015) with the following parameters; Min

Gene CPM = 10 reads in 3 samples, FDR cut-off =1, abs logFC=1) .

Whole genome analysis– DNA was extracted from leaf tissue of MvYL and SM using

the NUCLEON PHYTOPURE 50 PREPX0.1G Kit (Supplier Catalogue ID: RPN8510) with the

addition of an RNAse treatment using ThermoScientific RNAase T1 Catalogue #EN0541 .

Library preparation and sequencing was performed by Novogene using Illumina sequencing

technologies. All genome and transcriptome data are available on ENA, The accession number is

PRJEB75514.

Reads were aligned to the publicly available *Mimulus verbenaceus* (MvR=MVBL) and

Mimulus cardinalis (McR=CE10) reference genomes (http://mimubase.org/) using bwa version

0.7.17 (Li et al., 2009a) with subsequent processing using samtools version 1.7 (Li et al., 2009b)

and bcftools version 1.10.2 (Li, 2011).

The analysis of genomic differences described in Figure 4 were a result of variant calling

using bcftools against each line's respective red counterpart, following division of the assemblies

into 50kb windows using the bedtools makewindows function. This produced a table of variants
per window, per chromosome (Table S11). Figure 4 depicts this variant count with 2nd order
smoothing (50 neighbors).
The ITS sequences AY575418.1, AY575414.1, AY575439.1 were downloaded from
NCBI, and blasted against the reference sequences of Mimulus verbenaceus (MvR=MVBL) and
Mimulus cardinalis (McR=CE10) reference genomes (http://mimubase.org/). The blast hits of
ITS sequences that were full length and in the chromosomes were used to extract the fasta
sequence from the reference files using samtools faidx command. See Table S11 for fasta
sequences. We used Mafft (version 7.520) with default parameters to align the fasta sequences
(Katoh & Standley, 2013). The tree was built using fasttree (version 2.1.9) with default
parameters (Price et al., 2010). The tree was visualized using Interactive Tree Of Life (iTOL)
(Letunic et al., 2021).
Floral scent— We characterized floral volatile organic compounds (VOCs) for 3
technical replicates each of at least 5-10 plants per line, from plants growing in the glasshouse at
the John Innes Centre. Floral VOCs were collected using dynamic headspace collection as
follows. Pairs of freshly cut fully open flowers were placed in 50 mL glass beakers containing
approximately 40 mL of sterile ddH$_2$O, which were placed in oven roasting bags (Sainbury's,
UK), sealed and connected to scent traps with aluminum twist ties. Scent traps were composed of
modified glass Pasteur pipettes filled with 100mg Porapak Q (Merck/Sigma), contained within
pea-sized amounts of glass wool on either side, and were connected via plastic tubing to Spectrex
PAS-500 volatile pumps (Merck/Sigma), which were run at a flow rate of 100 mL/min for 24
hours to capture any circadian variation in emissions. Floral VOCs were then eluted from scent
traps using 600 uL of extraction solvent (10% acetone in HPLC-grade hexane) into 2mL screw-
top amber glass vials, which were stored at -20°C in a dedicated scent freezer before
concentration and GCMS analysis.
Compound identification, quantification, and analysis— Sample aliquots of 150ul were
first concentrated to 50ul prior to injection of 3ul into an Agilent GC-MS system (7890B GC
with 5977A/B MS) using a Gerstel MPS autosampler system with a splitless inlet held at 250C.
The oven temperature was held at 50C for 4 minutes, then raised at 5C/minute to 230C, where it
was held for 4 minutes. The column was a Phenomenex 7HG-6015-02-GGA (35m x 250um x
0.1um) and the carrier gas was helium at a flow rate of 1.0603 ml/minute. The MS was run in
scan mode for ion masses between 50 and 600, with the MS source held at 230C.
Data files were first analyzed using Agilent Unknowns software (v.10.1) to integrate
peaks using the default integrator settings and give the top three tentative NIST library
identifications, then processed further via custom shell and Perl scripts. We first excluded peaks
with an area under the curve of 1x10^5 due to the extremely sensitive nature of the default
Unknowns integrator, as well as excluding peaks that eluted after 30 minutes, which are less
likely to be volatile pollinator attractants due to their higher molecular weight. Peaks with the
same tentative library identification as a peak within 0.1 minutes in the blank sample and with
less than a 5-fold higher area in the sample of interest were excluded, as were peaks where all

three library hits contained silica or phthalate contaminants. Following this, remaining uncertain
peaks were compared to the nearest blank peaks (within 0.1 minute) by visual comparison of
spectra and retained or discarded as above. To further verify the identity of retained peaks, an
alkane ladder (Merck, 49452-U) was run at the same time as the samples and the Kovats
Retention Index of each peak was calculated and compared with published Kovats indices
available via NIST. Compounds with an available authentic reference standard available for
purchase (the majority of compounds in this study) were then verified by injection on our GCMS
system using the method above and floral sample integrated areas converted to ng/flower/hour
using a concentration curve run with each standard. Compounds without a standard available, but
with matching Kovats Retention Indices to the literature, had the areas converted using a
similarly structured reference compound. Compounds with no matching standard or Kovats
Retention Index are listed in Table 2 as "Unknowns" and quantified using a similarly structured
reference compound to the "best guess" of the NIST library. Scent data were visualized in R
v.4.2.2 (R Core Team 2022) using the vegan package v.2.6-4 (Oksanen et al., 2022) for NMDS
plotting and Welch's t-tests for comparisons between morphs within species for both individual
volatile and total volatile emissions.

Floral morphology and nectar—Approximately three fully open flowers each of 5-15
individuals of each line were measured using 150mm digital calipers (Linear Tools, RS
Components) for each morphological floral trait. Measured traits included corolla tube length
(measured from base of corolla to mouth opening), floral display height (vertical spread of petal
lobes as viewed from the front), display width (horizontal spread of petal lobes), tube opening
height (vertical space within mouth of corolla tube), and opening width (horizontal space within
mouth); these traits play important roles in pollinator attraction/visual signals and access to
nectar rewards. Herkogamy, the distance between the anthers and stigma of an individual flower,
was assessed by measuring the length of the stamens (from the receptacle to the anthers) and the
length of the pistil (from the receptacle to the stigma) of three flowers from five individual plants
of each line. Herkogamy was calculated as the difference between the length of the pistil and the
length of the anthers (allowing for a possible negative herkogamy value if stamens were longer
than the pistil).

For nectar measurement, at least three flowers per plant (5-15 plants per line) were
sampled on the first day of opening, to ensure consistent time for nectar production across lines.
Flowers were collected from plants growing in the glasshouse and immediately sampled for
nectar volume and sucrose concentration. Nectar was collected from the base of corollas into
microcapillary tubes (Cat. No. 2930210, 1.55 mm external diameter, Paul Marienfeld GmbH &
Co. KG, Germany), and the height of nectar in tubes was measured in mm using digital calipers,
which was then converted into uL (based on capillary tube dimensions (1.15mm internal
diameter and 100mm length, via manufacturer's specifications) and the geometric formula for
volume of a cylinder [height of nectar in mm/100*103.87=h/100*(π*0.5752)]. Concentration of
sucrose was then measured using a nectar refractometer (0 to 50% (BRIX) sugar, Bellingham
and Stanley UK Ltd.).

Multivariate differences in morphological traits (corolla length, display height and width, and opening height and width) were assessed among lines using a nonmetric multidimensional scaling (NMDS) ordination using Bray distances in R v.4.2.2 (R Core Team 2022) package vegan (Oksanen et al 2022), followed by a MANOVA of these traits by species and line nested within species. Differences among lines for each floral trait were assessed using a linear mixed model (the lmer() function in R v.4.2.2 (R Core Team 2022) package lme4 (Bates et al., 2015)) with line nested within species and individual plant included as a random effect, followed by Type II Wald chi-square tests using the Anova() function in R v.4.2.2 (R Core Team 2022) package car version 3.0-10 (Fox et al., 2020).

Floral trait integration analysis – Measured floral traits were divided into four categories: volatiles, color (including both individual pigments and reflectance principal components), morphology, and nectar. Within each category, all pairwise combination Pearson correlation coefficients (PCCs) were calculated between each individual trait pair. Absolute PCC values were compared within each category across lines using Type II Wald chi-square tests using the Anova() function in R v.4.2.2 (R Core Team 2022) package car (Fox et al., 2020) with Tukey's Honestly Significant Difference tests applied for volatiles, the only category where the Anova result was significant. No statistical test was performed for nectar, where only one PCC was calculated per line from the two measured traits. Overall correlations between total terpenoid emission and total carotenoid production were similarly analyzed via calculation of the PCC, with species grouped by color (red versus yellow).

For all-trait correlational analyses, Pearson correlation coefficients were calculated for all individual floral traits against one another using the rcorr() function in R v.4.2.2 (R Core Team 2022) package Hmisc (Harrell 2024). Principal components for reflectance spectrophotometry data were used in lieu of individual wavelength measurements for this analysis.

Pollinator perception of color and scent—To characterize how floral colors may be perceived by bee pollinators, we plotted the reflectance data of lower central petal lobes on trichromatic models of bee (*Apis mellifera*) visual systems using R v.4.2.2 (R Core Team 2022) R package pavo version 2.7.1 (Maia et al., 2019), which estimates level of contrast of a color signal against a vegetative (green) background.

To test whether the floral scents of different color morphs elicit different electrophysiological responses in bumblebees, we performed whole-extract electroantennography (EAG). We exposed antennae from nine naïve lab-reared *Bombus terrestris* ssp. *audax* individuals (see below for hive maintenance conditions; one antenna per individual) to floral scent extractions (concentrated down to half volume) of both yellow-flowered morphs (MvY and McY) and their conspecific red-flowered reference lines (MvR and McR respectively) and measured the electrophysiological response to these extracts, in addition to a positive control floral scent compound (0.5 ng/ul phenylacetaldehyde (PAA), a compound known to elicit a response to *Bombus terrestris* (Knauer and Schiestl, 2015)), a negative control

(a puff of air through a scent cartridge with no VOCs) and the extraction solution in which the
floral VOC blends were suspended (10% acetone in hexane).
Using micro-dissection scissors, single antennae were excised from live bees, which were
first chilled for 20 minutes at 4C, and the tip (last 1.5 flagellomeres) of the antenna was removed
using a micro-dissection scalpel under a dissecting microscope. The antenna was then placed
onto an antennal fork using electrode gel and placed into an MP-15 micromanipulator-electrode
holder mounted to a magnetic base plate. The antenna was allowed to equilibrate for 10 minutes
prior to recording commencing. Recording signals were sent to an IDAC-2 and from there to a
computer, where they were recorded using GcEad/2014 software. Stimuli were delivered via a
CS-55 stimulus controller. Continuous air flowed over the antenna, with pulses of the same air
flow strength controlled via a foot pedal. All EAG equipment and gel were supplied by
Ockenfels SYNTECH GmbH, Buchenback, Germany.
The order of scent stimulants was randomized (1 round of 6 exposures each, comprised
of 3 pulses 0.5 seconds apart, delivered every thirty seconds), apart from the positive control
(PAA) which was delivered in three rounds of 4 exposures and was always the first, fourth, and
final stimulant delivered to each antenna sample. This was done because antennae decrease in
responsiveness over time, so the positive control signal was used to normalize antennal responses
to other stimuli throughout the course of the EAG assay, and the final corrected amplitude value
was then scaled to the initial PAA response (following Byers et al., 2020). Additionally, the first
response to a novel stimulus (scent) often results in a disproportionately large peak due to
solvent blowoff, so the first response to each stimulus was removed as an outlier for each trial.
This resulted in a total of 1,230 recorded responses to stimuli (across nine trials using one
antenna from each of nine bees), which were measured and recorded using the software Gc-Ead
2014 v 1.2.5 (2014-05-03; SYNTECH, https://www.ockenfels-syntech.com/).
We tested whether the scaled and corrected amplitude of the antennal response varied
among stimuli using a linear mixed model (lmer()) with trial (individual antenna) included as a
random effect, along with a random effect term of stimuli nested within trial to account for the
repeated exposures of each stimuli to a single individual antenna. Pairwise comparisons among
stimuli were assessed using the function lsmeans() with Sattherwaite degrees of freedom and
lmer test limit set to 6000 in R v.4.2.2 (R Core Team 2022) package lmerTest version 0.9-38
(Kuznetsova et al., 2017).
To investigate whether other functional groups of novel pollinators may respond to floral
signals in these species, identical electroantennography experiments were also conducted with
seven virgin female *Manduca sexta* hawkmoths, which were reared on a standard artificial diet
(Frontiers Scientific Services, Newark, DE, USA). Phenylacetlaldehyde was again used as a
control, as it is known to be detectable by *Manduca sexta* (Fraser et al., 2003). Details of
antennal preparation (although no chilling was used prior to antenna removal), stimulus delivery,
and stimulus identity and statistical analysis were identical to the *Bombus terrestris audax*
experiments.

Bumblebee choice experiment—To assess whether bumblebees exhibited a preference
for visiting one floral color morph over another, we performed a pairwise choice experiment
using naïve *Bombus terrestris* workers. Hives of *Bombus terrestris* ssp. *audax* were sourced from
Agralan, Ltd. (Wiltshire, UK) and were maintained in a controlled environment room (CER) in
the Entomology Department at the John Innes Centre, Norwich, UK, which was kept at 22C
during the day and 20C at night with 14 hour day lengths. Hives were kept in a polycarbonate
cage and were given access to commercial bee pollen (organic raw bee pollen, Sevenhills
Wholefoods, Wakefield, UK) and 25% sucrose (Fischer Scientific) solution. The day (ca. 16:00-
17:00) preceding each experimental trial, sucrose solution was removed from the cage, in order
to motivate workers to forage during the trial the following day. For use in trials, workers were
collected from outside the hive and were kept in individual ventilated 50 mL Falcon tubes, kept
in darkness until use.
Open, receptive flowers of yellow- and red-flowered reference lines were cut from plants
maintained in the glasshouse and were immediately placed into damp floral foam (Cappstan UK)
in 15 mL Falcon tubes. For each within-species pairwise comparison, four flowers of the yellow
morph and 4 flowers of the red morph were randomly arranged in one of two 2x2 grid arrays on
the walls of a 65 x 80 x 78 mm polycarbonate flight cage. Comparisons were made only within-
species, so each trial consisted of a pairwise choice of either the red or yellow morph of *M.*
verbenaceus (MvR versus MvY) or of *M. cardinalis* (McR versus McY) presented in a 50:50
ratio. The background behind the floral arrays was covered in green poster roll paper (Meadow
Green, House of Card & Paper, UK) to simulate a vegetative background. The flight cage was
placed inside a CER kept at 20C during experiments.
For each trial, one naïve worker bee was placed in the flight cage with the randomized
array of flowers and their foraging behavior was recorded. A choice (floral visit) was recorded
any time the bee intentionally contacted the corolla of any flower; if the bee entered the throat of
the corolla tube (i.e., to attempt to access the nectar at the base of the corolla tube), this visit was
also recorded as a probe. The color morph of each choice was recorded, as well as the handling
time (how many seconds the bee handled the flower during each visit/choice). Each trial lasted
30 minutes, after which time, the bee was removed from the flight chamber and euthanized. If a
bee failed to forage after 20 minutes in the chamber, it was removed and euthanized. Flowers
that came in contact with bees during a trial were replaced before the next trials and the
arrangement of floral morphs in each array was re-randomized for each trial. The flight chamber
was cleaned with 70% ethanol between trials of different species to remove any lingering floral
volatiles.
Trials were performed until each species comparison had ten successful trials (i.e., ten
individual bees had made a choice). Preference of bees was assessed by comparing the observed
total number of flower choices made for each color morph to the number of choices expected
under the null hypothesis of no color preference (=50%) using a Chi-square goodness of fit test
(chisq.test() function in R), with a separate test run for each species, with choices pooled across
trials. We also tested for difference among color morphs of the first choice made by each bee

(i.e., the color of the flower each bee contacted first) in the same manner. In addition, because
preference could vary among individual worker bees, we also assessed preference in a second
way using generalized linear mixed models (GLMM) using the glmer() function in R v.4.2.2 (R
Core Team 2022) package lme4 version 1.1-27.1, which included a random effect for individual
bee. These GLMMs used one of the choice metrics (number of visits, number of probes, and
total handling time) as the response variable with floral color morph as the predictor, with a
Poisson error distribution useful for count data, and separate models were performed for each
species comparison.
876

ACKNOWLEDGEMENTS
The authors thank Yao-Wu Yuan for providing seeds for all focal lines, as well as original seed
sources provided by R. Vickery (McY=SM and MvY=MVYL), P. Beardsley (MvR=MVBL),
and H.D. Bradshaw (McR=CE10). We also thank J. Fant, A. Davies, D. Cohen and members of
the Fant lab group for thoughtful feedback on the manuscript, and we thank David Seung for use
of the spectrophotometer in his lab at the John Innes Centre. We would also like to thank
Baldeep Kular for metabolomic support. Finally, we are grateful to the Metabolomics,
Horticultural Services, Informatics, and Entomology Platforms at the John Innes Centre for their
support. This work was funded by start-up funds from the John Innes Centre to KJRPB, and by
the UK Biotechnology and Biological Sciences Research Council Institute Strategic Programmes
(Harnessing Biosynthesis for Sustainable Food and Health (HBio) (grant no. BB/X01097X/1)
and Building Robustness in Crops (BRiC) (grant no. BB/X01102X/1). We also thank anonymous
reviewers and the editor for their comments and suggestions.

DATA AVAILABILITY
Sequence data that support the findings of this study have been deposited with the European
Nucleotide Archive with the accession code PRJEB75514
(https://www.ebi.ac.uk/ena/browser/view/PRJEB75514). Metabolomic data and all other data
have been deposited with figshare (data DOI: XXX) and source data (excluding sequencing and
metabolomic data) are additionally provided with this paper. Unique seed materials used in this
study are available on request from the corresponding author.
898

899 **LITERATURE CITED**

Alarcón, R., Davidowitz, G., and Bronstein, J. L. 2008. Nectar usage in a southern Arizona
hawkmoth community. *Ecological Entomology* 33: 503-509.

Arendt, J., and D. Reznick. 2008. Convergence and parallelism reconsidered: what have we
learned about the genetics of adaptation? *Trends in Ecology & Evolution* 23: 26–32.

Arnold, S. J., M. E. Pfrender, and A. G. Jones. 2001. The adaptive landscape as a
conceptual bridge between micro- and macroevolution. In A. P. Hendry, and M. T.
Kinnison [eds.], Microevolution Rate, Pattern, Process, 9–32. Springer Netherlands,
Dordrecht.

Bates, D., M. Mächler, B. Bolker, and S. Walker. 2015. Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software* 67: 1–48.

Beardsley, P. M., A. Yen, and R. G. Olmstead. 2003. AFLP Phylogeny of *Mimulus* Section *Erythranthe* and the Evolution of Hummingbird Pollination. *Evolution* 57: 1397–1410.

Berardi, A. E., Betancourt Morejón, A. C., and Hopkins, R. 2022. Convergence without divergence in North American red-flowered *Silene*. *Frontiers in Plant Science* 13: 945806.

Berardi, A. E., Esfeld, K., Jäggi, L., Mandel, T., Cannarozzi, G. M., & Kuhlemeier, C. (2021). Complex evolution of novel red floral color in *Petunia*. *The Plant Cell*, 33(7), 2273-2295.

Bradshaw, H. D., Jr., and D. W. Schemske. 2003. Allele substitution at a flower colour locus produces a pollinator shift in monkeyflowers. *Nature* 426: 176–178.

Bradshaw, H. D., Jr., Wilbert, S. M., Otto, K. G., and Schemske, D. W. 1995. Genetic mapping of floral traits associated with reproductive isolation in monkeyflowers (*Mimulus*). *Nature* 376: 762-765.

Briscoe, A. D., and L. Chittka. 2001. The Evolution of Color Vision in Insects. *Annual Review of Entomology* 46: 471–510.

Butelli, E., L. Titta, M. Giorgio, H.-P. Mock, A. Matros, S. Peterek, E. G. W. M. Schijlen, et al. 2008. Enrichment of tomato fruit with health-promoting anthocyanins by expression of select transcription factors. *Nature Biotechnology* 26: 1301–1308.

Byers, K. J. R. P. 2021. "As if they discovered it by the scent": improving our understanding of the chemical ecology, evolution, and genetics of floral scent and its role in pollination. *American Journal of Botany* 108: 729-731.

Byers, K. J. R. P., and H. D. Bradshaw. 2021. Rational Design of a Novel Hawkmoth Pollinator Interaction in *Mimulus* Section *Erythranthe*. *Frontiers in Ecology and Evolution* 9.

Byers, K. J., Xu, S., & Schlüter, P. M. (2017). Molecular mechanisms of adaptation and speciation: why do we need an integrative approach?. *Molecular Ecology*, 26(1), 277-290.

Byers, K. J. R. P., H. D. Bradshaw, Jr., and J. A. Riffell. 2014a. Three floral volatiles contribute to differential pollinator attraction in monkeyflowers (*Mimulus*). *Journal of Experimental Biology* 217: 614–623.

Byers, K. J. R. P., J. P. Vela, F. Peng, J. A. Riffell, and H. D. Bradshaw, Jr. 2014b. Floral volatile alleles can contribute to pollinator-mediated reproductive isolation in monkeyflowers (*Mimulus*). *Plant Journal* 80: 1031-1042.

Byers, K. J. R. P., K. Darragh, J. Musgrove, D. A. Almeida, S. F. Garza, I. A. Warren, P. M. Rastas, et al. 2020. A major locus controls a biologically active pheromone component in *Heliconius melpomene*. *Evolution* 74: 349–364.

Castellanos, M. C., Wilson, P., & Thomson, J. D. (2004). 'Anti-bee' and 'pro-bird' changes during the evolution of hummingbird pollination in Penstemon flowers. *Journal of evolutionary biology* 17(4): 876-885.

Chittka, L. 1992. The colour hexagon: a chromaticity diagram based on photoreceptor excitations as a generalized representation of colour opponency. *Journal of Comparative Physiology A* 170: 533-543.

Cooley, A. M., Modlizewski, J. L., Rommel, M. L., and Willis, J. H. 2011. Gene duplication in *Mimulus* underlies parallel floral evolution via independent *trans*-regulatory changes. *Current Biology* 21: 700-704.

Cronk, Q., and I. Ojeda. 2008. Bird-pollinated flowers in an evolutionary and molecular context. *Journal of Experimental Botany* 59: 715–727.

Cueva del Castillo, R., and Fairbairn, D. J. 2012. Macroevolutionary patterns of bumblebee body size: detecting the interplay between natural and sexual selection. *Ecology and Evolution* 2: 46-57.

Dellinger, A. S. (2020). Pollination syndromes in the 21st century: where do we stand and where may we go?. *New Phytologist* 228(4): 1193-1213.

Esfeld, K., Berardi, A. E., Moser, M., Bossolini, E., Freitas, L., and Kuhlemeier, C. 2018. Pseudogenization and resurrection of a speciation gene. *Current Biology* 28: 3776-3786.

Fægri, K., and van der Pijl, L. 1979. The Principles of Pollination Ecology. Pergamon Press, Oxford, UK.

Fenster, C. B., Armbruster, W. S., Wilson, P., Dudash, M. R., & Thomson, J. D. (2004). Pollination syndromes and floral specialization. *Annu. Rev. Ecol. Evol. Syst.* 35: 375-403.

Fisher, R.A., 1934. Adaptation and mutations. School Science Review 15:294-301.

Fox, J., S. Weisberg, B. Price, D. Adler, D. Bates, G. Baud-Bovy, B. Bolker, et al. 2020. car: Companion to Applied Regression.

Fraser, A. M., W. L. Mechaber, and J. G. Hildebrand. 2003. Electroantennographic and Behavioral Responses of the Sphinx Moth *Manduca sexta* to Host Plant Headspace Volatiles. *Journal of Chemical Ecology* 29: 1813–1833.

Free, J. B. 1970. Effect of flower shapes and nectar guides on the behaviour of foraging honeybees. *Behaviour* 37: 269-285.

Goldsmith, K. M. and T. H. Goldsmith. 1982. Sense of smell in the Black-chinned Hummingbird. *Condor* 84: 237-238.

Grant, K. A. 1966. A hypothesis concerning the prevalence of red coloration in California hummingbird flowers. *The American Naturalist* 100: 85-97.

Harrell, Jr., F. 2024. Hmisc: Harrell Miscellaneous. R package version 5.1-3, https://CRAN.R-project.org/package=Hmisc.

Herrera, C. M., M. C. Castellanos, and M. Medrano. 2006. Geographical context of floral evolution: towards an improved research programme in floral diversification. Oxford

University Press.

Hirschberg, J. (2001). Carotenoid biosynthesis in flowering plants. *Current Opinion in Plant Biology*, 4(3), 210-218.

Ho, W. W., & Smith, S. D. (2016). Molecular evolution of anthocyanin pigmentation genes following losses of flower color. *BMC Evolutionary Biology* 16: 1-10.

Hoballah M. E., T. Gu:bitz, J. Stuurman, L. Broger, M. Barone, T. Mandel, A. Dell'Olivo, M. Arnold, and C. Kuhlemeier. 2007. Single gene-mediated shift in pollinator attraction in *Petunia*. *The Plant Cell* 19: 779-790.

Holton, T. A., & Cornish, E. C. (1995). Genetics and biochemistry of anthocyanin biosynthesis. *The Plant Cell*, 7(7), 1071.

Hopkins, R., and M. D. Rausher. 2011. Identification of two genes causing reinforcement in the Texas wildflower *Phlox drummondii*. *Nature* 469: 411-414.

Hopkins, R., and M. D. Rausher. 2012. Pollinator-Mediated Selection on Flower Color Allele Drives Reinforcement. *Science* 335: 1090–1092.

Jiang, N., Dillon, F. M., Silva, A., Gomez-Cano, L., & Grotewold, E. (2021). Rhamnose in plants-from biosynthesis to diverse functions. *Plant Science,* 302, 110687.

Johnson, S. D. 2006. Pollinator-driven speciation in plants. *In* L. D. Harder, and S. C. H. Barrett [eds.], Ecology and Evolution of Flowers, 295–310. Oxford University Press.

Kaczorowski, R. L., M. C. Gardener, and T. P. Holtsford. 2005. Nectar traits in *Nicotiana* section *Alatae* (Solanaceae) in relation to floral traits, pollinators, and mating system. *American Journal of Botany* 92: 1270–1283.

Katoh, K., & Standley, D. M. (2013). MAFFT multiple sequence alignment software version 7: improvements in performance and usability. Molecular biology and evolution, 30(4), 772-780.

Katsu, K., Suzuki, R., Tsuchiya, W., Inagaki, N., Yamazaki, T., Hisano, T., ... & Matsui, K. (2017). A new buckwheat dihydroflavonol 4-reductase (DFR), with a unique substrate binding structure, has altered substrate specificity. *BMC plant biology*, 17(1), 1-14.

Kay, K. M., and D. L. Grossenbacher. 2022. Evolutionary convergence on hummingbird pollination in Neotropical *Costus* provides insight into the causes of pollinator shifts. *New Phytologist* 236: 1572-1583.

Kay, K. M., and R. D. Sargent. 2009. The Role of Animal Pollination in Plant Speciation: Integrating Ecology, Geography, and Genetics. *Annual Review of Ecology, Evolution, and Systematics* 40: 637–656.

Kellenberger, R. T., Byers, K. J. R. P., De Brito Francisco, R. M., Staedler, Y. M., LaFountain, A. M., Schönenberger, J., Schiestl, F. P., and Schlüter, P. M. 2019. Emergence of a flower colour polymorphism by pollinator-mediated overdominance. *Nature Communications* 10: 63.

Knauer, A. C., and F. P. Schiestl. 2015. Bees use honest floral signals as indicators of reward when visiting flowers. *Ecology Letters* 18: 135–143.

Koski, M. H. 2020. The role of sensory drive in floral evolution. *New Phytologist* 227: 1012-

1024.

Kuznetsova, A., P. B. Brockhoff, and R. H. B. Christensen. 2017. lmerTest Package: Tests in Linear Mixed Effects Models. *Journal of Statistical Software* 82: 1–26.

LaFountain, A. M., H. A. Frank, and Y.-W. Yuan. 2015. Carotenoid composition of the flowers of *Mimulus lewisii* and related species: Implications regarding the prevalence and origin of two unique, allenic pigments. *Archives of Biochemistry and Biophysics* 573: 32–39.

Leonard, A. S. and D. R. Papaj. 2011. 'X' marks the spot: the possible benefits of nectar guides to bees and plants. *Functional Ecology* 25: 1293-1301.

Letunic, I., & Bork, P. (2021). Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display and annotation. Nucleic acids research, 49(W1), W293-W296.

(a)Li H, Durbin R: Fast and accurate short read alignment with burrows-wheeler transform. *Bioinformatics*. 2009, 25 (14): 1754-1760. 10.1093/bioinformatics/btp324.

(b) Li H, Handsaker B, Wysoker A, et al. The Sequence Alignment/Map format and SAMtools. *Bioinformatics*. 2009;25:2078–9.

Li H. A statistical framework for SNP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data. *Bioinformatics* (2011) 27(21) 2987-93.

Li, P., Li, Y. J., Zhang, F. J., Zhang, G. Z., Jiang, X. Y., Yu, H. M., & Hou, B. K. (2017). The *Arabidopsis* UDP-glycosyltransferases UGT79B2 and UGT79B3, contribute to cold, salt and drought stress tolerance via modulating anthocyanin accumulation. *The Plant Journal*, 89(1), 85-103.

Liang, M., W. Chen, A. M. LaFountain, Y. Liu, F. Peng, R. Xia, H. D. Bradshaw, and Y.-W. Yuan. 2023. Taxon-specific, phased siRNAs underlie a speciation locus in monkeyflowers. *Science* 379: 576–582.

Liao, Y., Smyth, G. K., & Shi, W. (2014). featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics*, 30(7), 923-930.

Lowry, D. B., Modliszewski, J. L., Wright, K. M., Wu, C. A., and Willis, J. H. 2008. The strength and genetic basis of reproductive isolating barriers in flowering plants. *Philosophical Transactions of the Royal Society B* 363: 3009-3021.

Lowry, D. B., J. M. Sobel, A. L. Angert, T.-L. Ashman, R. L. Baker, B. K. Blackman, Y. Brandvain, et al. 2019. The case for the continued use of the genus name *Mimulus* for all monkeyflowers. *TAXON* 68: 617–623.

Maia, R., H. Gruson, J. A. Endler, and T. E. White. 2019. pavo 2: New tools for the spectral and spatial analysis of colour in r. *Methods in Ecology and Evolution* 10: 1097–1107.

McGarvey D. J. and R. Croteau. 1995. Terpenoid metabolism. *The Plant Cell* 7: 1015-1026.

Morawo, T., Burrows, M., and Fadamiro, H. 2017. Electroantennogram response of the parasitoid, *Microplitis croceipes* to host-related odors: the discrepancy between relative abundance and level of antennal responses to volatile compound. *F1000*

Research 5: 2725.

Mostafa, S., Y. Wang, W. Zeng, and B. Jin. 2022. Floral Scents and Fruit Aromas: Functions, Compositions, Biosynthesis, and Regulation. *Frontiers in Plant Science* 13.

Nadeau, C. P., and M. C. Urban. 2019. Eco-evolution on the edge during climate change. *Ecography* 42: 1280–1297.

Nelson, T. C., A. M. Stathos, D. D. Vanderpool, F. R. Finseth, Y.-W. Yuan, and L. Fishman. 2021. Ancient and recent introgression shape the evolutionary history of pollinator adaptation and speciation in a model monkeyflower radiation (*Mimulus* section *Erythranthe*). *PLOS Genetics* 17: e1009095.

Nesom, G. L. and Egger, J. M. 2014. Review of the *Castilleja purpurea* complex (Orobanchaceae). *Phytoneuron* 15: 1-16.

Núñez, P., Méndez, M., and López-Rull, I. 2021. Can foraging hummingbirds use smell? A test with the Amazilia Hummingbird *Amazilia amazilia. Ardeola* 68: 433-444.

Oksanen, J., G. L. Simpson, F. G. Blanchet, R. Kindt, P. Legendre, P. R. Minchin, R. B. O'Hara, et al. 2022. vegan: Community Ecology Package.
Orr, H. A. 1998. The Population Genetics of Adaptation: The Distribution of Factors Fixed During Adaptive Evolution. Evolution 52: 935–949.

Orr, H. A. 2005. The genetic theory of adaptation: a brief history. Nature Reviews Genetics 6: 119–127.

Peng, F., K. J. R. P. Byers, and H. D. Bradshaw, Jr. 2017. Less is more: Independent loss-of-function *OCIMENE SYNTHASE* alleles parallel pollination syndrome diversification in monkeyflowers (*Mimulus*). *American Journal of Botany* 104: 1055-1059.

Powell, D. R. (2015). Degust: interactive RNA-seq analysis. Drpowell/Degust, 4(1), 4-1.

Price, M. N., Dehal, P. S., & Arkin, A. P. (2010). FastTree 2–approximately maximum-likelihood trees for large alignments. PloS one, 5(3), e9490.

Quattrocchio, F., Wing, J., van der Woude, K., Souer, E., de Vetten, N., Mol, J., and Koes, R. 1999. Molecular analysis of the *anthocyanin2* gene of *Petunia* and its role in the evolution of flower color. *Plant Cell* 11: 1433-1444.

R Core Team (2022). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Raguso, R. A. 2008. Wake up and smell the roses: the ecology and evolution of floral scent. *Annual Review of Ecology, Evolution, and Systematics* 39: 549-569.

Rasmont P., A. Coppee, D. Michez, T. De Meulemeester. 2008. An overview of the *Bombus terrestris* (L. 1758) subspecies (Hymenoptera: Apidae). *Annales de la Société entomologique de France (N.S.)* 44: 243-250.

Riffell, J. A., and Alarcón, R. 2013. Multimodal floral signals and moth foraging decisions. *PLOS ONE* 8:e72809.

Rivera, S., Vilaró, F., & Canela, R. (2011). Determination of carotenoids by liquid chromatography/mass spectrometry: effect of several dopants. *Analytical and*

bioanalytical chemistry, 400, 1339-1346.

Revelle, W. 2023. psych: Procedures for Psychological, Psychometric, and Personality Research.

Sánchez-Cabrera, M., Narbona, E., Arista, M., Ortiz, P. L., Jiménez-López, F. J., Fuller, A., ... & Whittall, J. B. (2023). Can a flower color ancestral polymorphism transcend speciation?. bioRxiv, 2023-11.

Schiestl, F. P., and S. D. Johnson. 2013. Pollinator-mediated evolution of floral signals. *Trends in Ecology & Evolution* 28: 307–315.

Schlüter, P. M., Xu, S., Gagliardini, V., Whittle, E., Shanklin, J., Grossniklaus, U., and Schiestl, F. P. 2011. Stearoyl-acyl carrier protein desaturases are associated with floral isolation in sexually deceptive orchids. *Proceedings of the National Academy of Sciences USA* 108: 5696-5701.

Sérino, S., L. Gomez, G. Costagliola, and H. Gautier. 2009. HPLC Assay of Tomato Carotenoids: Validation of a Rapid Microextraction Technique. Journal of Agricultural and Food Chemistry 57: 8753–8760.

Sheehan, H., Moser, M., Klahre, U., Esfeld, K., Dell'Olivo, A., Mandel, T., Metzger, S., Vandenbussche, M., Freitas, L., and Kuhlemeier, C. 2016. *MYB-FL* controls gain and loss of floral UV absorbance, a key trait affecting pollinator preference and reproductive isolation. *Nature Genetics* 48: 159-166.

Short, A. W., & Streisfeld, M. A. (2023). Ancient hybridization leads to the repeated evolution of red flowers across a monkeyflower radiation. *Evolution Letters* 7(5): 293-304.

Simonds, V., and Plowright, C. M. S. 2004. How do bumblebees first find flowers? Unlearned approach responses and habituation. *Animal Behaviour* 67: 379-386.

Smith, S. D. and Rausher, M. D. 2011. Gene loss and parallel evolution contribute to species differences in flower color. *Molecular Biology and Evolution* 28: 2799-2810.

Stankowski, S., & Streisfeld, M. A. (2015). Introgressive hybridization facilitates adaptive divergence in a recent radiation of monkeyflowers. *Proceedings of the Royal Society B: Biological Sciences* 282(1814): 20151666.

Stankowski, S., J. M. Sobel, and M. A. Streisfeld. 2017. Geographic cline analysis as a tool for studying genome-wide variation: a case study of pollinator-mediated divergence in a monkeyflower. *Molecular Ecology* 26: 107–122.

Strauss, S., and J. B. Whittall. 2006. Non-pollinator agents of selection on floral traits. *In* L. D. Harder, and S. C. H. Barrett [eds.], Ecology and Evolution of Flowers, 120–138. Oxford University Press.

Song, J., Shi, L., Li, D., Sun, Y., Niu, Y., Chen, Z., ... & Chen, S. (2012). Extensive pyrosequencing reveals frequent intra-genomic variations of internal transcribed spacer regions of nuclear ribosomal DNA.

Sun, Z., Gantt, E., & Cunningham, F. X. (1996). Cloning and functional analysis of the β-carotene hydroxylase of *Arabidopsis thaliana*. *Journal of Biological Chemistry*, 271(40), 24349-24352.

Sutherland, S. D., and R. K. Vickery. 1993. On the Relative Importance of Floral Color, Shape, and Nectar Rewards in Attracting Pollinators to *Mimulus*. *The Great Basin Naturalist* 53: 107–117.

Tamura K. and Nei M. (1993). Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. Molecular Biology and Evolution 10:512-526.

Thomson J. D. and P. Wilson. 2008. Explaining evolutionary shifts between bee and hummingbird pollination: convergence, divergence, and directionality. *International Journal of Plant Sciences* 169: 23-38.

Tripp E. A. and P. S. Manos. 2008. Is floral specialization an evolutionary dead end? Pollination system transitions in *Ruellia* (Acanthaceae). *Evolution* 62: 1712-1737.

van der Niet, T., and Johnson, S. D. 2012. Phylogenetic evidence for pollinator-driven diversification in angiosperms. *Trends in Ecology and Evolution* 27: 353-361.

van der Niet, T., R. Peakall, and S. D. Johnson. 2014. Pollinator-driven ecological speciation in plants: new evidence and future perspectives. *Annals of Botany* 113: 199–212.

Vickery, R. K. 1995. Speciation in *Mimulus*, or, Can a Simple Flower Color Mutant Lead to Species Divergence? *The Great Basin Naturalist* 55: 177–180.

Vickery, R. K., and P. K. Vickery. 1992. POLLINATOR PREFERENCES FOR YELLOW, ORANGE, AND RED FLOWERS OF MIMULUS VERBENACEUS AND M. CARDINALIS. *The Great Basin Naturalist* 52: 145–148.

Vickery, R. K., and P. K. Vickery. 1992. POLLINATOR PREFERENCES FOR YELLOW, ORANGE, AND RED FLOWERS OF *MIMULUS VERBENACEUS* AND *M. CARDINALIS*. *The Great Basin Naturalist* 52: 145–148.

Wang, L.; Lui, A.C.W.; Lam, P.Y.; Liu, G.; Godwin, I.D.; Lo, C. Transgenic expression of flavanone 3-hydroxylase redirects flavonoid biosynthesis and alleviates anthracnose susceptibility in sorghum. *Plant Biotechnol. J.* 2020, *18*, 2170–2172.

Waser, N. M., and D. R. Campbell. 2004. Ecological Speciation in Flowering Plants. *In* U. Dieckmann, M. Doebeli, J. A. J. Metz, and D. Tautz [eds.], Adaptive Speciation, 264–277. Cambridge University Press.

Waser, N. M., CaraDonna, P. J., & Price, M. V. (2018). Atypical flowers can be as profitable as typical hummingbird flowers. *The American Naturalist* 192(5): 644-653.

Wenzell, K. E., K. A. Skogen, and J. B. Fant. 2023. Range-wide floral trait variation reflects shifts in pollinator assemblages, consistent with pollinator-mediated divergence despite generalized visitation. *Oikos*: e09708.

Wessinger, C. A., A. M. Katzer, P. M. Hime, M. D. Rausher, J. K. Kelly, and L. C. Hileman. 2023. A few essential genetic loci distinguish *Penstemon* species with flowers adapted to pollination by bees or hummingbirds. PLOS Biology 21: e3002294.

Whittall, J. B., & Hodges, S. A. (2007). Pollinator shifts drive increasingly long nectar spurs in columbine flowers. *Nature* 447(7145): 706-709.

Wu, C. A., D. B. Lowry, A. M. Cooley, K. M. Wright, Y. W. Lee, and J. H. Willis. 2008.

Mimulus is an emerging model system for the integration of ecological and genomic studies. *Heredity* 100: 220–230.

Yoshida, K., Iwasaka, R., Shimada, N., Ayabe, S. I., Aoki, T., & Sakuta, M. (2010). Transcriptional control of the dihydroflavonol 4-reductase multigene family in *Lotus japonicus*. *Journal of plant research*, 123, 801-805.

Yuan, Y.-W. 2019. Monkeyflowers (*Mimulus*): new model for plant developmental genetics and evo-devo. *New Phytologist* 222: 694–700.

Yuan, Y.-W., J. M. Sagawa, L. Frost, J. P. Vela, and H. D. Bradshaw Jr. 2014. Transcriptional control of floral anthocyanin pigmentation in monkeyflowers (*Mimulus*). *New Phytologist* 204: 1013–1027.

Yuan, Y.-W., K. J. R. P. Byers, and H. Bradshaw. 2013. The genetic control of flower–pollinator specificity. *Current Opinion in Plant Biology* 16: 422–428.

900
901

Figure 1. (A) Focal floral morphs: *M. verbenaceus*, yellow form, MvYL (i) and red form, MvBL (ii); *M. cardinalis*, yellow form, SM (iii) and red form, CE-10 (iv). (B) Spectral reflectance of central petal lobes of focal lines. (C) Total anthocyanins extracted from corolla tissues of each floral morph, measured by absorbance at 525nm; (D) total carotenoids, measured as absorbance at 450nm, relative to average mass (g) of corolla per line.

Figure 2. Anthocyanins. (A) Individual anthocyanin levels in corollas of the four focal morphs as detected by UHPLC/MS. Abbreviations: Cyan = Cyanidin, Pel = Pelargonidin, Glc = Glucoside, Rha = Rhamnoside, Pen = Pentose, Ace = Acetate. (B) Core anthocyanin biosynthesis pathway. Genes responsible for each metabolic step are shown in bold next to each red arrow. Genes significantly differentially expressed (p<0.05) in each yellow-flowered morph when compared to their red counterparts are denoted by an arrow corresponding with their color ID in the key in (C) Expression of anthocyanin biosynthesis regulators NEGAN and PELAN as filtered gene counts.

Figure 3. Carotenoids. (A) Individual carotenoid levels in whole corollas of the four focal
morphs as detected by HPLC/MS. (B) Carotenoid biosynthesis pathway. Genes responsible for
each metabolic step are shown in bold next to each red arrow. Genes significantly differentially
expressed (p<0.05) in each yellow-flowered morph when compared to their red counterparts are
denoted by an arrow corresponding with their color ID in the key in (C) Expression of carotenoid
biosynthesis genes as filtered gene counts.

Figure 4. Chromosome maps. Whole genomes of yellow-flowered morphs MvY and McY were mapped to the published genomes of their respective red counterparts and the number of variants called using bcftools. Chromosomes were divided into 30kb windows with the number of variants detected per window plotted for each chromosome and the two species overlaid.

Figure 5. Floral scent emissions vary in total volatiles (both species) and global composition of VOCs (*M. cardinalis*) between red and yellow conspecific floral morphs. A: Comparison of total volatile emissions of the four species-color morph lines, with the y-axis on a log scale. B: NMDS plot of floral scent samples in *M. verbenaceus*. C: NMDS plot of floral scent samples in *M. cardinalis*. D: Floral scent composition in color morphs of *M. verbenaceus*. Only compounds contributing at least 1% of total emission in either morph are shown and area of the pie is relative to total emission of the yellow morph. E: Floral scent composition in color morphs of *M. cardinalis*, with compound choice and area as in (D). Note that no unified NMDS plot of *M. verbenaceus* and *M. cardinalis* is possible due to a lack of volatile sharing between species.

Figure 6. (A) NMDS ordination of floral morphological traits by line. Line centroid is labeled, with circles showing standard deviation. Individual trait loadings with $p < 0.05$ (envfit()) are shown. (B) Violin plots of selected morphology   traits by line, with mean +/- standard error).

959
Figure 7. (A) Floral color perception by bees. Central petal lobe reflectance data for each floral
morph plotted on a model of bee visual space, where the center point represents low visual
contrast to a green vegetative background. (B) EAG responses to scent stimuli, including
phenylacetaldehyde (PAA, a floral VOC known to be detectable by *Bombus terrestris*), air
(negative control), extraction solvent alone, and floral scent extractions of MVYL, MVBL, SM,
and CE10. Bee preference metrics in pairwise choice experiments between conspecific pairs of
floral color morphs (N= 10 trials per species comparison): (C) number of visits; (D) number of
flowers probed. (E) Bumblebees probing yellow morphs, often with poor contact to anthers and

968   stigma (left, arrows), and resulting damage to flowers (right). Note corolla tube split open from

969   bee entering corolla in *M. verbenaceus* (top photos).

Table 1. Summary of differences across traits in yellow floral morphs (MvY and McY) relative to their red conspecifics (MvR and McR, respectively). Plus sign (+) indicates a significant increase in yellow morphs compared to red; minus (-) indicates a significant decrease. Traits that did not differ significantly among morphs within species show "n.s." See text for details.

	M. verbenaceus	*M. cardinalis*
	MvY (compared to MvR)	**McY (compared to McR)**
Floral color		
Anthocyanins (total)	**-**	**-**
Cyanidin-glucoside	**absent**	**-**
Pelargonidin-Glucoside	**absent**	**absent in both**
Cyanidin-Glucoside-Rhamnose	**+ (absent in MvR)**	**-**
Pelargonidin-Glucoside-Rhamnose	**+ (absent in MvR)**	**-**
Pelargonidin-pentose	**absent**	**absent in both**
Isomer of Pel-Glc-Rha	**absent**	**absent in McY**
Cyanidin-Glucoside-Acetate	**absent**	**absent in both**
Pelargonidin-Glucoside-Acetate	**absent**	**absent in both**
Anthocyanin regulatory gene expression		
PELAN	**+**	**- (deleted)**

NEGAN	**absent in both**	+
Carotenoids (total)	+	+
zeaxanthin	**n.s.**	**-**
antheraxanthin	+	**n.s.**
violaxanthin	+	+
neoxanthin	+	+
mimulaxanthin	+	+
Carotenoid biosynthesis gene expression		
BCH	+	+
ZEP	+	+
Floral scent		
Total volatile emissions	+	+
Floral morphology		
corolla length	+	**-**
opening width	+	+
opening height	+	+
display width	**n.s.**	**n.s.**
display height	+	**n.s.**

Nectar

nectar volume   **n.s.**   **n.s.**

sucrose concentration   +   **n.s.**

Bumblebee preference

Flowers visited   +   +

Flowers probed   **n.s.**   +

Handling time   +   +

974
975
Table 2. Volatile emissions (ng/flower/hour ± standard error of the mean) of each volatile
identified from the four lines. Numbers in parentheses after each emission value indicate the
number of samples a compound was found in from that line (total sample numbers for each line
are in the table header). Under the Compound header, superscript letters: A: compound identity
validated using authentic reference standards; B: compound identity validated using published
Kovats Retention Indices, our calculated Kovats Retention Indices, and NIST Library spectrum
matching; C: compound identity could not be validated and compound is listed as an unknown .
Values in bold text differed significantly between red and yellow forms.
984

Compound Name	Type	Kovats Retention Index	Yellow *M. verbenaceus* MvY (n = 17)	Red *M. verbenaceus* MvR (n = 18)	Yellow *M. cardinalis* McY (n = 15)	Red *M. cardinalis* McR (n = 32)
Total emissions (summed across all compounds)	-	-	**675.830 ± 43.538**	**291.169 ± 19.487**	**5.036 ± 0.493**	**2.985 ± 0.240**
Cymene[A] (m-/o-/p-structure could not	aromatic	1021	**7.096±0.860 (16)**	**2.374±0.361 (17)**	Absent (0)	Absent (0)

be determined with standards)						
Cuminaldehyde[A]	aromatic	1236	**1.784±0.280 (15)**	**0.407±0.125 (10)**	Absent (0)	Absent (0)
Cuminol[A]	aromatic	1290	**1.037±0.211 (13)**	**0.071±0.042 (3)**	Absent (0)	Absent (0)
1-Octen-3-ol[A]	FAD	979	**3.342±0.397 (16)**	**0.299±0.180 (3)**	Absent (0)	Absent (0)
Tridecane[A]	FAD	1300	**0.496±0.144 (10)**	**1.699±0.413 (12)**	Absent (0)	Absent (0)
α-Thujene[A]	terpenoid	923	**55.323±5.055 (17)**	**12.770±2.420 (18)**	Absent (0)	Absent (0)
α-Pinene[A]	terpenoid	929	**254.941±18.331 (17)**	**126.776±8.250 (18)**	Absent (0)	Absent (0)
Camphene[A]	terpenoid	942	**2.132±0.288 (16)**	**0.580±0.044 (18)**	Absent (0)	Absent (0)
Sabinene[A]	terpenoid	971	**114.602±9.888 (17)**	**51.895±4.510 (18)**	Absent (0)	Absent (0)
β-Pinene[A]	terpenoid	972	**26.593±3.216 (17)**	**17.861±0.767 (18)**	Absent (0)	Absent (0)
β-Myrcene[A]	terpenoid	990	**3.630±0.267 (17)**	**1.365±0.176 (15)**	Absent (0)	Absent (0)
α-Terpinene[A]	terpenoid	1013	**1.670±0.278 (13)**	**0.395±0.114 (11)**	Absent (0)	Absent (0)
β-Phellandrene[A]	terpenoid	1025	**52.585±3.663 (17)**	**11.589±1.592 (18)**	Absent (0)	Absent (0)
Eucalyptol[A]	terpenoid	1026	**2.797±0.304 (17)**	**0.750±0.089 (18)**	Absent (0)	Absent (0)
γ-Terpinene[A]	terpenoid	1055	**2.050±0.405 (12)**	**0.125±0.071 (3)**	Absent (0)	Absent (0)
Sabinene	terpenoid	1063	**34.796±5.0**	**8.154±1.80**	Absent (0)	Absent (0)

hydrate[A]			**26 (15)**	**0 (11)**		
Terpinolene[A]	terpenoid	1085	**0.587±0.133 (13)**	**0.068±0.032 (4)**	Absent (0)	Absent (0)
Pinene oxide[A]	terpenoid	1092	**12.466±1.461 (17)**	**5.428±0.545 (17)**	Absent (0)	Absent (0)
Unknown monoterpenoid[C]	terpenoid	1095	**9.281±1.130 (17)**	**2.636±0.337 (17)**	Absent (0)	Absent (0)
β-Thujone[A]	terpenoid	1113	**1.146±0.190 (15)**	**0.311±0.067 (14)**	Absent (0)	Absent (0)
α-Campholenal[B]	terpenoid	1123	**5.080±0.513 (17)**	**1.864±0.267 (18)**	Absent (0)	Absent (0)
Nopinone[A]	terpenoid	1133	7.468±0.526 (17)	5.563±1.046 (18)	Absent (0)	Absent (0)
Sabinol[B]	terpenoid	1137	**2.111±0.397 (15)**	**Absent (0)**	Absent (0)	Absent (0)
(*E*)-Verbenol[A]	terpenoid	1142	**2.493±0.221 (17)**	**0.175±0.101 (3)**	Absent (0)	Absent (0)
Sabine ketone[B]	terpenoid	1153	21.484±2.918 (17)	14.803±2.303 (18)	Absent (0)	Absent (0)
Pinocarvone[B]	terpenoid	1157	**2.041±0.178 (17)**	**0.555±0.128 (12)**	Absent (0)	Absent (0)
Myrtenal isomer 1[A]	terpenoid	1181	**1.105±0.178 (14)**	**0.188±0.109 (3)**	Absent (0)	Absent (0)
Cryptone[B]	terpenoid	1182	**3.186±0.381 (17)**	**1.597±0.179 (18)**	Absent (0)	Absent (0)
α-Terpineol[A]	terpenoid	1188	**0.070±0.010 (16)**	**Absent (0)**	Absent (0)	Absent (0)
Myrtenal isomer 2[A]	terpenoid	1191	**5.490±0.436 (17)**	**1.115±0.402 (6)**	Absent (0)	Absent (0)
Verbenone[A]	terpenoid	1205	**4.426±0.784 (12)**	**0.268±0.184 (2)**	Absent (0)	Absent (0)

M. verbenaceus Unknown 1[C]	unknown	894	**1.047±0.219 (14)**	**0.139±0.045 (7)**	Absent (0)	Absent (0)
M. verbenaceus Unknown 2[C]	unknown	1176	**1.130±0.168 (15)**	**0.438±0.126 (9)**	Absent (0)	Absent (0)
M. verbenaceus Unknown 3[C]	unknown	1273	3.348±0.448 (16)	3.922±0.983 (15)	Absent (0)	Absent (0)
M. verbenaceus Unknown 4[C]	unknown	1312	1.710±0.228 (15)	1.375±0.173 (18)	Absent (0)	Absent (0)
M. verbenaceus Unknown 5[C]	unknown	1392	**23.187±3.656 (16)**	**0.521±0.357 (2)**	Absent (0)	Absent (0)
M. verbenaceus Unknown 6[C]	unknown	1396	1.517±0.460 (11)	1.256±0.212 (14)	Absent (0)	Absent (0)
M. verbenaceus Unknown 7[C]	unknown	1942	**0.537±0.207 (8)**	**9.071±1.868 (17)**	Absent (0)	Absent (0)
M. verbenaceus Unknown 8[C]	unknown	2028	**0.044±0.030 (2)**	**2.766±0.607 (16)**	Absent (0)	Absent (0)
Methyl salicylate[A]	aromatic	1191	Absent (0)	Absent (0)	**Absent (0)**	**0.626±0.047 (29)**
Hexyl 2-methylbutyrate[A]	FAD	1238	Absent (0)	Absent (0)	**1.888±0.132 (15)**	**0.343±0.032 (29)**
Phenethyl	FAD	1485	Absent (0)	Absent (0)	0.115±0.02	0.124±0.01

					8 (10)	5 (25)
2-methylbutyrate[A]						
Lavender lactone[B]	lactone	1041	Absent (0)	Absent (0)	**1.265±0.160 (15)**	**0.412±0.069 (22)**
Linalool oxide isomer 1[A]	terpenoid	1070	Absent (0)	Absent (0)	**0.105±0.023 (10)**	**0.360±0.036 (30)**
Linalool oxide isomer 2[A]	terpenoid	1086	Absent (0)	Absent (0)	**0.005 (1)**	**0.272±0.045 (27)**
Valencene[A]	terpenoid	1482	Absent (0)	Absent (0)	**0.421±0.093 (13)**	**0.093±0.043 (7)**
M. cardinalis unknown 1[C]	unknown	950	Absent (0)	Absent (0)	0.717±0.093 (13)	0.755±0.181 (13)
M. cardinalis unknown 2[C]	unknown	1052	Absent (0)	Absent (0)	**0.334±0.074 (10)**	**Absent (0)**
M. cardinalis unknown 3[C]	unknown	1687	Absent (0)	Absent (0)	**0.185±0.018 (14)**	**Absent (0)**