A comprehensive review of South Australia's Great Artesian Basin

spring and discharge wetlands biota

Beasley-Hall, P. G.[1,2]*, Hedges, B. A.[1], Cooper, S. J. B.[1,2], Austin, A .D.[1,2], Guzik, M.

T.[1,2]

[1] Invertebrate Systematics and Biodiversity Group, School of Biological Sciences, The

University of Adelaide, Adelaide, SA, Australia

[2] South Australian Museum, Adelaide, SA, Australia

****Correspondence**

Perry Beasley-Hall, School of Biological Sciences, The University of Adelaide,

Adelaide, SA, Australia

Email: perry.beasley-hall@adelaide.edu.au

Abstract

Context: The Great Artesian Basin (GAB) feeds thousands of springs in Australia's
arid centre, supporting relictual species not found elsewhere on Earth. Springs are
considerably threatened by ongoing water abstraction by industry. Robust management
plans are needed to prevent further extirpations of GAB taxa, but fundamental
biodiversity knowledge is lacking.
20

21   **Aims:** We aimed to characterise major organismal groups in South Australian GAB
22   springs and surrounding wetlands, their conservation and taxonomic status, and
23   potential biodiversity hotspots and connectivity of spring ecosystems.
24

25   **Methods:** Focusing on South Australia as a case study, we conducted a comprehensive
26   review of GAB spring biota based on the published scientific and grey literature.
27

28   **Key results:** Almost 500 taxa have been recorded from GAB springs, the majority
29   being invertebrates. Community composition is highly heterogeneous among spring
30   clusters and the true extent of spring biodiversity is far greater than currently known.
31

32   **Conclusions:** GAB springs have intrinsic value as refugia for both endemics and
33   cosmopolitan taxa. GAB invertebrates are poorly conserved and largely lacking in
34   taxonomic knowledge. We highlight several potential biodiversity hotspots that have
35   been overlooked in the literature.
36

Implications: Fundamental biodiversity information on the GAB is crucial for
decision-making in conservation management, for industry, and for Traditional
Custodians.

Additional keywords: Great Artesian Basin, groundwater-dependent ecosystems,
aquifers, springs, biodiversity, biogeography

Lay summary

The Great Artesian Basin (GAB) is Australia's largest freshwater resource. Springs fed

by the GAB support many species not found elsewhere on Earth, but conservation is

hindered by a lack of fundamental knowledge about the plants, animals, and fungi

reliant on these habitats. Using South Australia as a case study, we provide a

comprehensive review of GAB biodiversity in that state.

1   Introduction

The Great Artesian Basin (GAB) is Australia's largest groundwater resource, spanning over one fifth of the continent's area or almost 2 million km^2 (Habermehl, 2020). At the outer margins of the GAB where the confining layer is thin, pressurised water is often forced to the surface to form springs and associated wetlands. Thousands of such springs are found throughout Central Australia, with >5,000 individual spring vents (discrete discharge points of water) in South Australia (hereafter SA) (Arabana Aboriginal Corporation, 2021; Government of South Australia, 2023), >2,000 in Queensland (Queensland Department of Regional Development, Manufacturing and Water, 2023), and >400 in New South Wales (New South Wales Department of Planning, Industry and Environment, 2021). From an ecological and evolutionary standpoint, GAB-fed springs support wetlands representing "museums of biodiversity" housing plant and animal species not found elsewhere on Earth (Murphy et al., 2015). As relicts of the continent's mesic past, species endemic to GAB springs often have exceptionally small distributions. It is not uncommon for taxa to be restricted to a single cluster of springs, termed *ultra-short range endemics* (examples found in Gotch et al., 2008; Guzik et al., 2019, 2012; Harvey, 1990; King, 2009; King et al., 2014; Murphy et al., 2012, 2009).

The conservation and management of groundwater resources is lacking globally (Famiglietti, 2014). Springs fed by the GAB are no exception and are considerably threatened by a range of industrial and pastoral practices (Fairfax and Fensham, 2002; Lewis and Harris, 2020; Mudd, 2000). The sinking of more than 50,000 artificial boreholes and the direct abstraction of Basin water have led to substantial reductions in artesian pressure (hereafter *drawdown*) and spring flow (Beasley-Hall et al., 2023c; Gotch et al., 2016; Great Artesian Basin Coordinating Committee, 2019; Mudd, 2000). A complete cessation of flow has occurred for an estimated 800 springs across Australia (Andersen et al., 2016; Fensham et al., 2016), and declines of endemic fauna have been documented as a result of this degradation of spring habitats (Fensham et al., 2010). Springs and their biota also face threats in addition to drawdown, including grazing, trampling, and soiling of wetlands by livestock such as cattle (Fatchen, 2000), overabundant pest species (Kerezsy, 2015; Noack, 1994), climate change (Ordens et

al., 2020), and tourist activity (Witjira National Park Co-management Board, 2022).

Despite remediation efforts, drivers of spring flow remain poorly understood and net

water flows in GAB springs appear to still be reducing in certain regions of the Basin

(Green and Berens, 2013).

The development of a biotic inventory has been identified as an urgent action to

recover the community of native species dependent on GAB springs (Fensham et al.,

2010). Publicly available data can help to inform future management plans and early

warning systems that detect changes in these ecosystems of high biodiversity value

(Brack et al., 2015; Obura et al., 2019; Vaughan et al., 2001) and so improving their

accessibility is a clear priority. Further, industry stakeholders frequently rely on biotic

inventories to ensure they are fulfilling their environmental obligations. This is

particularly relevant for the GAB in the context of drawdown. Yet, basic taxonomic

and biological information for these communities is still lacking, inaccessible, or

disparate, making it difficult to accurately assess and monitor all spring species in the

face of rapid, human-driven change. The Queensland Government has published

information on metrics such as water quality, chemistry, flow rate, biodiversity, and

spring condition (Queensland Government, 2018), but such information is not public

for SA and New South Wales. This lack of taxonomic information is particularly

pronounced for endemic spring invertebrates (Rossini, 2020). The relationship between

environmental characteristics of springs and biodiversity is also not well understood

(Fensham et al., 2010; Green and Berens, 2013; Harris, 1992; Rossini et al., 2018),

meaning it is currently difficult to establish clear conservation priorities for GAB

spring taxa in certain locations over others. The compilation of baseline information on

these taxa in a centralised, accessible manner would represent a fundamental resource

to facilitate future conservation work on this ecological community. Here, we

conducted a literature review of species associated with springs in South Australia,

selected due to its high number of GAB springs (>60% of all active springs) and the

comparatively well-studied nature of its biodiversity (King, 2009; Murphy et al., 2015,

2013; Zeidler and Ponder, 1989). We specifically aimed to: 1) Characterise major

organismal groups in SA GAB springs and surrounding wetlands; 2) Document the

conservation and taxonomic status of these groups to highlight knowledge gaps; 3)

Identify hotspots of biodiversity in SA GAB springs; and 4) Assess connectivity of SA

GAB spring communities.

2   Materials and Methods

2.1   Literature review and database construction

We undertook a non-systematic review of all available information relevant to the
biodiversity of GAB springs in South Australia. A formal systematic searching strategy
was not possible as a large proportion of information on springs is present in
unpublished or grey literature such as government reports, internal publications from
mining companies, and museum records. We therefore relied heavily on repositories
such as the Government of South Australia's WaterConnect portal (South Australian
Department for Environment and Water, 2023) and The Atlas of Living Australia
(Belbin et al., 2021). We were specifically interested in data indicating the presence or
absence of species in or around spring wetlands as well as their occurrence extents, if
available, to gauge where taxa occurred and their degree of endemicity. The scope of
this review spanned the three spring supergroups in SA: Kati Thanda–Lake Eyre,
Munda / Lake Frome, and Dalhousie (Figure 1). **Supergroups** are clusters of **spring
complexes**, which themselves contain **spring groups** composed of individual **springs**.
Spring complexes share similar geomorphological characteristics and water chemistry,
whereas groups are clusters of springs sourcing from the same fault or structure (Lewis
et al., 2013). Individual springs are composed of permanent wetland vegetation with at
least one **vent**, a discrete discharge point of water (Gotch, 2013). Within these three
supergroups, we focused on 23 spring complexes containing 170 spring groups
(Beasley-Hall et al., 2023a for raw research data via FigShare). Whilst the number of
these spring clusters, as well as their naming conventions, have previously not been
standardised (Gotch et al., 2016), we selected these locations due to their widespread
use in conservation management by the Australian Government (Department of the
Environment, 2022; Lewis and White, 2013).

162 **Figure 1:** Springs fed by the Great Artesian Basin (GAB) in South Australia. Springs are
163 classified hierarchically as supergroups, the broadest classification (A, Dalhousie; B, Kati
164 Thanda–Lake Eyre; C, Munda / Lake Frome); complexes (labelled coloured circles); groups
165 (coloured points); springs; and finally vents, discrete water discharge points. The approximate
166 area of the GAB is shown in dark purple and the location of the SA GAB springs is indicated
167 by the dashed square.

170   We retrieved Aboriginal names for spring groups from a key anthropological
171 study (Hercus & Sutton, 1985) to ensure locations were referred to by a dual naming
172 convention whenever possible. The validity of these names was confirmed by The
173 Arabana Aboriginal Corporation in a personal communication to the authors. For
174 locations with dual names, but lacking "official" formatting (e.g., not used by

federal/state government legislation or publications), we followed the Australian

Government's style manual (Australian Government, 2023). We chose not to

standardise names across springs, groups, or complexes as they refer to specific

locations with distinct associated mythologies and histories (Hercus & Sutton, 1985).

Information below the spring group level, where present, was standardised to the group

level due to the rarity of such records and the inconsistency of spring vent naming

conventions. As GAB spring fauna are often morphologically cryptic yet genetically

distinct, we recorded species as separate taxa if explicitly indicated by their species

authorities based on genetic data following Rossini *et al.* (2018).

To supplement presence/absence records of spring taxa, we also gathered

information related to taxonomic status, conservation status, common names,

synonyms, and endemicity where available. Taxonomic status was coded into three

categories: taxa for which there is a corresponding formal taxonomic description

("described"), taxa awaiting taxonomic description ("undescribed"), or those for which

species-level occurrence records were not available (e.g., a family-only record;

"unidentified"). For taxa with species-level identifications, conservation status

information was retrieved from the Species Profile and Threats Database to capture

listings under the IUCN Red List of Threatened Species and Australian federal and

state environmental legislation (Department of Climate Change, Energy, the

Environment and Water, 2023). Finally, for each endemic taxon we noted whether it

occupied only one spring group, complex, or supergroup. Except for undescribed

species with well-established occurrence records, endemicity was not recorded for taxa

without a species-level identification.

2.2   Biodiversity metrics

For each spring group, we first transformed presence/absence records to weight them

by whether they represented resolution at the spring group level (hereafter confident

records) or those which referred to a taxon's presence within a certain complex but did

not supply spring group information (hereafter coarse records). To assess the inclusion

of uncertain occurrence information, we calculated metrics for our entire dataset and a

subset of the data only considering confident records. As the dataset included only

presence/absence records, we were limited in our choice of biodiversity metrics and

focused on species richness and endemicity. We calculated species richness values (hereafter taxon richness) by the number of putative taxa in each spring. Spring groups were ranked by the degree of endemicity of their biota using a modification of the scoring system developed by Fensham and Price (Fensham and Price, 2004). Originally applied to GAB flora in spring complexes in Queensland, the ranking has since been expanded to fauna across Australia (Rossini et al., 2018) and relies on the number of populations corresponding to the most widespread taxon in a spring dataset. The desert goby *Chlamydogobius eremius* (Zietz, 1896) is the most widespread SA GAB spring endemic, occurring across 24 known groups (Gotch et al., 2016; Rossini et al., 2018); we used this value as a proxy for the taxon's number of populations, although we acknowledge this may be an underestimate as spring groups do not necessarily share permanent wetlands. Rankings for each endemic taxon were first calculated by dividing 24 by the number of groups the taxon occurred in such that *C. eremius* (24/24) would receive the lowest ranking due to its comparably widespread distribution. As all endemic taxa assessed here do not occur beyond their respective supergroups, each taxon was then scored by whether it was further restricted to a single spring complex (+1) or group (+2). These scores, hereafter *endemicity rankings*, were summed for each spring group. We visualised these metrics using QGIS (QGIS Association, 2023). Spring groups were first mapped using latitude and longitude information corresponding to vents retrieved from the Government of South Australia's WaterConnect portal (Government of South Australia, 2023); for groups containing multiple vents, centroids were calculated to approximate their location. Circles corresponding to the above metrics per group were scaled using the Flannery method. To assess differences in community composition among spring groups, we calculated pairwise Jaccard distances using a binary matrix of presence/absence records using the *proxy* package in R v.4.3.0 (Meyer and Buchta, 2022; R Core Team, 2023). Principal components analysis was performed using the native R *stats* package and visualised using *ggplot2* (R Core Team, 2023; Wickham, 2016). Springs without occurrence records were excluded from the analysis, as were taxa known to occur in SA GAB springs generally but without specific location information. Finally, we produced rarefaction curves using the R package iNEXT (Hsieh and Chao, 2022) from our entire dataset and a subset of the data considering only confident records. Rarefaction was performed using the rarefaction and extrapolation models for species richness ($q = 0$), 95% confidence intervals, and 100 replications.

3 Results

3.1 Biodiversity of the SA GAB

The database we compiled based on our comprehensive literature review captured

3,463 occurrence records corresponding to 495 distinct and putatively distinct taxa. Of

these records, 2,300 were considered coarse—i.e., corresponding to the supergroup or

complex level, but not informing the presence of taxa at a specific spring group.

Invertebrates were by far the most speciose group in the dataset (42%, 207 taxa)

followed by vascular plants (22%, 111 taxa), vertebrates (21%, 102 taxa), algae (14%,

68 taxa; for the purposes of this paper, includes green, golden, and blue-green algae)

non-vascular plants (1%, 5 taxa), and fungi (0.4%, 2 taxa) (Figure 2). Sixty-five taxa

(13%) are known only from SA GAB springs, almost all of which are invertebrates.

Despite the species diversity dominance of invertebrates, these organisms are also the

most poorly known in GAB springs from a taxonomic standpoint. Just over one-third of

invertebrate taxa have formal taxonomic names (37%), whereas the remainder of the

fauna is either undescribed (14%) or has an unknown taxonomic status due to a lack of

species-level occurrence records (49%). For other taxonomic groups, taxonomic

description ranges from 50–100% (described), 0–1.5% (undescribed), and 0–50%

(unidentified). Further, apart from the Gastropoda, no invertebrate taxon has had its

conservation status assessed at the global, federal, or state level (Figure 2). Fifty-eight

of the 170 spring groups assessed in this study had no corresponding occurrence

records in the literature. A rarefaction curve derived from the dataset suggests the

artesian wetlands of SA have not been adequately surveyed, and if additional locations

were examined with equal sampling intensity, dozens of additional taxa would likely be

documented (Figure 3).

In addition to occurrence records in the dataset, we also identified several

records of potential local extinctions in the literature (Table 1). In all, we retrieved

absence records corresponding to one isopod taxon, which may represent multiple

species (*Phreatomerus latipes* (Chilton, 1922) Central, North, South haplotypes)

(Fensham et al., 2010; Guzik et al., 2012; Kinhill, 1997; Kinhill-Stearns, 1984), one

ostracod (*Ngarawa dirga* De Deckker, 1979) (Fensham et al., 2010; Kinhill, 1997;

McLaren et al., 1985), 15 snail taxa (*Fonscochlea accepta*, members of *F. aquatica*, *F.*

billakalina, *F. variabilis*, *Trochidrobia punicea*, *T. smithi* species complexes [all

Ponder, Hershler & Jenkins, 1989], *Sinumelon pedasum* Iredale, 1937) (Department of

the Environment, 2022; Fensham et al., 2010; Ponder et al., 1995; Rossini et al., 2018;

Zeidler and Ponder, 1989), and four fishes (the Dalhousie goby *Chlamydogobius*

gloveri Larson, 1995, Lake Eyre hardyhead *Craterocephalus eyresii* (Steindachner,

1883), spangled perch *Leiopotherapon unicolor* (Günther, 1859), Dalhousie gudgeon

Mogurnda thermophila Allen & Jenkins, 1999) (Department of the Environment, 2022;

Froese and Pauly, 2023; Gotch et al., 2016; Kodric-Brown et al., 2007; Rossini et al.,

2018; Zeidler and Ponder, 1989). To the best of our knowledge, these records

corresponded to likely local extinctions as opposed to complete species extinctions

(i.e., relevant taxa were present in at least one additional location). These likely

extirpations have occurred across 13 spring groups in the Dalhousie and Kati Thanda–

Lake Eyre supergroups (Table 1). Given the patchy sampling of GAB springs during

many surveys, we note that these data may be a result of stochastic sampling effort

rather than true extirpations and is therefore an opportunity for future research.

However, we note for certain taxa this evidence is more robust than others, e.g. the

discovery of empty snail shells at springs (Ponder et al. 1989).

Figure 2: Fauna, flora, and funga associated with SA GAB spring wetlands, with major organismal groups indicated by grey boxes. Groups lacking species-level records have an unknown conservation status and are indicated with n.d. (no data). * Rare and Extinct are specific to SA/TAS and NSW legislation, respectively, and only refer to populations/taxa within those states. Silhouette credits Maxime Dahirel, Armelle Ansart, Mathieu Pélissié, and Lafage via PhyloPic.

Figure 3: Extrapolation curves predicting an increase in taxon richness if additional spring groups were surveyed, calculated from all occurrence records in the dataset (A) and a subset of the dataset containing only confident records (B). As the curve does not flatten at the maximum number of known spring groups in the state (170), an increase in sampling effort *within* springs (as opposed to the sole sampling of additional springs) may also capture a more comprehensive representation of these communities. Dotted lines represent extrapolation beyond the taxon richness of our dataset (solid lines); shaded areas represent 95% confidence intervals.

3.2 Biodiversity metrics and community composition

Taxon richness was more heavily impacted by the exclusion of coarse occurrence records—as most data in this category are not spring endemics—whereas endemicity rankings were essentially unchanged between spring locations. Complexes may also be taxon-rich without containing a large number of endemics (e.g., groups within the Francis Swamp complex; Figure 4) and vice versa (e.g., Beresford Hill). Groups with high levels of richness values generally corresponded to the Coward, Dalhousie, Billa Kalina, and Hermit Hill complexes and those containing large numbers of isolated endemics included Dalhousie, Francis Swamp, Mount Denison, and Coward (Figure 4; (Beasley-Hall et al., 2023b) for raw research data via FigShare).

While there is some degree of overlap regarding taxon composition of spring

complexes, particularly in the Kati Thanda–Lake Eyre region (Figure 5), locations with

similar biodiversity metrics do not necessarily support the same biotic communities.

The Peake Creek, Lake Cadibarrawirracanna, Neales River, and Mount Dutton

complexes of the Kati Thanda–Lake Eyre supergroup are generally non-overlapping

and groups from within the Dalhousie complex/supergroup are easily differentiated

from other complexes (Figure 5). In contrast, groups from the Munda / Lake Frome

complex are nested within those from the Kati Thanda–Lake Eyre supergroup. Groups

belonging to the same complex are also generally more similar to one another than

those in different complexes. This apparent lack of connectivity is perhaps unsurprising

given the aridity of the surrounding landscape. Overall, spring groups within the

Dalhousie supergroup/complex are the most distinct in the dataset in that they support

taxa not known to be associated with other spring complexes, including highly isolated

endemics (Figures 4, 5).

Figure 4: Rankings of endemicity and overall biodiversity in SA GAB spring groups, with circle size indicating a greater value for a given metric. Larger circles indicate a higher proportion of endemics and/or more isolated endemic fauna (endemicity ranking, yellow) or a greater number of putative species (taxon richness, purple). Grey circles represent the approximate area of spring groups as shown in Figure 1. This visualisation was calculated from both coarse and confident records; raw data, including metrics calculated from only "confident" records, are available in Supplementary File S2.

4   Discussion

Here, we present a robust literature review of the biota associated with artesian

wetlands of South Australia. These data highlight several key trends: 1) invertebrates

are a poorly known component of GAB biodiversity relative to the rest of the biota; 2)

the composition of GAB biota differs considerably by location, challenging previous

conceptions of relevant biodiversity hotspots; and 3) the true extent of GAB

biodiversity is clearly far greater than is currently appreciated in the literature. We

discuss each of these findings below, as well as conservation implications for GAB

springs more broadly and how a centralised biodiversity resource for the GAB could be
implemented in future.
467

4.1   Biodiversity of South Australia's GAB springs
Almost 500 taxa are associated with GAB springs and their surrounding wetlands.
Invertebrates represent the largest proportion of this biota and the largest group of
endemics, with the majority of being insects, crustaceans, and arachnids (Figure 2).
Invertebrates are poorly documented from biodiversity, ecological, and conservation
standpoints relative to the remainder of GAB biota and are also overrepresented in
extinction records (Table 1). These groups may therefore be the most at risk of decline
due to difficulties in devising management strategies. In contrast, while vertebrates
were the third-most speciose group associated with the system (N = 102), almost all
were widespread species not of conservation concern (Fensham and Price, 2004)
(Figure 2). However, we note here that vertebrates such as waterbirds are nonetheless
dependent on the springs as breeding grounds (Badman, 1985), and may make more
extensive use of these habitats in non-flood years and/or as "stepping stone" habitats to
sustain migrations to nearby Kati Thanda–Lake Eyre. For example, the little grassbird
Poodytes gramineus (Gould, 1845) is not restricted to GAB springs, but nonetheless
has an obligate relationship with stands of the reed *Phragmites australis* in GAB
wetlands (Read, 1997).
485

Our data suggest a high degree of heterogeneity in the ecological communities
of the SA GAB and challenge past assumptions regarding the system's biodiversity
hotspots. It has long been recognised that certain SA GAB springs are more speciose
than others, but only recently has a quantitative view of candidate biodiversity hotspots
been put forward (Rossini et al., 2018), and without consideration of non-endemic taxa.
Springs at Dalhousie are one such hotspot proposed by Rossini *et al.* (2018), which we
support in our review here. We also propose the Francis Swamp, Mount Denison, and
Coward complexes as hotspots due to their contribution to the remaining bulk of
endemicity in the SA GAB. Indeed, these springs house a much more speciose and
isolated endemic biota than initially recognised based on published records
(Supplementary File S1). Yet, we note that not all taxon-rich locations are necessarily
rich in endemic species (Figure 4). Overall, our findings indicate the disturbance or

extinction of any given spring group could represent an enormous loss of biodiversity

when both endemic and non-endemic taxa are considered together. The biodiversity of

the South Australian GAB has not been adequately surveyed (Figure 3) and over two-

thirds of records we collated were coarse, meaning they lacked occurrence information

at the spring group level. While records are helpful in distinguishing spring group

locations and suggest considerable structure in taxa occupying different spring

complexes and supergroups (Figure 5), they nonetheless highlight a lack of detailed

past surveys and the sparseness of published information on GAB biodiversity.

The observed variability in the dataset may also be associated with sampling

bias. Spring groups with the highest endemicity rankings, such as those within the

Dalhousie and Coward complexes, are also those consistently associated with high

sampling effort in previous surveys (Badman, 1985; Greenslade, 1985; Kovac, 2003;

Mitchell, 1985; Noack, 1994; Ponder et al., 1995; Sokol, 1987; Zeidler and Ponder,

1989). The Kati Thanda–Lake Eyre and Dalhousie supergroups corresponded to over

2,000 occurrence records, whereas observations within the Munda / Lake Frome

supergroup were restricted to a single, coarse occurrence record of the copepod

Microcyclops dengizicus (Lepeschkin, 1900) (Zeidler and Ponder, 1989). Munda / Lake

Frome has clearly not been adequately sampled, although this does not necessarily

imply the location lacks endemic taxa yet to be characterised (McLaren et al., 1985;

Rossini, 2020). Indeed, widespread taxonomic groups expected to be present in springs,

such as the Hymenoptera and other cosmopolitan insect groups, were also poorly

represented in the dataset due to a presumed bias in sampling methods and a lack of

past taxonomic expertise. As such, we caution against the derivation of conservation

priorities from locations with high measures of metrics such as species richness alone,

but trends are nonetheless evident in this review regarding spring locations and

taxonomic groups that have been *under*sampled.

4.2   Conservation implications for South Australia's artesian springs

At least 14 taxa may have become locally extinct in the SA GAB springs (Table 1).

Evidence for extirpations includes survey work conducted prior to the extinction of

springs themselves or, conversely, the discovery of remains at extinct springs (e.g.,

snail shells). The bulk of these likely extirpations have occurred in spring groups that

have ceased to flow because of drawdown, namely Venable, Palura Pintjanha /

Priscilla, Marrinha / Hergott, Margaret, and Manda-wardunha / Mundowdna (Fensham

et al., 2010). The Venable and Palura Pintjanha / Priscilla spring groups became extinct

around 1990 following predictions by industry that mining activities would lead to a

partial, if not complete, reduction in artesian flow at certain GAB springs (Kinhill-

Stearns, 1983, 1982). Marrinha / Hergott ceased to flow in the mid-1980s as water was

withdrawn to supply the nearby town of Marree (McLaren et al., 1985, p. 198). Other

extirpations have been attributed to human modification of springs, the presence of

overabundant invasive and native species, and potentially insufficient sampling effort.

The apparent local extinction of the fish species, Lake Eyre hardyhead

(*Craterocephalus eyresii*) at Toondina and Old Nilpinna may have been caused by

spring excavations and competition from the invasive mosquitofish *Gambusia*

holbrooki Girard, 1859, respectively (Gotch et al., 2016). Extirpations of other fish

species at Dalhousie have been attributed to the overgrowth of the native reed

Phragmites as a result of decreased grazing pressure following the exclusion of

livestock (Kodric-Brown et al., 2007). We also note that there are active springs at

which only dead specimens of species have been collected, pointing to potential

additional extirpations: for example, in certain spring groups within the Neales River

and Francis Swamp complexes only empty shells of *Fonscochlea zeidleri* have been

collected, with no evidence of live individuals (Ponder et al., 1989).

It is important to establish conservation priorities given the potential for further

local, if not species-wide, extinctions in the SA GAB springs. Here, we ranked spring

groups by the number of endemic species supported by each spring group and their

degree of isolation. High-ranking locations mirror those proposed as conservation

priorities in past studies, corroborating our findings. For example, McLaren *et al.*

(1985) produced an inventory of fauna, flora, and funga to set out conservation

priorities based on species diversity, the presence of rare species, "naturalness" (i.e.,

extent of interference by humans and cattle grazing pressure) and perceived

vulnerability to degradation. Spring groups were then ranked by conservation priority.

High-ranking spring groups in this assessment include those found at Dalhousie and in

the Coward and Mt. Denison complexes of the Kati Thanda–Lake Eyre supergroup.

Further, Rossini *et al.* (2018) used a ranking system developed by Fensham and Price

(2004) (modified here to produce endemicity rankings) to conclude that groups within

the Dalhousie, Strangways, Francis Swamp, Billa Kalina, and Mt Denison complexes

were of high conservation significance relative to other springs. The highest-ranking

spring groups by endemicity in the current study presented here corresponded to all six

of the above complexes, often down to the spring group level (Supplementary File S2).

We stress that these trends do not suggest certain spring groups are more important

than others or that insignificant springs exist, as has been implied in past environmental

impact statements (Keane, 1997; Kinhill-Stearns, 1983). This is especially the case for

locations that could be perceived as having low biological importance, such as the

extinct Papu-ngaljuru / Primrose spring group, but have outstanding cultural

significance to Aboriginal peoples due to associated Dreaming stories and its use as a

major occupation site (Hercus and Sutton, 1985). Instead, our findings suggest that

certain spring groups harbour high numbers of endemics relative to other locations (but

remembering that insufficient sampling effort has occurred) and require targeted

conservation efforts to preserve these short-range taxa.

Several conservation programs are currently in place which aim to directly

improve the condition of SA GAB spring wetlands (Harris, 1992). Preliminary

evidence suggests several of these practices have had a positive impact on SA GAB

spring endemics. Springs are threatened by the presence of large-bodied and hard-

hooved livestock which can graze on or trample wetland vegetation, foul the water

through their faeces, and potentially severely reduce invertebrate biodiversity if cattle

stocking levels are high; almost all SA GAB springs are subject to such pressure,

particularly from cattle (Fensham et al., 2010; Gotch et al., 2016; Hutchinson and King,

1980; Kovac and Mackay, 2009). A combination of fencing and destocking has been

successful in restoring endemic biodiversity of SA GAB springs in some cases, e.g., the

endemic salt pipewort *Eriocaulon carsonii* subsp. *carsonii* F. Muell. in the Hermit Hill

complex (Fatchen and Fatchen, 1993) and aquatic invertebrates at springs that were

heavily damaged by stock (Kinhill-Stearns, 1984). The spring groups identified here as

harbouring a large number of endemic species could represent future candidates for

fencing and/or destocking initiatives.

As stated above, there is a large proportion of endemic taxa at several

complexes within the SA GAB, namely those contained within the Dalhousie and Kati

Thanda–Lake Eyre supergroups. The data presented here reiterates the need for

targeted surveys in these locations to not only gather additional biodiversity data, but
also to develop an understanding of population dynamics, habitat requirements, and
accidental human-mediated translocations of fauna, flora, and funga per the National
Recovery Plan (Fensham et al., 2010). The plan notably highlights the fact that the
construction of a robust biodiversity inventory for GAB springs has been hindered by a
lack of survey effort and taxonomic expertise. The utility of emerging environmental
DNA techniques to capture a holistic picture of SA GAB spring biodiversity in a non-
invasive manner would overcome several of these limitations (Beasley-Hall et al.,
2023c; Saccò et al., 2022; Vörös, 2017; West et al., 2020); in this regard, the database
developed as part of this study would assist in the selection of fieldwork locations for a
pilot study in this regard. Once taxa are established as occurring within the SA GAB,
monitoring efforts should be undertaken on a regular basis to develop deeper
understandings of ecological knowledge as opposed to presence/absence information.
612

Although taxa reliant on GAB-fed springs are protected as a single ecological
community under the EPBC Act, species-level listings are also far overdue. This is
particularly the case for invertebrates dependent on the GAB. The majority of GAB
molluscs and crustaceans meet the criteria to be listed as Critically Endangered
(Rossini, 2020), but almost 90% of SA GAB invertebrates recorded here lack
conservation listing at a global, federal, or state level, the bulk of which are either new
species awaiting formal description or those completely lacking species-level
identifications. This large knowledge gap is only exacerbated by a lack of
understanding of the relationship between spring characteristics and biodiversity
metrics (Fensham and Laffineur, 2022; Rossini et al., 2018) and indeed, relationships
among spring ecological and hydrogeological characteristics themselves (Green and
Berens, 2013; Love et al. (eds), 2013; Mudd, 2000). We hope that resources like the
database developed here will inform the prioritisation of certain spring locations and
taxonomic groups, ultimately leading to the listing of invertebrates and other spring
endemics under relevant legislation on a per-species basis.
628

629 **5   Conclusion**
630 Here, we have presented the first review of biota supported by the Great Artesian Basin
631 in South Australia. This dataset is an important resource we hope will facilitate future

studies on GAB spring endemics, investigations into their population dynamics, basic biology, and taxonomy, and ultimately facilitate the listing of relevant taxa under state and federal environmental legislation. Such resources are essential given the multitude of threats currently facing springs and their biota, with potential extirpations of populations of at least 13 species in the SA GAB springs to date. We have also highlighted springs of particular conservation concern which may assist in determining future conservation priorities for this system. Our dataset has highlighted three major points about GAB spring biodiversity. Firstly, the majority of taxa reliant on these wetlands are invertebrates, and these animals are also the most poorly known and conserved in the GAB. Secondly, ecological communities reliant on artesian springs in SA are largely non-overlapping, irrespective of whether only endemic taxa, or endemics as well as "incidental" species, are concerned. The extinction or considerable disturbance of any spring group is therefore likely to lead to a considerable loss of biodiversity and/or genetic diversity. Finally, the artesian springs of SA have not been adequately sampled by past survey efforts and a considerable proportion of taxa likely remain to be documented, particularly in understudied locations such as the Munda / Lake Frome supergroup. There also remains a dearth of occurrence records for certain taxa for which taxonomic expertise has been lacking in the past, such as the wasps and allies (Greenslade, 1985). We recommend that datasets such as these are made publicly available with the capacity to be modified and updated on an ongoing basis, embodying the gold standard of digital asset storage.

6 Statements and Declarations

6.1 Data Availability Statement

All raw data associated with this paper are available via FigShare (Beasley-Hall et al., 2023a, 2023b).

6.2   Conflicts of interest

The authors wish to disclose that authors PGBH and MTG, independent researchers affiliated with The University of Adelaide and South Australian Museum, received financial support from industry stakeholders when conducting this study. This funding did not influence the design, data collection, analysis, or reporting of this study. Research findings and conclusions expressed in this publication are based solely on the analysis of the data and the scientific merit of the paper.

6.3   Declaration of funding

Funding for PGBH was provided by contract research via BHP Group Ltd. Funding for BAH was provided by an Australian Government Research Training Program (RTP) Scholarship, the Roy and Marjory Edwards Scholarship as administered by Nature Foundation, and the Justin Costelloe Scholarship as administered by the Kati Thanda—Lake Eyre Basin Authority, Department of Agriculture, Water and the Environment. Funding for MTG was provided by the Australian Research Council (grant LP190100555) in partnership with Curtin University, The University of Adelaide, BHP Group Ltd., Rio Tinto Ltd., Chevron Australia Pty Ltd., Western Australian Museum, South Australian Museum, the Department for Biodiversity, Conservation and Attractions (WA), the Western Australian Biodiversity Science Institute, Department of Water and Environmental Regulation (WA).

6.4   Acknowledgements

The authors would like to acknowledge the Arabana Aboriginal Corporation, Travis Gotch, and The Friends of Mound Springs for their guidance and feedback.

7   References

Andersen, M., Barron, O., Bond, N., Burrows, R., Eberhard, S., Emelyanova, I., Fensham, R., Froend, R., Kennard, M., Marsh, N., Pettit, N., Rossini, R., Rutlidge, R., Valdez, D., Ward, D., 2016. Research to inform the assessment of ecohydrological responses to coal seam gas extraction and coal mining. Department of the Environment and Energy, Commonwealth of Australia.

Arabana Aboriginal Corporation, 2021. Submission to the inquiry into the destruction of 46,000-year-old caves at the Juukan Gorge in the Pilbara Region of Western Australia. https://www.aph.gov.au/DocumentStore.ashx?id=41e65957-4b2f-4245-bc55-f0198241c6ce&subId=690905, Canberra, Australia.

Australian Government, 2023. Australian place names | Style Manual. URL

https://www.stylemanual.gov.au/grammar-punctuation-and-conventions/names-and-
terms/australian-place-names (accessed 5.23.23).

Badman, F.J., 1985. Birds of the mound springs and bores south and west of Lake Eyre with
special reference to the Coward Spring area, in: South Australia's Mound Springs.
Nature Conservation Society of South Australia Inc., Hyde Park Press, Plympton,
South Australia.

Beasley-Hall, P.G., Hedges, B.A., Cooper, S.J.B., Austin, A.D., Guzik, M.T., 2023a. Database
collating information on the fauna, flora, and funga of South Australia's Great Artesian
Basin springs. https://doi.org/10.25909/24457105

Beasley-Hall, P.G., Hedges, B.A., Cooper, S.J.B., Austin, A.D., Guzik, M.T., 2023b. Species
richness and endemicity rankings for each South Australian spring group.
https://doi.org/10.25909/24457120

Beasley-Hall, P.G., Murphy, N.P., King, R.A., White, N.E., Hedges, B.A., Cooper, S.J.B.,
Austin, A.D., Guzik, M.T., 2023c. Time capsules of biodiversity: Future research
directions for groundwater-dependent ecosystems of the Great Artesian Basin. Front.
Environ. Sci. 10.

Belbin, L., Wallis, E., Hobern, D., Zerger, A., 2021. The Atlas of Living Australia: History,
current state and future directions. Biodivers. Data J. 9, e65023.
https://doi.org/10.3897/BDJ.9.e65023

Brack, W., Altenburger, R., Schüürmann, G., Krauss, M., López Herráez, D., van Gils, J.,
Slobodnik, J., Munthe, J., Gawlik, B.M., van Wezel, A., Schriks, M., Hollender, J.,
Tollefsen, K.E., Mekenyan, O., Dimitrov, S., Bunke, D., Cousins, I., Posthuma, L., van
den Brink, P.J., López de Alda, M., Barceló, D., Faust, M., Kortenkamp, A.,
Scrimshaw, M., Ignatova, S., Engelen, G., Massmann, G., Lemkine, G., Teodorovic, I.,
Walz, K.-H., Dulio, V., Jonker, M.T.O., Jäger, F., Chipman, K., Falciani, F., Liska, I.,
Rooke, D., Zhang, X., Hollert, H., Vrana, B., Hilscherova, K., Kramer, K., Neumann,
S., Hammerbacher, R., Backhaus, T., Mack, J., Segner, H., Escher, B., de Aragão
Umbuzeiro, G., 2015. The SOLUTIONS project: challenges and responses for present
and future emerging pollutants in land and water resources management. Sci. Total
Environ. 503–504, 22–31. https://doi.org/10.1016/j.scitotenv.2014.05.143

Department of Climate Change, Energy, the Environment and Water, 2023. Species Profile and
Threats Database. URL http://www.environment.gov.au/cgi-bin/sprat/public/sprat.pl
(accessed 4.26.23).

Department of the Environment, 2022. The community of native species dependent on natural
discharge of groundwater from the Great Artesian Basin in Community and Species
Profile and Threats Database. Australian Government Department of the Environment,
Canberra. Available from: http://www.environment.gov.au/sprat.

Fairfax, R. j., Fensham, R. j., 2002. In the Footsteps of J. Alfred Griffiths: a Cataclysmic
History of Great Artesian Basin Springs in Queensland, Australia. Aust. Geogr. Stud.
40, 210–230. https://doi.org/10.1111/1467-8470.00175

Famiglietti, J.S., 2014. The global groundwater crisis. Nat. Clim. Change 4, 945–948.
https://doi.org/10.1038/nclimate2425

Fatchen, T.J., 2000. Mound springs management planning: management issues, strategies, and
prescriptions for mound springs in Far North South Australia. Report prepared for the
South Australian Department of Environment and Heritage with the support of the
Natural Heritage Trust.

Fatchen, T.J., Fatchen, D.H., 1993. Dynamics of vegetation on mound springs in the Hermit
Hill region, northern South Australia. Prepared for WMC (Olympic Dam Operations)
Pty. Ltd., TJ Fatchen & Associates, Adelaide.

Fensham, R. j., Silcock, J. l., Powell, O., Habermehl, M. a., 2016. In Search of Lost Springs: A
Protocol for Locating Active and Inactive Springs. Groundwater 54, 374–383.
https://doi.org/10.1111/gwat.12375

Fensham, R.J., Laffineur, B., 2022. Response of spring wetlands to restored aquifer pressure in
the Great Artesian Basin, Australia. J. Hydrol. 612, 128152.
https://doi.org/10.1016/j.jhydrol.2022.128152

Fensham, R.J., Ponder, W.F., Fairfax, R.J., 2010. Recovery plan for the community of native
species dependent on natural discharge of groundwater from the Great Artesian Basin.
Report to Department of the Environment, Water, Heritage and the Arts, Canberra.
Queensland Department of Environment and Resource Management, Brisbane.
Fensham, R.J., Price, R.J., 2004. Ranking spring wetlands in the Great Artesian Basin of
Australia using endemicity and isolation of plant species. Biol. Conserv. 119, 41–50.
https://doi.org/10.1016/j.biocon.2003.10.019
Froese, R., Pauly, D., 2023. FishBase. URL www.fishbase.se (accessed 5.22.23).
Gotch, T., 2013. Spatial survey of springs, in: Allocating Water and Maintaining Springs in the
Great Artesian Basin, Volume IV: Spatial Survey and Remote Sensing of Artesian
Springs of the Western Great Artesian Basin. National Water Commission, Canberra.
Gotch, T., Keppel, M., Fels, K., McKenzie, J., 2016. Lake Eyre Basin Springs Assessment:
Ecohydrological conceptual models of springs in the western Lake Eyre Basin, South
Australia. DEWNR Tech. Rep. 201602 Gov. S. Aust. Dep. Environ. Water Nat.
Resour. Adel.
Gotch, T.B., Adams, M., Murphy, N.P., Austin, A.D., Gotch, T.B., Adams, M., Murphy, N.P.,
Austin, A.D., 2008. A molecular systematic overview of wolf spiders associated with
Great Artesian Basin springs in South Australia: evolutionary affinities and an
assessment of metapopulation structure in two species. Invertebr. Syst. 22, 151–165.
https://doi.org/10.1071/IS07045
Government of South Australia, 2023. Groundwater Data. WaterConnect. URL
https://www.waterconnect.sa.gov.au/Systems/GD/Pages/Default.aspx (accessed
4.20.23).
Great Artesian Basin Coordinating Committee, 2019. Great Artesian Basin Strategic
Management Plan 2019.
Green, G., Berens, V., 2013. Relationship between aquifer pressure changes and spring
discharge rates, in: Allocating Water and Maintaining Springs in the Great Artesian
Basin, Volume III: Groundwater Discharge of the Western Great Artesian Basin.
National Water Commission, Canberra.
Greenslade, P., 1985. Terrestrial invertebrates of the mound springs, bores, creek beds, and
other habitats, in: South Australia's Mound Springs. Nature Conservation Society of
South Australia Inc., Hyde Park Press, Plympton, South Australia.
Guzik, M.T., Adams, M.A., Murphy, N.P., Cooper, S.J.B., Austin, A.D., 2012. Desert Springs:
Deep Phylogeographic Structure in an Ancient Endemic Crustacean (Phreatomerus
latipes). PLOS ONE 7, e37642. https://doi.org/10.1371/journal.pone.0037642
Guzik, M.T., Stringer, D.N., Murphy, N.P., Cooper, S.J.B., Taiti, S., King, R.A., Humphreys,
W.F., Austin, A.D., Guzik, M.T., Stringer, D.N., Murphy, N.P., Cooper, S.J.B., Taiti,
S., King, R.A., Humphreys, W.F., Austin, A.D., 2019. Molecular phylogenetic analysis
of Australian arid-zone oniscidean isopods (Crustacea□: Haloniscus) reveals strong
regional endemicity and new putative species. Invertebr. Syst. 33, 556–574.
https://doi.org/10.1071/IS18070
Habermehl, M.A., 2020. Review: The evolving understanding of the Great Artesian Basin
(Australia), from discovery to current hydrogeological interpretations. Hydrogeol. J.
28, 13–36. https://doi.org/10.1007/s10040-019-02036-6
Harris, C.R., 1992. Mound Springs: South Australian Conservation Initiatives. Rangel. J. 14,
157–173. https://doi.org/10.1071/rj9920157
Harvey, M.S., 1990. A review of the water mite family Anisitsiellidae in Australia (Acarina).
Invertebr. Syst. 3, 629–646. https://doi.org/10.1071/it9890629
Hercus, L., Sutton, P., 1985. The assessment of Aboriginal cultural significance of mound
springs in South Australia, in: Heritage of the Mound Springs. Department of
Environment and Planning, South Australia.
Hsieh, T.C., Chao, K.H.M. and A., 2022. iNEXT: Interpolation and Extrapolation for Species
Diversity.
Hutchinson, K.J., King, K.L., 1980. The Effects of Sheep Stocking Level on Invertebrate
Abundance, Biomass and Energy Utilization in a Temperate, Sown Grassland. J. Appl.

Ecol. 17, 369–387. https://doi.org/10.2307/2402333

Keane, D., 1997. The sustainability of use of groundwater from the Great Artesian Basin with
particular reference to the south-western edge of the basin and impact on the Mound
Springs. Unpublished thesis, RMIT University.

Kerezsy, A., 2015. Development of a technique for quarantining Great Artesian Basin springs
from colonisation by the invasive fish Eastern Gambusia (Gambusia holbrooki). Ecol.
Manag. Restor. 16, 229–232. https://doi.org/10.1111/emr.12174

King, R., 2009. Two New Genera And Species Of Chiltoniid Amphipods (Crustacea:
Amphipoda: Talitroidea) From Freshwater Mound Springs In South Australia. Zootaxa
2293, 35–52. https://doi.org/10.5281/zenodo.191467

King, R., Leys, R., King, R.A., Leys, R., 2014. Molecular evidence for mid-Pleistocene
divergence of populations of three freshwater amphipod species (Talitroidea□:
Chiltoniidae) on Kangaroo Island, South Australia, with a new spring-associated genus
and species. Aust. J. Zool. 62, 137–156. https://doi.org/10.1071/ZO13099

Kinhill, 1997. Olympic Dam Expansion Project: Environmental impact statement. Prepared for
WMC (Olympic Dam Corporation) Pty. Ltd. Olympic Dam, South Australia.

Kinhill-Stearns, 1984. Olympic Dam Project supplementary environmental studies, Mound
Springs. Adelaide, Roxby Management Services Pty Ltd.

Kinhill-Stearns, 1983. Olympic Dam Project Supplementary to the Draft Environmental Impact
Statement. Prepared for Roxby Management Services Pty Ltd.

Kinhill-Stearns, 1982. Olympic Dam Project Draft Environmental Impact Statement. Prepared
for Roxby Management Services Pty Ltd.

Kodric-Brown, A., Wilcox, C., Bragg, J.G., Brown, J.H., 2007. Dynamics of Fish in Australian
Desert Springs: Role of Large-Mammal Disturbance. Divers. Distrib. 13, 789–798.

Kovac, K.-J., 2003. Bird surveys at Dalhousie Springs, in: Expedition Witjira: Interim Report,
12th - 26th July 2003. Scientific Expedition Group with National Parks and Wildlife
and Department of Environment and Heritage, Unley, South Australia.

Kovac, K.-J., Mackay, D.A., 2009. An experimental study of the impacts of cattle on spider
communities of artesian springs in South Australia. J. Insect Conserv. 13, 57–65.
https://doi.org/10.1007/s10841-007-9117-2

Lewis, M.M., White, D., 2013. Evaluation of remote sensing approaches, in: Volume IV:
Spatial Survey and Remote Sensing of Artesian Springs of the Western Great Artesian
Basin. National Water Commission, Commonwealth of Australia, Canberra.

Lewis, M.M., White, D., Gotch, T., 2013. Allocating Water and Maintaining Springs in the
Great Artesian Basin, Volume IV: Spatial Survey and Remote Sensing of Artesian
Springs of the Western Great Artesian Basin. National Water Commission, Canberra.

Lewis, S., Harris, C., 2020. Improving conservation outcomes for Great Artesian Basin springs
in South Australia. Proc. R. Soc. Qld. 126, 271–287.
https://doi.org/10.3316/informit.626794538781042

Love et al. (eds), 2013. Allocating Water and Maintaining Springs in the Great Artesian Basin,
Volume III: Groundwater Discharge of the Western Great Artesian Basin. National
Water Commission, Canberra.

McLaren, N., Wiltshire, D., Lesslie, R., 1985. Biological assessment of South Australian
mound springs, in: Heritage of the Mound Springs. Unpublished report prepared for
The South Australian Department of Environment and Planning, South Australia.

Meyer, D., Buchta, C., 2022. proxy: Distance and Similarity Measures.

Mitchell, B.D., 1985. Limnology of mound springs and temporary pools south and west of
Lake Eyre, in: South Australia's Mound Springs. Nature Conservation Society of South
Australia Inc., Hyde Park Press, Plympton, South Australia.

Mudd, G.M., 2000. Mound springs of the Great Artesian Basin in South Australia: a case study
from Olympic Dam. Environ. Geol. 39, 463–476.
https://doi.org/10.1007/s002540050452

Murphy, N.P., Adams, M., Austin, A.D., 2009. Independent colonization and extensive cryptic
speciation of freshwater amphipods in the isolated groundwater springs of Australia's
Great Artesian Basin. Mol. Ecol. 18, 109–122. https://doi.org/10.1111/j.1365-

294X.2008.04007.x

Murphy, N.P., Adams, M., Guzik, M.T., Austin, A.D., 2013. Extraordinary micro-endemism in
Australian desert spring amphipods. Mol. Phylogenet. Evol. 66, 645–653.
https://doi.org/10.1016/j.ympev.2012.10.013

Murphy, N.P., Breed, M.F., Guzik, M.T., Cooper, S.J.B., Austin, A.D., 2012. Trapped in desert
springs: phylogeography of Australian desert spring snails. J. Biogeogr. 39, 1573–
1582. https://doi.org/10.1111/j.1365-2699.2012.02725.x

Murphy, N.P., Guzik, M.T., Cooper, S.J.B., Austin, A.D., 2015. Desert spring refugia:
museums of diversity or evolutionary cradles? Zool. Scr. 44, 693–701.
https://doi.org/10.1111/zsc.12129

New South Wales Department of Planning, Industry and Environment, 2021. Hydrogeology
and ecology survey of the Great Artesian Basin springs in NSW — Survey
methodology and results. NSW Government.

Noack, D., 1994. Date palms and the exotic flora of Dalhousie Springs. South Aust. Geogr. J.
93, 81–89.

Obura, D.O., Aeby, G., Amornthammarong, N., Appeltans, W., Bax, N., Bishop, J., Brainard,
R.E., Chan, S., Fletcher, P., Gordon, T.A.C., Gramer, L., Gudka, M., Halas, J., Hendee,
J., Hodgson, G., Huang, D., Jankulak, M., Jones, A., Kimura, T., Levy, J., Miloslavich,
P., Chou, L.M., Muller-Karger, F., Osuka, K., Samoilys, M., Simpson, S.D., Tun, K.,
Wongbusarakum, S., 2019. Coral Reef Monitoring, Reef Assessment Technologies,
and Ecosystem-Based Management. Front. Mar. Sci. 6.

Ordens, C., McIntyre, N., Underschultz, J., Ransley, T., Moore, C., Mallants, D., 2020. Preface:
Advances in hydrogeologic understanding of Australia's Great Artesian Basin.
Hydrogeol. J. Online, 1–11. https://doi.org/10.1007/s10040-019-02107-8

Ponder, W. F., Hershler, R., Jenkins, B. J. 1989. An endemic radiation of hydrobiid snails from
artesian springs in northern South Australia: their taxonomy, physiology, distribution
and anatomy. Malacologia 31, 1–140.

Ponder, W.F., Eggler, P., Colgan, D.J., 1995. Genetic differentiation of aquatic snails
(Gastropoda: Hydrobiidae) from artesian springs in arid Australia. Biol. J. Linn. Soc.
56, 553–596. https://doi.org/10.1111/j.1095-8312.1995.tb01110.x

Ponder, W. F. 2022. The first galeommatoidean from inland waters: a new species of Arthritica
(Bivalvia, Lasaeidae) from artesian springs in South Australia, with comments on other
Australian members of the genus. Molluscan Research 43, 22–33.

QGIS Association, 2023. QGIS.org. QGIS Geogr. Inf. Syst. URL https://qgis.org (accessed
4.26.23).

Queensland Department of Regional Development, Manufacturing and Water, 2023.
Queensland's plan to protect our Great Artesian Basin. ArcGIS StoryMaps. URL
https://storymaps.arcgis.com/stories/259e2ef480944f799fb2465ba2821877 (accessed
4.20.23).

Queensland Government, 2018. Springs database. Open Data Portal. URL
https://www.data.qld.gov.au/dataset/springs (accessed 5.1.23).

R Core Team, 2023. R: A Language and Environment for Statistical Computing. R Foundation
for Statistical Computing, Vienna, Austria.

Read, J.L., 1997. Stranded on Desert Islands? Factors Shaping Animal Populations in Lake
Eyre South. Glob. Ecol. Biogeogr. Lett. 6, 431–438. https://doi.org/10.2307/2997352

Rossini, R.A., 2020. Current State and Reassessment of Threatened Species Status of
Invertebrates Endemic to Great Artesian Basin Springs. Proc. R. Soc. Qld. 126, 225–
248.

Rossini, R.A., Fensham, R.J., Stewart-Koster, B., Gotch, T., Kennard, M.J., 2018.
Biogeographical patterns of endemic diversity and its conservation in Australia's
artesian desert springs. Divers. Distrib. 24, 1199–1216.
https://doi.org/10.1111/ddi.12757

Saccò, M., Guzik, M.T., van der Heyde, M., Nevill, P., Cooper, S.J.B., Austin, A.D., Coates,
P.J., Allentoft, M.E., White, N.E., 2022. eDNA in subterranean ecosystems:
Applications, technical aspects, and future prospects. Sci. Total Environ. 820, 153223.

https://doi.org/10.1016/j.scitotenv.2022.153223

Sokol, A., 1987. Yabbies at Dalhousie Springs, northern South Australia: morphological
evidence for long-term isolation. Trans. R. Soc. S. Aust. Inc. 111, 207–210.

South Australian Department for Environment and Water, 2023. WaterConnect. URL
https://www.waterconnect.sa.gov.au (accessed 4.26.23).

Vaughan, H., Brydges, T., Fenech, A., Lumb, A., 2001. Monitoring Long-Term Ecological
Changes Through the Ecological Monitoring and Assessment Network: Science-Based
and Policy Relevant. Environ. Monit. Assess. 67, 3–28.
https://doi.org/10.1023/A:1006423432114

Vörös, J., 2017. Hunting the olm with eDNA. Herpetol. Rev. 48, 266.

West, K.M., Richards, Z.T., Harvey, E.S., Susac, R., Grealy, A., Bunce, M., 2020. Under the
karst: detecting hidden subterranean assemblages using eDNA metabarcoding in the
caves of Christmas Island, Australia. Sci. Rep. 10, 21479.
https://doi.org/10.1038/s41598-020-78525-6

Wickham, H., 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.

Witjira National Park Co-management Board, 2022. Witjira National Park management plan.
National Parks and Wildlife Service South Australia.

Zeidler, W., Ponder, W.F., 1989. The natural history of Dalhousie Springs. Special Publication,
South Australian Museum.

944
945
946
947

8   Tables

Table 1: Records of likely local extinctions of fauna in the SA GAB springs. To our

knowledge there are no records of floral extinctions in the SA GAB. ^ Taxon represents

multiple genetically distinct clades which we consider putative species, but which of

these were present in the below location(s) prior to their apparent extirpation is

unknown. KT–LE Sth = Kati Thanda–Lake Eyre South. * GAB spring endemic.

Taxon	Broad grouping	Spring group(s)	Spring complex(es)
Phreatomerus latipes^*	Isopod	Venable, Marrinha / Hergott	Hermit Hill, Marree
*Ngarawa dirga**	Ostracod	Venable, Manda-wardunha / Mundowdna	Hermit Hill, Marree
*Fonscochlea accepta**	Hydrobiid snail	Venable, Palura Pintjanha / Priscilla	Hermit Hill, KT–LE Sth
Fonscochlea aquatica^*	Hydrobiid snail	Margaret	Francis Swamp
*Fonscochlea billakalina**	Hydrobiid snail	Margaret	Francis Swamp
Fonscochlea variabilis^*	Hydrobiid snail	Venable, Palura Pintjanha / Priscilla	Hermit Hill, KT–LE Sth
Fonscochlea zeidleri^*	Hydrobiid snail	Venable, Palura Pintjanha / Priscilla, Centre Island	Hermit Hill, KT–LE Sth
Trochidrobia punicea^*	Hydrobiid snail	Venable, Palura Pintjanha / Priscilla	Hermit Hill, KT–LE Sth
Trochidrobia smithi^*	Hydrobiid snail	Margaret	Francis Swamp
Sinumelon pedasum	Camaenid snail	Irrwanjira / Errawanyera	Dalhousie
Chlamydogobius gloveri	Fish	Irrwanjira / Errawanyera, Frog Dreaming, Kirki / Dalhousie Proper, Cadni Dreaming	Dalhousie
Craterocephalus eyresii	Fish	Old Nilpinna, Thuntinha / Toondina	Peake Creek, Toondina
Leiopotherapon unicolor	Fish	Idnjundura / Kingfisher, Ilpikwa	Dalhousie
Mogurnda thermophila	Fish	Frog Dreaming	Dalhousie