Title: Metapopulations, the inflationary effect, and consequences for public health

Authors: Nicholas Kortessis[1], Gregory Glass[2,3], Andrew Gonzalez[4], Nick W. Ruktanonchai[5],

Margaret W. Simon[6], Burton Singer[3], and Robert D. Holt[6*]

Author Affiliations:

1. Department of Biology, Wake Forest University, Winston-Salem, NC, 27106;

2. Department of Geography, University of Florida, Gainesville, FL, 32611;

3. Emerging Pathogens Institute, University of Florida, Gainesville, FL, 32610;

4. Department of Biology, Quebec Centre for Biodiversity Science, McGill University,

Montreal, Quebec, Canada H3A 1B1;

5. Department of Population Health Sciences, Virginia Polytechnic University, Blacksburg,

VA, 24061;

6. Department of Biology, University of Florida, Gainesville, FL, 32611;

* Corresponding author; email: rdholt@ufl.edu

Author emails: NK: kortessn@wfu.edu; GG: gglass@epi.ufl.edu; AG:

andrew.gonzalez@mcgill.ca; NWR: nrukt00@vt.edu; MWS: margaret.w.simon@gmail.com;

BS: bhsinger@epi.ufl.edu; RDH: rdholt@ufl.edu;

Article Type: Special Feature

Keywords: Asynchrony; spatiotemporal heterogeneity; metapopulation; metacommunity;

inflationary effect; pandemic

Manuscript Elements: Word count: 10,134; Figures: 5; Tables: 1; Boxes 1

Author contributions: NAK and RDH conceived on the theme of the manuscript. NAK wrote

most of the initial draft, with original writing contributions from other authors. Editing of the

initial draft was done collectively by all authors.

Abstract

The metapopulation perspective is an important conceptual framework in ecology and

evolutionary ecology. Metapopulations are spatially distributed populations linked by dispersal.

Both metapopulation models and their community and ecosystem level analogues,

metacommunity and meta-ecosystem models, tend to be more stable regionally than locally and

display enhanced abundance because of the interplay of spatiotemporal heterogeneity and

dispersal (an effect that has been called the "inflationary effect"). We highlight the essential role

of spatiotemporal heterogeneity in metapopulation biology, sketch empirical demonstrations of

the inflationary effect, and provide a mechanistic interpretation of how the inflationary effect

arises and impacts population growth and abundance. We illustrate the effect with examples

from the spread of infectious disease. Namely, failure to recognize the full possible effects of

spatiotemporal heterogeneity likely enhanced the spread of COVID-19, a failure based on lack of

understanding of emergent population processes at large scales which may hamper control and

eradication of other infectious diseases. We finish by noting how the effects of spatiotemporal

heterogeneity have implicitly played roles in the history of ecology, ranging across

subdisciplines as diverse as natural enemy-victim dynamics, species coexistence, and

conservation biology. Seriously confronting the complexity of spatiotemporal heterogeneity

could push many of these subdisciplines forward.

Introduction

The concept of a metapopulation – a set of spatially distributed populations linked by dispersal – is central to discourse in ecology, evolution, and conservation biology. Metapopulation theory (Hanski and Gaggiotti, 2004)—and its community-level counterpart, metacommunity theory (Leibold, 2018)— provide a foundation for addressing important issues in ecology, including population persistence (Levins, 1969), species coexistence (Amarasekare, 2003; Hoopes et al., 2005), community assembly (Bergmann and Leveau, 2022; Leibold et al., 2022), and community stability (Loreau et al., 2003; Thompson et al., 2017). Metapopulation models have also helped in advancing our understanding of biogeographic patterns (Gonzalez et al., 1998) and evolutionary processes such as the ecology and evolution of range limits (Holt and Keitt, 2000), dispersal (Olivieri et al., 1995; Wang and Altermatt, 2019), and the interplay of selection and gene flow in adaptive evolution (Hanski et al., 2011; Ingvarsson, 2002). In applied contexts, metapopulations have been used to understand biocontrol (Levins, 1969), habitat loss, and habitat fragmentation (Ovaskainen and Hanski, 2004).

The origins of metapopulation theory come from observations by empirical ecologists such as Andrewartha and Birch (1954), Huffaker (1958) and den Boer (1968) on the often ephemeral and unstable nature of local populations. The formal theory, stemming back to the seminal paper of Levins (1969) for a single species, and later extended to interacting sets of species in metacommunities (e.g., Holt, 1997; Tilman et al., 1997), addressed the challenge of understanding the interplay of dispersal and environmental heterogeneities. The complex spatial and temporal structure of environments shapes the fitness of individuals and biology of species over both ecological and evolutionary time scales.

Like all valuable theory, metapopulation and metacommunity theory neglects many

complexities of the world to make other complexities manageable. Metapopulation models

simplify space into suitable habitat patches surrounded by non-habitat, akin to islands in an

ocean. Early representations of metapopulations simplified population growth by representing

any given patch as either containing a species (i.e., occupied) or not (i.e., unoccupied), a

simplification that persists in many modeling efforts today (e.g., occupancy-based models). By

ignoring details of spatial variation and the intricacies of local population growth,

metapopulation models facilitate the study of population and community dynamics at large

scales. Encompassing the full details of continuous space and population dynamics, even without

spatial and temporal environmental heterogeneities, often leads to unwieldy mathematical

problem (see e.g., Bolker and Pacala, 1997; Holmes et al., 1994). The simplifying assumptions

of metapopulation models make them tractable, but they nonetheless retain some essential

features of heterogeneities in space and time.

But not all. The simplicity of metapopulation models is, in some senses, deceptive. Our

purpose here in effect is to 'look under the hood' of such models and bring out key but largely

unrecognized assumptions crucial to their behavior. We will highlight how recognition of these

assumptions, especially the nature of environmental variation, can explain emergent population

properties at broad temporal scales – ignorance of which degraded the ability of public health

authorities to manage and control the SARS-CoV-2 pandemic around the globe.

First, we reflect on the history of theory related to metapopulations and metacommunities

and highlight the implicit role of spatiotemporal heterogeneity in their dynamics. Without

spatiotemporal heterogeneity, many of the foundational lessons from classic metapopulation

theory no longer hold. Then, we highlight how the presence of spatiotemporal heterogeneity

influences multi-species counterparts to metapopulations—metacommunity models. We next

explore a phenomenon that has received some attention in the theoretical literature but little

empirical attention—namely the inflationary effect (a term coined in Gonzalez and Holt, 2002).

The inflationary effect refers to the elevated growth rate (Kortessis et al., 2020) or population

abundance (Gonzalez et al., 1998) that emerges in spatiotemporally heterogeneous landscapes

coupled by dispersal, as compared to the same landscapes without temporal heterogeneity

(Gonzalez and Holt, 2002). Studies of the inflationary effect attempt to deal more explicitly with

heterogeneities in space and time and illustrate a phenomenon that we suggest is implicit in

many metapopulation models. We discuss how awareness of the inflationary effect enriches our

understanding of the spread of infectious disease, and how a lack of such awareness hampered

the effective management of the pandemic caused by SARS-CoV-2. We finish by identifying

future directions in ecology, evolution, and conservation biology that involve spatiotemporal

heterogeneity and the inflationary effect. Our emphasis throughout is on conceptual issues and

illustrative examples, not on mathematical details, although we provide key references in the

history of the literature that provide such details for interested readers.

Spatiotemporal heterogeneity

Metapopulation models embody a host of implicit assumptions about the nature of the

environment. Amongst them, perhaps the most consequential is that spatiotemporal

heterogeneity is influential for the dynamics of species. Spatiotemporal heterogeneity denotes

how the temporal pattern of conditions varies as one moves across space. Spatial models alone

cannot capture the effects of spatiotemporal heterogeneity. Nor can models with temporal

heterogeneity, assuming a spatially well-mixed population. Spatiotemporal heterogeneity is a

unique aspect of the environment revealed only when considering space and time together.

Though the founders of metapopulation theory certainly recognized this dimension of theory

(e.g., Hanski, 1998), an explicit discussion of this fundamental assumption is encountered

infrequently in the literature, and so deserves special attention.

Before delving into models, it is first worth being specific about spatiotemporal

heterogeneity: What is it, exactly? In essence, spatiotemporal heterogeneity occurs when patterns

of conditions relevant to fitness over time differ across space (or spatial patterning that changes

over time). Spatiotemporal heterogeneity can be expressed by a statistical decomposition of the

total variation in some fitness-affecting factor, $F(x,t)$, with both time t and location x coordinates

(illustrated in Fig. 1a). The total variance in this factor, $\sigma^2 = \mathrm{Var}(F)$, can be broken into three

parts: σ^2_S, variability in space that is present after averaging over time, σ^2_T, variability in time that

is present after averaging over space, and σ^2_{ST}, spatiotemporal variability (see Box 1 for a

mathematical description and Chesson, 1985; Johnson and Hastings, 2023; Melbourne et al.,

2007; Schreiber, 2023 for alternative, but conceptually similar definitions).

The term σ^2_{ST} in Box 1 quantifying heterogeneity arising jointly across space and time is

our focus. We will argue that this is not a merely 'residual', but has important consequences.

Spatiotemporal heterogeneity cannot be explained by spatial-only and temporal-only

perspectives of the world, separately or added up (Fig. 1a-b). Figure 1a visualizes the

decomposition. In a temporal-only perspective, conditions vary over time and are the same

across space. In a spatial-only perspective, conditions differ across space but are maintained over

time. Spatiotemporal variation is additional variation on top of these two sources and may

constitute anywhere from 0% of the total variability to 100%.

Spatiotemporal heterogeneity can be generated in many ways. One fundamental source of

spatiotemporal heterogeneity is interactions between physical variation in abiotic factors across

space and time. For example, temperature fluctuations differ among land slopes with different

aspects; daily and seasonal fluctuations in temperature can be greater on south-facing slopes in

the northern hemisphere, and on north-facing slopes in the southern hemisphere. In aquatic

metacommunities in landscapes with many lakes and ponds differing in area and depth, larger

and deeper water bodies are buffered from fluctuations in physical factors such as temperature,

compared to smaller and shallower water bodies.

In addition to physical factors, biotic factors can create strong spatiotemporal

heterogeneity. Any species interacting with a focal species—biotic resources, natural enemies,

competitors, and mutualists—can generate variation in that focal species' fitness. Fluctuations in

the abundance or activity levels of an interacting species can potentially lead to spatiotemporal

heterogeneity in the fitness of a focal species if such fluctuations play out differently in different

locations. Spatiotemporally varying abundances of species are often said to have spatially

asynchronous dynamics (Shoemaker et al., 2022). Asynchronous population dynamics are a form

of spatiotemporal heterogeneity because the dynamical patterns of abundance fluctuations differ

across space (although perfectly synchronous dynamics can exhibit spatiotemporal heterogeneity

if the magnitude of synchronized fluctuations differ across locations).

Asynchronous dynamics can arise even in homogeneous physical environments, due to

complex processes involving dispersal, chance events, and interactions with other species

(Gouhier et al., 2010; Keeling et al., 2000; Levin, 1974; Vasseur and Fox, 2009). For example,

chaotic dynamics could be a potent source of asynchrony even in physically homogeneous

landscapes because slight differences in initial conditions lead to large deviations in resulting

population trajectories among locations (Hassell et al., 1991; Holt, 1993). Regardless of the

source of asynchrony, asynchronous dynamics of one species (e.g., a generalist predator) can

drive spatiotemporal heterogeneity in the fitness of other species with which it interacts (e.g., one

species of that predator's prey).

Box 1. Mathematical description of spatiotemporal heterogeneity

To illustrate how to quantify spatiotemporal heterogeneity, consider a landscape with locations indexed by x and times indexed by t. Let $F(x,t)$ be a measure of some factor that influences the fitness or growth rate of individuals found in location x at time t.

A characterization of these environmental factors that does not include the effects of time is a *spatial-only* perspective. Consider the probability distribution for a fitness-factor in a location x, which we denote with the conditional variable $F|x$. A reasonable measure for the conditions in location x is $\tilde{F}_x = E_t[F|x]$, the time-average of the factor in location x ($E_t[\cdot]$ being the expectation with respect to time; we use a tilde to indicate a time average). The variation present in space of these values, $\sigma^2_S = \mathrm{Var}_x(\tilde{F}_x)$, measures *pure spatial variation* ($\mathrm{Var}_x(\cdot)$ is the variance with respect to space).

A characterization of variation ignoring space is a *temporal-only* perspective. A reasonable measure of the conditions at time t across the entire landscape is $\bar{F}_t = E_x[F|t]$, the spatial average conditions at time t ($E_x[\cdot]$ being the expectation with respect to space; we use an overbar to indicate a spatial average). The variation present across time, $\sigma^2_T = \mathrm{Var}_t(\bar{F}_t)$, gauges *pure temporal variation* in the factor affecting population growth ($\mathrm{Var}_t(\cdot)$ is the variance with respect to time).

With these definitions of pure spatial and pure temporal variance in hand, we can write the

total variation in the factor affecting population growth as

$$\mathrm{Var}(F) = \sigma^2_S + \sigma^2_T + \sigma^2_{ST},$$

where σ^2_{ST} measures *spatiotemporal variation*. We can define this spatiotemporal variation as

(see supplementary material S1 for derivation)

$$\sigma^2_{ST} = E_t[\mathrm{Var}_x(F|t)] - \mathrm{Var}_x(E_t[F|x])$$

$$= \text{Time average of the variance across space} - \text{pure spatial variance}.$$

The quantity σ^2_{ST} represents the amount of excess variation that occurs when we look at an

ensemble of spatial variation across time rather than when averaged over time. An equivalent

(and mathematically identical) representation quantifies the excess variation in time that

occurs when we look at an ensemble across space over a simple average across space (see

supplementary material S1 for details).

Metapopulations

One of the first approaches for dealing with spatiotemporal heterogeneity was to envision

a metapopulation, a set of spatially discrete local populations prone to extinction and connected

by dispersal. Metapopulations in the most general sense of the word are "populations of

populations" (Hanski, 1999; Levins, 1970). They are populations defined on a scale that

encompass sub-populations, the dynamics of which are determined by individuals present locally

but also by individuals that leave (emigrate) and join (immigrate).

It is worth distinguishing distinct types of metapopulation models. The term

metapopulation sometimes refers to models based solely on occupancy, and sometimes to

models of patchily distributed populations where the abundances across locations are modeled.

We will use the term "strict metapopulations", to refer to models whose dynamics are

characterized in terms of the qualitative state of the population (i.e., occupancy) and where

recurrent local extinctions and colonizations determine dynamics (Hanski, 1998). We will use

the term "patch models" to describe models in which dynamics are characterized by quantitative

measures of population states (e.g., local density) and whose dynamics depend explicitly on

birth, death, emigration, and immigration rates. Both models may include spatiotemporal

variation, but its inclusion in patch models is often explicit whereas it is implicit in strict

metapopulation models.

The distinction between "strict metapopulations" and "patch models" is to a degree an

arbitrary one based on choices used to describe local populations. All real populations are

spatially distributed with dynamics that could be described by abundance or by a simpler metric

of abundance: presence (abundance greater than zero) or absence (abundance equal to zero).

Indeed, any patch model could be rewritten as a strict metapopulation model, and there are many

types of local population dynamics that are consistent with any given strict metapopulation

model. They are, put simply, complementary modeling frameworks, each with their strengths

and limitations.

Simple models of spatiotemporal heterogeneity: spatial occupancy models

Levins (1969) introduced the concept of a metapopulation to deal precisely with the

complexities generated by modeling spatiotemporal heterogeneity[1]. (The term "metapopulation"

---

[1] While this is often overlooked in more contemporary descriptions, environmental heterogeneity was at the forefront of Levins' mind when developing his model. Levins' original article articulating the metapopulation model is titled "Some demographic and genetic consequences of environmental heterogeneity for biological control", the wording of which highlights the influence of environmental heterogeneity in Levins' thinking.

seems to have its first usage in Levins, 1970). He recognized that conditions differing across

space and time could cause populations sizes to fluctuate wildly, oftentimes causing extinctions

of local populations. But sites do not remain empty because reproductively viable individuals

from nearby, extant populations can immigrate.

The localization of extinctions and the subsequent potential for recolonization have

foundational roles in strict metapopulations. Classical, strict metapopulation models are based on

the following canonical form:

$$\frac{dp}{dt} = cp(1 - p) - ep, \tag{1}$$

where p is the proportion of patches on the landscape currently occupied, e is the rate at which

populations go locally extinct, and cp is the rate at which unoccupied patches again harbor a

viable population (c is the colonization rate per occupied patch). Such a model describes

landscapes with many patches whose extinction events are independent across patches, and

where recolonization events do not depend on the spatial position of empty patches (Hanski,

1998, 1997). Model (1) predicts that if the extinction rate is lower than the colonization rate (i.e.,

$e < c$), the fraction of occupied patches grows logistically to an equilibrium of $p^* = 1 - e/c$. Even

though local extinction is inevitable everywhere, frequent recolonizations allow for

metapopulation persistence.

A crucial assumption is that local extinctions are not strongly synchronized across space;

otherwise, recently extinct populations would have no sources for recolonization. This point has

been made before (Chesson, 1981; Hanski, 1998; Hoopes et al., 2005), but is rarely a central

point of discussion of metapopulations and so can be easily missed (although see Chesson, 1985;

Harrison and Quinn, 1989; Petchey et al., 1997). The assumption in Levins' model of no strong

synchronization thus reflects an implicit assumption of the presence of spatiotemporal

heterogeneity if environmental factors (rather than say pure demographic stochasticity) drive

those extinctions. Conditions in any one local population, at any one time, are not identical to all

other local populations during that time.

To illustrate this subtle assumption and its consequences for metapopulation dynamics,

consider a finite patch version of model (1). The model has n patches where we track occupancy

in each patch. Let $O_i(t)$ be the occupancy in patch i at time t ($O_i = 1$ if occupied and $O_i = 0$ if

not), and let $P(t) = \sum_i O_i(t)/n$ be the fraction of patches occupied at time t. To simulate the model,

we consider how occupancy changes in a small unit of time, Δt. In each unit of time, each

occupied patch goes extinct with probability $1 - \exp(-e\Delta t)$, and each unoccupied patch becomes

occupied with probability $1 - \exp(-cP(t)\Delta t)$. As the time steps get small, extinction occur at rate

e per occupied patch, and colonization occurs at rate $-cP(t)$ per unoccupied patch, just as in

model (1) (see supplementary material S2 for model details.)

Spatiotemporal variability can be controlled with the *joint probability* of extinctions and

colonizations across multiple patches each time step. If we assume extinction events in each

patch are independent in space and time, we recover a finite-patch version of equation (1). In that

case, any pattern of extinction-causing factors will have different patterns over time among

patches (see supplementary figure S1), an example of strong spatiotemporal heterogeneity. If

instead we assume extinction drivers in one patch correlate with drivers in other patches, patterns

of extinction over time would share at least a degree of similarity across patches, i.e., there is

decidedly less spatiotemporal heterogeneity. We can tune the level of spatiotemporal

heterogeneity with ρ_E, the correlation between any two patches in environmental conditions

affecting extinction (for simplicity, we assume this correlation applies to all pairs of patches).

Since all patches are otherwise the same, there is no spatial heterogeneity ($\sigma^2_S = 0$). When $\rho_E = 0$,

extinction events are independent across patches and all variation is spatiotemporal ($\sigma^2_{ST} > 0$; σ^2_T

= 0). When $\rho_E = 1$, extinction in one patch implies extinction in all, there is no spatiotemporal

variation ($\sigma^2_{ST} = 0$). In fact, in that case, all variation is temporal ($\sigma^2 = \sigma_T^2$).

With independent extinctions (i.e., $\rho_E = 0$), the classical, strict metapopulation assumption

is recovered, and the population reaches a relatively stable patch occupancy (Fig. 2, green line),

even in a finite patch model with relatively few patches ($n = 50$). By breaking the assumption of

independent extinction events across locations (i.e., reducing spatiotemporal heterogeneity),

stability of the metapopulation weakens (Figure 2, green line). Occupancy exhibits large

fluctuations because local extinctions, when they occur, tend to occur over a large spatial domain

with many local populations impacted simultaneously. Even highly occupied metapopulations

are vulnerable to drastic drops in occupancy because of highly synchronized extinction events.

The large fluctuations caused by eroding spatiotemporal heterogeneity greatly increase

the likelihood of metapopulation extinction. For example, a metapopulation of $n = 25$ patches

with independent extinction events (i.e., strong spatiotemporal heterogeneity) persists an average

of ~300,000 units of time (Fig. 3a, dark purple points). Removing this implied spatiotemporal

heterogeneity by increasing the correlation among patch extinction events reduces persistence

times by several orders of magnitude (Fig. 3a), despite little change in average occupancy (Fig.

3b). This simple model shows that the well-known stability implied by metapopulation models

results from the implicit assumption of strong spatiotemporal heterogeneity among patches.

Metacommunities: spatiotemporal heterogeneity in species interactions

A natural extension of metapopulation models is to consider other species. Patches now

represent local areas harboring communities connected by dispersal, the collection of which is a

metacommunity (a term first used by Wilson, 1992). Species' colonization and extinction rates in

metacommunities reflect some combination of the physical environment, interspecies

interactions, and demography. We distinguish "strict metacommunities", whose dynamics are

described in terms of extinction and colonization rates (i.e., occupancies, hence co-occupancies)

(see e.g., Hastings, 1980), and community patch models, whose dynamics are described in terms

of birth, death, immigration, and emigration rates of each constituent species within each patch

(see e.g., Holt and Chesson, 2016). Spatiotemporal heterogeneity in patch models—a rich and

growing area of community ecology (Driscoll and Lindenmayer, 2009; Leibold, 2018; Leibold et

al., 2022; Lu, 2021)—is considered below. Here we focus on strict metacommunities.

To illustrate issues that pertain more broadly to more complex multispecies models, we

use a simple model for a core component of any metacommunity – a specialist predator

interacting with its required prey species. A specialist predator-prey strict metacommunity model

is

$$\frac{dp}{dt} = c_p p(1 - p) - \left(e_p + c_q q\right)p$$

$$\frac{dq}{dt} = c_q qp - e_q q,$$

(2)

where p is the fraction of habitat patches occupied by prey alone, q is the fraction occupied by

both predators and prey, c_i is the colonization rate of patches of type i ($i = p,q$), and e_i is the

extinction rate of patches of type i (e.g., the prey alone, or it together with the predator).

Models such as (2) follow the same simple rules as a strict metapopulation model except

that now extinction and colonization rates depend on other species. Patches occupied by prey

alone change state independently of the predator (e.g., due to disturbance) with an extinction rate

of e_p and because they are colonized by the predator at a rate proportional to the availability of

prey-only patches and the prevalence of predators ($c_q p q$). We assume the predator greatly limits

prey numbers in co-occupied patches, so that patches occupied by the predator do not provide

any prey colonists to empty patches. The above model assumes that predators only go extinct

when their prey go extinct (although more elaborate models that relax these assumptions to

investigate other ways predators might influence prey extinction follow the same qualitative

rules; see Holt, 1997).

Metacommunity models often exhibit stability despite locally unstable interactions

between species in any patch (Gravel et al., 2016; Levin, 1974). Strong predator-prey dynamics

typically generate large-amplitude fluctuations in spatially and temporally homogeneous

environments (May, 1972). It is often difficult to sustain such dynamics experimentally (but see

Blasius et al. 2020 for a recent laboratory success). But metacommunity models such as (2)

exhibit stability despite such unstable local dynamics, where predators, once arriving at a patch,

eventually drive prey extinct locally (Gouhier et al., 2010; Hastings, 1977).

The stability exhibited in metacommunity models again reflects spatiotemporal

heterogeneity. To understand this, consider the predator first. Given sufficient availability of

prey patches to colonize, predator stability is ensured by the fact that extinctions occur

independently across locations, just as in metapopulation model (1). And a stable, suitable prey

source is ensured by the same argument, but including the fact that predators also show some

independence across patches in their colonization of prey-alone patches. Stated succinctly,

predator effects are spatiotemporally heterogeneous. Synchronous predator colonization or

extinction events would lead to large-scale oscillations comparable to those of Figure 2, with

high likelihood of extinction in realistic models.

For other metacommunity models including, for instance, infectious disease and

competitive interactions (discussed below), stability at the regional scale is also observed, and

relies on the implicit assumption that extinction and colonization events are independent across

patches, as in our stochastic metapopulation model (Figs. 2,3). Independence across patches

requires spatiotemporal heterogeneity be present and strong. One perspective about strict

metacommunity models, we suggest, is that some of the most interesting features about the

maintenance of diversity in metacommunities emerge from spatiotemporal heterogeneity.

Digging deeper: beyond models of occupancy

The stabilizing effects of spatiotemporal heterogeneity can be difficult to recognize in

strict metapopulation models. Such effects become clearer in patch models, where population

dynamics within patches are explicit in terms of births, deaths, emigration, and immigration.

Rather than comprehensively review the large and diffuse literature related to this theme, we

here follow one line of thought about the emergent effects of spatiotemporal variation.

Holt, Gonzalez, and other collaborators began exploring the consequences of

spatiotemporal heterogeneity in patch models in a series of papers beginning in the early 2000s.

In a simple but illuminating example, Gonzalez and Holt (2002) considered a single sink patch

supported by constant immigration from a source patch. It had long been recognized that

populations may exist in sinks given constant immigration from a source, and that the sink

population could be substantial given moderate rates of population decline (Holt, 1985; Pulliam,

1988; Runge et al., 2006; Van Horne, 1983). Gonzalez and Holt (2002) expanded on this theme

by envisioning that local sink conditions were sinks on average over time, but not all the time.

On average, conditions in the sink habitat were too poor to sustain a population without

immigration, but transient phases of locally favorable conditions might allow a population to

temporarily grow. Without immigration, the population may fluctuate and temporarily increase,

but it eventually, given its sink namesake, wanes away and goes locally extinct.

However, immigration prevents extinction during bad times, buffering declines to low

numbers. Immigration then allows the persistent sink population to capitalize on good times,

boosting time-average population sizes in the sink (sometimes by orders of magnitude), a

phenomenon they termed "the inflationary effect". Gonzalez and Holt (2002) demonstrated this

effect mathematically and used a microcosm experiment with source-sink populations of the

protozoan *Paramecium tetraurelia*, buffeted by temporally varying thermal conditions, to

rigorously test the phenomenon.

The inflationary effect of spatiotemporal variation on abundance works as follows and is

illustrated in Fig. 4. Conditions are sometimes poor for growth, and sometimes favorable for

growth. Good conditions cause large upswings in population size, particularly if growth rates are

positively autocorrelated, permitting runs of good conditions. Poor conditions decrease

population size, but rather than the population crashing (and erasing any increase in abundance),

persistent immigration into the sink maintains a minimum growth rate, keeping the population

from dipping too low. Thus, the effects of favorable conditions outweigh the effects of

unfavorable conditions, even if good and bad times are equally frequent. In essence, immigration

maintains a minimum local growth rate that, when paired with temporarily favorable conditions

for growth, greatly increases sink population size.

This increase in sink population size – an inflationary effect of abundance – is

fundamentally about spatiotemporal heterogeneity. The measure of spatiotemporal variance from

Box 1 applied to the model first investigated in Gonzalez and Holt (2002) reveals that $\sigma^2_{ST} =$

$\sigma_r^2/4$, where σ_r^2 is the variance in the sink growth rates (see supplementary material S3). Hence,

the inflationary effect of population size in Fig. 4 is present when $\sigma^2_{ST} > 0$ and absent when σ^2_{ST}

$= 0$ (black line in Fig. 4).

Further modeling studies demonstrated that the inflationary effect arises in many other,

more complex and biologically realistic models, and may show up in the metapopulation growth

rate as well as time-averaged abundance. The effect occurs in source-sink models with explicit

resource dynamics, density-dependence, demographic stochasticity, and competition between

species (Holt et al., 2003). Somewhat remarkably, it also applies to multiple sinks linked by

dispersal (so there is no persistent source habitat anywhere in the landscape at all) and can be

sufficiently strong in some cases to ensure the persistence of a regional landscape comprised

entirely of stochastic sinks (Roy et al., 2005).

This last point is worth reiterating. The inflationary effect can allow for the regional

persistence of populations living in a landscape of sinks. That is to say, in any given location on

the landscape, a population isolated from immigration cannot persist. But if some individuals

there disperse (not too many) to neighboring locations, such reshuffling of individuals can be

sufficient to allow for all populations to persist in perpetuity. This boost in regional growth

depends on spatiotemporal variation and asynchrony in growth conditions among patches, and it

is a specific case of paradoxical persistence of mixed dynamics systems with strong

heterogeneity (see the excellent review of such systems in Williams and Hastings 2011).

A microcosm study (Matthews and Gonzalez, 2007) prompted by this result convincingly

demonstrated that asynchrony could permit metapopulation persistence, even for just two patches

coupled by dispersal. Matthews and Gonzalez observed that two local populations of

Paramecium aurelia would quickly go extinct in both patches for treatments with synchronous

temperature fluctuations across patches, but populations persisted in treatments with

asynchronous temperature fluctuations across the two patches. They also showed that positive

autocorrelation boosted the inflationary effect on sink population sizes, a result that was

extended to host-parasite dynamics (Duncan et al., 2013) in metapopulations of *Paramecium*

caudatum and the bacterial parasite *Holospora undulata*. The key finding is that autocorrelated

and asynchronous variation in temperature allowed infected host populations to maintain sizes

equivalent to uninfected populations, despite negative demographic impacts on infected

individuals.

One might reasonably wonder whether such results are limited to small parameter ranges

with highly specific conditions, in which case the inflationary effect would merely be a

theoretical curiosity of scant real-world importance. Recent studies however have investigated

quite general models of population growth in discrete and continuous-time and find the

inflationary effect to be present over broad ranges of parameter space (Benaïm et al., 2023;

Katriel, 2022; Schreiber, 2010). Moreover, other studies have shown that the rates of dispersal

consistent with the inflationary effect are adaptive in both single species (Schreiber, 2012) and

multispecies models (Schreiber et al., 2023). These studies demonstrate a very general pattern:

spatiotemporal heterogeneity in growth rates can robustly allow for persistence of species and

increase their abundance. This should not come as too much of a surprise, as the lesson from

theories of metapopulations is that the metapopulation can persist despite no individual patch

persisting on its own, without the others. The inflationary effect is about just such persistence,

but the details of these models demonstrate the mechanism by which spatiotemporal

heterogeneity can be beneficial for population persistence, as we demonstrate below.

What is "the inflationary effect", and how does it work?

The clearest requirements for the inflationary effect were illustrated by Roy et al. (2005)

for abundance and Schreiber (2010) for growth rate, which we summarize here as they provide a

pathway to understand its mechanistic underpinnings. In both cases, the inflationary effect can be

thought of as the effect of spatiotemporal heterogeneity inflating some aspect of population

growth at the scale of the metapopulation, such that it is much larger than some simple average

of local conditions.

Prior studies illustrate that the inflationary effect needs (1) some (but not too much)

movement among patches, (2) variation in local growth rates over time, (3) a less than perfect

spatial correlation in growth rates, and (4) temporal autocorrelation in growth rates. Temporal

variation and low spatial correlation in growth rates together are signatures of spatiotemporal

heterogeneity. The combination of movement and temporal autocorrelation together allows

individuals in the metapopulation sufficient time to disperse across space so as to experience

spatiotemporal heterogeneity. Without movement, the effective landscape of an individual is a

single patch, not a metapopulation, despite the landscape having multiple patches. Without

temporal autocorrelations, environments shift before populations can build up in transiently

favorable locations. As populations grow, more individuals emigrate (in an absolute, not per-

capita sense), even if the locations into which they immigrate are currently poor for growth.

Given spatiotemporal heterogeneity, favorable conditions in one patch imply that emigrants from

there are likely to immigrate to a temporary sink, boosting its numbers. Temporary sinks

disproportionately benefit from immigration because it maintains minimum population sizes,

which enhances population (not per-capita) growth rate when the environment eventually turns,

as illustrated in the sink model with constant immigration above (Figure 4, see supplementary

material S3 for model details).

This qualitative description can be made more precise with a mathematical description of

the effect, which reveals how the mechanism operates on growth rates. Others have provided a

mathematical description for discrete-time (Johnson and Hastings, 2023; Schreiber, 2023, 2010),

but we present one for continuous-time where the math is (slightly) less painful. Consider a set

of n connected patches where N_i is the local density in a patch i, $r_i(t)$ is the local per capita

growth rate, and m_{ij} be the patch-specific per capita movement rates from patch j to i. The

dynamics in patch i can be written as

$$\frac{dN_i}{dt} = \underbrace{r_i(t)N_i}_{\text{Local Growth}} - \underbrace{\sum_{j\neq i}^{n} m_{ji}N_i}_{\text{Emigration}} + \underbrace{\sum_{j\neq i}^{n} m_{ij}N_j}_{\text{Immigration}}, \tag{3}$$

which allows for time- and location-dependent growth and movement rates.

To understand the role of spatiotemporal heterogeneity, we could ask how different

population growth would be in the absence of spatiotemporal heterogeneity. Hence, a reference

scenario in which spatial and temporal variation's effects are present, but the spatiotemporal

effects are not, is required. We use the scenario where the average growth rates across time

describe local growth in each patch (using the notation in Box 1, $r_i(t) = \tilde{r}_i$ for all t). This scenario

encapsulates the effects of temporal variability alone with the average growth rate and the effects

of spatial variability by allowing each population to have different average growth rates with

dispersal. This population reaches a stable spatial distribution, which is analogous to the stable

age distribution in stage-structured demographic models. Let $\tilde{v}_i = N_i/[n^{-1}\sum_j N_j]$ be the stable

relative density in patch i in the reference scenario. The growth rate in this reference scenario can

be found with the dominant eigenvalue of (3) (see supplementary material S4).

The amount to which the actual metapopulation growth rate deviates from this reference

scenario without spatiotemporal heterogeneity is

$$\text{Inflationary Effect} = E_t\big[\text{cov}_i\big(r_i(t), v_i(t)\big)\big] - \text{cov}(\tilde{r}_i, \tilde{v}_i), \qquad (4)$$

where $v_i(t) = N_i(t)/[n^{-1}\sum_j N_j(t)]$ is the relative density in location i at time t in the full model with

spatio-tempoal heterogeneity (derivation in supplementary material S4).

Equation (4) measures the average association over time between the density of

individuals across space and local growing conditions and how it compares to the association

between average conditions. As such, it measures the resulting ability of populations to "track"

good locations that shift in time and space in reference to the ability to find and persist in good

locations on average. Expression (4) depends on a time-average of a covariance, which may be

either positive or negative, and may lie above or below the covariance for the reference scenario,

implying that the inflationary effect could be positive, zero, or negative. At any given time, the

covariance is positive when individuals are concentrated in better than average spatial locations

and is negative if instead individuals in the metapopulation are concentrated in poorer than

average spatial locations. When population density is random with respect to local patch growth

rates, the covariance is zero. Negative covariances can arise, for instance, under directed passive

dispersal out of sources into sinks, as demonstrated by Keddy's (1982) study on the sea rocket in

Nova Scotia dunes, where seed dispersal is driven by winds from the ocean across the dunes.

Prior studies studying the inflationary effect have focused on situations where equation

(4) is positive, meaning that a randomly chosen individual in time and space experiences

conditions better than the average spatial conditions in time. These studies find that the average

individual experiences (somewhere) conditions in space that are (temporarily) favorable to

growth. These opportunities may not be present consistently, in any time or place, but are present

ephemerally. Equation (4) measures how dispersal capabilities of species interact with the

spatiotemporal structure of the environment. (Deriving explicit expressions for this quantity is

analytically challenging, however some approximations are possible; Schreiber, 2023.)

Studies on the inflationary effect typically assume passive forms of non-directional

dispersal, which leads the average individual to experience better than average conditions, even

without dispersal adaptively directed towards such locations. A positive covariance between

local growth rate and local relative abundance arises simply because populations tend to grow

more where conditions are temporarily good. Adaptive tracking of the environment will certainly

enhance the effect of spatiotemporal variation in fitness on population growth. For instance, if

organisms tend to show site fidelity in breeding, staying the next year if this year was good, and

local environments are positively autocorrelated in fitness over time, this tendency helps

populations persist in spatiotemporally varying environments (Schmidt, 2004). By contrast, we

suspect that some organisms with environmentally forced dispersal, such as happens with wind-

dispersed plant propagules and aquatic and marine organisms dispersing with water currents,

may experience limited (or no) benefits of the inflationary effect.

The relationship between inflationary effects on abundance and growth is as yet

undetermined. Inflated abundance in many cases implies inflated population growth at the

metapopulation scale, as is illustrated by the case of persistence in a landscape of stochastic

sinks. However, it is unclear whether inflated growth when rare always leads to inflated

abundance. We suspect that the answer lies in how density-dependence acts on the landscape to

influence how variation in fitness factors translates into long-term average abundance. Future

modeling work is necessary to understand the relationships between abundance and growth rate

inflation.

At this time, although the theory has been demonstrated in several laboratory experiments

with model species, we do not know how widespread the inflationary effect is in nature. This is

an open question for future empirical research. But the inflationary effect seems most likely to

occur for species living in environments with the characteristics outlined in Table 1. The

characteristics in Table 1 should be useful indicators for empirical study of the inflationary effect

in nature. The inflationary effect remains a niche concept in population and community ecology,

despite its apparent connection to the basic concepts of stability in metapopulation and

metacommunity models. We hope that this review, and the case studies from epidemiology

provided below, spark some interest in the concept, especially among empiricists.

Spatiotemporal heterogeneity and infectious disease

Our emphasis thus far has been conceptual with examples from free-living protozoans

and dune plants. Some others we have noted suggest application of the inflationary effect to host-

parasitoid dynamics (Duncan et al., 2013), hinting at application of the inflationary effect, and

the role of asynchrony and spatiotemporal heterogeneity, beyond the usual domain of the

ecological sciences to epidemiology. We here argue that the inflationary effect looms large in the

spread of infectious disease.

When considering infectious diseases, there are two scales at which one can envisage

metapopulation dynamics. First is the metapopulation perspective of hosts as patches. From the

perspective of an individual pathogen (e.g., a bacterium or virion), each individual host is an

enormous habitat patch or even heterogeneous landscape (Holt, 2000). An individual SARS-

CoV2 virion is about 0.1 microns in diameter and resides in a human host averaging 1.7m in

height. Were we to scale up the virion to the size of its host, and re-scale host size accordingly,

the host would be 29,000 km tall – more than twice the diameter of the Earth. At peak infection,

the estimated viral load in a single infected human host is 10^9 to 10^{11} viruses (Sender et al., 2021)

– there are about 10^{11} stars in our galaxy.

Transmission to uninfected hosts is akin to colonization of empty patches, and pathogen

clearance (or host death) is a form of local extinction for the pathogen. Susceptible-infectious-

recovered (SIR) models assume hosts are either healthy (but susceptible to infection), infectious

(and potentially unhealthy), or recovered from prior infection (and unable to be re-infected). This

is the epidemiological analogue of the metapopulation assumption that one only need track

pathogen occupancy (presence-absence) in a host habitat patch, rather than the details of within-

patch (or within-host) population dynamics. To see the role of spatiotemporal heterogeneity,

recognize that the usual form of SIR style models (and the constellation of similar models)

assumes an exponentially distributed timing for both transmission and infection clearance at the

individual level, meaning there are differences among hosts in the clearance (extinction) and

infection (colonization) times. In other words, epidemiological events across hosts are

asynchronous. Linking such within-host dynamics to among-host dynamics is a large challenge,

with many avenues yet to be explored (e.g., analogues of emigration, see Barfield et al., 2015).

This is an important theme, but here we instead go up in spatial scale, considering

metapopulations of hosts (see e.g., Cliff et al. 2000).

At the level of the hosts, there can be metapopulations of hosts coupled by host

movement (or possibly vector movement), which is our focus. Epidemiologists have long been

concerned with aspects of disease spread in spatially distributed host populations (Ch. 7 of

Keeling and Rohhani, 2007 and references therein), and with temporal variation in pathogen

transmission (e.g., seasonal outbreaks in a locality; Shone et al. 2006; Moore et al. 2014).

However, the impact of spatiotemporal variation along the line of our inquiry has been largely

neglected.

Asynchrony among host populations in pathogen transmission can arise for many

reasons. For instance, if a pathogen is transmitted by an arthropod vector along climatic

gradients, seasonal peaks in pathogen transmission might develop at different times across space

(Moore et al., 2014; Shone et al., 2006). This likely will be exacerbated for arthropod borne,

transovarially transmitted pathogens where infected adult females may infect most, if not all,

their progeny following dispersal. Spatiotemporal variation in the abundance of non-human hosts

(Childs, 2004) for multi-host pathogens could influence patterns of variation in human disease.

Human behavior is often key to infectious disease transmission, and spatiotemporal

variation in behavior can be a particularly potent source of spatiotemporal heterogeneous

patterns of disease transmission. Public health policies also lead to spatiotemporal heterogeneity

in transmission and asynchrony in control among polities or nation-states because differing

levels of intensity in control measures and public health infrastructure imply infectious diseases

are able to spread much more efficiently in some regions than in others. Asynchrony arises

because different localities experience different patterns of human behavior over time, driven

both by 'bottom-up' individual effects (e.g., differences among locations in social expectations

about the importance of non-pharmaceutical control measures such as mask-wearing, willingness

to avoid crowded venues and the like), and 'top-down' forces, such as imposed public health

measures.

In 2020, many of the present authors noticed the potential for broad scale asynchrony in

disease transmission to influence the spread of SARS-CoV-2, the etiological agent of COVID-

19, early during the COVID-19 pandemic. Given that SARS-CoV-2 was a novel and virulent

infectious agent, few therapeutic treatments were initially available. In the absence of known and

effective medical treatment, the main approach to slow and limit the spread of SARS-Cov-2—

thereby reducing stress on health care infrastructure—was the use of non-pharmaceutical

interventions (e.g., masking, social distancing, closing of non-essential business places,

lockdowns; hereafter NPIs). The timing, nature, and efficacy of NPIs varied across time and

space as social, economic, and epidemiological conditions changed. One simple example is that

government-ordered lockdowns of local non-essential businesses might respond to rises in local

reported case numbers. Such feedback interacts with existing heterogeneity in the social,

behavioral, economic, and demographic structure of populations to generate differential timing

of lockdowns. Uncoordinated decision making about the timing of NPIs could generate

substantial spatiotemporal heterogeneity in a fitness factor of SARS-Cov-2, the generalized

transmission rate.

To illustrate this idea, consider a version of the model used by Kortessis et al. (2020) to

explore potential consequences of asynchrony in the timing of lockdowns (see supplementary

material S5 for model details). In that model, there are two patches of human hosts linked by

movement. Disease spread in each patch follows the assumptions of an SIR model. All

epidemiological parameters are identical between patches except for $\beta(x,t)$, the generalized

transmission rate, which is indexed by patch location ($x = 1,2$) and time ($t \in [0,\infty)$). For

simplicity, we assume that $\beta(x,t)$ takes two values over time, β_0, the baseline transmission rate,

and $\beta_0(1 - \epsilon)$, where ϵ is the effectiveness of NPIs on reducing disease spread locally in

proportional terms. Furthermore, we consider that when a lockdown occurs, it persists for

duration T, before returning to "business as usual", with periodic bouts of lockdowns (i.e.,

"square-wave" temporal variation). To model differential timing of NPIs, we assume a delay

between the initiation of NPIs in the different patches and keep track over how much time the

two locations are in the same state (both "business as usual" or instead both under NPIs). We call

the fraction of time that the two patches are in the same state Ω, the NPI overlap.

The overlap in NPI controls the magnitude of spatiotemporal heterogeneity. Indeed, the

measure of spatiotemporal variance is $\sigma^2_{ST} = (1-\Omega)(\beta_0\epsilon/2)^2$, and on a scale proportional to the

total variation is $\sigma^2_{ST}/\sigma^2 = (1-\Omega)$ (see supplementary material S5). Hence, synchronized policies

($\Omega, = 1$) do not generate spatiotemporal variability, whereas asynchronous policies ($\Omega < 1$) do

generate spatiotemporal heterogeneity. When policies are exactly out of sync (i.e., $\Omega = 0$)

spatiotemporal heterogeneity accounts for all variability in pathogen transmission.

Pairing host movement between patches with less than complete overlap in NPIs greatly

elevates the rate of increase in new infections at the metapopulation scale. This is an instantiation

of the inflationary effect for infectious diseases, where aspects of human behavior plausibly

create spatiotemporal heterogeneity in conditions affecting the fitness of a respiratory virus. The

effect is larger for more effective NPIs (i.e., larger ϵ, generating greater variation in transmission

over time) and when the duration of application of NPIs is longer (larger T, details in

supplementary material S5). When NPIs do not completely overlap and some infectious

individuals travel between patches, a stochastic version of the model shows increasingly larger

waves of infection than with complete NPI overlap (Figure 5b).

Evidence from cell phone data shows human mobility changes during state-enforced

NPIs in the European Union (EU) were sufficient to promote the inflationary effect in the

presence of repeated, statewide lockdowns. Ruktanonchai et al. (2020) illustrated that if different

countries in the EU came out of lockdown at different times (thereby generating spatiotemporal

heterogeneity – as in fact was the case), then there would be many more infections and

hospitalizations from COVID-19 than if states came out of lockdown in a coordinated manner.

Others have modeled the timing of application of NPIs in light of this phenomenon and found it

to be quite general (Berestycki et al., 2023; Eksin et al., 2021; Wang and Wu, 2022; Zhang et al.,

2021). A general conclusion is that optimal local public health strategies locally depend on the

public health decisions made by nearby localities. In other words, spatiotemporal heterogeneity

in disease transmission, coupled with movement among locations, degraded the control of the

pandemic, with serious and devastating consequences for our species across the globe.

Discussion

Recent work on the inflationary effect clarifies and extends the intuition of empirical

ecologists many decades ago about the role of spatiotemporal variation in fitness, along with

dispersal, in promoting regional persistence of species. Andrewartha and Birch (1954, p. 657),

for instance, remark: "A natural population occupying any considerable area will be made up of

a number of … local populations or colonies. In different localities the trends may be going in

different directions at the same time." Later, they wrote admiringly of den Boer's idea of

spreading of risk (den Boer, 1968) and noted "The concept of spreading risk recognized the

importance of dispersal with the natural population that enables individuals to colonize new

localities" (Andrewartha and Birch, 1984, p. 175). They extensively quote den Boer (1981, p.

39), who suggests "The survival time of composite populations uninterruptedly inhabiting large

and heterogeneous areas, highly depends on the extent to which the numbers fluctuate unequally

in the different subpopulations." In other words, population persistence at broad scales may rest

on the interplay of dispersal and spatiotemporal variation in fitness.

The influence of spatiotemporal variation can be discerned even if populations are not

found in crisply defined habitat patches. den Boer (1981) studied ground beetle ecology over 35

years and showed that though dispersal occurred, it was mainly at a low level and over short

distances. The beetle *Pterostichus versicolor* fluctuated asynchronous across ten locations and

did not experience extinction. By contrast, the beetle *Calathus melanocephalus* fluctuated

synchronously, and did show local extinction. As Gotelli (2008, p. 96) notes for the persistent

species, "… at any point in time, some populations were increasing in size and acting as source

populations that prevented the extinction of other, declining sink populations." The other species

experienced synchrony in its fluctuations, so there were times when conditions were uniformly

bad for it everywhere. The data den Boer (1981) presents resembles qualitatively the patterns

shown in the simple metapopulation model of Figure 1.

A new generation of empirical research is needed to establish how widespread the

inflationary effect is in nature (see Table 1 for the necessary features), both in single species

metapopulation dynamics and in metacommunities where species interactions may contribute to

fluctuations in fitness over space and time.

Invasion biology

While initial studies of the inflationary effect highlighted persistence of a metapopulation

of sinks (Roy et al. 2005), a novel phenomenon highlighted in Kortessis et al. (2020) is that the

inflationary effect could also speed up the course of an invasion, even if metapopulation

persistence is not an issue. We suggest that this effect might contribute to a well-known

phenomenon in invasion biology – biphasic and accelerating patterns of expansion for invasive

species, including initial lag phases (Hui and Richardson, 2017, pp. 28–37 for many examples).

In the case of the common starling, *Sturnus vulgaris*, for instance, its spread was sluggish in the

first few decades after its introduction into North America (Okubo, 1988), and then it sped up.

Similarly, West Nile Virus was slow to disperse throughout the eastern USA and remained at

low incidence in the first couple years following introduction in New York City in 1999, but then

rapidly expanded in 2002 (Petersen and Hayes, 2008).

A variety of explanations for this pattern have been suggested (e.g., transitions among

different habitats, Okubo 1988; stratified diffusion, Shigesada et al. 1995). The inflationary

effect provides another viable explanation. If there are n locations with independently

fluctuating growth rates, and a species is introduced into one of those locations at random and

thereafter disperses, then its initial expected growth rate (prior to much dispersal) is simply the

arithmetic average growth rate over those locations. However, as time goes on and individuals

move among locations, then the covariance term in expression (4) grows, and if this is positive,

the invasion will accelerate. Spatial population data, especially for invasive insects, could be

used to assess the role of the inflationary effect during invasion.

Natural enemy-victim interactions

Effective predators tend to overeat their prey, and then suffer extinction themselves. The

'spreading of risk' among different populations, loosely coupled by dispersal, is a likely reason

for the persistence of strong natural-enemy interactions. In the classic experiments of Huffaker

(1958), the persistence of a predatory mite (*Typhlodromus occidentalis*) with a prey mite

(*Eotetranychus sexmaculatus*) in a mesocosm comprised of an array of oranges depended on

limited dispersal, and the fact that dynamics were asynchronous across space. This experiment

inspired an experiment with a bruchid beetle *Callosobruchus chinensis*, feeding on beans,

attacked by a pteromalid parasitoid *Anisopteromalus calandrae* (Hassell and May, 1988). In an

environment without spatial structure, the parasitoid rapidly overexploited its host, then went

extinct (allowing the host to rebound). When each beetle was put in its own patch, with restricted

dispersal among patches, the host-parasitoid interaction persisted indefinitely. Here, the physical

environment was homogeneous (by experimental design), and spatiotemporal variation in each

species' fitness arose entirely from its localized interaction with the other species and its own

direct density dependence.

It is likely that this effect explains the robust persistence of a wide range of strong natural

enemy-victim systems (Hassell, 2000). Because dispersal is often localized, the asynchrony

needed for spatiotemporal variation to be expressed depends upon spatial scale. The relevant

scale(s) of temporal fluctuations will vary with the generation time, demography, and movement

rates of the focal species (Bernhardt et al., 2020). For example, Wilson et al. (1998) examined

how the persistence of host-parasitoid and host-parasitoid-hyperparasitoid interactions depends

upon the size of the spatial lattice within which these dynamics played out. They found that food

chains of strongly interacting specialist species require larger arenas to persist. Adjacent areas

are more likely to be synchronous than are distant, far-flung locations. Island and patch size

effects on food chain length of specialists might thus reflect in part the importance of spatial

scaling of asynchrony in population dynamics among localities within defined spatial arenas

(Holt, 2010; e.g., islands, habitat fragments, Wilson et al., 1998).

Interspecific competition

The inflationary effect has consequences for competitive interactions and species

coexistence. A local habitat might be a sink for a species not because the abiotic environment is

poor but because of the presence of a superior competitor. Inferior species can nonetheless

persist because of recurrent immigration from a source. Temporal variability in local growth

conditions could inflate the abundance of the inferior competitor, thus increasing its impact on

the superior competitor. Long et al. (2007) showed experimentally in a laboratory microcosm

using bacterivorous protists that this effect could sometimes lead to the local extirpation of the

locally superior (on average) species.

The inflationary effect may also lie at the foundation of a classic mechanism of

coexistence in metacommunities: the competition-colonization tradeoff. Classic models (e.g.,

Tilman, 1994) show that with strongly asymmetric competition, multiple species can coexist

provided there are external sources of extinction that wipe out superior competitors. This

requires however that patches be colonized asynchronously. If the superior species is a sluggish

colonizer, there is a time period during which fitness of the subordinate in the empty patch once

it arrives is high, and far distant locations are unlikely to be colonized simultaneously by the

superior species. Once the superior species arrives, however, the fitness for the subordinate is

low, and it stays low until the superior species suffers extinction. Thus, the dynamics of the

superior species in effect generates a positively autocorrelated environment for the fitness of the

inferior species. This can promote coexistence, provided there is asynchrony among patches in

extinction and subsequent recolonization (Roy et al., 2005).

The inflationary effect also underpins the spatiotemporal dynamics of the spatial

insurance hypothesis (Loreau et al., 2003; Shanafelt et al., 2015; Thompson et al., 2017). This

hypothesis posits that biodiversity provides stability to ecosystem functioning across a

metacommunity. This effect stems from the interaction between dispersal, differences among

competing species in their use of the environment, and spatiotemporal environmental

709 fluctuations. If different metacommunities experience distinct temporal patterns of

710 environmental conditions, species differentially adapted to the environment are expected to

711 thrive in any metacommunity (as a temporary source) for a finite period before the

712 environmental state of the patch switches. This results in spatiotemporal shifts in the source or

713 sink status of patches across the metacommunity. Dispersal ensures that the species adapted to

714 the new environmental conditions locally are available to replace less adapted species as the

715 environment changes. As a result, biodiversity may enhance and buffer ecosystem processes by

716 virtue of periods of inflationary growth and spatial exchanges among local ecosystems, even

717 when such effects do not occur in a closed homogeneous system. Factors that synchronize

718 population dynamics across ecosystems (Reuman et al., in press) or constrain dispersal will

719 erode the multispecies inflationary dynamics that contribute to metacommunity stability.

721 *Genetic variation and evolutionary dynamics.*

722  Our focus has been on ecological consequences of spatiotemporal variation and the

723 inflationary effect, but there are doubtless evolutionary effects, as well. Here, we mention just a

724 few examples. The model of Loreau et al. (2003), which focuses on species, can be interpreted as

725 a model of asexual selection maintaining genetic diversity in a spatiotemporal environment

726 (Levene, 1953). Wieczynski and Vasseur (2016) demonstrated that the inflationary effect could

727 promote the maintenance of intraspecific genetic variation when different genotypes are

728 differentially superior at distinct times, and showed that this could increase overall population

729 size and promote population persistence. Expression (4) describing the inflationary effect in

730 terms of the covariance between local growth rate and relative abundance can be viewed as a

731 component of fitness that selection can act upon. If a behavioral variant arises that increased the

match between the spatial distribution of that variant, and its temporarily greater local growth

rate, that variant would have a selective advantage. This is one way of describing the fitness

advantage of habitat selection, a point illustrated by Altenberg (2012) in the context of the

evolution of dispersal.

The inflationary effect has been shown to promote the persistence and evolution of more

contagious viral strains. In essence, patches with high transmission rates foster genetic diversity

in the pathogen population. New variants can then spread to areas with low transmission rates

and establish themselves in communities where other, less infectious strains may not persist.

Overall, this creates an environment that both promotes the evolution of new variants, and the

persistence of highly contagious variants in areas where community transmission could

otherwise be stopped, as suggested in Europe through the early stages of the COVID-19

pandemic (Lemey et al., 2021). In the winter of 2020, the B.1.1.7 variant was suspected to first

appear in the United Kingdom and Spain, where intervention measures were mostly lifted.

Nonetheless, this variant was able to rapidly spread across the continent and persist in countries

still undergoing lockdown measures because of its increased contagiousness.

Conservation and management

Drivers of the inflationary effect can be leveraged to inform conservation and

management decisions. The greatest threats to biodiversity --- global change, habitat destruction,

overexploitation, among others (Bellard et al., 2022; Brook et al., 2008) --- could act to increase

or decrease the strength of inflationary effects operating in nature. For instance, enhanced

weather variability could decrease the spatial synchrony of local population growth (Hansen et

al., 2020), thus increasing the inflationary effect. That said, increased incidence of extreme

climatic events (e.g., droughts and heatwaves) at large spatial scales could synchronize

populations (a generalized Moran effect; Hansen et al. 2020, see also Reuman et al. in press),

thereby weakening the inflationary effect.

Scientists already recognize the importance of incorporating metapopulation processes

for pursuing conservation priorities (Akçakaya et al., 2007), but considering the inflationary

effect emphasizes the critical stabilizing nature of metapopulations. Facilitating movement

between local populations (e.g., facilitated by corridors) is well established as beneficial for

conserving species at risk of extinction. However, the inflationary effect suggests that too much

movement might degrade asynchronous population fluctuations (to at least some degree).

Furthermore, the timing of reintroduction or restocking of populations, might be more effective

if it maintains or generates asynchronous dynamics.

Similar ideas can be applied to management of fire-obligate communities, the complexity

of which requires a carefully planned decision-making framework (Kelly et al., 2015). Managed

burning regimes might be made more effective by thinking about the heterogeneity or

homogeneity in burns created across the landscape by land managers, which may affect decisions

about the acreage per burning event, the timing and spatial arrangement of burns, and burn

intensity. All such factors influence the spatiotemporal heterogeneity affecting the dynamics of

species in fire-prone systems that might benefit (or rely) on the inflationary effect. Similar

concepts apply in fisheries. Prescriptions about quotas might include considerations for how they

promote or degrade asynchrony. Lack of these considerations might foster intense fishing on a

few populations with high numbers, which synchronizes population dynamics across large

scales, reducing the so-called population portfolio (Stier et al., 2020) and eroding stability

provided by the inflationary effect.

Infectious disease and epidemiology

While we have emphasized the inflationary effect for the spread of SARS-CoV-2, it

likely plays a role for many other pathogens, such as measles. Prior to vaccination in the late

1960s, measles had biennial epidemics that were synchronized throughout England and Wales.

Bolker and Grenfell (1996) documented how widespread vaccination eliminated large epidemics,

which were the putative causative synchronizing factor, and so decoupled measles dynamics

across space. Such asynchrony was hypothesized to reduce the chances of extinction of measles

across all geographies simultaneously. Indeed, despite consistent declines in incidence of

measles for three decades after vaccine introduction, measles incidence is far above elimination

and eradication targets (Winter et al., 2022).

Bubonic plague (caused by the bacterium *Yersinia pestis*), which occasionally causes

localized epidemics in humans, appears to be maintained by the inflationary effect in rat

populations (Keeling and Gilligan, 2000). Locally abundant susceptible populations of rats

(wherein the pathogenic bacterium's fitness is high) temporarily cause large outbreaks that spill

over and infect humans. Epidemics in the rat population happen differently across space and are

localized in time, leading to a patchwork of populations in the landscape with differential fitness

from the perspective of the pathogen --- with knock-on consequences for human health.

A final thought

This paper is an invited contribution to a special feature of The American Naturalist on

the theme of neglected and misunderstood ideas in ecology. One of the lessons to emerge from

the recent pandemic is that governmental policymakers must seriously confront the complicated

dynamical issues we bring out here. However, the situation is not unique to the recent pandemic.

Pathogen eradication is extremely rare among human pathogens, a fact we believe occurs in part

because of the large heterogeneities present at the scale of human societies. For emerging

infectious diseases, the lesson is to address exponentially growing processes quickly and

aggressively before they get out of hand. Beyond this, we submit that ignorance of the emergent

effects of spatiotemporal variability in local growth rates, given dispersal among locations,

contributed substantially to the failure of governmental control of the Covid pandemic in its early

phases in many parts of the globe. In New York, for instance, there was a 'block-by-block'

control strategy (Guarino, 2020). In the United Kingdom, then Prime Minister Boris Johnson

emphasized local lockdowns and local 'whack-a-mole' strategies (BBC 2020).

This absence of coordinated control efforts, at spatial scales from local municipalities up

to entire continents, given that there was continued movement of asymptomatic, infectious

individuals among locations—estimated to be responsible for up to 35% of all infections (Sah et

al., 2021)—likely contributed to the pervasive failure of control. Failure to recognize the

population dynamical consequences of spatiotemporal variability present in landscapes, with

movement across those landscapes, perhaps fostered a great deal of human suffering. One can

only hope that nations, and indeed the world, learned something from this experience.

Acknowledgements

We thank Nick Gotelli, Rachel Germain, and Sebastian Schreiber for useful discussions. NK,

MWS and RDH thank NSF grants for support, and RDH thanks the University of Florida

Foundation for continued support. AG is supported by the Liber Ero Chair in Biodiversity

Conservation and an NSERC Discovery Grant.

Data Availability Statement

Code to reproduce the figures here and in the supplementary material can be found in the Github

repository: https://github.com/kortessis/Asynchrony-and-Inflationary-Effect. The code will be

archived in Zenodo pending acceptance of the manuscript for publication, following any

requested changes by reviewers and editors.

References

Akçakaya, H.R., Mills, G., Doncaster, C.P., 2007. The role of metapopulations in conservation, in: Key Topics in Conservation Biology. Blackwell Publishing, pp. 64–84.

Altenberg, L., 2012. The evolution of dispersal in random environments and the principle of partial control. Ecological Monographs 82, 297–333. https://doi.org/10.1890/11-1136.1

Amarasekare, P., 2003. Competitive coexistence in spatially structured environments: a synthesis. Ecology Letters 6, 1109–1122. https://doi.org/10.1046/j.1461-0248.2003.00530.x

Andrewartha, H.G., Birch, L.C., 1984. The Ecological Web: More on the Distribution and Abundance of Animals. University of Chicago Press.

Andrewartha, H.G., Birch, L.C., 1954. The distribution and abundance of animals. University of Chicago Press.

Barfield, M., Orive, M.E., Holt, R.D., 2015. The role of pathogen shedding in linking within- and between-host pathogen dynamics. Mathematical Biosciences, From Within-Host Dynamics to the Epidemiology of Infectious Disease 270, 249–262. https://doi.org/10.1016/j.mbs.2015.04.010

BBC, 2020. Boris Johnson's "Weston lockdown" claims criticised. BBC News.

Bellard, C., Marino, C., Courchamp, F., 2022. Ranking threats to biodiversity and why it doesn't matter. Nat Commun 13, 2616. https://doi.org/10.1038/s41467-022-30339-y

Benaïm, M., Lobry, C., Sari, T., Strickler, É., 2023. Untangling the role of temporal and spatial variations in persistence of populations. Theoretical Population Biology 154, 1–26. https://doi.org/10.1016/j.tpb.2023.07.003

Berestycki, H., Desjardins, B., Weitz, J.S., Oury, J.-M., 2023. Epidemic modeling with heterogeneity and social diffusion. J. Math. Biol. 86, 60. https://doi.org/10.1007/s00285-022-01861-w

Bergmann, G.E., Leveau, J.H.J., 2022. A metacommunity ecology approach to understanding microbial community assembly in developing plant seeds. Frontiers in Microbiology 13.

Bernhardt, J.R., O'Connor, M.I., Sunday, J.M., Gonzalez, A., 2020. Life in fluctuating environments. Philosophical Transactions of the Royal Society B: Biological Sciences 375, 20190454. https://doi.org/10.1098/rstb.2019.0454

Blasius, B., Rudolf, L., Weithoff, G., Gaedke, U., Fussmann, G.F., 2020. Long-term cyclic
persistence in an experimental predator–prey system. Nature 577, 226–230.
https://doi.org/10.1038/s41586-019-1857-0
Bolker, B., Pacala, S.W., 1997. Using Moment Equations to Understand Stochastically Driven
Spatial Pattern Formation in Ecological Systems. Theoretical Population Biology 52,
179–197. https://doi.org/10.1006/tpbi.1997.1331
Bolker, B.M., Grenfell, B.T., 1996. Impact of vaccination on the spatial correlation and
persistence of measles dynamics. Proceedings of the National Academy of Sciences 93,
12648–12653. https://doi.org/10.1073/pnas.93.22.12648
Brook, B.W., Sodhi, N.S., Bradshaw, C.J.A., 2008. Synergies among extinction drivers under
global change. Trends in Ecology & Evolution 23, 453–460.
https://doi.org/10.1016/j.tree.2008.03.011
Chesson, P.L., 1985. Coexistence of competitors in spatially and temporally varying
environments: A look at the combined effects of different sorts of variability. Theoretical
Population Biology 28, 263–287. https://doi.org/10.1016/0040-5809(85)90030-9
Chesson, P.L., 1981. Models for spatially distributed populations: The effect of within-patch
variability. Theoretical Population Biology 19, 288–325. https://doi.org/10.1016/0040-
5809(81)90023-X
Childs, J.E., 2004. Zoonotic viruses of wildlife: hither from yon, in: Calisher, C.H., Griffin, D.E.
(Eds.), Emergence and Control of Zoonotic Viral Encephalitides, Archives of Virology.
Supplementa. Springer, Vienna, pp. 1–11. https://doi.org/10.1007/978-3-7091-0572-6_1
Cliff, A.D., Haggett, P., Smallman-Raynor, and M.R., 2000. Island Epidemics, Oxford
Geographical and Environmental Studies Series. Oxford University Press, Oxford, New
York.
den Boer, P.J., 1981. On the survival of populations in a heterogeneous and variable
environment. Oecologia 50, 39–53. https://doi.org/10.1007/BF00378792
den Boer, P.J., 1968. Spreading of risk and stabilization of animal numbers. Acta Biotheor 18,
165–194. https://doi.org/10.1007/BF01556726
Driscoll, D.A., Lindenmayer, D.B., 2009. Empirical tests of metacommunity theory using an
isolation gradient. Ecological Monographs 79, 485–501. https://doi.org/10.1890/08-
1114.1
Duncan, A.B., Gonzalez, A., Kaltz, O., 2013. Stochastic environmental fluctuations drive
epidemiology in experimental host–parasite metapopulations. Proceedings of the Royal
Society B: Biological Sciences 280, 20131747. https://doi.org/10.1098/rspb.2013.1747
Eksin, C., Ndeffo-Mbah, M., Weitz, J.S., 2021. Reacting to outbreaks at neighboring localities.
Journal of Theoretical Biology 520, 110632. https://doi.org/10.1016/j.jtbi.2021.110632
Gonzalez, A., Holt, R.D., 2002. The inflationary effects of environmental fluctuations in source–
sink systems. Proceedings of the National Academy of Sciences 99, 14872–14877.
https://doi.org/10.1073/pnas.232589299
Gonzalez, A., Lawton, J.H., Gilbert, F.S., Blackburn, T.M., Evans-Freke, I., 1998.
Metapopulation Dynamics, Abundance, and Distribution in a Microecosystem. Science
281, 2045–2047. https://doi.org/10.1126/science.281.5385.2045
Gotelli, N.J., 2008. A Primer of Ecology, Fourth Edition. ed. Oxford University Press, Oxford,
New York.

Gouhier, T.C., Guichard, F., Gonzalez, A., 2010. Synchrony and Stability of Food Webs in
906   Metacommunities. The American Naturalist 175, E16–E34.
907   https://doi.org/10.1086/649579
908   Gravel, D., Massol, F., Leibold, M.A., 2016. Stability and complexity in model meta-
909   ecosystems. Nat Commun 7, 12457. https://doi.org/10.1038/ncomms12457
910   Guarino, B., 2020. New York's block-by-block lockdowns are curbing covid-19. But residents
911   aren't pleased. Washington Post.
912   Hansen, B.B., Grøtan, V., Herfindal, I., Lee, A.M., 2020. The Moran effect revisited: spatial
913   population synchrony under global warming. Ecography 43, 1591–1602.
914   https://doi.org/10.1111/ecog.04962
915   Hanski, I., 1999. Metapopulation Ecology, Oxford Series in Ecology and Evolution. Oxford
916   University Press, Oxford, New York.
917   Hanski, I., 1998. Metapopulation dynamics. Nature 396, 41–49. https://doi.org/10.1038/23876
918   Hanski, I., 1997. 4 - Metapopulation Dynamics: From Concepts and Observations to Predictive
919   Models, in: Hanski, I., Gilpin, M.E. (Eds.), Metapopulation Biology. Academic Press,
920   San Diego, pp. 69–91. https://doi.org/10.1016/B978-012323445-2/50007-9
921   Hanski, I., Gaggiotti, O., 2004. 1 - Metapopulation Biology: Past, Present, and Future, in:
922   Hanski, I., Gaggiotti, O.E. (Eds.), Ecology, Genetics and Evolution of Metapopulations.
923   Academic Press, Burlington, pp. 3–22. https://doi.org/10.1016/B978-012323448-
924   3/50003-9
925   Hanski, I., Mononen, T., Ovaskainen, O., 2011. Eco-Evolutionary Metapopulation Dynamics and
926   the Spatial Scale of Adaptation. The American Naturalist 177, 29–43.
927   https://doi.org/10.1086/657625
928   Harrison, S., Quinn, J.F., 1989. Correlated Environments and the Persistence of
929   Metapopulations. Oikos 56, 293–298. https://doi.org/10.2307/3565613
930   Hassell, M.P., 2000. The spatial and temporal dynamics of host-parasitoid interactions, Oxford
931   series in ecology and evolution. Oxford University Press, Oxford ;
932   Hassell, M.P., Comins, H.N., Mayt, R.M., 1991. Spatial structure and chaos in insect population
933   dynamics. Nature 353, 255–258. https://doi.org/10.1038/353255a0
934   Hassell, M.P., May, R.M., 1988. Spatial heterogeneity and the dynamics of parasitoid-host
935   systems. Annales Zoologici Fennici 25, 55–61.
936   Hastings, A., 1980. Disturbance, coexistence, history, and competition for space. Theoretical
937   Population Biology 18, 363–373. https://doi.org/10.1016/0040-5809(80)90059-3
938   Hastings, A., 1977. Spatial heterogeneity and the stability of predator-prey systems. Theoretical
939   Population Biology 12, 37–48. https://doi.org/10.1016/0040-5809(77)90034-X
940   Holmes, E.E., Lewis, M.A., Banks, J.E., Veit, R.R., 1994. Partial Differential Equations in
941   Ecology: Spatial Interactions and Population Dynamics. Ecology 75, 17–29.
942   https://doi.org/10.2307/1939378
943   Holt, G., Chesson, P., 2016. Scale-Dependent Community Theory for Streams and Other Linear
944   Habitats. The American Naturalist 188, E59–E73. https://doi.org/10.1086/687525
945   Holt, R. d., Keitt, T. h., 2000. Alternative causes for range limits: a metapopulation perspective.
946   Ecology Letters 3, 41–47. https://doi.org/10.1046/j.1461-0248.2000.00116.x
947   Holt, R.D., 2010. Towards a trophic island biogeography: reflections on the interface of island
948   biogeography and food web ecology., in: The Theory of Island Biogeography Revisited.
949   Princeton University Press, Princeton, New Jersey, pp. 143–185.

Holt, R.D., 2000. A biogeographical and landscape perspective on within-host infection
dynamics., in: Proceedings of the 8th International Symposium of Microbial Ecology.
Presented at the Atlantic Canada Society for Microbial Ecology, Halifax, Canada, pp.
538–588.
Holt, R.D., 1997. 7 - From Metapopulation Dynamics to Community Structure: Some
Consequences of Spatial Heterogeneity, in: Hanski, I., Gilpin, M.E. (Eds.),
Metapopulation Biology. Academic Press, San Diego, pp. 149–164.
https://doi.org/10.1016/B978-012323445-2/50010-9
Holt, R.D., 1993. Ecology at the mesoscale : The influence of regional processes on local
communities, in: Species Diversity in Ecological Communities. University of Chicago
Press, pp. 77–88.
Holt, R.D., 1985. Population dynamics in two-patch environments: Some anomalous
consequences of an optimal habitat distribution. Theoretical Population Biology 28, 181–
208. https://doi.org/10.1016/0040-5809(85)90027-9
Holt, R.D., Barfield, M., Gonzalez, A., 2003. Impacts of environmental variability in open
populations and communities: "inflation" in sink environments. Theoretical Population
Biology, Understanding the role of environmental vatiation in population and community
dynamics 64, 315–330. https://doi.org/10.1016/S0040-5809(03)00087-X
Hoopes, M.F., Holt, R.D., Holyoak, M., 2005. The effects of spatial processes on two species
interactions., in: Metacommunities: Spatial Dynamics and Ecological Communities. The
University of Chicago Press, pp. 35–67.
Huffaker, C., 1958. Experimental studies on predation: dispersion factors and predator-prey
oscillations. Hilgardia 27, 343–383.
Hui, C., Richardson, D.M., 2017. Invasion dynamics: the spread and impact of alien organisms.
Oxford University Press.
Ingvarsson, Pärk., 2002. A metapopulation perspective on genetic diversity and differentiation in
partially self-fertilizing plants. Evolution 56, 2368–2373. https://doi.org/10.1111/j.0014-
3820.2002.tb00162.x
Johnson, E.C., Hastings, A., 2023. Coexistence in spatiotemporally fluctuating environments.
Theor Ecol 16, 59–92. https://doi.org/10.1007/s12080-022-00549-7
Katriel, G., 2022. Dispersal-induced growth in a time-periodic environment. J. Math. Biol. 85,
24. https://doi.org/10.1007/s00285-022-01791-7
Keddy, P.A., 1982. Population ecology on an environmental gradient: Cakile edentula on a sand
dune. Oecologia 52, 348–355. https://doi.org/10.1007/BF00367958
Keeling, M.J., Gilligan, C.A., 2000. Metapopulation dynamics of bubonic plague. Nature 407,
903–906. https://doi.org/10.1038/35038073
Keeling, M.J., Rohhani, P., 2007. Modeling Infectious Diseases in Humans and Animals.
Keeling, M.J., Wilson, H.B., Pacala, S.W., 2000. Reinterpreting Space, Time Lags, and
Functional Responses in Ecological Models. Science 290, 1758–1761.
https://doi.org/10.1126/science.290.5497.1758
Kelly, L.T., Bennett, A.F., Clarke, M.F., McCarthy, M.A., 2015. Optimal fire histories for
biodiversity conservation. Conservation Biology 29, 473–481.
https://doi.org/10.1111/cobi.12384
Kortessis, N., Simon, M.W., Barfield, M., Glass, G.E., Singer, B.H., Holt, R.D., 2020. The
interplay of movement and spatiotemporal variation in transmission degrades pandemic

control. Proceedings of the National Academy of Sciences 117, 30104–30106. https://doi.org/10.1073/pnas.2018286117

Leibold, M.A., 2018. Metacommunity ecology, Monographs in population biology. Princeton University Press, Princeton, New Jersey.

Leibold, M.A., Rudolph, F.J., Blanchet, F.G., De Meester, L., Gravel, D., Hartig, F., Peres-Neto, P., Shoemaker, L., Chase, J.M., 2022. The internal structure of metacommunities. Oikos 2022. https://doi.org/10.1111/oik.08618

Lemey, P., Ruktanonchai, N., Hong, S.L., Colizza, V., Poletto, C., Van den Broeck, F., Gill, M.S., Ji, X., Levasseur, A., Oude Munnink, B.B., Koopmans, M., Sadilek, A., Lai, S., Tatem, A.J., Baele, G., Suchard, M.A., Dellicour, S., 2021. Untangling introductions and persistence in COVID-19 resurgence in Europe. Nature 595, 713–717. https://doi.org/10.1038/s41586-021-03754-2

Levene, H., 1953. Genetic Equilibrium When More Than One Ecological Niche is Available. The American Naturalist 87, 331–333. https://doi.org/10.1086/281792

Levin, S.A., 1974. Dispersion and population interactions. The American Naturalist 108, 207–228.

Levins, R., 1970. Extinction. Lecture Notes in Biomathematics 2, 75–107.

Levins, R., 1969. Some Demographic and Genetic Consequences of Environmental Heterogeneity for Biological Control1. Bulletin of the Entomological Society of America 15, 237–240. https://doi.org/10.1093/besa/15.3.237

Long, Z.T., Petchey, O.L., Holt, R.D., 2007. The effects of immigration and environmental variability on the persistence of an inferior competitor. Ecology Letters 10, 574–585. https://doi.org/10.1111/j.1461-0248.2007.01049.x

Loreau, M., Mouquet, N., Gonzalez, A., 2003. Biodiversity as spatial insurance in heterogeneous landscapes. Proceedings of the National Academy of Sciences 100, 12765–12770. https://doi.org/10.1073/pnas.2235465100

Lu, M., 2021. Complex relationships between beta diversity and dispersal in meta-community models. Ecography 44, 1769–1780. https://doi.org/10.1111/ecog.05937

Matthews, D.P., Gonzalez, A., 2007. The Inflationary Effects of Environmental Fluctuations Ensure the Persistence of Sink Metapopulations. Ecology 88, 2848–2856. https://doi.org/10.1890/06-1107.1

May, R.M., 1972. Limit Cycles in Predator-Prey Communities. Science 177, 900–902.

Melbourne, B.A., Cornell, H.V., Davies, K.F., Dugaw, C.J., Elmendorf, S., Freestone, A.L., Hall, R.J., Harrison, S., Hastings, A., Holland, M., Holyoak, M., Lambrinos, J., Moore, K., Yokomizo, H., 2007. Invasion in a heterogeneous world: resistance, coexistence or hostile takeover? Ecology Letters 10, 77–94. https://doi.org/10.1111/j.1461-0248.2006.00987.x

Moore, S.M., Eisen, R.J., Monaghan, A., Mead, P., 2014. Meteorological Influences on the Seasonality of Lyme Disease in the United States. The American Journal of Tropical Medicine and Hygiene 90, 486–496. https://doi.org/10.4269/ajtmh.13-0180

Okubo, A., 1988. Diffusion-type models for avian range expansion, in: Acta XIX Congress Internationalis Ornithologici. National Museum of Natural Sciences, University of Ottawa Press, pp. 1038–1049.

Olivieri, I., Michalakis, Y., Gouyon, P.-H., 1995. Metapopulation Genetics and the Evolution of Dispersal. The American Naturalist 146, 202–228. https://doi.org/10.1086/285795

Ovaskainen, O., Hanski, I., 2004. 4 - Metapopulation Dynamics in Highly Fragmented Landscapes, in: Hanski, I., Gaggiotti, O.E. (Eds.), Ecology, Genetics and Evolution of Metapopulations. Academic Press, Burlington, pp. 73–103. https://doi.org/10.1016/B978-012323448-3/50006-4

Petchey, O.L., Gonzalez, A., Wilson, H.B., 1997. Effects on population persistence: the interaction between environmental noise colour, intraspecific competition and space. Proceedings of the Royal Society of London. Series B: Biological Sciences 264, 1841–1847. https://doi.org/10.1098/rspb.1997.0254

Petersen, L.R., Hayes, E.B., 2008. West Nile Virus in the Americas. Medical Clinics of North America, New and Emerging Infectious Diseases 92, 1307–1322. https://doi.org/10.1016/j.mcna.2008.07.004

Pulliam, H.R., 1988. Sources, Sinks, and Population Regulation. The American Naturalist 132, 652–661. https://doi.org/10.1086/284880

Reuman, D.C., Castorani, M.C.N., Cavanaugh, K.C., Sheppard, L.W., Walter, J.A., Bell, T.W., in press. How environmental drivers of spatial synchrony interact. Ecography n/a, e06795. https://doi.org/10.1111/ecog.06795

Roy, M., Holt, R.D., Barfield, M., 2005. Temporal Autocorrelation Can Enhance the Persistence and Abundance of Metapopulations Comprised of Coupled Sinks. The American Naturalist 166, 246–261. https://doi.org/10.1086/431286

Ruktanonchai, N.W., Floyd, J.R., Lai, S., Ruktanonchai, C.W., Sadilek, A., Rente-Lourenco, P., Ben, X., Carioli, A., Gwinn, J., Steele, J.E., Prosper, O., Schneider, A., Oplinger, A., Eastham, P., Tatem, A.J., 2020. Assessing the impact of coordinated COVID-19 exit strategies across Europe. Science 369, 1465–1470. https://doi.org/10.1126/science.abc5096

Runge, J.P., Runge, M.C., Nichols, J.D., 2006. The Role of Local Populations within a Landscape Context: Defining and Classifying Sources and Sinks. The American Naturalist 167, 925–938. https://doi.org/10.1086/503531

Sah, P., Fitzpatrick, M.C., Zimmer, C.F., Abdollahi, E., Juden-Kelly, L., Moghadas, S.M., Singer, B.H., Galvani, A.P., 2021. Asymptomatic SARS-CoV-2 infection: A systematic review and meta-analysis. Proceedings of the National Academy of Sciences 118, e2109229118. https://doi.org/10.1073/pnas.2109229118

Schmidt, K.A., 2004. Site fidelity in temporally correlated environments enhances population persistence. Ecology Letters 7, 176–184. https://doi.org/10.1111/j.1461-0248.2003.00565.x

Schreiber, S.J., 2023. Partitioning the impacts of spatial-temporal variation in demography and dispersal on metapopulation growth rates.

Schreiber, S.J., 2012. The Evolution of Patch Selection in Stochastic Environments. The American Naturalist 180, 17–34. https://doi.org/10.1086/665655

Schreiber, S.J., 2010. Interactive effects of temporal correlations, spatial heterogeneity and dispersal on population persistence. Proceedings of the Royal Society B: Biological Sciences 277, 1907–1914. https://doi.org/10.1098/rspb.2009.2006

Schreiber, S.J., Hening, A., Nguyen, D.H., 2023. Coevolution of Patch Selection in Stochastic Environments. The American Naturalist 202, 122–139. https://doi.org/10.1086/725079

Sender, R., Bar-On, Y.M., Gleizer, S., Bernshtein, B., Flamholz, A., Phillips, R., Milo, R., 2021. The total number and mass of SARS-CoV-2 virions. Proceedings of the National Academy of Sciences 118, e2024815118. https://doi.org/10.1073/pnas.2024815118

Shanafelt, D.W., Dieckmann, U., Jonas, M., Franklin, O., Loreau, M., Perrings, C., 2015.
Biodiversity, productivity, and the spatial insurance hypothesis revisited. Journal of
Theoretical Biology 380, 426–435. https://doi.org/10.1016/j.jtbi.2015.06.017
Shigesada, N., Kawasaki, K., Takeda, Y., 1995. Modeling Stratified Diffusion in Biological
Invasions. The American Naturalist 146, 229–251. https://doi.org/10.1086/285796
Shoemaker, L.G., Hallett, L.M., Zhao, L., Reuman, D.C., Wang, S., Cottingham, K.L., Hobbs,
R.J., Castorani, M.C.N., Downing, A.L., Dudney, J.C., Fey, S.B., Gherardi, L.A., Lany,
N., Portales-Reyes, C., Rypel, A.L., Sheppard, L.W., Walter, J.A., Suding, K.N., 2022.
The long and the short of it: Mechanisms of synchronous and compensatory dynamics
across temporal scales. Ecology 103, e3650. https://doi.org/10.1002/ecy.3650
Shone, S.M., Curriero, F.C., Lesser, C.R., Glass, G.E., 2006. Characterizing Population
Dynamics of Aedes sollicitans (Diptera: Culicidae) Using Meteorological Data. Journal
of Medical Entomology 43, 393–402. https://doi.org/10.1093/jmedent/43.2.393
Stier, A.C., Olaf Shelton, A., Samhouri, J.F., Feist, B.E., Levin, P.S., 2020. Fishing,
environment, and the erosion of a population portfolio. Ecosphere 11, e03283.
https://doi.org/10.1002/ecs2.3283
Thompson, P.L., Rayfield, B., Gonzalez, A., 2017. Loss of habitat and connectivity erodes
species diversity, ecosystem functioning, and stability in metacommunity networks.
Ecography 40, 98–108. https://doi.org/10.1111/ecog.02558
Tilman, D., 1994. Competition and Biodiversity in Spatially Structured Habitats. Ecology 75, 2–
16. https://doi.org/10.2307/1939377
Tilman, D., Lehman, C.L., Yin, C., 1997. Habitat Destruction, Dispersal, and Deterministic
Extinction in Competitive Communities. The American Naturalist 149, 407–435.
Van Horne, B., 1983. Density as a Misleading Indicator of Habitat Quality. The Journal of
Wildlife Management 47, 893–901. https://doi.org/10.2307/3808148
Vasseur, D.A., Fox, J.W., 2009. Phase-locking and environmental fluctuations generate
synchrony in a predator–prey community. Nature 460, 1007–1010.
https://doi.org/10.1038/nature08208
Wang, Q., Wu, H., 2022. There exists the "smartest" movement rate to control the epidemic
rather than "city lockdown." Applied Mathematical Modelling 106, 696–714.
https://doi.org/10.1016/j.apm.2022.02.018
Wang, S., Altermatt, F., 2019. Metapopulations revisited: the area-dependence of dispersal
matters. Ecology 100, e02792. https://doi.org/10.1002/ecy.2792
Wieczynski, D.J., Vasseur, D.A., 2016. Environmental fluctuations promote intraspecific
diversity and population persistence via inflationary effects. Oikos 125, 1173–1181.
https://doi.org/10.1111/oik.02725
Williams, P.D., Hastings, A., 2011. Paradoxical persistence through mixed-system dynamics:
towards a unified perspective of reversal behaviours in evolutionary ecology. Proc Biol
Sci 278, 1281–1290. https://doi.org/10.1098/rspb.2010.2074
Wilson, D.S., 1992. Complex Interactions in Metacommunities, with Implications for
Biodiversity and Higher Levels of Selection. Ecology 73, 1984–2000.
https://doi.org/10.2307/1941449
Wilson, H.B., Hassell, M.P., Holt, R.D., 1998. Persistence and Area Effects in a Stochastic
Tritrophic Model. The American Naturalist 151, 587–595.
https://doi.org/10.1086/286144

Winter, A.K., Lambert, B., Klein, D., Klepac, P., Papadopoulos, T., Truelove, S., Burgess, C.,
Santos, H., Knapp, J.K., Reef, S.E., Kayembe, L.K., Shendale, S., Kretsinger, K., Lessler,
J., Vynnycky, E., McCarthy, K., Ferrari, M., Jit, M., 2022. Feasibility of measles and
rubella vaccination programmes for disease elimination: a modelling study. The Lancet
Global Health 10, e1412–e1422. https://doi.org/10.1016/S2214-109X(22)00335-7
Zhang, B., Liang, S., Wang, G., Zhang, C., Chen, C., Zou, M., Shen, W., Long, H., He, D., Shu,
Y., Du, X., 2021. Synchronized nonpharmaceutical interventions for the control of
COVID-19. Nonlinear Dyn 106, 1477–1489. https://doi.org/10.1007/s11071-021-06505-
0
1140

References cited only in the Supplementary Material

Casella, G. and R.J. Berger. 2001. Statistical Inference. Cengage Learning.
Ghosh, E. B., K. A. Brakora, and A. H. Lee. 2011. Evolution of ruminant headgear: a review.
Proceedings of the Royal Society B 278:2857–2865.
Kortessis, N. and P. Chesson. 2019. Germination variation facilitates the evolution of seed
dormancy when coupled with seedling competition. Theoretical Population Biology
130:60-73.
Kortessis, N. and P. Chesson. 2021. Character displacement in the presence of multiple trait
differences: Evolution of the storage effect in germination and growth. Theoretical
Population Biology 140:54-66.
1151

1152 **Tables**
1153
1154 Table 1. Features indicative of the inflationary effect in natural systems.

Empirical Pattern	Link to Inflationary Effect Measure (eqn 4)
Nonrandom distributions of individuals across space	Spatial variation in relative density, v_x
Temporal changes in the non-random distribution of individuals over space	Variation in local density across space and time, $v_x(t)$
Spatiotemporal variation in a fitness-affecting factor	Variation in local growth rates across space and time, $r_x(t)$
Undirected passive dispersal or active environmental tracking	Possibility that population is concentrated in locations and times of higher than average growth, (positive $\text{cov}(r_x(t), v_x(t))$, on average)

1155

1156　**Figures**

1157
Figure 1. A hypothetical landscape with temporal and spatial variability in some fitness factor.

(a) Spatial locations are aligned vertically and time points are aligned horizontally. Averaging

values across time (i.e., horizontally) gives $\tilde{F}_x$, a perspective that illuminates average differences

across locations. Averaging values across space (i.e., vertically) gives $\bar{F}_t$, a perspective that

illuminates average differences over time. The remaining variation is spatiotemporal and reflects

differences in the pattern of variation across spatial locations. (b) The decomposition of the total

variation across space and time in the fitness factor for the landscape in (a).

1165
Figure 2. Finite patch version of Levins' metapopulation model under different assumptions

about asynchrony of local patch extinctions. Dynamics under the conventional assumption that

environments influencing extinction are uncorrelated, i.e., asynchronous, with $\rho_E = 0$, are given

by the blue lines. Dynamics allowing for correlated extinctions, with $\rho_E = 0.75$, are shown by the

green lines. Simulation details found in supplement. For all dynamics, $e = 0.25$, and $c = 0.5$ such

that the infinite patch model predicts equilibrium occupancy of $p^* = 1 - e/c = 0.5$ (dashed lines).

1172

1173

Figure 3. Effect of correlated extinction events on metapopulation stability. (a) Metapopulation

extinction times for replicate simulations, and (b) corresponding conditional mean occupancy.

Squares connected by lines are the geometric mean extinction times in (a) and mean conditional

mean occupancy in (b) over all replicate simulations. The parameter ρ_E refers to the correlation

coefficient for all pairwise latent variables describing environmental variables determining

extinction in patches (details in the supplementary material S2). All simulations had 25 patches,

13 of which were initially occupied. There are 100 replicate simulations for each ρ_E value. Other

parameters: $e = 0.25$, and $c = 0.5$.

1185
1186

Figure 4. Sink population dynamics with fluctuating growth rate and a constant immigration.

Dynamics follow $N(t+1) = N(t)e^{r(t)} + I$, where $r(t)$ is the per-capita growth rate without

immigration, and I the density of immigrants. Here $r(t) = \bar{r} + \sigma_r X(t)$, where $X(t)$ is follows a

standard normal distribution. We chose $\bar{r} = -0.2$ and $I = 1 - e^{-0.2} \approx 0.18$ such that the equilibrium

density of the sink in a constant environment is 1 (heavy black line). All cases start with $N(t=0)$

= 1 and have identical sequences of $X(t)$. Further details in supplementary material S3.

1193

1194

1195
1196   Figure 5. The inflationary effect in a two-patch SIR model. (a) The metapopulation-scale growth

1197   rate and how it depends on the overlap in the application of NPIs and host dispersal. (b) Example

1198   dynamics from a stochastic version of the model. Parameters: infectious duration = 4.5 days, β_0 =

1199   0.375 d^{-1} (hence $R_0 \approx 1.7$), $T = 30$, and $\epsilon = 0.95$. In (b), $m = 0.025$ d^{-1}. Further details in

1200   supplementary material S5.