Molecular epidemiology of Cercospora leaf spot on resistant and susceptible sugar beet

hybrids

Chen Chen[1], Harald Keunecke[2], Enzo Neu[2], Friedrich J. Kopisch-Obuch[2], Bruce A. McDonald[1],

Jessica Stapley[1]

[1]Plant Pathology Group, Institute of Integrative Biology, ETH Zürich, 8092 Zürich, Switzerland

[2]KWS SAAT SE & Co. KGaA, Grimsehlstr. 31, 37574 Einbeck, Germany

Corresponding author: Chen Chen, chen.chen@usys.ethz.ch

Abstract

Cercospora leaf spot (CLS), caused by *Cercospora beticola*, is a major foliar disease impacting

sugar beet production worldwide. The development of new resistant sugar beet hybrids

provides a powerful tool to better manage the disease, but it is unclear how these hybrids

affect CLS epidemiology. We used a molecular epidemiology approach to dissect natural

epidemics of CLS affecting two susceptible and two resistant sugar beet hybrids with

different levels of resistance at two field sites. Infected plants were geotagged on a weekly

basis. Isolations of *C. beticola* were made from infected leaves and genotyped using six

simple sequence repeat loci to identify clones. We determined that CLS epidemics had a

later onset in plots planted to resistant hybrids, but once the pathogen established an

infection, there was little difference between resistant and susceptible hybrids in the

probability of localized spread and dispersal. We found that different clones often infected

the same leaf and that clusters of infected plants were often colonized by a mixture of

clones. There was little overall difference in genetic diversity between resistant and

susceptible hybrids, however genotypic evenness differed at one site. In this site we found

one genotype restricted to the resistant cultivars at a high frequency. We established that

infections were not randomly distributed across the fields and we found that a single clone

could spread over a distance of 100 m during a growing season.

Keywords: *Beta vulgaris*, *Cercospora beticola*, population genetics, cercospora leaf spot,

resistant cultivars

Introduction

Sugar beet (*Beta vulgaris* L.) contributes approximately one-fifth of global sucrose production annually (Rangel et al., 2020). The most destructive disease of sugar beet is the foliar disease Cercospora leaf spot (CLS) caused by the fungus *Cercospora beticola*. CLS affects the production of sugar beet in almost all sugar beet growing regions worldwide (Holtschulte, 2000). Under high disease pressure, the yield loss can exceed 40% (Shane and Teng, 1992). The main method of control is multiple applications of fungicides, however this has resulted in the emergence of widespread fungicide resistance in *C. beticola* populations (Bolton et al., 2013). Resistant cultivars are seen as an important means of control in addition to fungicides, with CLS-resistant cultivars providing protection under low to moderate disease pressure (Francis and Luterbacher, 2003; Vogel et al., 2018; Weiland and Koch, 2004; Wolf and Verreet, 2005). A new resistance gene named *BvCR4* has recently been identified and bred into sugar beets (Chen et al., 2023). This new gene has shown a high level of CLS resistance and could be a useful method of control.

As with the evolution of fungicide resistance, pathogens can evolve to overcome host resistance through selection for virulent strains. How quickly pathogens evolve virulence to resistant cultivars is strongly influenced by the genetic architecture of the resistance – single major resistance (R) genes with large effect are usually less durable compared to quantitative resistance encoded by several genes with smaller, additive effects (Brown, 2015). The evolution of virulence is also influenced by the amount of genetic diversity in the selected pathogen population, the deployment strategies used for the resistance genes, and the distribution of virulent pathogen genotypes across a heterogenous host population (Milgroom and Peever, 2003; Stam and McDonald, 2018).

The interaction between pathogen genotypes and host genotypes shapes both the development of epidemics at the field scale and the evolutionary landscape of the entire pathosystem at larger spatial and temporal scales. High genetic diversity within a pathogen population can promote the evolution of virulence (McDonald and Linde, 2002a), but the type and degree of host resistance can influence the distribution of genetic diversity in the corresponding pathogen population. When a new resistant cultivar carrying a major R gene is first deployed in the field, the corresponding pathogen population experiences a selection

bottleneck, because only genotypes that carry the compatible virulence factors can infect

the resistant cultivar. This selective bottleneck will likely result in a reduction in genetic

diversity in the pathogen population. If the resistance of the cultivar is quantitative,

selection is likely to have less effect on the overall genetic diversity, and genetic variation at

the selected loci can be maintained (Cowger and Brown, 2019). How quickly the pathogen

population and the corresponding genetic diversity can recover following this selection

bottleneck depends upon the life history and evolutionary potential of the pathogen - in

particular the frequency of sexual reproduction and the efficiency of dispersal of virulent

genotypes (Barrett et al., 2008; McDonald and Linde, 2002b). The subsequent population

recovery and dispersal are also likely to promote rapid evolution. The rate of sexual

reproduction in *C. beticola* is unknown as no sexual stage (teleomorph) has yet been

described, but the presence of both mating types at the *MAT* locus, together with high

genotypic diversity suggests that it undergoes regular sexual cycles (Bolton et al., 2012;

Groenewald et al., 2006, 2007b; Vaghefi et al., 2017a). The rapid evolution of fungicide

resistance in CLS indicates that *C. beticola* has a high evolutionary potential (Bolton et al.,

2013; Pethybridge et al., 2020; Trkulja et al., 2017), suggesting that *C. beticola* populations

could also adapt quickly to deployment of resistant host genotypes.

The epidemiology of CLS has been well studied (Imbusch et al., 2021; Rossi et al., 2000;

Tedford et al., 2018; Wolf and Verreet, 2005). *C. beticola* is polycyclic, with the primary

inoculum coming from four known sources, including overwintering structures called

pseudostromata which can persist on plant debris for two years (Khan et al., 2008),

windborne conidia blowing in from outside newly planted fields, conidia produced on

alternative hosts growing in or nearby new fields, and infested seeds (Spanner et al., 2022).

In as little as 7 days these first infections can reproduce asexually to form conidia that

establish secondary infections (Rangel et al., 2020). These conidia can be dispersed to

adjacent plants by rainsplash, wind and insects (Khan et al., 2008; Vereijssen et al., 2007;

Weiland and Koch, 2004), with the longest recorded dispersal distance exceeding 45 m

(Imbusch et al., 2021). On resistant hosts, the latent period did not change, but the spore

yield was reduced (Rossi et al., 2000). This finding illustrates that resistant hosts may

influence CLS epidemiology, which may in turn affect the emergence and distribution of

virulent genotypes.

Previous work established that *C. beticola* has high genetic diversity and a high evolutionary

potential, and that host resistance may influence aspects of CLS epidemiology, but few

studies have investigated the evolution of virulence in *C. beticola* populations or used

neutral genetic markers to understand how CLS epidemics develop in fields during a growing

season. Molecular epidemiology bridges the gaps between disease epidemiology and

pathogen population genetics. By including genetic markers into epidemiology studies, new

research questions can be addressed around the origins of epidemics, pathogen

movements, dispersal pathways and the spatio-temporal aspects of disease progression

(Archie et al., 2009; Desbiez et al., 2002; Xhaard et al., 2012).

Here we analyzed the molecular epidemiology of natural CLS epidemics occurring in

resistant and susceptible hybrids of sugar beet to better understand the evolutionary

potential of this pathogen on newly developed hybrids. The main objects were 1) to

determine the time of onset of CLS epidemics on different hybrids; 2) to measure the local

dispersal pattern for both primary and secondary inoculum of CLS; 3) to assess the genetic

diversity of *C. beticola* populations during epidemics across different spatial and temporal

scales, including: on the same leaves; on neighboring plants; and across hybrids within and

between fields; 4) to determine the degree of clonality on different host populations and

identify the most likely sources of primary and secondary inoculum in natural field

infections.

Methods

Field trials in Switzerland

Two trial sites were located in Switzerland: Rudolfingen, Trüllikon 47°38'18.6"N, 8°40'46.1"E

and Hendschiken, Othmarsingen 47°23'40.4"N, 8°13'41.1"E. At both sites, four sugar beet

hybrids were planted in March 2020 into single fields that had not been planted with sugar

beets in the previous three years. The four hybrids differed in their resistance to CLS,

including two highly resistant hybrids with the *BvCR4* gene – Hybrid_R2 and Hybrid_R3, and

two susceptible hybrids lacking *BvCR4*– Hybrid_S2 and Hybrid_S3 (Table 1). In Hendschiken,

12 rows of each hybrid were planted in each plot in a field covering 3600 m^2. In Rudolfingen,

18 rows of each hybrid were planted in each plot in a field covering 1950 m^2. Rows were

separated by approximately 45 cm in both locations (Figure S1). Sugar beets were harvested

in early November. No fungicides were applied at either field site and all CLS infection was

natural.

Temporal sampling strategy

Our sampling was designed to identify the naturally infected plants in each plot and

determine how natural infection spreads among plants. We systematically searched each

plot for infected plants once per week from July 21st to September 24th 2020. Individual

sugar beet plants served as sampling units and one leaf with CLS lesions per plant was

collected. The sampling design is shown in Figure S1. All sampled plants were marked with

flags to prevent resampling. In the early stages of the epidemic, we found single newly

infected plants surrounded by healthy plants, which we referred to as solitary infections. As

the epidemic progressed, the disease often spread from a single infected plant to

neighboring plants to form a cluster of infected plants, hereafter called a hotspot. For each

solitary infection, we recorded the GPS location (Garmin) and gave it a two-letter code (e.g.

AA) and the number 1 to indicate it was the first infected plant at that GPS position (e.g.

AA1). If the infected plant was adjacent to a previously identified and flag-labeled plant, it

was considered part of the same hotspot. Infected plants in the same hotspots were given

the same alphabetic code and a number to indicate the sampled plants were the 2nd, 3rd,

4th or 5th plant found in the hotspot (e.g. AA4 was the fourth plant sampled in the AA

hotspot). One leaf with CLS lesions was taken from each plant, with a preference for young

and otherwise healthy leaves with more than one clear CLS lesion. The excised leaves were

kept flat and cool until they were transported back to the laboratory for *C. beticola* isolation.

We sampled a maximum of five plants from each hotspot. We stopped sampling a hybrid

plot when it was no longer possible to find new solitary infected plants because hotspots

had coalesced. Our sampling design allowed us to track CLS epidemic progression and track

hotspot formation within and between plots planted to resistant and susceptible hybrids.

Leaf storage

In the field, the leaves were stored flat and cool until being transported back to the lab. To

store leaves flat, each leaf was placed on a labeled piece of paper and covered with a dry

paper towel. Multiple leaves were stacked between stiff cardboard sheets and secured with

strong rubber bands. The stacked leaves were then stored in a cool box with ice blocks to

maintain a low temperature during transport. Upon arrival at the laboratory, the leaves

were air dried at room temperature in a drying cupboard. Dried leaves were stacked and

placed in plastic bags with anhydrous silica gel and stored at 4°C until used for pathogen

isolation.

Pathogen isolation and DNA extraction

To obtain pure strains of *C. beticola*, spores were collected from the dried leaves. Three or

four pieces of leaf tissue with visible lesions were cut from each dried leaf. The tissue

samples were surface sterilized by immersing them in a 2% sodium hypochlorite solution for

3 minutes, followed by two rinses in sterile de-ionized water. The leaf pieces were placed on

dry paper towels to remove excess water, then were put into glass petri dishes on an

elevated sterile net placed on a piece of moist filter paper. The petri dishes were sealed with

parafilm and incubated at 24°C for 48-72 hours to induce sporulation. The induced lesions

were checked for the typical black pseudostromata and hyaline conidia under a dissecting

microscope (Olympus Schweiz AG). The spores were picked from sporulating lesions using

hand-made fine glass needles and transferred to fresh potato dextrose agar (PDA, Difco).

The plates were cultured under constant darkness at 24°C for 7-10 days. After a

characteristic *C. beticola* colony had formed, a 5-mm agar plug was excised from the edge of

each colony and transferred to a new PDA plate for 14 days to form a pure culture. Any

contaminated plates were discarded. For long-term storage of each pure culture, 4-6 agar

plugs were cut from the edges of a 2-week-old pure culture and put into cryotubes in

duplicate - with and without anhydrous silica gel, and stored at 4°C. For most leaves,

between 1-3 colonies were isolated from different lesions, but for some leaves, more than 3

strains were isolated to investigate pathogen diversity within the same leaf. In total, 1251 *C.*

beticola isolations were made. For practical reasons, not all isolates were transferred into

long-term storage. Instead, we aimed for a single isolation per leaf and preference was

given to isolates from hotpots, which were expected to be more virulent and capable of

dispersing to nearby healthy plants.

To obtain fresh mycelia for DNA extraction, we transferred hyphae from plates to liquid

culture by scraping small pieces of hyphae from the edge of each pure culture and placing

them into 150 mL flasks containing 50 mL potato dextrose broth (PDB, Difco). The liquid

cultures were incubated in the dark at 24°C on a shaker at 120 rpm for 10 days. The fungal

tissue was harvested, washed twice with sterilized water, and the supernatant was

discarded after centrifuging (Allegra X-12R, Beckman Coulter) at 3750 rpm for 10 mins. The

pellet was then transferred to 2 mL microcentrifuge tubes and lyophilized overnight. The

lyophilized pellets were ground into a fine powder using ~15 glass beads in a grinding

machine (Qiagen MM300 Retsch TissueLyser). A total of 560 DNA samples of isolates coming

from separate leaves were extracted using an automated DNA extraction robot, the

Kingfisher Flex (Thermo Fisher Scientific), using the recommended reagents and a protocol

for fungal samples (LGC Genomics GmbH). The resulting DNA samples were quantified using

a Nanodrop spectrophotometer (Thermo Fisher Scientific). We performed a polymerase

chain reaction (PCR) using species-specific primers (Knight and Pethybridge, 2020) to

confirm the samples were *C. beticola*. Only samples with a clear band at around 200 bp

(~85%) were retained.

Simple sequence repeats analysis

The six SSRs were amplified in 2 multiplexes (multiplex1: SSRCb1, SSRCb3, SSRCb21;

multiplex2: SSRCb22, SSRCb25, SSRCb27) (Groenewald et al., 2007a; Vaghefi et al., 2017b).

The SSR primers were blasted against the reference genome to ensure they were not

physically linked (Table S1). The PCR amplifications were performed using Qiagen Type-it

kits (Qiagen) in a total volume of 11 µL. The reaction mixture contained 20 ng of template

DNA, 0.4 µM of each forward and reverse primer, 4 µL Type-it mix (with buffer and MgCl$_2$), 1

µL Q-solution (1 U Taq polymerase), and 2 µl deionized water. The initial denaturation was

carried out at 95°C for 5 mins, followed by 30 cycles of denaturation at 95°C for 30 s,

annealing at 58°C or 56°C for 90 s, elongation at 72°C for 30 s and final extension at 60°C for

30 mins. The PCR amplicons were analyzed on an ABI 3730xl DNA Analyzer (Applied

Biosystems) at the Genetic Diversity Centre (GDC) of ETH Zürich. Multiple samples were

repeated at different stages to create technical replicates in order to estimate SSR

genotyping error rates and ensure consistent allele calling.

SSR allele size was scored using the R package *Fragman* (Covarrubias-Pazaran et al., 2016).

To reduce the genotyping error, allele fragment sizes of each SSR marker were visualized

and binned using the R package MsatAllele v 1.05 (Alberto, 2009). Fragment stutter patterns

varied between the SSRs. Visual inspection of the allele calling and consistency between

repeats was used to assess the accuracy of the MsatAllele automatic binning. Automatic

binning was accurate for three SSRs (SSRCb27, SSRCb3, SSRCb21) but manual adjustment of

the bin sizes was required for the rest. The consistency of allele calling was evaluated using

isolates that were replicated (n=51). In total we obtained SSR genotypes for 475 isolates.

Genotype calling was repeated for 20% of isolates chosen at random. Samples with multiple

allele calls were assumed to be a mixture of strains and were excluded from the dataset. All

individuals with missing SSR data were also removed, leaving 340 fully genotyped isolates

(158 from Hendschiken and 182 from Rudolfingen) for downstream analyses.

Statistical analysis

Statistical analysis was performed in R version 4.3.1 (R Core Team, 2023) and R Studio

version 2023.6.0.421 (Posit team, 2023). To investigate the localized spread between plants

to form a hotspot, we used a generalized linear model with a binomial distribution. To

compare across hybrids, we needed to control for the fact that CLS continued to spread

after we stopped sampling in some hybrid plots, meaning that any new infection cases that

were identified at the end of the sampling period were not followed to determine if they

spread to neighboring plants. Our solution was to right-censor the data by excluding the

new infection cases observed 1 and 2 weeks before the final sampling date in each hybrid

plot. This is a common practice when sampling ends before an event has been observed.

As the sampling period varied for each hybrid, the right-censor date also varied (Table 2).

The rate at which a hotspot developed was analyzed using a linear model. The pairwise

distances between geotagged infection sites were calculated using the R package *geosphere*

v1.5-18 (Hijmans, R. J., 2022). Whether the distributions of geotagged plants differed

between hybrids or between the beginning and the end of the season were assessed using

Kolmogorov-Smirnov (K-S) tests in the R package R4PDE (Del Ponte, 2023). Multi-locus

genotypes (MLGs) were identified and analyzed with the R package *poppr* v.2.9.3 (Kamvar et

al., 2015, 2014). We performed MLG analysis separately for each hybrid in each site to

compare the epidemics between them.

We performed clone-correction for each site separately as our earlier analyses indicated it

was unlikely to observe the same MLG between two sites (whole genome sequence data of

256 isolates found only one case of the same clone present at the two sites (Chen et al.,

2023)). The number of alleles, number of unique MLGs, the expected number of MLGs

controlling for differences in sample size (eMLG) and Nei's gene diversity, Simpson's

diversity and evenness were calculated using the R package *poppr* v.2.9.3 for each hybrid

and site. Bootstrapped confidence intervals were calculated for genetic diversity statistics

using function diversity_ci in *poppr*. Clonal fraction was calculated as (N–number of

MLGs)/N, where N equals the number of isolates.

Results

Onset and distribution of CLS disease

The first symptoms of *C. beticola* were observed in the susceptible hybrids, for Hybrid_S2 on

July 21st and one week later in Hybrid_S3. Three weeks later, the first infections appeared

on the resistant hybrids (Table 2). The appearance of the first *C. beticola* infections on

resistant hybrids with *BvCR4* varied between fields. In Hendschiken, the disease appeared

on Hybrid_R2 before Hybrid_R3, while in Rudolfingen, the first symptoms were observed

later in Hybrid_R2. GPS coordinates were used to map the locations of infected plants. The

spatial distribution of all infection cases is shown in Figure 1.

The accumulation of geotagged infection cases is shown in Figure 2. The sampling period

varied for each hybrid because the dates of disease onset differed (Table 2) and because we

stopped sampling when the hotspots coalesced and solitary infections could no longer be

clearly distinguished. Thus, we stopped sampling first in Hybrid_S2, then Hybrid_S3 and

then Hybrid_R2. Sampling continued until our final visit (September 24) in Hybrid_R3 as

hotspots had not coalesced. Infections continued to increase in all hybrids after we stopped

sampling. It should be noted that due to some initial sampling problems, we missed the

start of the epidemic in Hybrid_S3. Hence caution must be taken when comparing the

temporal spread in Hybrid_S3 with other hybrids. By the end of the season, 454 plants were

sampled (one leaf per plant) from 196 geotagged infections in Hendschiken, and 503 plants

were sampled from 209 geotagged infections in Rudolfingen.

Localized spread of CLS in each hybrid

Across all hybrids, about 80% of infection cases spread to at least one neighboring plant,

resulting in the formation of hotspots (Table 3). There was no significant difference between

fields or hybrids in the probability that CLS would spread from one solitary infected plant to

neighboring plants to form a hotspot of two or more infected plants, or to form a large

hotspot with at least five infected plants (hotspot: Field $F_{(1,209)}$ =1.23, p=0.26; Hybrid $F_{(3,206)}$

=4.84, p=0.1, Large hotspot Field $F_{(1,209)}$ =2.69, p=0.10; Hybrid $F_{(3,206)}$ =4.90, p=0.17). NB: the

numbers in Table 3 differ slightly compared to the total cases and hotspots (Figure 2) due to

the right-censoring of datasets.

To assess the rate of hotspot development in each hybrid, we calculated the rate of spread

as the number of infected plants divided by the number of days between sampling periods

for plants sampled in each hotspot. There was no difference in the rate of spread between

sites (Field $F_{(1,115)}$ =2.87, p=0.09), there was a significant effect of hybrid (Hybrid $F_{(3,115)}$ =6.5,

p<0.001) and no interaction (Field*Hybrid $F_{(3,115)}$ =2.17, p=0.09). In Hendschiken, the rate of

spread was slower in Hybrid_R3 and faster in Hybrid_R2 compared to Hybrid_S2

(Hybrid_R2: t=3.95, p=0.002, Hybrid_R3: t=-2.73, p=0.008). In Rudolfingen, there was no

difference in the rate of spread, although Hybrid_R3 tended to be slower (t=-1.82, p=0.07).

Spatial distribution of infections

We compared the distribution of geotagged infections at the beginning and the end of the

epidemic and across hybrids within each site. The first two sampling dates in each hybrid

plot were designated as the beginning of the epidemic and were used to investigate the

spatial distribution of the first CLS infections. The initial distributions of hotspots and

infection sites for most hybrids were randomly distributed (Table S2, p >0.10) with two

exceptions: Hybrid_S2 in Hendschiken and Hybrid_R2 in Rudolfingen. During the epidemic,

the minimum distance between infection sites decreased or remained constant, while the

maximum distance increased throughout each epidemic (Table 4). In some cases, the GPS

was not recorded for an infection site, resulting in smaller numbers in Table 4 compared to

Figure 2. The final distributions of hotspots and infection sites for most hybrids were

aggregated (non-random) (Table S2, p <0.10) with two exceptions: Hybrid_R2 in

Hendschiken and Hybrid_S2 in Rudolfingen, where the infection sites were randomly

distributed. Although the number of infection cases increased over the season, there was

little difference in the spatial distribution of geotagged infections at the beginning and at

the end of the season (Table S3, p >0.10, Figure S2).

We compared the distribution of geotagged infection sites in each hybrid and found that

Hendschiken Hybrid_R3 was often different from the other plots (Table S4, p <0.10). In

Rudolfingen, there were more differences; Hybrid_R2 was different from both susceptible

hybrids, Hybrid_R3 was also different from Hybrid_S3, but was similar to Hybrid_S2, while

Hybrid_S2 and Hybrid_S3 had different distributions.

Genetic and genotypic diversity at different hierarchical levels

The six SSR loci used in our study were polymorphic, resulting in 2 to 9 alleles per locus.

Allelic diversity after clone correction ranged from 0.23 to 0.83 (Table S5) and there was

some variation in the number of alleles for each SSR between hybrids (Table S6). There were

some alleles unique to each field; for example, the 183 bp allele at SSRCb22 was found only

in Hendschiken and the 373 bp allele at SSRCb27 was only detected in Rudolfingen.

We investigated genetic diversity at different hierarchical levels, including within and

between hybrids and fields, and within leaves and hotspots. In Hendschiken, 158 isolates

were distributed among 82 unique MLGs. In Rudolfingen, 182 isolates resulted in 77 unique

MLGs (Table 5). The genetic diversity ranged from 0.40 to 0.57 across different hybrid plots,

and the genotypic diversity ranged from 0.73 to 0.95 (Table 5). There was little difference in

Nei's gene diversity between sites or hybrids. The clonal fraction differed significantly among

the hybrids in Rudolfingen (χ^2 = 11.15, df=3, p=0.01), while it was similar among hybrids at

Hendschiken (χ^2= 3.38, df=3, p=0.33). In Rudolfingen, the clonal fraction was lower in

Hybrid_S3 and higher in Hybrid_R2. The frequency of MLGS differed in each site (Figure 3).

In Rudolfingen, there was greater skew in MLG frequencies; R_MLG77 appeared in 46

isolates (20 in Hybrid_R2 and 26 in Hybrid_R3) and was restricted to the resistant hybrids

with *BvCR4*. In Hendschiken, differences between the hybrids were less pronounced, with

the most frequent MLG (H_MLG75) appearing 13 times in the resistant hybrids with *BvCR4*

(Hybrid_R2 and Hybrid_R3) (3 and 10 times respectively). The evenness was significantly

higher in susceptible hybrids in Rudolfingen, and no difference was found in Hendschiken

(Figure 4).

Genotypic diversity within a leaf and hotspot

To investigate genotypic diversity within individual leaves, multiple isolations were made

from the same leaf for a total of 48 leaves (Table 6a). It was common to find more than one

genotype per leaf, with 23 out of the 48 (0.48) tested leaves having more than one genotype

on the same leaf. The maximum number of MLGs from one leaf was two, irrespective of the

number of isolations (2-4). Although there was not enough data to make statistical

comparisons between hybrids or sites, we commonly observed the coexistence of multiple

genotypes in resistant hybrids. Specifically, in Hendschiken 6 out of 14 (0.39) tested leaves

from resistant hybrids had multiple strains and 10 out of 16 (0.63) resistant leaves had

multiple strains in Rudolfingen. We also considered the genotypic diversity within a hotspot

(n= 48) (Table 6b). Just over half of the hotspots had more than one MLG. In resistant

hybrids, 8 out of 13 (0.61) tested hotspots had multiple strains in Hendschiken and 11 out of

19 (0.57) hotspots had multiple strains in Rudolfingen.

Genotypic diversity at the start of each epidemic

Genotypic diversity and the spatial distribution of isolates from the first two sampling dates

in each plot were used to infer the most likely sources of primary inoculum in each

epidemic. In Hybrid_S2 (July 21st and 27th) in Rudolfingen we genotyped 13 isolates coming

from seven hotspots and identified nine MLGs (clonal fraction (CF) = 0.31). In Hendschiken

we genotyped 14 isolates from six hotspots and identified seven MLGs (CF = 0.50, but five

isolates from the same hotspot had the same MLG). At the beginning of the epidemic, no

MLGs were shared between hotspots, indicating that different pathogen strains established

each hotspot. Only four out of the seven initial hotspots in Rudolfingen were geotagged,

with pairwise distances ranging from 18.4 to 64.7 meters. Three out of the six initial

hotspots in Hendschiken were geotagged, with pairwise distances ranging from 25.0 to 68.7

meters. For the first two sampling dates in Hybrid_S3, only one isolate was genotyped from

each site, and thus the distribution of MLGs cannot be compared.

For the resistant Hybrid_R2: In Rudolfingen, on the first sampling date (August 25th) 15

isolates from 11 hotspots were genotyped. These contained six MLGs (CF = 0.6) with

pairwise distances ranging from 1.3 to 49.2 m. At the beginning of this epidemic the

R_MLG77 genotype was found 10 times in eight hotspots that were separated by pairwise

distances ranging from 1.3 to 19.6 m. In Hendschiken, during the first sampling dates in

Hybrid_R2 (August 11th and 20th), five isolates were sampled from four hotspots composed

of four MLGs (CF = 0.2), with the pairwise distances ranging from 4.2 to 61.4 m. No MLGs

were shared between hotspots.

For the resistant Hybrid_R3: In Rudolfingen, on the first two sampling dates (August 20th and

25th), four isolates from four hotspots were genotyped and each had a unique MLG (CF = 0),

with pairwise distances ranging from 5.7 to 46.0 m. The frequent MLG R_MLG77 appeared

in one of these hotspots. In Hendschiken, 15 isolates sampled from eight hotspots belonged

to nine MLGs (CF = 0.4) (four isolates from the same hotspot had the same MLG), with

pairwise distances between hotspots ranging from 6.2 to 58.3 m. One MLG was shared

between two hotspots that were 7.8 m apart.

Distribution of isolates with the same MLGs over time

The primary CLS infections in each plot gave rise to asexual conidia that provided secondary

inoculum. The spatial distribution of the resulting clones with identical MLGs provides

insights into the dispersal of secondary inoculum during a growing season. Common MLGs,

observed ≥ 5 times in each field (eight in Rudolfingen and five in Hendschiken), were used to

calculate the dispersal distance of the same genotype (Table 7). The minimum pairwise

distance of 0 m was not included in the table. It was common to find the same MLG

separated by distances exceeding 10 meters. The maximum distance between isolates

sharing the same MLG was 108.5 m in Rudolfingen and 68.7 m in Hendschiken.

Discussion

The major objective of this study was to compare CLS epidemics on sugar beet hybrids

exhibiting different levels of resistance. We combined detailed epidemiological surveys with

geo-referenced sampling and SSR genetic data to understand the fine scale temporal and

spatial distribution of *C. beticola* genotypes on different resistant hybrids grown under the

same environmental conditions during a single season. The dynamics of different pathogen populations were assessed on their respective hybrids during each epidemic. Resistant hybrids showed delayed epidemics, however, the host genotype did not affect the likelihood of spread. There were no differences in genetic diversity between pathogen populations on different hosts, however we found high frequencies of some MLGs on resistant hosts. The distribution of hotspots tended to be aggregated and was similar at the beginning and end of each epidemic in each hybrid, with a few exceptions. We found that a single genotype could disperse over a hundred meters during an epidemic, indicating the possibility of long-distance dispersal of individual clones during a growing season.

Onset of the disease varied according to host genotype

CLS emerged at least 3 weeks later in the hybrids with *BvCR4* (Hybrid_R2 and Hybrid_R3) compared to the hybrids without *BvCR4* (Hybrid_S3 and Hybrid_S4). The delayed appearance of CLS in resistant hybrids at both sites suggests that the resistant hybrids suppressed the establishment of CLS early in the season, agreeing with observations from earlier field trials conducted by KWS (personal communication). Similar delays (2 to 4 weeks) in the onset of CLS on resistant sugar beet hybrids were also reported in previous studies (Weiland and Koch, 2004; Weltmeier et al., 2011; Wolf and Verreet, 2005). It is possible that we did not always detect the start of an epidemic due to the size of the plots and our sampling frequency. For example, for Hybrid_R2 in Rudolfingen and Hybrid_S2 in Hendschiken, we collected leaves from 43 and 22 plants respectively on the first day of sampling (Table 2), suggesting that the epidemic had already started in these plots in the days between our sampling visits. For the six remaining hybrid plots, the number of plants collected on the first sampling date was < 10, suggesting that we caught the beginning of those epidemics. If we consider only the six plots with less than 10 infected plants at the onset of the epidemic, the average distance between infected plants was 23.0 m in Hendschiken and 30.8 m in Rudolfingen. If we add the other two plots, the average distance between infected plants was 23.9 m in Hendschiken and 25.7 m in Rudolfingen. Our interpretation in both cases is that the primary inoculum was airborne. Given the high genetic diversity found on these first infected plants and the random distributions of primary infections across the fields in both locations (Table S2), we hypothesize that the primary inoculum was mainly airborne ascospores.

Localized spread was not strongly affected by host genotype

There was no effect of host genotype on the probability that a single infected plant would develop into a hotspot, with about 80% of infection cases spreading to neighboring plants (Table 3), but the speed of localized spread (involving 2-5 plants) varied between hybrids and fields. In Rudolfingen, there was no difference among hybrids in how quickly CLS spread to neighboring plants, but in Hendschiken, the spread was slower in Hybrid_R3 and faster in Hybrid_R2 compared to the susceptible Hybrid_S2. This field-dependent pattern may be attributed to differences in the distribution of the hybrid plots, in particular the susceptibility of the hybrid planted in a neighboring plot. Previous studies suggested that the level of host resistance did not influence the latent period, however, susceptible hybrids produced more conidia per unit of necrotic leaf area (Oerke et al., 2019). Thus, plots planted next to susceptible hybrids may experience higher local disease pressure. In Hendschiken, Hybrid_R2 was planted next to the susceptible Hybrid_S3, while in Rudolfingen, Hybrid_R2 was planted next to the resistant Hybrid_R3, which might have acted as a buffer zone that reduced the influx of conidia onto Hybrid_R2 (Figure S1). In addition to differences in local spore production, differences in local microclimates may also have affected epidemic development at each field site (Imbusch et al., 2021; Rossi et al., 2000; Vereijssen et al., 2006). It also is possible that our weekly sampling interval was not frequent enough to accurately monitor localized spread, as some hotspots spread from one to five plants between two successive time points.

To summarize, our findings suggest that the host genotype strongly influenced the emergence of CLS but had little impact on the probability that CLS would spread and form hotspots. This is in line with previous work that demonstrated that the latency of spore production was not influenced by host resistance (Rossi et al., 2000). If *C. beticola* strains succeeded in overcoming the resistance of a host and became established on a resistant plant, their ability to spread to neighboring plants did not differ from the speed of dispersal found on susceptible plants.

Infection sites were not randomly distributed in fields

At the end of the season, CLS infection sites were mostly aggregated (not randomly

distributed) with one exception (Table S2). The distributions of infections changed little

between the onset and the end of the epidemic, except for Hybrid_S2 and Hybrid_R2 in

Rudolfingen (Table S3). The minimum distance between infection cases was approximately

one meter, while the maximum distance reached 116.3 meters (Table 4). In line with earlier

findings, we observed higher levels of CLS in Hybrid_S2, Hybrid_S3 and Hybrid_R2 compared

to Hybrid_R3 at the end of the season in both sites. Lesions coalesced and caused wilting

and death of entire leaves, leading to new growth from the roots in hybrids S2, S3 and R2.

Hybrid_R3 did not exhibit obvious lesion coalescence, leaf death, or regrowth despite having

many plants with CLS. This may reflect the shorter time interval between disease onset and

final sampling in Hybrid_R3. Studies have shown that later infection results in lower yield

losses for resistant varieties (Kaiser et al., 2010), and that the delayed CLS onset observed in

resistant hybrids was positively correlated with sugar yield (Wolf and Verreet, 2009).

Genotypic variation and spatial distribution of genotypes

Earlier studies illustrated how pathogen populations with higher genetic diversity can have a

higher evolutionary potential to overcome R genes (McDonald and Linde, 2002b; Stam and

McDonald, 2018). In our study, the average Nei's gene diversity based on six SSR markers

ranged from 0.40 to 0.57 (Table 5). It is important to note that our estimates of genotypic

diversity were limited because only six SSR loci were used to identify putative clones. To

estimate how accurately the six SSRs identified clones, we compared the clones identified

using the SSR results with clones identified using whole genome sequences for 113 isolates

where we obtained both types of data (Chen et al., 2023). The six SSR loci correctly

identified clones or unique genotypes in 84% of cases (47 out of 58 in Rudolfingen and 48

out of 55 in Hendschiken). The incorrect assignments were a mix of wrongly identifying two

isolates as sharing the same genotype (underestimating genotypic diversity) and failing to

identify two isolates as being clones (overestimating genotypic diversity) at roughly equal

proportions (9.7% and 6.2% respectively). While this illustrates the potential errors of our

SSR genotypes, we expect that this error rate will be randomly distributed across hybrids, so

comparisons of relative genotypic diversity between hybrids remain valid.

The high genotypic diversity found in each population supports the hypothesis that *C.*

beticola populations undergo regular cycles of sexual recombination, as proposed in earlier

studies (Birla et al., 2012; Groenewald et al., 2007b, 2006). We detected high genotypic

diversity at small spatial scales, with 23 out of 48 tested leaves infected by more than one

MLG and 27 out of 48 tested hotspots containing more than one MLG. Other studies also

observed multiple SSR MLGs on a single leaf in table beet and Swiss chard (Vaghefi et al.,

2017c) and using other genetic markers (TAPD and DAMD) on sugar beet (Moretti et al.,

2006). The presence of multiple MLGs on a single leaf suggests that co-infection is common

and that multiple strains may contribute to the spread of disease within a CLS hotspot.

We hypothesized that virulent isolates would be more prevalent on *BvCR4* hybrids due to

their ability to overcome this resistance gene. If only a small fraction of isolates in the

primary inoculum are virulent and able to produce asexual secondary inoculum on the

resistant hybrids, we would expect these virulent clones to increase in frequency on

resistant hybrids during the epidemic, resulting in lower genotypic diversity and higher

clonal fractions on resistant hybrids compared to susceptible hybrids. Consistent with this

expectation, we observed a skewed frequency distribution of MLGs between hybrids with

and without *BvCR4* at both sites. In Rudolfingen, the clonal fraction was higher in the

resistant hybrids and evenness was strongly skewed (Figure 3): one MLG (R_MLG77)

appeared in 46 isolates exclusively in resistant Hybrid_R2 and Hybrid_R3. In Hendschiken,

the differences in frequency were less striking, but one MLG (H_MLG75) appeared in 13

isolates solely found on the *BvCR4* hybrids. A subset of the isolates collected from resistant

hybrids were tested for virulence against hybrids with and without *BvCR4* under field

conditions in a later study. Under experimentally elevated disease pressure that study

confirmed that the highest frequency MLGs were virulent strains that could overcome

BvCR4 (Chen et al., 2023). It also found that only half of the isolates collected from hybrids

with *BvCR4* were virulent against this gene. The lack of evidence for evolutionary

bottlenecks in our study may be attributed to the frequent coexistence of virulent and

avirulent strains within the same hybrid. Most of our temporal sampling was during the

epidemic growth, thus we did not capture the end-of-season dynamics which might show

stronger bottlenecks following clone competition. Transect or random sampling of the

pathogen populations at the end of the season may provide a better test of the

effectiveness of resistant hybrids as selection sieves.

Source of inoculum and spatial distribution of MLGs

One of our aims in this study was to identify the potential sources of primary inoculum for

CLS epidemics. We hypothesized that a CLS epidemic can be initiated by wind-dispersed

ascospores which are capable of long-distance dispersal over hundreds of meters or by

conidia on infected leaves from earlier crops that are moved by wind or watersplash. If the

primary inoculum in a plot originates from windborne ascospores coming from outside the

field, we would expect to find genetically different isolates sparsely and randomly

distributed across individual plants at the onset of an epidemic. Alternatively, if the primary

inoculum arises from locally produced asexual spores transmitted through watersplash from

crop debris or human activity (e.g. infested seeds or machinery), we would expect to find

the same genotype occurring simultaneously in different hotspots that might be aggregated

in the field (for the case of infested machinery). In plots planted with Hybrid_S2, at both

sites we observed unique genotypes in each hotspot, suggesting that unique genotypes led

to the establishment of each hotspot and that epidemics were initiated by genetically

diverse primary inoculum coming from outside the fields. Similar patterns were observed in

Hybrid_R3 at both sites and in Hybrid_R2 in Hendschiken, with no or few isolates shared

between hotspots. But a different pattern was found for the resistant Hybrid_R2 in

Rudolfingen. In this epidemic, the genotypic diversity of the first infections was lower, and

the dominant genotype R_MLG77 appeared simultaneously 10 times in 8 hotspots

distributed widely across the field. This suggests that asexual spores originating from a

neighboring unsampled border planted to Hybrid_R2 may have disproportionally affected

the genotypic diversity found in the sampled Hybrid_R2 plot. We postulate that the

R_MLG77 strain emerged from standing genetic variation and attained a relatively high

frequency in the border plots before being transmitted into the sampled Hybrid_R2 plot

(Figure S1). With the exception of Hybrid_R2 in Rudolfingen, we found that genetically

distinct primary inoculum was distributed widely across most plots (Table S2), as expected

for wind-dispersed ascospores originating from outside of the field.

Our experiment also aimed to better understand the dispersal pattern of secondary

inoculum within a season. In Rudolfingen, high frequency MLGs (≥5) were separated by

between 49 and 108 meters and in Hendschiken by between 9 and 68 m (Table 7). This is

the first study in the CLS pathosystem to measure the dispersal of individual genotypes

within a season. The dispersal distances we observe are unlikely to be due solely to water-

splash, as water-splash can only disperse *C. beticola* spores short distances (several meters)

(Carlson, 1967). Wind dispersal is a more likely mechanism, and although wind was already

recognized as an important mechanism, it was not clear before how far wind could disperse

C. beticola (Khan et al., 2008). Other possible dispersal mechanisms cannot be ruled out. It is

possible that insects could also disperse the spores, along with human activity, including

machine operations or spores moving on contaminated clothing during our weekly sampling

visits.

References

Alberto, F., 2009. MsatAllele_1.0: An R Package to Visualize the Binning of Microsatellite
Alleles. Journal of Heredity 100, 394–397. https://doi.org/10.1093/jhered/esn110

Archie, E.A., Luikart, G., Ezenwa, V.O., 2009. Infecting epidemiology with genetics: a new
frontier in disease ecology. Trends in Ecology & Evolution 24, 21–30.
https://doi.org/10.1016/j.tree.2008.08.008

Barrett, L.G., Thrall, P.H., Burdon, J.J., Linde, C.C., 2008. Life history determines genetic
structure and evolutionary potential of host–parasite interactions. Trends in Ecology
& Evolution 23, 678–685. https://doi.org/10.1016/j.tree.2008.06.017

Birla, K., Rivera-Varas, V., Secor, G.A., Khan, M.F.R., Bolton, M.D., 2012. Characterization of
cytochrome b from European field isolates of *Cercospora beticola* with quinone
outside inhibitor resistance. Eur J Plant Pathol 134, 475–488.
https://doi.org/10.1007/s10658-012-0029-y

Bolton, M.D., Rivera, V., Secor, G., 2013. Identification of the G143A mutation associated
with QoI resistance in *Cercospora beticola* field isolates from Michigan, United
States: G143A mutation associated with QoI resistance in *C. beticola*. Pest. Manag.
Sci. 69, 35–39. https://doi.org/10.1002/ps.3358

Bolton, M.D., Secor, G.A., Rivera, V., Weiland, J.J., Rudolph, K., Birla, K., Rengifo, J.,
Campbell, L.G., 2012. Evaluation of the potential for sexual reproduction in field
populations of *Cercospora beticola* from USA. Fungal Biology 116, 511–521.
https://doi.org/10.1016/j.funbio.2012.01.011

Brown, J.K.M., 2015. Durable resistance of crops to disease: a darwinian perspective. Annu.
Rev. Phytopathol. 53, 513–539. https://doi.org/10.1146/annurev-phyto-102313-
045914

Carlson, L.W., 1967. Relation of weather factors to dispersal of conidia of *Cercospora
beticola* (sacc.). JSBR 14, 319–323. https://doi.org/10.5274/jsbr.14.4.319

Chen, C., Keunecke, H., Bemm, F., Gyetvai, G., Neu, E., Kopisch-Obuch, F.J., McDonald, B.A.,
Stapley, J., 2023. Genome-wide association studies reveal a rapidly evolving
candidate avirulence effector in the Cercospora leaf spot pathogen *Cercospora
beticola*. https://doi.org/10.1101/2023.06.11.544483

Covarrubias-Pazaran, G., Diaz-Garcia, L., Schlautman, B., Salazar, W., Zalapa, J., 2016.
Fragman: an R package for fragment analysis. BMC Genet 17, 62.
https://doi.org/10.1186/s12863-016-0365-6

Cowger, C., Brown, J.K.M., 2019. Durability of quantitative resistance in crops: greater than
we know? Annu. Rev. Phytopathol. 57, 253–277. https://doi.org/10.1146/annurev-
phyto-082718-100016

Del Ponte, E., 2023. r4pde: Companion to R4PDE book.

Desbiez, C., Wipf-Scheibel, C., Lecoq, H., 2002. Biological and serological variability,
622   evolution and molecular epidemiology of Zucchini yellow mosaic virus (ZYMV,
623   Potyvirus) with special reference to Caribbean islands. Virus Research 85, 5–16.
624   https://doi.org/10.1016/S0168-1702(02)00013-8

625   Francis, S.A., Luterbacher, M.C., 2003. Identification and exploitation of novel disease
626   resistance genes in sugar beet. Pest Management Science 59, 225–230.
627   https://doi.org/10.1002/ps.569

628   Groenewald, M., Groenewald, J.Z., Harrington, T.C., Abeln, E.C.A., Crous, P.W., 2006. Mating
629   type gene analysis in apparently asexual Cercospora species is suggestive of cryptic
630   sex. Fungal Genetics and Biology 43, 813–825.
631   https://doi.org/10.1016/j.fgb.2006.05.008

632   Groenewald, M., Groenewald, J.Z., Linde, C.C., Crous, P.W., 2007a. Development of
633   polymorphic microsatellite and single nucleotide polymorphism markers for
634   *Cercospora beticola* (Mycosphaerellaceae). Molecular Ecology Notes 7, 890–892.
635   https://doi.org/10.1111/j.1471-8286.2007.01739.x

636   Groenewald, M., Linde, C.C., Groenewald, J.Z., Crous, P.W., 2007b. Indirect evidence for
637   sexual reproduction in *Cercospora beticola* populations from sugar beet. Plant
638   Pathology 57, 25–32. https://doi.org/10.1111/j.1365-3059.2007.01697.x

639   Hijmans, R. J., 2022. Geosphere: spherical trigonometry.

640   Holtschulte, B., 2000. Cercospora beticola worldwide distribution and incidence.

641   Imbusch, F., Liebe, S., Erven, T., Varrelmann, M., 2021. Dynamics of Cercospora leaf spot
642   disease determined by aerial spore dispersal in artificially inoculated sugar beet
643   fields. Plant Pathol. 70, 853–861. https://doi.org/10.1111/ppa.13337

644   Kaiser, U., Kluth, C., Märländer, B., 2010. Variety-specific epidemiology of *Cercospora*
645   *beticola* sacc. and consequences for threshold-based timing of fungicide application
646   in sugar beet. Journal of Phytopathology 158, 296–306.
647   https://doi.org/10.1111/j.1439-0434.2009.01618.x

648   Kamvar, Z.N., Brooks, J.C., Grünwald, N.J., 2015. Novel R tools for analysis of genome-wide
649   population genetic data with emphasis on clonality. Front. Genet. 6.
650   https://doi.org/10.3389/fgene.2015.00208

651   Kamvar, Z.N., Tabima, J.F., Grünwald, N.J., 2014. *Poppr*: an R package for genetic analysis of
652   populations with clonal, partially clonal, and/or sexual reproduction. PeerJ 2, e281.
653   https://doi.org/10.7717/peerj.281

654   Khan, J., Rio, L.E. del, Nelson, R., Rivera-Varas, V., Secor, G.A., Khan, M.F.R., 2008. Survival,
655   dispersal, and primary infection site for *Cercospora beticola* in sugar beet. Plant
656   Disease 92, 741–745. https://doi.org/10.1094/PDIS-92-5-0741

Knight, N.L., Pethybridge, S.J., 2020. An improved PCR assay for species-specific detection
and quantification of *Cercospora beticola*. Canadian Journal of Plant Pathology 42,
72–83. https://doi.org/10.1080/07060661.2019.1621380

McDonald, B.A., Linde, C., 2002a. The population genetics of plant pathogens and breeding
strategies for durable resistance. Euphytica 124, 163–180.
https://doi.org/10.1023/A:1015678432355

McDonald, B.A., Linde, C., 2002b. Pathogen population genetics, evolutionary potential, and
durable resistance. Annu. Rev. Phytopathol. 40, 349–379.
https://doi.org/10.1146/annurev.phyto.40.120501.101443

Milgroom, M.G., Peever, T.L., 2003. Population biology of plant pathogens: the synthesis of
plant disease epidemiology and population genetics. Plant Disease 87, 608–617.
https://doi.org/10.1094/PDIS.2003.87.6.608

Moretti, M., Karaoglanidis, G., Saracchi, M., Fontana, A., Farina, G., 2006. Analysis of
genotypic diversity in *Cercospora beticola* Sacc. field isolates. Ann. Microbiol. 56,
215–221. https://doi.org/10.1007/BF03175008

Oerke, E.-C., Leucker, M., Steiner, U., 2019. Sensory assessment of *Cercospora beticola*
sporulation for phenotyping the partial disease resistance of sugar beet genotypes.
Plant Methods 15, 133. https://doi.org/10.1186/s13007-019-0521-x

Pethybridge, S.J., Sharma, S., Hansen, Z.R., Vaghefi, N., Hanson, L.E., Kikkert, J.R., 2020.
Improving fungicide-based management of Cercospora leaf spot in table beet in New
York, USA. Canadian Journal of Plant Pathology 42, 353–366.
https://doi.org/10.1080/07060661.2019.1690048

Posit team, 2023. RStudio: Integrated Development Environment for R.

R Core Team, 2023. R: A Language and Environment for Statistical Computing.

Rangel, L.I., Spanner, R.E., Ebert, M.K., Pethybridge, S.J., Stukenbrock, E.H., De Jonge, R.,
Secor, G.A., Bolton, M.D., 2020. *Cercospora beticola*: The intoxicating lifestyle of the
leaf spot pathogen of sugar beet. Molecular Plant Pathology 21, 1020–1041.
https://doi.org/10.1111/mpp.12962

Rossi, V., Battilani, P., Chiusa, G., Giosuè, S., Languasco, L., Racca, P., 2000. Components of
rate-reducing resistance to Cercospora leaf spot in sugar beet: conidiation length,
spore yield. Journal of Plant Pathology.

Shane, W.W., Teng, P.S., 1992. Impact of cercospora leaf spot on root weight, sugar yield,
and purity of *beta vulgaris*. Plant Dis. 76, 812. https://doi.org/10.1094/PD-76-0812

Spanner, R., Neubauer, J., Heick, T.M., Grusak, M.A., Hamilton, O., Rivera-Varas, V., De
Jonge, R., Pethybridge, S., Webb, K.M., Leubner-Metzger, G., Secor, G.A., Bolton,
M.D., 2022. Seedborne *Cercospora beticola* Can Initiate Cercospora Leaf Spot from

Sugar Beet (*Beta vulgaris*) Fruit Tissue. Phytopathology® 112, 1016–1028.
https://doi.org/10.1094/PHYTO-03-21-0113-R

Stam, R., McDonald, B.A., 2018. When resistance gene pyramids are not durable-the role of
pathogen diversity: R-gene pyramid durability and pathogen diversity. Molecular
Plant Pathology 19, 521–524. https://doi.org/10.1111/mpp.12636

Tedford, S.L., Burlakoti, R.R., Schaafsma, A.W., Trueman, C.L., 2018. Relationships among
airborne *Cercospora beticola* conidia concentration, weather variables and
cercospora leaf spot severity in sugar beet (*beta vulgaris* l.). Canadian Journal of
Plant Pathology 40, 1–10. https://doi.org/10.1080/07060661.2017.1410726

Trkulja, N.R., Milosavljević, A.G., Mitrović, M.S., Jović, J.B., Toševski, I.T., Khan, M.F.R., Secor,
G.A., 2017. Molecular and experimental evidence of multi-resistance of *Cercospora*
beticola field populations to MBC, DMI and QoI fungicides. Eur J Plant Pathol 149,
895–910. https://doi.org/10.1007/s10658-017-1239-0

Vaghefi, N., Kikkert, J.R., Bolton, M.D., Hanson, L.E., Secor, G.A., Nelson, S.C., Pethybridge,
S.J., 2017a. Global genotype flow in *Cercospora beticola* populations confirmed
through genotyping-by-sequencing. PLoS ONE 12, e0186488.
https://doi.org/10.1371/journal.pone.0186488

Vaghefi, N., Kikkert, J.R., Bolton, M.D., Hanson, L.E., Secor, G.A., Pethybridge, S.J., 2017b. De
novo genome assembly of *Cercospora beticola* for microsatellite marker
development and validation. Fungal Ecology 26, 125–134.
https://doi.org/10.1016/j.funeco.2017.01.006

Vaghefi, N., Nelson, S.C., Kikkert, J.R., Pethybridge, S.J., 2017c. Genetic structure of
Cercospora beticola populations on beta vulgaris in New York and Hawaii. Sci Rep 7,
1726. https://doi.org/10.1038/s41598-017-01929-4

Vereijssen, J., Schneider, J.H.M., Jeger, M.J., 2007. Epidemiology of Cercospora leaf spot on
sugar beet: modeling disease dynamics within and between individual plants.
Phytopathology® 97, 1550–1557. https://doi.org/10.1094/PHYTO-97-12-1550

Vereijssen, J., Schneider, J.H.M., Stein, A., Jeger, M.J., 2006. Spatial pattern of Cercospora
leaf spot of sugar beet in fields in long- and recently-established areas. Eur J Plant
Pathol 116, 187–198. https://doi.org/10.1007/s10658-006-9046-z

Vogel, J., Kenter, C., Holst, C., Märländer, B., 2018. New generation of resistant sugar beet
varieties for advanced integrated management of Cercospora leaf spot in central
Europe. Front. Plant Sci. 9, 222. https://doi.org/10.3389/fpls.2018.00222

Weiland, J., Koch, G., 2004. Sugarbeet leaf spot disease (*Cercospora beticola* Sacc.)+. Mol
Plant Pathol 5, 157–166. https://doi.org/10.1111/j.1364-3703.2004.00218.x

Weltmeier, F., Mäser, A., Menze, A., Hennig, S., Schad, M., Breuer, F., Schulz, B., Holtschulte,
B., Nehls, R., Stahl, D.J., 2011. Transcript profiles in sugar beet genotypes uncover
timing and strength of defense reactions to *Cercospora beticola* infection.

Wolf, P.F.J., Verreet, J.-A., 2009. Empirical-deterministic prediction of disease and losses
caused by Cercospora leaf spots in sugar beets. JOURNAL FÜR KULTURFLANZEN 61
(5). S., 168–177.

Wolf, P.F.J., Verreet, J.A., 2005. Factors affecting the onset of Cercospora leaf spot
epidemics in sugar beet and establishment of disease-monitoring thresholds.
Phytopathology® 95, 269–274. https://doi.org/10.1094/PHYTO-95-0269

Xhaard, C., Barrès, B., Andrieux, A., Bousset, L., Halkett, F., Frey, P., 2012. Disentangling the
genetic origins of a plant pathogen during disease spread using an original molecular
epidemiology approach: genetic monitoring of a poplar rust epidemic. Molecular
Ecology 21, 2383–2398. https://doi.org/10.1111/j.1365-294X.2012.05556.x

Table 1. Sugar beet hybrids and their resistance to cercospora leaf spot.

Hybrids	Cercospora leaf spot resistance level
Hybrid_S2	Low quantitative resistance
Hybrid_S3	Moderate quantitative resistance
Hybrid_R2	*BvCR4* with low quantitative background resistance
Hybrid_R3	*BvCR4* with moderate quantitative background resistance

Table 2. The first appearance of CLS, the number of plants sampled at the first sampling date (No. plants), the number of days after the first appearance of symptoms on July 21st (onset delay), the total length of the sampling period, and the last sampling date (the right censored date is shown in parenthesis, see Methods) for each hybrid at each field site.

Location	Hybrid	First infection date	Onset delay	No. plants	Sampling period	Last sampling date (right censored date)
Hendschiken	Hybrid_S2	21-07-20	0	22	30	20-08-20 (11-08-20)
	Hybrid_S3	27-07-20	6	1	24	20-08-20 (11-08-20)
	Hybrid_R2	11-08-20	21	6	33	08-09-20 (03-09-20)
	Hybrid_R3	20-08-20	30	3	35	24-09-20 (03-09-20)
Rudolfingen	Hybrid_S2	21-07-20	0	7	30	20-08-20 (11-08-20)
	Hybrid_S3	27-07-20	6	2	24	20-08-20 (11-08-20)
	Hybrid_R2	25-08-20	35	43	14	08-09-20 (03-09-20)
	Hybrid_R3	20-08-20	30	4	35	24-09-20 (08-09-20)

Table 3. The number of infection cases (IC) and the proportion of these that developed into hotspots (≥2 infected plants) and the proportion that developed into large hotspots (≥5 infected plants) for each hybrid. The data has been right censored (see Methods).

Location	Hybrid	No. IC	Number of hotspots (proportion)	Number of large hotspots (proportion)
Hendschiken	Hybrid _S2	37	26 (0.70)	5 (0.14)
	Hybrid _S3	20	17 (0.85)	4 (0.20)
	Hybrid _R2	22	18 (0.82)	7 (0.32)
	Hybrid _R3	30	24 (0.80)	4 (0.13)
Rudolfingen	Hybrid _S2	23	19 (0.83)	9 (0.39)
	Hybrid _S3	18	17 (0.94)	6 (0.33)
	Hybrid _R2	29	26 (0.90)	9 (0.31)
	Hybrid _R3	31	23 (0.74)	5 (0.16)

Table 4. The number of geotagged infection sites and the minimum, mean and maximum pairwise distance (m) between infection sites at the beginning and the end of the sampling period in each hybrid.

Location	Hybrid	Pairwise distance at the start of epidemics				Pairwise distance at the end of epidemics			
		Count	Min	Mean	Max	Count	Min	Mean	Max
Rudolfingen	Hybrid _S2	14	2.2	28.2	73.7	41	1.3	36.7	110.7
	Hybrid _S3	4	3.8	54.7	98.6	31	1.3	47.8	116.3
	Hybrid _R2	15	1.3	19.7	62.0	37	1.3	29.2	108.3
	Hybrid _R3	8	4.2	33.9	94.5	88	0.2	37.2	110.0
Hendschiken	Hybrid _S2	14	1.3	24.7	72.8	37	0.8	26.0	72.8
	Hybrid _S3	8	1.3	20.0	51.1	24	1.3	23.2	64.6
	Hybrid _R2	6	4.2	29.7	57.3	41	1.3	27.2	76.0
	Hybrid _R3	9	3.1	22.5	58.3	66	0.2	21.6	68.5

Table 5. Indices of gene and genotype diversity based on six SSR loci for *Cercospora beticola* populations collected from four sugar beet hybrids in two fields in Switzerland. The table includes the number of strains (N), the number of multilocus genotypes (MLGs), the expected number of MLGs after rarefaction (eMLGs), the clonal fraction (CF=1-(MLGs/N), the Simpson's complement index (λ) and Nei's index of gene diversity before clone correction (He_0) and after clone correction (He_{cc}).

Location	Hybrid	N	MLGs	eMLGs	CF	λ	He_0	He_{cc}
Hendschiken	Hybrid_S2	30	20	15.6	0.33	0.91	0.56	0.60
	Hybrid _S3	22	16	16	0.27	0.92	0.51	0.54
	Hybrid _R2	43	28	17.3	0.35	0.95	0.52	0.54
	Hybrid _R3	63	37	16.2	0.41	0.94	0.56	0.58
	Total/Mean	**158**	**82**	**18.3**	**0.48**	**0.97**	**0.56**	**0.56**
Rudolfingen	Hybrid _S2	49	27	16.7	0.45	0.95	0.51	0.54
	Hybrid _S3	23	19	19	0.17	0.94	0.57	0.60
	Hybrid _R2	40	16	10.7	0.60	0.73	0.40	0.56
	Hybrid _R3	70	35	14.1	0.50	0.85	0.52	0.57
	Total/Mean	**182**	**77**	**16.3**	**0.58**	**0.92**	**0.55**	**0.57**

Table 6. Genotypic diversity within (a) a leaf and (b) a hotspot (HS). The first column indicates a frequency class relating to the number of isolations per leaf/hotspot. There are three classes: 2, 3 or 4 isolations were made per leaf leaf/hotspot. The table includes the number of leaves/hotspots that had only one MLG (all clones) and with more than one MLG. The proportions are provided in parenthesis.

(a)

Number of isolations per leaf	Number of leaves	Single MLGs Number (proportion)	Multiple MLGs Number (proportion)
2	41	21 (0.51)	20 (0.49)
3	6	3 (0.50)	3 (0.50)
4	1	1 (1.00)	0
Total	48	25 (0.52)	23 (0.48)

(b)

Number of isolations per HS	Number of HS	Single MLGs Number (proportion)	Multiple MLGs Number (proportion)
2	34	17 (0.50)	17 (0.50)
3	11	3 (0.27)	8 (0.73)
4	3	1 (0.33)	2 (0.67)
Total	48	21 (0.43)	27 (0.57)

Table 7. The median, mean and the maximum pairwise distance (m) between the most frequent MLGs that were found at least five times in a field site.

MLG	Count	Median	Mean	Maximum
R_MLG77	46	16.6	22.8	108.5
R_MLG56	7	43.6	38.9	100.5
R_MLG13	6	7.5	16.3	49.8
R_MLG42	6	58.4	48.0	107.8
R_MLG15	5	51.9	43.5	105.0
R_MLG20	5	14.9	28.5	74.0
R_MLG46	5	31.9	23.2	64.4
R_MLG55	5	19.2	30.0	77.1
H_MLG75	13	4.2	7.8	20.5
H_MLG51	11	15.7	18.9	46.7
H_MLG9	9	3.7	4.2	9.4
H_MLG77	8	15.0	13.5	36.7
H_MLG82	8	29.3	27.1	68.7

Figure 1. Spatial distribution of CLS in Rudolfingen (a) and Hendschiken (b). Each point represents a geotagged infection case. The size of the point indicates the number of sampled plants within the hotspot. The point color indicates the plant hybrid: green: Hybrid_S2; pink: Hybrid_S3; brown: Hybrid_R2; blue: Hybrid_R3. Plant hybrid was determined by counting the planted rows and using the different hybrid morphology. The dashed lines approximate the separation of the different hybrids, but in the actual field these lines were not perfectly straight and there was no physical separation. Thus, it can appear that one plot contains more than one hybrid (points with different colors), but this is an artifact of plotting and scaling the figure.

Figure 2. Accumulation of geotagged infection cases over time in Rudolfingen (a) and Hendschiken (b) in each hybrid. Sampling terminated at different dates in each hybrid due to rapid disease progression that resulted in complete infection of the entire experimental plot (see methods and Table 2). Sampling termination prior to end of field trial indicated by red circles (○).

Figure 3. Frequencies of multilocus genotypes (MLGs) appearing in different hybrids in Rudolfingen (a) and Hendschiken (b). Each bar refers to one MLG, the height of the bar indicates the frequency of the MLG in each hybrid.

Figure 4. Evenness of MLGs in each hybrid in Rudolfingen (a) and Hendschiken (b). Points are the values of evenness and the error bars indicate the bootstrapped confidence intervals.

| ● Hybrid_R2 | ● Hybrid_S2 |
| ● Hybrid_R3 | ● Hybrid_S3 |

Figure 1. Spatial distribution of CLS in Rudolfingen (a) and Hendschiken (b). Each point represents a geotagged infection case. The size of the point indicates the number of sampled plants within the hotspot. The point color indicates the plant hybrid: green: Hybrid_S2; pink: Hybrid_S3; brown: Hybrid_R2; blue: Hybrid_R3. Plant hybrid was determined by counting the planted rows and using the different hybrid morphology. The dashed lines approximate the separation of the different hybrids, but in the actual field these lines were not perfectly straight and there was no physical separation. Thus, it can appear that one plot contains more than one hybrid (points with different colors), but this is an artifact of plotting and scaling the figure.

Figure 2. Accumulation of geotagged infection cases over time in Rudolfingen (a) and Hendschiken (b) in each hybrid. Sampling terminated at different dates in each hybrid due to rapid disease progression that resulted in complete infection of the entire experimental plot (see methods and Table 2). Sampling termination prior to end of field trial indicated by red circles (○).

Figure 3. Frequencies of multilocus genotypes (MLGs) appearing in different hybrids in Rudolfingen (a) and Hendschiken (b). Each bar refers to one MLG, the height of the bar indicates the frequency of the MLG in each hybrid.

Figure 4. Evenness of MLGs in each hybrid in Rudolfingen (a) and Hendschiken (b). Points are the values of evenness and the error bars indicate the bootstrapped confidence intervals.