The mitochondrial mRNA stabilizing protein, SLIRP, regulates skeletal muscle mitochondrial

structure and respiration by exercise-recoverable mechanisms

Tang Cam Phung Pham[1,2], Steffen Henning Raun[2], Essi Havula[3], Carlos Henriquez-Olguín[1], Diana

Rubalcava-Gracia[4], Emma Frank[2], Andreas Mæchel Fritzen[1,2], Paulo R. Jannig[5], Nicoline Resen

Andersen[1], Rikke Kruse[6], Mona Sadek Ali[2], Jens Frey Halling[7], Stine Ringholm[7], Solvejg Hansen[1],

Anders Krogh Lemminger[1], Peter Schjerling[8,9], Maria Houborg Petersen[6], Martin Eisemann de

Almeida[6,10], Thomas Elbenhardt Jensen[1], Bente Kiens[1], Morten Hostrup[1], Steen Larsen[2,8,9,11], Niels

Ørtenblad[10], Kurt Højlund[6,12], Michael Kjær[8,9], Jorge L. Ruas[5], Aleksandra Trifunovic[13], Jørgen Frank

Pind Wojtaszewski[1], Joachim Nielsen[10], Klaus Qvortrup[2], Henriette Pilegaard[7], Erik Arne Richter[1],

Lykke Sylow[1,2]

[1] Department of Nutrition, Exercise and Sports, Faculty of Science, University of Copenhagen,
Copenhagen, Denmark

[2] Department of Biomedical Sciences, Faculty of Health and Medical Sciences, University of
Copenhagen, Copenhagen, Denmark

[3] Stem Cells and Metabolism Research Program, Faculty of Medicine, University of Helsinki, Finland

[4] Division of Molecular Metabolism, Department of Medical Biochemistry and Biophysics,
Karolinska Institutet, Stockholm, Sweden

[5] Molecular and Cellular Exercise Physiology, Department of Physiology and Pharmacology,
Karolinska Institutet, SE-17177 Stockholm, Sweden

[6] Steno Diabetes Center Odense, Odense University Hospital, Odense, Denmark

[7] Department of Biology, University of Copenhagen, Copenhagen, Denmark

[8] Institute of Sports Medicine Copenhagen, Department of Orthopaedic Surgery M, Bispebjerg
Hospital, Copenhagen, Denmark
[9] Center for Healthy Aging, Faculty of Health and Medical Sciences, University of Copenhagen,
Copenhagen, Denmark
[10] Department of Sports Science and Clinical Biomechanics, University of Southern Denmark,
Odense, Denmark
[11] Clinical Research Centre, Medical University of Bialystok, Bialystok, Poland
[12] Department of Clinical Research, University of Southern Denmark, Odense, Denmark

[13] Institute for Mitochondrial Diseases and Aging, Cologne Excellence Cluster on Cellular Stress

Responses in Aging-Associated Diseases (CECAD) and Center for Molecular Medicine (CMMC),

Medical Faculty, University of Cologne, D-50931 Cologne, Germany

Running title: Exercise reverses SLIRP-induced mitochondrial defects

Keywords: SRA stem-loop interacting RNA-binding protein, mitochondrial dysfunction, skeletal

muscle, exercise training, Drosophila

#Corresponding author: Lykke Sylow

38   Lykkesylow@sund.ku.dk

39   **Competing Interests.** The authors declare that there is no conflict of interest.

Summary and graphical abstract

The decline in mitochondrial function is associated with decreased muscle mass and strength in multiple conditions, including sarcopenia, type 2 diabetes, and cancer. Treatment of these diseases could include improving mitochondrial function, however, there are limited and equivocal data regarding the molecular cues that control muscle mitochondrial plasticity. Here we uncover the mitochondrial-mRNA-stabilizing protein SLIRP, in complex with LRPPRC, as a PGC-1α target that regulates mitochondrial structure, respiration, and mitochondrially-encoded-mRNA pools in skeletal muscle. Exercise training effectively counteracted mitochondrial defects induced by loss of LRPPRC/SLIRP, despite sustained low mitochondrially-encoded-mRNA pools, via a mechanism likely including increased mitoribosome translation capacity. In human skeletal muscle, exercise training robustly increased SLIRP and LRPPRC protein content across exercise modalities and sexes, yet this increase was less prominent in elderly and type 2 diabetes. Our work identifies a new mechanism of posttranscriptional mitochondrial regulation in skeletal muscle through mitochondrial mRNA stabilization. Moreover, it emphasizes exercise as an effective approach to alleviate mitochondrial dysfunction.

Introduction

Muscle mitochondrial homeostasis and function are vital for skeletal muscle physiology, primarily influenced by the ability to meet energy demands through oxidative phosphorylation (OXPHOS) in mitochondria[1]. Exercise training can remarkably upregulate myofibrillar protein synthesis, OXPHOS proteins, and mitochondrial density to improve muscle function and respiratory capacity[2]. Reduced muscle function adversely impacts health and quality of life[1,3-10] and is associated with increased all-cause mortality, which can be ameliorated by exercise training[11-14]. The beneficial effects of exercise training are more potent than any drug in preserving muscle mass and function, positioning exercise training as a frontline strategy for the prevention and treatment of sarcopenia, type 2 diabetes (T2D), and cancer[15-18]. Viewing mitochondrial control through the lens of exercise biology is a strategy to gain new mechanistic insights into mitochondrial regulation, which is currently poorly understood. Apart from exercise training regimens, there are no current treatments to target muscle functional decline.

OXPHOS proteins are of dual origin and transcribed in spatially distinct cellular compartments. Nuclear (n)DNA encodes for the majority of OXPHOS proteins. Yet, 13 essential OXPHOS proteins are encoded by the small circular high-copy number mitochondrial DNA (mtDNA)[19,20]. In non-muscle systems, a protein complex comprised of steroid receptor RNA activator stem-loop interacting RNA-binding protein (SLIRP) and the leucine-rich pentatricopeptide repeat containing protein (LRPPRC) mediates mitochondrial-mRNA (mt-mRNA) stability and polyadenylation of most mtDNA-encoded OXPHOS transcripts[21-25]. Stabilization of mt-mRNA is critical for protein translation by the mitochondrial ribosome (mitoribosome). Yet, the mitoribosome comprises 82 mitoribosomal proteins encoded by nuclear genes, and 12S and 16S mt-rRNAs, highlighting the requirement for intricate coordination between the processes of cytosolic and mitochondrial protein synthesis[22,23]. Despite the high mitochondrial abundance in skeletal muscle, the complex interaction between LRPPRC/SLIRP-mediated posttranscriptional processes, mitoribosomal translation, and mitochondrial function has not been studied before in skeletal muscle. Elucidating the underlying mechanisms for stabilizing mt-mRNA could not only facilitate our understanding of the fundamental energy metabolism in muscle, but also aid intervention strategies for diseases associated with skeletal muscle mitochondrial defects.

Mounting evidence suggests that improvements in mitochondrial metabolism contribute to the exercise-

induced functional benefits, as mitochondrial preservation protects against the age-associated decline

in skeletal muscle mass and performance[17,26]. Endurance exercise training potently increases

mitochondrial mass in skeletal muscle, in part due to the increased content of nDNA- and mtDNA-

encoded mitochondrial proteins[1,2]. Importantly, exercise training retains its ability to improve oxidative

capacity not only in healthy individuals but also in patients with mtDNA mutations, with common PGC-

1α-mediated mitochondrial adaptive responses shared between both groups[16]. These results suggest that

exercise training induces adaptations to reinforce the mitochondria's ability to efficiently respond to

increased energy demands, even in the presence of mtDNA mutations, that may negatively affect

mitochondrial transcription and translation. However, a notable knowledge gap persists in

understanding the role of mitochondrial posttranscriptional processes, specifically in mt-mRNA

stabilization and translation, in skeletal muscle biology and following exercise training. Yet, that

knowledge is needed to gain a comprehensive understanding of the adaptive responses to exercise

training.

Since SLIRP, a key player in mitochondrial posttranscriptional gene expression, is markedly

upregulated in mouse skeletal muscle by exercise training[27], we hypothesized that SLIRP would

regulate mitochondrial function in skeletal muscle at rest and in response to exercise training. Our

findings illuminate SLIRP´s role in regulating mt-mRNA transcript levels in skeletal muscle,

downstream of PGC-1α1. Knockout (KO) of SLIRP led to damaged and fragmented mitochondria

alongside lowered respiration. Intriguingly, exercise training was able to compensate for the absence of

SLIRP, leading to improvements in mitochondrial integrity and respiratory capacity. This implies the

activation of complex exercise-induced molecular signaling in skeletal muscle with mt-mRNA defects,

possibly through enhanced mitoribosomal translation.

Results

***Slirp* knockout caused defects in mitochondrial structure and respiratory capacity in mouse**

muscle.

SLIRP was present across a diverse array of tissues in mice and was highly abundant in energy-

demanding tissues such as liver, kidney, brown adipose tissue, heart, and skeletal muscle (Fig. 1A,

Supplementary Fig. 1A), in line with previously reported *Slirp* mRNA levels[28]. Among the five different

skeletal muscle tissues analyzed, SLIRP protein content was highest in the oxidative soleus compared

to glycolytic extensor digitorum longus (EDL, Fig. 1A, Supplementary Fig. 1A).

In skeletal muscle of global *Slirp* KO mice, SLIRP protein was undetectable and its binding partner

LRPPRC showed 90% reduction in protein content (Fig. 1B). To corroborate previous findings on their

co-stabilization in non-muscle tissue[21,22,24], we administered intramuscular injections of recombinant

adeno-associated viral serotype 6 (rAAV6):*Slirp*. As expected, we observed that the absence of

concomitant upregulation of LRPPRC, failed to increase total SLIRP protein content due to

downregulation of endogenous SLIRP protein (Fig. 1C). These findings show that SLIRP and LRPPRC

stabilize each other in skeletal muscle.

We next sought to mechanistically determine the necessity of SLIRP for regulating mitochondrial

structure and respiratory capacity in skeletal muscle. We assessed mitochondrial network structure

using the membrane potential probe, tetramethylrhodamine ethyl ester (TMRE+), in intact flexor

digitorum brevis (FDB) muscle fibers of *Slirp* KO and littermate control wild-type (WT) mice.

Interestingly, the finely interconnected mitochondrial network, evident in WT fibers, was disrupted in

Slirp KO muscle fibers, evident by a 30% increased mitochondrial fragmentation index (Fig. 1D).

Transmission electron microscopy (TEM) analysis of gastrocnemius muscle provided further insights

into the ultrastructural changes of mitochondrial morphology in *Slirp* KO (Fig. 1E). Half of the KO

gastrocnemius muscles displayed reduced density of the matrix, disarray of the cristae, and substantial

enlargement and vacuolation of intermyofibrillar (IMF) mitochondria (Fig. 1E). Quantitative analysis

of the mitochondria showed that the percentage of damaged IMF mitochondria was increased (Fig. 1F).

However, there were no significant alterations in the total number of IMF mitochondria (Fig. 1G), and

no changes were observed in either the number of damaged or the total number of subsarcolemmal

mitochondria (Fig. 1G) or any changes in number of damaged or total number of subsarcolemmal

mitochondria (not shown). The percentage of damaged mitochondria in *Slirp* KO muscle varied

between 2-40% of total mitochondria compared to <2% of damaged mitochondria in WT (Fig. 1E, F).

Other structural parameters of IMF mitochondria, such as mitochondrial area, aspect ratio, elongation,

convexity, perimeter, sphericity, and diameter did not display any significant changes in *Slirp* KO

muscles, likely due to heterogeneity in the extent of damage (Fig. 1F, Supplementary Fig. 1B-H).

In agreement with the observable abnormalities in mitochondrial structure, maximal respiratory

capacity was reduced in permeabilized gastrocnemius muscle fibers of *Slirp* KO mice compared to WT

(Fig. 1H). This was particularly evident when adding glutamate to assess maximal complex I linked

respiratory activity (-24%) and succinate to assess complex I+II linked respiratory capacity (-21%).

These results indicate an important role for SLIRP in skeletal muscle respiration, which contrast

findings in isolated mitochondria from liver and heart tissues, where *Slirp* KO did not compromise

mitochondrial respiration[22]. Additionally, in intact incubated soleus muscle, fatty acid oxidation tended

(p=0.0718) to be lower in *Slirp* KO mice (Fig. 1I). However, electrically-induced muscle contraction

increased fatty acid oxidation similarly in both genotypes, suggesting that fatty acid utilization for fuel

during muscle contraction does not depend on SLIRP (Fig. 1I). Moreover, we observed a 60-80%

reduction in mtDNA-encoded mRNA transcripts of *mt-Nd1*, *mt-Nd5/Nd6* (Complex I), *mt-CytB*

(Complex III), *mt-Co1* (Complex IV), and *mt-Atp6* (ATP synthase) in gastrocnemius muscle (Fig. 1J),

suggesting that SLIRP stabilizes mt-mRNA in skeletal muscle. Together, these results suggest that mt-

mRNA stabilization via SLIRP is required for proper mitochondrial network structure and morphology,

and respiration in mouse muscle.

SLIRP knockdown impaired locomotion and reduced lifespan in flies.

To determine the long-term consequences of SLIRP deficiency on the whole organism, we utilized the

UAS-GAL4 system[29] for muscle-specific (*Mef2*-GAL4>) knock-down (KD) of *SLIRP1* and *SLIRP2* in

Drosophila melanogaster. Only one *SLIRP* gene is present in mouse and human, while two *SLIRP*

genes exist in *D. melanogaster* (Flybase annotation symbols (http://flybase.org): CG33714 and

CG8021) likely to have originated from gene duplication events[30]. This consequently gives rise to two

fly orthologue proteins of the human and mouse SLIRP, denoted *SLIRP1* (CG33714) and *SLIRP2*

(CG8021)[30]. To test physical functionality, we subjected the flies to a negative geotaxis (climbing)

assay[31] and found that *SLIRP2* KD, but not *SLIRP1* KD, flies climbed 23% slower than control flies,

indicative of impaired locomotion (Fig. 1K). The detrimental effects of *SLIRP* KD in the muscle became

further apparent following starvation. Median fasting survival of *SLIRP1* KD flies was 41 h, compared

with 44 h for control and *SLIRP2* KD flies (Fig. 1L, p=0.0681). Muscle-specific *SLIRP* deficiency,

irrespective of orthologue, had a deleterious effect on life span, as both *SLIRP1* and *SLIRP2* KD

significantly reduced lifespan (Fig. 1M). Accordingly, while the median survival of control flies was

77 days, median survival of *SLIRP1* and *SLIRP2* KD flies was only 55 and 63 days, respectively.

Taken together these findings in the flies show that *SLIRP* KD-induced mitochondrial defects can have

detrimental long-term consequences such as impaired muscle function, starvation intolerance, and

reduced life span.

Muscle SLIRP protein content increases in response to exercise training, under the regulation of

PGC-1α1.

Having established critical functional roles of SLIRP in skeletal muscle mitochondrial morphology and

respiratory capacity with detrimental effects on lifespan, we next investigated the upstream regulation

of SLIRP protein content. Our recent work pinpointed SLIRP as a protein responsive to exercise

training[27]; yet the mechanisms governing its induction and functions remained elusive. Knowing that

exercise potently increases PGC-1α protein and mRNA levels in both mouse and human skeletal

muscle[32-35], and that PGC-1α is an important regulator of mitochondrial biogenesis, respiration, and

quality control[36], we explored the regulation of SLIRP by PGC-1α. The *Ppargc1a* gene encodes several

PGC-1α isoforms, including isoform 1 (PGC-1α1) and isoform 4 (PGC-1α4), with distinct regulation

and biological functions[37,38]. Overexpression (OE) of PGC-1α1 in mouse skeletal muscle[37,39], associated

with endurance exercise training-like adaptations[37,39,40], resulted in approximately 8.3-fold higher

muscle SLIRP protein content, without a concomitant change in mRNA levels (Fig. 2A). On the

contrary, PGC-1α4 OE[41], inducing resistance-type exercise training adaptations such as muscle

hypertrophy and strength[37], had no effect on SLIRP protein content in mouse gastrocnemius muscle

(Fig. 2B). This suggests that SLIRP is an unrecognized player in PGC-1α1-regulated oxidative

metabolism.

To test whether PGC-1α regulates *Slirp* mRNA in skeletal muscle following exercise, we determined

Slirp and *Ppargc1a* (Exon 3-5) mRNA levels in tibialis anterior (TA) muscle 3 h into recovery from

one exercise bout in mice lacking PGC-1α in skeletal muscle (PGC-1α mKO) and WT control

littermates. *Slirp* mRNA was not upregulated 3 h post-exercise (Fig. 2C), in contrast to the expected[35]

increase in *Ppargc1a* (+6.8-fold; Fig. 2C) in WT mice. Interestingly, *Slirp* mRNA levels were only

modestly 14% reduced in PGC-1α KO skeletal muscle (Fig. 2C), in line with the dissociation between

Slirp mRNA levels and SLIRP protein levels observed in PGC-1α1 OE skeletal muscle (Fig. 2A).

These findings align with reports describing only moderate correlations between mRNA and protein

levels following exercise[42-44]. Yet, with the interpretative limitations of a single post-exercise time point,

we aimed to acquire a more detailed record of the temporal changes in *Slirp* transcripts in recovery from

exercise. Accordingly, we subjected a cohort of WT mice to an acute exercise bout at 60% of their

maximal running capacity and harvested quadriceps muscles at rest, immediately after, 2h, 6h, and 24h

post-exercise. The phosphorylation of AMP-activated protein kinase (AMPK) at T172 served as a

validation of elevated metabolic stress during the exercise bout (Fig. 2E). We found that *Slirp* mRNA

levels were 1.4-fold elevated 6h into the recovery period in quadriceps muscle (Fig. 2D), a time-point

not included in our prior cohort (Fig. 2C), and partially concurrent with *PGC-1α1* mRNA levels that

were elevated 2h (+3.0-fold) and 6h post-exercise (+3.3-fold; Fig. 2D). Following a single bout of

exercise, we observed no changes to SLIRP protein content (Fig. 2E). These findings indicate that the

SLIRP promoter underlies PGC-1α1-regulated post-transcriptional control following acute exercise.

Given that SLIRP might be downstream of PGC-1α, we next turned to investigate the long-term

dependency of exercise training-induced regulation of SLIRP protein by PGC-1α. In contrast to acute

exercise, exercise training increased SLIRP protein content 4.8-fold in WT mice (Fig. 2F),

recapitulating our earlier observations in mice engaged in voluntary wheel running[27]. The effect of

exercise training on SLIRP protein content was markedly blunted by 70% in mice lacking PGC-1α in muscle compared to trained WT mice (Fig. 2F). Expectedly, in WT mice, exercise training potently upregulated the steady-state levels of multiple OXPHOS proteins including SDHB, UQCRC2, MTCO1 in skeletal muscle (Supplementary Fig. 1I-K). In mice lacking PGC-1α in muscle, SDHB (Supplementary Fig. 1I) and MTCO1 protein (Supplementary Fig. 1K) was 50-60% decreased in SED mice. Moreover, the effect of exercise training on SDHB (Supplementary Fig. 1I) and UQCRC2 protein content (Supplementary Fig. 1J) was blunted by 30% and 65%, respectively, in PGC-1α mKO compared to WT mice. In contrast to PGC-1α, the exercise-sensitive metabolic sensor AMPK[45] did not seem to regulate SLIRP expression, as SLIRP protein abundance was comparable to controls in muscle-specific double KO of AMPKα1 and AMPKα2 and WT mice[46,47] treated with or without the AMPK-activator AICAR for 4 weeks (Fig. 2G).

PGC-1α has marked effects on muscle endurance, partially attributed to its pivotal role in mitochondrial biogenesis, and respiration[39,48]. Our results suggest SLIRP as an unrecognized downstream target of PGC-1a in such exercise adaptations and oxidative metabolism.

SLIRP is dispensable for organismal adaptations to exercise training, yet, needed for improving blood glucose regulation after exercise training in male mice.

Having identified SLIRP as an exercise training responsive protein downstream of PGC-1a, we next determined SLIRP´s mechanistic involvement in exercise training-mediated organismal adaptations. To this end, we conducted a 10-week voluntary wheel-running training study in *Slirp* KO and WT littermate mice of both sexes commencing at ~18 weeks of age. The experimental design is shown in Fig. 3A.

The daily running distance recorded over 6 weeks of the 10-week study was 2.4 km/day on average for male mice of both genotypes, whereas it was 4.8 km/day for female exercise-trained (ET) WT mice and 3.0 km/day for female KO ET mice (Fig. 3B). Of note, the running distance observed in our study including 18-week old mice is shorter than those reported in other exercise training studies using ~ 10-week-old mice[27,49,50]. Despite the higher food intake in ET mice (Fig. 3C), exercise training lowered

body weight in both trained genotypes (Supplementary Fig. 2A). Moreover, irrespective of genotype,

exercise training reduced fat mass/body weight ratio (Supplementary Fig. 2B) and increased lean

mass/body weight ratio compared to sedentary (SED) groups (Supplementary Fig. 2C). Exercise

training had no effect on gastrocnemius or TA muscle weight relative to tibia length in either sex

(Supplementary Fig. 2D). However, we noted that female *Slirp* KO SED and both ET groups had

elevated heart weight relative to WT SED (Supplementary Fig. 2D).

Exercise capacity increased similarly with exercise training, independently of genotype (+1.2-fold for

WT ET, +1.3-fold for KO ET; Supplementary Fig. 2E) and sex (females shown in Supplementary Fig.

2G). Post-exercise blood lactate levels tended to be reduced (-1.7-fold; p=0.067) by ET in WT male

mice (Supplementary Fig. 2F). The training-induced lowering of blood lactate post-exercise was absent

in male and female *Slirp* KO mice (Supplementary Fig. 2F, H).

Exercise training is a powerful preventative treatment against many metabolic disorders[15], partially due

to its remarkable effects in improving glucose tolerance and insulin sensitivity documented in both

animal models and humans[27,49,51]. Accordingly, male WT ET mice displayed reduced fasting blood

glucose (Fig. 3D) and improved blood glucose control evidenced by a downshifted glucose response

curve compared to WT SED mice. This effect was not observed in trained male *Slirp* KO mice (Fig.

3E). Thus, trained *Slirp* KO mice displayed similar glucose tolerance as both SED groups. The

incremental area under the curve (iAUC) of blood glucose was similar between groups (Fig. 3F),

suggesting that the lowered plasma glucose excursion during the GTT was driven by lower basal

glucose levels. Yet, exercise training reduced the levels of plasma insulin in both WT and *Slirp* KO

mice, suggesting that insulin sensitivity was increased by exercise training in both groups (Fig. 3G).

There was no effect of exercise training or genotype on fasting glucose, glucose tolerance or insulin

levels in female mice (Fig. 3H-K).

Our results demonstrate that SLIRP is largely dispensable for organismal adaptations to exercise

training, yet, in male mice required for training-induced improvement in blood glucose regulation.

Exercise training reverses defects in muscle mitochondrial structure and function elicited by lack of SLIRP.

Exercise training can counteract mitochondrial damage, arising from excessive accumulation of reactive oxygen species (ROS) or impaired assembly of OXPHOS complex following mtDNA mutations[16,17,52-54]. As we established that loss of SLIRP had marked negative implications for mitochondrial structure and respiration in skeletal muscle (Fig. 1), we asked whether exercise training could rescue these defects.

Skeletal muscle mitochondrial network structure was qualitatively investigated in TMRE+ stained FDB fibers by confocal microscopy. Remarkably, exercise training completely rescued the derangements in mitochondrial network structure and mitochondrial fragmentation observed in *Slirp* KO SED mice, quantitatively corroborated by restoration of the fragmentation index (Fig. 4A).

Concomitantly with improved mitochondrial structure, exercise training restored the respiratory flux in *Slirp* KO mice to the level of WT muscle (Fig. 4B). WT mice did not improve respiratory flux with exercise training, which was consistent with no effect of exercise training on fatty acid oxidation of contracting isolated WT soleus muscle (Supplementary Fig. 2I) and aligns with other studies[55]. Exercise training in *Slirp* KO mice improved fatty acid oxidation in contracting *Slirp* KO soleus muscles (Supplementary Fig. 2I), in alignment with improved respiratory flux in *Slirp* KO muscles by exercise training. Thus, exercise training counteracted the structural alterations in *Slirp* KO SED mice and had a positive impact on mitochondrial respiratory capacity, suggesting restored mitochondrial function.

Together, these results support that exercise training exploits the remarkable adaptive plasticity that mitochondria retain even in the event of mitochondrial network disruptions and impaired respiratory flux in skeletal muscle.

Sustained reductions of mitochondrial transcript levels can be bypassed by exercise training.

We next aimed to investigate the molecular underpinnings for the defects in mitochondrial structure and function in *Slirp* KO and their correction by exercise training. First, we verified that both SLIRP (+1.4-fold; Fig. 4C) and concomitantly LRPPRC (+1.4-fold; Fig. 4D) were upregulated by exercise

training in the gastrocnemius muscle in WT, but not *Slirp* KO mice (Fig. 4C, D). These findings

reinforce the co-stabilizing relationship of SLIRP and LRPPRC existing not only at baseline[21,22,24,56] but

also during exercise training conditions.

SLIRP, together with LRPPRC, has been shown to maintain mt-mRNA stability and aid mitoribosomal

translation in non-muscle tissues[21-24,56]. As a proxy for mitochondrial function[57], we found that mtDNA

copy number, measured using quantitative PCR and primers for mt-*Nd1* (Supplementary Fig. 2J) or *mt-*

Nd5/mt-Nd6 (Supplementary Fig. 2K), were unaltered across groups or elevated only in *Slirp* KO SED

mice, suggesting a compensatory response to mitigate mitochondrial dysfunction by increasing mtDNA

copy number[57]. In contrast, the marked downregulation of mt-mRNA transcripts in SED *Slirp* KO

muscle, also shown in Fig. 1J, remained reduced in ET *Slirp* KO mice relative to WT ET mice (Fig.

4F). Thus, SLIRP is suggested to be crucial for mitochondrial transcript stability in skeletal muscle.

Despite that, exercise training still restored protein content of mtDNA-encoded OXPHOS subunits,

MT-ND3, MT-CO1 and MT-ATP6 (Fig. 4H, I, J). Thus, exercise training completely counteracted the

sustained reduction in the mt-mRNA levels of these proteins in *Slirp* KO ET mice. The mild reduction

of MT-ND3 protein, a subunit of Complex I, seen in *Slirp* KO muscle was similarly rescued by exercise

training (Fig. 4H). Interestingly, although the nuclear-encoded Complex I subunit NDUFB8 was

downregulated by 57% in the *Slirp* KO SED mice, exercise training markedly upregulated NDUFB8

protein content in *Slirp* KO mice, yet, below NDUFB8 protein content in WT ET mice (Fig. 4K). Other

nuclear-encoded OXPHOS subunits, such as SDHB (Complex II), UQCRC2 (Complex III) and ATP5A

(ATP synthase) were similarly upregulated by exercise training, independently of *Slirp* KO (Fig. 4L-N,

representative blots shown in Fig. 4O). Citrate synthase (CS) activity correlates with mitochondrial

content in human[58] and mouse skeletal muscle[59]. CS activity, indicative of mitochondrial content, was

increased similarly in *Slirp* KO and WT mice after exercise training (Supplementary Fig. 2L),

suggesting that improvements in mitochondrial content following exercise training are independent of

SLIRP.

These results underline the substantial plasticity that mitochondria retain despite a 60-80% reduction in

mt-mRNA pools in skeletal muscle. Our findings point towards unidentified mechanisms by which ET

improves translational efficiency to dictate final protein content.

Exercise training increases mitochondrial protein translation capacity by elevating mitoribosome

mass.

Enhanced mitochondrial protein translation mediates exercise training adaptations in muscle

mitochondrial function in both young and elderly individuals[43]. Thus, we aimed to assess pathways

associated with protein translation in mitochondria, facilitated by the resident mitoribomes. We

observed training-induced increases in 12S ribosomal RNA (12S rRNA) and MRPL11 protein content

(Fig. 5A, B), components of the mitoribosomes. MRPL11 protein content was 36% more elevated in

response to exercise training in *Slirp* KO ET than WT ET mice, mainly in male mice (Fig. 5B).

Conversely, exercise training did not induce changes in cytosolic ribosomal protein S6 protein content

(rpS6, Fig. 5C; representative blots in Fig. 5D), indicating a compartmentalized ribosomal response to

exercise training.

Given the predominant impact of *Slirp* KO on mitochondria, training-induced adaptive changes might

also manifest directly within the affected cellular compartment. Here, we investigated a mitochondrial

quality control system involving a mitochondrial scavenger enzyme, peroxiredoxin 3 (PRDX3).

PRDX3 plays a role in balancing redox environment in mitochondria[60,61], possibly contributing to the

preservation of factors critical for translation during conditions of mitochondrial dysfunction and

exercise training. Interestingly, PRDX3 protein content was 60% lower in *Slirp* KO SED compared to

WT SED mice, indicative of a lower mitochondrial oxidative stress defense capacity in *Slirp* KO

muscle. This was restored by exercise training, bringing PRDX3 up to levels observed in trained WT

mice (Fig. 5E). We further noted that ET lowered the PRDX3 dimer to monomer ratio in both

genotypes, indicative of a lower basal mitochondrial-derived peroxide accumulation likely due to an

enhancement of scavenger activity (Fig. 5F). The cytosolic peroxidoxin 2 (PRDX2) was unaffected by

genotype and exercise training (Fig. 5G, representative blots in Fig. 5H), further pointing towards highly

mitochondria-selective effects of exercise training and SLIRP.

Collectively, our findings indicate a complex interplay of spatially distinct molecular muscle adaptations in response to *Slirp* KO and ET. Our findings suggest that exercise training induces mechanisms to increase mitochondrial protein translation capacity. Mechanistically, exercise training elevated mitoribosome mass and restored oxidative stress defense systems to powerfully circumvent the *Slirp* KO-induced depletion of mtDNA-encoded transcript levels.

Elevation of muscle SLIRP and LRPPRC protein content in response to ET is conserved in human skeletal muscle.

Our intriguing results from mouse and fly models prompted us to test whether our results provide translational value for humans in health and disease. SLIRP and LRPPRC protein content were determined in human skeletal muscle in four independent exercise cohorts employing different exercise modalities in conjunction with or without aging or diabetes and in males and females. Participant characteristics and study designs for the different ET modalities have been published previously for all the clinical studies[62-65].

Skeletal muscle SLIRP and LRPPRC protein abundances were 70% increased following 14-weeks of controlled and supervised aerobic and strength exercise training intervention in healthy young women[62] (Fig. 6A-C). Also, a 6-week high-intensity interval training (HIIT) intervention elevated muscle protein content of SLIRP (+80%, Fig. 5E, representative blots in Fig. 5D) and LRPPRC (+50%, Fig. 6F) in healthy young men[65]. Thus, the exercise training response of SLIRP was highly consistent, independent of sex, and responsive to various exercise modalities. The 12-week progressive resistance exercise training[63] increased SLIRP and LRPPRC in the entire group of young and old individuals but had no significant effect on LRPPRC abundance in either young or elderly subjects separately (Fig. 6H, I, representative blots in Fig. 6G). The mtDNA-encoded transcript levels of *MT-ND6*, *MT-CYB*, *MT-CO1* and *MT-ATP6* were comparable across all groups (Fig. 6J-M), indicating that mitochondrial gene expression was not significantly affected by either aging or the training regimen in this study.

Exercise training-mediated improvements in mitochondrial content and function may be associated with improved glucose metabolism in individuals with obesity and T2D [66,67], although this association is not

always clear[68]. The training response of SLIRP and LRPPRC to HIIT[64] was conserved in individuals

who were lean (+150% and +54%, Fig. 5O) or obese (+85% and +45%, Fig. 6O). However, the exercise

response for LRPPRC, but not SLIRP, was blunted in patients with T2D in comparison to lean

individuals (Fig. 6P, representative blots in Fig. 6N), suggesting impaired sensitivity of muscle to some

mitochondrial adaptations to ET in T2D. At the mRNA level, mt-DNA encoded transcripts of *MT-ND1*,

MT-CYB, *MT-CO1* and *MT-ATP6* were reduced with HIIT across all groups (Fig. 6Q-T). Yet, protein

content of MT-CO2 (Fig. 6U) and other nuclear-encoded OXPHOS proteins (Supplementary Fig. 3A-

F) were significantly upregulated with HIIT. In line with previous findings[43], the upregulation in

OXPHOS protein abundance occurred despite lower levels of mRNA, and demonstrate a disassociation

between the transcriptome and proteome[69].

Together, we show that SLIRP and LRPPRC are robustly upregulated in skeletal muscle across various

exercise training modalities in men and women, lean and obese. However, the exercise training response

of LRPPRC was blunted in patients with T2D. This suggests that T2D is associated with decreased

sensitivity to some of the adaptive mitochondrial responses normally elicited by exercise training,

possibly influenced by exercise intensity or volume. Thus, our clinical data in humans provide strong

translational value of the SLIRP-deficient fly and mouse models and point towards a conserved

requirement of SLIRP for skeletal muscle health, and potentially the SLIRP-independent mechanisms

that compensate during exercise training.

Discussion

Our investigation led to six key findings that bridge substantial knowledge gaps concerning the role of

mitochondrial posttranscriptional mechanisms on skeletal muscle biology and their contributions to

exercise training adaptations. Our findings not only unravel a delicate interplay between the mt-mRNA-

stabilizing protein SLIRP, and the integrity of skeletal muscle mitochondria structure and performance,

but also underscore the potential of exercise training in promoting selective mitochondrial translation

capacity to circumvent these mechanisms.

First, lack of *Slirp* KO led to mitochondrial network disruption, mitochondrial fragmentation, and

reduced respiratory capacity in skeletal muscle in mice. Second, SLIRP deficiency impaired muscle

functionality, and reduced lifespan in flies. Third, SLIRP was identified as an exercise training-

responsive downstream target of PGC1-α. Fourth, SLIRP was needed for improving blood glucose

regulation after exercise training in male mice. Fifth, exercise training restored the mitochondrial

defects elicited by the lack of SLIRP on mitochondrial structure, respiration, and mtDNA-encoded

OXPHOS proteins, via mechanisms likely involving the upregulation of mitoribosome content and

enhanced antioxidant defense. Finally, we established a translational foundation for our findings by

substantiating the conservation of SLIRP´s response to exercise training in four independent human

cohorts integrating different exercise modalities, health, and age conditions.

Our first finding establishes SLIRP as an unrecognized player in regulating mitochondria content,

structure, and respiration in skeletal muscle. Our results are partially consistent with reports in heart,

liver, and kidney *Slirp* KO[21,22,56] showing that SLIRP forms a complex with LRPPRC and is crucial for

the maintenance of the mt-mRNA reservoir. Accordingly, our observation of mitochondria with

abnormal cristae in *Slirp* KO muscle aligns with transmission electron micrographs of *Lrpprc* KO heart

tissue[23]. However, despite the stark similarities, mitochondrial respiration was compromised in *Slirp*

KO skeletal muscle, but not in heart or liver[22], underscoring SLIRP's functional importance within

skeletal muscle relative to other tissues. In alignment, previous research further showed that *Slirp* KO

skeletal muscle fibers have reduced sarcoplasmic reticulum Ca^{2+} storage capacity[70]. Thus, our present

findings underscore the importance of SLIRP across various tissue types, with emphasis on its

importance to skeletal muscle mitochondrial morphology and respiration, and overall mitochondrial

health.

Our second finding that SLIRP depletion compromised climbing ability, impaired starvation tolerance,

and reduced lifespan in flies, illustrates the detrimental functional consequences of lacking SLIRP in

the muscle tissue. The effects of mitochondrial proteins on modulating life span are equivocal. Some

report that the depletion of specific OXPHOS genes reduces life span, whereas others report no effect

or an increase on life span in model organisms[71-73]. The "Mitochondrial Threshold Effect Theory"[74]

posits that mild mitochondrial dysfunction allows normal physiology in model organisms until a

threshold is reached, beyond which severe mitochondrial dysfunction can lead to premature aging,

developmental arrest or even death[75-77]. In support, heterozygous depletion of *Mclk1*, which encodes an

enzyme necessary for biosynthesis of the electron carrier coenzyme Q of OXPHOS, even increased

lifespan in mice, but homozygous depletion was embryonically lethal[77]. Our findings support an

important role for SLIRP and mitochondrial function in maintaining normal lifespan because we

observed reduced survival in muscle-specific SLIRP KD flies.

Our third finding was that PGC-1α1 regulated SLIRP, providing evidence for a new training-responsive

PGC-1α target in skeletal muscle. Interestingly, training-induced adaptations in mitochondrial proteins

can occur independently of PGC-1a1 as the main regulator[78]. In agreement with that, our fifth finding

was that the training-induced improvements in mitochondrial function could bypass SLIRP deletion.

These findings agree with other studies showing that exercise training can rescue mitochondrial

dysfunction and elicit improvements in glucose homeostasis in mice and humans[16,52,78,79]. Interestingly,

SLIRP was required for the training-induced improvements in blood glucose regulation in trained male

Slirp KO mice. While not explored in our study, this could be due to decreased intramyocellular insulin

signalling[80], reduced insulin-independent glucose uptake[18], capillarization, and/or blood flow[81]. Female

mice did not improve their glucose tolerance in response to exercise training. Sexual dimorphism in

response to metabolic challenges, including exercise training, has been frequently reported in mouse

models[82,83]. Yet, in both sexes, SLIRP is largely dispensable for exercise training-induced mitochondrial

and metabolic adaptations in skeletal muscle, even though SLIRP is crucial for mitochondrial transcript

stability in the basal state.

We were intrigued by the observation that exercise training could circumvent even 60-80% loss in

mitochondrial-mRNA pools to upregulate mtDNA-encoded OXPHOS proteins. In other words, the

beneficial effect of exercise training on mitochondrial oxidative phosphorylation and ATP provision is

independent of a correction of mt-mRNA transcript levels and LRPPRC/SLIRP protein content. These

findings give rise to the possibility that mt-mRNAs are produced in excess *in vivo*[22]. Excess mt-mRNA

may allow for rapid engagement of mitochondrial protein synthesis in the event of sudden changes in

energy demand. In response to exercise training, the mitoribosomal components, MRPL11 and *12S*

rRNA were upregulated, indicating increased capacity for mitochondrial protein synthesis. These

adaptations to exercise training may help sustain protein translation, even in the presence of low mt-

mRNA abundance. Exercise training also upregulated PRDX3, linked to peroxide scavenging. While

MRPL11, *12S rRNA* and PRDX3 may not directly control translation efficiency, their roles in the

mitoribosomal makeup and improved redox state can influence the overall cellular environment where

protein translation occurs. Albeit limited by not directly measuring mitochondrial translational rate, this

working hypothesis is supported by findings in human skeletal muscle showing that mitochondrial

protein translation is at the core of exercise training-induced benefits[43].

Finally, our results likely are relevant to humans, as SLIRP and LRPPRC were consistently upregulated

in human skeletal muscle in response to exercise training. The effect of resistance exercise training was

blunted for LRPPRC in patients with T2D. Interestingly, the age-related decline in muscle

mitochondrial protein synthesis[84] can be reversed by exercise training in elderly[43], corroborating our

findings that ET bypasses SLIRP to increase mitoribosomal translation. Importantly, our findings

demonstrate that the synthesis of mtDNA-encoded OXPHOS proteins is not limited by transcript

abundance in skeletal muscle – a feature that seems to be conserved as a fundamental mechanism[30,43].

Taken together, our findings not only unravel a new mechanism for the regulation of basal

mitochondrial function in skeletal muscle, but also highlight an incredible exercise training –stimulated

plasticity of mitochondria in skeletal muscle facing mitochondrial defects. Our findings underscore ET

as a therapeutic intervention to combat mitochondrial dysfunction in genetic and lifestyle-induced

muscle pathologies, including T2D.

Conclusions

In this work, we uncover a conserved role for mitochondrial transcript stability via SLIRP in basal

mitochondrial structure and respiratory capacity in skeletal muscle. We identify SLIRP as a novel

exercise-responsive downstream target of PGC-1α that is conserved in human skeletal muscle. Because

the defects in *Slirp* KO muscle could be reversed by ET, these findings add to the clinical relevance and

add impetus to the important health-promoting role of ET in pathologies characterized by mitochondrial

defects.

Material and methods

Fly maintenance

In standard conditions, the flies were maintained on a SYA medium containing 0.5% (w/v) agar, 2.4%

(v/v) nipagin, 0.7% (v/v) proprionic acid, 10% (w/v) dry baker's yeast and 5% (w/v) sucrose. The

experiments took place at +25°C, 65% humidity under a 12h:12h light:dark cycle. RNAi lines for

SLIRP1 (51019 GD), *SLIRP2* (23675 GD), and the GD library control line (w1118, 60000 GD) were

obtained from Vienna Drosophila Resource Center. Mef2-GAL4 line was obtained from Bloomington

Stock Center to generate muscle-specific *SLIRP* KD flies.

Negative geotaxis (climbing) assay

Drosophila has a natural tendency to climb upwards, also known as negative geotaxis. The negative

geotaxis, or climbing, assay was performed as previously described[85]. Briefly, 20 male *Drosophila* flies

aged 5-6 days were transferred into empty vials (25 x 95mm), and another empty vial was taped on top

to create a 25 x 190mm cylinder. Six cylinders were set side by side into an apparatus. The apparatus

was tapped down 5 times to make all the flies fall to the bottom of the cylinder. The climbing assay was

measured in technical triplicate for each biological replicate, with 90 seconds allowed for each climb.

The assay was recorded with a camera and videos were analysed to determine the time for half of the

flies in the vial to reach halfway point (95mm) in the cylinder.

Starvation assay using the Drosophila Activity Monitoring System (DAMS)

Four-day old male flies were individually housed in monitor tubes with an outside diameter of 5mm

(PPT5x65, Trikinetics, Waltham, MA). For starvation analysis, the tubes contained 1% agar in water.

32 tubes per monitor (DAM2 Drosophila Activity Monitor, Trikinetics, Waltham, MA) were set up,

one monitor per genotype. For the starvation analysis, flies were kept in the DAM system until the last fly had died.

Lifespan assay

For lifespan assays, flies were mated at a controlled density of 25 female and 10 male flies, respectively. One-day old male flies were collected into vials (10 male flies per vial, with 10 vials per genotype). The flies were kept on a SYA diet in Drosoflippers (http://www.drosoflipper.com/). Flies were flipped onto fresh food every second day and deaths were scored during the transfer.

Mice

PGC-1α1 transgenic mouse (PGC-1α1 OE)

The generation of the model, quality, and specificity of the overexpression has been described previously[39], of which we analyzed quadriceps muscle samples.

PGC-1α4 transgenic mouse (PGC-1α4 OE)

The generation of the model, quality, and specificity of the overexpression has been described previously[39], of which we analyzed gastrocnemius muscle samples.

Skeletal muscle-specific PGC-1α knockout (PGC-1α mKO)

The generation of the model, quality, and specificity of the KO has been described previously[48]. Male muscle-specific PGC-1α MKO mice and littermate control mice homozygous for loxP inserts (lox/lox) were generated by crossbreeding myogenin-Cre mice with loxP flanked-Pgc-1α mice.

For the acute exercise bout, Tibialis anterior muscles of PGC-1α MKO mice and littermate control mice were collected 3 hours after one bout of equal distance treadmill running (1.4 km) at 10° incline and 60% of their individual maximal running speed achieved by a graded treadmill running test[86].

For the ET intervention, mice were single-housed with or without access to in-cage running wheels from 8 to 20 weeks old. Running distance and duration were monitored by a regular cycle computer. PGC-1α MKO mice tended to run less than lox/lox, thus running wheels of lox/lox mice were occasionally blocked to ensure equal running distance. On average, mice ran 25 km/week. The running wheels were blocked for all mice 24 hours prior to euthanization and harvest of quadriceps muscle in the morning in the fed state.

AAV6 Vector construction and preparation

The AAV6 vector for SLIRP overexpression and ultra-purified eGFP control AAV6 virus were manufactured by VectorBuilder Inc. (Shenandoah, Texas, USA; AAV6SP(VB190219-1010jvy)-C).

Transfection of tibialis anterior muscle using rAAV6 vector

Viral particles were diluted in Gelofusine (B. Braun, Germany) to a dosage of 5×10^9 vector genomes at a volume of 30 uL per injection. For muscle-specific delivery of rAAV6 vectors, 12-week-old mice were placed under general anaesthesia (2% isoflurane in O_2) and mice were injected intramuscularly with rAAV6:SLIRP in the TA muscle of one leg and control rAAV6:eGFP (empty vector) in the contralateral leg. Muscles were harvested 4 weeks after rAAV6 administration.

Time course of acute exercise

Twelve-week old female C57BL/6J mice (n=8 for each time point) were subjected to an acute exercise bout (1h running, 60% of maximal running intensity, 15° incline). Quadriceps muscle was harvested from rested and exercised mice immediately after, 2h, 6h, and 24 h after the acute exercise bout.

SLIRP knockout mouse (*Slirp* KO mice)

The generation of the model, quality, and specificity of the KO has been described previously[22]. Sperm of homozygous *Slirp* KO mice kindly provided by Nils-Göran Larsson and re-derived offspring was backcrossed to C57BL/6N background in our own animal facilities.

All mice were maintained under a 12:12 h light/dark photocycle at $22 \pm 2°C$ with nesting material. The female mice were group-housed (except during the ET intervention), whereas the male mice were

single-housed. All mice received a rodent chow diet (Altromin no. 1324; Chr. Pedersen, Denmark) and water *ad libitum*.

Mouse genotyping of *Slirp* KO and WT mice was performed as previously described [87] using qPCR on DNA from ear punches with the following primers: WT; 5'-AGAAGGGAAT CCACAGGATA GGACA-3' and 5'-GCTTTATTCC TAGTGCTGGC CTTGTT-3', KO; 5'-AGAAGGGAAT CCACAGGATA GGACA-3' and 5'-CGCCGTATAA TGTATGCTAT ACGAAGTT-3'.

10-week ET intervention in *Slirp* KO mice

For 10-week voluntary wheel-running exercise training interventions, 16–18-week-old **Slirp** KO and littermate mice were randomized into test groups with or without access to in-cage running wheels (Tecniplast activity cage, wheel diameter: 23 cm; Tecniplast, Buguggiate VA, Italy). The experimental design is schematically illustrated in Fig. 3A. Running distance was monitored by a regular cycle computer prior to the metabolic tests for 6 weeks. Running wheels were locked 12 h prior to terminal procedures to avoid effects of acute exercise bouts.

Body composition

Total, fat, and lean body mass was measured by nuclear magnetic resonance using an EchoMRI™ (USA).

Glucose tolerance test (GTT) and plasma insulin analysis

We subjected the *Slirp* KO and WT mice to a GTT at 7 weeks of the training intervention study (as illustrated in Fig. 3A). The GTT was executed after a 5 h fasting period (7:00 a.m.–12:00 p.m.). Resting blood samples were taken from the tail 30 min prior to the intraperitoneal injection of d-mono-glucose (2 g/kg body weight). Tail blood glucose was measured after 0, 20, 40, 60, and 90 min of injection.

To measure glucose-stimulated plasma insulin concentration at time-point (min) 0 and 20, tail vein blood samples were collected in a capillary-tube (50 µl), centrifuged at 14,200 g for 5 min at 4 °C, plasma collected and stored at -80°C. Insulin concentration was determined in duplicates using the Ultra-Sensitive Mouse Insulin ELISA Kit (#80-INSTRU-E10; ALPCO Diagnostics) to the

573   manufacturer's instructions. The incremental area under the curve (iAUC) from the basal blood glucose

574   concentration was determined using the trapezoid rule.

Exercise capacity tests

Slirp KO and WT mice were acclimatized to the treadmill on 3 consecutive days by running at a speed

of 0.16 m/s and incline of 10° for 5 min the first day and 10 min the following days. Prior to the running

test and with a 1h delay, 2 blood samples were drawn from the tail before the test for pre-exercise blood

glucose and blood lactate measurements. Afterwards, the mice ran at 0.16 m/s for 5 minutes followed

by a gradual increase in speed every minute with 0.02 m/s until exhaustion. Once the mouse reached its

maximal running capacity, 2 blood samples were immediately drawn from the tail for post-exercise

blood glucose and blood lactate measurements. The test was stopped when the mouse failed to keep up

with the treadmill despite motivational efforts by the researcher.

TMRE staining in live fibers

Flexor digitorum brevis muscles (FDB) were dissected and then incubated for 2 hours in serum-free α-

MEM (22571-020, Gibco) containing 1.9 mg/ml collagenase type 1 from clostridium histolyticum

(C0130, Sigma-Aldrich) and 1 mg/ml bovine serum albumin (9048-46-8, Sigma-Aldrich) at 37° C on

a rotator. After collagenase treatment, muscles were incubated in α-MEM containing 10% fetal bovine

serum (26050-70, Gibco) and subjected to mechanical dissociation using fire-polished Pasteur pipettes.

Single muscle fibers were seeded in 35x14mm glass-bottom microwell dishes (P35G-1.5-14-C, MatTek

Corporation) coated with 4 uL of Engelbreth-Holm-Swarm murine sarcoma ECM gel (E1270, Merck).

Muscle fibers were kept in α-MEM containing 5% fetal bovine serum in a cell incubator (37°C, 5%

CO_2) for at least 16 h prior to the experiments.

For determination of mitochondrial membrane potential (ΔΨmitochondrial) and mitochondrial network

morphology, the fibers were incubated in 20nM tetramethylrhodamine and ethyl ester (TMRE+, Life

Technologies) dissolved in Krebs Ringer buffer (145 mM NaCl, 5 mM KCl, 1 mM $CaCl_2$, 1 mM $MgCl_2$,

5.6 mM glucose, 20 mM HEPES, pH 7.4) for 30 minutes before imaging. Confocal images were

collected using a C-Apochromat ×40, 1.2 NA water immersion objective lens on an LSM 980 confocal

microscope (Zeiss) driven Zeiss Zen Blue 3.

Mitochondrial network analysis was performed semi-automatically in ImageJ (National Institute of

Health, USA). At least two nucleus-free regions per fiber were analyzed blindly. Before segmentation,

the background was subtracted, and the pixels were averaged using a value of mean=2. The images

were segmented, and particle analyses revealed the relative area of the TMRE+ signal compared to the

total area (% mitochondrial area). The fragmentation index was calculated as the number of objects in

relation to the total area covered by the dye. The ImageJ 'Red Hot' lookup table was used to visualize

the images.

Transmission electron microscopy analysis

A small longitudinal section (<2 mm) of the red portion of the gastrocnemius muscle tissue was fixed

by immersion 2% glutaraldehyde in 0.05M Phosphatebuffer, pH 7.4 and stored at 4°C. The samples

were rinsed four times in 0.1 M sodium cacodylate buffer, pH 7.4, and post-fixed with 1% osmium

tetroxide (OsO_4) and 1.5% potassium ferrocyanide [$K_4Fe(CN)_6$] in 0.1 M sodium cacodylate buffer, pH

7.4 for 90 min at 4°C. The samples were then rinsed twice and dehydrated through a graded mixture

of alcohol at 4–20°C, infiltrated with graded mixtures of propylene oxide and Epon at 20°C, and

embedded in 100% Epon at 30°C, as previously described (Nielsen et al., 2011). The samples were cut

in 60 nm longitudinal sections using a Leica Ultracut UCT ultramicrotome. The sections were

contrasted with uranyl acetate and lead citrate, and subsequently examined and image recorded in a

CM100 TEM (Philips, Eindhoven, The Netherlands) equipped with a Veleta camera and the iTem

software package (Olympus, Hamburg, Germany) with a resolution of 2048 x 2048 pixels.

A mean of eight fibers per sample (7-9) was included from the sectioned samples, and from each fiber,

24 images were obtained at x13,500 magnification. The imaging was performed in a randomized

systematic order including 12 images from the subsarcolemmal (SS) region, and 6 from both the

superficial and central region of the intermyofibrillar (IMF) space. The Z-disc width was measured once

on all IMF images and the mean from all fibers within one sample was calculated to determine the fiber

type. The categorization of fiber types was based on previous reports from observations[47]. The images

were analyzed by two genotype-blinded investigators and all further analyses were performed by the

same blinded investigator. The quantification of mitochondrial morphology was done using the Radius

EM Imaging Software (Emsis GmbH, Radius 2.0).

Several measurements were included in this study: 1) Mitochondrial number: Total number of

mitochondrial profiles (count). 2) Mitochondrial volume fraction: IMF mitochondrial volume is

annotated as percentage of mitochondria covering the IMF space. 3) Damaged mitochondria:

Mitochondria were categorized as damaged when they were swollen, vacuolated, or empty.

Mitochondrial respiration in gastrocnemius muscle

In a subset of mice, mitochondrial respiratory capacity was measured in permeabilized gastrocnemius

skeletal muscle fibers as previously described[88]. In brief, gastrocnemius muscle was rinsed from fat and

connective tissue and separated into small fiber bundles. Fiber bundles were permeabilized with saponin

(50 μg/ml) in BIOPS buffer for 30 min, followed by a 20 min wash in MiR05 buffer on ice.

Mitochondrial respiration was measured in duplicate under hyperoxic conditions at 37°C using high

resolution respirometry (Oxygraph-2k, Oroboros Instruments, Innsbruck, Austria). The following

protocol was applied: Leak respiration was assessed by addition of malate (5 mM) and pyruvate (5

mM), followed by adding three different concentrations of ADP (0.01 mM, 0.25 mM and 5 mM; for

the final concentration of ADP, 3 mM of magnesium (Mg) was added as well) to measure complex I

linked respiratory capacity. 10 mM Glutamate was added to measure maximal complex I linked

respiratory capacity followed by 10 mM succinate to measure complex I+II linked respiratory capacity.

Finally, 5 μM Antimycin A was added to inhibit complex III in the electron transport chain. Pooled data

for both sexes are shown as no sex-specific differences were detected.

Contraction-stimulated fatty acid oxidation in isolated soleus muscles

647   Contraction-stimulated exogenous palmitate oxidation in isolated soleus muscle from *Slirp* KO and WT

648   mice was measured as previously described [89]. In brief, excised soleus muscles from mice anesthetized

649   with pentobarbital were mounted at resting tension (∼5 mN) in 15 ml vertical incubation chambers with

650   a force transducer (Radnoti, Monrovia, CA) containing 30°C carbonated (95% O_2 and 5% CO_2) Krebs-

Henseleit Ringer buffer (KRB), pH=7.4, supplemented with 5 mM glucose, 2% fatty acid-free BSA, and 0.5 mM palmitate. After ~20 min of pre-incubation, the incubation buffer was refreshed with KRB additionally containing [1–14C]-palmitate (0.0044 MBq/ml; Amersham BioSciences, Buckinghamshire, U.K.). To seal the incubation chambers, mineral oil (Cat. No. M5904, Sigma–Aldrich) was added on top. Exogenous palmitate oxidation was measured simultaneously at rest and during 25 min contractions (18 trains/min, 0.6 s pulses, 30 Hz, 60 V). After incubation, incubation buffer and muscles were collected to determine the rate of palmitate oxidation as previously described[88,90,91]. Palmitate oxidation was determined as CO_2 production (complete FA oxidation) and acid-soluble metabolites (representing incomplete FA oxidation). As no difference was observed in complete and incomplete FA oxidation between genotypes, palmitate oxidation is presented as a sum of these two forms.

Lysate preparation and immunoblotting

Lysate preparation and immunoblotting of vastus lateralis muscles samples of HIIT in young, healthy men were performed as described by Hostrup et al.[92], whereas lysate preparation and immunoblotting of vastus lateralis muscles of HIIT in individuals, who were lean, obese or T2D were performed as described by Kruse et al[93].

To preserve and assess PRDX3 dimer to monomer ratio, freshly harvested mouse gastrocnemius muscle tissue was incubated for 10 min in ice-cold 100 mM N-Ethylmaleimide (NEM) diluted in PBS. NEM was subsequently aspirated and the tissue homogenized for 1 min at 30 Hz using a TissueLyser II bead mill (QIAGEN, USA) in ice-cold homogenization buffer [10% glycerol, 1% NP-40, 20 mM sodium pyrophosphate, 150 mM NaCl, 50 mM Hepes (pH 7.5), 20 mM β-glycerophosphate, 10 mM NaF, 2 mM phenylmethylsulfonyl fluoride, 1 mM EDTA (pH 8.0), 1 mM EGTA (pH 8.0), 2 mM Na_3VO_4, leupeptin (10 μg ml^{-1}), aprotinin (10 μg ml^{-1}), and 3 mM benzamidine]. Lysate preparation and immunoblotting of all remaining mouse tissues and human vastus lateralis muscle samples of aerobic and strength ET in women[62], and resistance training in the young and elderly cohorts[94] were performed as described in[27]. Immunoblotting of the NEM-treated samples was performed under non-denaturing conditions. The primary antibodies used are listed in Supplementary Table 2.

Coomassie Brilliant Blue staining was used as a control to assess total protein loading and transfer

efficiency[95] by quantifying the whole lane and for each sample set, a representative membrane from the

immunoblotting is shown. The same Coomassie brilliant blue staining is presented for proteins analyzed

when derived from the same sample set. Band densitometry was carried out using Image Lab (version

4.0). For each set of samples, a standard curve was loaded to ensure quantification within the linear

range for each protein probed for. For immunoblotting in Fig. 5, pooled data for both sexes are shown

as no sex-specific differences were detected, unless specified.

Supplementary Table 2: List of primary antibodies used during Western blotting analysis

Protein	Manufacturer	Identifier	Dilution
p-AMPK T172	Cell Signaling	Cat# 2531; RRID:AB_330330	1:1000, 2% Skim milk
ATP5F1B	ThermoFisher	Cat# A-21351; RRID:AB_221512	1:2000, 2% Skim milk
Hexokinase II	Cell Signaling	Cat# 2867; RRID:AB_2232946	1:1000, 2% Skim milk
LRPPRC	Abcam	Cat# ab97505; RRID:AB_10688419	1:1000, 2% Skim milk
MRPL11	Cell Signaling	Cat# 2066; RRID:AB_2145598	1:1000, 2% Skim milk
MT-ATP6	ThermoFisher Scientific	Cat# PA5-109432; RRID:AB_2854843	1:1000, 3% BSA
MT-CO2	Abcam	Cat# ab110413; RRID:AB_2629281	1:5000, 3% Skim milk
MT-ND1	Abcam	Cat# ab181848; RRID:AB_2687504	1:1000, 2% Skim milk
MT-ND3	Abcam	Cat# ab192306; RRID:AB_2943449	1:1000, 2% Skim milk
PRDX2	Abcam	Cat#ab109367; RRID:AB_10862524	1:1000, 2% Skim milk
PRDX3	Abcam	Cat#ab73349; RRID:AB_1860862	1:1000, 2% Skim milk
rpS6	Cell Signaling	Cat# 2217, RRID:AB_331355	1:1000, 2% Skim milk
SLIRP (mouse)	Santa Cruz Biotechnology	Cat# sc-514508; RRID: AB_2943449	1:500, 2% Skim milk
SLIRP (human)	Thermo Fisher Scientific	Cat# 703632; RRID:AB_2784591	1:2000, 2% Skim milk
Total OXPHOS Human WB Antibody Cocktail	Abcam	Cat# ab110411; RRID:AB_2756818	1:1000, 2% Skim milk
Total OXPHOS Rodent WB Antibody Cocktail	Abcam	Cat# ab110413; RRID:AB_2629281	1:5000, 2% Skim milk

RNA extraction and RT–qPCR

Fig. 1J, 4F: RNA was extracted from ~20 mg of pulverized whole mouse gastrocnemius muscle using TRIzol™ reagent (Invitrogen) following the manufacturer's instructions (tissue homogenization was performed using the MP Bio lysis system) and treated with the TURBO DNA-free™ Kit (Ambion) to remove contaminating DNA. For RT–qPCR expression analysis, cDNA was reversed transcribed from 0.85 μg total RNA using the High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems) in the presence of RNase Block (Agilent). The qPCR was performed in a QuantStudio 6 Flex Real-Time PCR System (Life Technologies), using TaqMan™ Universal Master Mix II, with UNG (Applied Biosystems) to quantify mitochondrial transcripts (mitochondrial-rRNAs and mt-mRNAs). The gene expression levels were determined using the ΔΔCt method, comparing the Ct values of mitochondrial transcripts to that of the beta-actin reference gene for normalization.

Fig. 2A, C, D: Total RNA was extracted from pulverized quadriceps muscle using an adapted guanidinium thiocyanate-phenol-chloroform extraction method. Reverse transcription to cDNA was performed as previously described[50]. Real-time qPCR was performed in triplicate with QuantStudio 7 Flex Real-Time PCR System (Applied Biosystems, Waltham, MA). Cycle threshold (Ct) was converted to a relative amount using a standard curve derived from a serial dilution of a representative pooled samples run together with the samples of interest. Beta-actin or Hprt mRNA was used for normalization of target mRNA levels.

Fig. 6J-M: Total RNA extraction from vastus lateralis muscle and reverse transcription to cDNA was performed as previously described[63]. The qPCR was performed in a QuantStudio 6 Flex Real-Time PCR System (Life Technologies), using TaqMan™ Universal Master Mix II (Applied Biosystems) to quantify mt-mRNAs. The gene expression levels were determined using the ΔΔCt method, comparing the Ct values of mitochondrial transcripts to that of the 18S rRNA reference gene for normalization.

Fig. 6Q-T: Total RNA was extracted from skeletal muscle biopsies using TRI Reagent (Sigma-Aldrich) following the manufacturer's instructions. cDNA was reversed transcribed using the High-Capacity

cDNA Reverse Transcription Kit (Applied Biosystems), while the qPCR was performed on an Aria Mx

(Agilent) using a TaqMan™ Universal Master Mix II (Applied Biosystems). The gene expression levels

were determined using the ΔΔCt method with the mRNA levels being normalized to the geometric

mean of *PPIA* and *B2M*.

Primers or Taqman probes or Taqman expression assays used for mRNA levels measured in Fig. 2,

Fig. 4 and Fig. 6 are shown in Supplementary Table 3.

Supplementary Table 3: List of primers or Taqman probes used for RT-qPCR

Primers used for Fig. 2C	
PGC-1a_Ex3-5_F	AGCCGTGACCACTGACAACGAG
PGC-1a_Ex3-5_R	GCTGCATGGTTCTGAGTGCTAAG
Slirp_F	GTCGGCCTATTGCGTTTGTC
Slirp_R	ATGTTCTCTCAGCTCACTCGC
Hprt F	AGTCCCAGCGTCGTGATTAG
Hprt R	TTTCCAAATCCTCGGCATAATGA

Primers/ Taqman gene expression assays used for Fig. 2A, D; Fig. 4F	
Slirp	Mm01296845_m1
Full length Ppargc1 F	TCAAGCCAAACCAACAACTTTATCT
Full length Ppargc1 R	GGTTCGCTCAATAGTCTTGTTCTCA
Ppargc1 probe	CACCAAATGACCCCAAGGGTTCCC
12S rRNA	
18S rRNA	Mm03928990_g1
ActB	Mm01205647_g1
mt-Nd1	Mm04225274_s1
mt-Nd6	Mm04225325_g1
mt-CytB	Mm04225271_g1
mt-Co1	Mm04225243_g1
mt-Atp6	Mm03649417_g1
Taqman gene expression assays used for Fig. 6	
MT-ND1	Hs02596873_s1
MT-ND6	Hs02596879_g1
MT-CYB	Hs02596867_s1
MT-CO1	Hs02596864_g1
MT-ATP6	Hs02596862_g1
PPIA	Hs99999904_m1
B2M	Hs99999907_m1

DNA isolation and mtDNA quantification

Total DNA was extracted from ~20 mg mouse gastrocnemius muscle using the Dneasy Blood and Tissue Kit (Qiagen) according to the manufacturer's instructions and treated with RNase A. Levels of mtDNA were measured by qPCR using 2.5 ng of DNA in a QuantStudio 6 Flex Real-Time PCR System using TaqMan™ Universal Master Mix II, with UNG. The mt-Nd1 and mt-Nd6/Nd5 TaqMan gene expression assays were used. The 18S probe was used for normalization.

Aerobic and strength ET intervention in healthy young women

A detailed description of the study participants, research design and methods has been previously published[63]. In the present study, we included vastus lateralis muscle lysate for immunoblotting obtained from 9 healthy young women (age, 33 ± 6 years; body mass index (BMI), 23.2 ± 2.6 kg m^{-2}) before and after 14-week of aerobic and strength exercise training.

High-intensity interval training intervention in healthy young men

A detailed description of the study participants, research design and methods has been previously published[65]. In the present study, we included vastus lateralis muscle lysate for immunoblotting obtained from 6 healthy young men before and after 6-week HIIT training.

Progressive resistance exercise training in male young and older individuals

A detailed description of the study participants, research design and methods has been previously published [49]. The original study reported no additive effect of vitamin D intake during the 12 weeks of resistance exercise training on muscle hypertrophy or muscle strength [62]. Accordingly, in the present study, samples from young or older participants were considered as one group, irrespective of vitamin D intake. We included vastus lateralis muscle samples (~10 mg wet weight) for immunoblotting obtained from a subset of the original sample set due to lack of sample material. For the young participants, we included 10 samples before and 6 samples after resistance exercise training. For the older participants we included 11 before and 9 samples after resistance exercise training.

High-intensity interval training in middle-aged male individuals, who were lean, obese or obese with type 2 diabetes

A detailed description of the study participants, research design and methods has been previously published[64]. In short, 15 middle-aged men with T2D and obesity, 15 age-matched glucose-tolerant men with obesity, and 18 age-matched glucose-tolerant lean men were recruited. All participants, except four (two men with T2D and obesity and two lean men) completed the study. The training protocol consisted of 8-weeks with three weekly training sessions consisting of supervised HIIT in combination with biking and rowing. HIIT-sessions consisted of 5 x 1 min exercise blocks interspersed with 1 min rest, and shifted between blocks on cycle and rowing ergometers. The volume was increased from two to five blocks during the 8 weeks. In the present study, we included vastus lateralis muscle samples that were obtained 4-5 days after the last HIIT session but 48 hours after the last physical activity (a VO_2 max test). The number of samples included for immunoblotting as follows: 16 before and after HIIT from lean glucose-tolerant men,15 before and after HIIT from glucose-tolerant men with obesity, and 13 before and after HIIT from men with T2D and obesity.

Ethical committee approval

Clinical experiments were approved by the Ethics Committee of Copenhagen or by the Regional Scientific Ethical Committees for Southern Denmark and performed in accordance with the Helsinki Declaration II. Studies including human study participants are described in Ref[62-65]. All mouse experiments complied with the European Convention for the protection of vertebrate animals used for experimental and other scientific purposes (No. 123, Strasbourg, France, 1985; EU Directive 2010/63/EU for animal experiments) and were approved by the Danish Animal Experimental Inspectorate (License number: 2016-15-0201-01043).

Graphical illustrations

Graphical illustrations were created in ©BioRender - biorender.com, as indicated.

Statistical methods

Statistical analyses were performed using GraphPad Prism 9 (version 9.3.1). Results are presented as
mean ± SEM with individual values shown, when feasible. Statistical differences were analyzed by
ordinary one-way ANOVA, repeated/ordinary two-way ANOVA, or Log-rank (Mantel-Cox) test, and
Mann-Whitney test as applicable. Dunnett's multiple comparisons test, or Šídák's multiple comparisons
test were used to evaluate significant interactions in ANOVAs. As statistical tests varied according to
the dataset being analyzed, the respective tests utilized are specified within the figure legends.
$0.05 \leq p < 0.1$ were considered a tendency and p values < 0.05 were considered significant.

Acknowledgements

We acknowledge the technical assistance of Betina Bolmgren and Irene Nielsen, Martin Thomassen,
and Roberto Meneses-Valdes (Department of Nutrition, Exercise and Sports, Faculty of Science,
University of Copenhagen, Denmark), Anja Jokipii-Utzon (Institute of Sports Medicine, Bispebjerg
Hospital, Copenhagen, Denmark), and Michala Carlsson (Department of Biomedical Sciences,
University of Copenhagen, Denmark). We thank Elise Needham and Vivian Shang for their assistance
with the *Drosophila* experiments (Charles Perkins Centre, The University of Sydney, Australia) and
Milica Popovic with the sample preparation for proteomic analysis (Cologne Excellence Cluster on
Cellular Stress Responses in Aging-Associated Diseases (CECAD), University of Cologne, Germany).
We acknowledge Cristiano di Benedetto for preparing muscle samples for TEM analysis (Core Facility
for Integrated Microscopy, Faculty of Health and Medical Sciences, University of Copenhagen,
Denmark).

Funding

789 The study was supported by the Novo Nordisk Foundation (grant NNF16OC0023418,
790 NNF18OC0032082, and NNF20OC0063577 to L.S.; grant NNF22OC0074110 to A.M.F.), by
791 Independent Research Fund Denmark to L.S. (#0169-00013B), by the European Union's Horizon 2020
792 research and innovation programme (Marie Skłodowska-Curie grant agreement No 801199 to T.C.P.P.
793 and E.A.R.), by the large instrument grant of the German Research Foundation (DFG

CRediT authorship contribution statement

Tang Cam Phung Pham: Conceptualization, Methodology, Validation, Formal analysis, Investigation,

Writing - Original Draft, Visualization, Project administration, Funding acquisition; **Steffen Henning**

Raun: Conceptualization; Investigation, Writing - Review & Editing; **Essi Havula**: Methodology,

Investigation, Writing - Review & Editing; **Carlos Henriquez-Olguín:** Methodology, Investigation,

Writing - Review & Editing; **Diana Rubalcava-Gracia:** Methodology, Investigation, Writing - Review

& Editing; **Emma Frank:** Methodology, Investigation, Writing - Review & Editing; **Andreas Mæchel**

Fritzen: Methodology, Investigation, Writing - Review & Editing; **Paulo Jannig:** Methodology,

Investigation, Writing - Review & Editing; **Nicoline Resen Andersen:** Investigation, Writing - Review

& Editing; **Rikke Kruse:** Investigation, Writing - Review & Editing; **Mona Sadek Ali:** Investigation,

Writing - Review & Editing; **Jens Frey Halling:** Resources, Writing - Review & Editing; **Stine**

Ringholm: Resources, Writing - Review & Editing; **Solvejg Hansen:** Resources, Writing - Review &

Editing; **Anders Krogh Lemminger:** Resources, Writing - Review & Editing; **Maria Houborg**

Petersen: Resources, Writing - Review & Editing; **Martin Eisemann de Almedia:** Resources, Writing

- Review & Editing; **Thomas E. Jensen:** Resources, Writing - Review & Editing; **Bente Kiens:**

Resources, Writing - Review & Editing; **Morten Hostrup:** Resources, Writing - Review & Editing;

Steen Larsen: Methodology, Investigation, Writing - Review & Editing; **Niels Ørtenblad:** Resources,

Writing - Review & Editing; **Kurt Højlund:** Resources, Writing - Review & Editing; **Peter**

Schjerling: Methodology, Investigation, Resources, Writing - Review & Editing; **Michael Kjær:**

Resources, Writing - Review & Editing; **Jorge Ruas:** Resources, Writing - Review & Editing;

Aleksandra Trifunovic: Resources, Writing - Review & Editing; **Jørgen Wojtaszewski;** Resources,

Writing - Review & Editing; **Joachim Nielsen:** Methodology, Investigation, Writing - Review &

Editing; **Klaus Qvortrup:** Resources, Writing - Review & Editing; **Henriette Pilegaard:** Resources,

Writing - Review & Editing; **Erik Arne Richter:** Resources, Investigation, Writing - Review &

Editing, Supervision, Funding acquisition; **Lykke Sylow**: Conceptualization, Methodology, Validation,

Formal analysis, Investigation, Writing - Original Draft, Project administration, Supervision, Funding

acquisition.

Figure Legends

Fig 1. *Slirp* knockout led to defects in mitochondrial structure and function, and reduced

lifespan

(A) SLIRP protein content across different wild-type tissues (n=4, female C57BL/6J, 12 weeks of age)

assessed by immunoblotting.

(B) SLIRP and LRPPRC protein content in tibialis anterior muscle of *Slirp* knockout (KO) and

littermate wildtype (WT) mice, and recombinant adeno-associated virus serotype 6 encoding SLIRP

(rAAV6:SLIRP) and rAAV6:eGFP as control mice.

(C) SLIRP and LRPPRC protein content in tibialis anterior muscle of recombinant adeno-associated

virus serotype 6 encoding SLIRP (rAAV6:SLIRP) and rAAV6:eGFP in the contralateral leg as control.

(D) Confocal microscopy of mitochondrial network structure using a mitochondrial membrane potential

probe (TMRE+) in flexor digitorum brevis muscle fibers of *Slirp* KO and WT mice (n=4-5, 7-10 fibers

per mouse). Corresponding fragmentation index are presented as super plots[96]; small symbols represent

each fiber, large symbols represent the mean of fibers from each mouse that are color- and symbol-

coded for each sex (○ (blue) male, ◇ (red) female).

(E) Transmission electron microscopy images of *Slirp* KO and WT gastrocnemius muscle.

(F and G) Quantification of percentage of damaged intramyofibrillar (IMF) mitochondria within total

mitochondrial and relative volume of mitochondria. Small symbols refer to damaged or total

mitochondria per fiber, large symbols of identical symbol refer to average of fibers per biological

replicate (female *Slirp* KO and WT gastrocnemius muscle, n=4/group).

(H) Mitochondrial respiration in of *Slirp* KO and WT gastrocnemius muscle (n=4-5/group, ○ (blue)

male, ◇ (red) female).

(I) Fatty acid oxidation in isolated *Slirp* KO and WT soleus muscle at rest and in response to contraction

(n=3-6/group, ○ (blue) male, ◇ (red) female).

(J) RT-qPCR analysis of mitochondrial transcript levels in gastrocnemius of male *Slirp* KO and WT

mice (n=6-7/group).

(K-M) Climbing time during climbing assay, starvation tolerance and life span of control (*Mef2-*

GAL4>GD control 6000), *SLIRP1* (*Mef2-GAL4*>*SLIRP1* RNAi 51019 GD) and *SLIRP2* (*Mef2-GAL4*>

SLIRP2 RNAi 23675 GD) knockdown flies.

Data are shown as mean ± SEM, including individual values, where applicable.

Geno, main effect of genotype; substrate, main effect of substrate addition; Substrate X Geno,

interaction between genotype and substrate; Contraction, main effect of contraction. *p<0.05,

**p<0.01, ***p<0.001, as per Unpaired Student´s t test (D, J), Mann Whitney test (F, G), Two-way

RM ANOVA with Šídák's multiple comparisons test (H, I), ordinary one-way ANOVA with Dunnett's

multiple comparisons test (K), Log-rank (Mantel-Cox) test (L, M).

Fig 2. SLIRP is upregulated by ET and is a target of PGC-1α

(A) RT-qPCR analysis of *Slirp* transcript levels relative to *Actb* levels and Western blot analysis of

SLIRP protein abundance in quadriceps with skeletal-muscle-specific transgenic expression of PGC-

1α1 (α1 OE; n=8-11/group)

(B) Western blot analysis of SLIRP protein abundance in gastrocnemius with skeletal-muscle-specific

transgenic expression of PGC-1α4 (α4 OE; n=6-9/group).

(C) RT-qPCR analysis of *Slirp* and *Ppargc1α* transcript levels relative to *Hprt* levels in gastrocnemius

of muscle-specific PGC-1α knockout (PGC-1α mKO) and littermate control (WT) mice at rest and 3h

after acute exercise bout (n=5-8/group).

(D) RT-qPCR analysis of *Slirp* and *Ppargc1α* transcript levels relative to *Actb* levels in WT quadriceps

at rest, immediately after, 2 h, 6 h or 24 h after acute exercise bout (n=8/group).

(E) Western blot analysis of SLIRP and pAMPK T172 protein abundance in WT quadriceps at rest, immediately after, 2 h, 6 h or 24 h after acute exercise bout (n=8/group).

(F) Western blot analysis of SLIRP protein abundance in quadriceps of sedentary (SED) and 12-week ET PGC-1α mKO and WT mice (n=9-10/group).

(G) Western blot analysis of SLIRP protein abundance in quadriceps of muscle-specific AMPKα1 and -α2 double KO (mDKO) and WT mice treated with/without AICAR (n=4-7/group).

Data are means ± SEM, including individual values. Geno, main effect of genotype; Ex, main effect of acute exercise; Geno x Ex, interaction between genotype and acute exercise. *p<0.05, ***p<0.001, as per Unpaired Student´s t test (A), Two-way ANOVA with Šídák's multiple comparisons test (C, F, G), and ordinary one-way ANOVA with Dunnett's multiple comparisons test (D, E).

Fig. 3. SLIRP is dispensable for organismal adaptations to ET, yet, needed for improving blood glucose regulation after ET in male mice

(A) Experimental design of 10-week exercise training (ET) intervention.

(B) Average running distance of male and female ET *Slirp* KO or littermate control (WT) mice measured for 6 weeks (n=6-11/group, ○ (blue) male, ◇ (red) female)

(C) Average food intake of male and female sedentary (SED) and ET *Slirp* KO or WT mice measured for 2 weeks after 4 weeks of running (n=3-8, ○ (blue) male, ◇ (red) female).

(D-K) Blood glucose levels following a 4-hour fasting period before the glucose tolerance test (GTT). Glucose tolerance of male and female SED and ET *Slirp* KO or WT mice after 7-weeks of ET. iAUC of glycemic excursion in response to bolus of glucose 2 g kg^{-1} body weight (BW). Insulin response before (0 min) and following (20 min) the oral glucose challenge (n=6-11/group, ○ (blue) male, ◇ (red) female).

Data are means ± SEM, including individual values where applicable.

Geno, main effect of genotype; ET, main effect of exercise training; Acute exercise, main effect of acute

exercise bout; Time X Geno ET, interaction between glucose bolus and genotype in the ET groups.

*$p<0.05$, **$p<0.01$; (E) WT ET vs. KO ET, *$p<0.05$; WT SED vs. WT ET, #$p<0.05$, as per Unpaired

Student´s t test (B), Two-way RM ANOVA with Šídák's multiple comparisons test (E, I), Two-way

ANOVA (C, D, F, G, H, J, K).

Fig 4. Exercise training reverses SLIRP-induced defects in muscle mitochondrial structure and

respiration despite sustained reductions of mitochondrial transcripts

(A) Confocal microscopy of mitochondrial network structure using a mitochondrial membrane potential

probe (TMRE+) in flexor digitorum brevis muscle fibers of sedentary (SED) and 10-week ET *Slirp*

knockout (KO) mice and littermate controls (WT) (n=3-4, 7-10 fibers per mouse), and corresponding

fragmentation index.

(B) Mitochondrial respiration measured by Oroboros respirometry system in gastrocnemius of SED and

ET *Slirp* KO and WT (n=2-6/per group, ○ male, ◇ female). SED data is also depicted in Fig. 1H.

(C and D) Western blot analysis of SLIRP and LRPPRC protein abundance in gastrocnemius of SED

and 10-week ET *Slirp* KO and WT mice (n=6-11/group, ○ (blue) male, ◇ (red) female).

(E) Schematic of mtDNA- and nuclear DNA-encoded oxidative phosphorylation (OXPHOS) proteins

analysed.

(F) RT-qPCR analysis of mitochondrial transcript levels in gastrocnemius of male SED and ET *Slirp*

KO and WT mice (n=6-11/group). SED data is also depicted in Fig. 1H.

(G-O) Western blot analysis of MT-ND1 (CI), MT-ND3 (CI), MT-CO1 (CIV), MT-ATP6 (CV),

NDUFB8 (CI), SDHB (CII), UQCRC2 (CIII) and ATP5A (CV), and representative blots in

gastrocnemius of SED and ET *Slirp* KO and WT mice (n=6-11/group, ○ (blue) male, ◇ (red) female).

Data are means ± SEM, including individual values. Geno, main effect of genotype; ET, main effect of

exercise training; Geno x ET, interaction between genotype and exercise training; Substrate, main effect

of substrate. *$p<0.05$, **$p<0.01$, ***$p<0.01$ as per Two-way ANOVA with Šídák's multiple

comparisons test (A, B, C, D, F-N).

Fig 5. Exercise training increases mitochondrial protein translation capacity by elevating

mitoribosome mass

(A-H) RT-qPCR analysis of 12S rRNA (A) and Western blot analysis of MRPL11 (B), rpS6 (C),

PRDX3 (E), PRDX3 dimer/monomer ratio (F), PRDX2 (G), and representative blots in gastrocnemius

of sedentary (SED) and exercise trained (ET) *Slirp* knockout (KO) and control littermate (WT) mice

(n=6-11/group, ○ (blue) male, ◇ (red) female). Data are means ± SEM, including individual values.

Geno, main effect of genotype; ET, main effect of exercise training; *p<0.05, **p<0.01, as per Two-

way ANOVA (A-C, E-G).

Fig 6. In human skeletal muscle SLIRP and LRPPRC protein content are increased by exercise

training

(A-C) Western blot analysis of SLIRP (B) and LRPPRC (C) protein in vastus lateralis of healthy young

women (n=9/group, (○ Pre ET, □ Post ET)) after a 14-week controlled aerobic and strength exercise

training (ET) intervention[62].

(D-F) Immunoblotting of SLIRP (E) and LRPPRC (F) protein in vastus lateralis of healthy young men

(n=6/group, (○ Pre ET, □ Post ET)) after a 6-week high intensity interval training[65].

(G-M) Immunoblotting of SLIRP (H) and LRPPRC (I) protein and RT-qPCR analysis of mitochondrial

transcript levels (J-M) in vastus lateralis of male young and older individuals after 12-week progressive

resistance training[63] (n=6-11/group, (○ Pre ET, □ Post ET)).

(N-U) Immunoblotting of SLIRP (O), LRPPRC (P) and MT-CO2 (U) protein and RT-qPCR analysis

of mitochondrial transcript levels (Q-T) in vastus lateralis of male individuals, who were lean, obese or

type 2 diabetic after high-intensity interval training[64] (n=13-16/group○ pre ET, □ post ET). Con, control;

OB, obese; T2D, Type 2 diabetes.

Data presented as individual before-and-after values. ET, main effect of exercise training, ET x T2D,

Interaction between exercise training and presence of type 2 diabetes in Con and T2D group; *p<0.05,

**$p < 0.01$, ***$p < 0.001$, as per paired Student´s t test (B, C, E, F), Mixed-effects model (H-M, Q-T),

Two-way RM ANOVA with Šídák's multiple comparisons test (O, P, U).

Supplementary Fig. 1

(A) Quantification of SLIRP protein content across different WT tissues (WT samples of n=4, 12 weeks

of age) assessed by immunoblotting.

(B-H) Transmission electron microscopy quantification of the following parameters of mitochondrial

area in the intermyofibrillar (IMF) region (B), aspect ratio (C), elongation (D), convexity (E), perimeter

(F), sphericity (G), and diameter (H). Small circles refer to each fiber, large squares in similar color

refer to average of fibers per biological replicate, (female, n=4).

(I-M) Western blot analysis of SDHB (CII), UQCRC2 (CIII), MT-CO1 (CIV), and ATP5A (CV)

protein abundance in quadriceps of sedentary (SED) and 12-week exercise-training (ET) PGC-1α mKO

and control littermates (WT) mice (subset of total cohort, male, n=3). Representative blots. Coomassie

was used as loading control.

Data are means ± SEM, including individual values. ET, main effect of exercise training; Geno, main

effect of genotype. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, as per Unpaired Student´s t test (B-D, F-H), Mann

Whitney test (E), Two-way ANOVA (I-L).

Supplementary Fig. 2

(A-C) Body weight, fat mass relative to body weight, lean mass relative to body weight of male and

female SED and ET *Slirp* KO or WT mice after 6-weeks of ET.

(D) Gastrocnemius, tibialis anterior (TA) and heart weight relative to tibia length of female SED and

ET *Slirp* KO or WT mice after 10-weeks of ET.

(E-H) Exercise capacity of male and female SED and ET *Slirp* KO or WT mice after 8-weeks of ET

and corresponding blood lactate levels before and after exercise bout.

(I) Fatty acid oxidation in isolated soleus of male and female SED and ET *Slirp* KO or WT mice after

10-weeks of ET. WT SED male/female, n=3/4, WT ET male/female, n=4/3, KO SED male/female,

n=4/9, KO ET male/female, n=4/7. SED data is also depicted in Fig. 1H.

(J-L) RT-qPCR analysis of mtDNA levels and citrate synthase activity in gastrocnemius of SED and

ET *Slirp* KO and WT mice.

Data are means ± SEM, including individual values where applicable. ET, main effect of exercise

training; Geno, main effect of genotype; Acute exercise, main effect of acute exercise; Geno in ET

group, main effect of genotype within exercise training group; Contraction, main effect of contraction.

*p<0.05, **p<0.01, ***p<0.001, as per Two-way ANOVA (A-E, G, J-L), Two-way RM ANOVA with

Šídák's multiple comparisons test (F, H, I).

Supplementary Fig. 3

(A-F) Western blot analysis of NDUFB8 (CI), SDHB (CII), UQCRC2 (CIII), ATP5A (CV) and ATP5B

(CV) protein in vastus lateralis of male individuals, who were lean (pre/post, n=16/16), obese (pre/post,

n=15/15) or type 2 diabetic (pre/post n=13/13) after high-intensity interval training[64]. Coomassie was

used as control. Data presented as individual before-and-after values. ET, main effect of exercise

training. *p<0.05, **p<0.01, ***p<0.001, Two-way RM ANOVA (A-E).

References

1.   Romanello, V., and Sandri, M. (2021). The connection between the dynamic remodeling of
the mitochondrial network and the regulation of muscle mass. Cell Mol Life Sci *78*, 1305-
1328. 10.1007/s00018-020-03662-0.
2.   Holloszy, J.O. (1967). Biochemical adaptations in muscle. Effects of exercise on mitochondrial
oxygen uptake and respiratory enzyme activity in skeletal muscle. J Biol Chem *242*, 2278-
2282.
3.   Distefano, G., Standley, R.A., Zhang, X., Carnero, E.A., Yi, F., Cornnell, H.H., and Coen, P.M.
(2018). Physical activity unveils the relationship between mitochondrial energetics, muscle
quality, and physical function in older adults. J Cachexia Sarcopenia Muscle *9*, 279-294.
10.1002/jcsm.12272.
4.   Gonzalez-Freire, M., de Cabo, R., Bernier, M., Sollott, S.J., Fabbri, E., Navas, P., and Ferrucci,
L. (2015). Reconsidering the Role of Mitochondria in Aging. J Gerontol A Biol Sci Med Sci *70*,
1334-1342. 10.1093/gerona/glv070.
5.   Janssen, I., Heymsfield, S.B., and Ross, R. (2002). Low relative skeletal muscle mass
(sarcopenia) in older persons is associated with functional impairment and physical
disability. J Am Geriatr Soc *50*, 889-896. 10.1046/j.1532-5415.2002.50216.x.
6.   Short, K.R., Bigelow, M.L., Kahl, J., Singh, R., Coenen-Schimke, J., Raghavakaimal, S., and Nair,
K.S. (2005). Decline in skeletal muscle mitochondrial function with aging in humans. Proc
Natl Acad Sci U S A *102*, 5618-5623. 10.1073/pnas.0501559102.
7.   Haines, M.S., Leong, A., Porneala, B.C., Meigs, J.B., and Miller, K.K. (2022). Association
between muscle mass and diabetes prevalence independent of body fat distribution in
adults under 50 years old. Nutr Diabetes *12*, 29. 10.1038/s41387-022-00204-4.
8.   Sugimoto, K., Tabara, Y., Ikegami, H., Takata, Y., Kamide, K., Ikezoe, T., Kiyoshige, E.,
Makutani, Y., Onuma, H., Gondo, Y., et al. (2019). Hyperglycemia in non-obese patients with
type 2 diabetes is associated with low muscle mass: The Multicenter Study for Clarifying
Evidence for Sarcopenia in Patients with Diabetes Mellitus. J Diabetes Investig *10*, 1471-
1479. 10.1111/jdi.13070.
9.   Yoon, J.W., Ha, Y.C., Kim, K.M., Moon, J.H., Choi, S.H., Lim, S., Park, Y.J., Lim, J.Y., Kim, K.W.,
Park, K.S., and Jang, H.C. (2016). Hyperglycemia Is Associated with Impaired Muscle Quality
in Older Men with Diabetes: The Korean Longitudinal Study on Health and Aging. Diabetes
Metab J *40*, 140-146. 10.4093/dmj.2016.40.2.140.
10.   Ebhardt, H.A., Degen, S., Tadini, V., Schilb, A., Johns, N., Greig, C.A., Fearon, K.C.H.,
Aebersold, R., and Jacobi, C. (2017). Comprehensive proteome analysis of human skeletal
muscle in cachexia and sarcopenia: a pilot study. J Cachexia Sarcopenia Muscle *8*, 567-582.
10.1002/jcsm.12188.
11.   Leitzmann, M.F., Park, Y., Blair, A., Ballard-Barbash, R., Mouw, T., Hollenbeck, A.R., and
Schatzkin, A. (2007). Physical activity recommendations and decreased risk of mortality. Arch
Intern Med *167*, 2453-2460. 10.1001/archinte.167.22.2453.
12.   Lee, D.H., Rezende, L.F.M., Joh, H.K., Keum, N., Ferrari, G., Rey-Lopez, J.P., Rimm, E.B.,
Tabung, F.K., and Giovannucci, E.L. (2022). Long-Term Leisure-Time Physical Activity Intensity
and All-Cause and Cause-Specific Mortality: A Prospective Cohort of US Adults. Circulation
146, 523-534. 10.1161/CIRCULATIONAHA.121.058162.
13.   Leong, D.P., Teo, K.K., Rangarajan, S., Lopez-Jaramillo, P., Avezum, A., Jr., Orlandini, A.,
Seron, P., Ahmed, S.H., Rosengren, A., Kelishadi, R., et al. (2015). Prognostic value of grip
strength: findings from the Prospective Urban Rural Epidemiology (PURE) study. Lancet *386*,
266-273. 10.1016/S0140-6736(14)62000-6.
14.   Ruiz, J.R., Sui, X., Lobelo, F., Morrow, J.R., Jr., Jackson, A.W., Sjostrom, M., and Blair, S.N.
(2008). Association between muscular strength and mortality in men: prospective cohort
study. BMJ *337*, a439. 10.1136/bmj.a439.

15. Pedersen, B.K., and Saltin, B. (2015). Exercise as medicine - evidence for prescribing exercise
as therapy in 26 different chronic diseases. Scand J Med Sci Sports *25 Suppl 3*, 1-72.
10.1111/sms.12581.
16. Adhihetty, P.J., Taivassalo, T., Haller, R.G., Walkinshaw, D.R., and Hood, D.A. (2007). The
effect of training on the expression of mitochondrial biogenesis- and apoptosis-related
proteins in skeletal muscle of patients with mtDNA defects. Am J Physiol Endocrinol Metab
293, E672-680. 10.1152/ajpendo.00043.2007.
17. Safdar, A., Bourgeois, J.M., Ogborn, D.I., Little, J.P., Hettinga, B.P., Akhtar, M., Thompson,
J.E., Melov, S., Mocellin, N.J., Kujoth, G.C., et al. (2011). Endurance exercise rescues
progeroid aging and induces systemic mitochondrial rejuvenation in mtDNA mutator mice.
Proc Natl Acad Sci U S A *108*, 4135-4140. 10.1073/pnas.1019581108.
18. Sylow, L., Kleinert, M., Richter, E.A., and Jensen, T.E. (2017). Exercise-stimulated glucose
uptake - regulation and implications for glycaemic control. Nat Rev Endocrinol *13*, 133-148.
10.1038/nrendo.2016.162.
19. Boengler, K., Heusch, G., and Schulz, R. (2011). Nuclear-encoded mitochondrial proteins and
their role in cardioprotection. Biochim Biophys Acta *1813*, 1286-1294.
10.1016/j.bbamcr.2011.01.009.
20. Couvillion, M.T., Soto, I.C., Shipkovenska, G., and Churchman, L.S. (2016). Synchronized
mitochondrial and cytosolic translation programs. Nature *533*, 499-503.
10.1038/nature18015.
21. Sasarman, F., Brunel-Guitton, C., Antonicka, H., Wai, T., Shoubridge, E.A., and Consortium, L.
(2010). LRPPRC and SLIRP interact in a ribonucleoprotein complex that regulates
posttranscriptional gene expression in mitochondria. Mol Biol Cell *21*, 1315-1323.
10.1091/mbc.E10-01-0047.
22. Lagouge, M., Mourier, A., Lee, H.J., Spahr, H., Wai, T., Kukat, C., Silva Ramos, E., Motori, E.,
Busch, J.D., Siira, S., et al. (2015). SLIRP Regulates the Rate of Mitochondrial Protein
Synthesis and Protects LRPPRC from Degradation. PLoS Genet *11*, e1005423.
10.1371/journal.pgen.1005423.
23. Ruzzenente, B., Metodiev, M.D., Wredenberg, A., Bratic, A., Park, C.B., Camara, Y.,
Milenkovic, D., Zickermann, V., Wibom, R., Hultenby, K., et al. (2012). LRPPRC is necessary
for polyadenylation and coordination of translation of mitochondrial mRNAs. EMBO J *31*,
443-456. 10.1038/emboj.2011.392.
24. Chujo, T., Ohira, T., Sakaguchi, Y., Goshima, N., Nomura, N., Nagao, A., and Suzuki, T. (2012).
LRPPRC/SLIRP suppresses PNPase-mediated mRNA decay and promotes polyadenylation in
human mitochondria. Nucleic Acids Res *40*, 8033-8047. 10.1093/nar/gks506.
25. Baughman, J.M., Nilsson, R., Gohil, V.M., Arlow, D.H., Gauhar, Z., and Mootha, V.K. (2009). A
computational screen for regulators of oxidative phosphorylation implicates SLIRP in
mitochondrial RNA homeostasis. PLoS Genet *5*, e1000590. 10.1371/journal.pgen.1000590.
26. Romanello, V., and Sandri, M. (2015). Mitochondrial Quality Control and Muscle Mass
Maintenance. Front Physiol *6*, 422. 10.3389/fphys.2015.00422.
27. Kleinert, M., Parker, B.L., Jensen, T.E., Raun, S.H., Pham, P., Han, X., James, D.E., Richter, E.A.,
and Sylow, L. (2018). Quantitative proteomic characterization of cellular pathways
associated with altered insulin sensitivity in skeletal muscle following high-fat diet feeding
and exercise training. Sci Rep *8*, 10723. 10.1038/s41598-018-28540-5.
28. Hatchell, E.C., Colley, S.M., Beveridge, D.J., Epis, M.R., Stuart, L.M., Giles, K.M., Redfern, A.D.,
Miles, L.E., Barker, A., MacDonald, L.M., et al. (2006). SLIRP, a small SRA binding protein, is a
nuclear receptor corepressor. Mol Cell *22*, 657-668. 10.1016/j.molcel.2006.05.024.
29. Brand, A.H., and Perrimon, N. (1993). Targeted gene expression as a means of altering cell
fates and generating dominant phenotypes. Development *118*, 401-415.
10.1242/dev.118.2.401.

30. Baggio, F., Bratic, A., Mourier, A., Kauppila, T.E., Tain, L.S., Kukat, C., Habermann, B.,
Partridge, L., and Larsson, N.G. (2014). Drosophila melanogaster LRPPRC2 is involved in
coordination of mitochondrial translation. Nucleic Acids Res 42, 13920-13938.
10.1093/nar/gku1132.
31. Inagaki, H.K., Kamikouchi, A., and Ito, K. (2010). Methods for quantifying simple gravity
sensing in Drosophila melanogaster. Nat Protoc 5, 20-25. 10.1038/nprot.2009.196.
32. Baar, K., Wende, A.R., Jones, T.E., Marison, M., Nolte, L.A., Chen, M., Kelly, D.P., and
Holloszy, J.O. (2002). Adaptations of skeletal muscle to exercise: rapid increase in the
transcriptional coactivator PGC-1. FASEB J 16, 1879-1886. 10.1096/fj.02-0367com.
33. Chinsomboon, J., Ruas, J., Gupta, R.K., Thom, R., Shoag, J., Rowe, G.C., Sawada, N.,
Raghuram, S., and Arany, Z. (2009). The transcriptional coactivator PGC-1alpha mediates
exercise-induced angiogenesis in skeletal muscle. Proc Natl Acad Sci U S A 106, 21401-21406.
10.1073/pnas.0909131106.
34. Pilegaard, H., Saltin, B., and Neufer, P.D. (2003). Exercise induces transient transcriptional
activation of the PGC-1alpha gene in human skeletal muscle. J Physiol 546, 851-858.
10.1113/jphysiol.2002.034850.
35. Kjobsted, R., Pedersen, A.J., Hingst, J.R., Sabaratnam, R., Birk, J.B., Kristensen, J.M., Hojlund,
K., and Wojtaszewski, J.F. (2016). Intact Regulation of the AMPK Signaling Network in
Response to Exercise and Insulin in Skeletal Muscle of Male Patients With Type 2 Diabetes:
Illumination of AMPK Activation in Recovery From Exercise. Diabetes 65, 1219-1230.
10.2337/db15-1034.
36. Halling, J.F., and Pilegaard, H. (2020). PGC-1alpha-mediated regulation of mitochondrial
function and physiological implications. Appl Physiol Nutr Metab 45, 927-936.
10.1139/apnm-2020-0005.
37. Ruas, J.L., White, J.P., Rao, R.R., Kleiner, S., Brannan, K.T., Harrison, B.C., Greene, N.P., Wu, J.,
Estall, J.L., Irving, B.A., et al. (2012). A PGC-1alpha isoform induced by resistance training
regulates skeletal muscle hypertrophy. Cell 151, 1319-1331. 10.1016/j.cell.2012.10.050.
38. Jannig, P.R., Dumesic, P.A., Spiegelman, B.M., and Ruas, J.L. (2022). SnapShot: Regulation
and biology of PGC-1alpha. Cell 185, 1444-1444 e1441. 10.1016/j.cell.2022.03.027.
39. Lin, J., Wu, H., Tarr, P.T., Zhang, C.Y., Wu, Z., Boss, O., Michael, L.F., Puigserver, P., Isotani, E.,
Olson, E.N., et al. (2002). Transcriptional co-activator PGC-1 alpha drives the formation of
slow-twitch muscle fibres. Nature 418, 797-801. 10.1038/nature00904.
40. Arany, Z. (2008). PGC-1 coactivators and skeletal muscle adaptations in health and disease.
Curr Opin Genet Dev 18, 426-434. 10.1016/j.gde.2008.07.018.
41. Handschin, C., Choi, C.S., Chin, S., Kim, S., Kawamori, D., Kurpad, A.J., Neubauer, N., Hu, J.,
Mootha, V.K., Kim, Y.B., et al. (2007). Abnormal glucose homeostasis in skeletal muscle-
specific PGC-1alpha knockout mice reveals skeletal muscle-pancreatic beta cell crosstalk. J
Clin Invest 117, 3463-3474. 10.1172/JCI31785.
42. Miller, B.F., Konopka, A.R., and Hamilton, K.L. (2016). The rigorous study of exercise
adaptations: why mRNA might not be enough. J Appl Physiol (1985) 121, 594-596.
10.1152/japplphysiol.00137.2016.
43. Robinson, M.M., Dasari, S., Konopka, A.R., Johnson, M.L., Manjunatha, S., Esponda, R.R.,
Carter, R.E., Lanza, I.R., and Nair, K.S. (2017). Enhanced Protein Translation Underlies
Improved Metabolic and Physical Adaptations to Different Exercise Training Modes in Young
and Old Humans. Cell Metab 25, 581-592. 10.1016/j.cmet.2017.02.009.
44. Liu, Y., Beyer, A., and Aebersold, R. (2016). On the Dependency of Cellular Protein Levels on
mRNA Abundance. Cell 165, 535-550. 10.1016/j.cell.2016.03.014.
45. Richter, E.A., and Ruderman, N.B. (2009). AMPK and the biochemistry of exercise:
implications for human health and disease. Biochem J 418, 261-275. 10.1042/BJ20082055.
46. Fentz, J., Kjobsted, R., Birk, J.B., Jordy, A.B., Jeppesen, J., Thorsen, K., Schjerling, P., Kiens, B.,
Jessen, N., Viollet, B., and Wojtaszewski, J.F. (2015). AMPKalpha is critical for enhancing

1133		skeletal muscle fatty acid utilization during in vivo exercise in mice. FASEB J *29*, 1725-1738. 10.1096/fj.14-266650.
1135	47.	Lantier, L., Fentz, J., Mounier, R., Leclerc, J., Treebak, J.T., Pehmoller, C., Sanz, N., Sakakibara, I., Saint-Amand, E., Rimbaud, S., et al. (2014). AMPK controls exercise endurance, mitochondrial oxidative capacity, and skeletal muscle integrity. FASEB J *28*, 3211-3224. 10.1096/fj.14-250449.
1139	48.	Handschin, C., Chin, S., Li, P., Liu, F., Maratos-Flier, E., Lebrasseur, N.K., Yan, Z., and Spiegelman, B.M. (2007). Skeletal muscle fiber-type switching, exercise intolerance, and myopathy in PGC-1alpha muscle-specific knock-out animals. J Biol Chem *282*, 30014-30021. 10.1074/jbc.M704817200.
1143	49.	Raun, S.H., Henriquez-Olguin, C., Karavaeva, I., Ali, M., Moller, L.L.V., Kot, W., Castro-Mejia, J.L., Nielsen, D.S., Gerhart-Hines, Z., Richter, E.A., and Sylow, L. (2020). Housing temperature influences exercise training adaptations in mice. Nat Commun *11*, 1560. 10.1038/s41467-020-15311-y.
1147	50.	Brandt, N., Kotowska, D., Kristensen, C.M., Olesen, J., Lutzhoft, D.O., Halling, J.F., Hansen, M., Al-Soud, W.A., Hansen, L., Kiilerich, P., and Pilegaard, H. (2018). The impact of exercise training and resveratrol supplementation on gut microbiota composition in high-fat diet fed mice. Physiol Rep *6*, e13881. 10.14814/phy2.13881.
1151	51.	Wojtaszewski, J.F., and Richter, E.A. (2006). Effects of acute exercise and training on insulin action and sensitivity: focus on molecular mechanisms in muscle. Essays Biochem *42*, 31-46. 10.1042/bse0420031.
1154	52.	Memme, J.M., and Hood, D.A. (2020). Molecular Basis for the Therapeutic Effects of Exercise on Mitochondrial Defects. Front Physiol *11*, 615038. 10.3389/fphys.2020.615038.
1156	53.	Carter, H.N., Pauly, M., Tryon, L.D., and Hood, D.A. (2018). Effect of contractile activity on PGC-1alpha transcription in young and aged skeletal muscle. J Appl Physiol (1985) *124*, 1605-1615. 10.1152/japplphysiol.01110.2017.
1159	54.	Lanza, I.R., Short, D.K., Short, K.R., Raghavakaimal, S., Basu, R., Joyner, M.J., McConnell, J.P., and Nair, K.S. (2008). Endurance exercise as a countermeasure for aging. Diabetes *57*, 2933-2942. 10.2337/db08-0349.
1162	55.	Schaefer, P.M., Rathi, K., Butic, A., Tan, W., Mitchell, K., and Wallace, D.C. (2022). Mitochondrial mutations alter endurance exercise response and determinants in mice. Proc Natl Acad Sci U S A *119*, e2200549119. 10.1073/pnas.2200549119.
1165	56.	Spahr, H., Rozanska, A., Li, X., Atanassov, I., Lightowlers, R.N., Chrzanowska-Lightowlers, Z.M., Rackham, O., and Larsson, N.G. (2016). SLIRP stabilizes LRPPRC via an RRM-PPR protein interface. Nucleic Acids Res *44*, 6868-6882. 10.1093/nar/gkw575.
1168	57.	Filograna, R., Mennuni, M., Alsina, D., and Larsson, N.G. (2021). Mitochondrial DNA copy number in human disease: the more the better? FEBS Lett *595*, 976-1002. 10.1002/1873-3468.14021.
1171	58.	Larsen, S., Nielsen, J., Hansen, C.N., Nielsen, L.B., Wibrand, F., Stride, N., Schroder, H.D., Boushel, R., Helge, J.W., Dela, F., and Hey-Mogensen, M. (2012). Biomarkers of mitochondrial content in skeletal muscle of healthy young human subjects. J Physiol *590*, 3349-3360. 10.1113/jphysiol.2012.230185.
1175	59.	Halling, J.F., Ringholm, S., Olesen, J., Prats, C., and Pilegaard, H. (2017). Exercise training protects against aging-induced mitochondrial fragmentation in mouse skeletal muscle in a PGC-1alpha dependent manner. Exp Gerontol *96*, 1-6. 10.1016/j.exger.2017.05.020.
1178	60.	Cox, A.G., Winterbourn, C.C., and Hampton, M.B. (2009). Mitochondrial peroxiredoxin involvement in antioxidant defence and redox signalling. Biochem J *425*, 313-325. 10.1042/BJ20091541.
1181	61.	Lee, K.P., Shin, Y.J., Cho, S.C., Lee, S.M., Bahn, Y.J., Kim, J.Y., Kwon, E.S., Jeong, D.Y., Park, S.C., Rhee, S.G., et al. (2014). Peroxiredoxin 3 has a crucial role in the contractile function of

skeletal muscle by regulating mitochondrial homeostasis. Free Radic Biol Med *77*, 298-306.
10.1016/j.freeradbiomed.2014.09.010.

62. Hansen, S.L., Bojsen-Moller, K.N., Lundsgaard, A.M., Hendrich, F.L., Nilas, L., Sjoberg, K.A.,
Hingst, J.R., Serup, A.K., Olguin, C.H., Carl, C.S., et al. (2020). Mechanisms Underlying Absent
Training-Induced Improvement in Insulin Action in Lean, Hyperandrogenic Women With
Polycystic Ovary Syndrome. Diabetes *69*, 2267-2280. 10.2337/db20-0062.

63. Agergaard, J., Trostrup, J., Uth, J., Iversen, J.V., Boesen, A., Andersen, J.L., Schjerling, P., and
Langberg, H. (2015). Does vitamin-D intake during resistance training improve the skeletal
muscle hypertrophic and strength response in young and elderly men? - a randomized
controlled trial. Nutr Metab (Lond) *12*, 32. 10.1186/s12986-015-0029-y.

64. Petersen, M.H., de Almeida, M.E., Wentorf, E.K., Jensen, K., Ortenblad, N., and Hojlund, K.
(2022). High-intensity interval training combining rowing and cycling efficiently improves
insulin sensitivity, body composition and VO(2)max in men with obesity and type 2 diabetes.
Front Endocrinol (Lausanne) *13*, 1032235. 10.3389/fendo.2022.1032235.

65. Hostrup, M., Lemminger, A.K., Thomsen, L.B., Schaufuss, A., Alsoe, T.L., Bergen, G.K., Bell,
A.B., Bangsbo, J., and Thomassen, M. (2023). High-Intensity Training Represses FXYD5 and
Glycosylates Na,K-ATPase in Type II Muscle Fibres, Which Are Linked with Improved Muscle
K(+) Handling and Performance. Int J Mol Sci *24*. 10.3390/ijms24065587.

66. Hey-Mogensen, M., Hojlund, K., Vind, B.F., Wang, L., Dela, F., Beck-Nielsen, H., Fernstrom,
M., and Sahlin, K. (2010). Effect of physical training on mitochondrial respiration and reactive
oxygen species release in skeletal muscle in patients with obesity and type 2 diabetes.
Diabetologia *53*, 1976-1985. 10.1007/s00125-010-1813-x.

67. Phielix, E., Meex, R., Moonen-Kornips, E., Hesselink, M.K., and Schrauwen, P. (2010). Exercise
training increases mitochondrial content and ex vivo mitochondrial function similarly in
patients with type 2 diabetes and in control individuals. Diabetologia *53*, 1714-1721.
10.1007/s00125-010-1764-2.

68. Karakelides, H., Irving, B.A., Short, K.R., O'Brien, P., and Nair, K.S. (2010). Age, obesity, and
sex effects on insulin sensitivity and skeletal muscle mitochondrial function. Diabetes *59*, 89-
97. 10.2337/db09-0591.

69. Schwanhausser, B., Busse, D., Li, N., Dittmar, G., Schuchhardt, J., Wolf, J., Chen, W., and
Selbach, M. (2011). Global quantification of mammalian gene expression control. Nature
473, 337-342. 10.1038/nature10098.

70. Gineste, C., Youhanna, S., Vorrink, S.U., Henriksson, S., Hernandez, A., Cheng, A.J., Chaillou,
T., Buttgereit, A., Schneidereit, D., Friedrich, O., et al. (2022). Enzymatically dissociated
muscle fibers display rapid dedifferentiation and impaired mitochondrial calcium control.
iScience *25*, 105654. 10.1016/j.isci.2022.105654.

71. Copeland, J.M., Cho, J., Lo, T., Jr., Hur, J.H., Bahadorani, S., Arabyan, T., Rabie, J., Soh, J., and
Walker, D.W. (2009). Extension of Drosophila life span by RNAi of the mitochondrial
respiratory chain. Curr Biol *19*, 1591-1598. 10.1016/j.cub.2009.08.016.

72. Dell'agnello, C., Leo, S., Agostino, A., Szabadkai, G., Tiveron, C., Zulian, A., Prelle, A.,
Roubertoux, P., Rizzuto, R., and Zeviani, M. (2007). Increased longevity and refractoriness to
Ca(2+)-dependent neurodegeneration in Surf1 knockout mice. Hum Mol Genet *16*, 431-444.
10.1093/hmg/ddl477.

73. Dillin, A., Hsu, A.L., Arantes-Oliveira, N., Lehrer-Graiwer, J., Hsin, H., Fraser, A.G., Kamath,
R.S., Ahringer, J., and Kenyon, C. (2002). Rates of behavior and aging specified by
mitochondrial function during development. Science *298*, 2398-2401.
10.1126/science.1077780.

74. Rossignol, R., Faustin, B., Rocher, C., Malgat, M., Mazat, J.P., and Letellier, T. (2003).
Mitochondrial threshold effects. Biochem J *370*, 751-762. 10.1042/BJ20021594.

75. Hwang, A.B., Jeong, D.E., and Lee, S.J. (2012). Mitochondria and organismal longevity. Curr
Genomics *13*, 519-532. 10.2174/138920212803251427.

76. Xu, C., Hwang, W., Jeong, D.E., Ryu, Y., Ha, C.M., Lee, S.V., Liu, L., and He, Z.M. (2018). Genetic inhibition of an ATP synthase subunit extends lifespan in C. elegans. Sci Rep *8*, 14836. 10.1038/s41598-018-32025-w.

77. Liu, X., Jiang, N., Hughes, B., Bigras, E., Shoubridge, E., and Hekimi, S. (2005). Evolutionary conservation of the clk-1-dependent mechanism of longevity: loss of mclk1 increases cellular fitness and lifespan in mice. Genes Dev *19*, 2424-2434. 10.1101/gad.1352905.

78. Leick, L., Wojtaszewski, J.F., Johansen, S.T., Kiilerich, K., Comes, G., Hellsten, Y., Hidalgo, J., and Pilegaard, H. (2008). PGC-1alpha is not mandatory for exercise- and training-induced adaptive gene responses in mouse skeletal muscle. Am J Physiol Endocrinol Metab *294*, E463-474. 10.1152/ajpendo.00666.2007.

79. Larsen, S., Skaaby, S., Helge, J.W., and Dela, F. (2014). Effects of exercise training on mitochondrial function in patients with type 2 diabetes. World J Diabetes *5*, 482-492. 10.4239/wjd.v5.i4.482.

80. Sylow, L., Tokarz, V.L., Richter, E.A., and Klip, A. (2021). The many actions of insulin in skeletal muscle, the paramount tissue determining glycemia. Cell Metab *33*, 758-780. 10.1016/j.cmet.2021.03.020.

81. Richter, E.A., Sylow, L., and Hargreaves, M. (2021). Interactions between insulin and exercise. Biochem J *478*, 3827-3846. 10.1042/BCJ20210185.

82. Cartee, G.D. (2021). Sexual Dimorphic Effects of Exercise Training on Subcutaneous White Adipose Tissue of Mice. Diabetes *70*, 1242-1243. 10.2337/dbi21-0014.

83. Holcomb, L.E., Rowe, P., O'Neill, C.C., DeWitt, E.A., and Kolwicz, S.C., Jr. (2022). Sex differences in endurance exercise capacity and skeletal muscle lipid metabolism in mice. Physiol Rep *10*, e15174. 10.14814/phy2.15174.

84. Rooyackers, O.E., Adey, D.B., Ades, P.A., and Nair, K.S. (1996). Effect of age on in vivo rates of mitochondrial protein synthesis in human skeletal muscle. Proc Natl Acad Sci U S A *93*, 15364-15369. 10.1073/pnas.93.26.15364.

85. Ganetzky, B., and Flanagan, J.R. (1978). On the relationship between senescence and age-related changes in two wild-type strains of Drosophila melanogaster. Exp Gerontol *13*, 189-196. 10.1016/0531-5565(78)90012-8.

86. Correia, J.C., Kelahmetoglu, Y., Jannig, P.R., Schweingruber, C., Shvaikovskaya, D., Zhengye, L., Cervenka, I., Khan, N., Stec, M., Oliveira, M., et al. (2021). Muscle-secreted neurturin couples myofiber oxidative metabolism and slow motor neuron identity. Cell Metab *33*, 2215-2230 e2218. 10.1016/j.cmet.2021.09.003.

87. Geng, T., Li, P., Okutsu, M., Yin, X., Kwek, J., Zhang, M., and Yan, Z. (2010). PGC-1alpha plays a functional role in exercise-induced mitochondrial biogenesis and angiogenesis but not fiber-type transformation in mouse skeletal muscle. Am J Physiol Cell Physiol *298*, C572-579. 10.1152/ajpcell.00481.2009.

88. Jeppesen, J., Maarbjerg, S.J., Jordy, A.B., Fritzen, A.M., Pehmoller, C., Sylow, L., Serup, A.K., Jessen, N., Thorsen, K., Prats, C., et al. (2013). LKB1 regulates lipid oxidation during exercise independently of AMPK. Diabetes *62*, 1490-1499. 10.2337/db12-1160.

89. Eisenberg, B.R. (1983). Quantitative ultrastructure of mammalian skeletal muscle. In Handbook of Physiology, Skeletal Muscle, L.D. Peachey, R.H. Adrian, and S.R. Geiger, eds. (American Physiological Society), pp. p. 73–112. doi:10.1002/cphy.cp100103.

90. Pesta, D., and Gnaiger, E. (2012). High-resolution respirometry: OXPHOS protocols for human cells and permeabilized fibers from small biopsies of human muscle. Methods Mol Biol *810*, 25-58. 10.1007/978-1-61779-382-0_3.

91. Dzamko, N., Schertzer, J.D., Ryall, J.G., Steel, R., Macaulay, S.L., Wee, S., Chen, Z.P., Michell, B.J., Oakhill, J.S., Watt, M.J., et al. (2008). AMPK-independent pathways regulate skeletal muscle fatty acid oxidation. J Physiol *586*, 5819-5831. 10.1113/jphysiol.2008.159814.

92. Hostrup, M., Lemminger, A.K., Stocks, B., Gonzalez-Franquesa, A., Larsen, J.K., Quesada, J.P., Thomassen, M., Weinert, B.T., Bangsbo, J., and Deshmukh, A.S. (2022). High-intensity

interval training remodels the proteome and acetylome of human skeletal muscle. Elife *11*.
10.7554/eLife.69802.
93. Kruse, R., Pedersen, A.J., Kristensen, J.M., Petersson, S.J., Wojtaszewski, J.F., and Hojlund, K.
(2017). Intact initiation of autophagy and mitochondrial fission by acute exercise in skeletal
muscle of patients with Type 2 diabetes. Clin Sci (Lond) *131*, 37-47. 10.1042/CS20160736.
94. Kruse, R., Petersson, S.J., Christensen, L.L., Kristensen, J.M., Sabaratnam, R., Ortenblad, N.,
Andersen, M., and Hojlund, K. (2020). Effect of long-term testosterone therapy on molecular
regulators of skeletal muscle mass and fibre-type distribution in aging men with subnormal
testosterone. Metabolism *112*, 154347. 10.1016/j.metabol.2020.154347.
95. Welinder, C., and Ekblad, L. (2011). Coomassie staining as loading control in Western blot
analysis. J Proteome Res *10*, 1416-1419. 10.1021/pr1011476.
96. Lord, S.J., Velle, K.B., Mullins, R.D., and Fritz-Laylin, L.K. (2020). SuperPlots: Communicating
reproducibility and variability in cell biology. J Cell Biol *219*. 10.1083/jcb.202001064.

1298

Fig.1

Fig. 2

A

♂Littermates (WT) or *Slirp* knockout (KO)

- **sedentary WT** (WT SED; 6♀, 6♂)
- **sedentary KO** (*Slirp* KO SED; 11♀, 6♂)
- **exercise-trained WT** (WT ET; 6♀, 7♂)
- **exercise-trained KO** (*Slirp* KO ET; 10♀, 7♂)

| 18 | 24 | 25 | 26 | 28 | Age (weeks) |

START/ MRI — 6 wk-post ET MRI — GTT — Exercise capacity — Tissue Dissection

B Average running distance

male   females

km day⁻¹

*

WT *Slirp* KO   WT *Slirp* KO

C Average food intake

ET, p<0.001

males   females

g day⁻¹

SED ET SED ET SED ET SED ET
WT *Slirp* KO   WT *Slirp* KO

D Fasting glucose ♂

Geno, p<0.05
ET, p<0.001

mM

SED ET SED ET
WT *Slirp* KO

E Glucose tolerance test ♂

Time x Geno ET, p<0.05

blood glucose (mM)

*

WT SED   WT ET
KO SED   KO ET

time (min)

F iAUC ♂

mmol x min

SED ET SED ET
WT *Slirp* KO

G Plasma insulin during GTT ♂

ET, p<0.01

ng mL⁻¹

0 min
20 min

SED ET SED ET
WT *Slirp* KO

H Fasting glucose ♀

mM

SED ET SED ET
WT *Slirp* KO

I Glucose tolerance test ♀

Time, p<0.001

blood glucose (mM)

WT SED   WT ET
KO SED   KO ET

time (min)

J iAUC ♀

mmol x min

SED ET SED ET
WT *Slirp* KO

K Plasma insulin during GTT ♀

ng mL⁻¹

0 min
20 min

SED ET SED ET
WT *Slirp* KO

Fig.3

Fig.4

Fig.5

Fig. 6

Supplementary Fig.2

HIIT in individuals with a lean body, obesity, and T2D

Supplementary Fig. 3