1 **From sequence to molecules: Feature sequence-based genome mining**

2 **uncovers the hidden diversity of bacterial siderophore pathways**

Shaohua Gu[1,2*], Yuanzhe Shao[2*], Karoline Rehm[3], Laurent Bigler[3], Di Zhang[1], Ruolin He[1],

Ruichen Xu[4], Jiqi Shao[1], Alexandre Jousset[5], Ville-Petri Friman[6], Xiaoying Bian[7], Zhong Wei[5#],

Rolf Kümmerli[8#], Zhiyuan Li[1,2#]

[1] Center for Quantitative Biology, Academy for Advanced Interdisciplinary Studies, Peking

University, Beijing, 100871, China

[2] Peking-Tsinghua Center for Life Sciences, Academy for Advanced Interdisciplinary Studies,

Peking University, Beijing, 100871, China

[3] University of Zurich, Department of Chemistry, Winterthurerstr. 190, 8057 Zurich, Switzerland

[4] School of Life Science, Shandong University, Qingdao, Shandong 266237, China

[5] Jiangsu Provincial Key Lab for Organic Solid Waste Utilization, Key lab of organic-based

fertilizers of China, Nanjing Agricultural University, Nanjing, P R China

[6] University of Helsinki, Department of Microbiology, 00014, Helsinki, Finland

[7] Helmholtz International Lab for Anti-infectives, State Key Laboratory of Microbial Technology,

Shandong University, Qingdao, Shandong 266237, China

[8] University of Zurich, Department of Quantitative Biomedicine, Winterthurerstr. 190, 8057

Zurich, Switzerland

[*] These authors contributed equally to this article.

[#]Corresponding authors (email: weizhong@njau.edu.cn; rolf.kuemmerli@uzh.ch;

zhiyuanli@pku.edu.cn)

Abstract (146 words count)

Microbial secondary metabolites are a rich source for pharmaceutical discoveries and play

crucial ecological functions. While tools exist to identify secondary metabolite clusters in

genomes, precise sequence-to-function mapping remains challenging because neither function

nor substrate specificity of biosynthesis enzymes can accurately be predicted. Here we

developed a knowledge-guided bioinformatic pipeline to solve these issues. We analyzed 1928

genomes of *Pseudomonas* bacteria and focused on iron-scavenging pyoverdines as model

metabolites. Our pipeline predicted 188 chemically different pyoverdines with nearly 100%

structural accuracy and the presence of 94 distinct receptor groups required for the uptake of

iron-loaded pyoverdines. Our pipeline unveils an enormous yet overlooked diversity of

siderophores (151 new structures) and receptors (91 new groups). Our approach, combining

feature sequence with phylogenetic approaches, is extendable to other metabolites and

microbial genera, and thus emerges as powerful tool to reconstruct bacterial secondary

metabolism pathways based on sequence data.

Introduction

Rapid advancements in sequencing technologies have revolutionized our view on microbial communities. While amplicon sequencing provides information on community composition and diversity, shotgun and whole genome sequencing allow us to reliably anticipate evolutionary and ecological relationships between microbes and to obtain functional information on communities. Computational models assessing the metabolic capacity of individual members, or an entire consortium, have become very popular and powerful[1-3]. The major focus of such modelling approaches is typically on the primary metabolism of bacteria, as genes involved in core metabolic pathways are highly conserved and can be identified with relative ease[2,4]. Conversely, analysis of the secondary metabolites has attracted less attention, even though they include compounds such as antibiotics, toxins, siderophores, biosurfactants, all known to have important implications for microbial community assembly[5,6] and to be important sources for pharmaceutical discoveries[7-9] .

There are multiple challenges that currently prevent a detailed unravelling of secondary metabolism of bacteria based on genome data[10-12]. First, most secondary metabolites are produced by pathways comprised of modular enzymes such as non-ribosomal peptide synthetases (NRPSs) or polyketide synthases (PKS)[13,14]. Locating complete biosynthesis clusters and identifying all enzyme-encoding genes is challenging from highly fragmented metagenomic sequences or draft genomes with a high number of contigs. Second, functional predictions for coding regions within a cluster rely on homologous comparisons with experimentally characterized genes. Such information is often restricted to a limited number of model organisms, meaning that only a small portion of the existing secondary metabolism

pathways is covered by current data bases. Finally, given the complex multi-modular

biosynthesis machineries, it is challenging to precisely predict the secondary metabolites

produced even with accurately annotated NRPS or PKS clusters. The main challenge is that a

large pool of non-proteinogenic amino acids is used as substrates and the specificity of an

enzyme's A domain, connecting these unusual amino acids, is often poorly understood[15]. As a

result, new computational methods are needed to accurately reconstruct bacterial secondary

metabolism from sequence data.

Here, we present a new bioinformatic pipeline that overcomes these challenges. We

specifically focus on a particular class of secondary metabolites (iron-scavenging siderophores)

as a case study to develop a bioinformatic workflow that predicts the chemical structure of the

produced metabolites with nearly 100% accuracy. Our pipeline is based on improved gene

annotation combined with a phylogeny- and feature sequence-based substrate prediction

techniques (Figure 1). In comparison with the currently available databases and bioinformatic

tools[16], the main advancement of our method is the more accurate prediction of synthesized

products based on NRPS clusters identified in genome data. In addition, we show that our

workflow is capable of correcting previous mis-annotations and is extendable to other

secondary metabolites (e.g., toxins, biosurfactants, virulence factors) and microbial genera.

Among siderophores, we focus on NRPS machineries that are responsible for the

biosynthesis of pyoverdines, a class of chemically diverse siderophores with high iron affinity,

produced by *Pseudomonas* bacteria[17,18]. While each *Pseudomonas* strain produces a single

type of pyoverdine, an enormous structural diversity has been described across strains and

species[19-22]. Pyoverdine types differ in their peptide backbone, meaning that the diversity

should be mirrored in NRPS enzyme diversity and their selectivity for the different amino acid

substrates[18]. Based on this knowledge, our pipeline entails the following steps (Figure 1): (i)

identification of the complete sequences of pyoverdine synthetase genes from fragmented draft

genomes, (ii) building the pyoverdine biosynthesis machinery *in silico* by extracting the feature

sequences for substrate specificity from motif-standardized NRPSs, and (iii) predicting the

precise chemical structure of pyoverdines followed by empirical verification.

**Figure 1 Scheme depicting our new genome mining pipeline to precisely predict the
biosynthesis, the molecular structure and the uptake machinery of pyoverdines, a family
of iron-scavenging siderophores produced by members of the *Pseudomonas* genus.** The
grey rounded outer rectangle represents a bacterial cell. The red and blue arrow-shaped boxes stand for
the synthetase and receptor genes for pyoverdines, respectively. Synthetase genes are transcribed and
translated to form the *n*-modular NRPS enzymes. These enzymes synthesize the peptide backbone of
pyoverdine through an assembly line using their repeating module units, with the A domain being
responsible for substrate selection and the E domain for chirality. The *n*-substrate siderophores are then
exported to the extracellular space for iron chelation. Membrane-embedded TonB-dependent receptors
recognize the ferri-siderophore complex and import it into the cell. Bold black text and black arrows

describe our multi-step computational methods developed to reconstruct the entire process from genome sequence data. First, the annotation pipeline was improved (from antiSMASH[23]) to extract the complete sequence of pyoverdine synthetase genes from draft genomes. Second, NRPSMotifFinder was used to define A- and E-domains and to determine the exact motif-intermotif structure of the pyoverdine assembly line. Third, intermotif regions most indicative of substrate specificity were used to develop a phylogeny-focused method for precise product prediction. Fourth, a sequence-region-based annotation method was combined with genome architecture features to identify the FpvA, receptors responsible for ferri-pyoverdine import.

An additional element of iron acquisition involves specific receptors that recognize and trigger the uptake of iron-loaded siderophores. Pyoverdine receptors are annotated as FpvA and it is known that receptor diversity matches pyoverdine diversity[22,24]. Moreover, FpvA belongs to the family of TonB-dependent receptors and a single *Pseudomonas* species often has multiple gene copies encoding different receptors. This poses an additional bioinformatic challenge: how to find the gene encoding the specific pyoverdine receptor among several potential receptor genes? To overcome this, we develop an algorithm that focuses on receptor sequence regions involved in pyoverdine recognition and translocation across the outer membrane with supervised learning methods that reliably locate *fpvA* genes in fragmented genomes based on these regions (Figure 1). Not only did our method unveil a substantial number of previously unrecognized pyoverdine receptors, but it also revealed that certain receptor groups are widely distributed among *Pseudomonas* strains., opening the possibility of pyoverdine sharing and exploitation across the strain and species boundaries. Altogether, our bioinformatic pipeline uses knowledge-guided insights empowered by supervised learning to construct a first systematic sequence-to-function mapping of a family of secondary metabolites (pyoverdine) and their corresponding receptors. Our analysis unveils a yet unrecognized extraordinary diversity of iron-scavenging machineries in pseudomonads.

Results

Section 1: Improved annotation pipeline reveals a vast reservoir of pyoverdine

synthetase genes

Figure 2 Improved annotation pipeline reveals a vast diversity of pyoverdine synthetase

genes. a. Improved annotation pipeline based on the raw annotation from antiSMASH[23]. **b.** The

annotation pipeline was applied to 9599 *Pseudomonas* genomes (94% draft genomes). Genomes could

be separated into three categories. Yellow: genomes without pyoverdine cluster. Green: genomes with a

complete pyoverdine cluster. Red: genomes with incomplete pyoverdine synthetase cluster. The red

category involved genomes with truly incomplete clusters (lacking Flu or Pep synthetic genes) or genomes

with likely truncated synthetic genes at the edge of contigs. **c.** Distributions of the sequence length (upper

panel) and the number of A domains (lower panel) across all the genomes with a complete synthetase

cluster. **d.** Workflow applied to separate the 9599 *Pseudomonas* genomes into the three categories

described in b and removing of redundant genomes with high phylogenic similarity and showing high

similarity in pyoverdine synthetases. Red star indicates the start of the workflow. **e.** Phylogenetic tree

depicting the relationship among the 1928 non-redundant *Pseudomonas* strains (1664 producers and 264

non-producers) based on the concatenated alignment of 400 single-copy conserved genes in their

genomes. The inner ring depicts the taxonomical classification including the four most prevalent species.

The outer ring shows the number of A domains present in the pyoverdine synthetase assembly line in

each strain.

The first step of our bioinformatic pipeline was to improve the annotation of pyoverdine

synthetase genes. The pyoverdine molecules is composed of a conserved fluorescent

chromophore (Flu) and a peptide chain (Pep), which are both synthesized by NRPS enzymes[25].

There exist already tools, such as antiSMASH that can find and annotate NRPS clusters in

microbial genomes[23]. However, antiSMASH (and other popular annotation platforms[26,27]) rely

on accurate gene predictions, which are typically problematic for fragmented genomes.

Consequently, while antiSMASH can recognize and annotate certain genes of an NRPS cluster,

the precise reconstruction of a complete NRPS assembly line often fails. This is particularly

problematic because most available genomes are drafts and any analysis suffers from the

unavoidable issue of incomplete or misannotation of gene fragments.

To overcome these issues, we developed an improved four-step annotation pipeline

starting with the raw annotation of the pyoverdine cluster obtained from antiSMASH[23] (Figure

2a). First, we implemented a NRPS Hidden Markov Model (HMM) to re-annotate and extract

the entire nucleotide sequence of the pyoverdine synthetase cluster[28], including the genes

missed by antiSMASH. For this step, the nucleotide sequences were converted into amino acid

sequences to avoid erroneous gene predictions typically associated with antiSMASH. Second,

we assembled the entire re-annotated pyoverdine coding region into a single sequence with

defined start (co-enzyme A ligase, CAL) and end (thioesterase, TE) markers. In most cases

(~90%), there are two separate biosynthetic gene clusters for Flu and Pep. Their ordering is

easy as Flu always precedes Pep. In the remaining cases (~10%), there is either one or three

biosynthetic gene clusters. No ordering is required for the former. For the latter, we put Flu first

and the cluster with TE domain at the end. Third, we used NRPSMotifFinder to identify the C,

A, T, E and TE motifs that determine the NRPS structure of pyoverdine[15]. Finally, we applied a

safety measure to ensure that the recovered NRPS assembly line is complete (contains both

Flu and Pep) and is not truncated, which can occur when a synthetase coding region is at the

edge of a contig. Consequently, we dismissed all pyoverdine biosynthesis clusters located

within 100 bp proximity to contigs' edges and either lacked Flu or Pep synthetic genes.

Next, we applied our improved pyoverdine biosynthesis annotation pipeline to 9599

Pseudomonas genomes (including 613 complete and 8986 draft genomes) retrieved from the

Pseudomonas Genome Database[29]. We found the pyoverdine biosynthesis machinery in 97%

of the genomes (Figure 2b), indicating that the machinery is ubiquitous in *Pseudomonas*.

However, since 94% of the analyzed genomes were in draft form, the pyoverdine biosynthesis

machinery was likely truncated (i.e., on the edge of the contig) in 63.4% (6087) of the genomes.

These genomes were excluded from further analysis. Around 3.1% of retained genomes (293)

with high assembly completeness were missing pyoverdine synthetic genes, indicating that

these *Pseudomonas* strains were not able to produce pyoverdine ('non-producers'). The rest

of the genomes (33.5%; 3219 genomes) were classified as 'producers' with complete

pyoverdine NRPS assembly lines that meet all our quality controls. For these 3219 genomes,

we used NRPSMotifFinder to find boundaries between the various biosynthesis domains and

to determine the amino acid length of the cluster and the number of A domains. The lengths of

pyoverdine synthetic clusters ranged between 7690 and 21333 amino acids, and the number

of A domains per synthetase ranged between 6 and 17, with a total of 35,281 A domains being

present across all strains (Figure 2c and Figure S1). Overall, our analysis pipeline unveiled a

vast diversity of pyoverdine synthetases that goes far beyond of what has previously been

described in the literature.

Finally, we utilized the PhyloPhlAn3 pipeline[30] to conduct a phylogenetic analysis based

on 400 conserved genes with the 293 non-producers and the 3219 producers. We first removed

redundant non-producers by retaining the most integrative genome among strains with high

phylogenic similarity (>95% identity). Then, we removed redundant producers by retaining the

most integrative genome among strains with high phylogenic and pyoverdine synthetase

similarity (Figure 2d, >95% identity). This data cleaning yielded a total of 1928 *Pseudomonas*

strains (403 complete and 1525 incomplete genomes), segregating into 1664 pyoverdine

producers and 264 non-producers. The phylogenetic tree revealed that all major *Pseudomonas*

species clades were present in our data set (Figure 2e). Moreover, the number of A domains

varied widely among species and even between strains within species. For example, the

number of *Pseudomonas aeruginosa* A domains ranges between 7 and 14. In summary, by

improving the synthetase annotation method, we successfully obtained 1664 highly reliable

pyoverdine synthetases (with a total of 18,292 A domains) and 264 non-producers.

Section 2: Phylogeny-focused substrate prediction for pyoverdine A domains

Our next goal was to precisely predict the molecular structure of the pyoverdines produced by the 1664 strains with complete synthetase gene clusters. The first essential step towards this goal was to reliably predict the substrate selectivity of all A domains in the NRPS assembly line. The A domain of each module selects for a single substrate among 22 proteinogenic and hundreds of non-proteinogenic amino acids[31,32]. Moreover, whenever an E domain exists downstream of an A domain, the chirality of the amino acid incorporated into the peptide chain gets modified from L to D. Thus, the modularity combined with the selectivity of A domains can promote an enormous diversity of pyoverdine molecule structures. To date, at least 73 pyoverdine structures have been reported (Supplementary_table1). Among those, we could identify 13 cases for which both structures (obtained from structural analysis experiments, we wanted to avoid any form of biases that bioinformatic pre-predictions could introduce to downstream analyses) and completely sequenced synthetase genes were available (Supplementary_table2). In order to make reliable molecule predictions, two challenges must be addressed: (i) the extraction of relevant information from A domain sequences for which the substrate is known, and (ii) the effective application of this information to predict specificity of A domain sequences for which the substrate is unknown.

To address the first challenge, we built our analysis on the NRPS assembly lines of the 13 known pyoverdines identified above to extract relevant information from the A domain sequences. From this dataset, we could identify 101 A domains that are linked to 13 experimentally confirmed amino acid substrates (Supplementary_table3). We next performed a multisequence alignment of the 101 A domains to determine (i) the "feature sequence",

defined as the most informative sequence for substrate selectivity and (ii) the "feature sequence

distance", which allows to reliably differentiate between two different substrates. To this end,

we tested three different A domain regions, three different sequence similarity measurements

and seven different clustering methods for their predictive power (Figure 3a). We found that the

full A domain sequence is not informative for substrate prediction (Figure 3b, left panel). Instead,

our analysis indicated that information-rich positions start with motif A4 and end before motif

A5, consistent with the known role of the A domain pocket in substrate selectivity[15]. Overall, the

sequence region from motifs A4 to A5 (termed "Amotif4-5"), in conjunction with Jukes-Cantor

distance and Ward linkage clustering, performed best in accurately distinguishing between

different substrates and maintaining homogeneity for identical substrates (Figure 3b). Utilizing

these parameters, we conducted a comprehensive analysis of all A domains and their

respective substrate information within the *Pseudomonas* NRPS biosynthetic gene cluster

available in the MIBiG database[33] (Supplementary_table4). To our delight, our method can find

and correct previously misclassified substrate information of A domains in the MIBiG database

(Figure S2 and Supplementary_table5). This demonstrates the effectiveness of performing

substrate predictions for the A domain based on feature sequences and the selected parameter

combination.

To address the second challenge, we developed a "phylogeny-focused method" to apply

the above-derived feature sequence distance approach to the 18,292 discovered A domains.

We realized that a direct construction of a phylogenetic tree including all 18,292 query A

domains and the 101 reference A domains would be computationally too demanding and

impossible to scale up. Furthermore, such an approach would result in phylogeny-interference

issues, where domains would cluster not only based on their substrate similarities but also

based on overall species relatedness[15]. To minimize the effect of phylogeny and speed up

calculations, we took each of the 18,292 query A domains and identified the two most similar A

domain clusters among our 101 reference A domains. We then compared the feature distance

between each query A domain and the two most similar reference A domains and assigned a

substrate to the query A domain using the following rules (Figure 3c). (1) If the feature sequence

distance is below the 0.7 threshold (corresponding to 50% identity) for only one of the two

reference A domains, then the substrate of the query A domain is matched to the substrate of

the more similar (lower distance) reference A domain. (2a) If the feature sequence distance is

below the 0.7 threshold for both reference A domains, then we considered the relative difference

of the query A domain towards the two reference A domains. If the relative difference is larger

than 0.2, the query A domain is matched to the substrate of the more similar reference A domain.

(2b) If the relative difference is smaller than 0.2, the substrate of the query A domain cannot

unambiguously be determined and is thus matched with both reference substrates. (3) If the

feature distance is above the 0.7 threshold (below 50% identity) for both reference A domains,

then the substrate of the query A domain is marked as "unknown". For most query A domains,

rule (1) could be applied (17880 cases), whereas rules (2) and (3) had to be used rarely (133

and 279 cases, respectively). We applied our methodology termed "phylogeny-focused method"

to all following substrate and pyoverdine structure predictions.

Section 3: Experimental validation of the annotation and prediction pipeline

To test whether our bioinformatic pipeline can reliably predict pyoverdine structures, we

sequenced the genomes of 20 pyoverdine-producing *Pseudomonas* strains previously isolated

from soil and water[34]. We applied our annotation and prediction pipeline to generate predicted

pyoverdine structures for all 20 strains harboring a total of 237 A domains. Notably, none of the

predicted structures matched any of the 13 reference pyoverdine structures, and 9 out of the

20 structures were predicted to be novel (not yet characterized) pyoverdines. We then used the

UHPLC-HR-MS/MS method developed in 2022[35] to elucidate the chemical structures of these

pyoverdines. Six out of the 20 structures had already been elucidated in our previous work[35],

while we here used the same method to elucidate the chemical structure of the remaining 14

pyoverdines (see Supplementary Materials for MS/MS profiles).

We found a near-perfect match (94.4%) between the predicted and the observed

pyoverdine chemical structures and were able to accurately assign amino acids in 151 out of

160 cases (Figure 3d). Our method demonstrated a substantial improvement compared to

commonly used approaches or platforms integrating several methods (Supplementary_table6),

such as   antiSMASH 7.0[16], which could only accurately assign 94 out of 160 amino acids

(accuracy: 58.8%). The nine non-matching cases in our analysis segregated into three groups.

In three cases (1.9%), our algorithm could not distinguish between the substrates Lysine and

Ornithine, as these two amino acids are highly similar both in terms of their chemical structures

and corresponding A domain sequences. This is the only sensitivity issue that is associated

with our approach. In four cases (2.5%), our technique assigned an "unknown" substrate to

amino acids that turned out to be valine, citrulline and histidine. Indeed, these three amino acids

were not yet present in our reference dataset and therefore no correct prediction was possible.

While citrulline has never been reported in pyoverdines before, valine has been found in a

pyoverdine structure but no sequence data from the corresponding strain was available. These

cases show that our analysis pipeline can be used to highlight new substrates. Once

experimentally verified, the new A domains and their substrates can expand the reference

dataset, allowing targeted improvement of our phylogeny-focused prediction technique. Finally,

there were only two cases (0.8%) that represented true mismatches between observed and

predicted amino acids. Altogether, our phylogeny-focused method is highly accurate in

predicting pyoverdine peptide structures and in identifying unknown substrates in

Pseudomonas.

To assess the extendibility of our pipeline, we chose *Burkholderiales* as a test case,

leveraging a manually curated dataset that we built up from the literature (see Methods for

detail). This dataset contains 203 A-domain sequences and corresponding substrate

information for 34 NRPS metabolites (e.g., toxins, biosurfactants, virulence factors) across 7

genera of *Burkholderiales* (Supplementary_table7). We further integrated experimentally

validated   biosynthetic   gene   cluster   data   from   *Burkholderiales*   in   MIBiG   3.0[33]

(Supplementary_table8) into our manual dataset leading to a total of 302 A-domains. We then

split the available data into a training set (124 A domains, Supplementary_table12) and a test

set (178 A domains, Supplementary_table9). As for the pyoverdines, we first trained our

algorithm on the training data set and then applied it to the test data set. We observed that our

pipeline consistently yields high prediction accuracy within *Burkholderiales* (83%), which is

considerably higher than the accuracy obtained with antiSMASH 7.0[16], (67%, Figure S3 and

Supplementary_table9). This outcome underscores the robustness of our developed prediction

algorithm, highlighting its potential extension to other NRPS secondary metabolites and

microbial genera.

Figure 3 Phylogeny-focused substrate prediction for pyoverdine synthetase assembly lines. a. Information from 101 reference A domains with known amino acid substrates were used to

develop an algorithm that predicts substrates from A domain sequence data with high accuracy. The challenge is to group the variable A domains into clusters that predict the same substrate (captured by the silhouette index). To find the most distinctive algorithm, we combined different feature sequences of A domains (Amotif) with different distance and linkage methods in our hierarchical clustering analyses. The best performing path is shown in pink. **b.** Heatmap showing the hierarchically clustered distances of the 101 reference A domains as a function of the feature sequence used. Color bar in the lower right shows the sequence distance. Left panel: complete A domain sequences. Middle panel: Amotif3-6 sequences. Right panel: Amotif4-5 sequences. The heatmaps show that hierarchical clustering, reliably associating sequence distances with substrates, worked best with the Amotif4-5. The experimentally validated substrates are shown on top of each heatmap and consist of 13 amino acids and 2 amino acid derivatives: Ser (Serine), Lys (Lysine), Thr (Threonine), Ala (Alanine), Gly (Glycine), Gln (Glutamine), Arg (Arginine), Asp (Aspartic acid), Glu (Glutamic acid), Tyr (Tyrosine), Orn (Ornithine), Rsc (succinic acid derivatives), Dab (2,4-diaminobutyric acid), OHAsp (Aspartic acid derivative), and OHOrn (a general term for three ornithine derivatives: FoOHOrn, AcOHOrn, cOHOrn). **c.** Phylogeny-focused substrate prediction pipeline for query A domains (grey circle) based on Amotif4-5 feature sequence comparisons. X1 and X2 represent the feature distance between the query A domain and two closest reference A domains (blue and green circles), respectively. Three rules are used, based on the feature sequence distances X1 and X2 and a threshold value of 0.7 (50% similarity), to make substrate predictions for the query A domain. There are three possible outcomes: unambiguous substrate prediction (blue or green squares), ambiguous substrate prediction (dual-colored squares), and no prediction ("unknown). **d.** Phylogenetic tree of 20 *Pseudomonas* strains and visualization of their predicted and actual pyoverdine structures to validate our phylogeny-focused substrate prediction pipeline. Strains marked in red font indicate cases with novel (not yet characterized) pyoverdines structures. 151 out of the 160 substrates (94.4%) were correctly predicted. The nine inconsistencies are boxed in blue (Lysine and Ornithine are indistinguishable), in dashed black (detection of "unknown" substrates), and in red (true mismatches). Note that our prediction pipeline (as any other pipeline) cannot distinguish between modified variants of the same amino acid, for example, we cannot distinguish the three derivatives of ornithine (FoOHOrn, AcOHOrn, cOHOrn).

Section 4: Application of the annotation and prediction pipelines to a full dataset

After successful validation, we applied our bioinformatic pipeline to the 1664 complete NRPS assembly lines annotated in our genome analysis (Figure 2). Across all assembly lines, we were able to predict the substrates of 17,880 A domains (97.75%) without ambiguity, whereas 133 A domains (0.73%) were associated with two different substrates, and 279 A domains (1.52%) predicted an unknown substrate (similar to the cases of valine and citrulline above). After considering the presence/absence of an E domain in each module, we derived the structures of 1664 pyoverdines according to our methods in section 2 (Figure 4).

Species

■ *P. aeruginosa*
■ *P. fluorescens*
■ *P. syringae*
■ *P. putida*
■ *P. sp*
■ *Others*

Substrate

■ OHOrn ■ Gln ■ Lys ■ OHAsp ■ Ala ■ Rsc ■ Orn ■ Tyr
■ Ser ■ Arg ■ Thr ■ Asp ■ Gly ■ Glu ■ Dab ■ Unknow

Figure 4 Predicted pyoverdine structural diversity based on our developed algorithm mapped onto the phylogenetic tree of all 1928 (non-redundant) *Pseudomonas* strains. In this figure, the strains are arranged based on the phylogenetic tree without showing the tree itself. The stacked boxes in the outermost circle show the predicted structure of pyoverdines, whereby each color represents a specific amino acid substrate. Strains without boxes represent non-producers (n = 264). Boxes with two colors indicate cases of ambiguous (dual) substrate prediction. The inner circle shows the taxonomic species classification following Figure 2e. Because the allocation of strains to species names is often imprecise, we divided the 1928 strains by their phylogenetic distance into 18 clades (color shadings in inner-most circle), out of which 13 contained more than one strain. Lines within the inner-most circle link strains from different clades that share the same pyoverdine structures, whereby line colors represent the shared unique pyoverdines. The bending of the lines represents the phylogenetic sequence distances of the connected strain pairs.

Our prediction yielded 188 different pyoverdine molecules, out of which only 37 structures

had been previously reported. While these 37 reported structures were highly abundant across

strains (1103 out of 1664), our pipeline was powerful in identifying many of the rarer pyoverdine

variants. Agreeing with previous studies, we observed that the fluorophore is highly conserved

among the 188 predicted structures. Moreover, our analysis found that 13 amino acids form the

main structural substrates of all the 188 pyoverdine structures, with most of the variation being

attributable to different substrate combinations, peptide lengths, and substrate chirality (Figure

4).

Notably, pyoverdine structural diversity was not strongly linked to phylogeny because the

same pyoverdine structure could be found in completely unrelated species, while closely related

species often had different pyoverdine structures (Figure 4). These observations suggest that

there may be both frequent recombination and horizontal gene transfer of pyoverdine

synthetase clusters between species. Taken together, the bioinformatics methods developed in

our study can predict a suit of secondary metabolites (pyoverdines) from sequence data with

high accuracy, revealing an unprecedented richness and evolutionary history of siderophores

within pseudomonads and the discovery of 151 novel pyoverdine candidate variants.

Section 5: Development of a region-based identification method for annotation of the

FpvA receptors

In pseudomonads, iron-loaded pyoverdines are recognized by FpvA, a TonB-dependent

receptor, that transports the ferri-pyoverdine into the periplasm[19,36-38]. The protein structure of

characterized FpvA variants consists of three domains: The Secretin and TonB N-terminus short

domain (STN), the Plug domain (Plug), and the TonB-dependent receptor domain (TonB)[19].

While these domains are conserved across FpvA variants and other siderophore receptors,

there is substantial variation at the sequence level. This makes it challenging to reliably identify

FpvA receptors from sequence data by homologous search. As an example, we were unable

to find FpvA genes (with a 60% identity threshold) by homologous search in several genomes

although they had complete pyoverdine biosynthesis machineries. Moreover, there are many

other TonB-dependent receptors with fairly high sequence identity to FpvA but that transport

other siderophores than pyoverdine (e.g. FpvB, 55% identity, transporting pyoverdine,

ferrichrome and ferrioxamine B[39]). Therefore, it is imperative to develop a new comprehensive

method for identifying FpvA receptors in *Pseudomonas* genomes.

We started our approach by comparing the sequences of 35 reported siderophore

receptors, including 21 FpvA and 6 FpvB from The Pseudomonas Genome Database

(https://www.pseudomonas.com/), and 8 TonB-dependent siderophore receptor sequences

often found in *Pseudomonas* genomes from Uniprot (https://www.uniprot.org/), encoding

receptors for the uptake of heterologous siderophores (Supplementary_table10). We found that

all receptor sequences share a similar length of around 800 amino acids (FpvA and FpvB

sequences: 809 ± 10 amino acids). We then used the complete sequences to calculate the

pair-wise distances by global alignment before applying hierarchical clustering (Figure 5a). We

found substantial divergence between FpvA variants to an extent that was comparable to the

distance between FpvA and other siderophore receptors. Moreover, FpvB variants clustered

with FpvA variants, showing that FpvA identification based on full sequence distances is

unachievable. We hence focused on the three typical receptor domains (TonB, Plug, and STN,

retrieved from the Pfam database) and applied Profile Hidden Markov Models (pHMM) to each

reference sequence to calculate the pHMM probability scores for each domain. The probability

scores (calculated as the log-odd ratios for emission probabilities and log probabilities for state

transitions) had reasonably high scores but no distinction was apparent between the three

receptor classes (Figure 5b).

Figure 5 A sequence-region-based identification pipeline for annotating FpvA receptors.
a. Heatmap displaying the hierarchically clustered sequence distances (p-distance calculation method, identity (%) = (1-sequence distance) * 100) of 35 reference siderophore receptors identified in *Pseudomonas* spp., based on full sequences. No clear discrimination between FpvA, FpvB and other receptors is possible. The order of receptors is consistent across panels (**b**), (**e**), and (**f**). **b.** The pHMM scores of the three standard receptor domains (STN, Plug, and TonBDR) vary across the 35 reference sequences (A: FpvA, B: FpvB and NA: others), but do not allow to distinguish between receptor groups. **c.** FpvA region-based conservation scores from a multi-alignment of the 35 reference sequences mapped to the FpvA sequence of strain *P. aeruginosa* PAO1. All residues within the top 10% of the conservation score are denoted with black dots. For each region flanked by two black dots, we calculated the FpvA identification score (heatmap), representing the ability to distinguish FpvA from non-FpvA receptors. **d.** Mapping of the two regions with the highest FpvA identification scores R1(dark red) and R2 (orange) to the crystal structure of FpvA from PAO1 conjugated with pyoverdine (PDB 2IAH). **e.** Heatmap showing the hierarchically clustered sequence distances of 35 reference siderophore receptors based on the R1 sequence region. A clear discrimination between FpvA/FpvB and other receptors emerges. **f.** Heatmap showing the hierarchically clustered sequence distances of 35 reference siderophore receptors based on the R2 sequence region. A clear discrimination between FpvA and FpvB receptors emerges. **g.** The pHMM scores of regions R1 and R2 for the 35 siderophore reference receptors are contrasted against each other, yielding a clear separation between FpvA, FpvB and other receptors. Dashed lines indicate the pHMM threshold scores used for later analysis. **f.** Flowchart showing all steps involved in the FpvA annotation from genome sequence data. The red star indicates the start of the workflow.

We next asked whether there are specific regions within the receptor sequences that are characteristic of FpvA. To address this, we conducted a multiple sequence alignment (MSA) with all 35 reference receptor sequences and mapped them onto the sequence of the well-characterized FpvA of *P. aeruginosa* PAO1 (Figure 5c). MSA allows to identify conserved sites (Figure 5c, black dots representing the top 10% most conserved sites) that are shared by the majority of the reference sequences. We then used these conserved sites to partition the MSA into variable regions which were flanked by two conserved sites. For each variable region, we assessed its predictive power to differentiate FpvA from non-FpvA sequences. For this we defined the "FpvA identification score" analogous to the intercluster-vs-intracluster Calinski-Harabasz variance ratio, as

$$I_{\mathrm{FpvA}} = d_{\mathrm{A:non}}/d_{\mathrm{A:A}}$$

where $d_{A:A}$ is the sequence distance among all 21 FpvA sequences, and $d_{A:non}$ is the sequence

distance between all 21 FpvA and the 14 non-FpvA sequences.

Our analysis yielded two locations with noticeably high FpvA identification scores (Figure

5c). The region with the highest FpvA identification score (referred to as R1) locates at the

intersection of the Plug domain and the barrel structure of the TonB domain (Figure 5d, between

258 Gly and 309 Gly in the PAO1 FpvA). According to the sequence distance matrix, the R1

region allows to distinguish heterologous siderophore receptors from FpvA and FpvB receptors

(Figure 5e). The region with the second highest FpvA identification score (referred to as R2)

was located in the C-terminal signaling domain (Figure 5d, between 59 Leu and 86 Lys in the

PAO1 FpvA). The sequence distance matrix revealed that R2 allows to distinguish FpvB from

FpvA receptors (Figure 5f).

We then defined two informative pHMM scores based on (i) the alignment of the 21 FpvA

sequences in the R1 region, termed R1(FpvA), and (ii) the alignment of the 6 FpvB sequences

in the R2 region, termed R2(FpvB). Running R1(FpvA) and R2(FpvB) against all 35 reference

sequences revealed a clear separation between the three receptor categories (Figure 5g).

Along the R1(FpvA) axis, FpvA and FpvB reference sequences have high R1 scores (minimal

score 77.0) that separate them from other siderophore receptors (maximal score 38.1),

whereas FpvAs references have substantially lower R2 scores (maximal 20.2) than FpvBs

(minimal 49.0) along the R2(FpvB) axis.

Based on these insights, we developed a decision flow chart for annotating FpvAs in

Pseudomonas genomes (Figure 5h). First, we considered sequences as Fpv-like receptors that

share similar properties to the ones identified in our reference database. Particularly, protein

coding sequence (CDS) length had to be between 750 and 850 amino acids and the phMM

scores for the three typical receptor domains STN, Plug, and TonB must be greater than 25, 50,

and 80, respectively (Figure 5b, red dashed lines). Second, we used the pHMM threshold

scores obtained for R1(FpvA) and R2(FpvB) (Figure 5g) to differentiate other siderophore

receptors (R1(FpvA) score < 50) from FpvB receptors (R1(FpvA) score > 50 and R2(FpvB)

score > 30) and FpvA receptors (R1(FpvA) score > 50 and R2(FpvB) score < 30). Our method

effectively identifies FpvA receptors from sequence data and can be readily applied to the entire

Pseudomonas dataset.

Section 6: Application of the receptor annotation pipeline to the full dataset.

The region-based receptor identification pipeline was applied to all 1928 *Pseudomonas*

genomes. The analysis identified 4547 FpvAs, 615 FpvBs, and 9139 other TonB-dependent

Fpv-like receptors across the dataset (Figure 6a). The 4547 FpvA sequences clustered

hierarchically into 114 groups, defined by an identity threshold of 60%. When comparing to the

21 reference FpvAs (Figure 6b), we found that 2293 FpvA sequences have close homologues

in the reference data base, while 2254 FpvA sequences lack such close homologues (sequence

identity < 50%). These latter sequences, termed as "dissimilar to reference", may represent

novel subtypes of FpvA receptors that could not be found by simple homology search. Our

analysis further shows that many strains have more than one FpvA receptor.

We then asked whether the 4547 FpvAs are found in proximity of pyoverdine Pep

synthetase genes as it is commonly the case for cognate FpvA receptors [40]. We thus calculated

the proximity between pyoverdine Pep genes and the Fpv-like receptor genes by counting the

number of base pairs between the two coding regions. All TonB-dependent receptors that have

not been classified as FpvAs were more than 20 kb away from the Pep genes (Figure 6c). In

contrast, 92% of the nearest FpvA genes were indeed located within 20 kb of their pyoverdine

Pep genes (Figure 6d, called proximate receptors). These proximate receptors encompassed

both those with close (66%) and more distant (34%) resemblance to the reference receptor

types. Overall, this proximity analysis confirmed that our region-based gene identification

method can reliably identify FpvA receptors.

Next, we explored the diversity among FpvA receptors in greater detail by focusing on the

1,534 strains with proximate receptors located within 20 kb of the pyoverdine Pep genes (Figure

6d). We utilized these high-confidence FpvAs for sequence feature extraction. When

considering the whole gene sequences, these receptors segregated into 44 groups according

to single-linkage clustering with an identity threshold of 60% (Figure S4a). To investigate which

sequence regions were the most informative for reliable clustering, we used a similar approach

as with FpvAs detection by quantifying the "group identification score" for variable regions

flanked by highly conserved sites. The higher the score, the stronger a region's capacity to

discriminate between FpvA groups. We found that the four regions with the top discrimination

capacities all located near the Plug domain surrounding the pyoverdine transmission channel

(Figure 6e). The plug domain is known to undergo conformational changes and is involved in

pyoverdine selectivity and import[41,42], suggesting that the four high-score regions are

responsible for pyoverdine specificity.

Figure 6 Application of the receptor annotation pipeline to the full database. a. Applying
the receptor annotation pipeline to the genomes of the 1928 non-redundant *Pseudomonas* strains yields
14301 Fpv-like receptors, which segregate into 4547 FpvA receptors (red box), 615 FpvB receptors, and
9139 other receptors, based on the pHMM score thresholds for regions R1 and R2. The heatmap indicates
receptor density. **b.** Sequence distance matrix between the 35 reference sequences (y-axis) and the 4547

annotated FpvA sequences in the full database (x-axis). Database sequences were ordered by hierarchically clustering and segregated into 114 groups. 2254 of the annotated FpvA sequences have low sequence identity < 60% compared to the reference receptors, pointing at many novel subtypes of FpvA receptors. **c.** Genomic distance (in base pairs) between each Fpv-like receptor sequence and its pyoverdine peptide synthetase gene (Pep) for annotated FpvA receptors (upper panel), FpvB receptors (middle panel) and other receptors (lower panel). **d.** Distribution of the genomic distance between each FpvA receptor and its nearest pyoverdine peptide synthetase depending on whether the annotated FpvA receptor has high sequency similarity (blue, ≥ 50%) or low sequence similarity (yellow, < 50%) with at least one of the 21 reference FpvAs. **e.** FpvA region-based conservation scores from a multi-alignment of all the annotated FpvA receptors that are proximate (< 20 kbp) to the pyoverdine synthetase cluster mapped to the FpvA sequence of strain *P. aeruginosa* PAO1. All residues within the top 10% of the conservation score are denoted with black dots. For each region flanked by two black dots, we calculated the group identification score (heatmap, lower panel), representing the ability of the region to distinguish between different groups of FpvA receptors. Four regions in the plug domains had a particularly high group identification score (called the feature sequence). They are mapped to the crystal structure of FpvA from PAO1 conjugated with pyoverdine (PDB 2IAH, upper panel). All four regions surround the pyoverdine transmission channel and are shown in the respective heatmap color. **f.** Heatmap showing the hierarchically clustered distances between the 4547 annotated FpvA receptors based on the feature sequence (comprising the four groups with the highest identification scores). The analysis identifies 94 receptor groups with a 70% identity threshold. **e.** The diversity of FpvA receptors along the 13 phylogeny clades containing more than 1 strain. Receptor diversity was calculated by the Shannon entropy, similar to the alpha-diversity in microbial community.

Based on the above insights, we concatenated the four high-score regions (from 168 Pro to 295 Ala in PAO1) into a single "feature sequence". The feature sequence could characterize 98% of the distance matrix compared to the whole sequence (1534 FpvAs, r=0.98, Figure S4a-b) and substantially reduced within-group distance. We applied the concatenated feature sequence approach to all the 4547 annotated FpvAs to calculate the sequence distance matrix. Single-linkage clustering with an identity threshold of 70% revealed a total of 94 groups, out of which 43 groups contained more than 10 members (Figure 6f). The diversity of receptors is hence much larger than currently anticipated as only 3 groups of FpvAs have previously been reported. Finally, we calculated the diversity of receptor FpvAs for each of the 13 phylogenetic clades with more than one strain by the Shannon entropy, which is similar to the alpha-diversity

in microbial community (Figure 6g). We noticed large differences in FpvA diversity across the

clades and species: clades with *P. aeruginosa* and *P. syringae* species had lower FpvAs

diversity (1.55 and 1.60) than clades containing *P. putida* and *P. fluorescens* species (4.82 and

3.77). Taken together, the region-based identification method developed in our study can

reliably mine pyoverdine receptors from genome data, revealing an undiscovered FpvA

diversity that is unequally distributed across the different phylogenetic clades of pseudomonads.

We further extended the search for FpvA receptors, using our pipeline in Figure 5h, to

the >30,000 complete bacterial genomes in NCBI. We found 31,936 receptors to pass the R1

and R2 thresholds (Figure S5), and surprisingly found that FpvA receptors seem to be widely

dispersed across 12,944 strains spanning 13 phyla, 22 classes, 54 orders, 139 families, and

468 genera, and 2,598 species (Figure S6 and Supplementary_table11). While *Pseudomonas*

remains the genus with the highest distribution of FpvAs, other genera with notable distributions

comprise *Escherichia*, *Salmonella*, *Enterobacter*, *Acinetobacter*, *Xanthomonas*, *Bordetella*, and

Achromobacter. While confirmatory experimental work is still pending, these analyses suggest

that FpvA receptors for pyoverdine uptake may be widely distributed throughout the bacterial

kingdom.

Discussion

The rapid expansion of sequencing data offers exciting opportunities for microbiology[43-45]. One

key challenge of current research in the field is to infer biological functions of microbial

communities from genome sequence data[46-48]. While this endeavor is increasingly successful

for biological functions involving the primary metabolism and the associated complex metabolic

flux, reconstructing aspects of the secondary metabolism is much more challenging. The main

issue is that neither the function of a secondary metabolite enzyme nor the resulting metabolite

can be precisely predicted from gene sequence data. In our study, we tackled this challenge

and developed a bioinformatic pipeline to reconstruct the complete secondary metabolism

pathway of pyoverdines, a class of iron-scavenging siderophores produced by *Pseudomonas*

spp. These secondary metabolites are synthetized by a series of non-ribosomal peptide

synthetases and require a specific receptor (FpvA) for uptake. We combined knowledge-guided

learning with phylogeny-based methods to predict with high accuracy: (i) the full pyoverdine

assembly line, (ii) the substrate specificity for each enzyme within the assembly lines, (iii) the

complete chemical structure of pyoverdines, and (iv) the FpvA receptors from genome

sequences. After validation, we tested our pipeline with sequence data from 1664

phylogenetically distinct *Pseudomonas* strains and were able to determine 18,292 enzymatic A

domains involved in pyoverdine biosynthesis, to reliably predict 97.8% of their substrates, and

to identify 188 different pyoverdine molecule structures and 4547 FpvA receptor variants

belonging to 94 distinct groups. The uncovered diversity is stunning and goes far beyond

currently known levels of variation (73 pyoverdines and 3 FpvA groups).

We show that knowledge-guided learning is an extremely powerful tool to predict enzyme,

metabolite, and receptor properties. The establishment of our entire pipeline is based on only

101 previously known enzymatic A domains (from 13 known pyoverdine assembly lines) and

21 FpvA receptor sequences. Even with this limited amount of information, we were able to

predict the substrates of almost all the 18,292 enzymatic A domains and to identify 4547 FpvA

receptors from the sequence data. A key insight from our knowledge-guided learning is that

comparisons based on the full gene sequences (e.g., for pyoverdine synthetase or receptor)

are typically non-informative and unsuitable for obtaining functional information. This is because

overall diversity does not stand for functional diversity, meaning that A domains recognizing the

same substrate can diverge substantially in their full sequences. The same holds true for

receptor sequences: whole-sequence alignments can neither accurately identify FpvA

receptors nor reliably separate them into functional groups. Instead, it is imperative to extract

informative feature sequences that are defined as sequence stretches within a gene whose

diversity is tightly linked to variation in its functioning. We successfully extracted and applied

feature sequence comparisons for both A domain substrate prediction and FpvA identification.

It is important to note that a knowledge-guided pipeline does not have to be perfect right from

the start. For example, our pipeline for pyoverdine structure prediction returned unknowns for

several amino acid positions within the PEP. Our experimental verifications then revealed new

substrates such as valine and citrulline. This information can then be used to refine our

prediction algorithm in a feedback loop.

Another main advantage of our bioinformatic pipeline is that it can be applied to draft

genomes. This reflects a major improvement compared to existing annotation tools such as

antiSMASH[23], which typically has difficulties in recognizing NRPS structures in fragmented

genome assemblies. However, draft genomes are the most common data source in

microbiology. While our pipeline shows high performance, we need to acknowledge that we still

lose many genomes (6087 out of 9599, 63.4%). The reason for the loss is that the pyoverdine

biosynthesis machinery is large, which increases the probability that it is positioned at the end

of a contig. We decided to exclude those cases because the annotated biosynthesis machinery

might be truncated and thus incomplete. Thus, the high loss rate of draft genomes is rather due

to limitations in sequence quality (too many short contigs) and not due to a limitation of our

bioinformatic pipeline. We believe that this limitation may disappear in the future as long-read

sequencing technologies are quickly becoming cheaper and more reliable.

We further show that knowledge-guided learning combined with a phylogeny-focused

approach is a powerful tool for predicting the substrate specificity of A domains of synthetases.

It outperforms currently known bioinformatics prediction tools of NRPS substrates such as

antiSMASH[23]. Most current algorithms[49-53] perform poorly when applied to pyoverdines,

particularly when encountering non-proteogenic amino acids. The high accuracy of our

algorithm can largely be attributed to our reference set, composing only 13 pyoverdines from

Pseudomonas spp., yet capturing most of the substrate diversity. Similarly accurate

predictions based on a handful of known substrates among closely related species were

observed in several fungal NRPS systems[54]. It is worth noting that when the algorithm output

is "unknown," it signifies the presence of uncharacterized A domains that are not yet

incorporated into the reference data set. This should prompt researchers to pay attention to

these A domains, and like in our case, subject them to further experimental investigation. This

approach helped us discover new substrates (valine, citrulline) and link them to the

corresponding A domains. The novel substrates identified through our structural assessment

and mass spectrometry experiment can subsequently be used to enhance the precision of our

phylogeny-centered substrate prediction technique in the future, creating a progressive

feedback loop of expanding knowledge. Taken together, supervised learning based on a few

known compounds produced by species from the same genus can outperform generalized

prediction algorithms trained on many products from a diverse set of microbes for NRPS

substrate predictions.

Our results show that both pyoverdine and receptor diversity has been vastly

underestimated. While considerable pyoverdine diversity (n=73) has already been captured in

previous studies, here we predicted 151 putatively new variants. On the receptor side, the

uncovered novel diversity is more dramatic. One reason for this is that research on receptors

has mainly focused on the pathogen *P. aeruginosa*[19-22,55]. For this species, three different

pyoverdine types were described[21] together with three structurally different FpvA receptor types

that each recognize one of the pyoverdine types[22]. While our study confirmed that *P. aeruginosa*

strains (n = 554) indeed have only 3 pyoverdine-receptor types, we further discovered 91 new

FpvA groups among environmental *Pseudomonas* spp. Our findings raise the question why

there are so many different pyoverdine and receptor variants. One potential explanation is that

the benefit of specific siderophores could be context-dependent and locally adapted to the

multitude of different environmental conditions pseudomonads are exposed to. For example,

experimental work has revealed that pyoverdines can be cooperatively shared among strains

with matching receptors[37,56], or conversely, pyoverdines can serve as competitive agents by

locking away iron from species that have non-matching receptors[57]. Given that bioavailable iron

is limited in most natural and host-associated habitats[58-60], the unraveled functional diversity is

likely a direct evolutionary consequence of the struggle and competition of microbes for iron.

While experimental work is often restricted to a low number of strains, we propose that our

bioinformatic pipeline can be used to predict pyoverdine-mediated interaction networks across

thousands of strains and across different habitats. We have addressed this point in a

complementary study[61].

We are confident that our pipeline can be expanded to study iron competition in multi-

species communities and perhaps also in plant-microbe ecosystems, as siderophores exist

ubiquitously and are shared among microbes and certain hosts[62]. To move further, a key

question is whether our knowledge-guided approach can be applied to other microbes and

important secondary metabolites, such as antibiotics, toxins, biosurfactants and pigments? Our

preliminary investigation, focused on the curated dataset in *Burkholderiales* as a test case,

provides a highly promising response to this query. The outcomes from this evaluation

predominantly showcased the method's potential for extension to diverse microbial genera and

metabolites, given sufficiently accurate dataset. Meanwhile, we observed a notable decline in

accuracy (from 94% to 83%) when compared to pyoverdine. This indicates that our algorithm

possibly needs refinement, particularly concerning specific secondary metabolites or microbial

genera. As soon as sufficient case-by-case knowledge on a specific system is available, the

annotation strategies together with the feature sequence extraction and the phylogeny-focused

approach developed in our paper can be applied. It is worth noting that our current algorithm

does not predict tailoring reactions. While tailoring reactions are crucial for predicting the final

NRPS product structure, this challenge is not addressed by any existing pipelines and remains

an area for future research. In our study, it is important to note that the specificity of pyoverdines

is mainly determined by the backbone composition, with tailoring reactions playing a relatively

minor role[17]. For most of the secondary metabolites listed above, there are no receptors as the

compounds have purely extra-cellular functions, which substantially simplifies the development

of bioinformatic pipelines. In the long run, it will certainly be possible to automate the steps

implemented in our workflow so that the algorithms can be applied to a large set of secondary

metabolites when fed with an appropriate training set.

Method

Construction of phylogeny tree

The phylogenetic tree depicted in Figure 1e was constructed utilizing the PhyloPhlAn3

pipeline[30]. PhyloPhlAn is a comprehensive pipeline that encompasses the identification of

phylogenetic markers, multiple sequence alignment, and the inference of phylogenetic trees. In

this analysis, we employed 400 universal genes defined by PhyloPhlAn as our selected

phylogeny markers. Subsequently, the taxonomic cladogram was generated using the iTOL

web tool (http://huttenhower.sph.harvard.edu/galaxy/).

Application of our pipeline to identify errors in the substrate information associated with

the A domain of *Pseudomonas* in the MIBiG database

We downloaded all A domain and substrate data for all NRPS secondary metabolism instances

for *Pseudonomas* within MIBiG 3.0[33](Supplementary_table4). Following our pipeline, we

initiated the process by utilizing NRPSMotifFinder to extract all Amotif4-5 sequences of the A

domains. Subsequently, we calculated the Jukes-Cantor sequence distance between Amotif4.5

pairs. Finally, employing Ward linkage clustering, we conducted a cluster analysis on all

Amotif4.5 sequences, resulting in the generation of Figure S2. We observed that within the

same cluster, substrates often exhibit some subtle impurities. We speculate that such clusters

might contain some errors in substrate information. Consequently, we focused on these

potential errors, manually searched for the respective original literature, and verified that the A

domain and its corresponding substrate information entered in MIBiG indeed contain errors

(Supplementary_table5).

Application of our pipeline to predict the structural composition of diverse NRPS

metabolites across various genera of *Burkholderiales*

To evaluate the extendibility of our bioinformatic pipeline to various metabolites and microbial

genera, we utilized a previously curated dataset of A domain sequences with experimentally

confirmed substrates for *Burkholderiales* [63] (Supplementary_table7). This manually curated

dataset, encompassing 7 genera, 34 secondary metabolite products, 203 A domains, and 25

substrates within *Burkholderiales*, served as a valuable resource and as a test set to validate

the extendibility of our bioinformatic approach. Simultaneously, we compiled a training set by

aggregating A domain and substrate data from all NRPS secondary metabolism instances

within *Burkholderiales*, utilizing experimentally validated biosynthetic gene cluster data from

MIBiG 3.0[33] (Supplementary_table8). This initial training set covered 6 genera, 19 secondary

products, 99 A domains, and 21 substrates. To ensure the inclusivity of all substrates from the

test set, we randomly selected an A domain from each substrate in the manually curated dataset,

incorporating it into our training set. Consequently, the finalized training set comprised 124 A

domains (Supplementary_table12), while the test set encompassed 178 A domains

(Supplementary_table9). Then, we concurrently employed our pipeline and antiSMASH[16] to

predict the corresponding substrate based on the sequence data of the A domain in the test set.

Subsequently, we compared these predictions with the actual substrate information, calculating

the accuracy rate for the two algorithms.

735

736 **Application of our pipeline to annotate FpvA receptors across the bacterial domain**

737 To explore the distribution of FpvA receptors across the bacterial domain, we acquired all

738 complete bacterial genomes (33,207) from NCBI (https://www.ncbi.nlm.nih.gov). Employing the

739 pipeline depicted in Figure 5h to examine the CDS of these genomes, we identified a total of

740 357,790 TonB-dependent receptors, of which 31,936 and 623 matched the criteria for being

741 considered as FpvA and FpvB receptors, respectively. Subsequently, leveraging the taxonomy

742 information from NCBI, we conducted a comparative analysis of the distribution differences

743 between FpvA and FpvB among various species. The findings suggest that FpvA exhibit a

744 widespread distribution across different species, whereas FpvB was exclusive to the genus

745 *Pseudomonas*, with a maximum occurrence of one copy per genome.

746
747

748 **Data Availability**

749 The source code and parameters used are available in the supplementary material. The

750 sequences of the 20 Pseudomonas strains are available on ENA accession PRJEB76792.

751 One-click online notebook for the annotation and substrate prediction can be found here:

752 https://drive.google.com/drive/folders/1JsfyPUGDTFo8BDDZk8JLSvKry8emzMhr?usp=drive_

753 link , following a more detail explanation on the code.

754

755 **Acknowledgements**

756 We thank Richard Allen and Simon Maréchal for the handling of the genome sequencing of the

20 *Pseudomonas* strains used for the structure validation experiment. We further thank Vera

Vollenweider for sample preparation for the experiment elucidating pyoverdine structures.

Funding

This work was supported by the National Key Research and Development Program of China

(No. 2021YFF1200500, 2021YFA0910700), National Natural Science Foundation of China (No.

42107140, No. 32071255, No.41922053, No. T2321001), National Postdoctoral Program for

Innovative Talents (No. BX2021012). R.K. is supported by a grant from the Swiss National

Science Foundation no. 310030_212266. V-P.F. is supported jointly by a grant from UKRI, Defra,

and the Scottish Government, under the Strategic Priorities Fund Plant Bacterial Diseases

program (BB/T010606/1), Research Council of Finland, and The Finnish Research Impact

Foundation.

Author contributions

Shaohua Gu performed the majority of computational analysis in this research and drafted the

manuscript. Yuanzhe Shao built the pipeline for retrieving synthetase sequence information and

developed the phylogeny-centered method for predicting substrates. Karoline Rehm and

Laurent Bigler performed the pyoverdine structure elucidation experiments. Di Zhang depicted

the circular plot. Ruolin He standardized all NRPS structures to the motif-intermotif format. Jiqi

Shao and Ruichen Xu assisted in revising the manuscript. Xiaoying Bian provided curated

dataset of A domain sequences with experimentally confirmed substrates from *Burkholderiales*.

Xiaoying Bian, Alexandre Jousset and Ville-Petri Friman offered insightful comments and

assisted in revising and writing of the manuscript. Rolf Kümmerli and Zhong Wei oversaw the

project, designed experiments and revised the manuscript. Zhiyuan Li conceptualized and

oversaw the project, conducted the analysis of the receptor annotated method, and revised the

manuscript.

Competing interests

The authors declare no competing interests.

References

1   Zengler, K. & Palsson, B. O. A road map for the development of community systems

789   (CoSy) biology. *Nature Reviews Microbiology* **10**, 366-372, doi:10.1038/nrmicro2763

790   (2012).

791   2   Gu, C., Kim, G. B., Kim, W. J., Kim, H. U. & Lee, S. Y. Current status and applications

792   of genome-scale metabolic models. *Genome Biology* **20**, 121, doi:10.1186/s13059-

793   019-1730-3 (2019).

794   3   García-Jiménez, B., Torres-Bacete, J. & Nogales, J. Metabolic modelling approaches

795   for describing and engineering microbial communities. *Computational and Structural*

796   *Biotechnology Journal* **19**, 226-246, doi:10.1016/j.csbj.2020.12.003 (2021).

797   4   Colarusso, A. V., Goodchild-Michelman, I., Rayle, M. & Zomorrodi, A. R. Computational

798   modeling of metabolism in microbial communities on a genome-scale. *Current Opinion*

799   *in Systems Biology* **26**, 46-57, doi:10.1016/j.coisb.2021.04.001 (2021).

800 5  Scherlach, K. & Hertweck, C. Mediators of mutualistic microbe–microbe interactions.

801   *Natural Product Reports* **35**, 303-308, doi:10.1039/C7NP00035A (2018).

802 6  Trivedi, P., Leach, J. E., Tringe, S. G., Sa, T. & Singh, B. K. Plant–microbiome

803   interactions: from community assembly to plant health. *Nature Reviews Microbiology*

804   **18**, 607-621, doi:10.1038/s41579-020-0412-1 (2020).

805 7  Price-Whelan, A., Dietrich, L. E. P. & Newman, D. K. Rethinking 'secondary' metabolism:

806   physiological roles for phenazine antibiotics. *Nature Chemical Biology* **2**, 71-78,

807   doi:10.1038/nchembio764 (2006).

808 8  Thirumurugan, D., Cholarajan, A., Raja, S. & Vijayakumar, R. An introductory chapter:

809   secondary metabolites. (2018).

810 9  Chaturvedi, K. S., Hung, C. S., Crowley, J. R., Stapleton, A. E. & Henderson, J. P. The

811   siderophore yersiniabactin binds copper to protect pathogens during infection. *Nature*

812   *Chemical Biology* **8**, 731-736, doi:10.1038/nchembio.1020 (2012).

813 10  Penn, K. *et al.* Genomic islands link secondary metabolism to functional adaptation in

814   marine Actinobacteria. *The ISME Journal* **3**, 1193-1203, doi:10.1038/ismej.2009.58

815   (2009).

816 11  Yee, D. A. *et al.* Genome mining for unknown–unknown natural products. *Nature*

817   *Chemical Biology* **19**, 633-640, doi:10.1038/s41589-022-01246-6 (2023).

818 12  Lautru, S., Deeth, R. J., Bailey, L. M. & Challis, G. L. Discovery of a new peptide natural

819   product by Streptomyces coelicolor genome mining. *Nature Chemical Biology* **1**, 265-

820   269, doi:10.1038/nchembio731 (2005).

13   Andryukov, B., Mikhailov, V. & Besednova, N. The Biotechnological Potential of

Secondary Metabolites from Marine Bacteria. *Journal of Marine Science and*

Engineering **7** (2019).

14   Kautsar, S. A., Blin, K., Shaw, S., Weber, T. & Medema, M. H. BiG-FAM: the biosynthetic

gene cluster families database. *Nucleic Acids Research* **49**, D490-D497,

doi:10.1093/nar/gkaa812 (2021).

15   He, R. *et al.* Knowledge-guided data mining on the standardized architecture of NRPS:

Subtypes, novel motifs, and sequence entanglements. *PLOS Computational Biology*

19, e1011100, doi:10.1371/journal.pcbi.1011100 (2023).

16   Blin, K. *et al.* antiSMASH 7.0: new and improved predictions for detection, regulation,

chemical structures and visualisation. *Nucleic Acids Research* **51**, W46-W50,

doi:10.1093/nar/gkad344 (2023).

17   Ringel, M. T. & Brüser, T. The biosynthesis of pyoverdines. *Microb Cell* **5**, 424-437,

doi:10.15698/mic2018.10.649 (2018).

18   Hopkinson, B. M. & Morel, F. M. M. The role of siderophores in iron acquisition by

photosynthetic marine microorganisms. *BioMetals* **22**, 659-669, doi:10.1007/s10534-

009-9235-2 (2009).

19   Cobessi, D. *et al.* The Crystal Structure of the Pyoverdine Outer Membrane Receptor

FpvA from Pseudomonas aeruginosa at 3.6Å Resolution. *Journal of Molecular Biology*

347, 121-134, doi:10.1016/j.jmb.2005.01.021 (2005).

20   Diggle, S. P. & Whiteley, M. Microbe Profile: Pseudomonas aeruginosa: opportunistic

pathogen and lab rat. *Microbiology (Reading)* **166**, 30-33, doi:10.1099/mic.0.000860

(2020).

21   Meyer, J.-M. *et al.* Use of Siderophores to Type Pseudomonads: The Three

845   Pseudomonas Aeruginosa Pyoverdine Systems. *Microbiology* **143**, 35-43,

846   doi:10.1099/00221287-143-1-35 (1997).

847   22   Bodilis, J. *et al.* Distribution and evolution of ferripyoverdine receptors in Pseudomonas

848   aeruginosa. *Environmental Microbiology* **11**, 2123-2135, doi:10.1111/j.1462-

849   2920.2009.01932.x (2009).

850   23   Blin, K. *et al.* antiSMASH 5.0: updates to the secondary metabolite genome mining

851   pipeline. *Nucleic Acids Research* **47**, W81-W87, doi:10.1093/nar/gkz310 (2019).

852   24   Kümmerli, R. Iron acquisition strategies in pseudomonads: mechanisms, ecology, and

853   evolution. *BioMetals*, doi:10.1007/s10534-022-00480-8 (2022).

854   25   Meyer, J. M. Pyoverdines: pigments, siderophores and potential taxonomic markers of

855   fluorescent Pseudomonas species. *Arch Microbiol* **174**, 135-142,

856   doi:10.1007/s002030000188 (2000).

857   26   Xu, Z., Park, T. J. & Cao, H. Advances in mining and expressing microbial biosynthetic

858   gene clusters. *Crit Rev Microbiol*, 1-20, doi:10.1080/1040841x.2022.2036099 (2022).

859   27   Keller, N. P. Fungal secondary metabolism: regulation, function and drug discovery.

860   *Nature Reviews Microbiology* **17**, 167-180, doi:10.1038/s41579-018-0121-1 (2019).

861   28   Bateman, A. *et al.* The Pfam protein families database. *Nucleic Acids Research* **32**,

862   D138-D141, doi:10.1093/nar/gkh121 (2004).

863   29   Winsor, G. L. *et al.* Enhanced annotations and features for comparing thousands of

864   Pseudomonas genomes in the Pseudomonas genome database. *Nucleic Acids*

Research **44**, D646-D653, doi:10.1093/nar/gkv1227 (2016).

30   Asnicar, F. *et al.* Precise phylogenetic analysis of microbial isolates and genomes from

metagenomes using PhyloPhlAn 3.0. *Nature Communications* **11**, 2500,

doi:10.1038/s41467-020-16366-7 (2020).

31   Felnagle, E. A. *et al.* Nonribosomal peptide synthetases involved in the production of

medically relevant natural products. *Mol Pharm* **5**, 191-211, doi:10.1021/mp700137g

(2008).

32   Süssmuth, R. D. & Mainz, A. Nonribosomal Peptide Synthesis—Principles and

Prospects. *Angewandte Chemie International Edition* **56**, 3770-3821,

doi:10.1002/anie.201609079 (2017).

33   Terlouw, B. R. *et al.* MIBiG 3.0: a community-driven effort to annotate experimentally

validated biosynthetic gene clusters. *Nucleic Acids Research* **51**, D603-D610,

doi:10.1093/nar/gkac1049 (2023).

34   Butaitė, E., Kramer, J., Wyder, S. & Kümmerli, R. Environmental determinants of

pyoverdine production, exploitation and competition in natural Pseudomonas

communities. *Environmental Microbiology* **20**, 3629-3642, doi:10.1111/1462-

2920.14355 (2018).

35   Rehm, K., Vollenweider, V., Kümmerli, R. & Bigler, L. A comprehensive method to

elucidate pyoverdines produced by fluorescent Pseudomonas spp. by UHPLC-HR-

MS/MS. *Analytical and Bioanalytical Chemistry* **414**, 2671-2685, doi:10.1007/s00216-

022-03907-w (2022).

36   Butaitė, E., Baumgartner, M., Wyder, S. & Kümmerli, R. Siderophore cheating and

cheating resistance shape competition for iron in soil and freshwater Pseudomonas

communities. *Nature Communications* **8**, 414, doi:10.1038/s41467-017-00509-4

(2017).

37   Kramer, J., Özkaya, Ö. & Kümmerli, R. Bacterial siderophores in community and host

interactions. *Nature Reviews Microbiology* **18**, 152-163, doi:10.1038/s41579-019-

0284-4 (2020).

38   Jin, Z. *et al.* Conditional privatization of a public siderophore enables Pseudomonas

aeruginosa to resist cheater invasion. *Nature Communications* **9**, 1383,

doi:10.1038/s41467-018-03791-y (2018).

39   Chan, D. C. K. & Burrows, L. L. Pseudomonas aeruginosa FpvB

is a high-affinity transporter for xenosiderophores ferrichrome and ferrioxamine B.

bioRxiv, 2022.2009.2020.508722, doi:10.1101/2022.09.20.508722 (2022).

40   González, J. *et al.* Loss of a pyoverdine secondary receptor in Pseudomonas

aeruginosa results in a fitter strain suitable for population invasion. *The ISME Journal*

15, 1330-1343, doi:10.1038/s41396-020-00853-2 (2021).

41   Schalk, I. J., Mislin, G. L. & Brillet, K. Structure, function and binding selectivity and

stereoselectivity of siderophore-iron outer membrane transporters. *Curr Top Membr* **69**,

37-66, doi:10.1016/B978-0-12-394390-3.00002-1 (2012).

42   Greenwald, J. *et al.* FpvA bound to non-cognate pyoverdines: molecular basis of

siderophore recognition by an iron transporter. *Molecular Microbiology* **72**, 1246-1259,

doi:10.1111/j.1365-2958.2009.06721.x (2009).

43   Almeida, A. *et al.* A new genomic blueprint of the human gut microbiota. *Nature* **568**,

499-504, doi:10.1038/s41586-019-0965-1 (2019).

44 Schloss, P. D. & Handelsman, J. Metagenomics for studying unculturable

microorganisms: cutting the Gordian knot. *Genome Biology* **6**, 229, doi:10.1186/gb-

2005-6-8-229 (2005).

45 Handelsman, J. Metagenomics: Application of Genomics to Uncultured

Microorganisms. *Microbiology and Molecular Biology Reviews* **68**, 669-685,

doi:10.1128/MMBR.68.4.669-685.2004 (2004).

46 Tsilimigras, M. C. B. & Fodor, A. A. Compositional data analysis of the microbiome:

fundamentals, tools, and challenges. *Annals of Epidemiology* **26**, 330-335,

doi:10.1016/j.annepidem.2016.03.002 (2016).

47 Weiss, S. *et al.* Correlation detection strategies in microbial data sets vary widely in

sensitivity and precision. *ISME J* **10**, 1669-1681, doi:10.1038/ismej.2015.235 (2016).

48 Faust, K. & Raes, J. Microbial interactions: from networks to models. *Nature Reviews*

Microbiology **10**, 538-550, doi:10.1038/nrmicro2832 (2012).

49 Khayatt, B. I., Overmars, L., Siezen, R. J. & Francke, C. Classification of the

Adenylation and Acyl-Transferase Activity of NRPS and PKS Systems Using

Ensembles of Substrate Specific Hidden Markov Models. *PLOS ONE* **8**, e62136,

doi:10.1371/journal.pone.0062136 (2013).

50 Minowa, Y., Araki, M. & Kanehisa, M. Comprehensive Analysis of Distinctive Polyketide

and Nonribosomal Peptide Structural Motifs Encoded in Microbial Genomes. *Journal*

of Molecular Biology **368**, 1500-1517, doi:10.1016/j.jmb.2007.02.099 (2007).

51 Prieto, C., García-Estrada, C., Lorenzana, D. & Martín, J. F. NRPSsp: non-ribosomal

peptide synthase substrate predictor. *Bioinformatics* **28**, 426-427,

doi:10.1093/bioinformatics/btr659 (2012).

52 Röttig, M. *et al.* NRPSpredictor2—a web server for predicting NRPS adenylation

domain specificity. *Nucleic Acids Research* **39**, W362-W367, doi:10.1093/nar/gkr323

(2011).

53 Zierep, P. F., Ceci, A. T., Dobrusin, I., Rockwell-Kollmann, S. C. & Günther, S. SeMPI

2.0-A Web Server for PKS and NRPS Predictions Combined with Metabolite Screening

in Natural Product Databases. *Metabolites* **11**, doi:10.3390/metabo11010013 (2020).

54 Fan, J. *et al.* Biosynthetic diversification of peptaibol mediates fungus-mycohost

interactions. *bioRxiv* (2022).

55 Smith, E. E., Sims, E. H., Spencer, D. H., Kaul, R. & Olson, M. V. Evidence for

diversifying selection at the pyoverdine locus of Pseudomonas aeruginosa. *J Bacteriol*

187, 2138-2147, doi:10.1128/jb.187.6.2138-2147.2005 (2005).

56 Gu, S. *et al.* Competition for iron drives phytopathogen control by natural rhizosphere

microbiomes. *Nature Microbiology* **5**, 1002-1010, doi:10.1038/s41564-020-0719-8

(2020).

57 Figueiredo, A. R. T., Özkaya, Ö., Kümmerli, R. & Kramer, J. Siderophores drive invasion

dynamics in bacterial communities through their dual role as public good versus public

bad. *Ecology Letters* **25**, 138-150, doi:10.1111/ele.13912 (2022).

58 Andrews, S. C., Robinson, A. K. & Rodríguez-Quiñones, F. Bacterial iron homeostasis.

FEMS Microbiology Reviews **27**, 215–237, doi:10.1016/s0168-6445(03)00055-x

(2003).

59   Boyd, P. W. & Ellwood, M. J. The biogeochemical cycle of iron in the ocean. *Nat Geosci*

3, 675-682, doi:10.1038/ngeo964 (2010).

60   Emerson, D., Roden, E. & Twining, B. The microbial ferrous wheel: iron cycling in

terrestrial, freshwater, and marine environments. *Frontiers in Microbiology* **3**,

doi:10.3389/fmicb.2012.00383 (2012).

61   Gu, S. *et al.* Siderophore synthetase-receptor gene coevolution reveals habitat- and

pathogen-specific bacterial iron interaction networks. *bioRxiv*, 2023.2011.2005.565711,

doi:10.1101/2023.11.05.565711 (2024).

62   Ruolin, H. *et al.* SIDERITE: Unveiling Hidden Siderophore Diversity in the Chemical

Space Through Digital Exploration. *bioRxiv*, 2023.2008.2031.555687,

doi:10.1101/2023.08.31.555687 (2023).

63   Chen, H. *et al.* Biosynthesis and engineering of the nonribosomal peptides with a C-

terminal putrescine. *Nature Communications* **14**, 6619, doi:10.1038/s41467-023-

42387-z (2023).

    Supplementary figures

    Figure S1 Clustering heatmap of sequence distance matrix between A domains from 1664
    strains and 101 reference A domains. The color on top indicates the substrate of reference A
    domains.

Figure S2 Clustering heatmap of sequence distance matrix between 318 Amotif4-5 from *Pseudomonas* strains in the MIBiG database (Supplementary_table4). MIBiG is a standard annotation database of biosynthetic gene clusters. We retrieved all secondary metabolites produced by members of the *Pseudomonas* genus from MIBiG. The colors on top of the heatmap indicate the substrate of the A domains. We focus on the domains with closed sequence distance but with distinct substrate specificity (marked with red text). After checking the original reference, theses domains were labeled incorrectly in MIBiG (Supplementary_table5).

Figure S3 Comparison of prediction accuracy in the structural composition of secondary metabolites between our approach and antiSMASH in *Burkholderiales*. Each data point in the violin plot symbolizes a distinct secondary metabolite. The median is denoted by a red circle, while the mean is represented by a blue '×'.

988
989
990
991

Figure S4 Comparison of clustering and grouping effects of complete sequences (a) and feature sequences (b) of all proximate-receptors.

994
995

Figure S5 Applying the receptor annotation pipeline to >30,000 complete bacterial genomes in NCBI based on the pHMM score thresholds for regions R1 and R2. The heatmap indicates receptor density.

998
999
1000

1001

1002

1003 **Figure S6** The distribution of 31,936 FpvAs spans various taxonomic categories. Taxa with an
1004 abundance exceeding 1/500 are highlighted with colors, while others remain white. Taxa
1005 surpassing an abundance of 1/50 will have their specific names marked.

1006

1007