Full Title

Interactive visualization of whole eukaryote genome alignments using NCBI's Comparative

Genome Viewer (CGV)

Short Title

Visualization of genome alignment in NCBI CGV

Authors

Sanjida H Rangwala*, Dmitry V Rudnev, Victor V Ananiev, Andrea Asztalos, Barrett Benica,

Evgeny A Borodin, Nathan Bouk, Vladislav I Evgeniev, Vamsi K Kodali, Vadim Lotov, Eyal Mozes,

Dong-Ha Oh, Marina V Omelchenko, Sofya Savkina, Ekaterina Sukharnikov, Joël Virothaisakun,

Terence D. Murphy, Kim D Pruitt, Valerie A. Schneider

National Center for Biotechnology Information, National Library of Medicine, National Institutes

of Health (NIH), Bethesda, MD 20894, USA.

* Corresponding author

E-mail: rangwala@nih.gov

Abstract

We report a new visualization tool for analysis of whole genome assembly-assembly

alignments, the Comparative Genome Viewer (CGV) (https://ncbi.nlm.nih.gov/genome/cgv/).

CGV visualizes pairwise same-species and cross-species alignments provided by NCBI using

assembly alignment algorithms developed by us and others. Researchers can examine the

alignments between the two assemblies using two alternate views: a chromosome ideogram-

based view or a 2D genome dotplot. Whole genome alignment views expose large structural

differences spanning chromosomes, such as inversions or translocations. Users can also

navigate to regions of interest, where they can detect and analyze smaller-scale deletions and

rearrangements within specific chromosome or gene regions. RefSeq or user-provided gene

annotation is displayed in the ideogram view where available. CGV currently provides

approximately 700 alignments from over 300 animal, plant, and fungal species. CGV and related

NCBI viewers are undergoing active development to further meet needs of the research

community in comparative genome visualization.

Introduction

Comparative genome visualization

Comparative genomics leverages shared evolutionary histories among different species to answer basic biological questions and understand the causes of disease. As sequencing costs have dropped and assembly algorithms have improved, there has been tremendous growth in the number of high-quality genome assemblies available in public archives, and the diversity in the organisms they represent. These data now make it possible to use comparative genomics approaches to explore more elements of biology and reveal the need for different types of analysis tools to support this exploration. The NIH Comparative Genomics Resource (CGR) maximizes the impact of eukaryotic research organisms and their genomic data to biomedical research (1). CGR facilitates reliable comparative genomics analyses for all eukaryotic organisms through community collaboration and a National Center for Biotechnology Information (NCBI) genomics toolkit. As part of CGR, we have created the Comparative Genome Viewer (CGV), a web-based visualization tool to facilitate comparative genomics research.

Graphical visualization of genomic data can illuminate relationships among different data types and highlight differences and anomalies; for instance, areas of a genome that are depleted in gene annotation, or have unusually high repeat content, or are more variable between species. Interactive genome browsers have become particularly valuable in recent years in helping biologists navigate large sequence datasets and more easily find genomic locations of interest to their specific research interest question. These visualizations can display molecular data that

can help resolve competing hypotheses and also expose patterns that can spur additional

research questions.

Linear genome browsers, such as the Genome Data Viewer (GDV) at NCBI (2), the UCSC

Genomics Browser (3), and JBrowse (4), display molecular data as "tracks" laid out in parallel

and anchored on a single genome assembly. Users can navigate to different chromosome

regions and view gene annotation, repeats, and other types of data. Displays of known

sequence variants (e.g., dbVar or dbSNP data) can aid in the analysis of gene or sequence level

variation in a discrete genome region. While these browsers can be powerful tools to analyze

many different types of data from a single genome concurrently, they are more limited in their

ability to support comparative genome analysis. In particular, linear genome browsers cannot

easily show whether a genome region that appears evolutionarily conserved has been

rearranged (e.g., inverted or translocated) in one genome relative to another one, since the

data is only displayed relative to a single genomic region in a single genome at a time.

Different types of two-dimensional visualizations have been proposed to facilitate analysis of

larger scale genome structural differences between two or more genomes. These visuals

include two-dimensional line graphs (also known as dotplots) (5, 6), circular diagrams (i.e.,

Circos plots) (7), and linear genome browsers that stack one assembly on top of another (8-10).

Different types of visuals have advantages and disadvantages. Circos plots can show multiple

datasets in one graphic but can be visually challenging to interpret and usually do not support

views of sub-genomic regions. Dotplots can allow zooming to view chromosome or sub-

chromosome regions but cannot easily or elegantly display gene or other annotation in the

same visual. In order to better serve different research questions, some groups provide a choice

of multiple different types of visuals for genome comparisons (11-13).

Genome comparison data

Broadly, there are two types of whole genome comparison data that can be displayed in a

comparative genome visualization tool. The first type of data is locations of gene orthologs.

Orthology is typically determined using a protein homology-based method (e.g., BLASTP) in

consideration with local gene order conservation (14, 15). This type of data can lend itself to

straightforward "beads on string" visualizations that allows researchers to easily determine

how syntenic gene regions have evolved across related species (15-17).

The second type of comparison data is whole genome assembly alignments (e.g., Mauve (18),

LASTZ (19)), which are sequence-based and include both genic and intergenic regions. Whole

genome alignments can be much more complex than simple gene ortholog locations but have

the advantage of including alignments in regulatory regions and other regions not annotated as

genes.

Here we introduce a new viewer tool at NCBI, the Comparative Genome Viewer (CGV), that is a

key element of CGR. The main view of CGV takes the "stacked linear browser" approach —

chromosomes from two assemblies are laid out horizontally with colored bands connecting

regions of sequence alignment. Initial usability research with conceptual prototypes revealed

that this type of visual was the easiest to interpret for scientists from a broad range of research

expertise in genomics. We display whole genome pairwise assembly-assembly alignments in

CGV. These sequence-based alignments can be used to analyze gene synteny conservation but

can also expose similarities in regions outside known genes e.g., ultraconserved regions that

may be involved in gene regulation. Because CGV is a web-based application, researchers do

not have to install or configure software or generate their own comparison files before they can

begin using it for their research. Below we describe some of the features of CGV and provide

examples of how visualization in this tool can generate insights into genome structure and

evolution.

Results

Overview of CGV

We developed a web application, the Comparative Genome Viewer (CGV)

(https://ncbi.nlm.nih.gov/genome/cgv/), to aid in comparing genome structures between two

eukaryotic assemblies. CGV facilitates analyses of genome variation and evolution between

different strains or species or strains, as well as evaluation of assembly quality between older

and newer assemblies from same species.

Alignments are generated at NCBI using BLAST (20) or LASTZ-based algorithms (19), or imported

from the UCSC Genomics Institute (https://hgdownload.soe.ucsc.edu/downloads.html) or from

other research groups (e.g., HPRC, https://humanpangenome.org/). Shorter alignments are

merged where possible; however, because of repeats and gaps in the alignments, even very

similar chromosomes may be broken down into many alignment segments. Refer to **Materials**

and Methods for more detail on how we generate whole-genome assembly alignments and

load them into the viewer.

The CGV home page provides a menu where users can select from available species and

assembly combinations (Fig 1A). We frequently add new alignments as high-profile assemblies

become available and in response to requests from the scientific community.   As of October

2023, we provided a selection of almost 700 alignments from over 300 eukaryotic species (Fig

1B).

Fig 1. Overview of CGV.

(A) CGV selection menu. (B) Taxonomic distribution of species represented by alignments in CGV.

(C) CGV ideogram view of whole genome assembly alignment. Buttons in the lower right provide download access

for complete alignment data or an SVG image of the current alignment view.

https://www.ncbi.nlm.nih.gov/genome/cgv/browse/GCF_015227675.2/GCF_000001635.27/27835/10116 (D) CGV

zoomed to a chromosome-by-chromosome view, with information box shown. i. flip orientation ii. zoom in/out iii.

pan left/right iv. view assembly in Genome Data Viewer (GDV) v. information panel.

https://www.ncbi.nlm.nih.gov/genome/cgv/browse/GCF_015227675.2/GCF_000001635.27/27835/10116#NC_05

1343.1:44657546-50471623/NC_000075.7:41365809-52285446/size=10000

(E) CGV search interface and sample search results. (F) Filtering and configure options for ideogram view.

CGV's main view (the "ideogram view") displays pairwise alignments as colored connectors

linking the chromosomes in the two assemblies (Fig 1C). The view is filtered by default to show

only reciprocal best hits between assemblies to facilitate the analysis of orthologous genomic

regions. The researcher can choose to show the non-best placed alignments to expose close

sequence duplications or homologues (Fig 1F). Users of CGV can also filter alignments in view

by size (e.g., to only show large alignments blocks) or by orientation (e.g., to only show regions

that have undergone a potential inversion). The complete alignment data in GFF3 and human-

readable formats like XLSX can be downloaded from the viewer for a researcher's own use.

Users can select a chromosome from each assembly to zoom to the alignments for this

chromosome. They can navigate further within this chromosome comparison using the zoom

in/out and pan buttons or by using the mouse to pinch-zoom or drag to pan. Users can zoom

directly to a particular region of a chromosome by dragging their cursor over the coordinate

ruler or the ideogram for either assembly. Double-clicking on a selected alignment segment will

synchronously zoom both the top and bottom assembly on the aligned coordinates, so that

they are stacked on top of one another (Fig 1D).

Where available, RefSeq or assembly-submitter provided gene annotation is displayed on the

chromosomes (Fig 1D). Similarities in gene order denote regions of synteny, while discrepancies

can point to evolutionarily or biologically significant differences, or assembly errors, particularly

if evaluating different assemblies from the same species. Researchers can use the search

feature in CGV to find their gene of interest by name or keyword, and subsequently navigate to

the location of the gene in the viewer (Fig 1E). If the gene region is aligned, the viewer will

simultaneously navigate to the aligned location, which may contain the gene's known or

putative ortholog on the second assembly. The "flip" button allows the user to reverse one

chromosome to see inverted alignments displayed in the same relative orientation, which may

aid in the detection of discrepancies in gene annotation in regions that are locally syntenic

between the two assemblies. Once a user has completed their analysis of a region of interest,

they can export the image as an SVG to adapt for use in publications and presentations.

We provide additional information for each alignment segment in a pop-over panel (Fig 1D).

This panel reports the chromosome scaffold accession and sequence coordinates of the

alignment on each assembly, as well as the percent identity, number of gaps and mismatches,

and alignment length. While the ideogram view in CGV does not display specific nucleotide

bases, users can open another panel from the right-click menu that shows the alignment

sequence. They can also download the alignment FASTA file of a particular alignment segment

for downstream analysis, such as BLAST search or primer design. Moreover, researchers can

also navigate from CGV to NCBI's genome browser, the Genome Data Viewer (GDV) (2). GDV

can display the assembly-alignment data viewed in CGV as a linear track alongside additional

data mapped onto a genome assembly, such as detailed transcript and CDS annotation,

repeats, GC content, variation data, or user-provided annotations. Zooming to a location within

GDV can reveal granular differences in nucleotide sequence or gene exon or CDS annotation

between the two assemblies.

In addition to the main ideogram-based view, the Comparative Genome Viewer also provides a

two-dimensional dotplot view of the pairwise genome alignment (Fig 2A). The dotplot shows

aligned sequence locations in one assembly on the X-axis plotted against aligned locations on

the second assembly on the Y-axis. Alignments in the reverse orientation are plotted with an

opposite slope and in a different color (purple) than alignments in the same orientation (green),

making it easier to identify inversions and inverted translocations. The CGV dotplot shows both

reciprocal best-placed and non-best placed alignments. As a result, compared to the ideogram

view, this plot may more easily expose differences in copy number between two assemblies,

such as segmental duplications or differences in genome or chromosome ploidy. Users can

select a pair of chromosomes in the whole genome plot (i.e., a "cell" in the plot) and zoom into

them on a full screen, where smaller alignment segments may be more easily interpretable (Fig

2B). Once a researcher has discovered a chromosome pair of interest, they can navigate back to

the ideogram view to conduct even more fine-grained analysis, including examining gene

annotation and observing very short alignment segments that were beyond the resolution of

the dotplot.

Fig 2. CGV dotplot view of *Xenopus laevis* and *Xenopus tropicalis* alignment.

https://ncbi.nlm.nih.gov/genome/cgv/plot/GCF_000004195.4/GCF_017654675.1/38475/8355

(A) Full genome dotplot. (B) Dotplot of chromosome 8 of *Xenopus tropicalis* vs chromosome 8S of *Xenopus laevis*.

Analysis using CGV: Conservation of linkage groups with local rearrangement of synteny

CGV can aid in detecting unusual patterns in genome evolution in different taxa. Researchers

had previously observed that genomes from Drosophila species conserve gene content within

linkage groups, known as Muller elements, corresponding to chromosomes or large sub-

chromosome regions. Within these Muller elements, gene order can be reshuffled extensively

in one species relative to another (21). For alignment between *D albomicans* vs *D*

melanogaster, the CGV ideogram view shows restriction of alignment from each chromosome

in one genome to a particular chromosome or chromosome region (i.e., linkage group) in the

other genome (Fig 3A). However, within a chromosome-chromosome pair, sequence alignment

is broken into many small fragments whose relative order is not conserved. This fragmentation

of alignment is more clearly visible in the CGV dotplot, which shows that pairwise alignments

are restricted to a single chromosome pair, but appear in a scattered pattern, suggesting that

the sequence and gene order has been extensively rearranged within chromosomes (Fig 3B).

Fig 3. CGV shows conservation of linkage groups in the absence of conservation of gene order.

(A) CGV ideogram view of alignment between *Drosophila albomicans* and *Drosophila melanogaster* genomes.

Alignments are restricted to a single chromosome or chromosome region.

https://ncbi.nlm.nih.gov/genome/cgv/browse/GCF_009650485.2/GCF_000001215.4/40865/0 (B)CGV dotplot view

of alignment between *Drosophila albomicans* and *Drosophila melanogaster* demonstrates that sequence order is

"scrambled" within linkage groups, as demonstrated by a scatter pattern indicating many short rearranged

alignments. https://www.ncbi.nlm.nih.gov/genome/cgv/plot/GCF_009650485.2/GCF_000001215.4/40865/0 (C)

CGV dotplot view of alignment between starfish species *Luida sarsii* and *Asteria rubens* with similar scatter pattern

to Drosophila alignments.

https://www.ncbi.nlm.nih.gov/genome/cgv/plot/GCA_949987565.1/GCF_902459465.1/41045/0 (D) CGV

ideogram view of alignment between chromosome 1 of *Luida sarsii* and chromosome 1 of *Asteria rubens*. These

chromosomes align to each other across their length, but the alignment is broken into multiple short segments

which are extensively rearranged.

https://www.ncbi.nlm.nih.gov/genome/cgv/browse/GCA_949987565.1/GCF_902459465.1/41045/0#OX465101.1/

NC_047062.1/size=1,firstpass=0 (E) CGV dotplot view of alignment between starfish species *Luida sarsii* and

Patiria pectinifera with similar scatter pattern to Drosophila alignments.

https://www.ncbi.nlm.nih.gov/genome/cgv/plot/GCA_949987565.1/GCA_029964075.1/41165/0 (F) CGV dotplot

view of alignment between starfish species *Plazaster borealis* and *Pisaster ochraceus*. Alignments show less scatter

and more of a diagonal slope, indicating more conservation of sequence order between these two species'

genomes. https://www.ncbi.nlm.nih.gov/genome/cgv/plot/GCA_021014325.1/GCA_010994315.2/41175/466999

We observed a similar pattern when comparing some genomes from different starfish species

using CGV. When looking at pairwise alignments between *Asterias rubens*, *Patiria pectinifera*,

and *Luida sarsii* species, sequences from a chromosome from one genome mainly align to a

single other chromosome in the other species. However, within a pairwise chromosome

alignment, the sequence order is rearranged, resulting in a scatter pattern in the dotplot (Fig

3C, E). The ideogram view can show the alignment fragmentation and rearrangement in more

granular detail (Fig 3D). We also noted that some starfish pairs show more conservation of

location synteny (22) (Fig 3F), consistent with measured sequence distance based on shared k-

mers (Mash distance is 0.128 compared to other pairs with a Mash distance >0.3).

Bhutkar et al (21) speculated that the need to keep certain genes in the same regulatory

environment may result in conservation of genes within linkage groups, even in the absence of

selective pressure to maintain the gene order. More recently, conservation of macrosynteny

with extensive small-scale sequence rearrangement was also detected in comparisons between

other invertebrate species, such as cephalopods, cnidarians, jellies, and sponges (16, 23, 24).

These rearrangements were used to parse the phylogenetic relationships within this clade. We

demonstrate here that CGV can aid researchers in detecting and analyzing this phenomenon in

starfish and other evolutionarily varied taxa.

Analysis using CGV: Detection of amylase family expansion

CGV can uncover potential copy number differences in segmental gene families. These

differences may appear as gaps in the alignment in otherwise syntenic gene regions. Segmental

insertions or deletions may be too small to be apparent on the whole genome or whole

chromosome alignment but can be detected when searching and navigating to a gene of

interest.

Initial analysis of the complete human telomere-to-telomere CHM13 genome revealed an

expansion of amylase genes on chromosome 1 compared to the GRCh38 reference assembly

(25). This expansion can be validated in CGV: a search for 'amylase' in the alignment between

GRCh38 and T2T-CHM13v2 assembly finds six matches to this gene name in the GRCh38 and

twelve in the CHM13 assembly (Fig 4A). Zooming out in the region of the *AMY1A* gene on

chromosome 1 reveals a nearby sequence segment in the CHM13 assembly that is not aligned

to GRCh38 (Fig 4B). This region contains numerous annotated loci that lack official

nomenclature (i.e., LOC); six of these loci are described as 'alpha-amylase'. Therefore, there are

at least six additional alpha-amylase genes in CHM13 genome compared to the GRC reference.

It is possible that copy number of this gene is variable in humans; it is also possible that the GRC

reference genome represents fewer than the typical number of gene copies. While these two

human assemblies are likely to be high quality, other assemblies in other species may be of

lower quality, and differences in copy number observed in CGV may reflect assembly or

annotation errors.

Fig 4. **CGV can help uncover gene duplications and rearrangements in closely related genomes.**

(A) Gene search of an alignment between two human assemblies in CGV finds twelve amylase gene family

members in the human T2T-CHM113v2.0 assembly and six amylase gene family members in GRCh38.p14. (B) CGV

view showing that T2T-CHM13v2.0 contains an insertion relative to GRCh38.p14, which appears as an unaligned

region on chromosome 1. This insertion contains additional alpha-amylase family members. A popup label (tooltip)

indicates one of these additional family members.

https://www.ncbi.nlm.nih.gov/genome/cgv/browse/GCF_009914755.1/GCF_000001405.40/23025/9606#NC_060

925.1:103415704-103764412/NC_000001.11:103566852-103915505/size=1000,firstpass=0 (C) CGV view showing

that chromosome 2 of *Canis lupus familiaris* (dog) UMICH_Zoey_3.1 align to chromosomes 2, 15, 23, and 25 of

Dog10K_Boxer_Tasha.

https://ncbi.nlm.nih.gov/genome/cgv/browse/GCF_005444595.1/GCF_000002285.5/17685/9615#NC_049262.1:6

542815-78714085//size=10000 (D) UMICH_Zoey_3.1 assembly chromosome 2 alignment to Dog10K_Boxer_Tasha

chromosome 25 contains the *MALRD1* gene, which is annotated as *LOC608668* in the Tasha assembly (boxed in

red). Gene synteny is not conserved outside of the region of assembly-assembly alignment.

Analysis using CGV: Possible gene translocation between two dog assemblies

For closely related strains or species, CGV can help uncover and validate structural anomalies,

such as where gene order synteny has been disrupted. Visual inspection of the whole genome

CGV ideogram view of alignment between two dog genomes – the Great Dane Zoey

(UMICH_Zoey_3.1) and the boxer Tasha (Dog10K_Boxer_Tasha) – indicated a region that

aligned to chromosome 2 in the Zoey assembly and chromosome 25 in the Tasha assembly (Fig

4C). This region contains the *MALRD1* gene in the Zoey assembly, which is shown to align to a

MALRD1 homolog annotated as *LOC608668* in Tasha (Fig 4D). CGV alignments indicate that

LOC608668 is likely the Tasha *MALRD1* gene; there are no better alignments to *MALRD1*

detected in CGV or by an independent BLAST search of the Tasha genome.

Zooming out in the aligned region of *MALRD1* indicates that gene synteny is not conserved

outside of the *MALRD1* gene region (Fig 4D). It appears that this gene has been rearranged

from chromosome 2 on the Zoey assembly to chromosome 25 on Tasha. It is also possible that

the Tasha genome may have been misassembled in this region, and the *MALRD1* gene

sequence is properly situated on chromosome 2 within the otherwise conserved syntenic block.

A researcher would need to further examine the quality of the Tasha assembly in this region to

distinguish these possibilities, for instance, by examining the sequencing reads for the Tasha

assembly or viewing a HiC map of assembly structure.

If this anomaly represents a true difference between the two genomes, it could prove

biologically significant. The translocation of *MALRD1* may have placed it in a different gene

regulatory environment in the boxer (Tasha) genome, which could possibly result in different

levels or patterns of gene expression. The human ortholog of *MALRD1* was shown to be

involved in bile acid metabolism (26). This gene region was also genetically linked to

Alzheimer's disease (27). Therefore, if valid and not assembly artifacts, differences like these

could provide insight into human health.

Discussion

We describe here a new visualization tool for eukaryotic assembly-assembly alignments, the

Comparative Genome Viewer (CGV). We developed this web application with a view toward

serving both expert genome scientists as well as organismal biologists, students, and educators.

Users of CGV do not have to generate their own alignments or configure the software using

command line tools. Instead, they can select from our menu of available alignments, access a

view immediately in a web application, and start their analysis. We are continuing to add new

alignments regularly and invite researchers to contact us if assemblies or organisms of interest

are missing. We continue to do periodic outreach to the community to help us improve our

visual interfaces so that they are simple, intuitive, and accessible.

CGV displays whole genome sequence alignments provided by NCBI; users cannot currently

upload their own alignment data or choose assemblies to align in real time. There are both

technical and scientific considerations to allowing researchers to select and align assemblies

themselves. Currently, whole genome assembly-assembly alignments take several hours to

days, using up to one thousand CPU processing hours per pairwise alignment of larger

genomes, such as those for mammalian or plant assemblies. Moreover, whole genome

alignments are difficult to generate past a certain genetic distance (i.e., Mash > 0.3). Not only is

alignment between distantly related genomes computationally expensive, but the alignments

themselves may be of limited research value. These alignments will likely have sparse and short

segments that may correspond only to the most highly conserved coding sequence (CDS) (Fig 5;

S1 Table). We suggest protein similarity or gene orthology-based alignments as more

appropriate for comparisons between distantly related organisms. At present, we manually vet

requested alignments to make sure that assemblies are complete, of high assembly quality (e.g.

high scaffold N50 or BUSCO scores (28)), and a reasonable evolutionary distance. This review

ensures that alignments will be useful to both the original requester and others in the research

community.

Fig 5. Genome and CDS coverage of assembly-assembly alignments relative to Mash distance.

Percentages of total target genome or CDS nucleotides covered by the ungapped alignments are plotted against

the Mash distance between the pair of genomes. (A) Alignments between the human GRCh38.p14 assembly and

other vertebrates. (B) Alignments of fall army worm (FAW, *Spodoptera frugiperda*, a major insect agricultural pest)

and related insects. At lower Mash distances, the whole genome alignments cover most of the genome and CDS.

At Mash distances greater than 0.25 or 0.3, the alignment covers less than 20% of the genome overall, and

between 30% and 60% of the CDS. Refer to S1 Table for the data used to populate these graphs.

S1 Table. Alignment coverage at different Mash distances for selected assembly pairs.

Many research questions in comparative biology may be best answered by simultaneously

visualizing alignments among more than two assemblies. We are exploring user needs and

existing tools when it comes to multigenome alignment visualization, such as visualization of

pangenome data. Our lessons from developing CGV will prove valuable in this upcoming

initiative.

Materials and Methods

Preparation of assembly-assembly alignments

Genome assemblies are aligned using a two-phase pipeline first described in Steinberg et al

(29), with adaptations for cross-species alignments. In the first phase, initial alignments are

generated using BLAST (20) or LASTZ (19), or imported from a third-party source such as UCSC

(30). In the second phase, alignments are merged and ranked to distinguish reciprocal-best

alignments from additional alignments that are locally best on one assembly but not the other.

The resulting alignment set is omnidirectional and can be used to project information from

query to subject assembly or vice versa.

In the first phase, for both BLAST and LASTZ, repetitive sequences present in the query and

target assemblies are soft-masked using WindowMasker (31). The default parameters are

usually suitable for aligning assembly pairs within the same species. However, more aggressive

masking is required when aligning cross-species assemblies. The masking rate is adjusted with

the parameter `t_thres_pct` set to 99.5 (default, for BLAST same-species), 98.5 (for BLAST

cross-species), or 97.5 (LASTZ cross-species), or lower for some genomes with extensive and

diverse repeat composition. The 97.5 - 98.5 values typically result in a masking percentage

similar to RepeatMasker (http://www.repeatmasker.org) with a species-specific repeat library,

with the advantage of not needing to define repeat models beforehand.

Genome assemblies are aligned using either BLAST or LASTZ. The selection of the aligner and

specific parameters depends on the level of similarity between the assemblies. We use Mash

(32) to compute the approximate distance between two assemblies (Fig 5). BLAST is employed

for aligning pairs of assemblies belonging to the same species, as well as cross-species assembly

pairs with a Mash distance of less than 0.1. An exemplar BLAST command is:

`blastn -evalue 0.0001 -gapextend 1 -gapopen 2 -max_target_seqs`

`250 -soft_masking true -task megablast -window_size 150 -`

`word_size 28`

A BLAST `word_size` of 28 is used for pairs of assemblies with Mash distances below 0.05,

such as human and orangutan, while a `word_size` of 16 is used to enhance sensitivity for

more distant cross-species pairs with Mash distances ranging from 0.05 to 0.1, such as human

and rhesus macaque.

Assembly pairs with Mash distances exceeding 0.1, such as human and mouse assemblies, are

aligned using LASTZ. The `make_lastz_chains` pipeline (19) is employed to generate

alignments between query and target assemblies in UCSC chain format. The default parameters

are often adequate to produce satisfactory alignments for many assembly pairs, though some

distant assembly pairs (e.g., Mexican tetra-medaka) warrant changes such as the use of a

different substitution matrix (`BLASTZ_Q=HoxD55`).

Alignments generated with the `make_lastz_chains` pipeline or precomputed alignments

imported from UCSC are in chain format. The UCSC chainNet pipeline (33) is run on query x

target and target x query alignment sets separately so that the alignments are 'flattened' in a

way that the reference sequences are covered only once by the alignments in each set, and the

two chainNet outputs are concatenated.

In the second phase, alignments are converted to NCBI ASN.1 format and processed further for

the NCBI's CGV and GDV browsers. In this phase, the full set of BLAST or LASTZ-derived

alignments are processed to merge neighboring alignments and split and rank overlapping

alignments to identify a set of best alignments (S1A Figure). Merging is accomplished on a

sequence-pair-by-sequence-pair basis, and ranking is accomplished globally for the assembly

pair. The process is designed to find a dominant diagonal among a set of potentially conflicting

alignments.

S1 Figure. Merging, sorting, and ranking assembly-assembly alignments.

(A) Flowchart showing that adjacent alignment segments are merged. Subsequently, alignments are split once

again at large gaps. (B) Flowchart showing how overlapping alignments are separated, ranked, and re-merged.

Reciprocal best-placed alignments are designated as "first pass", while the non-best placed alignment is

designated "second pass".

Merging involves the following steps. First, when applicable, alignments based on common

underlying sequence components of the assemblies (e.g., the same BAC component used in

both human GRCh37 and GRCh38) are identified and merged into the longest and most

consistent stretches possible. Second, adjacent alignments are merged if there are no

conflicting alignments. Third, alignments are split on gaps using a default threshold of 50 bp (for

the same or closer species) or 50 kb (for more distant species), or longer than 5% of the

alignment length. Alignments are also split at any point where they intersect with overlapping

alignments (S1A Figure). Duplicate or low-quality alignments fully contained within higher-

quality alignments are dropped.

After merging and splitting, alignments are subsequently processed using a sorting and ranking

algorithm (S1B Figure). Alignments are sorted based on a series of properties, including the use

of common components, assembly level (alignment to chromosomes preferred over alignment

to unplaced scaffolds), total sequence identity, and alignment length. The alignments are then

scanned twice, once each on the query and subject sequence ranges, to sort out reciprocal

best-placed (also referred to as "first pass" or reciprocity=3) and non-best placed (also referred

to as "second pass" or reciprocity=1 or 2) alignment sets. Finally, all alignments in each

reciprocity are merged again to stitch together adjacent alignments with no conflicting

alignments into the longest representative stretches.

Assembly-assembly alignments are stored in an internal database available for rendering in

NCBI's CGV and GDV browsers. The alignment data are publicly available in GFF3 and ASN.1

formats at https://ftp.ncbi.nlm.nih.gov/pub/remap/.

For display in CGV, assembly alignment batches are filtered to keep only alignments that

contain chromosome scaffolds as both anchors and targets, since non-chromosomal scaffolds

are not displayed in this viewer.   Alignments are converted into a compact binary format

designed to keep only the synteny data required for display. This preparation step is done by

programs written in C++ and bash scripts that tie them together.

Technical architecture of CGV

CGV operates on a two-tier model, with a front end implemented using HTML/JavaScript

running in the user's web browser and a backend running at NCBI. The graphical rendering is

done on the front end using modern WebGL technologies. The main advantage of this approach

is speed and fluidity of the user interface since most of the alignment data needed to be

rendered is sent to the front end at the initial load and there are no additional roundtrips to the

server when the user interacts with the page (e.g., panning or zooming). Using front end

graphical rendering also reduces the network traffic between NCBI and the end user, which

makes CGV more responsive.

The backend of the CGV application resolves internal alignment identifiers to an alignment data

file that the front end can use for generating graphical images. The back end is implemented as

an industry-standard gRPC service written in C++ and running in a scalable NCBI service mesh

(linkerd, namerd, consul). When a CGV view is initially loaded, our gRPC service requests the

alignment data needed for the particular page. On graphical pages, gRPC resolves an alignment

identifier to a URL with prepared synteny/alignment data and the page loads the file at this URL

into a WASM module which is written in C++ and compiled with Emscripten. The WASM module

serves this data on demand to the page's JavaScript code which uses it for building the image

and all user interactions.

In parallel, the list of assembly-assembly alignments and their metadata is sent to the selection

menu (i.e., "Set up your view") on the CGV landing page. This allows the selection menu to

report scientific and common species names, assembly accessions, and assembly names.

The alignment selection menu on the CGV landing page is a traditional web form page. We

utilize the NCBI version of United States Web Design System (USWDS) design standards and

components (https://www.ncbi.nlm.nih.gov/style-guide) in order to unify graphical design with

other US government pages.

On the more graphically intensive ideogram and dotplot pages, graphical rendering is done in

the user's web browser using WebGL using d3.js or pixi.js libraries, which allows for efficient

interactivity, scalability, and fluidity of user interaction. Other elements of the page use

jQuery/extJS and USWDS components. CGV reuses chromosome ideograms initially developed

for NCBI's Genome Data Viewer (2).

Gene annotations shown in CGV are obtained from NCBI's public databases using NCBI Entrez

Programming Utilities (E-utilities) (https://www.ncbi.nlm.nih.gov/books/NBK25501/).

Annotation-build specific gene search is provided by an NCBI Datasets

(https://www.ncbi.nlm.nih.gov/datasets/) gRPC service.

Design of CGV Application

A philosophy of user-centered design, which puts user needs at the forefront of decision-

making, was an integral element in the development of the Comparative Genome Viewer

(CGV). Participants for user research were recruited from members of the genomics research

community who provided their contact information through feedback links on the CGV

application and other sequence analysis tools at NCBI. Some of these researchers were

previously familiar with CGV, while others had little or no experience with this tool. Data from

user research testing sessions was compiled and analyzed for patterns in behavior, thereby

allowing the team to validate that the design was moving in a direction that facilitated analysis

of sequence alignment data. To date, we have conducted user research with over 30 different

experts in the field of comparative genomics. We also evaluate the application for Section 508

compliance, which also helps insures CGV performs well on mobile devices and is accessible to

users with limited or no visibility.

Acknowledgments

We thank Anne Ketter and Emily W Davis for help with project planning and coordination. We

thank Guangfeng Song for aid with user experience research and analysis.

Thanks to Wayne Matten, Michelle Formica-Frizzi, and others in the NCBI customer service

team for marketing, webinars, and video tutorials. Anatoliy Kuznetsov and Andrei Shkeda

conducted early technical design work that influenced the architecture of the CGV application.

Deanna M. Church and Mike DiCuccio participated in initial design and testing of the assembly

alignment protocol. Finally, we thank members of the greater genomics research community

who have participated in usability sessions and provided feedback and alignment requests to

CGV. CGV was developed as a part of the National Institutes of Health's Comparative Genomics

Resource (CGR) (https://www.ncbi.nlm.nih.gov/comparative-genomics-resource/). This work

was supported by the National Center for Biotechnology Information of the National Library of

Medicine (NLM), National Institutes of Health.

References

1.   Bornstein K, Gryan G, Chang ES, Marchler-Bauer A, Schneider VA. The NIH Comparative

Genomics Resource: addressing the promises and challenges of comparative genomics on human

health. BMC Genomics. 2023;24(1):575.

2.   Rangwala SH, Kuznetsov A, Ananiev V, Asztalos A, Borodin E, Evgeniev V, et al. Accessing NCBI

data using the NCBI Sequence Viewer and Genome Data Viewer (GDV). Genome Res. 2021;31(1):159-69.

3.   Nassar LR, Barber GP, Benet-Pages A, Casper J, Clawson H, Diekhans M, et al. The UCSC Genome

Browser database: 2023 update. Nucleic Acids Res. 2023;51(D1):D1188-D95.

4.   Buels R, Yao E, Diesh CM, Hayes RD, Munoz-Torres M, Helt G, et al. JBrowse: a dynamic web

platform for genome visualization and analysis. Genome Biol. 2016;17:66.

5.   Cabanettes F, Klopp C. D-GENIES: dot plot large genomes in an interactive, efficient and simple

way. PeerJ. 2018;6:e4958.

6.   Haug-Baltzell A, Stephens SA, Davey S, Scheidegger CE, Lyons E. SynMap2 and SynMap3D: web-

based whole-genome synteny browsers. Bioinformatics. 2017;33(14):2197-8.

7.   Krzywinski M, Schein J, Birol I, Connors J, Gascoyne R, Horsman D, et al. Circos: an information

aesthetic for comparative genomics. Genome Res. 2009;19(9):1639-45.

8.   Wang H, Su Y, Mackey AJ, Kraemer ET, Kissinger JC. SynView: a GBrowse-compatible approach

to visualizing comparative genome data. Bioinformatics. 2006;22(18):2308-9.

9.   Xiao Z, Lam HM. ShinySyn: a Shiny/R application for the interactive visualization and integration

of macro- and micro-synteny data. Bioinformatics. 2022;38(18):4406-8.

10.   Richardson JE, Baldarelli RM, Bult CJ. Multiple genome viewer (MGV): a new tool for

visualization and comparison of multiple annotated genomes. Mamm Genome. 2022;33(1):44-54.

11.   Diesh C, Stevens GJ, Xie P, De Jesus Martinez T, Hershberg EA, Leung A, et al. JBrowse 2: a

modular genome browser with views of synteny and structural variation. Genome Biol. 2023;24(1):74.

12.   Albert VA, Krabbenhoft TJ. Navigating the CoGe Online Software Suite for Polyploidy Research.

Methods Mol Biol. 2023;2545:19-45.

13.   Lee J, Hong WY, Cho M, Sim M, Lee D, Ko Y, et al. Synteny Portal: a web-based application portal

for synteny block analysis. Nucleic Acids Res. 2016;44(W1):W35-40.

14.   Wang Y, Tang H, Debarry JD, Tan X, Li J, Wang X, et al. MCScanX: a toolkit for detection and

evolutionary analysis of gene synteny and collinearity. Nucleic Acids Res. 2012;40(7):e49.

15.   Lovell JT, Sreedasyam A, Schranz ME, Wilson M, Carlson JW, Harkess A, et al. GENESPACE tracks

regions of interest and gene copy number variation across multiple genomes. Elife. 2022;11.

16.   Schultz DT, Haddock SHD, Bredeson JV, Green RE, Simakov O, Rokhsar DS. Ancient gene linkages

support ctenophores as sister to other animals. Nature. 2023;618(7963):110-7.

17.   Muffato M, Louis A, Poisnel CE, Roest Crollius H. Genomicus: a database and a browser to study

gene synteny in modern and ancestral genomes. Bioinformatics. 2010;26(8):1119-21.

18. Darling AC, Mau B, Blattner FR, Perna NT. Mauve: multiple alignment of conserved genomic

sequence with rearrangements. Genome Res. 2004;14(7):1394-403.

19. Kirilenko BM, Munegowda C, Osipova E, Jebb D, Sharma V, Blumer M, et al. Integrating gene

annotation with orthology inference at scale. Science. 2023;380(6643):eabn3107.

20. Zhang Z, Schwartz S, Wagner L, Miller W. A greedy algorithm for aligning DNA sequences. J

Comput Biol. 2000;7(1-2):203-14.

21. Bhutkar A, Schaeffer SW, Russo SM, Xu M, Smith TF, Gelbart WM. Chromosomal rearrangement

inferred from comparisons of 12 Drosophila genomes. Genetics. 2008;179(3):1657-80.

22. Lee Y, Kim B, Jung J, Koh B, Jhang SY, Ban C, et al. Chromosome-level genome assembly of

Plazaster borealis sheds light on the morphogenesis of multiarmed starfish and its regenerative capacity.

Gigascience. 2022;11.

23. Simakov O, Bredeson J, Berkoff K, Marletaz F, Mitros T, Schultz DT, et al. Deeply conserved

synteny and the evolution of metazoan chromosomes. Sci Adv. 2022;8(5):eabi5884.

24. Albertin CB, Medina-Ruiz S, Mitros T, Schmidbaur H, Sanchez G, Wang ZY, et al. Genome and

transcriptome mechanisms driving cephalopod evolution. Nat Commun. 2022;13(1):2427.

25. Vollger MR, Guitart X, Dishuck PC, Mercuri L, Harvey WT, Gershman A, et al. Segmental

duplications and their variation in a complete human genome. Science. 2022;376(6588):eabj6965.

26. Reue K, Lee JM, Vergnes L. Regulation of bile acid homeostasis by the intestinal Diet1-FGF15/19

axis. Curr Opin Lipidol. 2014;25(2):140-7.

27. Zubenko GS, Hughes HB, 3rd. Predicted gene sequence C10orf112 is transcribed, exhibits tissue-

specific expression, and may correspond to AD7. Genomics. 2009;93(4):376-82.

28. Simao FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. BUSCO: assessing genome

assembly and annotation completeness with single-copy orthologs. Bioinformatics. 2015;31(19):3210-2.

29.   Steinberg KM, Schneider VA, Graves-Lindsay TA, Fulton RS, Agarwala R, Huddleston J, et al.

Single haplotype assembly of the human genome from a hydatidiform mole. Genome Res.

2014;24(12):2066-76.

30.   Blanchette M, Kent WJ, Riemer C, Elnitski L, Smit AF, Roskin KM, et al. Aligning multiple genomic

sequences with the threaded blockset aligner. Genome Res. 2004;14(4):708-15.

31.   Morgulis A, Gertz EM, Schaffer AA, Agarwala R. WindowMasker: window-based masker for

sequenced genomes. Bioinformatics. 2006;22(2):134-41.

32.   Ondov BD, Treangen TJ, Melsted P, Mallonee AB, Bergman NH, Koren S, et al. Mash: fast

genome and metagenome distance estimation using MinHash. Genome Biol. 2016;17(1):132.

33.   Kent WJ, Baertsch R, Hinrichs A, Miller W, Haussler D. Evolution's cauldron: duplication,

deletion, and rearrangement in the mouse and human genomes. Proc Natl Acad Sci U S A.

2003;100(20):11484-9.

Fig1

A

B

Reset to genome view

Fig2

Fig3

A

B

C

D

Fig4

A

B

Fig5