Concurrent response and action effect representations across the

somatomotor cortices during novel task preparation.

Ana F. Palenciano*[1], Carlos González-García[1], Jan De Houwer[2], Baptist Liefooghe[3], and

Marcel Brass[4,5]

[1] Mind, Brain and Behavior Research Center, University of Granada, 18071, Granada, Spain

[2] Department of Experimental Clinical and Health Psychology, Ghent University, 9000,

Ghent, Belgium

[3] Department of Psychology, Utrecht University, 3508, Utrech, The Netherlands

[4] Berlin School of Mind and Brain, Department of Psychology, Humboldt University of Berlin,

10117, Berlin, Germany

[5] Department of Experimental Psychology, Ghent University, 9000, Ghent, Belgium

Corresponding author. E-mail address: palencianoap@ugr.es (Ana F. Palenciano)

Abbreviated tittle: Response and action effect coding for novel tasks.

Number of pages: 52

Number of figures: 6

Number of words: abstract, 250; introduction, 649; discussion, 1500.

Conflict of interest statement: The authors declare no competing financial interests.

Acknowledgment: This research was supported by grant G00951N of the Flemish

Government attributed to BL and JDH. AFP was supported by Grant PAIDI21_00207 of the

Andalusian Autonomic Government. CG-G was supported by Grant IJC2019-040208-I and

Project PID2020-116342GA-I00 funded by MCIN/AEI/10.13039/501100011033, and Grant

RYC2021-033536-I funded by MCIN/AEI/10.13039/501100011033 and by the European

Union NextGeneration EU/PRTR. JDH was supported by Methusalem funding from the

Special Research Fund (BOF) of Ghent University (reference number: BOF22/MET_V/002).

MB was supported by an Einstein Strategic Professorship of the Einstein Foundation Berlin

(EPP-2018-483) and by the Deutsche Forschungsgemeinschaft (DFG, German Research

Foundation) under Germany's Excellence Strategy – EXC 2002/1 "Science of Intelligence"

(project number: 390523135). BL was supported by the Utrecht University Focus Area on

Human-Centered Artificial Intelligence.

Abstract

Instructions allow us to fulfill novel and complex tasks on the first try. This skill has been

linked to preparatory brain signals that encode upcoming demands in advance, facilitating

novel performance. To deepen insight into these processes, we aimed to explore whether

instructions pre-activated task-relevant motoric and perceptual neural states. Critically, we

addressed whether these representations anticipated activity patterns guiding overt

sensorimotor processing, which could reflect that internally simulating the novel task

facilitates the preparation. To do so, we collected functional magnetic resonance imaging

data while participants encoded and implemented novel stimulus-response associations.

Participants also completed localizer tasks designed to isolate the neural representations of

the mappings-relevant motor responses, perceptual consequences, and stimulus

categories. Using canonical template tracking, we identified whether and where these

sensorimotor representations were pre-activated. We found that response-related templates

were encoded in advance in regions linked with action control, entailing not only the

instructed responses but also their action effects. This result was particularly robust in

primary motor and somatosensory cortices. While, following our predictions, we found a

systematic decrease in the irrelevant stimulus templates' representational strength

compared to the relevant ones, this difference was due to below-zero estimates linked to

the irrelevant category activity patterns. Overall, our findings reflect that instruction

processing relies on the sensorimotor cortices to anticipate motoric and kinesthetic

representations of prospective action plans, suggesting the engagement of motor imagery

during novel task preparation. More generally, they stress that the somatomotor system

could participate with higher-level frontoparietal regions during anticipatory task control.

Keywords: *Novel instructed behavior, cognitive control, task preparation, fMRI, multivariate*

analyses

Significance Statement

By using instructions, humans can successfully implement novel stimulus-response

mapping on the first try. Here, we investigated whether this skill entailed anticipating the

mappings-relevant motoric and perceptual neural states before task execution. Using

tailored pattern analyses and localizer tasks, we isolated a series of task-relevant

sensorimotor neural representations and addressed whether they were pre-activated during

novel mapping preparation. We found widespread anticipation of the response-related

activity patterns, which not only entailed the relevant motor responses but also their

perceptual consequences. This finding was particularly robust in primary somatomotor

areas. Overall, our results stress the contribution of lower-level cortices during instructed

performance, suggesting that when no experience with a task is available, simulating the

prospective sensorimotor processing may facilitate optimal preparation.

1 Introduction

By using symbolic information, humans convey and communicate task procedures that they have never implemented before. Thanks to this ability, we can acquire novel and arbitrary stimulus-response (S-R) mappings directly from instructions, highlighting the adaptability of human learning processes. This complex skill has been associated with anticipatory brain signals across a set of frontal and parietal areas (Cole et al., 2010; Demanet et al., 2016; Dumontheil et al., 2011; Hartstra et al., 2011, 2012; Palenciano, González-García, Arco, & Ruz, 2019; Ruge & Wolfensteller, 2010) that substantially overlaps with the Multiple Demand Network (MDN; Duncan, 2010; Fedorenko et al., 2013), a system linked to the implementation of cognitive control. These areas encode in advance behaviorally relevant instruction information, such as the targeted stimuli, the relevant responses, or the rules binding both (Cole, Reynolds, et al., 2013; González-García et al., 2017, 2021; Muhle-Karbe et al., 2017; Palenciano, González-García, Arco, Pessoa, et al., 2019). Accordingly, current models emphasize the role of the MDN in assembling novel task representations from instructions (Brass et al., 2017; Cole, Laurent, et al., 2013). Critically, these representations are characterized by their abstract, multidimensional nature (Bourguignon et al., 2018; Cole et al., 2011; Cole, Reynolds, et al., 2013; Palenciano, González-García, Arco, Pessoa, et al., 2019). How this higher-order task coding ultimately shapes the downstream sensorimotor processing and impacts instruction implementation remains elusive to date.

To address this issue, we aimed to investigate the contribution of lower-level, sensorimotor neural representations to novel task preparation. In this regard, it has been suggested that internally simulating novel tasks through imagery may facilitate optimal preparation (Moran & O'Shea, 2020). This view, based on frameworks like the motor simulation theory (Jeannerod, 1994, 2001), emphasizes that imagery is driven by an internal replay engaging action representations shared with overt performance (Grush, 2004; Guillot & Collet, 2005;

Hardwick et al., 2018; Jeannerod, 2001). Hence, by means of imagery, novel instructions

could pre-activate sensorimotor neural states overlapping with those driving task execution.

That would bridge the above-described abstract task coding found in the MDN with the later

behavioral performance. Supporting this idea, previous behavioral results have evidenced

an association between novel task implementation and imagery (Liefooghe et al., 2021;

Theeuwes et al., 2018). Robust evidence also emphasizes that just encoding instructions

induces automatic response activation before task execution (Liefooghe et al., 2012, 2013;

Liefooghe & De Houwer, 2018; Meiran et al., 2015a; Meiran & Cohen-Kdoshay, 2012).

Moreover, imposing secondary motor demands during instruction preparation disrupts

execution in an effector-specific fashion (Palenciano et al., 2021), suggesting that optimal

anticipatory control may rely on task-relevant response representations. At the neural level,

functional magnetic resonance imaging (fMRI) data have shown heightened mean activity

across motor regions during instruction encoding and preparation (Hartstra et al., 2011,

2012; Ruge & Wolfensteller, 2010). Electroencephalography findings have also evidenced

that instructions elicit neurophysiological signatures of response pre-activation (Everaert et

al., 2014; Meiran et al., 2015a), namely, lateralized readiness potentials (Gratton et al.,

1988). Nonetheless, although all these results converge on the relevance of lower-level

motoric processes for novel task preparation, the underlying neural computations have not

been systematically explored.

To fill this gap, we addressed whether encoding instructions pre-activated motoric and

perceptual neural representations that share content and format with those involved during

sensorimotor processing. To do so, we collected fMRI data while participants encoded and

executed pairs of novel S-R mappings. Then, they completed a series of localizer tasks

seeking to isolate the neural activity patterns of three mappings' components: the effector-

specific motor responses, their perceptual consequences, and the targeted stimulus

categories. Employing canonical template tracking (Palenciano et al., 2022), we estimated

standard neural representations (or templates) from the localizer task data and assessed

their representational strength during the S-R mapping paradigm. Our main prediction was

that the mappings-relevant sensorimotor templates would be anticipatorily encoded in a set

of regions of interest (ROIs) located across the motor, somatosensory, and visual brain

systems.

2   Materials and methods

2.1.   Participants

Thirty-five participants completed the experiment (27 women, 8 men; mean age = 23 years,

SD = 3 years). All of them were right-handed, with normal or corrected-to-normal vision, and

MRI compatible. Most of the participants were native Dutch speakers (N = 33), except two

of them, who were fluent English speakers. All of them signed a consent form approved by

the UZ Ghent Ethics Committee. The participation was compensated with 36.25 EURO. The

data of four participants were discarded due to excessive movement. Another participant

was unable to finish the MRI session. The data of the remaining 30 participants were

included in all the analyses, with the exception of the Action Effects analysis (see Section

2.6), where data from three additional participants needed to be excluded due to technical

issues with the tactile stimulators.

2.2.   Apparatus, stimuli, and protocol

Novel S-R mapping paradigm

In the main paradigm, participants encoded pairs of novel S-R mappings to later implement

one of them upon target presentation. For that purpose, we generated a pool of unique

associations, all consisting of a picture of an animal, and a verbal label indicating the

response linked to that stimulus. The mappings' response involved either the left index, the

left middle, the right index, or the right middle finger. The response labels were presented

either in Dutch ("*L wijs*", "*L middel*", "*R wijs*", "*R middel*") or in English ("*L index*", "*L middle*",

"*R index*", "*R middle*") depending on the participant's preferred language, and were shown

in font Arial, size 50. The pictures were obtained from a previous database containing 770

animate images (Brady et al., 2013; Brodeur et al., 2014; Griffin et al., 2007; Konkle et al.,

2010) and completed with stimuli from the Animal Images Database (Possidónio et al.,

2019), the C.A.R.E. database (Russo et al., 2018) and available sources with Creative

Commons license. All the stimuli were resized to 200 by 200 pixels, and appeared on

greyscale, with the background removed and the animal centered on the canvas. Four

animal categories were used: canines, felines, birds, and sea animals. For each mapping,

we displayed the picture on the left side of the screen, and the response label, on the right

side. In each trial, two mappings were shown, one in the upper part of the screen, and

another on the lower part.

For each participant, we created a pool of 640 different S-R mappings, arranged in 320

pairs. In each trial, two out of the four possible responses and two out of the four possible

animal categories were used. We ensured an equal representation of the six pair-wise

responses combination ([left index, left middle], [left index, right index], [left index, right

middle], [left middle, right index], [left middle, right middle], and [right index, right middle])

and the six pair-wise combinations of stimulus categories ([cat, dog], [cat, bird], [cat, fish],

[dog, bird], [dog, fish], and [bird, fish]). We further controlled the mappings among response

and stimulus pairs (i.e.: trials with [cat-left index / dog right-middle] were equally frequent as

trials with [bird-left middle / fish-right index]), to avoid that some categories were more

frequently linked to some responses by chance. The location of the mappings on the screen

(upper or lower part) was also counterbalanced across category and response conditions.

The sequence of events within a trial is depicted in **Fig. 1A**. First, the participants encoded

the mappings during 2s, followed by a preparation interval displaying a fixation cross, and

with a jittered duration (mean = 2.39s, SD = 1.26s, range = [0.6, 5s], values extracted from

a pseudo-logarithmic distribution). Then, a probe stimulus was presented during a window

of 1.75s where participants responded. In 90% of the trials, probes corresponded either to

the upper or the lower mapping picture (with a 50% probability each). In the remaining 10%

of trials (*catch trials*), a novel picture – not in any of the two mappings – was presented, and

participants were asked to respond with the left and right index and middle fingers

simultaneously. Catch trials were included to ensure that both S-R mappings were equally

well encoded. Catch probes were also animal pictures but always belonged to one of the

two categories that were not shown in the current mappings. Finally, a fixation was

presented during a jittered inter-trial interval (ITI), with the same temporal properties as the

previous one.

Before the MRI session, participants practiced the S-R mapping task until they achieved an

85% of accuracy. To ensure that the mappings from the practice session were not further

employed, we used a different set of stimuli composed of inanimate items (musical

instruments and vehicles). The stimuli from the practice session were not further employed

in the experiment. Once inside the MRI scanner, participants completed four task runs, each

containing 80 trials. They could take a self-paced pause between runs. Overall, the duration

of the S-R mapping task was approximately 45 minutes.

Localizer tasks

After completing the main paradigm, the participants remained on the scanner and

completed three localizer tasks, designed to isolate the representations of the mappings'

motor response, action effects, and stimulus categories. The localizers' experimental design

aimed to maximize the similarity between the targeted component and the S-R mapping

paradigm while keeping the remaining task features as independent as possible. This was

done to avoid confounding specific template activation with more general motoric or

perceptual processes (Palenciano et al., 2023). Each localizer block lasted approximately 5

minutes, and the order in which they were presented was counterbalanced across

participants.

In the motor response localizer (**Fig. 1B, left panel**), participants responded to cues that

explicitly indicated the required key presses. The cues were composed of four rectangles

(200 by 100 pixels) located along the horizontal axis of the screen, each indicating a

response with a different finger: from left to right, with the left index, left middle, right index,

or right middle finger. The cues were displayed against a black background, and to indicate

the required response, the corresponding rectangle was filled in white. We employed these

cues instead of verbal labels (as in the S-R mapping task) to avoid semantic processing

during the localizer. Instead, we aimed to capitalize on the motor component, by generalizing

across instructed response modalities (Liefooghe et al., 2012). Each trial of this localizer

consisted of the presentation of the response cue for 1.5s, where participants responded,

and a jitter ITI with a fixation cross (same temporal properties as above). Participants

completed 80 trials, 20 for each response condition.

For the action effect localizer (**Fig. 1B, middle panel**), we applied single-tap tactile

stimulation on the tip of the participants' left and right index and middle fingers, separately.

To do so, we employed four MRI-compatible piezo-electric stimulation devices (mini-PTS,

http://www.dancerdesign.co.uk/), each covering an area of 1cm^2 of the participants'

fingertips. The stimulation pulses were generated with single sawtooth waves, with 40ms of

duration. We attached the stimulators to two foam structures, one per hand, in an ergonomic

position for the index and middle fingers. The intensity of the stimulation was adjusted to the

participants' sensitivity at the beginning of the MRI session. In the localizer, the stimulation

was applied while the cues from the motor response localizer were displayed on the screen.

Nonetheless, the cues were not informative about the stimulated finger and all the rectangles

appeared filled in black. This was done to keep the two response-related localizers as similar

as possible perceptually. Each trial consisted of the pulse of stimulation (40ms) together

with the presentation of the uninformative cues for 1.5s, both synchronized at the onset, and

a jittered ITI (same temporal properties as above). To facilitate the participants' engagement

during the block, we added a secondary, memory task in 16% of the trials. In these trials,

the stimulation was not delivered, and the cues appeared filled in color (pink, yellow, blue,

or green). All participants saw the 4 colors in a randomized order, with each color being

shown in 4 trials. The participants were instructed to pay attention and to remember the

colors displayed, ignoring their appearance order. When the localizer finished, the

participants verbally reported the colors that they remembered. The task consisted of 96

trials, and the stimulation was applied in 80 of them (20 trials per stimulation condition).

Finally, participants completed a stimulus category localizer (**Fig. 1B, right panel**), where

they discriminated between animal and non-animal items. For each participant, we randomly

selected 80 animal pictures (20 per category) and 20 inanimate ones (10 musical

instruments, 10 vehicles) from the same database as in the S-R mapping task. All pictures

were 200 by 200 pixels, in greyscale, with the background removed and the item centered

on the image. In each trial, the stimulus was presented for 1.5s, followed by a jittered ITI

(same temporal properties as above). Participants needed to respond simultaneously with

their two index and middle fingers only when an inanimate stimulus appeared, which

happened in 20% of the trials. In the remaining 80% of trials, an animal stimulus from one

of four categories used in the mappings was shown, and no response was required. With

this approach, we aimed to isolate the perceptual processing of the four animal categories,

in the absence of any motor preparation or execution. The localizer consisted of 100 trials,

80 displaying an animal picture (20 per animal category).

All tasks were programmed in Psychopy (v3; Peirce et al., 2019) and were presented on a

screen located at the back of the scanner and projected onto a mirror system. Responses

were collected using a pair of MRI-compatible joysticks.

2.3. Experimental design

In the S-R mapping paradigm, we manipulated two within-subjects variables: the mappings'

relevant responses ([left index, left middle], [left index, right index], [left index, right middle],

[left middle, right index], [left middle, right middle], and [right index, right middle]), and

stimulus category ([cat, dog], [cat, bird], [cat, fish], [dog, bird], [dog, fish], and [bird, fish]).

We collected 48 trials per response and per category condition. For experimental control,

we fully crossed these two variables, resulting in 36 different mappings pairs, sampled

through eight observations each. However, all our analyses treated independently the

response and category manipulations (e.g.: in the response-related analyses, we collapsed

across stimulus categories within response conditions). Hence, our analyses were based on

48 observations per cell.

In the motor response and action effects localizers, we had one within-subject manipulation:

the effector (left index, left middle, right index, and right middle) used to respond or stimulate.

In the stimulus category localizer, we manipulated within subjects the stimulus animacy

(animate, and inanimate items) and the animal category of the animate stimulus (cat, dog,

bird, and fish). In the three localizers, we collected 20 observations per condition of interest.

A. Novel S-R Mapping paradigm:

Encoding [2s]   Preparation interval [0.6-5s, jittered, mean: 2.266s]   Probe [1.5s]   ITI [0.6-5s, jittered, mean: 2.266s]

B. Localizer tasks:

MOTOR RESPONSE — *Press left index* / *Press right index* / *Press left middle* / *Press right middle*

ACTION EFFECTS — [16% of trials] *Remember the color!*

STIMULUS CATEGORY — *Cat: Don't respond!* / *Dog: Don't respond!* / *Bird: Don't respond!* / *Fish: Don't respond!* [20% of trials] *Inanimate: Respond!*

Figure 1. Experimental paradigms. **A.** Trial's sequence of events during the S-R mapping paradigm.

B. Localizer tasks. In the motor response localizer (left panel), explicit cues indicated the required

response (either with the left or right index or middle finger). In the action effect localizer (middle

panel), stimulation was applied separately in each of these four fingers. The visual display consisted

of uninformative cues, except for 16% of trials which entailed a secondary, color memory task. In the

stimulus category localizer (right panel), pictures of animals belonging to the four categories used in

the mappings were interspersed with pictures of inanimate items. Participants responded only to the

infrequent (20%) inanimate stimuli.

2.4.   MRI data acquisition and preprocessing

The MRI data was collected with a 3T Siemens Magnetom Trio scanner at the Ghent

University Hospital (UZ Ghent, Belgium) with a 64-channel head coil. Whole-brain functional

images were obtained with a T2* EPI sequence [TR = 1730 ms, TE = 30 ms, flip angle=66°,

FOV = 210 mm, image matrix = 84 × 84, voxel size = 2.5 x 2.5 x 2.5 mm, distance factor =

0%, 50 slices, slice acceleration factor = 2, slice orientation according to the participants'

AC-PC line]. Four 400-volumes runs were collected for the S-R mapping task, and three

185-volumes runs for the localizers. Before the functional runs, we collected anatomical T1-

weighted images using a MP-RAGE sequence [TR = 2250 ms, TE = 4.18 ms, flip angle=9°,

FOV = 256 mm, image matrix = 256 x 256, voxel size = 1 x 1 x 1 mm, 176 slices]. We also

collected phase and magnitude field map images [TR = 520 ms, TE1 = 4.92 ms,

TE2 = 7.38 ms, flip angle=60°, FOV = 210 mm, image matrix = 70 × 70, voxel

size = 3 × 3 × 2.5 mm, distance factor=0%, 50 slices], to correct for inhomogeneities of the

magnetic field. Overall, participants remained inside the scanner for 75 minutes.

MRI Data preprocessing and GLM estimation (see section 2.6) were carried out on SPM12

(http://www.fil.ion.ucl.ac.uk/spm/software/spm12/). The functional images were spatially

realigned, corrected for magnetic field inhomogeneities, and slice-time corrected. The T1

anatomical images were defaced, coregistered to the participants' mean functional volume,

and segmented. For univariate analyses, the functional images were normalized to the MNI

space using the forward deformation fields derived from the T1 segmentation and smoothed

with an 8mm FWHM Gaussian kernel. The multivariate analyses were performed on

unnormalized and unsmoothed images to avoid distorting the finer-grained activity patterns.

The normalization and smoothing were performed on the results maps. The T1

segmentation also generated inverse deformation fields which were used to convert the

ROIs masks, defined in the MNI space, into the participants' native anatomical space.

Four participants showed sustained, excessive movement (>2.5mm) and were removed

from the sample. However, 3 additional participants displayed punctual movements larger

than our voxel size but restricted on time. For those participants, the volumes affected by

movement (less than 1% of the acquired images) were masked in further analyses using

regressors generated with the ART toolbox (http://web.mit.edu/swg/software.htm).

2.5.   Statistical analyses: behavioral data.

We analyzed the effect of the mappings' relevant responses and stimulus categories on

response accuracy and speed, separately. This was done in an exploratory fashion, to

assess whether the conditions were behaviorally balanced. We carried out repeated-

measures ANOVAs, using as within-subject factor either the response or the category

condition. A Greenhouse-Geisser correction was used when sphericity was violated. Post

hoc pair-wise *t*-tests were carried out to identify conditions where performance deviated from

the rest. A Bonferroni-Holm correction for multiple comparisons was applied. Catch trials

were excluded from the analyses. Only correct trials were included in RT ANOVAs.

2.6.   Statistical analyses: fMRI data

Univariate analyses

We first conducted a series of univariate analyses to provide an overview of the regions

engaged by both the S-R mapping paradigm and the localizer tasks. For the S-R mapping

task, we fitted a GLM with regressors covering the mapping encoding stage, the preparation

interval, and the probe, separately for the six response conditions. Each regressor was

defined using the event duration, except the probe, which was modeled with the trial RT.

The ITI was not defined in the model and thus contributed to the implicit baseline. As

nuisance regressors, we included errors, catch trials, and the six movement parameters

derived from the realignment. Within participants, we carried out three one-sample *t*-tests to

contrast each event of interest (encoding, preparation, and probe) against the baseline,

collapsing across conditions. The outputs were entered into group-level *t*-tests to detect

significant BOLD signal increases induced by our paradigm's events. The results were

corrected at *p* < .05 using a cluster-wise FWE multiple-comparison correction (from an initial,

uncorrected threshold of *p* < .001 and k = 10).

The three localizers were analyzed in separate GLMs. For the motor response localizer, we

defined a regressor for each response condition, locked to the onset of the response cue

and with duration fixed to the trials' RT. Errors were included as a nuisance regressor. For

the action effects localizer, the GLM included a regressor for each stimulation condition,

locked and with the same duration as the stimulation pulse. Trials from the memory task

were included as a nuisance regressor. Finally, for the stimulus category localizer, we

modeled a regressor per animal category, with the same onset and duration as the stimulus

presentation. Inanimate target trials and errors were included as nuisance regressors. The

localizers GLMs also included the six motion parameters. We computed one-sample *t*-tests

to detect BOLD signal increases linked to the localizers' event of interest (response, tactile

stimulation, and target presentation) against the implicit baseline. We followed the same

statistical inference strategy as with the S-R mapping task.

Canonical template tracking

The main goal of this study was to assess whether the motoric and visual representations

guiding overt performance and perception were covertly engaged during novel mapping

preparation. To do so, we employed a canonical template tracking procedure (Palenciano

et al., 2023; for other examples, see González-García et al., 2021; Wimber et al., 2015).

Specifically, we estimated a series of canonical neural representations (or templates) from

each localizer data. These templates aimed to capture the corresponding motoric or

perceptual component representation in the same format as engaged during overt

processing (i.e., pressing a key, experiencing the action effect, or perceiving the stimulus

category). Then, we assess the representational strength of these templates during the S-R

mapping paradigm, by computing their similarity with the multivoxel activity patterns evoked

by the novel mappings' preparation interval.

First, we fitted a set of GLMs to estimate the multi-voxel activity patterns during the S-R

mapping task (**Fig. 2A**). To maximize the number of data points being fed into the template

tracking procedure, we estimated the signal in a trial-by-trial fashion. We used a least-

square-sum (LSS) approach (Arco et al., 2018; Mumford et al., 2012, 2014) to reduce the

collinearity. For each trial, we estimated a separate GLM with the following regressors: (1)

the preparation interval of the current trial; (2) the encoding stage, collapsing across all

correct trials; (3) the preparation stage, excluding the current trial but collapsing across the

remaining correct trials; (4) the probe event, collapsing across all correct trials. All trial-wise

GLMs further included nuisance regressors modelling, (5) errors and (6) catch trials (using

the same onset and duration as in the univariate GLM defined above), and the six movement

parameters. We iterated the estimation procedure across all the trials, obtaining a beta map

for the preparation interval of each pair of novel mappings. Next, we performed one-sample

t-tests comparing each beta map against the baseline, to normalize noise from a univariate

perspective (Kriegeskorte et al., 2008; Walther et al., 2016). The activity patterns used for

the template tracking were directly extracted from the resulting trial-wise *t*-maps.

Next, we estimated the canonical neural templates using the three localizers' data (**Fig. 2B**).

To maximize the robustness of the templates, we estimated standard GLMs that provided a

single beta map per localizer condition. Within each localizer, the four conditions' beta

images (corresponding to the four effectors in the case of the motor response and action

effects localizers, and to the four animal categories in the stimulus category localizer) were

contrasted separately against the baseline. The templates were extracted from the resulting

t-maps.

Finally, we estimated the similarity between the templates and each activity pattern from the

mappings' preparation interval (**Fig. 2C**). In each trial from the S-R mapping task, two out of

the four possible motor responses, and two out of the four possible stimulus categories,

were prepared in advance. This design allowed distinguishing between two subsets of

templates regarding a given trial: relevant (i.e., prepared) and irrelevant (i.e., not prepared)

ones. Hence, we estimated the similarity between the main task data and both types of

templates and used the irrelevant similarity measurements as baseline. Critically, the use of

this baseline condition allowed us to infer the specificity of the motoric and perceptual

information encoded during the preparation (Palenciano et al., 2022). To compare the

similarity estimates from several templates simultaneously, while accounting for the shared

variance among them, we used a multiple regression framework (e.g., Palenciano et al.,

2019).

We first carried out the template tracking separately for each localizer task and on a set of

ROIs (see *ROI selection* section). For each region, we first normalized (i.e., z-scored) each

activity pattern from the S-R mapping task and each canonical template, ensuring that all of

them had a mean of 0 and a standard deviation of 1. That sought to mitigate the impact of

univariate activity differences on our canonical template results. Then, we fitted a series of

multiple regressions in which the multi-voxel activity pattern from a trial's preparation interval

was predicted using the four localizer templates as regressors. That provided beta

coefficients that reflected the similarity between relevant and irrelevant templates and the

main task data. Then, we averaged the beta weights across templates and trials to obtain a

single relevant and irrelevant value per ROI. At the group level, we carried out one-tailed,

paired-sample Wilcoxon signed rank tests to assess the hypothesis that the beta weights

from relevant templates were higher than the irrelevant ones. This comparison offered an

estimate of the representational strength of specific templates while controlling for more

general motor preparation or visual expectation processes. Furthermore, one-sample

Wilcoxon tests were performed to assess whether relevant and irrelevant beta weights were

different from zero. It is important to note that the later results could be contaminated by the

positive bias present in similarity estimates, and hence, were only interpreted to qualify the

comparison against the irrelevant (or baseline) condition. In all cases, we controlled for

multiple comparisons (i.e., number of ROIs) using a Bonferroni-Holm correction (Holm,

1979).

Since the motor response and the action effects localizers tapped into a common component

of the S-R mappings, we performed an additional analysis including templates from the two

localizers simultaneously. With this approach, we aimed to disentangle the contribution of

passive perception of the action effects, from a richer action representation conveying motor

planning and execution. The procedure was identical to the one specified above, with the

exception that the multiple regressions included eight regressors (one per effector condition

and localizer) instead of four. After averaging, we obtained four estimates per ROI,

corresponding to relevant and irrelevant motor response and action effects templates. Next,

we carried out a repeated measures ANOVA including the ROI, the localizer task, and the

template condition (relevant, irrelevant) as within-subject factors. We carried out post hoc

comparisons to explore the three-way interaction between ROI, Localizer, and Template,

using a Bonferroni-Holm multiple comparison correction.

Finally, we ran an additional exploratory analysis to better understand which aspects of the

mappings' instructed responses were captured by our canonical templates. In this regard,

the novel mappings involved different response effectors (index, middle fingers) and

different laterality (left-hand, right-hand responses). Due to the pervasive effects of response

laterality, we aimed to address whether specific response effector representations could be

also detected with the canonical template procedure. While the experiment was designed to

control for laterality, by balancing it across the response conditions, we further investigated

this issue by repeating our main analysis in a subset of novel mappings where only the

templates' effector information (and not the laterality) was relevant. First, we extracted the

relevant and irrelevant beta weights from S-R mappings that entailed the same effector from

the two hands (i.e., the conditions [left index, right index] and [left middle, right middle]) and

estimated the activation of the relevant effector templates in comparison with the irrelevant

ones. In each of the targeted conditions, the relevant and irrelevant templates would differ

only regarding the response effector being equated regarding response laterality. We carried

out this comparison in all the ROIs examined in the main analysis. Second, to provide a

deeper insight into the processes implemented in the primary motor and somatosensory

cortex, we carried out an additional test on these ROIs using the same subset of trials but

further differentiating the laterality of the templates. Specifically, each canonical template

was estimated using data from the contralateral hemisphere ROIs. For instance, to predict

the activity patterns of mappings involving [left index, right index], we used data from the left

M1 to estimate the similarity with the right index (relevant) and right middle (irrelevant)

templates, and data from the right M1 to estimate the similarity with the left index (relevant)

and left middle (irrelevant) templates. This way, we ensured that our results were driven by

specific effector representations engaged in the corresponding, contralateral primary cortex.

Nonetheless, it is important to stress the post hoc nature of these analyses, for which the

experiment design was not optimized. As a result, only 33% of the observations from the S-

R mapping task were included, with a corresponding negative impact on the analysis's

statistical power. Still, these results helped qualify the planned analyses here reported.

Figure 2. Canonical template tracking. **A.** Estimation of trial-by-trial activity patterns from the preparation interval of the S-R mapping paradigm. In the trial shown as an example (Trial_i), the prepared mapping responses entailed the right index and middle fingers. **B.** Canonical template estimation. From each localizer, we estimated a single activity pattern per condition. Here we illustrate the motor response localizer, with templates corresponding to the four response conditions. The templates relevant to the mapping are shown in bold. **C.** A multiple regression was fitted to predict each trial's activity pattern. The regressors entailed the four (two relevant, in bold, and two irrelevant) canonical templates, together with a constant and an error term. The inference was carried out on the comparison between averaged beta weights from relevant and irrelevant templates. **D.** ROIs used in the motor response and action effect analyses. **E.** ROIs used in the stimulus category analyses.

ROI selection

We used two sets of a-priori ROIs, one for the motor response and action effects templates,

and another one for the stimulus category ones. In the former case, we selected regions

associated with action control and somatosensory processing. Specifically, we focused on

the primary motor (M1) and somatosensory cortex (S1), the ventral and dorsal portion of the

premotor cortex (PMv and PMd, respectively), the supplementary and pre-supplementary

motor areas (SMA and preSMA, respectively), and the inferior and superior parietal lobe

(IPL and SPL, respectively). A post hoc ROI was placed on the posterior insula (pInsula)

due to its involvement during the tactile stimulation processing (see **Fig. 4**). The impact of

the stimulus category templates was assessed in regions processing both physical stimulus

attributes and more abstract, categorical information. Specifically, we used as ROI the

primary visual cortex (V1), the early visual cortices from V2 to V4 (V2-V4), the lateral

occipital complex (LOC), and a broader region covering the ventral pathway until the

inferotemporal cortex (VentralVisual).

The M1, PMv, PMd, SMA, preSMA, and S1 ROIs were obtained from the Human Motor Area

Template (HMAT; Mayka et al., 2006), while the IPL, SPL, V1, V2-V4, LOC, and

VentralVisual masks were extracted from the HCP-MMP1.0 parcellation (Glasser et al.,

2016). All ROIs covered the bilateral portions of the corresponding region, except for the M1

and S1, for which we used separated ROIs for the left and right hemispheres to account for

their contralateral functional organization. This was done to avoid capturing broader

interhemispheric differences in univariate signal that would remain after the z-score

correction. Finally, we used MarsBar (http://www.sourceforge.net/projects/marsbar) for the

pInsula, building two 10mm radius spherical ROIs located in the bilateral activation peaks

found in the univariate results from the action effect localizer.

For each localizer, we included a control ROI that was unrelated to the templates of interest

and where no effects were expected. Specifically, we used V1 in the motor response and

action effects analyses, and M1 in the stimulus-related analysis.

Searchlight analyses

To complement the ROI results, we additionally performed exploratory whole-brain analyses

using a searchlight procedure. We iterated a 5-voxel radius sphere across every location of

the brain, performing multiple regressions with the S-R mapping task activity patterns and

templates estimated from the current sphere data. The averaged beta coefficients for the

relevant and irrelevant templates were assigned to the voxel at the center of the sphere. As

a result, we obtained two beta maps per participant and localizer task, one for the relevant

and another for the irrelevant templates. Next, within participants, we subtracted the

irrelevant beta map from the relevant one. The resulting images were normalized and

smoothed, and entered into group-level one-sided, one-sample *t*-tests to detect significantly

greater beta weights linked to relevant templates, in comparison with irrelevant ones.

Control analysis

Finally, we aimed to ensure that our results were not driven by differences in the reliability

of the canonical templates employed. The experimental design sought to eliminate such

confound by balancing each localizer's set of templates (e.g., the four response effectors'

templates) between the relevant and irrelevant conditions. Nonetheless, we computed two

reliability measurements per template (González-García et al., 2021; Palenciano et al.,

2023). First, to estimate the overall univariate noise, we computed the Signal-to-Noise Ratio

(SNR), dividing the absolute mean t value from a given template by its standard deviation.

Then, to assess the stability of the templates' informational content, we used a

correlationability index (González-García et al., 2021). We split each localizer data into odd

and even trials and fitted new GLMs, obtaining two (instead of a single) canonical templates

per localizer condition. The correlationability index consisted of the Pearson correlation

coefficient between each condition's pair of templates. To identify reliability differences

among templates, we run one-way repeated measures ANOVAs separately for each

measurement, localizer, and ROI. The template condition was used as a single factor. Post

hoc paired-sample t-tests (Bonferroni-corrected for multiple comparisons) were computed

whenever the main effect of the template condition was significant.

3. Results

3.1.   Behavioral results.

Performance was in line with previous studies using novel S-R associations. On average,

participants responded accurately on 89% of trials of the S-R mapping task (SD = 6%), with

a mean RT of 7 ms (SD = 99 ms). On catch trials, the average accuracy was 84% (SD =

22%), and the mean RT, 943 ms (SD = 134 ms). Regarding the localizers' performance, in

the motor response task, participants responded correctly in 98% of trials (SD = 1.7%), with

an average RT of 523 ms (SD = 58 ms). In the action effects localizer, all the participants

recalled correctly the four colors used in the secondary memory task. Finally, in the stimulus

category localizer, the mean accuracy was 99% (SD = 12%), with an average of 0.2% of

false alarms (SD = 0.5%) and 4% of misses (SD = 8%). The average RT was 942 ms (SD

= 14 ms).

We explored whether the behavioral performance was modulated by the mappings' relevant

responses and stimulus categories. The response condition effect (**Fig. 3**) was not

significant neither accuracy rates, $F(3.4, 99.9) = 0.83$, $p = .49$, $\eta_p^2 < .01$, nor in RTs, $F(5,$

$145) = 1.36$, $p = .24$, $\eta_p^2 = .05$, supporting that performance was equivalent across response

pairings. Regarding the stimulus category (**Fig. 3**), however, we found a significant effect on

both accuracy rates, $F(5, 145) = 3.84$, $p = .003$, $\eta_p^2 = .12$, and RTs, $F(5, 145) = 2.38$, $p =$

$.04$, $\eta_p^2 = .08$. Post hoc paired-sample t-test showed that, in general, participants performed

worst mappings that included the category pair [bird-fish]. Accuracy was lower in this

condition than in the [cat-dog] pair (T = 3.43, *p* = .01) and the [cat-fish] pair (T = 3.90, *p* =

.002), with an additional marginal difference with the [cat-bird] pair (T = 2.84, *p* = 0.07). RTs

were marginally slower in [bird-fish] trials than in [dog-fish] ones (T = 2.79, *p* = 0.09). More

qualitatively, the overall pattern reflected a general impoverishment in the [bird-fish] category

pairings than in the other conditions.

Figure 3. Boxplots displaying the behavioral performance in the novel S-R Mapping paradigm. **A.**

RTs (left) and percentage of correct responses (right) across the six response conditions. **B.** RTs

(left) and percentage of correct responses (right) across the six stimulus category conditions. LI = left

index; RI = right index; LM = left middle; RM = right middle.

Finally, to explore whether the S-R mappings combination further modulated behavioral

performance we carried out additional repeated measures ANOVA including the response

and the category condition as factors. In line with the above-described results, the effect of

category was significant for both RT, $F(4.31, 124.84) = 2.54$, *p* = .03, η_p^2 = .08, and accuracy

rates, $F(4.22, 122.23) = 3.84$, *p* = .003, η_p^2 = .12. Contrary, neither the response condition

nor the interaction was significant (all *p values* > .20).

3.2.   fMRI results

Univariate analyses

As a first step, we aimed to characterize the regions engaged by the different stages of the

S-R mapping paradigm. Replicating previous findings, encoding novel S-R associations was

associated with increased activity broadly in the MDN and the visual and motor systems

(**Fig. 4A**). A wide cluster ([38, -66, -16], k = 57033) covered the early and late visual cortices,

extending dorso-medially into the precuneus and anteriorly into the inferotemporal cortex.

The cluster also included S1, M1, PMC, and SMA. Within the MDN, it covered the lateral

DLPFC, incurring into the anterior insula/frontal operculum (aI/fO) in the left hemisphere,

and the SPL, IPL, and IPS. It also covered part of the cerebellum and the thalamus. A

second cluster ([32, 22, 4], k = 539) covered the right aI/fO. Conversely, during the

mappings' preparation interval (**Fig. 4B**), only a restricted set of regions showed a consistent

activity increase. Specifically, we found significant clusters bilaterally in the aI/fO (left: [-30,

20, -2], k = 867, right: [34, 22, -2], k = 825), the inferior frontal junction (left: [-42, 24, 28], k

= 258), right: [40, 30, 30], k = 442), the preSMA ([6, 20, 50], k = 1323), and the bilateral IPS

and IPL (left: [-32, -54, 42], k = 376; right: [36, -54, 42], k = 387). Finally, probe processing

(**Fig. 4C**) substantially engaged the visual and motor system and the MDN, as well as

subcortical nuclei. An extended cluster ([36, 2, 10], k = 104250) overlapped with the

activations found during the encoding, also including the premotor cortices, the preSMA, the

middle cingulum, the middle and posterior region of the insula, the temporal operculum, the

rostrolateral prefrontal cortex, and the basal ganglia.

Figure 4. Univariate analysis results. **A.** *t* maps showing the increases in activity associated with the encoding (left panel), preparation (middle panel), and implementation (right panel) of the novel S-R mappings. **B.** *t* maps showing the increases in activity associated with the three localizers' event of interest: the response in the motor response localizer (in blue), the tactile stimulation in the action effects localizer (in green), and the animate picture visualization in the stimulus category localizer (yellow). All maps are statistically thresholded at $p < .05$.

With a similar approach, we described the regions engaged by the three localizers' events of interest (**Fig. 4D**). As expected, the motor response localizer induced an increase in activity in regions associated with action control. A wide cluster ([20, -88, -12], k = 32523) bilaterally covered the motor cortices, including M1, premotor cortices, and SMA, and extended into the S1, SPL, and IPL. As expected from the presence of visual stimulation

during this localizer, the cluster also included early and late visual cortices. Two additional

clusters were found in the basal ganglia and posterior insula in both hemispheres (left: [-46,

0, 12], k = 2418; right [28, 0, 2], k = 1223). Conversely, the tactile stimulation applied in the

action effect localizer was linked to increased activity in the pInsula, bilaterally (left: [-46, -

20, 14, k = 950), right: [50, -24, 22], k = 1046). This task also recruited the early visual cortex

([24, -90, 4], k = 4269). Finally, the stimulus category localizer was associated with

heightened activity in a cluster covering the bilateral early and late visual cortex and the right

SPL and IPL (left: [-38, -84, -2], k = 7009; right: [40, -82, -4], k = 8761), and another cluster

located in the left SPL and IPL ([-22, -48, 40], k = 575). Increased activity was also found in

the bilateral hippocampus (left: [-22, 30, 0], k = 224; right: [22, -30, -2], k = 273).

Canonical template tracking: ROI analyses

To address the main goal of this work, we conducted a series of multivariate, ROI-based

analyses which informed on whether the preparation to implement novel S-R mappings

activated templates of the instructed responses and stimuli. On one hand, we explored how

the novel mappings' responses were represented, using templates estimated from the motor

response and the action effects localizers. On the other, we also addressed the anticipatory

encoding of the mappings' stimuli, using templates from the stimulus category localizer.

Response-related analyses

The results obtained using the motor response templates are displayed in **Fig. 5A** and in

the extended data table **Fig. 5-1**. First, we assessed whether the relevant motor response

templates were activated to a greater degree than the irrelevant ones, using the latter as

baseline. The results supported this hypothesis in most of the regions explored. The paired-

sample Wilcoxon tests were significant in the left and right M1 and S1, SMA, PMd, IPL, SPL,

and pInsula. Additional one-sample tests showed that the relevant templates' beta

coefficients were greater than zero in all these regions and the PMv. Finally, after visually

inspecting our data, we observed that some regions' irrelevant templates were linked to

below-zero beta coefficients. The corresponding one-sample tests on irrelevant template

data evidenced significant below-zero beta weights in the left and right M1 and S1, and the

pInsula. Contrary, significant above-zero irrelevant templates' beta weights were found in

the PMv. As expected, neither test was significant in the control ROI, V1.

Next, we applied the same analytical approach with the templates obtained from the action

effects localizer (**Fig. 5B**, **Fig 5-1**). Like the motor response, the beta weights from relevant

and irrelevant action effect templates differed in most of the ROIs. The effect was significant

in the right M1, the left and right S1, the SMA, PMd, IPL, SPL, and pInsula. The relevant

templates were linked to statistically significant above-zero beta weights across all these

regions and the left M1. Finally, we found below-zero beta coefficients linked to irrelevant

action effects templates in the left S1 and pInsula. Again, neither test was significant in V1.

Figure 5. Boxplots showing the averaged beta weights for relevant and irrelevant canonical templates. **A.** Motor response localizer results. **B.** Action effects localizer results. **C.** Template tracking results after including the motor response and the action effect templates in the same multiple regressions. **D.** Stimulus category localizer results.

Considering the equivalent pattern of results obtained with the two response-related

localizers and the nature of both tasks, our findings could also reflect that the motor response

and action effects templates captured overlapping representations, and consequently,

shared variance from the mappings' activity patterns. To estimate the unique variance

explained by each localizer, we ran an additional analysis including the templates from both

tasks simultaneously in the multiple regressions. The resulting averaged beta coefficients

are displayed in **Fig. 5C.** To explore these results, we ran repeated measures ANOVA on

the beta weights, using ROI (left M1, right M1, left S1, right S1, SMA, preSMA, PMv, PMd,

IPL, SPL pInsula), Localizer (motor response, action effects), and Template (relevant,

irrelevant) as factors. We found significant main effects of Template, $F(1, 26) = 216.65$, $p <$

$.001$, $\eta^2_p = .893$, and ROI, $F(5.80, 150,80) = 2.76$, $p = .015$, $\eta^2_p = .096$, but not of Localizer,

$F(1, 26) = 0.25$, $p = .619$, $\eta^2_p = .010$. Nonetheless, the interaction between Localizer and

Template was significant, $F(1, 26) = 19.60$, $p < .001$, $\eta^2_p = .430$, supporting that the localizer

task modulated the relationship between relevant and irrelevant templates. To address the

anatomical extent of this effect, we next focused on the significant three-way Localizer x

Template x ROI interaction, $F(4.89, 127.07) = 18.20$, $p < .001$, $\eta^2_p = .412$, and ran post hoc

comparisons contrasting relevant and irrelevant templates separately for each localizer and

ROI (**Fig. 5-2**). These tests showed a generalized increase in beta coefficients in relevant

than irrelevant templates in the two localizer tasks. Significant differences involving motor

response and action effect templates were found in left and right M1 and S1, SMA, PMd,

IPL, SP, and pInsula. The test was also significant for the action effect templates in the

preSMA. Overall, even when the templates' effect was numerically different depending on

the localizer (as evidenced by the two-way and three-way significant interactions), it

persisted after including the two localizers' data simultaneously in the analysis, suggesting

that both sets of templates captured independent aspects of the S-R mapping task data.

Finally, we explored the response attributes that the motor response and action effects

templates were capturing. The visual inspection of the motor response (**Fig. 5-5**) and action

effect (**Fig. 5-6**) beta weights across the six response conditions highlighted the robust effect

of response laterality, especially, in primary and secondary somatomotor areas. That could

be identified in trials where the division between relevant and irrelevant templates was

mapped into different laterality (i.e., in the mappings' response conditions [left index, left

middle] and [right index, right middle]). To rule out that our results were uniquely driven by

differences between right- and left-hand response representations, we first compared

relevant and irrelevant templates in a subset of trials in which only the effector information

was relevant for the analysis. The results from the motor response localizer (**Fig. 5-4A**)

showed greater beta weights for relevant than irrelevant templates in the right S1 (z = 2.76,

p = .002, $p_{corrected}$ = .032). A similar pattern was found in the right M1 (z = 2.24, p = .013,

$p_{corrected}$ = .124) and the SPL (z = 1.71, p = .044, $p_{corrected}$ = .394), although these tests did

not survive the correction for multiple comparisons. Regarding the action effects templates

(**Fig. 5-4B**), we found a significant effect in the left S1 (z = 2.80, p = .003, $p_{corrected}$ = .028).

Next, we aimed to shed some light on the representations instantiated in M1 and S1. Since

these two regions have clear contralateral neural architecture, we analyzed separately the

two hemispheres' data. Specifically, we used data from the contralateral hemisphere of

specific mappings and templates to perform the analysis. The tests contrasting relevant and

irrelevant templates remained significant in S1 when using data from either the motor

response (**Fig. 5-4C**) or the action effect localizers (**Fig. 5-4D)**. Specifically, we found a

significant effect of the motor response templates in the right S1 (z = 2.67, p = .004, $p_{corrected}$

= .016), and the action effects templates in the left S1 (z = 2.97, p = .002, $p_{corrected}$ = .002).

An equivalent pattern was found in the left and right M1, although neither test survived the

statistical correction (Motor response, right M1: z = 1.60, p = .054, $p_{corrected}$ = .188; Action

effects, left M1: z = 1.50, p = .066, $p_{corrected}$ = .081).

        Stimulus-related analyses

We further aimed to address whether the canonical templates obtained during the

perception of the instructed stimuli were reactivated during the novel mappings' preparation.

To do so, we followed the same procedure as in the response-related analyses, using

templates built from the stimulus category localizer data. The results are displayed in **Fig.**

5D and **Fig. 5-3**. The paired-sample Wilcoxon tests were significant in all the regions

explored, i.e., in V1, V2-4, LOC, and VentralVisual. Nonetheless, we did not find above-zero

beta coefficients linked to relevant templates in either region. Instead, the effect was driven

by below-zero beta weights associated with the irrelevant templates. The latter effect was

significant in the V2-4, LOC, and VentralVIsual ROIs. In these analyses, no effects were

detected in the control ROI, M1.

Since the behavioral analyses showed that one of the mappings' stimulus category

conditions (the [bird-fish] combination) was linked to generally impoverished performance,

we further explored the template tracking findings across conditions to ensure that the

unexpected, below-zero irrelevant beta weights were not driven by an outlier pair of

categories. Nonetheless, the pattern found in the main results seemed stable across

mappings' conditions and ROIs (**Fig. 5-7**), suggesting that these findings were not driven by

trials from a particular condition.

Template tracking: searchlight

To complement the ROIs' results, we carried out an exploratory searchlight analysis. That

provided whole-brain maps displaying regions where the beta weights were higher for

relevant than irrelevant canonical templates. The results, displayed in **Fig. 6**, overlapped

substantially with the two sets of ROIs selected a priori for this work. When the motor

response localizer templates were used, we found a wide, bilateral cluster ([32, -14, 56], k

= 27155) which covered the motor cortices (M1, PMd and SMA), S1, the IPL and SPL, and

the posterior insula. The cluster also extended into the precuneus. A second cluster ([0, -62,

-20], k = 5390) covered the cerebellum bilaterally. Regarding the action effects, we found a

significant cluster ([44, -22, 58], k = 13826) covering the SMA and the right M1 and PMd,

and extended into the right S1, IPL, SPL and pInsula. A second cluster involved the left S1,

IPL and SPL ([-52, -24, 50], k = 1200). A significant cluster was also found in the cerebellum

([-20, -60, -26], k = 1718. Finally, the stimulus category whole-brain map included a

significant, bilateral cluster ([50, -76, -2], k = 18133) extending from the early visual cortices

into the fusiform gyrus and the inferotemporal cortex, and which also covered part of the

LOC.

● Motor Response ● Action Effects ● Stimulus Category

Figure 6. Searchlight results. Brain maps displaying clusters with significantly higher beta

weights from relevant than irrelevant templates. The motor response localizer results are

shown in blue, the action effects ones in green, and the stimulus category, in yellow.

Control analysis

We aimed to ensure that our results were not biased by differences in reliability across the

canonical templates. To do so, we computed and compared (using repeated measures

ANOVAs) the SNR and the correlationability across the four template conditions from each

localizer. This analysis was carried out, separately, for each of the ROIs explored in the main

analysis. In general terms, these results supported that these two reliability measurements

were equivalent across template conditions (**Fig. 5-8**). We only found significant differences

regarding the SNR in action effect templates in the IPL, $F(3, 78) = 2.85$, $p < .043$, $\eta^2_p = .099$,

driven by greater SNR in the right index templates than the right middle finger template. The

fact that no other significant result was found in the remaining 67 tests (even when no

statistical correction for multiple comparisons was applied) strongly suggests that the

findings here reported were not contaminated by differences in the templates' reliability.

4. Discussion

In this work, we aimed to characterize the preparatory neural states leading to novel S-R

mapping execution. In contrast to previous literature, which has focused on abstract task-

level representations in frontoparietal MDN regions, here we addressed the contribution of

lower-level stimulus and response coding across the sensorimotor systems. Critically, we

aimed to detect activity patterns with a shared encoding format between novel S-R mapping

preparation and overt performance. Using canonical template tracking, we isolated the

mappings' relevant sensorimotor activity patterns and estimated their representational

strength during task preparation. Our findings showed that the response-related

components were indeed pre-activated during novel mapping preparation, entailing the

presence of both motor response and action effect templates. Nonetheless, we obtained

less interpretable results regarding the stimulus-related templates, with non-relevant

stimulus representations displaying systematically smaller, below-zero similarity estimates,

while the representational strength of the relevant categories did not differ from zero.

Overall, our findings support that novel instruction preparation anticipates prospective motor

and kinesthetic representations while leaving open the role of stimulus neural codes.

Our work revealed that the anticipatory encoding of response-related representations was a

rather widespread phenomenon across regions previously associated with action control

(Haggard, 2008; Rizzolatti & Luppino, 2001), detecting this effect in M1, SMA, PMd, S1, the

IPS, and the posterior insular cortex. Still, our results highlighted two areas, M1 and S1, that

showed numerically larger and more consistent effects. While these two regions were not

engaged in a univariate fashion during the preparation, we found robust above-zero

representational strength for the relevant templates, accompanied by below-zero values for

irrelevant ones. Moreover, when we addressed whether the response effector information

was retained after removing the impact of laterality, we replicated the same pattern of

results, stressing the specificity of these representations. Overall, our findings converge on

the relevance of downstream processing, especially in M1 and S1, during higher-level novel

task preparation. Previous literature had already pointed in that direction, showing

heightened mean activity across the somatomotor system during instruction processing

(Hartstra et al., 2011, 2012; Ruge & Wolfensteller, 2010). Nonetheless, the univariate

approach employed in these studies left unaddressed the computations implemented in

these regions. In this work, we showed for the first time that instructions pre-activated

concrete motoric and kinesthetic neural states in M1 and S1. Furthermore, we evidenced

that these regions' encoding space was shared with sensorimotor execution, offering a

deeper insight into their representational format (Kriegeskorte & Kievit, 2013).

Taken together, these findings follow the predictions from simulation-based frameworks

(Grush, 2004; Jeannerod, 1994, 2001) and suggest that the task-relevant sensorimotor

processing was internally anticipated before the execution. Hence, they point toward the

involvement of motor imagery when preparing for novel tasks, in agreement with theoretical

(Brass et al., 2017; Moran & O'Shea, 2020) and empirical work (Liefooghe et al., 2021;

Theeuwes et al., 2018). In this regard, recent behavioral evidence showed that novel task

preparation relied on the pre-activation of effector-specific response representations, since

blocking their access (via dual-task demands) robustly impaired performance (Palenciano

et al., 2021). Our results seem to capture the neural mechanism underpinning that

behavioral effect, evidencing the presence of these low-level response representations, and

further characterizing the different action-related information that could be anticipated via

motor simulation. Also supporting this interpretation, the regions that showed anticipatory

template instantiation have been linked to motor imagery in the past (Hardwick et al., 2018;

Hétu et al., 2013). Moreover, some of them encode movements in a common

representational space that generalizes between imagery and overt execution (Zabicki et

al., 2017), as we found in our data. Nonetheless, it is important to stress that a similar

scenario has been described in the broader action control field. There, it has been suggested

that preparatory signals push the motor system closer to execution-induced states during

action planning (Churchland et al., 2010; Shenoy et al., 2013), with recent neuroimaging

data supporting a representational overlap between motor planning and execution in M1

and S1 (Ariani et al., 2020). That leaves open whether our findings reflect a specific,

imagery-driven mechanism engaged in novel scenarios, or instead, a rather general tool at

the service of anticipatory action control.

Our findings also highlighted the role of task-relevant action effects during instruction

preparation. This result is highly relevant for theoretical frameworks based on ideomotor

principles, emphasizing that action planning is the consequence of pre-activating the

intended perceptual effects linked to actions (Hommel, 2009; Hommel et al., 2001; Shin et

al., 2010). Agreeing with this view and previous neuroimaging findings (Waszak et al., 2012),

we showed that task-relevant kinesthetic outputs were anticipated before mapping

execution. Nonetheless, our work extended this research by identifying finer-grained action

effect representations, and more critically, by evidencing this effect for the first time in more

volatile, novel task scenarios. Furthermore, our findings showed that the motor response

and action effects templates captured independent facets of the preparatory activity

patterns. This result is less straightforward to interpret from the above-mentioned ideomotor

perspective, which would predict a greater overlap between both sets of templates. In

opposition, our findings could reflect that information about the motor responses and their

kinesthetic consequences may be segregated and multiplexed across these regions'

representational spaces. Alternatively, they could reflect that each set of templates could

also contain non-shared response information. For instance, the response templates could

capture additional proprioceptive information beyond the tactile feedback. Future research

will be key to describing in greater detail the nature of these representations as well as their

functional role during preparation.

Contrary to the response-related results, we obtained a less clear pattern regarding the

stimulus templates. We found the expected relationship between prepared and non-

prepared target categories, with smaller representational strength for the latter than the

former. ROI and whole-brain analyses showed that this pattern spread across the visual

stream, with numerically greater effects in higher-order areas sensitive to abstract category

information, such as the LOC (Grill-Spector et al., 1999). However, while the irrelevant

category templates were consistently linked to negative estimates, the relevant ones were

not statistically different from zero. While this finding may relate to the nature of visual

imagery (which may differ from the motor domain), we believe that this pattern could also

be driven by the design of the category localizer task. Since participants responded to the

broader animate-inanimate distinction, the individual templates' animal categories remained

task-irrelevant. Although this decision allowed us to isolate the stimulus processing from

unwanted motor confounds, it may have hindered the category information conveyed by the

templates. Having said that, the consistent results linked to the irrelevant templates suggest

that this information was somehow anticipated. However, although it could be tempting to

interpret this finding as the suppression of irrelevant category representations during task

preparation, the similarity measurements computed (i.e., non-cross-validated; Walther et al.,

2016) require that relative instead of absolute activation estimates are considered

(Palenciano et al., 2022; Walther et al., 2016). Hence, no inference should be performed

based only on the irrelevant categories' below-zero representational strength. Instead, we

encourage caution when interpreting these results and emphasize that more compelling

evidence is required to understand the significance of stimulus-related representations in

this context.

Overall, our results highlighted the action-oriented nature of proactive novel task control,

and more specifically, the involvement of downstream motor and somatosensory regions.

This view could seem in conflict with previous literature which has stressed the role of

cognitive control-related MDN areas during novel instruction processing (Cole et al., 2010;

Demanet et al., 2016; Dumontheil et al., 2011; Hartstra et al., 2011, 2012; Palenciano,

González-García, Arco, & Ruz, 2019; Ruge & Wolfensteller, 2010). Therefore, the main

frameworks explaining instructed performance have focused on how these frontoparietal

regions flexibly generate task representation from scratch (Brass et al., 2009, 2017; Cole,

Laurent, et al., 2013). Here we also supported this notion, by showing heightened averaged

activity during mapping preparation across the MDN. Nonetheless, our methodology

enabled us to identify a co-existent preparatory mechanism outside this network and

anchored in lower-level cortices. We believe that our findings reflect a different but

complementary facet of novel task processing that entails the anticipatory tuning of

sensorimotor systems via motor and kinesthetic template instantiation. In this regard, our

results left open a key question: how do the higher-order task representations previously

identified in the MDN, and the sensorimotor coding here described, interact? Related to that,

current working memory models state that the abstract, multidimensional representations

held in the MDN (especially, in the lateral prefrontal cortex) lead to the activation of more

specific content encoded in the sensory regions (Christophel et al., 2017; Curtis &

D'Esposito, 2003; Sreenivasan et al., 2014). Following this framework, one could expect that

the previously documented instruction representations in the MDN precede and influence

the engagement of the concrete sensorimotor templates here reported. Future studies

exploiting temporally resolved brain data will be crucial to address this hypothesis and bridge

both sets of findings. That would enrich the current understanding of instruction processing,

and more generally, help to build more holistic frameworks of anticipatory task control that

integrate higher-order and downstream cognitive processes.

5. Conclusions

In this work, we addressed whether novel S-R mapping preparation entailed the pre-

activation of sensorimotor representations that overlapped with those guiding task

execution. By combining fMRI recording and tailored pattern analyses, we evidenced the

anticipatory engagement of templates conveying the mappings' motor responses and their

perceptual consequences. While this effect was widespread across frontoparietal regions

linked to action control, it was particularly robust in the primary somatosensory and motor

cortices. In contrast, we did not obtain evidence supporting an equivalent pattern regarding

stimulus-related perceptual templates. Overall, our findings stress that motoric and

kinesthetic representations anchored in the sensorimotor systems are proactively engaged

by instructions, suggesting that motor imagery may underpin preparation in novel contexts.

More broadly, these results suggest that the anticipatory and flexible tuning of the

sensorimotor interface systems could be a key component for current models of cognitive

control.

Credit author statement

AFP: Conceptualization, Methodology, Data collection, Data analysis, Writing - original draft,

Writing - review & editing. CG-G: Conceptualization, Methodology, Writing - review & editing.

JDH: Conceptualization, Writing - review & editing, Funding acquisition. BL:

Conceptualization, Writing - review & editing, Funding acquisition. MB: Conceptualization,

Writing - review & editing, Funding acquisition.

Data and code Accessibility

The raw fMRI BIDS dataset is available in the following OpenNeuro repository:

https://openneuro.org/datasets/ds004829. The processed individual and group-level MRI

and behavioral data, together with the scripts used for data analyses, are available on OSF:

https://osf.io/dq6v5/

References

Aitken, F., Turner, G., & Kok, P. (2020). Prior Expectations of Motion Direction Modulate

Early Sensory Processing. *Journal of Neuroscience*, *40*(33), 6389–6397.

https://doi.org/10.1523/JNEUROSCI.0537-20.2020

Arco, J. E., González-García, C., Díaz-Gutiérrez, P., Ramírez, J., & Ruz, M. (2018).

Influence of activation pattern estimates and statistical significance tests in fMRI

decoding analysis. *Journal of Neuroscience Methods*, *308*, 248–260.

https://doi.org/10.1016/j.jneumeth.2018.06.017

Ariani, G., Pruszynski, J. A., & Diedrichsen, J. (2022). Motor planning brings human primary

somatosensory cortex into action-specific preparatory states. *ELife*, *11*, e69517.

https://doi.org/10.7554/ELIFE.69517

Ariani, G., Wurm, M. F., & Lingnau, A. (2015). Decoding Internally and Externally Driven

Movement Plans. *Journal of Neuroscience*, *35*(42), 14160–14171.

https://doi.org/10.1523/JNEUROSCI.0596-15.2015

Bourguignon, N. J., Braem, S., Hartstra, E., De Houwer, J., & Brass, M. (2018). Encoding of

Novel Verbal Instructions for Prospective Action in the Lateral Prefrontal Cortex:

Evidence from Univariate and Multivariate Functional Magnetic Resonance Imaging

Analysis. *Journal of Cognitive Neuroscience*, *30*(8), 1170–1184.

https://doi.org/10.1162/jocn

Brady, T. F., Konkle, T., Alvarez, G. A., & Oliva, A. (2013). Real-world objects are not

represented as bound units: Independent forgetting of different object details from

visual memory. *Journal of Experimental Psychology: General*, *142*(3), 791–808.

https://doi.org/10.1037/A0029649

Brass, M., Liefooghe, B., Braem, S., & De Houwer, J. (2017). Following new task

instructions: Evidence for a dissociation between knowing and doing. *Neuroscience &*

Biobehavioral Reviews, *81*, 16–28. https://doi.org/10.1016/j.neubiorev.2017.02.012

Brass, M., Wenke, D., Spengler, S., & Waszak, F. (2009). Neural correlates of overcoming

interference from instructed and implemented stimulus-response associations. *Journal*

of Neuroscience, *29*(6), 1766–1772. https://doi.org/10.1523/JNEUROSCI.5259-

08.2009

Braver, T. S. (2012). The variable nature of cognitive control: A dual-mechanisms

framework. *Trends in Cognitive Sciences*, *16*(2), 106–113.

https://doi.org/10.1016/j.tics.2011.12.010.The

Brodeur, M. B., Guérard, K., & Bouras, M. (2014). Bank of Standardized Stimuli (BOSS)

Phase II: 930 New Normative Photos. *PLOS ONE*, *9*(9), e106953.

https://doi.org/10.1371/JOURNAL.PONE.0106953

Christophel, T. B., Klink, P. C., Spitzer, B., Roelfsema, P. R., & Haynes, J.-D. (2017). The

Distributed Nature of Working Memory. *Trends in Cognitive Sciences*, *21*(2), 111–124.

https://doi.org/10.1016/j.tics.2016.12.007

Churchland, M. M., Cunningham, J. P., Kaufman, M. T., Ryu, S. I., & Shenoy, K. V. (2010).

Cortical Preparatory Activity: Representation of Movement or First Cog in a Dynamical

Machine? *Neuron*, *68*(3), 387–400. https://doi.org/10.1016/J.NEURON.2010.09.015

Cole, M. W., Bagic, A., Kass, R., & Schneider, W. (2010). Prefrontal Dynamics Underlying

Rapid Instructed Task Learning Reverse with Practice. *Journal of Neuroscience*,

30(42), 14245–14254. https://doi.org/10.1523/JNEUROSCI.1662-10.2010

Cole, M. W., Etzel, J. A., Zacks, J. M., Schneider, W., & Braver, T. S. (2011). Rapid Transfer

of Abstract Rules to Novel Contexts in Human Lateral Prefrontal Cortex. *Frontiers in*

Human Neuroscience, *5*, 142. https://doi.org/10.3389/fnhum.2011.00142

Cole, M. W., Laurent, P., & Stocco, A. (2013). Rapid instructed task learning: A new window

into the human brain's unique capacity for flexible cognitive control. *Cognitive, Affective*

and Behavioral Neuroscience, *13*(1), 1–22. https://doi.org/10.3758/s13415-012-0125-

7

Cole, M. W., Reynolds, J. R., Power, J. D., Repovs, G., Anticevic, A., & Braver, T. S. (2013).

Multi-task connectivity reveals flexible hubs for adaptive task control. *Nature*

Neuroscience, *16*(9), 1348–1355. https://doi.org/10.1038/nn.3470

Curtis, C. E., & D'Esposito, M. (2003). Persistent activity in the prefrontal cortex during
working memory. *Trends in Cognitive Sciences*, *7*(9), 415–423.
https://doi.org/10.1016/S1364-6613(03)00197-9

Demanet, J., Liefooghe, B., Hartstra, E., Wenke, D., De Houwer, J., & Brass, M. (2016).
There is more into 'doing' than 'knowing': The function of the right inferior frontal sulcus
is specific for implementing versus memorising verbal instructions. *NeuroImage*, *141*,
350–356. https://doi.org/10.1016/j.neuroimage.2016.07.059

Dumontheil, I., Thompson, R., & Duncan, J. (2011). Assembly and Use of New Task Rules
in Fronto-parietal Cortex. *Journal of Cognitive Neuroscience*, *23*(1), 168–182.
https://doi.org/10.1162/jocn.2010.21439

Duncan, J. (2010). The multiple-demand (MD) system of the primate brain: mental programs
for intelligent behaviour. *Trends in Cognitive Sciences*, *14*(4), 172–179.
https://doi.org/10.1016/j.tics.2010.01.004

Everaert, T., Theeuwes, M., Liefooghe, B., & De Houwer, J. (2014). Automatic Motor
Activation by Mere Instruction. *Cognitive, Affective and Behavioral Neuroscience*, *14*,
1300–1309. https://doi.org/10.3758/s13415-014-0294-7

Fedorenko, E., Duncan, J., & Kanwisher, N. (2013). Broad domain generality in focal regions
of frontal and parietal cortex. *Proceedings of the National Academy of Sciences of the*
United States of America, *110*(41), 16616–16621.
https://doi.org/10.1073/pnas.1315235110

Feuerriegel, D., Blom, T., & Hogendoorn, H. (2021). Predictive activation of sensory
representations as a source of evidence in perceptual decision-making. *Cortex*, *136*,
140–146. https://doi.org/10.1016/J.CORTEX.2020.12.008

Gale, D. J., Flanagan, J. R., & Gallivan, J. P. (2021). Human somatosensory cortex is

modulated during motor planning. *Journal of Neuroscience, 41*(27), 5909–5922.

https://doi.org/10.1523/JNEUROSCI.0342-21.2021

Gallivan, J. P., McLean, D. A., Flanagan, J. R., & Culham, J. C. (2013). Where One Hand

Meets the Other: Limb-Specific and Action-Dependent Movement Plans Decoded from

Preparatory Signals in Single Human Frontoparietal Brain Areas. *Journal of*

Neuroscience, 33(5), 1991–2008. https://doi.org/10.1523/JNEUROSCI.0541-12.2013

Gallivan, J. P., McLean, D. A., Valyear, K. F., Pettypiece, C. E., & Culham, J. C. (2011).

Decoding action intentions from preparatory brain activity in human parieto-frontal

networks. *Journal of Neuroscience, 31*(26), 9599–9610.

https://doi.org/10.1523/JNEUROSCI.0080-11.2011

Glasser, M. F., Coalson, T. S., Robinson, E. C., Hacker, C. D., Harwell, J., Yacoub, E.,

Ugurbil, K., Andersson, J., Beckmann, C. F., Jenkinson, M., Smith, S. M., & Van Essen,

D. C. (2016). A multi-modal parcellation of human cerebral cortex. *Nature, 536*(7615),

171–178. https://doi.org/10.1038/nature18933

González-García, C., Arco, J. E., Palenciano, A. F., Ramírez, J., & Ruz, M. (2017).

Encoding, preparation and implementation of novel complex verbal instructions.

NeuroImage, 148, 264–273. https://doi.org/10.1016/J.NEUROIMAGE.2017.01.037

González-García, C., Formica, S., Liefooghe, B., & Brass, M. (2020). Attentional

prioritization reconfigures novel instructions into action-oriented task sets. *Cognition,*

194, 104059. https://doi.org/10.1016/j.cognition.2019.104059

González-García, C., Formica, S., Wisniewski, D., & Brass, M. (2021). Frontoparietal action-

oriented codes support novel instruction implementation. *NeuroImage, 226*, 117608.

https://doi.org/10.1016/j.neuroimage.2020.117608

Gratton, G., Coles, M. G., Sirevaag, E. J., Eriksen, C. W., & et al. (1988). Pre- and
poststimulus activation of response channels: a psychophysiological analysis. *Journal*
of Experimental Psychology. Human Perception and Performance, *14*(3), 331–344.
https://doi.org/10.1037//0096-1523.14.3.331

Griffin, G., Holub, A., & Perona, P. (2007). *Caltech-256 Object Category Dataset*.
https://resolver.caltech.edu/CaltechAUTHORS:CNS-TR-2007-001

Grill-Spector, K., Kushnir, T., Edelman, S., Avidan, G., Itzchak, Y., & Malach, R. (1999).
Differential processing of objects under various viewing conditions in the human lateral
occipital complex. *Neuron*, *24*(1), 187–203. https://doi.org/10.1016/S0896-
6273(00)80832-6

Grush, R. (2004). The emulation theory of representation: Motor control, imagery, and
perception. *Behavioral and Brain Sciences*, *27*(3), 377–396.
https://doi.org/10.1017/S0140525X04000093

Guillot, A., & Collet, C. (2005). Contribution from neurophysiological and psychological
methods to the study of motor imagery. *Brain Research Reviews, 50*(2), 387–397.
https://doi.org/10.1016/j.brainresrev.2005.09.004

Haggard, P. (2008). Human volition: towards a neuroscience of will. *Nature Reviews*
Neuroscience, 9(12), 934–946. https://doi.org/10.1038/nrn2497

Hardwick, R. M., Caspers, S., Eickhoff, S. B., & Swinnen, S. P. (2018). Neural correlates of
action: Comparing meta-analyses of imagery, observation, and execution.
Neuroscience and Biobehavioral Reviews, 94, 31–44.
https://doi.org/10.1016/j.neubiorev.2018.08.003

Hartstra, E., Kühn, S., Verguts, T., & Brass, M. (2011). The implementation of verbal

instructions: An fMRI study. *Human Brain Mapping*, *32*(11), 1811–1824.

https://doi.org/10.1002/hbm.21152

Hartstra, E., Waszak, F., & Brass, M. (2012). The implementation of verbal instructions:

Dissociating motor preparation from the formation of stimulus-response associations.

NeuroImage, *63*(3), 1143–1153. https://doi.org/10.1016/j.neuroimage.2012.08.003

Haxby, J. V., Connolly, A. C., & Guntupalli, J. S. (2014). Decoding Neural Representational

Spaces Using Multivariate Pattern Analysis. *Annual Review of Neuroscience*, *37*(1),

435–456. https://doi.org/10.1146/annurev-neuro-062012-170325

Haynes, J. D., & Rees, G. (2006). Decoding mental states from brain activity in humans.

Nature Reviews Neuroscience, 7, 523–534. https://doi.org/10.1038/nrn1931

Hétu, S., Grégoire, M., Saimpont, A., Coll, M. P., Eugène, F., Michon, P. E., & Jackson, P.

L. (2013). The neural network of motor imagery: An ALE meta-analysis. *Neuroscience*

and Biobehavioral Reviews, 37(5), 930–949.

https://doi.org/10.1016/j.neubiorev.2013.03.017

Hommel, B. (2009). Action control according to TEC (theory of event coding). *Psychological*

Research, *73*(4), 512–526. https://doi.org/10.1007/s00426-009-0234-2

Hommel, B., Müsseler, J., Aschersleben, G., & Prinz, W. (2001). The Theory of Event

Coding (TEC): A framework for perception and action planning. *Behavioral and Brain*

Sciences, *24*(5), 849–878. https://doi.org/10.1017/S0140525X01000103

Jeannerod, M. (1994). The representing brain: neural correlates of motor intention and

imagery. *Behavioral and Brain Sciences*, *17*(2), 187–245.

https://doi.org/10.1017/S0140525X00034026

Jeannerod, M. (2001). Neural simulation of action: A unifying mechanism for motor
cognition. *NeuroImage*, *14*(1), S103–S109. https://doi.org/10.1006/nimg.2001.0832

Kok, P., Mostert, P., & De Lange, F. P. (2017). Prior expectations induce prestimulus
sensory templates. *Proceedings of the National Academy of Sciences of the United*
States of America, *114*(39), 10473–10478. https://doi.org/10.1073/pnas.1705652114

Konkle, T., Brady, T. F., Alvarez, G. A., & Oliva, A. (2010). Conceptual distinctiveness
supports detailed visual long-term memory for real-world objects. *Journal of*
Experimental Psychology: General, *139*(3), 558–578.
https://doi.org/10.1037/A0019165

Kriegeskorte, N., & Kievit, R. A. (2013). Representational geometry: Integrating cognition,
computation, and the brain. *Trends in Cognitive Sciences, 17*(8), 401–412.
https://doi.org/10.1016/j.tics.2013.06.007

Kriegeskorte, N., Mur, M., & Bandettini, P. (2008). Representational similarity analysis –
connecting the branches of systems neuroscience. *Frontiers in Systems Neuroscience*,
2, 4. https://doi.org/10.3389/neuro.06.004.2008

Liefooghe, B., & De Houwer, J. (2018). Automatic effects of instructions do not require the
intention to execute these instructions. *Journal of Cognitive Psychology*, *30*(1), 108–
121. https://doi.org/10.1080/20445911.2017.1365871

Liefooghe, B., De Houwer, J., & Wenke, D. (2013). Instruction-based response activation
depends on task preparation. *Psychonomic Bulletin and Review*, *20*(3), 481–487.
https://doi.org/10.3758/s13423-013-0374-7

Liefooghe, B., Jim, A., & De Houwer, J. (2021). Automatic effects of covert practice.

Quarterly Journal of Experimental Psychology, *74*(10), 1697-1708.

https://doi.org/10.1177/17470218211007138

Liefooghe, B., Wenke, D., & De Houwer, J. (2012). Instruction-based task-rule congruency

effects. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, *38*(5),

1325–1335. https://doi.org/10.1037/a0028148

Mayka, M. A., Corcos, D. M., Leurgans, S. E., & Vaillancourt, D. E. (2006). Three-

dimensional locations and boundaries of motor and premotor cortices as defined by

functional brain imaging: a meta-analysis. *NeuroImage*, *31*(4), 1453–1474.

https://doi.org/10.1016/J.NEUROIMAGE.2006.02.004

Meiran, N., & Cohen-Kdoshay, O. (2012). Working memory load but not multitasking

eliminates the prepared reflex: Further evidence from the adapted flanker paradigm.

Acta Psychologica, *139*(2), 309–313. https://doi.org/10.1016/j.actpsy.2011.12.008

Meiran, N., Pereg, M., Kessler, Y., Cole, M. W., & Braver, T. S. (2015a). Reflexive activation

of newly instructed stimulus–response rules: evidence from lateralized readiness

potentials in no-go trials. *Cognitive, Affective and Behavioral Neuroscience*, *15*(2),

365–373. https://doi.org/10.3758/s13415-014-0321-8

Meiran, N., Pereg, M., Kessler, Y., Cole, M. W., & Braver, T. S. (2015b). The power of

instructions: Proactive configuration of stimulus–response translation. *Journal of*

Experimental Psychology: Learning, Memory, and Cognition, *41*(3), 768–786.

https://doi.org/10.1037/xlm0000063

Miller, E. K., & Cohen, J. D. (2001). An Integrative Theory of Prefrontal Cortex Function.

Annual Review of Neuroscience, *24*(1), 167–202.

https://doi.org/10.1146/annurev.neuro.24.1.167

Moran, A., & O'Shea, H. (2020). Motor Imagery Practice and Cognitive Processes. *Frontiers*

in Psychology, *11*: 394. https://doi.org/10.3389/fpsyg.2020.00394

Muhle-Karbe, P. S., Duncan, J., De Baene, W., Mitchell, D. J., & Brass, M. (2017). Neural

Coding for Instruction-Based Task Sets in Human Frontoparietal and Visual Cortex.

Cerebral Cortex, *27*(3), 1891–1905. https://doi.org/10.1093/cercor/bhw032

Mumford, J. A., Davis, T., & Poldrack, R. A. (2014). The impact of study design on pattern

estimation for single-trial multivariate pattern analysis. *NeuroImage*, *103*, 130–138.

https://doi.org/10.1016/J.NEUROIMAGE.2014.09.026

Mumford, J. A., Turner, B. O., Ashby, F. G., & Poldrack, R. A. (2012). Deconvolving BOLD

activation in event-related designs for multivoxel pattern classification analyses.

NeuroImage, *59*, 2636–2643. https://doi.org/10.1016/j.neuroimage.2011.08.076

Palenciano, A. F., González-García, C., Arco, J. E., Pessoa, L., & Ruz, M. (2019).

Representational organization of novel task sets during proactive encoding. *Journal of*

Neuroscience, *39*(42) 8386-8397. https://doi.org/10.1523/JNEUROSCI.0725-19.2019

Palenciano, A. F., González-García, C., Arco, J. E., & Ruz, M. (2019). Transient and

Sustained Control Mechanisms Supporting Novel Instructed Behavior. *Cerebral*

Cortex, *29*(9), 3948–3960. https://doi.org/10.1093/cercor/bhy273

Palenciano, A. F., González-García, C., De Houwer, J., Brass, M., & Liefooghe, B. (2021).

Exploring the link between novel task proceduralization and motor simulation. *Journal*

of Cognition, *4*(1). https://doi.org/10.5334/JOC.190/METRICS/

Palenciano, A. F., Senoussi, M., Formica, S., & González-García, C. (2023). Canonical

template tracking: Measuring the activation state of specific neural representations.

Frontiers in Neuroimaging, *1*, 52. https://doi.org/10.3389/FNIMG.2022.974927

Peirce, J., Gray, J. R., Simpson, S., MacAskill, M., Höchenberger, R., Sogo, H., Kastman,

E., & Lindeløv, J. K. (2019). PsychoPy2: Experiments in behavior made easy. *Behavior*

Research Methods, *51*(1), 195–203. https://doi.org/10.3758/s13428-018-01193-y

Possidónio, C., Graça, J., Piazza, J., & Prada, M. (2019). Animal Images Database:

Validation of 120 Images for Human-Animal Studies. *Animals*, *9*(8), 475.

https://doi.org/10.3390/ani9080475

Rizzolatti, G., & Luppino, G. (2001). The Cortical Motor System. *Neuron*, *31*(6), 889–901.

https://doi.org/10.1016/S0896-6273(01)00423-8

Ruge, H., & Wolfensteller, U. (2010). Rapid formation of pragmatic rule representations in

the human brain during instruction-based learning. *Cerebral Cortex*, *20*(7), 1656–1667.

https://doi.org/10.1093/cercor/bhp228

Russo, N., Hagmann, C. E., Andrews, R., Black, C., Silberman, M., & Shea, N. (2018).

Validation of the C.A.R.E. stimulus set of 640 animal pictures: Name agreement and

quality ratings. *PLOS ONE*, *13*(2), e0192906.

https://doi.org/10.1371/journal.pone.0192906

Shenoy, K. V., Sahani, M., & Churchland, M. M. (2013). Cortical Control of Arm Movements:

A Dynamical Systems Perspective. *Annual Review of Neuroscience, 36,* 337–359.

https://doi.org/10.1146/ANNUREV-NEURO-062111-150509

Shin, Y. K., Proctor, R. W., & Capaldi, E. J. (2010). A Review of Contemporary Ideomotor

Theory. *Psychological Bulletin*, *136*(6), 943–974. https://doi.org/10.1037/A0020541

Smulders, F. T. Y., & Miller, J. O. (2011). The Lateralized Readiness Potential. *The Oxford*

Handbook of Event-Related Potential Components.

https://doi.org/10.1093/OXFORDHB/9780195374148.013.0115

Sreenivasan, K. K., Curtis, C. E., & D'Esposito, M. (2014). Revisiting the role of persistent

neural activity during working memory. *Trends in Cognitive Sciences*, *18*(2), 82–89.

https://doi.org/10.1016/j.tics.2013.12.001

Theeuwes, M., Liefooghe, B., De Schryver, M., & De Houwer, J. (2018). The role of motor

imagery in learning via instructions. *Acta Psychologica*, *184*, 110–123.

https://doi.org/10.1016/j.actpsy.2017.05.002

Walther, A., Nili, H., Ejaz, N., Alink, A., Kriegeskorte, N., & Diedrichsen, J. (2016). Reliability

of dissimilarity measures for multi-voxel pattern analysis. *NeuroImage*, *137*, 188–200.

https://doi.org/10.1016/j.neuroimage.2015.12.012

Waszak, F., Cardoso-Leite, P., & Hughes, G. (2012). Action effect anticipation:

Neurophysiological basis and functional consequences. *Neuroscience & Biobehavioral*

Reviews, *36*(2), 943–959. https://doi.org/10.1016/J.NEUBIOREV.2011.11.004

Wimber, M., Alink, A., Charest, I., Kriegeskorte, N., & Anderson, M. C. (2015). Retrieval

induces adaptive forgetting of competing memories via cortical pattern suppression.

Nature Neuroscience, *18*(4), 582–589. https://doi.org/10.1038/nn.3973

Zabicki, A., De Haas, B., Zentgraf, K., Stark, R., Munzert, J., & Krüger, B. (2017). Imagined

and executed actions in the human motor system: Testing neural similarity between

execution and imagery of actions with a multivariate approach. *Cerebral Cortex*, *27*(9),

4523–4536. https://doi.org/10.1093/cercor/bhw257

1129

Extended data

Figure 5-1

Results from the Wilcoxon Signed-Ranked tests using the motor response and action effects templates.

	Motor response localizer									Action effects localizer								
	Relevant vs. irrelevant			*Relevant against 0*			*Irrelevant against 0*			*Relevant vs. irrelevant*			*Relevant against 0*			*Irrelevant against 0*		
	z	p	p_{corr}	z	p	p_{corr}	z	p	p_{corr}	z	p	p_{corr}	z	p	p_{corr}	z	p	p_{corr}
leftM1	4.78	<.001	<.001	4.72	<.001	<.001	-4.68	<.001	<.001	1.93	.026	.056	3.22	.001	.002	0.19	.848	1
rightM1	4.78	<.001	<.001	4.70	<.001	<.001	-4.78	<.001	<.001	4.02	<.001	<.001	4.40	<.001	<.001	-2.55	.011	.188
leftS1	4.78	<.001	<.001	4.76	<.001	<.001	-4.00	<.001	<.001	3.54	<.001	<.001	3.63	<.001	<.001	-1.80	.072	.201
rightS1	4.69	<.001	<.001	4.72	<.001	<.001	-4.47	<.001	<.001	4.43	<.001	<.001	4.47	<.001	<.001	-3.46	<.001	.002
SMA	4.77	<.001	<.001	4.43	<.001	<.001	-2.05	.041	.057	3.52	<.001	<.001	3.92	<.001	<.001	0.77	.442	1
preSMA	0.45	.325	.678	0.57	.572	.507	0.85	.393	1	2.01	.022	.056	1.27	.203	.258	-0.02	.980	1
PMv	0.91	.183	.678	3.40	<.001	.001	3.75	<.001	<.001	0.76	.225	.210	2.23	.025	.083	1.11	.269	1
PMd	4.26	<.001	<.001	4.45	<.001	<.001	-0.69	.491	1	3.71	<.001	<.001	3.92	<.001	<.001	0.82	.414	1
IPL	3.17	<.001	<.001	3.03	.002	.002	-0.24	.813	1	3.50	<.001	<.001	3.00	.003	.005	-0.62	.532	1
SPL	3.78	<.001	<.001	3.51	<.001	<.001	-1.55	.120	.243	2.68	.003	.003	3.82	<.001	<.001	1.61	.107	.512
pInsula	4.67	<.001	<.001	3.90	<.001	<.001	-2.52	.012	.037	4.36	<.001	<.001	3.24	.001	.006	-3.46	<.001	.002
V1	-0.82	.795	.825	-1.45	.147	.357	-0.79	.428	1	-0.28	.609	.684	1.13	.258	.394	1.58	.113	.686

Extended data

Figure 5-2

Results of the posthoc comparisons from the repeated measures ANOVA including the motor response and action effects templates simultaneously.

	Motor response			Action effects		
	z	p	p_{corr}	z	p	p_{corr}
left M1	4.53	<.001	<.001	2.51	.006	.011
right M1	4.53	<.001	<.001	4.23	<.001	<.001
left S1	4.53	<.001	<.001	3.57	<.001	<.001
right S1	4.43	<.001	<.001	4.26	<.001	<.001
SMA	4.53	<.001	<.001	3.33	<.001	<.001
preSMA	0.80	.210	.707	2.49	.006	.018
PMv	0.88	.190	.707	0.80	.210	.648
PMd	4.05	<.001	<.001	3.40	<.001	<.001
IPL	3.23	<.001	<.001	3.40	<.001	<.001
SPL	3.59	<.001	.001	2.97	.002	.004
pInsula	4.43	<.001	<.001	3.98	<.001	<.001
V1	-2.05	.775	.816	-1.36	.097	.157

Extended data

Figure 5-3

Results from the Wilcoxon Signed-Ranked tests using the stimulus category templates.

	Relevant vs. irrelevant			Relevant against 0			Irrelevant against 0		
	z	p	p_{corr}	z	p	p_{corr}	z	p	p_{corr}
V1	3.27	<.001	<.001	0.75	.452	.707	-2.73	.006	.012
V2-4	4.55	<.001	<.001	-1.31	.192	1	-4.64	<.001	<.001
LOC	4.77	<.001	<.001	-2.97	.039	.673	-4.78	<.001	<.001
VentralVisual	4.67	<.001	<.001	-1.24	.213	1	-4.78	<.001	<.001
M1	-0.88	.812	.590	1.08	.280	.157	1.32	.185	.252

Extended data

Figure 5-4. Boxplots showing the averaged beta weights for relevant and irrelevant canonical templates capturing the mappings' response effector information. **A,B.** Results when only [Left Index, Right Index] and [Left Middle, Right Middle] trials are included in the analysis, using the motor response (**A**) and action effects (**B**) templates. **C,D.** Results using activity patterns from the M1 and S1 ROIs contralateral to the mappings' and canonical templates' laterality condition, using data from the motor response (**C**) and action effects (**D**) localizer.

Extended data

Figure 5-5. Bar plots displaying the averaged beta weight for relevant and irrelevant motor response templates, across mappings' conditions (shown in the X axes) and ROIs. These data were shown for illustrative purposes only. Hence, no statistical information is represented in this figure.

Extended data

Figure 5-6. Bar plots displaying the averaged beta weight for relevant and irrelevant action effects templates, across mappings' conditions (shown in the X axes) and ROIs. These data were shown for illustrative purposes only. Hence, no statistical information is represented in this figure.

Extended data

Figure 5-7. Bar plots displaying the averaged beta weight for relevant and irrelevant stimulus category templates, across mappings' conditions (shown in the X axes) and ROIs. These data were shown for illustrative purposes only. Hence, no statistical information is represented in this figure.

Figure 5-8. Canonical templates' reliability measurements (SNR, correlationability) in the motor response (A), action effects (B), and stimuli category (C) localizers, across ROIs. Asterisks denote statistically significant differences (*p* < .05).