1

## Perilipin membrane-association determines
## lipid droplet heterogeneity in differentiating adipocytes

Mario Majchrzak[2], Ozren Stojanovic[1], Dalila Ajjaji[3], Kalthoum Ben M'barek[3], Mohyeddine Omrane[3], Abdou Rachid Thiam[3], Robin W. Klemm[1,2,4,5]

[1]Department of Physiology Anatomy and Genetics, University of Oxford, Oxford OX1 3PT, UK
[2]Department of Molecular Life Sciences, University of Zurich, 8057 Zurich, CH
[3] Laboratoire de Physique de l'École Normale Supérieure, ENS, Université PSL, CNRS, Sorbonne Université, Université de Paris, F-75005 Paris, France
[4]Correspondence : Robin.Klemm@dpag.ox.ac.uk
[5]Lead contact

**Summary**

The storage of fat within lipid droplets (LDs) of adipocytes is critical for whole-body health. Acute fatty acid uptake by adipocytes leads to the formation of at least two LD-classes marked by distinct members of the perilipin (PLIN) protein-family. How LD-heterogeneity arises is an important yet unresolved cell biological problem. Here, we show that an unconventional integral membrane-segment targets the adipocyte specific LD-surface factor PLIN1 to the endoplasmic reticulum (ER) and binds with high affinity to the first LD-class. The other PLINs do not insert into the ER and remain cytoplasmic until fatty acid influx recruits them to a newly forming LD-population. Preventing ER-localization turns PLIN1 into a cytoplasmic LD-protein, reduces its LD-affinity and switches its LD-class specificity. Our results explain how differences in organelle targeting and disparities in lipid-affinity of LD-surface factors establish LD-heterogeneity. These findings have direct implications for mechanism-based interference with fatty-liver disease, type-2 diabetes and obesity.

**Introduction**

Adipocytes deposit an excess of dietary nutrients as triacylglycerols (TAGs) in large lipid droplets (LDs) and release fatty acids to feed other tissues during fasting[1–3]. Deficiencies in LD biogenesis and turnover, or defects in factors important for LD surface assembly are associated with several metabolic diseases such as lipodystrophies, fatty liver disease, type-2 diabetes, and obesity[4–6].

LDs form at the membranes of the endoplasmic reticulum (ER) when neutral lipids, such as TAGs, accumulate between the two leaflets of the lipid bilayer[7–9]. With increasing TAG amounts, they phase-separate from the phospholipids in the membrane to form lenses or TAG-inclusions, which eventually bud out into the cytoplasm giving rise to LDs. As a result, the LD-surface is surrounded by a phospholipid monolayer and a set of LD specific surface proteins[10–14].

Early models in the field suggested that all LDs within a cell have the same lipid and protein composition and carry out similar functions. Accumulating evidence suggests, however, that LDs are highly heterogenous, and can specialize into different sub-populations which are morphologically, chemically, and functionally distinct[14–21].

One way by which cells can generate different LD-classes is determined by the pathways that synthesize the lipids in the hydrophobic core of the LD. For example, fly fat bodies contain a population of large LDs in the center of the cell, which is mostly filled with TAGs derived from *de novo* lipogenesis, i.e. the synthesis of fatty acids from non-lipid precursors such as glucose[22,23]. A class of smaller LDs in the periphery of the cell seems more specialized in storing TAGs made from fatty acids that are taken up from the outside[22,23]. Consistent with the work in flies, fatty acid feeding of mammalian adipocytes has long been known to lead to the formation of two distinct LD-classes[10,24,25]. As in flies the central LD population is formed mainly by *de novo* lipogenesis [26], whereas acute fatty acid influx appears to trigger the formation of a second LD class in the periphery of the cell[10,24,25]. However, the mechanisms that lead to the formation of different LD populations remain poorly understood, and current models of LD-biogenesis do not explain how LD-surface factors can be targeted to a specific LD population in the cell.

In general, protein targeting to LDs occurs by two major pathways[14,27,28]. Class I (or ERTOLD) proteins are synthesized as monotopic integral membrane proteins at the ER, and subsequently move to the LD surface (**Fig. 1A**, class I). The dual localization to the ER and the LDs is facilitated by special membrane anchors that often appear to fold into hairpin-like conformations and are characterized by signature residues such as internal prolines and tryptophans as well as positively charged residues[13,28].
Class II (or CYTOLD) proteins, on the other hand, bind directly from the cytoplasm to the LD surface (**Fig. 1A,** class II). They do not possess integral membrane segments and are instead targeted to the LD by amphipathic domains that detect packing defects in the LD monolayer where the hydrophobic LD core is exposed to the cytoplasm[14,29–34].

The perilipins (PLINs) are currently viewed as the prototypical class II proteins, comprising a family of evolutionarily conserved LD-surface factors carrying out important structural and regulatory functions[25,35–38]. The family defining N-terminal PAT domain and the following 11-mers both contain amphipathic regions, which are critical for LD-targeting (**Fig. S1A**)[31,36,39,40]. All five perilipin family members are thought to target the LD directly from the cytoplasm and should

therefore freely mix on the same LD class. In contrast to this expectation, the LD sub-populations
that form in adipocytes upon fatty acid feeding are marked by specific PLINs (**Fig. 1B**) [10,24,25]. An
affinity-based LD binding "hierarchy" among the different PLIN-family members is likely key to
the formation of the two LD classes[33,39,41–44]. However, the molecular mechanisms leading to
class-specific localization of LD surface proteins are not well understood, and the biochemical
characterization of the responsible targeting domains remains incomplete[37].

**Results**

**Acute fatty acid feeding induces the formation of distinct LDs in adipocytes**

To investigate how perilipins (PLINs) target LDs in adipocytes, we differentiated 3T3-L1
cells into white adipocytes and visualized endogenously expressed PLIN1, 2, 3, and 4 after fatty
acid feeding by immunofluorescence (PLIN5 is not expressed in 3T3-L1 adipocytes [11,26]). Our data
showed that adipocytes form two separate LD populations under these conditions (**Fig. 1B-F**)
confirming previous observations[25,35,38,39,41]. The LDs in the centre of the cell were almost
exclusively labelled with PLIN1, and the second population localizing more to the periphery was
preferentially bound by PLIN2, 3, and 4 (Fig. 1C, D, and E).  While the central population was
specifically marked by PLIN1 (Fig. 1 C-F), the peripheral LD population was more heterogenous,
and was either exclusively marked by PLIN2, 3, and 4, or additionally contained some PLIN1,
which we therefore classified as a mixed LD population (**Fig. 1C-E** (zoom-ins), **F** (MIX)).

Consistent with our results in adipocytes and the established LD binding hierarchy in other
cells[25,42], we next showed that ectopically expressed PLIN1 excluded PLIN2, 3, or 4 from LDs when
co-transfected in COS7 cells (**Fig. S1A-D**). As predicted, the high LD-affinity of PLIN1 was
dependent on features C-terminally of the 11-mer domain (**Fig. S1A**). A construct only consisting
of the N terminal PAT and 11-mer regions (PLIN1 1-192) was not able to exclude PLIN2 from the
LD surface (**Fig. S1E**). In fact, in most of the cells expressing both constructs, PLIN2 excluded the
PLIN1 truncation mutant (PLIN1 1-192) from the LD surface indicating higher affinity (**Fig. S1E**).
Together these data raised the question of how class-II LD proteins can separate onto two
different LD-populations within the same cell, and which domains in PLIN1 are required for this
behaviour.

**PLIN1 is an integral membrane protein dually localizing to the ER and LDs**

To determine the molecular basis of LD targeting by the PLINs, we examined PLIN1 and 2
by biochemistry and microscopy experiments expressing them alone in otherwise untreated
COS7 cells. As expected, PLIN1 localized to endogenous LDs but was, surprisingly, never seen in
the cytoplasm. Instead, it always bound to the ER (**Fig. 2A, Fig. S2B**). In agreement with data from
others[44], our results indicated that PLIN1 does not behave as a prototypical class-II LD protein.
PLIN2, however, showed normal class-II behaviour accumulating in the cytoplasm upon
overexpression and covering all endogenous LDs (**Fig. 2B, S2A, S2D**). After addition of fatty acid
(Na$^+$ oleate), both PLIN1 and PLIN2 targeted efficiently to LDs (**Fig. 2C, and D**). Only around 10 %
of the cells showed some residual ER-staining for PLIN1 (**Fig. 2E**).

Since PLIN1 does not contain any predicted membrane segments (**Fig. S2D, S2E**), and
exhibits similar overall hydrophobicity as PLIN2 (**Fig. S2F, S2G**), we initially expected that it would
bind the ER as a peripheral membrane protein or would associate with membranes through a
receptor in the ER. In support of this view, PLIN1 sedimented both as an untagged and GFP-fusion
protein together with the endogenously expressed integral ER membrane protein calnexin (CNX)
to the membrane pellet (**Fig. 2F**, PLIN1 untagged; GFP-PLIN1, P100, consistent with ER in[44]).
However, when we subjected the membrane pellet (P100) to alkaline extraction with $Na_2CO_3$-
buffer, PLIN1 quantitatively remained in the alkaline resistant membrane fraction (**Fig. 2F**, P100E)
behaving like an integral membrane protein. PLIN2, on the other hand was absent from the
membrane fractions and as expected for a canonical class-II protein, it co-fractionated with the
soluble cytoplasmic marker glyceraldehyde-3-phosphate dehydrogenase (GAPDH, **Fig. 2F**, S100).
To exclude possible artefacts, we additionally carried out flotation experiments. To this
end, we prepared a ribosome stripped membrane fraction from cells expressing PLIN1 and
loaded it to the bottom of a sucrose step gradient. As shown in **Fig. 2G**, PLIN1 moved efficiently
out of the bottom fraction and floated together with ER membranes to an interface fraction
marked by CNX (5-35% sucrose). When we solubilized the membrane pellet with detergent,
neither PLIN1 nor CNX floated (**Fig. 2H**). Finally, PLIN1 and CNX both remained soluble when we
mixed the cell lysate with detergent before high-speed centrifugation (**Fig. 2I**).
Taken together, the imaging and biochemical data indicated that PLIN1 is an integral ER
membrane protein and does not belong to the cytoplasmic class-II proteins. The targeting
pathway to the LD might therefore be more complex than currently envisaged, probably involving
the ER as an intermediate step. In support of this notion, we found that PLIN1 moved efficiently
onto LDs when we blocked translation with cycloheximide 1h before LD induction (**Fig. S2H-J**).
The ER-pool re-distributed efficiently to the LDs (**Fig. 2J**).
152

**PLIN1 contains an unconventional integral membrane segment and a peripheral membrane binding motif**

To explore the molecular basis of the subcellular localization of PLIN1 in more detail, we
next aimed at identifying the domains that are required for membrane targeting. We, therefore,
carried out a systematic structure-function analysis using fluorescence microscopy and
subcellular fractionation followed by alkaline extraction to probe for membrane integration.
We started the analysis with a minimal N-terminus of the mouse PLIN1 sequence (PLIN1
1-192) containing the PAT and 11-mer regions that were previously shown to be sufficient for LD
targeting (**Fig. S3A**)[31,39,42]. As expected, neither of these two domains nor a polybasic region
enriched in prolines downstream of the 11-mer region was involved in ER targeting. In fact, the
N-terminal PLIN1 portion until position 237 remained completely soluble (**Fig. 3A**, S100) and was
found in the cytoplasm as well as on the endogenous LDs (**Fig. 3B, S3B**).
However, a construct extended until residue 281 (PLIN1 1-281) pelleted efficiently to the
membrane fraction P100, and was alkaline resistant (**Fig. 3A**, P100E). This construct also localized
to the ER as determined by microscopy, indicating that the region between residues 238 and 280
contained an integral membrane segment (iMS) (**Fig. 3A** green domain, and **S3B**). Interestingly,
the corresponding C-terminal portion (PLIN1 281-517) was found in the supernatant S100 (**Fig.
3A**) but localized to the ER as shown by microscopy (**Fig. 3D**), suggesting that PLIN1 contains an

additional peripheral membrane segment (pMS) that contributes to ER binding. Computational
analysis of the C-terminal portion predicted a strong amphipathic region in a segment between
residues 380 and 400 (**Fig. 3A**, yellow). However, a construct containing this motif (Plin1 371-517)
remained completely soluble in the cytoplasm (**Fig. 3A, and 3E**). Further extension of the C-
terminal part just until the beginning of the 4-helix bundle (PLIN1 328-517) led to ER localization
(**Fig. 3F**), but this construct remained soluble in the fractionation experiments (**Fig. 3A**), indicating
that the potential pMS is positioned in this first section of the middle domain (**Fig. 3A**, orange).
To narrow down the two membrane binding regions further, we turned to hydropathy
plot analysis (**Fig. S2F**). A first segment with moderate hydrophobicity was found between
tryptophan 238 and proline 280 of the mouse PLIN1 sequence, corresponding to the region in
which we identified the iMS (**Fig. S2F and 3G**). Although slightly less hydrophobic than the first
segment, a second region between residues 348 and 370 coincided with the peripheral
membrane segment (pMS, **Fig. S2F and 3G**).
To test the function of these domains in ER targeting we deleted them from the PLIN1
sequence. PLIN1ΔΔ lacking both the iMS and the pMS (**Fig. 3G**, PLIN1 Δ238-280, Δ348-370) was
completely soluble, and showed no membrane association after high-speed membrane
sedimentation or in the flotation experiments (**Fig. 3H**). Additionally, this mutant localized to the
cytoplasm in all cells that expressed the construct (**Fig. 3I**). Although having lost its ability to bind
the ER, PLIN1ΔΔ was still found on LDs after fatty acid feeding, but did not exclude PLIN2
anymore, which was now often found on the same LDs (**Fig. 3J**). In conclusion, removal of the
iMS and the pMS converted PLIN1 into a class II LD-protein, targeting LDs directly from the
cytoplasm (**Fig. 3K**).
193
In agreement with these results, addition of the iMS to PLIN2, right downstream of the
11-mer region (PLIN2-iMS (insertion at position 251)), was sufficient for membrane integration
(**Fig S3C, S3D, S3E**). ER-insertion of PLIN2-iMS simultaneously increased LD affinity, and permitted
efficient competition with PLIN1 for LD formation at the ER (**Fig. S3E**). However, PLIN2-iMS did
not mix with PLIN1 on the same LDs, but tended to segregate onto a separate LD class (**Fig. S3E**),
indicating a critical function for the iMS in the nucleation of different LD populations at the ER.
200
Taken together our data show that PLIN1 behaves as an integral ER membrane protein
that can move to LDs showing characteristic of a class-I rather than a class-II LD-protein. Both the
iMS and the pMS are sufficient for ER targeting, but membrane integration, and the generation
of different LD-classes depend on the iMS. Removing the ability to integrate into the ER converts
PLIN1 into a conventional class II LD-protein. The ER-binding deficient PLIN1-mutant mixes better
with class II-PLINs on the same LDs indicating that it has reduced LD affinity.

**The iMS is critical in determining PLIN1 LD-binding kinetics**

To investigate this possibility biochemically, we analysed the monolayer affinity of PLIN1 and PLIN1ΔΔ *in vitro*, using an assay that mimics the LD surface (**Fig. 4A**,[30]). Shrinkage of the aqueous compartments in this buffer-in-oil system leads to compression of the artificial LD-monolayer at the oil-buffer interface. The proteins fall off the monolayer with progressive reduction of the interface area and accumulate proportionally to their off-rates in the buffer compartment[30]. In agreement with the microscopy and fractionation experiments (**Fig. 3G-K**, and [39,41]), full-length PLIN1 remained efficiently bound to the interface of the buffer compartment, indicating that the high LD affinity is determined by low off-rates from the monolayer (**Fig. 4, B**). In comparison, PLIN1ΔΔ readily fell off the interface when the aqueous phase began to shrink, showing faster off-rates and lower affinity to the monolayer than full-length PLIN1 (**Fig. 4A, B**). The progressive accumulation in the buffer phase with increasing monolayer compression further showed that PLIN1 excluded PLIN1ΔΔ from the LD surface mimic, confirming the results in cells (**Fig. 4A, B**).

To gain insights into the corresponding on-reaction of the binding equilibrium, we used fluorescence recovery after photobleaching experiments in cells. Deletion of either the iMS (PLIN1ΔiMS) alone or together with the pMS (PLIN1ΔΔ) made recovery on the LD faster by more than an order of magnitude compared to a mutant in which the iMS is present and only the pMS is removed (PLIN1ΔpMS) (**Fig. 4C**, still frames of time series **Fig. S4A**). Together these results suggest that PLIN1 binds LDs with both low off and on rates, which is determined by the iMS. LD-binding and exchange via the classical amphipathic segments on the other hand appears to be faster, but results, overall, in lower LD-affinity.

These data suggested that differences in membrane targeting and LD-affinity by the PLINs may provide a simple explanation for the formation of the two LD populations in adipocytes: PLIN1 is targeted to the ER via the iMS where it binds to the emerging LDs first, excluding the low-affinity PLINs from the surface of this LD class. However, due to its high affinity, PLIN1 is "trapped" on this first LD population and is therefore not able to re-distribute fast enough to the second LD-class which acutely appears upon fatty acid feeding. Since PLIN2, 3 and 4 are excluded from the LDs that are covered by PLIN1, they remain available in the cytoplasm and can readily bind to the surface of these newly synthesized LDs. Despite their lower LD-affinity, the class-II PLINs would label the acutely emerging LD-population first.

**Perturbation of PLIN1 membrane targeting leads to a LD-class switch in adipocytes**

To test these interpretations of the *in vitro* experiments directly in adipocytes, we stably integrated GFP-PLIN1 and GFP-PLIN1ΔΔ as inducible transgenes into our preadipocytes (3T3-L1 cells). Consistent with the results above, PLIN1ΔΔ remained soluble in fractionation and alkaline extraction experiments (**Fig. 4D**), and full-length PLIN1 indeed behaved like an integral membrane protein sedimenting to the alkaline resistant membrane pellet, marked here by the endogenously expressed ER membrane protein CLIMP63 (**Fig. 4D and 4E**). PLIN1 localized to LDs after fatty acid feeding and as in COS7 cells moved from the ER to LDs even when translation was shut off with cycloheximide before LD induction (**Fig. S4B-E**).

When we induced expression of the transgenes in adipocytes, 6 days post differentiation, PLIN1ΔΔ behaved like a class-II LD protein mostly remaining cytoplasmic and being largely

excluded from the big LDs which formed during differentiation (**Fig. 4E and F**). Flotation
experiments with extracts from the same adipocytes revealed that endogenous PLIN1 was
expressed and floated with the LDs to the top fraction (**Fig. 4G**). PLIN1ΔΔ, was additionally found
abundantly in the soluble middle fraction, confirming the microscopy experiments (**Fig. 4E and**
**F**). Although being effectively excluded from the LDs covered by endogenous PLIN1, PLIN1ΔΔ
remains LD binding competent in cells and *in vitro* (**Fig. 3J, Fig. 4A**). PLIN1ΔΔ probably gained
access to the LD core via its amphipathic PAT domain when the LD surface was disrupted during
preparation of the cell extracts by mechanical lysis and was therefore also seen in the LD fraction
on top of the gradient (**Fig. 4G**).
Importantly, PLIN1ΔΔ efficiently moved from the cytoplasm to the peripheral LD
population which forms after fatty acid feeding (**Fig. 4H and F**). Consistent with the results shown
for PLIN2-4 in **Fig. 1**, PLIN1ΔΔ remained largely excluded from the LDs that were covered by
endogenously expressed full-length PLIN1 (**Fig. 4H and F**).
271

272         Taken together, these data suggest that removing the iMS and pMS not only reduced LD
273     affinity and turned PLIN1ΔΔ into a class-II LD-protein, but also leads to a change in LD-class
274     specificity. Moreover, these results are consistent with a model by which the high LD affinity of
275     PLIN1 depends on the membrane targeting segments, particularly the iMS (**Fig. S4F**).
276

Finally, we examined the localization of endogenously expressed PLIN1 during adipocyte
differentiation. As expected, six days post differentiation, PLIN1 localized almost exclusively to
the surface of LDs (**Fig. 4I**). However, at earlier days in the time course, e.g. at day 3 post
differentiation, the endogenous PLIN1 pool was additionally abundantly present on the
peripheral ER (**Fig. 4J**). Since the ER of adipocytes is structurally less defined than in other
cells[10,43,44], and the staining of the membrane is less clear than in other cells, it was critical to
additionally probe the membrane association biochemically. Consistent with the imaging results,
the PLIN1 membrane pool was completely alkaline-resistant, and pelleted together with the
integral ER marker CNX to the bottom of the tube (**Fig. 4L**). At this stage of the differentiation,
PLIN1 is likely produced in excess and is stock-piled in the ER to provide a readily available pool
of building blocks for rapid expansion of the LD-surface as the cells synthesize TAGs and the LDs
grow at high rate.
Interestingly, when we stimulated lipolysis in these cells triggering breakdown of the LDs,
PLIN1 efficiently moved back to the ER (**Fig. 4K**) and remained integral to the membrane as shown
by alkaline extraction (**Fig. 4K-L**). The interpretation that PLIN1 moves in these cells between the
LDs and the ER was further supported by flotation experiments, showing that stimulation of
lipolysis re-distributed PLIN1 away from the remaining floating LDs into the membrane pellet at
the bottom of the tube (**Fig. 4M**).
295
296
297
298

## Discussion

Here, we have shown that ER membrane integration and high LD-affinity of PLIN1 are necessary for the formation of LD heterogeneity in adipocytes. In contrast to its previous classification as a cytoplasmic, class-II LD protein (CYTOLD proteins), we have found that PLIN1 in fact behaves like an integral ER membrane protein. We show that an unconventional integral membrane segment (iMS) is necessary and sufficient for integration into the ER membrane and mediates at the same time high LD-affinity.

Our imaging and fractionation experiments with early adipocytes revealed that endogenously expressed PLIN1 is abundantly present in the ER and readily moves to the LD-class that emerges during the differentiation time course. Due to its high affinity, PLIN1 excludes the other PLINs from the surface of this first LD-class, already from the earliest stages in the biosynthesis process. The high LD-affinity is likely necessary to protect the *de novo* synthesized TAGs from turnover by lipases and is therefore required for efficient LD-growth. However, in later stages of adipocyte differentiation, the high LD-affinity has a "trade-off": starting from around day 6, the cellular PLIN1 pool is almost completely found on the surface of the existing LDs and the ER pool is reduced to a minimum. Under these conditions the PLIN1-LDs mostly grow through fusion mediated by CIDE-C and for a so far unknown reason are not able to incorporate new TAGs at high rates. During an acute burst of TAG synthesis, the cell rather forms a second LD population. Since the high LD-affinity "traps" PLIN1 on the surface of the first LD-class it is unavailable for association with the rapidly expanding surface of the second LD-class that form upon fatty acid uptake. This second LD-population is, therefore, primarily covered with the other PLIN-family members thus forming a separate LD class – the class-II LDs. Although class-II PLINs have lower affinity for LDs, they bind faster to the newly emerging LDs because they move onto the expanding LDs directly from the cytoplasm and do not have to take a de-tour through the ER.

Whether PLIN1 LD-targeting follows the same pathway as conventional class-I proteins (ERTOLD proteins) remains to be clarified (**Fig. S4F, ER→LD**). The length of the hydrophobic region within the iMS of around 30 amino acids (**Fig. S4G**) corresponds to sizes of known hairpin motifs, but the iMS exhibits surprisingly low hydrophobicity and does, therefore, not satisfy the basic definition of a conventional ER membrane anchor[45–47]. The low hydrophobicity is probably also the reason why the iMS is computationally not predicted as an integral membrane segment. However, the iMS contains some features that are found in other membrane segments that move between the ER and the LDs[13], such as prolines at the centre of the motif (position 14 and 18 in the iMS of mouse PLIN1) as well as large hydrophobic residues like two internal tryptophanes (**Fig. S4G**). The iMS also contains positively charged residues but instead of being in the core of the motif (**Fig. S4G**)[13], they flank the iMS, perhaps ensuring the correct membrane topology of PLIN1 or contributing to the regulation of LD access. Consistent with *in vitro* studies[48] our data indicate that the iMS favours the LD surface over the ER bilayer. Its high LD affinity might even reflect the ability to directly bind TAGs, which would possibly explain why it is readily sequestered to the large LDs of adipocytes and its abundance in the ER is low. In early adipocytes PLIN1 likely reaches the LDs from the ER on the class-I route using physical connections between the ER and LDs called "bridges"[27,49]. In later adipocytes which contain larger LDs, access may depend on other pathways which were only recently identified[50]. Another possibility may involve a cytoplasmic intermediate (**Fig. S4F ER→cytoplasm→LD**) as proposed by others[44]. However,

unless this potential cytoplasmic form of PLIN1 is for an unknown reason quantitatively sedimenting to the alkaline resistant membrane fraction, our data seem to exclude this route.

Why the PLIN1 covered LD-class does not efficiently expand upon fatty acid feeding and a second LD-population is required to buffer rapid TAG synthesis, is currently unknown. PLIN1 is, however, essential for adipose tissue function and cannot be replaced by the other PLINs: Loss of PLIN1 function in mice and humans can cause lipodystrophy characterized by dysfunctional or absent white adipose tissue[4,51]. PLIN1 ablation in mice leads to increased basal adipose lipolysis preventing efficient build-up of LDs[52,53]. Complete loss of PLIN1 function in humans is metabolically detrimental rendering the affected patients insulin-resistant and often causes hypertriglyceridemia, cardiovascular problems and fatty liver disease[53,54]. Partial loss of PLIN1 or PLIN1 haploinsufficiency can, surprisingly, have beneficial effects on metabolic health[55]. According to recent reports some shorter PLIN1 truncation mutants in fact improve metabolic profiles and reduce risk for cardiovascular disease[56,57]. A controlled perturbation of PLIN1 membrane targeting might therefore open new avenues for mechanism-based therapy of fatty liver disease, type-2 diabetes, and obesity.

**Acknowledgements**

We thank Pedro Carvalho (University of Oxford) for critical reading of the manuscript and Lucas Pelkmans, University of Zurich, for generous support with research infrastructure.

A.R.T. was supported by **ANR-MOBIL** and **ANR-21-CE11-LIPRODYN**. R.W.K acknowledges support from the Swiss National Science Foundation (**SNF 31003A_159793**), the Helmut Horten Foundation, and E.P. Abraham Cephalosporin fund. This publication arises from research funded by the John Fell Oxford University Press Research Fund and a grant from Diabetes UK (**22/0006453)**. R.W.K. acknowledges support from the imaging facility at the Dunn School of Pathology, University of Oxford.

**Author contribution:**

**Conceptualization,** R.W.K; **Methodology,** M.M., R.W.K., A.R.T; **Formal analysis,** M.M., O.S., D.A., K.B., M.O, A.R.T, R.W.K.; **Investigation,** M.M., O.S., D.A., K.B., M.O, A.R.T, R.W.K.; **Resources,** A.R.T. and R.W.K; **Writing – Original draft ,** R.W.K.; **Writing – Review & Editing,** M.M., O.S., A.R.T., R.W.K., **Visualization;** M.M., O.S., A.R.T., R.W.K.; **Supervision,** A.R.T. and R.W.K.; **Funding Acquisition;** A.R.T and R.W.K.

**Declaration of Interests**

The authors declare no competing interests.

**References**

381 **References**
382
383
1. Rosen, E.D., and Spiegelman, B.M. (2006). Adipocytes as regulators of energy balance and
glucose homeostasis. Nature *444*, 847 853. 10.1038/nature05483.
2. Morigny, P., Boucher, J., Arner, P., and Langin, D. (2021). Lipid and glucose metabolism in
white adipocytes: pathways, dysfunction and therapeutics. Nat Rev Endocrinol *17*, 276–295.
10.1038/s41574-021-00471-8.
3. Grabner, G.F., Xie, H., Schweiger, M., and Zechner, R. (2021). Lipolysis: cellular mechanisms
for lipid mobilization from fat stores. Nat Metabolism *3*, 1445–1465. 10.1038/s42255-021-
00493-6.
4. Robbins, A.L., and Savage, D.B. (2015). The genetics of lipid storage and human
lipodystrophies. Trends Mol Med *21*, 1 6. 10.1016/j.molmed.2015.04.004.
5. Samuel, V.T., and Shulman, G.I. (2012). Mechanisms for Insulin Resistance: Common Threads
and Missing Links. Cell *148*, 852–871. 10.1016/j.cell.2012.02.017.
6. Guilherme, A., Virbasius, J.V., Puri, V., and Czech, M.P. (2008). Adipocyte dysfunctions linking
obesity to insulin resistance and type 2 diabetes. Nat Rev Mol Cell Bio *9*, 367 377.
10.1038/nrm2391.
7. Olzmann, J.A., and Carvalho, P. (2019). Dynamics and functions of lipid droplets. Nat Rev Mol
Cell Bio *20*, 137 155. 10.1038/s41580-018-0085-z.
8. Walther, T.C., Chung, J., and Farese, R.V. (2017). Lipid Droplet Biogenesis. Annu Rev Cell Dev
Bi *33*, 491–510. 10.1146/annurev-cellbio-100616-060608.
9. Pol, A., Gross, S.P., and Parton, R.G. (2014). Review: biogenesis of the multifunctional lipid
droplet: lipids, proteins, and sites. J Cell Biology *204*, 635 646. 10.1083/jcb.201311051.
10. Wolins, N.E., Quaynor, B.K., Skinner, J.R., Schoenfish, M.J., Tzekov, A., and Bickel, P.E.
(2005). S3-12, Adipophilin, and TIP47 package lipid in adipocytes. J Biol Chem *280*, 19146
19155. 10.1074/jbc.m500978200.
11. Wolins, N.E., Quaynor, B.K., Skinner, J.R., Tzekov, A., Croce, M.A., Gropler, M.C., Varma, V.,
Yao-Borengasser, A., Rasouli, N., Kern, P.A., et al. (2006). OXPAT/PAT-1 is a PPAR-induced lipid
droplet protein that promotes fatty acid utilization. Diabetes *55*, 3418 3428. 10.2337/db06-
0399.
12. Kassan, A., Herms, A., Fernández-Vidal, A., Bosch, M., Schieber, N.L., Reddy, B.J.N., Fajardo,
A., Gelabert-Baldrich, M., Tebar, F., Enrich, C., et al. (2013). Acyl-CoA synthetase 3 promotes
lipid droplet biogenesis in ER microdomains. J Cell Biology *203*, 985 1001.
10.1083/jcb.201305142.
13. Olarte, M.-J., Kim, S., Sharp, M.E., Swanson, J.M.J., Farese, R.V., and Walther, T.C. (2020).
Determinants of Endoplasmic Reticulum-to-Lipid Droplet Protein Targeting. Dev Cell.
10.1016/j.devcel.2020.07.001.
14. Dhiman, R., Caesar, S., Thiam, A.R., and Schrul, B. (2019). Mechanisms of protein targeting
to lipid droplets: A unified cell biological and biophysical perspective. Semin. cell Dev. Biol. *108*,
4–13. 10.1016/j.semcdb.2020.03.004.
15. Wilfling, F., Wang, H., Haas, J.T., Krahmer, N., Gould, T.J., Uchida, A., Cheng, J.-X., Graham,
M., Christiano, R., Fröhlich, F., et al. (2013). Triacylglycerol synthesis enzymes mediate lipid

droplet growth by relocalizing from the ER to lipid droplets. Dev Cell *24*, 384 399.
10.1016/j.devcel.2013.01.013.

16. Eisenberg-Bord, M., Mari, M., Weill, U., Rosenfeld-Gur, E., Moldavski, O., Castro, I.G., Soni,
K.G., Harpaz, N., Levine, T.P., Futerman, A.H., et al. (2018). Identification of seipin-linked factors
that act as determinants of a lipid droplet subpopulation. J Cell Biol *217*, 269 282.
10.1083/jcb.201704122.

17. Chitraju, C., Mejhert, N., Haas, J.T., Diaz-Ramirez, L.G., Grueter, C.A., Imbriglio, J.E., Pinto, S.,
Koliwad, S.K., Walther, T.C., and Jr, R.V.F. (2017). Triglyceride Synthesis by DGAT1 Protects
Adipocytes from Lipid-Induced ER Stress during Lipolysis. Cell Metab *26*, 407 418.e3.
10.1016/j.cmet.2017.07.012.

18. Thiam, A.R., and Beller, M. (2017). The why, when and how of lipid droplet diversity. J Cell
Sci *130*, 315–324. 10.1242/jcs.192021.

19. Straub, B.K., Stoeffel, P., Heid, H., Zimbelmann, R., and Schirmacher, P. (2008). Differential
pattern of lipid droplet-associated proteins and de novo perilipin expression in hepatocyte
steatogenesis. Hepatology *47*, 1936–1946. 10.1002/hep.22268.

20. Hsieh, K., Lee, Y.K., Londos, C., Raaka, B.M., Dalen, K.T., and Kimmel, A.R. (2012). Perilipin
family members preferentially sequester to either triacylglycerol-specific or cholesteryl-ester-
specific intracellular lipid storage droplets. J Cell Sci *125*, 4067 4076. 10.1242/jcs.104943.

21. Herms, A., Bosch, M., Ariotti, N., Reddy, B.J.N., Fajardo, A., Fernández-Vidal, A., Alvarez-
Guaita, A., Fernández-Rojo, M.A., Rentero, C., Tebar, F., et al. (2013). Cell-to-cell heterogeneity
in lipid droplets suggests a mechanism to reduce lipotoxicity. Curr Biol *23*, 1489 1496.
10.1016/j.cub.2013.06.032.

22. Henne, M., Goodman, J.M., and Hariri, H. (2019). Spatial compartmentalization of lipid
droplet biogenesis. Biochimica Et Biophysica Acta Mol Cell Biology Lipids *1865*, 158499.
10.1016/j.bbalip.2019.07.008.

23. Ugrankar, R., Bowerman, J., Hariri, H., Chandra, M., Chen, K., Bossanyi, M.-F., Datta, S.,
Rogers, S., Eckert, K.M., Vale, G., et al. (2019). Drosophila Snazarus Regulates a Lipid Droplet
Population at Plasma Membrane-Droplet Contacts in Adipocytes. Dev Cell *50*, 557-572.e5.
10.1016/j.devcel.2019.07.021.

24. Heid, H., Rickelt, S., Zimbelmann, R., Winter, S., Schumacher, H., Dörflinger, Y., Kuhn, C., and
Franke, W.W. (2014). On the formation of lipid droplets in human adipocytes: the organization
of the perilipin-vimentin cortex. Plos One *9*, e90386. 10.1371/journal.pone.0090386.

25. Wolins, N.E., Brasaemle, D.L., and Bickel, P.E. (2006). A proposed model of fat packaging by
exchangeable lipid droplet proteins. Febs Lett *580*, 5484 5491. 10.1016/j.febslet.2006.08.040.

26. Freyre, C.A.C., Rauher, P.C., Ejsing, C.S., and Klemm, R.W. (2019). MIGA2 Links
Mitochondria, the ER, and Lipid Droplets and Promotes De Novo Lipogenesis in Adipocytes. Mol
Cell *76*, 811-825.e14. 10.1016/j.molcel.2019.09.011.

27. Kory, N., Farese, R.V., and Walther, T.C. (2016). Targeting Fat: Mechanisms of Protein
Localization to Lipid Droplets. Trends Cell Biol *26*. 10.1016/j.tcb.2016.02.007.

28. Olarte, M.-J., Swanson, J.M.J., Walther, T.C., and Farese, R.V. (2022). The CYTOLD and
ERTOLD pathways for lipid droplet–protein targeting. Trends Biochem. Sci. *47*, 39–51.
10.1016/j.tibs.2021.08.007.

29. Prévost, C., Sharp, M.E., Kory, N., Lin, Q., Voth, G.A., Farese, R.V., and Walther, T.C. (2018).
Mechanism and Determinants of Amphipathic Helix-Containing Protein Targeting to Lipid
Droplets. Dev Cell *44*, 73 86.e4. 10.1016/j.devcel.2017.12.011.

30. Kory, N., Thiam, A.-R., Farese, R.V., and Walther, T.C. (2015). Protein Crowding Is a
Determinant of Lipid Droplet Protein Composition. Dev Cell *34*, 351–363.
10.1016/j.devcel.2015.06.007.

31. Rowe, E.R., Mimmack, M.L., Barbosa, A.D., Haider, A., Isaac, I., Ouberai, M.M., Thiam, A.R.,
Patel, S., Saudek, V., Siniossoglou, S., et al. (2016). Conserved Amphipathic Helices Mediate
Lipid Droplet Targeting of Perilipins 1-3. J Biol Chem *291*, 6664 6678. 10.1074/jbc.m115.691048.

32. Čopič, A., Antoine-Bally, S., Giménez-Andrés, M., Garay, C.T., Antonny, B., Manni, M.M.,
Pagnotta, S., Guihot, J., and Jackson, C.L. (2018). A giant amphipathic helix from a perilipin that
is adapted for coating lipid droplets. Nat Commun *9*, 1 16. 10.1038/s41467-018-03717-8.

33. Giménez-Andrés, M., Emeršič, T., Antoine-Bally, S., D'Ambrosio, J.M., Antonny, B., Derganc,
J., and Čopič, A. (2021). Exceptional stability of a perilipin on lipid droplets depends on its polar
residues, suggesting multimeric assembly. Elife *10*, e61401. 10.7554/elife.61401.

34. Chorlay, A., Monticelli, L., Ferreira, J.V., M'barek, K.B., Ajjaji, D., Wang, S., Johnson, E., Beck,
R., Omrane, M., Beller, M., et al. (2019). Membrane Asymmetry Imposes Directionality on Lipid
Droplet Emergence from the ER. Dev Cell *50*, 25-42.e7. 10.1016/j.devcel.2019.05.003.

35. Bickel, P.E., Tansey, J.T., and Welte, M.A. (2009). PAT proteins, an ancient family of lipid
droplet proteins that regulate cellular lipid stores. Biochimica Et Biophysica Acta Bba - Mol Cell
Biology Lipids *1791*, 419 440. 10.1016/j.bbalip.2009.04.002.

36. Londos, C., Brasaemle, D.L., Schultz, C.J., Segrest, J.P., and Kimmel, A.R. (1999). Perilipins,
ADRP, and other proteins that associate with intracellular neutral lipid droplets in animal cells.
Semin Cell Dev Biol *10*, 51 58. 10.1006/scdb.1998.0275.

37. Kimmel, A.R., and Sztalryd, C. (2016). The Perilipins: Major Cytosolic Lipid Droplet–
Associated Proteins and Their Roles in Cellular Lipid Storage, Mobilization, and Systemic
Homeostasis*. Annu Rev Nutr *36*, 471 509. 10.1146/annurev-nutr-071813-105410.

38. Brasaemle, D.L. (2007). Thematic review series: adipocyte biology. The perilipin family of
structural lipid droplet proteins: stabilization of lipid droplets and control of lipolysis. J Lipid Res
*48*, 2547 2559. 10.1194/jlr.r700014-jlr200.

39. Garcia, A., Subramanian, V., Sekowski, A., Bhattacharyya, S., Love, M.W., and Brasaemle,
D.L. (2004). The amino and carboxyl termini of perilipin a facilitate the storage of
triacylglycerols. J Biol Chem *279*, 8409 8416. 10.1074/jbc.m311198200.

40. Choi, Y.M., Ajjaji, D., Fleming, K.D., Borbat, P.P., Jenkins, M.L., Moeller, B.E., Fernando, S.,
Bhatia, S.R., Freed, J.H., Burke, J.E., et al. (2023). Structural insights into perilipin 3 membrane
association in response to diacylglycerol accumulation. Nat. Commun. *14*, 3204.
10.1038/s41467-023-38725-w.

41. Garcia, A., Sekowski, A., Subramanian, V., and Brasaemle, D.L. (2003). The central domain is
required to target and anchor perilipin A to lipid droplets. J Biol Chem *278*, 625 635.
10.1074/jbc.m206602200.

42. Ajjaji, D., M'barek, K.B., Mimmack, M.L., England, C., Herscovitz, H., Dong, L., Kay, R.G.,
Patel, S., Saudek, V., Small, D.M., et al. (2019). Dual binding motifs underpin the hierarchical
association of perilipins1–3 with lipid droplets. Mol Biol Cell *30*, 703–716. 10.1091/mbc.e18-08-
0534.

43. Wolins, N.E., Skinner, J.R., Schoenfish, M.J., Tzekov, A., Bensch, K.G., and Bickel, P.E. (2003).
Adipocyte protein S3-12 coats nascent lipid droplets. J Biol Chem *278*, 37713 37721.
10.1074/jbc.m304025200.
44. Skinner, J.R., Harris, L.-A.L.S., Shew, T.M., Abumrad, N.A., and Wolins, N.E. (2013). Perilipin 1
moves between the fat droplet and the endoplasmic reticulum. Adipocyte *2*, 80 86.
10.4161/adip.22864.
45. Shibata, Y., Hu, J., Kozlov, M.M., and Rapoport, T.A. (2009). Mechanisms shaping the
membranes of cellular organelles. Annu Rev Cell Dev Bi *25*, 329 354.
10.1146/annurev.cellbio.042308.113324.
46. Parton, R.G., and Pozo, M.A. del (2013). Caveolae as plasma membrane sensors, protectors
and organizers. Nat Rev Mol Cell Bio *14*, 98 112. 10.1038/nrm3512.
47. Schrul, B., and Kopito, R.R. (2016). Peroxin-dependent targeting of a lipid-droplet-destined
membrane protein to ER subdomains. Nat Cell Biol *18*, 1 26. 10.1038/ncb3373.
48. Caillon, L., Nieto, V., Gehan, P., Omrane, M., Rodriguez, N., Monticelli, L., and Thiam, A.R.
(2020). Triacylglycerols sequester monotopic membrane proteins to lipid droplets. Nat
Commun *11*, 3944. 10.1038/s41467-020-17585-8.
49. Wilfling, F., Thiam, A.R., Olarte, M.-J., Wang, J., Beck, R., Gould, T.J., Allgeyer, E.S., Pincet, F.,
Bewersdorf, J., Farese, R.V., et al. (2014). Arf1/COPI machinery acts directly on lipid droplets
and enables their connection to the ER for protein targeting. Elife *3*, e01607.
10.7554/elife.01607.
50. Song, J., Mizrak, A., Lee, C.-W., Cicconet, M., Lai, Z.W., Tang, W.-C., Lu, C.-H., Mohr, S.E.,
Farese, R.V., and Walther, T.C. (2022). Identification of two pathways mediating protein
targeting from ER to lipid droplets. Nat Cell Biol, 1–14. 10.1038/s41556-022-00974-0.
51. Mann, J.P., and Savage, D.B. (2019). What lipodystrophies teach us about the metabolic
syndrome. J Clin Invest *129*, 4009–4021. 10.1172/jci129190.
52. Tansey, J.T., Sztalryd, C., Gruia-Gray, J., Roush, D.L., Zee, J.V., Gavrilova, O., Reitman, M.L.,
Deng, C.-X., Li, C., Kimmel, A.R., et al. (2001). Perilipin ablation results in a lean mouse with
aberrant adipocyte lipolysis, enhanced leptin production, and resistance to diet-induced
obesity. Proc. Natl. Acad. Sci. *98*, 6494–6499. 10.1073/pnas.101042998.
53. Gandotra, S., Lim, K., Girousse, A., Saudek, V., O'Rahilly, S., and Savage, D.B. (2011). Human
frame shift mutations affecting the carboxyl terminus of perilipin increase lipolysis by failing to
sequester the adipose triglyceride lipase (ATGL) coactivator AB-hydrolase-containing 5
(ABHD5). J Biological Chem *286*, 34998–35006. 10.1074/jbc.m111.278853.
54. Garg, A. (2011). Clinical review#: Lipodystrophies: genetic and acquired body fat disorders. J
Clin Endocrinol Metabolism *96*, 3313 3325. 10.1210/jc.2011-1159.
55. Melvin, A., Stears, A., and Savage, D.B. (2019). Recent developments in lipodystrophy. Curr.
Opin. Lipidol. *30*, 284–290. 10.1097/mol.0000000000000613.
56. Koprulu, M., Zhao, Y., Wheeler, E., Dong, L., Rocha, N., Li, C., Griffin, J.D., Patel, S., Streek,
M.V. de, Glastonbury, C.A., et al. (2021). Identification of rare loss of function genetic variation
regulating body fat distribution. J. Clin. Endocrinol. Metab. *107*, dgab877-.
10.1210/clinem/dgab877.
57. Patel, K.A., Burman, S., Laver, T.W., Hattersley, A.T., Frayling, T.M., and Weedon, M.N.
(2022). PLIN1 Haploinsufficiency Causes a Favorable Metabolic Profile. J. Clin. Endocrinol.
Metab. *107*, e2318–e2323. 10.1210/clinem/dgac104.

58. Krogh, A., Larsson, B., Heijne, G. von, and Sonnhammer, E.L. (2001). Predicting

transmembrane protein topology with a hidden Markov model: application to complete

genomes. J Mol Biol *305*, 567 580. 10.1006/jmbi.2000.4315.

59. Jayasinghe, S., Hristova, K., and White, S.H. (2001). Energetics, stability, and prediction of

transmembrane helices. J Mol Biol *312*, 927 934. 10.1006/jmbi.2001.5008.

**Figure 1**

**Figure 1**
**Fatty acid feeding leads to the formation of at least two distinct LD-classes in adipocytes.**
**(A)** Class-I LD proteins are monotopic membrane proteins that move from the ER (membrane
brown, ER lumen blue) to the LD surface. Class-II LD proteins bind to LDs directly from the
cytoplasm (pink).
**(B)** Segregation of class-II proteins (e.g. perilipins) to different LD-classes.
**(C)** Day 6 3T3-L1 derived adipocytes were fed for 3h with 750 µM $Na^{2+}$oleate. Endogenously
expressed PLIN1 and PLIN2 were visualized by immunofluorescence. Scale bar, 10 µm.
**(D)** As in (C) but PLIN1 and PLIN3 were visualized.
**(E)** As in (C) but PLIN1 and PLIN4 were visualized.
**(F)** Quantification of the results in (C)-(E). n=9; n=9; n=8, for cells co-stained with PLIN1 and 2 or
PLIN1 and 3 or PLIN1 and 4, respectively. LD counts per replicate >100; values are expressed as
mean LD in each class ± SEM.
574

**Figure 2**

**Figure 2**
**PLIN1 is an integral membrane protein which localizes to the ER and LDs.**
**(A)** Mouse PLIN1 tagged at its N-terminus with GFP (GFP-PLIN1) was expressed in COS7 cells.
The ER is visualized with an antibody specific to the luminal ER protein calreticulin (CRT). LDs
are stained with LipidToxRed. Scale bar, 10 μm.
**(B)** As in (A) but the cells were transfected with GFP-PLIN2.
**(C)** As in (A) but 250 μM Na$^+$oleate was added for 6h prior to fixation.
**(D)** As in (C) but the cells were transfected with GFP-PLIN2.
**(E)** Quantification of results in (A) and (C) using supervised machine learning to identify cells
with GFP-PLIN1 on the ER. n = 3 independent experiments with >1500 cell per replicate; values
are expressed as the mean fraction of cells with PLIN1 on the ER ±SD.
**(F)** COS7 cells were transfected with untagged PLIN1, GFP-PLIN1 or GFP-PLIN2, and subjected to
fractionation followed by alkaline extraction. **Input,** cell lysate; **S100, P100,** supernatant and
pellet of a 100k x g centrifugation step, **S100E, P100E,** supernatant and pellet of 100k x g
centrifugation after alkaline extraction with Na$_2$CO$_3$. **GAPDH,** glycerolaldehyde-3-phosphate
dehydrogenase (cytoplasm); **CNX,** calnexin (integral ER-membrane protein).
**(G)** As in (F), but P100 was subjected to flotation on a sucrose step gradient with the indicated
concentrations.
**(H)** As in (G), but P100 was incubated with Triton X-100 before flotation.
**(I)** As in (H), but the cell lysate (input) was treated with Triton X-100 before centrifugation at
100k x g.
**(J)** PLIN1 is an integral membrane protein with a membrane anchor (blue) allowing dual
targeting to the ER and the LD surface (unknown pathway indicated as "?").
599

**Figure 3**

**Figure 3**

**Identification of two ER targeting segments in PLIN1.**

**(A)** Structure-function analysis of PLIN1 by fractionation and alkaline extraction using the indicated truncation constructs and immune-blotting with a anti GFP-antibody. **PAT,** perilipin, adipohilin/ADRP, TIP-47 domain (purple); **11-mer,** LD targeting region (pink); **HS1, 2** hydrophobic segment 1 and 2 (green, orange, resp.); **AT,** acidic tract (red); **AS,** amphipathic segment (yellow); **C-term,** carboxy terminus with regulatory elements (grey). Classification explained in Fig. S3B.

**(B)** COS7 cells expressing a GFP-tagged N-terminal portion of PLIN1 (1-237). The ER was visualized by immunofluorescence with an antibody to calreticulin (CRT). The nuclei were stained with DAPI. Scale bar, 10 µm.

**(C)** As in (B) but with a different truncation construct (PLIN1 1-281).

**(D)** As in (B) but with cells expressing the C-terminal truncation construct PLIN1 281-519.

**(E)** As in (D) but with PLIN1 371-517.

**(F)** As in (D) but with PLIN1 328-517.

**(G)** Same illustrations as in (A) but the integral membrane segment (iMS, 238-280) and the peripheral membrane segment (pMS, 348-370) are highlighted. PLIN1ΔΔ lacks the iMS and the pMS.

**(H)** As in (A) but with a construct lacking the iMS and the pMS (GFP-PLIN1ΔΔ). Membranes in the P100 fraction were in parallel subjected to alkaline extraction (+$Na_2CO_3$) and flotation experiments.

**(I)** As in (B) but with PLIN1ΔΔ and the LDs were stained with LipidToxDeepRed.

**(J)** COS7 cells were co-transfected with PLIN2 N-terminally tagged with mCherry2, and GFP tagged PLIN1ΔΔ. The cells were treated with 250 µM $Na^+$oleate 6h prior to fixation.

**(K)** Deletion of the iMS and pMS results in a switch of LD-protein class with PLIN1 and PLIN1ΔΔ behaving as class-I and class-II LD proteins, respectively. Domain description as in (A).

**Figure 4**

**Figure 4**
**Membrane association determines LD-class specificity by tuning LD binding kinetics**
**(A)** GFP-PLIN1ΔΔ and mCherry-PLIN1 full-length were reconstituted in an buffer-in oil system
and their localization was monitored by microscopy as the aqueous phase progressively shrinks.
**(B)** Ratio of fluorescence intensity at the oil-buffer interface and in the buffer phase at 4
different surface compression ratios for PLIN1ΔΔ (green) and PLIN1 full length (red). N=3
independent experiments; mean ± SD.
**(C)** Fluorescence recovery after photobleaching in oleate treated Huh7 cells expressing GFP-
PLIN1ΔΔ, ΔiMS or ΔpMS. Scale bar, 10 μm. N = 3; results are expressed as mean ± SD.
**(D)** GFP-PLIN1ΔΔ or GFP-PLIN1 (full-length) were expressed in 3T3-L1 cells from a stably
integrated inducible transgene and subjected to fractionation and alkaline extraction. **Input,**
cell lysate; **S100, P100,** supernatant and pellet of a 100k x g centrifugation step, **S100E, P100E,**
supernatant and pellet of 100k x g centrifugation after alkaline extraction with 100 mM $Na_2CO_3$.
**GAPDH,** glyceraldehyde-3-phosphate dehydrogenase (cytoplasm); **CLIMP-63,** integral ER-
membrane protein.
**(E)** As in (D) but the cells were differentiated into adipocytes and expression of PLIN1ΔΔ was
induced 6 days post differentiation by the addition of 500 ng/ml anhydrotetracycline 6h prior to
fixation. LDs were stained with LipidToxDeepRed and visualized together with GFP-PLIN1ΔΔ by
fluorescence microscopy. Scale bar, 10 μm.
(**F**) Percentage of the cells in (E) and (H) in which GFP-PLIN1ΔΔ is either on peripheral LDs or in
the cytoplasm, N = 100 cells per group.
**(G)** As in (E) but the adipocytes were subjected to flotation. Endogenously expressed PLIN1 and
PLIN1ΔΔ were visualized by immuno-blotting with antibodies specific to the C-term of PLIN1
recognizing both proteins.
**(H)** As in (E) with addition of oleate 6h before fixation. In contrast to (F) GFP-PLIN1ΔΔ is now
found on peripheral LDs. PLIN1ΔΔ remains excluded from the central LDs covered by endogenous
PLIN1. On the right, additional staining with anti-PLIN1 antibody (red) recognizing both
endogenous PLIN1 on large LDs (red) and the PLIN1ΔΔ mutant on the peripheral LDs
(GFP+red=yellow).
(**I**) Localization of endogenous PLIN1 (green) in 3T3-L1 adipocytes at day 6 post differentiation;
line plot of PLIN1 intensity (normalized to maximum), as drawn in the inset.. Scale bar: 10 μm for
whole cell image, 4 m for inset 4 μm.
(J) As in (I), but at day 3 post differentiation endogenous PLIN1 is on the ER. Arrowheads mark
terminal points of ER network
(K) As in (J), but after addition of 10 μM forskolin to stimulate lipolysis for 6h prior to fixation.
(L) Fractionation and alkaline extraction of adipocytes as in (J) and (K); labels as in (D); **CNX,**
calnexin (integral ER-membrane protein).
(M) Flotation experiment of the cells as in (J) and (K), done as in (G).

**Figure S1**

**Figure S1**

**PLIN1 excludes PLIN2, 3 and 4 from LD binding in COS7 cells.**

**(A)** Schematic representation of perilipin (PLIN) 1-4. The amino terminal PAT (perilipin,

adipophilin/ADRP domain) and 11-mer regions are important for LD targeting. PLIN1-3 contain a

predicted four helix bundle. The PLINs were grouped into a protein family based on the sequence

similarity found in the PAT and 11-mer domains as well as their localization to the LD surface.

PLIN4 does not contain a four-helix bundle and is much larger than the other family members.

**(B)** COS7 cells co-expressing PLIN1 tagged at its N-terminus with GFP (GFP-PLIN1) and PLIN2

carrying an N-terminal mCherry2 tag (mCherry-PLIN2). 250 µM Na$^+$oleate was added to the

growth medium 6h prior to fixation. Lipid droplets were stained with LipidToxDeepRed (LipidTox).

The cells were automatically segmented and assigned by a support vector machine based

classifier into 4 groups, in which PLIN1 was exclusively found on the LD surface and PLIN2 was

cytoplasmic **(PLIN1)**, cells that showed only PLIN2 LD staining **(PLIN2)**, cells in which various levels

of PLIN2 was found on LDs together with PLIN1 **(MIX)**, and cells in which neither of the two

proteins was found on the LD **(NONE)**. Intensity and texture of PLIN1 and PLIN2 channels as well

as correlation with LipidTOXDeepRed were used as training features. n = 3 independent

experiments ; the results are expressed as mean ±SD. Scale bar, 10µm.

**(C)** As in (B) but PLIN3 tagged at its N-terminus with c-myc (myc-PLIN3) was co-expressed with

GFP-PLIN1. n = 3 independent experiments ; the results are expressed as mean ±SD.

**(D)** As in (C) but PLIN4 tagged at its N-terminus with mCherry2 (mCherry-PLIN4) was co-expressed

with GFP-PLIN1. n = 3 independent experiments ; the results are expressed as mean ±SD.

**(E)** COS7 cells co-expressing PLIN1 truncated after the 11-mer at residue 192 tagged at its N-

terminus with a GFP (GFP-PLIN1-1-192) and PLIN2 tagged at its N-terminus with mCherry2

(mCherry-PLIN2).

**Figure S2**

**Figure S2**

**Subcellular targeting of PLIN1 and PLIN2;**

**TMHMM predictions and hydropathy plot analysis for PLIN1 and PLIN2;**

**PLIN1 LD targeting does not require translation.**

**(A)** Schematic illustration of image stack and the middle plane shown in (B) and (C).

**(B)** COS7 cells expressing PLIN1 tagged at its N-terminus with GFP. Lipid droplets were stained

with LipidToxRed and the ER was visualized with an antibody specific to the endogenously

expressed luminal ER protein calreticulin (CRT).

**(C)** As in (B) but cells expressed PLIN2 with an N-terminal GFP tag (GFP-PLIN2).

**(D)** Topology analysis of PLIN1 using TMHMM 2.0 (Trans Membrane prediction Hidden Markov

Model[58] https://services.healthtech.dtu.dk/service.php?TMHMM-2.0).

No integral membrane domains are predicted. **Outside,** segment in cytoplasm; **Inside,** segment

inside the ER lumen; **transmembrane,** integral membrane segment .

**(E)** As in (D) but for PLIN2.

**(F)** Hydropathy plot analysis for PLIN1 with membrane protein explorer (MPEX)[59].

**(G)** As in (F) but for PLIN2.

**(H)** COS7 cells expressing PLIN1 tagged at its N-terminus with GFP (GFP-PLIN1). Lipid droplets

were stained with LipidToxRed (LipidTox) and nuclei were visualized with DAPI. Cells were

either treated with DMSO as a control or with 100μg/ml cycloheximide 7h prior to fixation.

Scale bar, 10 μm.

**(I)** As in **(H)** but the cells were additionally treated with 200 μM Na$^+$oleate for 6h prior to

fixation (1h after addition of DMSO or 100μg/ml cycloheximide). The results are expressed as %

of cells with GFP-PLIN1 on LDs.

**(J)** Quantification of (H) and (I). The results are expressed as % of cells with GFP-PLIN1 on ER;

DMSO: N=20 (-OA), N=28 (+OA); cycloheximide: N=27 (+OA), N=20 (-OA).

**Figure S3**

 **Figure S3**
**Summary of all constructs used in the structure function analysis of PLIN1.**
**(A)** Structure-function analysis of PLIN1 by fractionation and alkaline extraction using the
indicated truncation constructs.
**Input,** cell lysate; **S100, P100,** supernatant and pellet of a 100k x g centrifugation step, **S100E,**
**P100E,** supernatant and pellet of 100k x g centrifugation after alkaline extraction with $Na_2CO_3$.
Note that PLIN1 1-517 (full length) is also shown in Fig. 2F (GFP-PLIN1) and the constructs 1-
237, 1-281 as well as 281-517, 328-517, and 371-517 are shown in main Fig. 3A. PLIN1ΔΔ is also
shown in Fig. 3H.
**(B)** Schematic illustration of the domains in PLIN1 and the tested truncation constructs. The
table summarizes the results in (A) classifying the construct as soluble when found in S100 and
in the cytoplasm by microscopy, as peripheral when found in S100, but seen on the ER by
microscopy, and integral when present in the alkaline resistant membrane pellet P100E. ER,
and cytoplasm is the localization as determined by microscopy. **PAT,** perilipin, adipohilin/ADRP,
TIP-47 domain (purple); **11-mer,** LD targeting region (pink); **HS1, 2** hydrophobic segment 1 and
2 (green, orange, resp.); **AT,** acidic tract (red); **AS,** amphipathic segment (yellow); **C-term,**
carboxy terminus with regulatory elements (grey). Classification of the constructs as soluble,
cytoplasmic, peripheral or integral ER membrane proteins.
**(C)** Fractionation and alkaline extraction experiments with GFP-PLIN2 (also shown in Fig 2F, for
simplicity marked as PLIN2) and GFP-PLIN2 containing the iMS core from PLIN1 inserted after
amino acid at position 251. For comparison with fractions in Fig. 2F the cytoplasmic marker
**GAPDH,** glycerolaldehyde-3-P-dehydrogenase (from Fig. 2F) and the ER membrane protein
marker **CNX,** calnexin (from Fig. 2F) are shown.
**(D)** As in (B) but PLIN2 is shown. Classification and definitions as in (B).
**(E)** COS7 cells were transfected with PLIN1 tagged at its N-terminus with mCherry2 (mCherry-
PLIN1) and PLIN2 containing the iMS core from PLIN1 inserted after amino acid at position 251
(GFP-PLIN1+iMS (251). The cells were additionally treated with 250 µM $Na^+$oleate 6h prior to
fixation when indicated. The illustration shows that PLIN1 and PLIN2+iMs segregate
preferentially to distinct LD classes showing that the iMS is critical in class specific LD nucleation
at the ER. Domain colours are as indicated in (B) and (D).

**Figure S4**

756
**Figure S4.**
**Recruitment of PLIN1 from the ER to the LD surface is independent of translation**
**(A)** Huh7 cells expressing the indicated GFP-tagged PLIN1 constructs were subjected to
photobleaching and the recovery of the GFP signal within the boxed area was plotted over
time. **Related to Fig. 4C.**
**(B)** 3T3-L1 cells expressing PLIN1 tagged at its N-terminus with GFP (GFP-PLIN1) from a stably
integrated inducible trans-gene. Expression was induced with 500 ng/ml anhydrotetracycline
(AHT) over-night (total 20h). Lipid droplets were stained with LipidToxRed (LipidTox) and nuclei
were visualized with DAPI. Scale bar, 10 µm.
**(C)** As in (B) but the cells were additionally treated with Na$^+$oleate 3h prior to fixation.
**(D)** As in (C) but translation was shut off by treating the cells with 100 µg/ml cycloheximide 1h
before the addition of Na$^+$oleate.
**(E)** Validation of GFP-PLIN1 expression and translation shut off by cycloheximide. To induce
expression of GFP-PLIN1, the cells were treated with 500ng/ml anhydrotetracycline (AHT) 6h
prior to cell harvest. Translation was inhibited with the indicated concentrations of
cycloheximide 1h before treatment with AHT. As a negative control AHT was omitted (-AHT).
**(F)** PLIN1 is targeted to the ER membrane (brown) by the iMS (green) and the pMS (orange).
PLIN1 likely moves to the LD using bridge-connections or by an unknown cytoplasmic
intermediate that carries lipids to the LD (indicated by "?"). PLIN1ΔΔ binds to the LD directly
from the cytoplasm as expected for a prototypical class-II protein.
**(G)** Alignment of mouse, rat and human PLIN1 iMS. The regions correspond to mouse PLIN1
residues 238-280. The sequences were retrieved from Uniprot Q8CGN5, P43884, O60240
respectively, aligned using ClustalOmega https://www.ebi.ac.uk/Tools/msa/clustalo/ and
analysed in JalView https://www.jalview.org/ using the T-coffee algorithm for determination of
the consensus sequence shown below the sequence conservation bar graph.
**(H)** LD-binding rates of class-II proteins are faster than for class-I proteins. The steady state
PLIN1 ER-pool is low, slowing down the LD on-reaction. When bound to the LD, class-II proteins
exchange faster than class-I proteins. The resulting high affinity to LDs leads to the gradual
replacement of class-II by class-I proteins from the newly formed LDs.
786
787

# Experimental procedures

## STAR Methods

### Lead contact and materials availability
Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Robin Klemm (Robin.Klemm@dpag.ox.ac.uk).

### Data and code availability
Original/source data for in the paper is available upon request from the lead contact and in Supplementary Data.

### Method Details
### Cell lines

COS-7 cells were a kind gift from Tom A. Rapoport (Harvard Medical School, Boston, USA). COS7 is a male African green monkey kidney fibroblast cell line. HEK293T cells were obtained from Dharmacon (#TLP5918). HEK293T is a female human embryonic kidney cell line. 3T3-L1 pre-adipocytes were from ATCC (CL-173). 3T3-L1 is a continuous sub-strain of male murine 3T3 (Swiss albino) cells, developed through clonal isolation.

### Cell culture

COS-7 cells were maintained in Dulbeccos's Modified Eagle Medium with 4.5 g/l glucose and L-glutamine (DMEM, Thermo Fischer Scientific, #41965062), supplemented with 10% fetal bovine serum (FBS; Sigma-Aldrich, #F7524). HEK293T cells were maintained in DMEM containing 10% FBS, 25 mM HEPES (Sigma-Aldrich, H4034), 1 mM sodium pyruvate (Thermo-Fisher), 2 mM glutamine (Sigma-Aldrich) and 100 µM non-essential amino acids (Thermo-Fisher). All cells were kept as sub-confluent cultures and were sub-cultured regularly. All cell lines were periodically tested for mycoplasma and grown according to ATCC guidelines. Cells were incubated at 37°C and 5% $CO_2$.

### Pre-adipocyte maintenance

3T3-L1 pre-adipocytes were obtained from ATCC (CL-173) and cultured in DMEM with 4.5 g/l glucose and L-glutamine (Thermo Fischer Scientific, #41965062), containing 10% calf serum (CS; Sigma-Aldrich, #12133C). Cells were grown as sub-confluent cultures in 10 cm tissue culture dishes (TPP) and sub-cultured regularly.

### Adipocyte differentiation procedure

For the differentiation of 3T3-L1 pre-adipocytes into white adipocytes, the cells were first grown to confluence and kept as a confluent culture for 48 hours. For differentiation in 96-well format, cells were seeded at a density of 10,000 cells per well of a 96-well plate (Greiner). Typically, the cells reached confluence the next day. After 48 hours of confluence differentiation was induced by the addition of adipocyte differentiation medium consisting of DMEM supplemented with 10% FBS, 172 nM bovine insulin (Sigma-Aldrich, #I0516), 500 µM 3-

Isobutyl-1-methylxanthine (IBMX, Sigma-Aldrich #I5879), and 1 µM dexamethasone
(Gbiosciences, #API-04). Cells were kept in differentiation medium for 48 hours. The
differentiation medium was then replaced by DMEM containing 10% FBS and 172 nM bovine
insulin. Cells were kept in the insulin medium for another 48 hours. At day four after induction
of differentiation the insulin medium was replaced by DMEM containing 10% FBS. The medium
was then refreshed every two days. Typically, cells were used for experiments at day 6 or day 8
after start of differentiation.
834

**Method details**

**Transfection and subcellular fractionation and alkaline extraction**

$1.1 \times 10^6$ COS-7 cells were seeded on a 10 cm dish the one day before transfection. The next
morning cells were transfected with 10 µg of the plasmid of interest using GeneJuice
transfection reagent (Merck, #70967-3) according to the manufacturer's instructions. After 6
hours of incubation at 37°C the medium was replaced with fresh DMEM containing 10% FBS.
The cells were then incubated at 37°C/5% $CO_2$ overnight. The next morning the cells were
washed once with ice-cold PBS and then harvested in 1 mL PBS with a cell scraper. The cells
were briefly pelleted at 2000 g for 5 min at 4°C, and the supernatant was discarded. The cells
were resuspended in 1 mL PBS containing protease inhibitor cocktail (cOmplete™, Mini, EDTA-
free Protease Inhibitor Cocktail; Merck, #11836153001) (PBSI) and lysed by 20 passages
through a bead-homogenizer (Isobiotec, 16 µm bead). A small aliquot for subsequent analysis
by SDS-PAGE was taken ("Input") and the remaining lysate was clarified by centrifuging for 10
minutes at 4°C and 2000 g. The supernatant was transferred to a 1.5 mL ultracentrifuge tube
(Beckman-Coulter, #357448) and the membranes were pelleted at 100,000 g for one hour at
4°C in a TLA 100.3 rotor (Beckman-Coulter). Afterwards the supernatant was taken off and
stored at -20°C ("S100") and the pellet was resuspended in 50 µL PBSI. A small aliquot of 25 µL
was stored at -20°C for analysis by SDS-PAGE ("P100") and the remaining 25 µL were subjected
to alkaline extraction by adding 75 µL of 250 mM $Na_2CO_3$ (pH 11) (Sigma Aldrich,
#A0634292404) for 15 min on ice. The sample was transferred to a 250 µL ultracentrifuge tube
(5 x 20 mm Beckman-Coulter, #342630) and a 50 µL of 10% sucrose cushion was placed below
the sample at the bottom of the tube. The samples were then centrifuged for 1 h at 100,000 g
and 4°C in a TLS 55 rotor (Beckman-Coulter) with the appropriate adapters (#358614). The top
200 µL were carefully taken off and stored at -20°C ("S100E") and the pellet was resuspended in
50 µL PBS and stored at -20°C ("P100E"). The samples were subsequently analyzed by SDS-PAGE
and western blotting to PVDF membranes (see respective sections).

**Denaturing SDS polyacrylamide gel electrophoresis (PAGE) and immunoblotting**

SDS-PAGE analysis was carried out either with 10% gels (TGX FastCast acrylamide Kit 10%; Bio-
Rad) or precast 4-15% gradient gels (Mini-PROTEAN TGX precast gel; Bio-Rad). The protein
concentration was determined using the Pierce 660 nm protein assay (Thermo-Fisher). Samples
were denatured by the addition of SDS sample buffer (62.5 mM Tris-HCl, pH 6.8, 10% glycerol,
2% SDS, 0.002% bromophenol blue, 5% β-mercaptoethanol) and boiled for 10 minutes at 95°C.
For western blotting the proteins were transferred to polyvinylidene difluoride (PVDF)

membranes using the Trans-Blot Turbo RTA Mini PVDF Transfer Kit (Bio-Rad) and the Trans-Blot
Turbo Transfer System (Bio-Rad). Membranes were blocked for one hour in 3% BSA/TBST (20
mM Tris-HCl, pH 7.4, 137 mM NaCl, 0.1% Tween 20) at room temperature. The incubation with
the primary antibody was carried out in 3% BSA/TBST either for 1h at room temperature or
overnight at 4°C. After extensive washing with TBST the membranes were incubated with the
secondary HRP-conjugated antibodies diluted in 5% skim milk/TBST for 1h at room
temperature. The membranes were again extensively washed with TBST and the ECL reagent
(GE Healthcare) was mixed and added for 1 min afterwards. Images were acquired using Vilber-
Lourmat instrument and its software.

### Lipid droplet flotation

One 10 cm dish of day 6 3T3-L1 adipocytes was used. The cells were washed with ice-cold PBS
and subsequently harvested by scraping in PBS. The cells were centrifuged at 600 g, 4°C for 5
minutes. The supernatant was discarded, and the pellet containing the cells was resuspended in
1 mL PBS supplemented with protease inhibitor (cOmplete™, Mini, EDTA-free Protease
Inhibitor Cocktail; Merck). The cells were lysed by 20 passages through a bead homogenizer (16
µm bead, Isobiotec) and the lysate was clarified at 600 g, 4°C for 10 minutes. The supernatant
was transferred to a new 1,5 mL tube and supplemented with 60% sucrose/PBS to a final
concentration of 12% sucrose. 600 µL of the lysate were transferred to an ultracentrifuge tube
(5 x 41 mm, Beckman-Coulter, #344090) and centrifuged in an MLS-50 rotor (Beckman-Coulter)
at 100,000 g, 4°C for 1h. After the centrifugation the tubes were sealed, and the lower
boundary of the floating lipid droplet fraction was marked. The tubes were flash frozen in liquid
nitrogen and the top and were collected by cutting the frozen tube. The middle fraction was
collected after thawing and the remaining pelleted fraction was washed once with PBS and
then resuspended in PBS. All samples were then stored at -20°C until further analysis by SDS-
PAGE.

### Membrane flotation

Flotation experiments were carried out using five 10 cm dishes COS7 cells. Cells were washed
with ice-cold PBS and subsequently harvested by scraping in PBS. The cells were briefly pelleted
at 600 g, 4°C for 5 minutes. The supernatant was discarded, and the pellet was resuspended in
1 mL PBS supplemented with protease inhibitor (cOmplete™, Mini, EDTA-free Protease
Inhibitor Cocktail; Merck). The cells were lysed by approx. 20 passages through a bead
homogenizer (Isobiotec, 16 µm bead) and the lysate was cleared from debris at 600 g, 4°C for
10 minutes. 100 µL supernatant was collected and stored at -20°C; the remaining supernatant
was transferred to a 1.5 mL ultracentrifuge tube (Beckman-Coulter) and centrifuged in a
TLA 100.3 rotor (Beckman-Coulter) at 100,000 g, 4°C for one hour. The supernatant was
transferred to a new 1.5 mL tube and stored at -20°C for later analysis. The pellet was
resuspended in 36 µL PBS supplemented with protease inhibitor and 10 µL were stored for later
analysis at -20°C. The remaining 26 µL were supplemented with 150 µL of 60% sucrose (w/v). To
remove the ribosomes from the membranes 24 µL puromycin (10 mg/mL) were added to a final
concentration of 2,5 M puromycin and 45% sucrose. The reaction was incubated at RT for 30
minutes. 300 µL of 35% sucrose (w/v) were added to a 600 µL ultracentrifuge tube (Beckman-
Coulter) and the resuspended pellet fraction in 45% sucrose (w/v) was loaded to the bottom of

the tube. Finally, 150 µL of 5% sucrose (w/v) were pipetted on top, the interphases were
marked, and the sample centrifuged in a MLS 50 rotor (Beckman-Coulter) at 150,000 g, 4°C for
3h. The tubes were subsequently frozen in liquid nitrogen and the fractions separated by
cutting the tube at the interface marks using a clean scalpel. The fractions that were collected
are the top fraction with 5% sucrose, the interphase between 5% and 35% sucrose, the 35%
sucrose fraction, the interphase between 35% and 45% sucrose and the bottom fraction with
45% sucrose. The protein concentration of the fractions was determined, and equal amounts
analyzed by SDS-PAGE.

**Indirect immunofluorescence**

For indirect immunofluorescence experiments cells were trypsinized and transferred either to
Ø 18 mm cover glasses (No. 1; VWR) in 12-well tissue culture plates (TPP) or to 96-well plates
(Greiner) and cultured using DMEM supplemented with 10% of the respective serum (calf
serum CS for 3T3-L1 or FBS for COS7 and differentiated adipocytes). Differentiated adipocytes
were cultured for at least two days after trypsinization to ensure proper settling of the cells.
Cells were fixed with 4% paraformaldehyde (PFA; Electron Microscopy Sciences) at 37°C for 30
minutes. After washing three times with PBS, cells were permeabilized for 15 minutes either
with 0.1% Triton-X100 in PBS, or, when PLIN2 or PLIN3 or 4 were stained in adipocytes, 0.2%
saponin and 1% CS in PBS. Cells were again washed and incubated with primary antibodies in
blocking solution (PBS supplemented with 1% CS) for 1 h at RT. After washing, the cells were
incubated with fluorescently labeled secondary antibodies in blocking solution for 1h at RT. For
lipid droplet visualization HCS LipidTOX™ Green/Red/Deep Red dyes (Thermo-Fisher,
#H34475/#H34476/3H34477, respectively) were used at 1:500 dilution in PBS. Nuclei were
counterstained using DAPI (Thermo-Fisher) at a concentration of 0.4 µL/mL for 10 minutes at
RT. Cell outline staining, if applied, was done using Atto 490LS NHS-Ester (ATTO-TEC) at a
dilution of 1:800,000 in carbonate buffer for five minutes at RT.
Cover glasses were mounted onto Superfrost Excell glass slides (Thermo-Fisher) using
Fluoromount G (SouthernBiotech). Image acquisition was performed as described under
"Microscopy".

**Determination of protein concentration**

Protein concentration was determined in 96-well plates using the Pierce™ 660nm Protein Assay
(Thermo-Fisher) according to the manufacturer's manual. A standard-curve was produced with
a BSA solution ranging from 50-2000 µg/mL.
943
**Microscopy**

Confocal fluorescent microscopy was done on a Nikon Eclipse Ti-E microscope with enhanced
CSU-W1 spinning disk (Microlens-enhanced dual Nipkow disk confocal scanner, wide view
type), a Nikon CFI PlanApo 100x oil immersion objective (NA 1.49), and a sCMOS PCO Edge 5.5
camera (PCO, 2560x2160 pixels).
For automated confocal microscopy a Yokogawa CellVoyager 7000 spinning disk confocal
microscope with enhanced CSU-X1 spinning disk (Microlens-enhanced dual Nipkow disk
confocal scanner, wide view type), a 40x air objective (NA 1,15) or 60x water immersion (NA
1,2), and three Neo sCMOS cameras (Andor, 2560x2160 pixels) were used. 108 sites were

acquired per well of a 96-well plate with seven confocal planes of 1 µm thickness per site,
which were maximum intensity projected before saving. The signals for UV (405 nm) and far-
red (640 nm) as well as green (488 nm) and Atto490LS (675 nm) signals were acquired in dual-
camera mode. Red signal (561 nm) was acquired separately.
Images for parts of Fig. 4, S4 and S2H-I were captured on an Olympus SpinSR10 spinning disc
confocal system fitted with an Olympus IX-83 frame, a Yokogawa CSU-W1 SoRa super-
resolution spinning disc module, a Photometrics Prime BSI camera and a 60× objective (1.5 NA,
UPLAPOHR60x), using Olympus cellSens Dimension software. When indicated the images were
post-processed using an OSR filter (standard) and deconvolved (constrained iterative, maximum
likelihood, 5 iterations) using Olympus cellSens Dimension software (version 3.1.1).
963

**Automated image analysis**

The acquired images were analyzed using CellProfiler (https://cellprofiler.org/). When
necessary the cell profiler pipelines were run on the high-performance cloud computing system
"ScienceCloud" at the University of Zurich. Analysis pipelines were composed of generic or
partially customized modules as described in (Battich, Stoeger, & Pelkmans, 2015). Individual
cells were segmented with nuclei as primary objects that were identified based on DAPI signal,
and cell outlines were determined from Atto490LS signal using iterative segmentation based on
a watershed algorithm and related to nuclei as secondary objects. After cell segmentation the
background of all stainings of interest was subtracted and single-cell based features like area,
shape, intensities and textures were extracted using the modules "MeasureObjectAreaShape",
"MeasureObjectIntensity", and "MeasureTexture". Subsequent analysis was done with Matlab
(Mathworks).

**Single cell classification by support vector machines (SVMs)**

Single cells were classified using user supervised machine learning with a custom Matlab script
CellClassifier (Rämö, Sacher, Snijder, Begemann, & Pelkmans, 2009). The classifier is based on
support vector machine algorithm (SVM). In general, the classifiers were trained to identify and
remove cells that were miss-segmented using nuclear area, nuclear shape, intensity, and
textures as training features. Cells which overlapped with the border of an individual imaging
site were automatically identified and removed. Furthermore, experiment specific SVMs were
trained using intensity, channel correlation and texture as training features.
984

985    **Cloning of truncations and deletion mutants**

986    All truncation- and deletion-mutants were cloned by restriction ligation into the mEGFP-C1
987    using *Hind*III and *Kpn*I restriction sites. The mEGFP-C1 plasmid was a kind gift from Michael
988    Davidson (Addgene plasmid #54759).
989

**Cloning of GFP-PLIN2-251InsIMs**

Cloning of the PLIN2 domain swap mutant carrying the core iMS (PLIN1 amino acids 238-268) of
PLIN1 was done by insertion of the iMS sequence into mEGFP-C1-PLIN2 after amino acid 251 of
the PLIN2 sequence. The insertion was done using the Q5 Site-Directed Mutagenesis Kit (NEB)

and design of respective primers using the NEBaseChanger online tool (http://
nebasechanger.neb.com/).

**Generation of stable cell line inducibly expressing GFP-PLIN1**

The stable 3T3-L1 cells with inducibe GFP-PLIN1 or GFP-PLIN1ΔΔ (Δ238-280-Δ348-370),
respective transgenes were generated by lentiviral gene transfer using the pINDUCER20
construct and Gateway™ cloning technology. The pInducer20 was a kind gift from Stephen
Elledge (Addgene #44012). The respective transgenes were first sub-cloned by TOPO cloning
into a Gateway™ entry vector (pENTR™/D-TOPO™; Thermo-Fisher) and subsequently
recombined into the pINDUCER20 vector using the Gateway™ LR Clonase™ II Enzyme mix
(Thermo-Fisher). Production of lentiviruses and transduction of cells was performed as
described below.
The cell lines were subsequently established by selecting positively transduced cells using G418
selection (1 mg/mL) for several passages.
Expression of the respective transgene was usually induced with 500 ng/mL
anhydrotetracycline (ATC) for the indicated time-frames.

1010

**Preparation of sodium oleate**

To prepare a solution of sodium oleate, 30 mL of double-distilled water were warmed up in the
microwave to approx. 40°C and 400 µL of 5 M NaOH were added. While constantly stirring, 400
µL of oleic acid (Sigma-Aldrich) were slowly added and the solution was stirred for another five
minutes (stock concentration 41 mM). If micelles where still visible, additional NaOH was added
until all micelles disappeared. The solution was sterile filtered through a 0.22 µm sterile filter,
aliquoted and stored at -20°C. Using 147 µL of the solution per 25 mL of culture medium
resulted in a final concentration of 250 µM.

1019

**Cell treatments**

Lipolysis was induced in differentiating adipocytes by addition of 1 mM forskolin in the medium
(or vehicle, DMSO) to the final concentration of 10µM, for 6 hours. Oleate uptake was done by
adding sodium oleate to the medium in the indicated concentration for 6 hours. Protein
synthesis was inhibited by incubating cells in the medium containing cycloheximide (or DMSO
control) to final concentration of 100µg/ml for the indicated time.

1026

1027　**Cover glasses acid wash**

1028　Cover glasses (Ø 18 mm; VWR No. 1) were placed in a ceramic rack and submerged in 2 M HCl
1029　for 15 minutes. The coverslips were rinsed under tap water for 30 minutes and then washed
1030　three times with double-distilled water. Finally, the coverslips were rinsed in ethanol, sterilized
1031　under UV-light for 30 minutes and stored in a closed container in a tissue culture hood.

1032

**Production of lentivirus for gene transfer**

Lentivirus for gene transfer and integration of pINDUCER constructs was produced in HEK 293T
cells (Dharmacon). The cells were cultured in DMEM supplemented with 10% FBS, 25 mM

HEPES (Sigma-Aldrich, H4034), 1 mM sodium pyruvate (Thermo-Fisher), 2 mM glutamine
(Sigma-Aldrich) and 100 µM non-essential amino acids (Thermo-Fisher) on 10 cm dishes until
reaching approx. 70% confluency. The cells were transfected using the calcium phosphate
method with the three components necessary for the lentiviral packaging: 10 µg of the
pINDUCER expression vector containing the construct of interest, 6.5 µg psPAX2 (Addgene #
12260), the viral genome packaging plasmid, and 3.5 µg pMD2.G (Addgene # 12259), encoding
for the viral envelope components were gently mixed with $H_2O$ to a total volume of 597 µl. This
was mixed with 682 µl 2x HBSS (Hank's Balanced Salt Solution), and 85 µl 2M $CaCl_2$ were added
followed by 30 min incubation at RT. The entire transfection mix was added dropwise to the
cells and incubated for approx. 12h at 37˚C, 5% $CO_2$. After that the medium was replaced by
DMEM supplemented with 5% FBS. The media containing the lentiviral particles was collected
at 24, 48, and 60 hours after changing the medium. The individual fractions were sterile filtered
through a 0.45 µm filter, pooled and stored at 4°C. The virus particles were then concentrated
to approx. 200 µL final volume using an Amicon Ultra-15 centrifugal filter unit (Merck) with a
molecular weight cut-off of 100 kDa (Amicon "Ultra – 15 centrifugal filter"; 100 kDa MW-cut
off). The filtrate was brought to 1 mL final volume with DMEM supplemented with 10% CS and
immediately used for transduction of 3T3-L1 cells as described below.
1053
**Transduction of 3T3-L1 preadipocytes with lentivirus**

3T3-L1 cells were grown in 6-well plates until 50-60% confluency before the medium was
replaced by the concentrated virus mixture supplemented with 8 µg/mL polybrene (Sigma-
Aldrich, #107689). After overnight incubation the medium was replaced by DMEM
supplemented with 10% CS. Cells were further selected or subjected to single clone dilution as
described below.
1060
**Single cell classification by support vector machines (SVMs)**

Single-cell data obtained after automated image analysis was further analyzed using
CellClassifier (Rämö, Sacher, Snijder, Begemann, & Pelkmans, 2009), a classifier software based
on SVMs for supervised machine learning. In general, for all datasets SVMs were trained to
identify and remove cells that were miss-segmented based on features like nuclear area,
nuclear shape, intensity, and texture features. Also, datasets were automatically cleansed of
cells which touch the border of an individual imaging site. Furthermore, experiment specific
SVMs were trained e. g. for identifying transfected cells or cells which show a specific
phenotype.
1070
**In vitro experiments**

The purification protocol for LDs from cells expressing fluorescently tagged LD proteins was the
same as in Ajjaji et al. 2019. Briefly, cells from 5 x 15 cm dishes were harvested, washed once in
ice-cold PBS, and lysed using a 30G needle in 1ml 20 mM Tris-EDTA buffer containing complete
protease and phosphatase inhibitor tablets at pH 7.5. To isolate LDs, 1ml of respective cell
lysates were mixed with 1 ml 60% sucrose in 20 mM Tris-EDTA buffer supplemented with
protease inhibitors, overlaid with 20%, 10% and 0% in buffered sucrose on top of one another
in 5 ml Ultra-Clear centrifuge tubes (Beckman). Gradients were centrifuged for 16 h at 100,000

x $g$ and 4 °C, using an SW60 rotor in a Beckman L8-70 centrifuge, and 300 µL were collected
from the top as the LD fraction.

1081

1082 In vitro experiments were performed in HKM buffer: 50 mM Hepes, 120 mM $K^+$OAc, and 1 mM
1083 $MgCl_2$ (in Milli-Q water) at pH 7.4. To create buffer-in-oil drops, a buffer-diluted LD fraction was
1084 mixed with an excess of triolein by vortexing, as previously done in Ajjaji et al. 2019. The
1085 shrinkage of the buffer compartment was based on water evaporation of the aqueous drops,
1086 with the proteins at their surface, during imaging for 10 to 15 min on PDMS-coated glass slides.

1087

1088

**Fluorescence recovery after photo-bleaching FRAP experiments**

For FRAP experiments, we bleach the signal on a collection of drops and monitor the signal
recovery. The background signal, e.g. from the cytosol, is removed from the recorded signal,
which was at the end normalized by intrinsic bleaching of non-bleached areas. We used
GraphPad Prism to fit the FRAP recovery curves with a non-linear regression and the
exponential « one-phase association model ». The characteristic recovery time is extracted
from the model.

1096

1097 **Transfections**

1098 When indicated Huh7 cells (60-70% confluence) were exposed for 1hr to 500 µM oleic acid
1099 coupled to BSA (1% v:v) to induce LD formation and then cells were transfected with 3 µg of
1100 plasmid DNA/mL using Polyethylenimine HCl MAX (Polysciences, Inc) following the
1101 manufacturer's instructions.

1102

1103 **Quantification and statistical analysis**

1104 Number of samples and replicates in an experiment, as well as dispersion (standard deviation,
1105 SD, standard error of the mean, SEM), are reported for each figure in the respective legends.
1106 Quantification of experimental data are described in relevant Method sections.

1107

1108

1109 **KEY RESOURCES TABLE**

| REAGENT or RESOURCE | SOURCE | IDENTIFIER |
|---|---|---|
| **Antibodies** | | |
| Mouse monoclonal c-Myc, 1:200 | Santa Cruz | Cat#: sc-40 |
| Rabbit polyclonal Calreticulin, 1:1000 | Abcam | Cat#: ab2907 |
| Mouse monoclonal FLAG, 1:200 | Sigma-Aldrich | Cat#: F3165 |
| Mouse monoclonal HA, 1:200 | Roche | Cat#: 11583816001 |
| Goat polyclonal PLIN1, 1:200 | Abcam | Cat#: ab61682 |
| Rabbit monoclonal PLIN1, 1:200 (Fig4 I-M) | Cell Signaling | Cat#: #9349 |
| Rabbit polyclonal PLIN2, 1:200 | Abcam | Cat#: ab52355 |
| Rabbit polyclonal PLIN3, 1:200 | Millipore | Cat#: ABS482 |
| Rabbit polyclonal PLIN4, 1:200 | Sigma-Aldrich | Cat#: ABS526 |
| Rabbit polyclonal Calnexin (CNX), 1:200 | Proteintech | Cat#: 10427-2-AP |
| Rabbit polyclonal CKAP4 (CLIMP63), 1:200 | Bethyl Laboratories | Cat#: A302-257A |
| Donkey anti-goat IgG Alexa Fluor 488, 1:500 | Thermo-Fisher | Cat#: A11055 |
| Donkey anti-goat IgG Alexa Fluor 568, 1:500 | Thermo-Fisher | Cat#: A11057 |
| Donkey anti-goat IgG Alexa Fluor 647, 1:500 | Thermo-Fisher | Cat#: A21447 |
| Donkey anti-mouse IgG Alexa Fluor 488, 1:500 | Thermo-Fisher | Cat#: A21202 |
| Donkey anti-mouse IgG Alexa Fluor 568, 1:500 | Thermo-Fisher | Cat#: A10042 |
| Donkey anti-mouse IgG Alexa Fluor 647, 1:500 | Thermo-Fisher | Cat#: A31571 |
| Donkey anti-rabbit IgG Alexa Fluor 488, 1:500 | Thermo-Fisher | Cat#: A21206 |
| Donkey anti-rabbit IgG Alexa Fluor 568, 1:500 | Thermo-Fisher | Cat#: A10042 |
| Donkey anti-rabbit IgG Alexa Fluor 647, 1:500 | Thermo-Fisher | Cat#: A31573 |
| Donkey anti-goat HRP, 1:2500 | Bethyl | Cat#: A50-201P |
| Donkey anti-mouse HRP, 1:5000 | Promega | Cat#: W4021 |
| Donkey anti-rabbit HRP, 1:5000 | Promega | Cat#: W4011 |
| | | |
| **Biological samples** | | |
| N/A | | |
| **Chemicals, peptides, and recombinant proteins** | | |
| DMEM | GIBCO | Cat#41965-039; CAS/RRID: N/A |
| Fetal bovine serum | Sigma-Aldrich | Cat#F7524; CAS/RRID: N/A |
| Calf serum | Sigma-Aldrich | Cat#12133C; CAS/RRID: N/A |
| OptiMEM | GIBCO | Cat#11058-021; CAS/RRID: N/A |
| Non-essential amino acids | Sigma | Cat#M7145; CAS/RRID: N/A |
| Sodium Pyruvate | GIBCO | Cat#11360 070; CAS: 113-24-6 |
| Puromycin | Invivogen | Cat#ant-pr-1; CAS: 58-58-2 |
| Geneticin G418 | Invivogen | Cat#ant-gn-1; CAS: 108321-42-2 |
| Paraformaldehyde | Electron Microscopy Sciences | Cat#15711; CAS: 50-00-0 |
| Triton X-100 | Thermo Fisher Scientific | Cat#28314; CAS: 9002-93-1 |

| | | |
|---|---|---|
| Hexadimethrine bromide | Sigma-Aldrich | Cat#107689; CAS: 28728-55-4 |
| Oleic Acid | Sigma-Aldrich | Cat#O1383-5G; CAS: 112-80-1 |
| HCS LipidTOX Red Neutral Lipid Stain | Thermo-Fisher | Cat#H34476; CAS: N/A |
| HCS LipidTOX Deep Red Neutral Lipid Stain | Thermo Fisher Scientific | Cat#H34477; CAS: N/A |
| Fluoromount-G | SouthernBiotech | Cat#0100-01; CAS: 0100-01 |
| Dexamethasone | G-Biosciences | Cat#API-04; CAS: 50-02-2 |
| 3-Isobutyl-1-methylxanthine | Sigma-Aldrich | Cat#I5879; CAS: 28822-58-4 |
| Insulin | Sigma-Aldrich | Cat#I0516-5ML; CAS: 11070-73-8 |
| Cycloheximide | Sigma-Aldrich | Cat#C7698; CAS: 66-81-9 |
| Forskolin | Sigma-Aldrich | Cat#F3917-10MG; CAS: 66575-29-9 |
| Atto490LS NHS Ester | Atto-tec | Cat#AD 490LS-31; CAS/RRID: N/A |
| Alexa Fluor 647 NHS Ester | Life Technologies | Cat#SE-af647; CAS/RRID: N/A |
| cOmplete, Mini Protease Inhibitor Cocktail | Sigma Aldrich | Cat#11836153001; CAS/RRID: N/A |
| Ionic Detergent Compatibility Reagent for Pierce | Thermo Fisher Scientific | Cat#22663; CAS/RRID: N/A |
| D-Glucose [U-14C] | Hartmann analytic | Cat#MC 144W; CAS/RRID: N/A |
| Doxycycline hyclate | Sigma-Aldrich | Cat#D9891-1G; CAS: 24390-14-5 |
| Chloroform | Rathburn Chemicals | Cat#RH1009; CAS: 67-66-3 |
| Methanol | Biosolve Chemicals | Cat#13687802; CAS: 67-56-1 |
| Propanol | Sigma-Aldrich | Cat#650447; CAS: 67-63-0 |
| | | |
| **Critical commercial assays** | | |
| Pierce 660nm Protein Assay Reagent | Thermo Fisher Scientific | Cat#22660; CAS/RRID: N/A |
| GFP-Trap Magnetic Agarose | ChromoTek | Cat# gtma-10 |
| | | |
| **Deposited data** | | |
| Mouse 3T3-L1 cells | ATCC | Cat#CL-173; RRID: CVCL_0123 |
| Monkey COS-7 cells | Tom A. Rapoport, Harvard Medical School, Boston, USA | N/A; RRID: CVCL_0224 |
| Human HEK293T cells | Dharmacon | Cat#TLP5918; RRID: CVCL_0063 |
| | | |
| **Experimental models: Cell lines** | | |
| COS-7 | Tom A. Rapoport (Harvard Medical School, Boston, USA) | |

| 3T3-L1 | ATCC | CL-173 |
|---|---|---|
| HEK293T | Dharmacon | |
| | | |

**Experimental models: Organisms/strains**

| N/A | | |
|---|---|---|

**Oligonucleotides**

| Cloning primers for GFP-PLIN1 mutations | This study | Supplementary Table |
|---|---|---|
| | | |

**Recombinant DNA**

| mEGFP-N1 | Michael Davidson Lab | Addgene 54767 |
|---|---|---|
| mEGFP-C1 | Michael Davidson Lab | Addgene 54759 |
| mCherry2-C1 | Michael Davidson Lab | Addgene 54563 |
| mCherry2-N1 | Michael Davidson Lab | Addgene 54517 |
| pINDUCER20 | Stephen Elledge Lab | Addgene 44012 |
| pINDUCER21 | Stephen Elledge & Thomas Westbrook Lab | Addgene 46948 |
| APEX2-GBP | Rob Parton Lab | Addgene 67651 |
| psPAX2 | Didier Trono Lab | Addgene 12260 |
| pMD2.G | Didier Trono Lab | Addgene 12259 |
| GFP-PLIN1 | This study | N/A |
| GFP-PLIN2 | This study | N/A |
| GFP-PLIN3 | This study | N/A |
| GFP-PLIN1 1-92, 1-237, 1-281, 1-371 | This study | N/A |
| GFP-PLIN1 192-517, 237-517, 281-517, 328-517, 347-517, 371-517 | This study | N/A |
| GFP-PLIN1-ΔΔ (Δ238-280 Δ348-370) | This study | N/A |
| GFP-PLIN2+251InsiMS | This study | N/A |
| cDNA Mouse PLIN1 | Dharmacon | MMM1013-202799812 |
| cDNA Mouse PLIN3 | Dharmacon | MMM1013-202764453 |
| | | |

**Software and algorithms**

| MATLAB | | https://www.mathworks.com/products/matlab.html |
|---|---|---|
| CellProfiler | Carpenter et al., 2006 | https://github.com/pelkmanslab/CellProfilerPelkmans |
| CellProfiler Pelkmans Lab | Battich et al., 2015 | https://github.com/pelkmanslab/CellProfilerPelkmans/tree/master/Modules |
| ImageJ | National Institutes of Health, USA | https://imagej.net/Welcome RRID: SCR_003070 |
| | | |

**Other**

| Round microscope cover glasses, 18 mm, No. 1 | VWR | Cat#631-1580; CAS/RRID: N/A |
|---|---|---|
| Nikon Eclipse Ti E with CSU-W1 spinning disk | Nikon | |
| Yokogawa CellVoyager 7000 with CSU-X1 spinning disk | Yokugawa | |
| Leica SP5 confocal laser scanning microscope | Leica | |

1110