Title: Crossroads of assembling a moss genome: navigating contaminants and horizontal gene

transfer in the moss *Physcomitrellopsis africana*

Vidya S. Vuruputoor,[1] Andrew Starovoitov,[1] Yuqing Cai,[2] Yang Liu,[2] Nasim Rahmatpour,[1]

Terry A. Hedderson,[3] Nicholas Wilding,[4] Jill L. Wegrzyn,[1,5]* Bernard Goffinet[1]*

[1]Department of Ecology and Evolutionary Biology, University of Connecticut, Storrs, CT 06269, USA

[2]State Key Laboratory of Agricultural Genomics, BGI-Shenzhen, Shenzhen, China; Key Laboratory of

Southern Subtropical Plant Diversity, Fairy Lake, China

[3]Bolus Herbarium, Department of Biological Sciences, University of Cape Town, Private Bag, 7701,

Rondebosch, South Africa

[4]Université de La Réunion, UMR PVBMT, BP 7151, chemin de l'IRAT, 97410 Saint-Pierre, La Réunion,

France; Missouri Botanical Garden, P.O. Box 299, St. Louis, MO 63166-0299, USA

[5]Institute for Systems Genomics, University of Connecticut, Storrs, CT 06269, USA

ORCID numbers:

VV: 0000-0002-5836-7054

YC: 0009-0009-3715-2482

YL: 0000-0002-5942-839X

TH: 0000-0002-3537-6599

NW: 0000-0003-4029-5387

JW:0000-0001-5923-0888

BG: 0000-0002-2754-3895

* Corresponding authors: Department of Ecology and Evolutionary Biology, University of Connecticut,

Storrs, CT 06269, USA. Email: jill.wegrzyn@uconn.edu and bernard.goffinet@uconn.edu

Running head: *Physcomitrellopsis* genome

Abstract

The first chromosome-scale reference genome of the rare narrow-endemic African moss

Physcomitrellopsis africana is presented here. Assembled from 73x nanopore long reads and

163x BGI-seq short reads, the 414 Mb reference comprises 26 chromosomes and 22,925 protein-

coding genes (BUSCO: C:94.8%[D:13.9%]). This genome holds two genes that withstood

rigorous filtration of microbial contaminants, have no homolog in other land plants and are thus

interpreted as resulting from two unique horizontal gene transfers from microbes. Further,

Physcomitrellopsis africana shares 176 of the 273 published HGT candidates identified in

Physcomitrium patens, but lacks 98 of these, highlighting that perhaps as many as 91 genes were

acquired in *P. patens* in the last 40 million years following its divergence from its common

ancestor with *P. africana*. These observations suggest rather continuous gene gains via HGT

followed by potential losses, during the diversification of the Funariaceae. Our findings

showcase both dynamic flux in plant HGTs over evolutionarily "short" timescales, alongside

enduring impacts of successful integrations, like those still functionally maintained in extant

Physcomitrellopsis africana. Furthermore, this study describes the informatic processes

employed to distinguish contaminants from candidate HGT events.

Keywords: Funariaceae; reference genome; *Physcomitrium*; Bryophyta; HGT; contamination

Article Summary

The first draft genome of the rare South African endemic moss *Physcomitrellopsis Africana* is

presented. The 414 Mb assembly contains 22,925 genes, including two uniquely horizontally

transferred genes, but lacks 97 of the microbial genes previously identified in the closely related

model, *Physcomitrium patens* - highlighting the dynamic role of HGT in the evolution of these

moss genomes and loss. This study presents best practices for contamination detection and new

insights into HGT identification.

Introduction

Horizontal gene transfer (HGT), also referred to as lateral gene transfer (Zhaxybayeva and

Doolittle 2011) is the lateral movement of genetic material between distant branches of the tree

of life. This process is ubiquitous among bacteria, facilitating rapid adaptation through exchange

of ecologically important genes (Aminov 2011). While HGT is less common in eukaryotes than

in prokaryotes, it plays a role in shaping eukaryotic evolution with about 0.04-6.49% of

eukaryotic genes originating from HGT from microbes (Van Etten and Bhattacharya 2020).

During the evolution of land plants, increasing interactions with rhizosphere microbes,

particularly bacteria and mycorrhizal fungi, have enabled occasional horizontal transfer of

functional genes between these distantly related lineages (Martin *et al.* 2017). For example, the

gene for Killer Protein 4 (KP4) was likely acquired by mosses through HGT from ascomycete

fungi (Guan *et al*. 2023). Similarly, the HET domain, common in fungal heterokaryon

incompatibility genes and involved in self/non-self-recognition, was identified as a truncated

form in *Physcomitrium patens* and may play a role in defenses against microbial pathogens (Sun

et al. 2020). Van Etten and Bhattacharya (2020) suggested that rather than being merely

anecdotal, HGT has been an important evolutionary force driving the adaptations of land plants

to new habitats and stressors throughout their evolution, as previously already hypothesized by

Yue *et al.* (2012).

Accurately detecting the taxonomic origin of genes and confirming horizontal gene

transfer (HGT) events is challenging. Many reported cases of HGT in published genomes have

been proven to be artifacts resulting from contamination that had gone undetected. For example,

the initial report that 17% of the genes in the genome of the tardigrade *Hypsibius dujardini*

originated from HGT (Boothby *et al.* 2015), has subsequently been revised to only 0.2%

(Koutsovoulos *et al.* 2016). Similarly, claims of HGT in mammals also turned out to be

erroneous due to bacterial contamination (Douvlataniotis *et al.* 2020), highlighting the need for

careful analyses and experimental validation. Systematic screening of 43 arthropod genomes by

Francois *et al.* (2020) revealed extensive bacterial contaminants, often outnumbering true

horizontal gene transfer events. For example, in the bumblebee *Bombus impatiens*, most

contaminating genes were concentrated on just 30 contaminant scaffolds. Based on the size and

number of contaminating sequences, it was concluded that the genome of the symbiont

Candidatus Schmidhempelia bombi was co-assembled with that of its host. Strategies to

confidently identify HGT, including taxonomic assignment of candidates via

BLASTp/DIAMOND, phylogenetic analysis with candidate and donor proteins, synteny analysis

of flanking genes, and quantitative PCR validation, have been presented to address these issues

(Francois *et al.* 2020).

Physcomitrellopsis africana inhabits transitional zones between grassland and forests in

coastal habitats in South Africa. It is currently the sole species of the genus, which likely should

also include several species of the paraphyletic genus *Entosthodon* (Wilding 2015). Its genome

of 414 MB, the first genome of an African bryophyte, is presented here. This resource

complements those available for *Funaria hygrometrica* and *Physcomitrium patens* (Rensing *et*

al. 2008; Kirbis *et al.* 2022) and thereby constitutes a fundamental resource to further the

reconstruction of the evolution of the genome of the model, *Physcomitrium patens* (Rensing *et*

al. 2020). Through rigorous contamination screening and verification, two genes emerge as

candidates resulting from unique HGT events. Informatically validating HGT events in

Physcomitrellopsis africana yields evolutionary insights while underscoring the need for

stringent standards to support HGT conclusions from genomics data.

Materials and Methods

Sample collection and culturing

A population of *Physcomitrellopsis africana* was sampled in October 2010 from a coastal forest

in the Eastern Cape Province of South Africa (Dwesa National Park, along trail to chalets behind

campground, coordinates 32˚18.222' S 28˚49.666', at ± sea level). The voucher specimens

collected by Goffinet (collection numbers 10326 and 10329, with J. Goffinet, T Hedderson and

N Wilding) are deposited in the CONN herbarium under accession numbers CONN00235389

and CONN00235388, respectively. Specimen 10326 (culture and long read library DNA #5074)

provided DNA for long read sequencing, whereas 10329 (RNA and BGI-Seq DNA #5075)

provided RNA and DNA for short read sequencing. Sterile cultures were first established on

Knop medium using spores from a single operculate capsule. The gametophytes were harvested,

ground, and spread on a rich sandy loam soil in PlantCon tissue culture containers (MP

Biomedicals, Solon, OH, USA), and maintained in a growth chamber under 16 h of daylight at

about 24℃.

Genomic DNA and RNA extraction

Gametophytic tissue consisting of stems and leaves, of *Physcomitrellopsis africana* was

harvested from fresh soil cultures under a dissecting microscope and ground in liquid nitrogen.

DNA was extracted by following a modified protocol by Young (2022). The quality of the DNA

sample 5074 was assessed by quantitative PCR prior to sequencing, yielding a DIN score of 7.0

and concentration of 40.9 ng/μL.

DNA for short read sequencing was extracted using the NucleoSpin Plant midi DNA

extraction kit, following the manufacturer's protocol (Macherey-Nagel, Düren, Germany). DNA

quality was evaluated using a Qubit® 3.0 Fluorometer (Thermo Fisher Scientific, USA). Total

RNA was extracted from approximately 1 g of fresh gametophytic tissue using the RNeasy Plant

Mini Kit (Qiagen, Valencia, CA, USA).

Genome and transcriptome library preparation and sequencing

DNA was prepared for long-read PromethION sequencing through a DNA repair step,

generating blunt ends, and ligating sequencing adapters followed by priming of the flow cell, as

described in the Oxford Nanopore Technologies Amplicons by Ligation (SQK-LSK109)

protocol. The HMW DNA library was sequenced on an Oxford Nanopore PromethION (Center

for Genome Innovation, UConn) using a FLO-PRO0002 flow cell. . The two short-read DNA

libraries were prepared following the methods used by Yu et al (2020), and was sequenced (150

bp PE) on two lanes of a BGISEQ-500 (BGI-Shenzhen, China). .

Approximately 1 μg of RNA was used to generate a paired-end library with an insert

fragment size of 200–300 bp of the corresponding cDNA. RNA purification, reverse

transcription, library construction and sequencing were performed at WuXi NextCode (Shanghai,

China). The captured coding regions of the transcriptome from total RNA were prepared using

the TruSeq® RNA Exome Library Preparation Kit. The two RNA libraries were sequenced on

one lane of an Illumina HiSeq 2000 (100 bp PE) at the WuXi NextCode (Shanghai, China).

Quality control of genomic and transcriptomic reads

The genomic short reads were first assessed with FASTQC v.0.11.7 (Andrews 2010). In

preparation for assembly with Haslr and Wengan, Sickle v.1.33 with a minimum quality score

threshold of 30 (-q) and a minimum length of 50 bp (-l) was employed for read trimming.

Nanoplot v.1.21.0 was used to quantify and assess the quality of the nanopore reads. To detect

potential contamination, the long reads were aligned against the pre-indexed bacterial, human

and viral databases with a metagenomic classifier, Centrifuge v.1.0.4-beta (p+h+v; min-hit

length was increased to 50 bp) (Kim *et al.* 2016). Reads aligning to the database were removed.

The quality of the transcriptomic short reads was assessed with FASTQC v.0.11.7 (Andrews

2010). The reads were trimmed with Sickle v1.33 (Joshi and Fass 2011) with a minimum quality

score threshold of 30 (-q) and a minimum length of 40 bp (-l).

Genome size estimation

A single lane of short-read genomic data from accession 5075 was employed to estimate the

genome size. The k-mer distribution was calculated using Jellyfish v2.2.6 (Marçais and

Kingsford 2011) and size estimates were processed with GenomeScope v2.0 (Ranallo-Benavidez

et al. 2020; Supplementary Fig. S1).

Transcriptome assembly

The transcriptome was independently assembled to provide protein-level evidence for the

structural annotation of the genome, using Trinity v2.6.6 (Grabherr *et al.* 2011), with a minimum

contig length of 300bp. Contigs with minimal read support, post assembly, were removed

(FPKM > 0.5) with RSEM v1.3.0 (Li and Dewey 2011). Transdecoder v3.0.1 (Haas and

Papanicolaou 2011) was used to translate the remaining contigs into Open Reading Frames

(ORFs) and remove sequences without a viable frame. To aid Transdecoder in the identification

of ORFs, searches against the Pfam database were performed with HMMER v3.1b2 (Zhang and

Wood 2003). Transdecoder annotated the putative transcripts as complete, partial, and internal.

Those without a defined start and stop codon (defined as internals) were removed

(split_frames.py). The final set of peptide sequences were functionally annotated with EnTAP

v0.8.0 (Hart *et al.* 2020) against NCBI's nr protein database and the UniProt/Swiss-prot

reference databases. EnTAP was run with contaminant filters that included bacteria, archaea,

fungi, and insecta. Transcripts with high confidence alignments to these organisms were

removed (contam_removal.py).

Genome assembly

Hybrid genome assembly, integrating both long and short-read data, was conducted with

MaSuRCA v.4.0.3 (Zimin *et al*. 2013), Wengan v.0.2 (Di Genova *et al*. 2021), and Haslr v.0.8a1

(Haghshenas *et al.* 2020). Additionally, a separate assembly using only long-reads as input was

conducted with Flye v.2.5 (with three polishing iterations) (Kolmogorov *et al.* 2019).

The Flye, Wengan, and Haslr assemblies were polished following long-read alignment

with Medaka v1.3.2 (github.com/nanoporetech/medaka). Post assembly with Wengan, the

assembly was filtered to remove scaffolds less than 3 Kb. To further improve the accuracy of the

assemblies, the hybrid assembly generated by MaSuRCA was polished with the short reads using

Pilon v.1.24 (Walker *et al*. 2014). Subsequently, the selected MaSuRCA assembly was processed

with Purge Haplotigs v.1.0 (Roach *et al*. 2018).

The Purge Haplotigs pipeline categorized the assembled scaffolds into four coverage

levels based on the distribution of mapped reads. This categorization enabled the identification

and removal of redundant sequences exhibiting low coverage, presumed to represent erroneous

duplicates given the haploid genome. Specific cutoff values of 0, 7, and 65 for the coverage

levels were selected to delineate scaffolds to be retained or discarded, based on read depths

(Supplementary Fig. S1). The term "allele" is used to describe these redundant sequences for

convenience, although technically inaccurate for a haploid genome. The coverage analysis and

purging allowed isolation of the primary genome sequence from duplications and artifacts

generated during assembly.

To evaluate the quality of the assemblies, QUAST v5.2.0 (Gurevich *et al*. 2013) and

BUSCO v4.1.2 (viridiplantae_odb10:425 single copy orthologs) (Manni *et al.* 2021) were

employed. Each assembly was also evaluated with Merqury v1.3 (Rhie *et al*. 2020).

Genome annotation

Repeat library construction and masking

The repeat library for the final MaSuRCA assembly was generated using RepeatModeler v.2.01

(Flynn *et al.* 2020) with the long terminal repeat (LTR) discovery pipeline enabled. The genome

was then soft-masked with RepeatMasker v.4.0.9-p2 using the consensus repeat library from

RepeatModeler (Smit *et al.* 2013-2015).

Structural and functional genome annotation

The RNA reads were aligned to the soft-masked MaSuRCA assembly with HISAT2 v.2.1.0

(Kim *et al.* 2019) to provide evidence for protein-coding gene prediction. Two gene prediction

analyses were run on the soft-masked assembly using BRAKER v.2.1.5 (Brůna *et al.* 2021), one

with RNA-Seq alignment evidence and one with protein evidence originating from *de novo*

assembled transcriptome. Gene predictions from both BRAKER runs were integrated with

TSEBRA v.1.0.3 (Gabriel *et al.* 2021). From this point, separate assessments were conducted on

the RNA-Seq evidence gene predictions (BRAKER) and the final TSEBRA gene predictions to

select the best approach. Putative genes were removed from both sets if they did not contain a

complete protein domain. This filter was applied with Interproscan v.5.35-74.0 (Jones *et al.*

2014) using the Pfam database v32.0 (Finn *et al*. 2014). It is worth noting that mono-exonic

genes can be the result of fragmented annotations and the target metric of 0.2 (mono:multi-exon

gene ratio) is often achieved through protein domain filters (Vuruputoor *et al.* 2022). Metrics for

the gene predictions were generated with AGAT (Dainat 2020) and BUSCO. After assessment,

the filtered BRAKER gene predictions were selected for functional annotation with EnTAP

v.0.10.8 (Hart *et al.* 2020). Functional annotation reports from EnTAP (both sequence similarity

search and EggNog taxonomy scope classifications) allowed for the identification of non-target

species scaffolds in the assembly (Huerta-Cepas *et al*. 2019).

Assembly-level contaminant filtering

Using the functional annotation results from EnTAP, contaminated scaffolds were removed.

Scaffolds with a length of 10 Kb or less, and with 40% or more of their total genes classified as

archaea, bacteria, or fungi, were removed. Additionally, scaffolds greater than or equal to 10 Kb

with 55% or more genes classified as archaea, bacteria, or fungi, were also excluded. The final

annotation was then assessed for the annotation rate using EnTAP, the mono:multi ratio using

AGAT, and BUSCO completeness.

Horizontal gene transfer candidate identification

To identify candidate HGTs in *Physcomitrellopsis africana*, protein sequence similarity searches

were conducted with Diamond v2.1.8 (Buchfink *et al.* 2021). The protein sequences of

Physcomitrellopsis africana were aligned against "donor" databases, which included sequences

from bacteria, fungi, archaea, and metazoa from NCBI's nr database. Additionally, the same

proteins were aligned to "recipient" databases containing sequences from Streptophyta,

Tracheophyta, Embryophyta, Viridiplantae, and Spermatophyta (Ma *et al*. 2022). Although these

categories are not fully exclusive, each database was utilized separately to systematically assess

presence across plants at different evolutionary divergence points.

To identify candidate genes representing putative horizontal gene transfer (HGT) events

unique to *Physcomitrellopsis africana*, the following criteria were utilized: genes were required

to have between one and four significant sequence alignments (E-value <1e-5) to microbial

donor databases, while exhibiting no significant sequence similarity to plant recipient databases.

This range of one to four microbial alignments was selected to capture potential HGTs while

avoiding ubiquitous domains shared across many microbes. The lack of hits to plant databases

was intended to enrich for *Physcomitrellopsis africana*-specific sequences, rather than those

conserved across plants through vertical inheritance. At this stage, any scaffolds containing only

bacterial or fungal genes, without any plant-related genes, were removed from the assembly.

To assess the validity of the two proposed HGT candidates, the target proteins were

independently aligned to the reference genome with Miniprot v0.7. To examine independent

transcriptomic support, the RNA reads were aligned to the genome with HISAT2 v2.1.0 and

assembled via StringTie2 v2.2.1 (Kovaka *et al*. 2019).

Analyzing HGT candidates from Physcomitrium patens

The 273 putative horizontally transferred genes (HGTs) previously identified in *Physcomitrium*

patens (Yue *et al*. 2012; Ma *et al.* 2022) were independently searched against the

Physcomitrellopsis africana and *F. hygrometrica* protein sets using DIAMOND v2.1.8

(Buchfink *et al.* 2021). DIAMOND searches were conducted with an E-value cutoff of 1e-5 and

max target sequences set to 1. Hits against *Physcomitrellopsis africana* and *F. hygrometrica*

were collected and merged to generate a summary table with *Physcomitrium patens* HGTs and

the respective top hits in each species.

Scaffolding the genome to chromosome scale

A single HiC library was prepared for *Physcomitrellopsis africana* using Proximo Hi-C kit (Phase Genomics) and sequenced as paired-end reads on an Illumina NovaSeq platform (Center for Genome Innovation, UConn). The reads were aligned to the MaSuRCA genome assembly using BWA v0.7.17 but did not provide sufficient coverage due to contamination. Instead, the *Physcomitrium patens* T2T reference genome assembly (Bi *et al.* 2024) was used to scaffold the *Physcomitrellopsis africana* assembly to chromosome-scale with RagTag v2.1.0 under default parameters, except a minimum fragment size threshold of 1 Kb was specified.

Comparative genome analyses

A comparative analysis of the protein-coding gene space was conducted with OrthoFinder v.2.5.1 (Emms and Kelly 2019) with *F. hygrometrica* (Kirbis *et al.* 2022) and *Physcomitrium patens* v3 (Lang *et al.* 2018). To provide a preliminary estimate of gene family size dynamics, gene counts from each species in the assembled orthogroups were categorized as neutral, expanded, or contracted. The first and third quartiles were calculated for the distinct gene counts within a gene family for each species. If the number of genes from a species was lower than the first quartile or higher than the third quartile, then the gene family was categorized as "contracted" or "expanded", respectively. If the number of genes did not fit with either of these two criteria, then the gene family was considered "neutral". The longest gene for each orthogroup was used to assign functional attributes to all genes in the group from the original EnTAP annotation. If the longest gene did not originate from *Physcomitrellopsis africana*, then the functional annotation was derived from either *Physcomitrium patens* or *F. hygrometrica*.

GOSeq (Young *et al.* 2010) enrichment analysis was performed in R v4.2.0. GO terms were extracted for each gene from the EnTAP run. Enrichment analysis was investigated separately for the *Biological Process* and *Molecular Function* GO categories. Paralogs of Light Harvesting Complex (LHC), Serine/Threonine-protein kinase (STN) 7, and STN8 were identified with Diamond v2.8.1 (E-value <1e-5).

Whole genome duplication analysis

Chromosome-scale genomes of *F. hygrometrica* and *Physcomitrium patens* were assessed with the reference genome generated for Ph*yscomitrellopsis africana*, with wgd (v.2.0.19) to characterize whole genome duplication events (Chen and Zwaenepoel 2023). Each species was

compared against itself. Nucleotide sequences (CDS) were used as input to Blast & Markov

clustering (MCL) (Altschul *et al.* 1997; van Dongen 2000). A Ks distribution was constructed

using the 'ksd' subcommand and 'mcl' output (Yang 2007; Katoh and Toh 2008; Price *et al*.

2010). Next, a collinearity analysis was performed with 'syn' using the structural genome

annotations (Proost *et al*. 2011). Gaussian mixture models, via 'mix', were used to generate Ks

distributions to aid in component interpretation. Model fit was evaluated with the Bayesian and

Akaike information criterion (BIC/AIC).

Results and Discussion

Sequencing and quality control of genomic reads

The single ONT run generated 16 million reads (N50: 7,518 bp; 127X coverage: 404 Mb; 116X

coverage: 440 Mb; Supplementary Table S1). Centrifuge filtering reduced this set to

approximately 14M reads (N50: 4,561 bp; 80X coverage: 404 Mb; 73X coverage: 440 Mb). The

primary contaminants included bacteria from *Xanthomonadaceae* (10.68%) followed by

Bradyrhizobiaceae (0.7%). The short-read genomic libraries (100 bp PE) generated 529 M reads.

Following trimming with Sickle, 360 M reads remained (178X coverage: 404 Mb; 163X

coverage: 440 Mb; Supplementary Table S2). Coverage estimates are provided for both the

original estimate (404 Mb) and the final assembled genome size (440 Mb).

Transcriptome assembly

A total of 36.58M RNA short reads were generated. Following quality filtering using

Trimmomatic, the dataset was reduced to 31.84M reads. The *de novo* assembly process with

Trinity yielded 143,150 contigs. After expression filtering via RSEM, the number of contigs was

reduced to 123,373. Identifying open reading frames (ORFs) in the contigs resulted in 110,852

successfully translated transcripts. The average sequence length of these translated ORFs was

803 bp, with an N50 value of 1,038 bp. To enhance the quality of the transcriptome assembly,

internal sequences and putative contaminants were removed, resulting in 74,997 total transcripts.

After removing internal sequences and contamination, the total unique sequences with a

sequence similarity search alignment are 51,982 (69.3%). An assessment of contamination

revealed that 8,720 transcripts, 16.78% of the transcriptome, originated from various sources

such as amoeba (0.06%), bacteria (0.64%), fungi (15.93%), or insecta (0.12%), as identified by

EnTAP. A further 15,500 (18.5%) sequences remained unannotated. The final BUSCO score of

the remaining transcripts was C:82%[D:35.7%]. A total of 30,657 (36.6%) transcripts aligned to

Physcomitrium patens proteins.

Genome assembly

Initial assembly: Multiple genome assembly approaches were employed to generate

comprehensive draft assemblies of the *Physcomitrellopsis africana* genome (Fig. 2). The long-

read approach, Flye, assembled 538.68 Mb in 8,388 scaffolds, with an N50 of 152.58 Kb. The

BUSCO completeness score was C:84.0%[D:12.2%] and the Merqury QV score was 21.1.

Among the hybrid approaches, Haslr produced a 295.98 Mb reference distributed across 12,738

scaffolds, with an N50 of 52.29 Kb. The BUSCO score was C:94.1%[D:13.2%], and the QV

score was 11.9. Wengan assembled a total of 466.64 Mb across 10,516 scaffolds, with an N50 of

119.42 Kb a BUSCO score of C:96.4%[D:18.8%], and a QV score of 27.4. Finally, MaSuRCA

assembled 506.22 Mb across 3,571 scaffolds, with an N50 of 381.33 Kb, (BUSCO: C:96.4%

[15.5%], and a QV score of 31.0).

Polishing and improving genome assembly:

The Medaka polished Flye assembly had a genome size of 530.57 Mb in 8,386 scaffolds, and the

N50 was decreased to 145.17 Kb. The BUSCO dropped to C: 90:3%[D:11.5%]. The QV score

decreased to 19.6. Polishing of the Haslr assembly resulted in a genome size of 295.73 Mb,

across 12,737 scaffolds. The N50 remained almost unchanged at 52.31 Kb, and the BUSCO

score reduced to C:90.6%[D:9.4%]. The QV score more than doubled to 25.2. Filtering for

scaffolds less than 3 Kb, followed by Medaka, resulted in a smaller genome size for the Wengan

assembly at 434.41 Mb across 7,639 scaffolds. The N50 increased to 132.40 Kb. The BUSCO

score was reduced to C:93.2%[D:11.8%], and the QV increased slightly to 27.5.

Polishing with Pilon had very minimal influence on the completeness, as expected, and

substantial impact on the accuracy of the MaSuRCA assembly. The polished MaSuRCA hybrid

assembly size increased slightly to 507.10 Mb across 3,590 scaffolds, with a slightly decreased

N50 of 378.43 Kb. The BUSCO completeness remained the same at C:96.4%[D:15.5%], and the

QV score increased to 35.6.

Refinement of genome assemblies with Purge-haplotigs: The MaSuRCA assembly was selected

for further refinement. This decision was based on the overall quality assessed by Merqury,

BUSCO completeness, and overall contiguity. The MaSuRCA assembly was refined with Purge-

haplotigs, which reduced the assembly length to 502.34 Mb across 3,237 scaffolds with an N50

value of 382.25 Kb. The BUSCO completeness score remained the same (e.g., 96.4%) and the

QV score was minimally reduced to 35.5. At 502.34 Mb, the assembled genome is ~100 Mb

longer than the k-mer based estimate (440 Mb, Supplementary Fig. S2).

Genome annotation

Repeat identification: RepeatModeler produced a library containing 580 unique repeats that were

used to softmask 50.22% of the final assembly (Supplementary Table S3). Of the repeats,

36.53% were composed of *Ty3/Gypsy* and 1.46% of *Ty1/Copia*. This is similar to the pattern in

Physcomitrium patens, wherein approximately 57% of the genome is composed of repeat

elements, with long terminal repeats (LTRs), particularly the *Gypsy* family, accounting for 48%

of the masked genome (Lang *et al.* 2018). These findings align with observations by Kirbis *et al.*

(2022), suggesting a common pattern regarding the activation of *Gypsy* elements in

Physcomitrellopsis africana and *Physcomitrium patens*. In contrast, *F. hygrometrica*, which

diverged from *Physcomitrium patens* 60 to 80 MYA (Medina *et al.* 2018; Bechteler *et al.* 2023),

exhibits a lower overall repeat estimate of 35%. Here, *Gypsy* elements contribute less to the LTR

content, i.e., roughly 10%, whereas *Copia* elements contribute 17%.

Protein-coding gene identification: The *Physcomitrellopsis africana* RNA-Seq library had an

overall alignment rate of 78.43%, likely due to contaminant content in the source extraction.

Regardless, a substantial set of 37M reads were retained, exceeding the minimum needed for

prediction. The first BRAKER2 predictions, using RNA-Seq evidence alone, generated 60,917

protein-coding genes with a BUSCO completeness score of C:95.8%[D:15.3]. The mono:multi

exonic ratio was 0.52. With only protein evidence, BRAKER2 generated 37,752 gene predictions

with a BUSCO score of C:46.1%[D:16.0]. Merging these predictions, as recommended by

TSEBRA, resulted in a set of 45,737 genes with a BUSCO score of C:91.3%[D:15.1], and a

mono:multi exonic ratio of 1.06. The gene prediction set generated with RNA-Seq alignment

evidence exclusively was selected for further refinement due to its higher BUSCO score and

lower mono:multi ratio compared to the merged TSEBRA transcripts. Through protein domain

filtering (InterProScan filter), the number of mono-exonic genes was further reduced from

20,081 to 12,696, producing a total of 23,561 genes (BUSCO: C:88.5%[D:13.8%]; mono:multi

ratio: 0.08).

Based on the functional annotations of the gene space, we removed 831 scaffolds and

1250 genes related to contaminants. This also reduced the assembly length to 440 Mb in 2,406

scaffolds with an N50 of 363 Kb, and an assembly BUSCO score of C:96%[D:13.9%]. This

substantial reduction led to an assembly within ~40 Mb of the original k-mer-based estimate

(e.g., 404 Mb).

The annotated protein-coding gene space in the contaminant-filtered assembly included

1,708 mono-exonic and 21,853 multi-exonic genes. A BUSCO score of C:93.8%[D:13.6%] and

mono:multi ratio of 0.08 was reported. Reciprocal BLAST conducted by EnTAP produced an

annotation rate of 78%, of which 93.6% aligned to *Physcomitrium patens* (Table 1; File S1). This

contrasts with the annotation rate of 50% in *F. hygrometrica* (Kirbis *et al*. 2022), attributed to

significant divergence time from *Physcomitrium patens,* and the lack of closely related species in

public genomic databases (Kirbis *et al*. 2020; Rahmatpour *et al*. 2021). The higher annotation

rate in this study suggests that many genes found in *Physcomitrium patens,* but not *F.*

hygrometrica, may have been acquired in the ancestor of the *Physcomitrellopsis-Entosthodon-*

Physcomitrium clade (Medina *et al.* 2019).

Scaffolding the genome assembly to chromosome-scale

The T2T assembly of *Physcomitrium patens* (V4; Bi *et al*. 2024) was used to scaffold the

Physcomitrellopsis africana genome assembly. The final assembly contained 440.65 Mb in 583

scaffolds with an N50 length of 15.04 Mb. The 26 largest pseudomolecules, corresponding to the

Physcomitrium patens V4 haploid chromosome number, covered 414.21 Mb or 94% of the 440

Mb assembly. From here on, we will refer to these pseudomolecules as chromosomes, numbered

according to their lengths. The final genome assembly had a BUSCO completeness of C:94.8%

[S:80.9%, D:13.9%], F:1.2%, M:4.0%, n:425, and a total number of 22,925 protein coding genes.

While most of the annotated gene space was retained, the BUSCO duplication score remained at

13%, compared to 11–15% for the *F. hygrometrica* and *Physcomitrium patens* genomes (Kirbis

et al. 2022). Of the 557 contigs/scaffolds that did not contribute to the 26 chromosomes, 185

(33.34%) contained at least one protein coding gene annotation from the first annotation. Among

those, three (1.6%) contained at least one gene that was annotated as a contaminant but did not

meet the original thresholds that removed full scaffolds.

Comparative genome analysis

The comparison of the genomes of *Physcomitrellopsis africana*, *F. hygrometrica*, and

Physcomitrium patens, reveals that they share 13.5K orthogroups, but also hold 187, 939, and

958 unique orthogroups, respectively. *F. hygrometrica* and *Physcomitrellopsis africana*

exclusively share more (1053) orthogroups than *Physcomitrellopsis africana* and *Physcomitrium*

patens (745) (Fig. 3A), despite the latter sharing a more recent unique common ancestor (Fig. 1).

Thus, whereas Funariaceae share a rather conserved architecture of their vegetative body even

after at least 60 MY of divergence, their gene space varies considerably (Kirbis *et al.* 2022).

Such differentiation was previously noted based on highly diverging transcriptomes

(Rahmatpour *et al.* 2019), and interpreted as reflective of strong ecophysiological adaptations,

which may be the driving force in bryophyte evolution (Glime 1990). Signatures of two ancient

whole genome duplication (WGD) events were found in both *Physcomitrium patens* and *F.*

hygrometrica. Previous studies have shown that genes related to regulation were preferentially

retained after the first WGD in *Physcomitrium patens* (Lang *et al.* 2018). Our analysis of

Physcomitrellopsis africana also retrieved two peaks, in agreement with the two WGD events in

F. hygrometrica and *Physcomitrium patens*. The Ks peaks were at 0.56 and 0.92, and the AIC

and BIC values accorded with two WGD events. To assess the gene expansion and contraction

analysis, of the putative gene families from the *Physcomitrellopsis africana* annotation, 809 and

1,694 were, respectively, categorized as expanded and contracted (File S2). Enrichment analysis

examined through Gene Ontology's *Biological Process* category revealed 167 expanded terms

and six contracted terms. In contrast, within the *Molecular Function* GO category, 62 were

contracted, and no terms were significantly expanded.

Although expanded *Biological Process* GO terms were excessively broad, orthogroup

analysis revealed a pattern of contraction in GO terms pertaining to oxidoreductase activity,

FMN binding, and serine protease activity (Fig. 3B). Whether this unique suite of downregulated

categories diagnoses photosynthetic properties of *Physcomitrellopsis africana* only or of the

expanded genus sensu Wilding (2015) remains to be tested. The observed changes in GO term

enrichment related to oxidoreductase activity, FMN binding, and serine protease activity suggest

potential differences in photosynthetic processes between *Physcomitrellopsis africana* and other

moss species.

The complement of light-harvesting complex (LHC) genes is expanded in *Physcomitrium*

patens compared to algae and vascular plants (Alboresi *et al.* 2008; Iwai *et al.* 2018). LHC

proteins bind chlorophylls and carotenoids to facilitate light absorption and energy transfer to the

reaction centers of Photosystems I and II. The LHC genes are classified into two groups: Lhca

encodes antenna proteins for PSI (LHCI) whereas Lhcb encodes antenna proteins for PSII

(LHCII). Although ancestral land plants contain several LHC homologs, further expansion

occurred in *Physcomitrium patens* after whole genome duplication events (Alboresi *et al.* 2008;

Rensing *et al.* 2008; Zimmer *et al*. 2013; Sun *et al.* 2023;). This led to a larger repertoire of LHC

genes compared to the alga *Chlamydomonas reinhardtii* and the vascular plant *Arabidopsis*

thaliana. Specific Lhca and Lhcb paralogs are present in multiple copies in the *Physcomitrium*

patens genome compared to one to two copies in *C. reinhardtii* and *A. thaliana*. The major

antenna proteins encoded by Lhcbm also show greater redundancy and diversity in

Physcomitrium patens (Iwai *et al*. 2018; Sun *et al*. 2023).

Comparing the genomes of *Physcomitrellopsis africana, F. hygrometrica* and

Physcomitrium patens reveals both conservation and divergence of LHC genes. For example,

Physcomitrellopsis africana has eight distinct Lhca genes, whereas *Physcomitrium patens* has 12

and *F. hygrometrica* has seven. Similarly, whereas *Physcomitrellopsis africana* has seven

distinct Lhcb genes, *Physcomitrium patens* has 12, and *F. hygrometrica* has nine. Finally,

Physcomitrellopsis africana has two distinct Lhcbm genes, *Physcomitrium patens* has 14, and *F.*

hygrometrica has eight. However, echoing Kirbis *et al*. (2022), more light-harvesting complex

(LHC) gene duplicates were retained in *Physcomitrium patens* than in *F. hygrometrica* and

Physcomitrellopsis africana (Supplementary Fig. S3, Kirbis *et al*. 2022). Whereas

Physcomitrium patens retained multiple LHC paralogs of possible WGD origin or gene

duplications, *F. hygrometrica* and *Physcomitrellopsis africana* may have lost some of this gene

redundancy. This pattern of differential retention is further supported by assessing the number of

paralogs and orthologs for each LHC gene family across the three genomes (Supplementary

Table S4). Although some copies of LHC genes were lost in *F. hygrometrica* and

Physcomitrellopsis africana, key STN7 and STN8 kinases involved in photosynthetic

acclimation are conserved and retained in all three genomes, suggesting retention of core light

signaling components.

460 Contaminant filtering and the identification of horizontal gene transfer events

In plants, horizontal transfer of genes, can occur at all phylogenetic depths, via

introgression following hybridization between plant species (Pfennig 2021) to acquisition of

genetic material from other domains (i.e., bacteria) or phyla of eukaryotes (e.g., fungi) (Soucy *et*

al. 2015; Husnik and McCutcheon 2018). Identifying such transfers and hence the donors, is

challenged by the occurrence of the plant's microbiome (Koutsovoulos *et al*. 2016; Francois *et*

al. 2020) and hence requires critical screening of sequencing outputs. Filtering before and after

the assembly of *Physcomitrellopsis africana* was employed, illustrating that iterative approaches

at all stages enhanced the final quality of the reference. Here, metagenomic tools (Centrifuge;

Kim *et al*. 2016), optimized for long reads, identified contaminants before assembly. Since this

initial filter did not include the short reads or assess fungal contributions in either sequence set,

further filtering was conducted after assembly and annotation. An estimated 22% of the genes

were of fungal origin and 31% were of bacterial origin. Scaffolds that contained numerous genes

originating from algae, bacteria, or fungi (ABF) were removed. Separately, the annotated gene

space was aligned to a set of "donor" and "recipient" databases. This identified 31 potential

horizontal gene transfer (HGT) events. Following the set of best practices outlined by Francois *et*

al. (2020), an additional set of 10 scaffolds was found to be contaminated as the flanking genes

of the candidates were not of plant origin. This resulted in their removal (HGT filter). Two

unique HGT candidates survived these filters, and the final scaffolding (Fig. 4).

The first HGT candidate, *Pa1_19801.1* (alignment to C-type lectin mannose-binding

isoform-like XP_032077963.1). C-type lectins are the most frequent binding sites across all

plants and animals. In vertebrates, CTLs have a function of pathogen recognition. HGTs in the

context of plant defense and immunity are not new and have been identified before (Yue *et al.*

2012; Ma *et al*. 2020). In fact, Ma *et al*. (2020) identified a lectin that was acquired from

bacteria. The second candidate, *Pa1_12964.1*, aligns to a hypothetical protein found in *Endogone*

sp. FLAS-F59071 (RUS16308.1). While there is no information on the function of this gene,

transfers from fungi are not surprising. As mentioned previously, fungal killer proteins KP4 have

been acquired by mosses and play an important role in protonemal development (Guan *et al*.

2023).

These two candidates were manually validated by comparing alignments of de novo

assembled transcriptomes. Both candidates are directly flanked by well-annotated plant (moss)

genes. Additionally, aligning the full set of *Physcomitrium patens* and *Ceratodon purpureus*

proteins to the *Physcomitrellopsis africana* scaffolds produced no alignments in proximity to the

candidates, further suggesting that these HGT candidates are specific to *Physcomitrellopsis*

africana (Fig. 5).

Analyzing HGT candidates from Physcomitrium patens: The genomes of *Physcomitrellopsis*

africana and *F. hygrometrica* were screened for the 273 putative horizontally transferred genes

(HGTs) previously identified in *Physcomitrium patens* (Yue *et al*. 2012; Ma *et al*. 2022). Ninety-

one of these genes (33%) were not found in the other two species, whereas 15 (5%) were shared

only by *Physcomitrellopsis africana* and *Physcomitrium patens*, and seven (3%) only by

Physcomitrium patens and *F. hygrometrica* (Supplementary Fig. S4). The greater number of

shared HGTs between *Physcomitrium patens* and *Physcomitrellopsis africana* likely reflects

their more recent divergence (at least 40 MYA) compared to that of *Physcomitrium patens* and

F. hygrometrica (at least 60 MYA) (Medina *et al*. 2018; Bechteler *et al*. 2023). These findings

suggest that horizontal gene transfer may have played a role in the evolution of mosses,

potentially facilitating their adaptation to diverse environments over their long evolutionary

history dating back 500 million years (Bechteler *et al*. 2023). The life cycle of mosses, with

stages where cells are exposed to microbes, such as the egg or zygote within the open

archegonium or motile sperm cells, could provide opportunities for the uptake of foreign genetic

material from bacteria or fungi (Yue *et al*. 2012; Huang 2013). These vulnerable stages raise the

hypothesis that putative microbial symbionts or associates of mosses may have been sources of

horizontally transferred genes, potentially aiding mosses in adapting to harsh environmental

conditions encountered as some of the earliest terrestrial plants.

Data availability

All scripts and data are described in doi.org/10.5281/zenodo.11094703. NCBI BioProject ID

PRJNA1020579 contains the genomic short reads and nanopore long reads (SRR26596311,

SRR26596310 and SRR26666216) and the RNA reads (SRR26586950), as well as the whole

genome assembly (GCA_036785485.1). The *de novo* assembled transcripts and annotation files
are available at dx.doi.org/10.6084/m9.figshare.25724079.

Acknowledgements

521 The authors would like to thank the Institute for Systems Genomics (ISG) at the University of
522 Connecticut, including the Center for Genome Innovation for sequencing support and the
523 Computational Biology Core for software support and access to high-performance computing.

Funding

525 This study was made possible through the US National Foundation grants DEB-0919284
526 (fieldwork), DEB-1753811 to BG, and DBI-1943371 to JW.

Conflict of interest

528 The authors declare no conflict of interest.

Author contributions

530 J.W. & B.G. designed the study. B.G., N.W. & T.H. conducted fieldwork to sample the wild
531 population. N. R., Y. C. & Y. L. generated genomic and transcriptomic data. A.S. & V.S.V.
532 conducted all analyses. V.S.V., J.W. & B.G. wrote the paper. All authors approved of the final
533 version.

Literature Cited

Alboresi, A., Caffarri, S., Nogue, F., Bassi, R., Morosinotto, T. 2008. In silico and biochemical
analysis of Physcomitrella patens photosynthetic antenna: identification of subunits which
evolved upon land adaptation. PloS One. *3*(4): e2033. doi: 10.1371/journal.pone.0002033.
Altschul, S. F., Madden, T. L., Schäffer, A. A., Zhang, J., Zhang, Z., Miller, W., Lipman, D. J.
1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search
programs. Nucleic Acids Research. 25(17): 3389–3402. doi: 10.1093/nar/25.17.3389.
Aminov, R. I. 2011. Horizontal gene exchange in environmental microbiota. Front Microbiol.
2:158. doi: 10.3389/fmicb.2011.00158.
Andrews, S. 2010. FastQC: a quality control tool for high throughput sequence data. Available

online. Retrieved May, 17, 2018.

Bechteler, J., Peñaloza-Bojacá, G., Bell, D., Burleigh, J. G., McDaniel, S. F., Christine Davis, E.,

Sessa, E. B., Bippus, A., Christine Cargill, D., Chantanoarrapint, S., Draper, I., Endara, L.,

Forrest, L. L., Garilleti, R., Graham, S. W., Huttunen, S., Jauregui Lazo, J., Lara, F.,

Larraín, J., … Villarreal A, J. C. 2023. Comprehensive phylogenomic time tree of

bryophytes reveals deep relationships and uncovers gene incongruences in the last 500

million years of diversification. Am J Bot. 110(11): e16249. doi: 10.1002/ajb2.16249

Bi, G., Zhao, S., Yao, J., Wang, H., Zhao, M., Sun, Y., Hou, X., Hass, F. B., Varshney, D.,

Prigge, M., Rensing, S., Jiao, Y., Ma, Y., Yan, J., Dai, J. 2024. Near telomere-to-telomere

genome of the model plant Physcomitrium patens. Nat Plants. 10:327–343. doi:

10.1038/s41477-023-01614-7.

Boothby, T. C., Tenlen, J. R., Smith, F. W., Wang, J. R., Patanella, K. A., Nishimura, E. O.,

Tintori, S. C., Li, Q., Jones, C. D., Yandell, M., Messina, D. N., Glasscock, J., Goldstein, B.

2015. Evidence for extensive horizontal gene transfer from the draft genome of a tardigrade.

Proc Natl Acad Sci U S A. 112(52):15976–15981. doi: 10.1073/pnas.1510461112.

Brůna, T., Hoff, K. J., Lomsadze, A., Stanke, M., Borodovsky, M. 2021. BRAKER2: automatic

eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS supported by a

protein database. NAR Genom Bioinform. 3(1):lqaa108. doi: 10.1093/nargab/lqaa108.

Buchfink, B., Reuter, K., Drost, H.-G. 2021. Sensitive protein alignments at tree-of-life scale

using DIAMOND. Nat Methods. 18(4), 366–368. doi: 10.1038/s41592-021-01101-x.

Chen, H., Zwaenepoel, A. 2023. Inference of Ancient Polyploidy from Genomic Data. *In*

Polyploidy: Methods and Protocols (pp. 3–18). New York, NY: Springer US.

Dainat J. 2020. Another Gff analysis toolkit to handle annotations in any GTF/GFF format

(v0.8.0). Zenodo. doi:10.5281/zenodo.3552717

Di Genova, A., Buena-Atienza, E., Ossowski, S., Sagot, M.-F. 2021. Efficient hybrid de novo

assembly of human genomes with WENGAN. Nat Biotechnol. 39(4):422–430. doi:

Douvlataniotis, K., Bensberg, M., Lentini, A., Gylemo, B., Nestor, C. E. 2020. No evidence for

DNA N 6-methyladenine in mammals. Sci Adv. 6(12):eaay3335. doi:

10.1126/sciadv.aay3335.

Emms, D. M., Kelly, S. 2019. OrthoFinder: phylogenetic orthology inference for comparative

genomics. Genome Biol. 20(1):238. doi: 10.1186/s13059-019-1832-y.

Finn, R. D., Bateman, A., Clements, J., Coggill, P., Eberhardt, R. Y., Eddy, S. R., Heger, A.,

Hetherington, K., Holm, L., Mistry, J., Sonnhammer, E. L. L., Tate, J., Punta, M. 2014.

Pfam: the protein families database. Nucleic Acids Research. 42(Database issue):D222–

D230. doi: 10.1093/nar/gkt1223.

Flynn, J. M., Hubley, R., Goubert, C., Rosen, J., Clark, A. G., Feschotte, C., Smit, A. F. 2020.

RepeatModeler2 for automated genomic discovery of transposable element families. Proc

Natl Acad Sci U S A. 117(17):9451–9457. doi: 10.1073/pnas.1921046117.

Francois, C. M., Durand, F., Figuet, E., Galtier, N. 2020. Prevalence and implications of

contamination in public genomic resources: A case study of 43 reference arthropod

assemblies. G3. 10(2):721–730. doi: 10.1534/g3.119.400758.

Husnik, F., McCutcheon, J.P. 2018. Functional horizontal gene transfer from bacteria to

eukaryotes. Nat Rev Microbiol. 16(2):67–79. doi: 10.1038/nrmicro.2017.137.

Gabriel, L., Hoff, K. J., Brůna, T., Borodovsky, M., Stanke, M. 2021. TSEBRA: transcript

selector for BRAKER. BMC Bioinformatics. 22(1):566. doi: 10.1186/s12859-021-04482-0.

Glime, J. M. 1990. The ecology column: Introduction. The Bryological Times. 55:5–7.

Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., Adiconis, X.,

Fan, L., Raychowdhury, R., Zeng, Q., Chen, Z., Mauceli, E., Hacohen, N., Gnirke, A.,

Rhind, N., di Palma, F., Birren, B. W., Nusbaum, C., Lindblad-Toh, K., … Regev, A. 2011.

Full-length transcriptome assembly from RNA-Seq data without a reference genome. Nat

Biotechnol. 29(7): 644–652. doi: 10.1038/nbt.1883.

Guan, Y., Ma, L., Wang, Q., Zhao, J., Wang, S., Wu, J., Liu, Y., Sun, H., Huang, J. 2023.

Horizontally acquired fungal killer protein genes affect cell development in mosses. Plant J.

113(4):665–676. doi: 10.1111/tpj.16060.

Gurevich, A., Saveliev, V., Vyahhi, N., Tesler, G. 2013. QUAST: quality assessment tool for

genome assemblies. Bioinformatics. 29(8):1072–1075. doi: 10.1093/bioinformatics/btt086.

Haas, B. Papanicolaou, A. 2012) TransDecoder (Find Coding Regions within Transcripts)

[WWW Document]. https://transdecoder.github.io/.

Haghshenas, E., Asghari, H., Stoye, J., Chauve, C., Hach, F. 2020. HASLR: Fast Hybrid

Assembly of Long Reads. iScience. 23(8):101389. doi: 10.1016/j.isci. 2020.101389.

Hart, A. J., Ginzburg, S., Xu, M. S., Fisher, C. R., Rahmatpour, N., Mitton, J. B., Paul, R.,

Wegrzyn, J. L. 2020. EnTAP: Bringing faster and smarter functional annotation to non-

model eukaryotic transcriptomes. Mol Ecol Resour. 20(2):591–604. doi: 10.1111/1755-0998.13106.

Huang, J. 2013. Horizontal gene transfer in eukaryotes: the weak-link model. Bioessays. 35(10):868–875. doi: 10.1002/bies.201300007.

Huerta-Cepas, J., Szklarczyk, D., Heller, D., Hernández-Plaza, A., Forslund, S. K., Cook, H., Mende, D. R., Letunic, I., Rattei, T., Jensen, L. J., von Mering, C., Bork, P. 2019. eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated orthology resource based on 5090 organisms and 2502 viruses. Nucleic Acids Research. 47(D1):D309–D314. doi: 10.1093/nar/gky1085.

Iwai, M., Grob, P., Iavarone, A. T., Nogales, E., Niyogi, K. K. 2018. A unique supramolecular organization of photosystem I in the moss Physcomitrella patens. Nat Plants. 4(11):904–909. doi: 10.1038/s41477-018-0271-1.

Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H., Maslen, J., Mitchell, A., Nuka, G., Pesseat, S., Quinn, A. F., Sangrador-Vegas, A., Scheremetjew, M., Yong, S.-Y., Lopez, R., Hunter, S. 2014. InterProScan 5: genome-scale protein function classification. Bioinformatics. 30(9):1236–1240. doi: 10.1093/bioinformatics/btu031.

Joshi, N. A. Fass J. N. 2011. Sickle: A sliding-window, adaptive, quality-based trimming tool for FastQ files (Version 1.33). https://github.com/najoshi/sickle.

Katoh, K., Toh, H. 2008. Recent developments in the MAFFT multiple sequence alignment program. Brief Bioinform. 9(4):286–298. doi: 10.1093/bib/bbn013.

Kim, D., Song, L., Breitwieser, F. P., Salzberg, S. L. 2016. Centrifuge: rapid and sensitive classification of metagenomic sequences. Genome Res. 26(12):1721–1729. doi: 10.1101/gr.210641.116.

Kim, D., Paggi, J. M., Park, C., Bennett, C., Salzberg, S. L. 2019. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol. 37(8):907–915. doi: 10.1038/s41587-019-0201-4.

Kirbis, A., Rahmatpour, N., Dong, S., Yu, J., van Gessel, N., Waller, M., Reski, R., Lang, D., Rensing, S. A., Temsch, E. M., Wegrzyn, J. L., Goffinet, B., Liu, Y., Szövényi, P. 2022. Genome dynamics in mosses: Extensive synteny coexists with a highly dynamic gene space. bioRxiv p. 2022.05.17.492078. doi: 10.1101/2022.05.17.492078.

Kirbis, A., Waller, M., Ricca, M., Bont, Z., Neubauer, A., Goffinet, B., Szövényi, P. 2020.

Transcriptional landscapes of divergent sporophyte development in two mosses,

Physcomitrium (Physcomitrella) patens and Funaria hygrometrica. Front Plant Sci. 11:747.

doi: 10.3389/fpls.2020.00747.

Kolmogorov, M., Yuan, J., Lin, Y., Pevzner, P. A. 2019. Assembly of long, error-prone reads

using repeat graphs. Nat Biotechnol. 37(5):540–546. doi: 10.1038/s41587-019-0072-8

Koutsovoulos, G., Kumar, S., Laetsch, D. R., Stevens, L., Daub, J., Conlon, C., Maroon, H.,

Thomas, F., Aboobaker, A. A., Blaxter, M. 2016. No evidence for extensive horizontal gene

transfer in the genome of the tardigrade Hypsibius dujardini. Proc Natl Acad Sci U S A.

113(18):5053–5058. doi: 10.1073/pnas.1600338113.

Kovaka, S., Zimin, A. V., Pertea, G. M., Razaghi, R., Salzberg, S. L., Pertea, M. 2019.

Transcriptome assembly from long-read RNA-seq alignments with StringTie2. Genome

Biol. 20(1):278. doi: 10.1186/s13059-019-1910-1.

Lang, D., Ullrich, K. K., Murat, F., Fuchs, J., Jenkins, J., Haas, F. B., Piednoel, M., Gundlach,

H., Van Bel, M., Meyberg, R., Vives, C., Morata, J., Symeonidi, A., Hiss, M., Muchero, W.,

Kamisugi, Y., Saleh, O., Blanc, G., Decker, E. L., … Rensing, S. A. 2018. The

Physcomitrella patens chromosome-scale assembly reveals moss genome structure and

evolution. Plant J. 93(3):515–533. doi: 10.1111/tpj.13801.

Li, B., Dewey, C. N. 2011. RSEM: accurate transcript quantification from RNA-Seq data with or

without a reference genome. BMC Bioinformatics 12, 323. doi: 10.1186/1471-2105-12-323.

Ma, J., Wang, S., Zhu, X., Sun, G., Chang, G., Li, L., Hu, X., Zhang, S., Zhou, Y., Song, C.-P.,

Huang, J. 2022. Major episodes of horizontal gene transfer drove the evolution of land

plants. Molecular Plant. 15(5):857–871. doi: 10.1016/j.molp.2022.02.001.

Manni, M., Berkeley, M. R., Seppey, M., Zdobnov, E. M. 2021. BUSCO: Assessing genomic

data quality and beyond. Current Protocols. 1(12):e323. doi: 10.1002/cpz1.323.

Marçais, G., Kingsford, C. 2011. A fast, lock-free approach for efficient parallel counting of

occurrences of k-mers. Bioinformatics. 27(6):764–770. doi: 10.1093/bioinformatics/btr011.

Martin, F. M., Uroz, S., Barker, D. G. 2017. Ancestral alliances: Plant mutualistic symbioses

with fungi and bacteria. Science. 356(6340):ea.ad4501. doi: 10.1126/science.aad4501.

Medina R, Johnson MG, Liu Y, Wilding N, Hedderson TA, Wickett N, Goffinet B. 2018.

Evolutionary dynamism in bryophytes: Phylogenomic inferences confirm rapid radiation in

the moss family Funariaceae. Mol. Phylogenet Evol. 120:240–247. doi:

10.1016/j.ympev.2017.12.002.

Medina, R., M. G. Johnson, Y. Liu, N.J. Wickett, A.J. Shaw B. Goffinet. 2019. Phylogenomic

delineation of Physcomitrium (Bryophyta: Funariaceae) based on nuclear targeted exons

and their flanking regions rejects the retention of Physcomitrella, Physcomitridium and

Aphanorrhegma. J Syst Evol. 57:404–417. doi: 10.1111/jse.12516.

Pfennig KS. 2021. Biased hybridization and its impact on adaptive introgression. Trends Ecol

Evol. 36(6): 488–497. doi: 10.1016/j.tree.2021.02.010.

Price, M. N., Dehal, P. S., Arkin, A. P. 2010. FastTree 2 – Approximately maximum-likelihood

trees for large alignments. PloS One. 5(3):e9490. doi: 10.1371/journal.pone.0009490.

Proost, S., Fostier, J., De Witte, D., Dhoedt, B., Demeester, P., Van de Peer, Y., Vandepoele, K.

2011. i-ADHoRe 3.0—fast and sensitive detection of genomic homology in extremely large

data sets. Nucleic Acids Research. 40(2):e11–e11. doi: 10.1093/nar/gkr955.

Rahmatpour, N., Perera, N. V., Singh, V., Wegrzyn, J. L., Goffinet, B. 2021. High gene space

divergence contrasts with frozen vegetative architecture in the moss family Funariaceae.

Mol Phylogenet Evol. 154:106965. doi: 10.1016/j.ympev.2020.106965.

Ranallo-Benavidez, T. R., Jaron, K. S., Schatz, M. C. 2020. GenomeScope 2.0 and Smudgeplot

for reference-free profiling of polyploid genomes. Nat Comm. 11(1):1432. doi:

10.1038/s41467-020-14998-3.

Rensing, S. A., Goffinet, B., Meyberg, R., Wu, S.-Z., Bezanilla, M. 2020. The moss

Physcomitrium (Physcomitrella) patens: A model organism for non-seed plants. Plant Cell.

32(5):1361–1376. doi: 10.1105/tpc.19.00828.

Rensing, S. A., Lang, D., Zimmer, A. D., Terry, A., Salamov, A., Shapiro, H., Nishiyama, T.,

Perroud, P.-F., Lindquist, E. A., Kamisugi, Y., Tanahashi, T., Sakakibara, K., Fujita, T.,

Oishi, K., Shin-I, T., Kuroki, Y., Toyoda, A., Suzuki, Y., Hashimoto, S.-I., … Boore, J. L.

2008. The Physcomitrella genome reveals evolutionary insights into the conquest of land by

plants. Science. 319(5859):64–69. doi: 10.1126/science.1150646.

Rhie, A., Walenz, B. P., Koren, S., Phillippy, A. M. 2020. Merqury: reference-free quality,

completeness, and phasing assessment for genome assemblies. Genome Biol. 21(1):245.

doi: 10.1186/s13059-020-02134-9.

Roach, M. J., Schmidt, S. A., Borneman, A. R. 2018. Purge Haplotigs: allelic contig

reassignment for third-gen diploid genome assemblies. BMC Bioinformatics. 19(1):460.

doi: 10.1186/s12859-018-2485-7.

Smit, A. F. A., Hubley, R., Green, P. 2013-2015. RepeatMasker Open-4.0. RepearMasker.

http://www.repeatmasker.org

Soucy, S.M., Huang, J., Gogarten, J.P., 2015. Horizontal gene transfer: building the web of life.

Nat Rev Genet. 16(8):472–482. doi: 10.1038/nrg3962.

Sun, G., Bai, S., Guan, Y., Wang, S., Wang, Q., Liu, Y., Liu, H., Goffinet, B., Zhou, Y., Paoletti,

M., Hu, X., Haas, F. B., Fernandez-Pozo, N., Czyrt, A., Sun, H., Rensing, S. A., Huang, J.

2020. Are fungi-derived genomic regions related to antagonism towards fungi in mosses?

New Phytol. 228(4):1169–1175. doi: 10.1111/nph.16776.

Sun, H., Shang, H., Pan, X., Li, M. 2023. Structural insights into the assembly and energy

transfer of the Lhcb9-dependent photosystem I from moss Physcomitrium patens. Nat

Plants. 9(8):1347–1358. doi: 10.1038/s41477-023-01463-4.

van Dongen, S. M. 2000. Graph clustering by flow simulation. Doctoral dissertation. Utrecht

University, Utrecht, The Netherlands.

Van Etten, J., Bhattacharya, D. 2020. Horizontal gene transfer in Eukaryotes: Not if, but how

much? Trends Genet. 36(12):915–925. doi: 10.1016/j.tig.2020.08.006.

Vuruputoor, V. S., Monyak, D., Fetter, K. C., Webster, C., Bhattarai, A., Shrestha, B., Zaman,

S., Bennett, J., McEvoy, S. L., Caballero, M., Wegrzyn, J. L. 2022. Welcome to the big

leaves: best practices for improving genome annotation in non-model plant genomes.

Applications in Plant Sciences. 11(4):e11533. doi: 10.1002/aps3.11533

Walker, B. J., Abeel, T., Shea, T., Priest, M., Abouelliel, A., Sakthikumar, S., Cuomo, C. A.,

Zeng, Q., Wortman, J., Young, S. K., Earl, A. M. 2014. Pilon: an integrated tool for

comprehensive microbial variant detection and genome assembly improvement. PloS One.

9(11):e112963. doi: 10.1371/journal.pone.0112963.

Wilding, N. 2015. Systematics, biogeography and morphological evolution in Entosthodon

Schwägr. (Bryopsida, Funariaceae) with a revision of the genus in Africa. Doctoral

dissertation. University of Cape Town, South Africa.

Yang, Z. 2007. PAML 4: Phylogenetic Analysis by Maximum Likelihood. Mol Biol Evol.

24(8):1586–1591. doi: 10.1093/molbev/msm088.

Young, L. A. 2022. Relationships among AA-genome Chenopodium diploids and a whole

genome assembly of the North American species, C. watsonii. Doctoral dissertation.

Brigham Young University, Utah, USA.

Young, M. D., Wakefield, M. J., Smyth, G. K., Oshlack, A. 2010. Gene ontology analysis for

RNA-seq: accounting for selection bias. Genome Biol. 11(2):R14. doi: 10.1186/gb-2010-

11-2-r14.

Yu, J., Li, L., Wang, S., Dong, S., Chen, Z., Patel, N., Goffinet, B., Chen, H., Liu, H., Liu, Y.

2020. Draft genome of the aquatic moss Fontinalis antipyretica (Fontinalaceae, Bryophyta).

GigaByte. 2020:gigabyte8. doi: 10.46471/gigabyte.8.

Yue, J., Hu, X., Sun, H., Yang, Y. and Huang, J., 2012. Widespread impact of horizontal gene

transfer on plant colonization of land. Nat Commun. 3(1):1152. doi: 10.1038/ncomms2148.

Zhaxybayeva, O., Doolittle, W. F. 2011. Lateral gene transfer. Curr Biol. 21(7):R242-R246.

Zhang, Z., Wood, W. I. 2003. A profile hidden Markov model for signal peptides generated by

HMMER. Bioinformatics 19(2):307–308. doi: 10.1093/bioinformatics/19.2.307.

Zimin, A. V., Marçais, G., Puiu, D., Roberts, M., Salzberg, S. L., Yorke, J. A. 2013. The

MaSuRCA genome assembler. Bioinformatics. 29(21):2669–2677. doi:

10.1093/bioinformatics/btt476.

Zimmer, A. D., Lang, D., Buchta, K., Rombauts, S., Nishiyama, T., Hasebe, M., ... Reski, R.

2013. Reannotation and extended community resources for the genome of the non-seed

plant Physcomitrella patens provide insights into the evolution of plant gene structures and

functions. BMC Genomics. 14, 1–20.a. doi: 10.1186/1471-2164-14-498.

Fig. 1. a) *Physcomitrellopsis africana* exhibits a reduced architectural complexity of the

sporophyte similar to that observed in *Physcomitrium patens*, namely a sessile, aperistomate and

cleistocarpous sporangial capsule. b) The known geographic distribution of *Physcomitrellopsis*

africana, a rare narrow endemic to the Eastern Cape Region in South Africa. **C** Phylogenetic

relationships and chronology of the evolution of *Physcomitrellopsis* based on Medina *et al*.

(2018).

Fig. 2. Workflow and statistics for read QC, genome assembly, polishing, and haplotype

phasing. Short reads generated from BGI-seq were subject to trimming and the long reads (ONT)

were filtered for contaminants with Centrifuge. Filtered and trimmed reads were utilized as input

for four different genome assembly software tools: Flye (long-read only), Haslr, Wengan, and

MaSuRCA. The green box indicates the selection of the MaSuRCA assembly for further

analysis. The final assembly was polished using Pilon and phased with Purge haplotigs. A total

of 831 scaffolds were removed as a result of contaminant filtering using EnTAP following

structural genome annotation. Ten additional scaffolds were removed after HGT candidate

assessment. The assembly was then scaffolded against the *Physcomitrium patens* V4 genome

using RagTag, capturing 94% of the total assembly across 26 chromosomes. Summary statistics

derived from Quast (N50 - scaffold/contig length at which 50% of the genome is contained in

scaffolds/contigs of this size or greater), BUSCO, and Merqury (QV - Quality Value, an

assessment of assembly contiguity) are displayed for each process.

Fig. 3. a) The total shared and unique orthogroups among *Funaria hygrometrica*,

Physcomitrellopsis africana, and *Physcomitrium patens*. b) Enriched *Molecular Function* GO

terms for gene families in a comparative analysis between *Physcomitrellopsis africana*,

Physcomitrium patens, and *F. hygrometrica,* showed contraction in the GO terms related to

oxidoreductase activity, as well as serine peptidase activity and FMN binding for

Physcomitrellopsis africana. The size of each bubble represents the number of gene families

within an orthogroup, and the gradient of the color denotes the significance level of enrichment-

the darker green denotes more significance.

Fig. 4. a) Contamination versus horizontal gene transfer (HGT) in the *Physcomitrellopsis*

africana genome. Parameters used for removing a set of contaminated scaffolds from the draft

genome based on functional characterization of the annotated gene space. Scaffolds with a length

of 10 kb or less, and those with 40% or more of their total genes classified as archaea, bacteria,

or fungi were removed. Additionally, scaffolds with a length greater than or equal to 10 Kb and

having 55% or more genes classified as archaea, bacteria, or fungi were also excluded. In total,

831 scaffolds were removed because of this filtering process. Sixty-three scaffolds contained at

least one gene annotated as a contaminant but did not meet the original thresholds for removing

full scaffolds. When the chromosome-scale assembly was generated with RagTag scaffolding,

three of these scaffolds were not placed in the 26 chromosomes. b) Identification of HGT

candidates via sequence similarity comparisons of *Physcomitrellopsis africana* proteins against

both "donor" databases (archaea, bacteria, fungal, and metazoan) and "recipient" databases

(Streptophyta, Tracheophyta, and Spermatophyta). Proteins with >1 and <4 hits against all donor

databases, and no hit against recipient databases were labeled as HGT candidates. This analysis

was conducted on the contaminated scaffolds removed in A, confirming their contamination

status. On the post-filtered scaffolds, there were some putative HGTs that were flanked by

contaminants. These scaffolds were also removed from the assembly (additional 10), resulting in

the retention of two HGT candidates in the final analysis.

Fig. 5. Integrated Genome Viewer (IGV) screens depicting five tracks of the *Physcomitrellopsis*

africana genome. Track 1 shows the protein-coding structural annotation in context to the

genome. Track 2 displays genome-guided transcript assemblies via StringTie2. Track 3

illustrates the alignment of the Horizontal Gene Transfer (HGT) candidate (nr database) to the

genome. Tracks 4 and 5 show the alignments of *Physcomitrium patens* and *Ceratodon purpureus*

proteins onto chromosomes 2 and 26. a) and b) show the HGT candidates *Pa1_04828.1* and

Pa1_22336.1, where the HGT candidate alignment validates the presence of a protein. In both

cases, independent RNA assemblies, via StringTie2, were not able to generate a supporting

model. None of the other moss proteins align to the HGT candidates. c) highlights the example

of *Pa1_15362.1*, where the StringTie2 transcript and the protein alignment reveal that the gene

spans two gene models, indicating a false identification as an HGT candidate. Further

transcriptomic evidence is required to modify the annotation model and establish the validity of

these gene models.

Table 1. Genome annotation statistics for *Physcomitrellopsis africana*.

Annotation Method	Total genes	BUSCO (viridiplantae)	Mono: multi ratio	Annotation rate
BRAKER (RNA)	60,917	C:95.8%[D:15.3]	0.52	65%
BRAKER (protein)	37,752	C:46.1%[D:16.0]	0.95	73%
TSEBRA (BRAKER RNA + BRAKER protein)	45,737	C:91.3%[D:15.1]	1.02	74%

BRAKER (RNA) + InterProScan filter + scaffold contam filter)	23,561	C:93.8%[D:13.6%]	0.08	78%
BRAKER (RNA) + InterProScan filter + scaffold contam filter + HGT filter)	23,535	C:88.5%[D:11.3%]	0.07	83%
Scaffolded to 26 chromosomes with RagTag	22,925	C:86.4%[D:11.1%]	0.07	75%

815
816

(a)

(b)

(c)

Physcomitrium

Entosthodon

Physcomitrellopsis

Afoninia

Funaria

Pyramiduloidae

Funarioidae

80   60   43   30   0 mya

(a)

Contaminated scaffolds

Gene   Gene   Gene

Filter parameters
<=10kb **length** and =>40% **classified ABF**
>=10kb **length** and =>55% **classified ABF**

Filtered scaffolds

Gene   Gene   Gene

Scaffolds reduced from 3,237 to 2,396

Overall # of scaffolds removed: 831 + an additional 10 scaffolds

Borderline scaffolds: 63

Chromosome-scale assembly

Gene   Gene   Gene

26 chromosomes

Total borderline scaffolds removed: 3

(b)

Donor databases

diamond

>1 and <4 hits against donor

0 hits against recipients

Recipient databases

HGT candidates

Gene   Gene   Gene

Confirmation of contamination: 11 HGT candidates on contaminated scaffolds

Gene   Gene   Gene

Additional 10 scaffolds removed: 17 HGT candidates on filtered scaffolds flanked by contaminants

Gene   Gene   Gene

2 HGT candidates on filtered scaffolds flanked by plant genes

The HGT candidates were retained in the chromosome-scale assembly

(a)

Funaria hygrometrica 939

Physcomitrium patens 958

1697

13583

1053

Physcomitrellopsis africana 187

(b)

GO molecular function

amino acid binding
enzyme binding
FMN binding
Iron ion binding
oxidoreductase activity
peptidase activity
serine hydrolase activity
serine-type endopeptidase activity
serine-type peptidase activity
structural constituent ribosome
structural molecule activity

Contracting

Total genes
50
100
150
200

-log10(FDR)
0.015
0.010
0.005
0.000

(a)

(b)

(c)

Physcomitrium

Entosthodon

Physcomitrellopsis

Afoninia

Funaria

Pyramiduloideae

80 60 43 30 0 mya

Funarioideae