WTAP is Transcriptionally Regulated by p65 and Promotes Inflammation through m^{6}A Modification and Phase Separation

Yong Ge[1,2,†], Rong Chen[1,2,†], Tao Ling[1,2], Biaodi Liu[1], Jingrong Huang[1,2], Youxiang Cheng[1,2], Yi Lin[1,2], Hongxuan Chen[1], Xiongmei Xie[1], Guomeng Xia[1], Guanzheng Luo[1], Shaochun Yuan[1,2,4,✉] & Anlong Xu[1,3,✉]

[1]Guangdong Province Key Laboratory of Pharmaceutical Functional Genes, MOE Key Laboratory of Gene Function and Regulation, State Key Laboratory of Biocontrol, School of Life Sciences, Sun Yat-Sen University, Guangzhou, China.

[2]Southern Marine Science and Engineering Guangdong Laboratory (Zhuhai), Zhuhai, China.

[3]School of Life Sciences, Beijing University of Chinese Medicine, Beijing, China.

[4]Laboratory for Marine Biology and Biotechnology, Pilot National Laboratory for Marine Science and Technology, Qingdao, China.

†These authors contributed equally.

✉Correspondence: lssxal@mail.sysu.edu.cn (A.X.); yuanshch@mail.sysu.edu.cn (S.Y.)

Abstract

Emerging evidence has linked dysregulation of N^6-methyladenosine (m^{6}A) to inflammation and

inflammatory diseases, but the underlying mechanism remains lacking. Here, we found that high

m^{6}A modification in a variety of hyperinflammatory states is p65-dependent, because WTAP, a

key component of the writer complex, is transcriptionally regulated by p65 and its overexpression

can lead to higher m^{6}A modification. Mechanistically, upregulated WTAP is more prone to phase

separation to facilitate the aggregation of "writer" complex to nuclear speckles and the deposition

of m^{6}A marks onto transcriptionally active inflammatory transcripts, thereby accelerating

proinflammatory response. Furthermore, myeloid deficiency of WTAP attenuates the severity of

LPS-induced sepsis and DSS-induced IBD. Thus, the proinflammatory effect of WTAP is a

general risk-increasing mechanism, and interrupting the assembly of m^{6}A writer complex by

targeting the phase separation of WTAP to reduce the global m^{6}A level may be a potential and

promising therapeutic strategy for alleviating hyperinflammation.

Introduction

Inflammation is usually a physiological healing response that is triggered by noxious stimuli and conditions, such as infection and tissue injury (Cao, 2016). Moderate inflammation is essential for the pathogen clearance, tissue repair and regeneration. However, dysregulated inflammation disrupts immune homeostasis, which may lead to the development of a variety of inflammatory diseases, such as chronic inflammation, metabolic disorders, autoimmune diseases (ADs) and cancer (Gong *et al*, 2020; Greten & Grivennikov, 2019). To maintain immune homeostasis and prevent harmful outcomes, the activation or induction of proinflammatory regulators and effectors must be precisely controlled at multiple levels, including the transcriptional (Medzhitov & Horng, 2009), posttranscriptional (Carpenter *et al*, 2014) and posttranslational (Liu *et al*, 2016) levels, and these are thus potential targets for the treatment of inflammatory diseases (Banerjee *et al*, 2017; Yeung *et al*, 2018). The therapeutic strategies for inflammatory diseases have been established and even become invaluable by developing antagonists or inhibitors against principal inflammatory effectors (NF-κB, STAT3 and JAK) and multifunctional proinflammatory cytokines (IL-6, TNF-α, IL-17A and IL-23) (Bradley, 2008; Hunter & Jones, 2015), such as specific antibodies against IL-6(R) (e.g., tocilizumab and siltuximab) and TNF-α (e.g., infliximab, etanercept and adalimumab), which have been used in the treatment of many autoimmune diseases (Kang *et al*, 2019). However, due to pleiotropy, the therapeutic effects of targeting specific cytokines such as IL-6 or IL-17A greatly vary among different inflammatory diseases (Stone *et al*, 2017). In contrast, small-molecule inhibitors targeting the NF-κB and STAT3 signalling pathways lead to significant side effects that alter the homeostasis of the immune system as well as the nonimmune cells (Siegel *et al*, 2011; Zhang *et al*, 2017). Therefore, in-depth research into how inflammation is precisely regulated remains particularly important for the development of new treatments against excessive inflammatory responses.

In the past two decades, studies on immune regulation have mainly focused on posttranslational modifications of proteins, such as phosphorylation and ubiquitination. Recently, emerging evidence has strongly indicated that immune signalling can trigger dynamic alterations to the epitranscriptome, which orchestrates the regulation of immune response (Li & Rana, 2022; O'Connell *et al*, 2015). Among those dynamic epitranscriptomics alterations, m^6A modification is the most abundant and reversible RNA modification and can influence pre-mRNA splicing,

stability, translation, location and transport (Fu *et al*, 2014). The m^{6}A modification is mediated by

a methyltransferase complex called "writer" and removed by demethylase, namely "eraser". In

addition to playing roles in distinct biological processes such as embryonic development,

hematopoiesis and cancer, the roles of m^{6}A modification in inflammatory regulation have recently

attracted intense attention (Luo *et al*, 2021; Shulman & Stern-Ginossar, 2020). For example,

depleting METTL3, the key methyltransferase in the "writer" complex, reduces the m^{6}A

abundance on *TRAF6* mRNA and decreases the expression of TRAF6 by entrapping the

transcripts in the nucleus, which inhibits the activation of the NF-κB and mitogen-activated

protein kinase (MAPK) signalling pathways (Zong *et al*, 2019). In contrast, another study revealed

that METTL3 is significantly elevated in patients with rheumatoid arthritis (RA) and that high

expression of METTL3 attenuates the inflammatory response in macrophages which are

stimulated with LPS (Wang *et al*, 2019). Not only METTL3, but also inconsistent results have

been obtained in functional studies of demethylases. For instance, knocking-down the

demethylase FTO leads to an increased overall m^{6}A abundance, STAT3 phosphorylation and

secretion of proinflammatory cytokines (Tang *et al*, 2022), whereas the knockdown of ALKBH5

inhibits cell viability and production of inflammatory cytokines (Zhu & Lu, 2020). In addition to

the m^{6}A writers and erasers, m^{6}A readers, such as YTHDF2, play different or even opposite roles

in regulating different inflammatory events. For example, a previous study showed that YTHDF2

limits the decay of *KDM6B* mRNA in a m^{6}A-dependent manner and promotes the demethylation

of histone 3 lysine 27 trimethylation (H3K27me3), which is required for the transcription of

certain proinflammatory cytokines (Wu *et al*, 2020). However, another study indicated that

YTHDF2 is an inflammatory suppressor that downregulates m^{6}A-modified transcripts in

inflammatory response and thereby protects hematopoietic stem cells from excessive

proinflammatory signals (Mapperley *et al*, 2021). Thus, although notable evidence clearly points

to m^{6}A modification in modulating inflammatory responses, its effects appear to depend greatly on

the disease status or specific target genes.

Recently, due to the finding of aberrant RNA methylation in inflammatory diseases (Luo *et al*.,

2021) and cancers (Jiang *et al*, 2021), small-molecule inhibitors, which target specific "writer" or

"eraser" to reprogram the m^{6}A epitranscriptome, have attracted considerable attention and have

been proven to be feasible (Barbieri & Kouzarides, 2020). Notably, a research group identified a

selective METTL3 catalytic inhibitor STM2457, that can suppress the growth of acute myeloid

leukemia (AML) (Yankova *et al*, 2021). Moreover, two FTO inhibitors (FB23 and FB23-2) have

been designed to bind FTO directly and inhibit FTO demethylase activity, and exert a selective

inhibitory effect on the proliferation of AML cells (Huang *et al*, 2019). In addition to the

aforementioned inhibitors used to treat AML, potential inhibitors of m^{6}A methylase and

demethylase remain in the research stage for the treatment of other inflammatory diseases.

Because m^{6}A methylation contributes to pluripotency, identification of the intrinsic principles

underlying the mechanism through which m^{6}A regulates inflammation and inflammatory diseases

is extremely important before a therapeutic approach can be developed to target the m^{6}A mark.

The investigation of other general regulators, in addition to m^{6}A methylase and demethylase

inhibitors, may provide another method for developing effective and even safer treatments for

inflammatory diseases.

By revealing the basic principle of m^{6}A modification in inflammatory regulation, we report a new

function of the m^{6}A "writer" protein Wilms' tumour 1-associating protein (WTAP) in controlling

inflammatory responses and associated diseases. We found that WTAP is an NF-κB p65-

stimulated gene and that its expression is significantly upregulated in response to a variety of

inflammatory stimuli and in many inflammatory diseases. Mechanistically, after an increase in its

concentration, WTAP spontaneously undergoes phase separation, which facilitates the aggregation

of the "writer" complex to nuclear speckles and the deposition of m^{6}A onto transcriptionally active

proinflammatory genes. Hence, WTAP enhances the protein synthesis of many m^{6}A-modified

proinflammatory transcripts, including IL6ST, IL18R1, IL15RA, IL-1A, IL12B and CXCL11, in

response to inflammatory stimuli, which accelerates the process of inflammatory responses and

aggravates the severity of many inflammatory diseases, such as LPS-induced sepsis, IBD and

psoriasis. Thus, our study here identified WTAP as a risk factor in inflammatory responses and

thereby provides mechanistic insights into WTAP as a novel and potential therapeutic target for

preventing excessive inflammation.

Results

Hyperinflammation is accompanied by elevated WTAP levels in many inflammatory diseases

Although some evidence has indicated that m^{6}A modification plays an essential role in inflammation and inflammatory diseases (Cui *et al*, 2022; Xu *et al*, 2022), the intrinsic principles remain unclear. By analyzing data obtained from the Gene Expression Omnibus (GEO) dataset (GSE13887/137268/69063/97779/166388/208303, Appendix Table S1), we found that the expression of WTAP is commonly upregulated in patients with SLE, asthma, sepsis, rheumatoid arthritis, psoriasis, or Crohn's disease (Fig EV1A-F). To verify this phenomenon, we then performed statistical analyses of previously published microarray datasets (GSE19315/198326/2411/2638/227851/189847, Appendix Table S1) and further identified that only the mRNA abundance of WTAP was significantly increased after LPS stimulation in THP-1 cells (Fig EV1G) and human macrophages (Fig EV1H). Similar results were obtained from the lung tissue of mice with LPS-induced lung injury (Fig EV1I). In addition, we found that the mRNA abundance of WTAP also increased in TNF-α stimulated human microvascular endothelial cells (HMECs) (Fig EV1J), *Mycobacterium tuberculosis* (H37Rv)-infected THP-1 cells (Fig EV1K) and *Salmonella typhimurium* (SL1344)-infected macrophages (Fig EV1L). These observations suggest that the overexpression of WTAP is a common phenomenon in hyperinflammation states.

To verify the abovementioned observations, THP-1 cells were treated with the Toll-like receptor 4 (TLR4) agonist LPS, an ideal reagent to activate the inflammatory signalling cascade *in vivo* (Triantafilou & Triantafilou, 2002). RNA-sequencing (RNA-seq) analysis was then performed and the results confirmed that only the mRNA abundance of *WTAP* was significantly upregulated in the LPS-stimulated THP-1 cells (Fig 1A). Further qRT–PCR (Fig 1B-D) or immunoblotting (Fig 1E-G) analyses conducted with THP-1 cells, peripheral blood mononuclear cells (PBMCs) and mouse bone marrow-derived macrophages (BMDMs) after LPS stimulation confirmed that the expression of WTAP was upregulated. To determine whether the upregulated expression of WTAP is ubiquitous upon specific inflammatory stress, we treated THP-1 cells and BMDMs with different TLR agonists or heat-killed bacteria, such as CL097 (a TLR7/8 ligand), Pam3CSK4 (a TLR1/2 ligand), heat-killed *Salmonella typhimurium* (HKST) and heat-killed *Listeria monocytogenes* (HKLM), and found that the mRNA and protein abundance of WTAP were both

significantly increased (Fig 1H-M). Thus, high WTAP expression is positively correlated with

hyperinflammation states, implying a potential role of WTAP in the regulation of inflammation.

WTAP is an NF-κB p65-stimulated gene

Protein abundance can be regulated at the transcriptional, translational, or posttranslational level.

To reveal the specific mechanism underlying the upregulation of WTAP at both the mRNA and

protein levels upon inflammatory stimulation, bioinformatics analyses using the Promoter 2.0

prediction server (http://www.cbs.dtu.dk/services/promoter/), CpG plot

(http://www.ebi.ac.uk/Tools/seqstats/emboss_cpgplot/) and JASPAR (http://jaspar.genereg.net/)

were performed to identify the region between −800 and +250 in the genomic sequence of human

WTAP containing the TATA box, CAAT box and GC box, which are characteristic of promoters.

Further analyses of transcription factor-binding sites revealed that the *WTAP* promoter contains

NF-κB p65-, C/EBPβ- and IRF3-binding motifs, and among these, the p65-binding motifs exhibit

the highest prevalence and are distributed in a relatively concentrated and overlapping region (Fig

2A). Similar results were obtained from an analysis of the mouse *Wtap* promoter region (Fig 2B).

As shown in Fig 2A and B, we constructed a series of reporter plasmids containing the wide-type

(WT) WTAP promoter with NF-κB p65-, C/EBPβ-, IRF3-binding motifs or mutated or deleted

p65-binding motifs based on the pGL3 basic construct. These reporter plasmids were then

cotransfected with increasing amounts of NF-κB p65-, C/EBPβ- or IRF3-expressing plasmids into

293T cells. The results revealed that that NF-κB p65 but not IRF3, increased the expression of the

respective *WTAP* promoter reporter in a dose-dependent manner (Figs 2C and EV2A and B).

However, deletion or mutation of the core p65-binding sites in the *WTAP* promoter inhibited the

expression of the reporter gene (Fig 2D). Similar results were obtained using reporter plasmids

that contained the mouse *Wtap* promoter (Fig EV2C). Furthermore, the treatment of THP-1 cells

or PBMCs with the p65 inhibitor PG490 (triptolide) inhibited the induction of WTAP by LPS or

Pam3CSK4 (Figs 2E-H and EV2D and E). Using $RELA^{-/-}$ (encoding p65) THP-1 cells generated

using the CRISPR-mediated genome editing approach (Fig EV2F), we found that the upregulation

of WTAP was inhibited in $RELA^{-/-}$ THP-1 cells after LPS, Pam3CSK4, HKST or HKLM

stimulation (Figs 2I-L and EV2G and H). DNA pull-down assays confirmed the direct binding of

Flag-tagged p65 to biotin-labelled p65-binding probes of the *WTAP* promoter *in vitro* (Fig 2M and

N). In addition, C/EBPβ can also slightly activate the expression of WTAP (Fig EV2A), and DNA

pull-down assays confirmed the direct binding of C/EBPβ to the *WTAP* promoter (Fig EV2I).

Because p65 can further activate some inducible transcription factors such as ATF3, C/EBPδ and

C/EBPβ, to enhance the LPS-induced transcriptional response (Medzhitov & Horng, 2009), we

hypothesized that although C/EBPβ may play some role in the transcription of *WTAP*, the

upregulation of *WTAP* predominantly depends on the activation of NF-κB p65.

WTAP positively regulates the expression of proinflammatory cytokines

To determine the significance of WTAP in inflammatory responses, we first generated *WTAP*-

knockout THP-1 cells using the CRISPR/Cas9 approach. Two sgRNAs were designed to target the

second and third exons (E2 and E3) of *WTAP*. sgRNA1 caused two bases insertion, whereas

sgRNA2 caused one base deletion in the *WTAP* genome locus (Fig EV3A and B), leading to the

introduction of a new stop codon upstream of the CDS and resulting in an early termination of

WTAP translation. However, with open reading frame (ORF) prediction, it was found that the cells

may skip these mutations by alternative translation initiation to produce alternative WTAP

isoforms with 376 and 319 aa, respectively (Fig EV3C). A subsequent immunoblotting assay

using a monoclonal antibody (mAb) purchased from Abcam confirmed the extremely low

expression of these two isoforms (Fig EV3D). Thus, the cell lines were designated as "$WTAP^{\Delta1\text{-}20}$

THP-1 cells" and "$WTAP^{\Delta1\text{-}77}$ THP-1 cells". In addition to generating KO THP-1 cells, we also

generated the *Wtap* conditional KO (CKO) mice by crossing $Wtap^{\text{flox/flox}}$ mice with mice

expressing Cre recombinase under the control of the lysozyme 2 promoter (*LyzM*-Cre) (Fig

EV3E). Genomic sequencing confirmed that the third exon was fully removed after crossing,

which also resulted in early termination of WTAP translation (Fig EV3F and G). Similarly, ORF

prediction showed that gene-edited BMDMs may also produce alternative WTAP isoforms with

319 aa (Fig EV3G). Because the WTAP mAb purchased from Abcam did not recognize mouse

WTAP, we used a polyclonal antibody (pAb) purchased from Proteintech to detect the WTAP

isoforms in mice, and the results showed that only WT WTAP could be identified (Fig EV3H and

I).

It has been demonstrated that WTAP possesses a nuclear localization signal (NLS) at its N

terminus (Fig EV3C), and mutation of this signal can affect the entry of the protein into the

nucleus and its function (Scholler et al, 2018). Since the NLS is missing in both alternative

isoforms, their entry to the nucleus may be affected. The nuclear-cytoplasmic extraction assays

confirmed the extremely low concentrations of WTAP$^{\Delta 1\text{-}20}$ and WTAP$^{\Delta 1\text{-}77}$ both in the cytoplasm

and nucleus (Fig EV3J). Moreover, LC–MS/MS assays indicated that ectopic expression of WT

WTAP but not the two alternative isoforms in $WTAP$ KO 293T cells could increase the global m^{6}A

modification level (Fig EV3K). Similarly, the m^{6}A modification levels of $WTAP^{\Delta 1\text{-}20}$ and $WTAP^{\Delta 1\text{-}}$

^{77} THP-1 cells were significantly lower than those of wild-type cells (Fig EV3L). Although we

could not conclude that the WTAP$^{\Delta 1\text{-}20}$ and WTAP$^{\Delta 1\text{-}77}$ isoforms are not functional at all, we found

that their expression is extremely low and that their function is likely severely impaired due to the

lack of NLS, therefore, the status of $WTAP^{\Delta 1\text{-}20}$ and $WTAP^{\Delta 1\text{-}77}$ THP-1 cells should be very close to

that of cells in which the protein is completely knocked out. Moreover, the m^{6}A abundance in

BMDMs from $LyzM$-Cre$^+$ $Wtap^{\Delta 1\text{-}77}$ mice was decreased significantly, consistent with the results

observed in THP-1 cells (Fig EV3M). Thus, although we do not have another antibody to further

confirm whether the WTAP$^{\Delta 1\text{-}77}$ is present in $Wtap$ KO BMDMs, we can assume that even if

WTAP$^{\Delta 1\text{-}77}$ exists, its expression should be extremely low and that its function is impaired in CKO

mice. Additionally, no differences in the proportion of major immune cell populations were

observed between $Wtap^{fl/fl}$ and $LyzM$-Cre$^+$ $Wtap^{\Delta 1\text{-}77}$ mice in steady state (Fig EV4A-C), indicating

that the depletion of $Wtap$ in myeloid cells did not affect macrophage development and

maturation.

After verifying the status of WTAP KO THP-1 cells and CKO BMDMs, we next performed RNA-

seq analyses with WT and $WTAP^{\Delta 1\text{-}77}$ THP-1 cells before and after LPS treatment. KEGG

enrichment analyses showed that the transcripts downregulated in $WTAP^{\Delta 1\text{-}77}$ THP-1 cells were

mainly enriched in the cytokine production and inflammatory signalling pathways (Fig 3A and B).

In contrast, KEGG analyses of the genes upregulated in $WTAP^{\Delta 1\text{-}77}$ THP-1 cells showed that there

were enriched in a few pathways that are not associated with inflammatory responses (Fig EV5A).

To confirm the results obtained by RNA-seq, qRT–PCR analyses were performed and revealed

marked reductions in $IL6$, $CCL2$, $CCL8$ and $CXCL8$ expression in $WTAP^{\Delta 1\text{-}77}$ THP-1 cells after

stimulation with LPS (Fig 3C). Similar results were obtained with $WTAP^{\Delta 1\text{-}77}$ THP-1 cells treated

with Pam3CSK4, HKST or HKLM (Fig EV5B). Consistently, $WTAP^{\Delta 1\text{-}77}$ THP-1 cells exhibited a

significant reduction in IL-6 secretion in response to the aforementioned stimuli compared with

the WT cells (Fig 3D). Furthermore, ectopic expression of WTAP (Fig EV5C) substantially

facilitated the expression of IL-6 under inflammatory stimulation (Fig EV5D). These results

confirmed that WTAP accelerates inflammatory responses by promoting the expression of many

proinflammatory cytokines during multiple inflammatory stimuli.

WTAP aggravates LPS-induced sepsis in mice

Because WTAP was upregulated after LPS stimulation and because its deficiency can reduce

inflammatory responses in macrophages in THP-1 cells, we next examined the biological effects

of WTAP on the progression of inflammatory diseases in *LyzM*-Cre$^+$ *Wtap*$^{\Delta 1\text{-}77}$ mice. BMDMs first

isolated from *Wtap*$^{\text{fl/fl}}$ and *LyzM*-Cre$^+$ *Wtap*$^{\Delta 1\text{-}77}$ mice were treated with LPS and then used for

RNA-seq analyses. The results showed that WTAP depletion in BMDMs exhibited a significant

reduction in the expression of proinflammatory cytokines upon LPS stimulation (Fig 3E), which is

consistent with the trend observed in THP-1 cells. As mentioned above, the *WTAP* mRNA

abundance was significantly higher in the PBMCs of sepsis patients than in those of healthy ones

(Fig EV1C). A retrospective study of publicly available data also revealed that WTAP is a risk

gene for sepsis. Specifically, its expression was significantly correlated with 28-day cumulative

mortality (Zhang *et al*, 2020). Thus, we further investigated the biological effects of WTAP using

LyzM-Cre$^+$ *Wtap*$^{\Delta 1\text{-}77}$ and *Wtap*$^{\text{fl/fl}}$ mouse models of LPS-induced sepsis. Consistent with the

results obtained from LPS-induced macrophages, the LPS-triggered expression of *Il6*, *Ccl2* and

Ccl8 in lung and colon tissues was decreased in *LyzM*-Cre$^+$ *Wtap*$^{\Delta 1\text{-}77}$ mice compared with *Wtap*$^{\text{fl/fl}}$

mice (Fig 3F-H), whereas the expression of *Il1a*, *Il1b* and *Tnfa* showed no significant differences

(Fig EV6A and B). The same trend was obtained with RNA-seq data from colon samples (Fig 3I).

Notably, *LyzM*-Cre$^+$ *Wtap*$^{\Delta 1\text{-}77}$ mice showed less lung inflammation and fewer pathological

characteristics of lung injury (Fig 3J) and were therefore more tolerant to LPS-induced fatal sepsis

(Fig 3K) than their WT counterparts. Moreover, the serum concentration of IL-6 was significantly

reduced in *LyzM*-Cre$^+$ *Wtap*$^{\Delta 1\text{-}77}$ mice compared with their *Wtap*$^{\text{fl/fl}}$ littermates (Fig 3L). Similarly,

after intraperitoneal injection of *Pseudomonas aeruginosa* (*P. aeruginosa*) or *Listeria*

monocytogenes (*L. monocytogenes*), the induced expression of *Il6*, *Ccl2* and *Ccl8* in lung (Figs

3M and EV6C and D) and colon (Fig EV6E-G) tissues, as well as the secretion of IL-6 into serum

(Fig 3N), were decreased in *LyzM*-Cre$^+$ *Wtap*$^{\Delta 1\text{-}77}$ mice. Collectively, these data suggested that the

increased expression of WTAP aggravates LPS- or bacteria-induced inflammatory responses *in*

vivo.

WTAP promotes activation of the STAT3 signalling axis through the m^{6}A modification to

accelerate inflammatory responses

As a key m^{6}A "writer", WTAP is required for anchoring METTL3 and other cofactors to nuclear

speckles to modulate m^{6}A modification of RNA (Ping *et al*, 2014). To determine whether WTAP

affects the inflammatory responses through m^{6}A modification, we first quantified m^{6}A abundance

in cells by LC–MS/MS assays. The results showed that the total m^{6}A level was observably

increased in WTAP-overexpressing 293T cells and reduced in *WTAP* KO 293T cells (Fig EV7A

and B). In particular, the overall m^{6}A modification level increased in WT cells but not WTAP-

deficient cells after LPS stimulation (Fig EV7A and B). These data suggested that upregulation of

WTAP can significantly improve the overall m^{6}A modification level during inflammatory stimuli,

which is consistent with the previous conclusion that WTAP is responsible for the observed

increase in m^{6}A modification upon bacterial infection (Wu *et al.*, 2020).

We then performed methylated RNA immunoprecipitation sequencing (MeRIP-Seq) analyses to

map mRNA transcripts with different m^{6}A peaks accompanied by different mRNA levels in WT

and *WTAP$^{Δ1-77}$* THP-1 cells or *Wtap$^{Δ1-77}$* BMDMs after LPS stimulation. According to the MeRIP-

Seq data, the consensus m^{6}A core motifs were enriched in the m^{6}A peaks in all the samples (Fig

EV7C). A peak distribution analysis showed that the m^{6}A sites were enriched in both exons and

3'UTRs (Fig EV7D), with the highest enrichment near the stop codon (Fig EV7E and F).

Moreover, the MeRIP-Seq analysis revealed that the m^{6}A peaks of nearly 5,000 or more than

1,500 transcripts were lower in *WTAP$^{Δ1-77}$* THP-1 cells or in *Wtap$^{Δ1-77}$* BMDMs compared with

those in WT cells upon LPS treatment, respectively (Fig 4C). These identified transcripts were

enriched in many inflammatory signalling pathways, including the NF-κB (TNF-

α/TLR/NLR/CLR-mediated), JAK-STAT and MAPK pathways (Fig EV7G and H). We further

compared the transcripts with reduced m^{6}A marks after WTAP deletion between THP-1 cells and

BMDMs treated with LPS and identified 989 overlapping transcripts (Fig 4C). KEGG enrichment

analyses revealed that these overlapping transcripts were closely related to bacterial infection and

inflammatory signalling pathways (Fig 4D), such as *IL6ST, IL15RA, IL18R1, TYK2, JAK2,*

RIPK2, *IL1A*, *IL12B* and *CXCL11* (Fig 4C).

Both RNA-seq and MeRIP-seq indicated that WTAP is closely related to inflammatory pathways,

such as the NF-κB, JAK-STAT and MAPK pathways. We then explored the effect of WTAP on the

activation of these pathways. As shown in Fig 4E and F, the phosphorylation level of STAT3, but

not that of p65 or p38, was significantly decreased in *WTAP*^{Δ1-77} THP-1 cells in response to LPS or

HKST treatment. To further confirm this effect, the same treatments were performed with

BMDMs from *Wtap*^{fl/fl} or *LyzM*-Cre⁺ *Wtap*^{Δ1-77} mice, and similar results were obtained (Fig 4G

and H). Consistently, i.p. injection of LPS upregulated the phosphorylation level of STAT3 (Fig 4I

and J). However, such induction was significantly reduced in the lung (Fig 4I) and colon (Fig 4J)

tissues from *LyzM*-Cre⁺ *Wtap*^{Δ1-77} mice. We further found that the LPS-induced mRNA expression

of *IL6*, *CCL2*, *CCL8* and *CXCL8* (Fig 4K), but not that of *TNFA*, *IL1B* and *LTA* (Fig EV7I), was

markedly reduced after pretreatment with SC144 (Xu *et al*, 2013), a gp130 (IL6ST) inhibitor that

abrogates the IL-6/STAT3 signalling axis. Interestingly, although the expression of many

proinflammatory cytokines was reduced in *WTAP*^{Δ1-77} THP-1 cells, the expression and secretion of

IL-1β, TNF-α or LTA (TNF-β) showed no significant change compared with those of WT THP-1

cells in response to the indicated stimuli (Figs 4L and EV7J-L), which is consistent with the fact

that p65 activation is not affected by WTAP. Together, these data suggested that WTAP-mediated

m⁶A modification is deeply involved in the regulation of bacterial infection and inflammatory

response in both human and mice and may accelerate inflammatory responses by enhancing the

STAT3 signalling axis.

WTAP promotes the protein output of proinflammatory genes through m⁶A modification

Consistent with our experimental results, many m⁶A-modified genes that were found to be

regulated by WTAP based on the sequencing data are involved in the activation and regulation of

the STAT3 signalling axis, such as IL6ST, TYK2, JAK2, IL15RA and IL18R1 (Figs 4C and

EV8A-I). Since the transcription and secretion of IL-6 is significantly reduced in our analyses and

its contribution to the activation of STAT3 may be predominant in acute inflammatory response,

we then further measured the protein abundance of the identified key components involved in the

IL-6/STAT3 signalling axis, including IL6ST, IL6R, TYK2, JAK1, JAK2 and STAT3 (Hirano,

2021). The results showed that the protein abundance of IL6ST but not the other tested

components was reduced in $WTAP^{A1-77}$ THP-1 cells and $Wtap^{A1-77}$ BMDMs compared with control

cells (Figs 5A and EV9A). Moreover, the upregulation of IL6ST induced by LPS was inhibited in

the $WTAP^{A1-77}$ THP-1 cells (Fig 5B). Consistently, the expression of IL6ST was significantly

reduced on the cell surface of $Wtap^{A1-77}$ macrophages (Fig 5C). Notably, the analysis of mouse

models of LPS-induced sepsis showed that the increased protein expression of IL6ST and WTAP

(Fig EV9B and C) was reduced in the lung and colon tissues (Fig 5D) from $LyzM$-Cre^{+} $Wtap^{A1-77}$

mice. We next generated $IL6ST^{-/-}$ THP-1 cells using the CRISPR/Cas9 approach, and as shown in

$WTAP^{A1-77}$ THP-1 cells, the phosphorylated STAT3 level was significantly decreased in $IL6ST^{-/-}$

THP-1 cells after stimulation with LPS (Fig 5E). The expression of IL-6 was also reduced in

$IL6ST^{-/-}$ THP-1 cells in response to different inflammatory stimuli (Fig 5F).

MeRIP-Seq revealed that the m^{6}A peaks obtained with high confidence were distributed in regions

near the stop codon in $IL6ST$ transcripts (Fig EV8A). We then designed two gene-specific primer

pairs, F1/R1 and F2/R2 (Appendix Table S2), to measure the change in m^{6}A abundance at sites 1

and 2 of $IL6ST$ transcripts via MeRIP-qPCR assay (Fig 5G). The results showed that the m^{6}A

marks enriched at both site 1 and site 2 in the $IL6ST$ transcripts were markedly decreased in

$WTAP^{A1-77}$ THP-1 cells compared with control cells (Fig 5H). For verification, we mutated the

putative m^{6}A-modified adenosine by replacing it with a thymine in the $IL6ST$ mRNA and inserted

the wild-type (WT) or mutated (Mut) UTRs into a reporter gene plasmid (psiCHECKTM-2) (Fig

5I). Luciferase reporter assay data showed that the luciferase activities of the Mut reporter were

significantly weaker than those of the WT reporter (Fig 5J). Moreover, the luciferase activities of

the WT reporter were weakened in $WTAP$ KO 293T cells (Fig 5K). Ectopic expression of Flag-

tagged WTAP in $WTAP$ KO 293T cells the enhanced luciferase activities of the WT reporter (Fig

EV9D). All these data indicated that the detected m^{6}A mark sites in $IL6ST$ transcripts are direct

substrates of WTAP and crucial for maintaining the output of the IL6ST protein.

Because m^{6}A modification can affect many aspects of gene expression, including nuclear export,

splicing, 3'-end processing, decay and translation (Zaccara et al, 2019), we evaluated the mRNA

expression of $IL6ST$ and found that deleting WTAP did not affect the $IL6ST$ mRNA abundance or

stability (Fig EV9E and F). Due to the decay of $IL6ST$ mRNA may not be affected by WTAP, we

next measured the translation efficiency of IL6ST through polysome profiling (Fig EV9G). We

calculated the proportion of mRNAs in polysome fractions via qRT–PCR and found that the

distribution of *IL6ST* mRNA shifted to the lighter fraction in *WTAP[A1-77]* THP-1 cells after

stimulation with LPS for 4 h (Fig 5L). However, this difference was not observed in *GAPDH*

mRNA (Fig EV9H). Taken together, WTAP can facilitate the expression of IL6ST by enhancing

the translation efficiency via m^{6}A modification. Functional interpretation of m^{6}A modification is

realized by RNA-binding proteins called "readers", mainly YTHDF1/2/3 and YTHDC1/2, which

can influence the degradation or translation of m^{6}A-modified RNA (Shi *et al*, 2017; Zhao *et al*,

2020). Because YTHDF1 and YTHDF3 are the major m^{6}A readers that promote the translation of

m^{6}A-modified mRNA, we sought to determine whether YTHDF1 or YTHDF3 targets cellular

IL6ST transcripts to regulate their translation. Immunoblot analyses showed that the deletion of

YTHDF1 or YTHDF3 in THP-1 cells using a CRISPR-mediated genome editing approach

reduced the protein expression of IL6ST (Fig EV9I). To verify the direct binding of YTHDF1 and

YTHDF3 to m^{6}A-methylated *IL6ST* mRNA, we synthesized biotin-labelled RNA probes based on

the distribution of the m^{6}A peak on *IL6ST* transcripts (Fig EV9J) and performed an RNA pull-

down assay followed by immunoblotting of the isolated proteins. The results revealed that

methylated *IL6ST* transcripts strongly interacted with YTHDF1 and YTHDF3 (Fig EV9K).

Furthermore, RNA immunoprecipitation (RIP) analyses with antibodies against YTHDF1 or

YTHDF3 followed by qRT–PCR revealed that the amount of *IL6ST* mRNA bound to YTHDF1 or

YTHDF3 was significantly increased after stimulation with LPS but reduced after WTAP

depletion (Fig EV9L). These results suggested that the m^{6}A marks on *IL6ST* mRNA mediated by

WTAP obviously promoted the binding of YTHDF1 and YTHDF3 to *IL6ST* mRNA. Polysome

profiling assays showed that deletion of YTHDF1 or YTHDF3 in THP-1 cells efficiently

decreased the protein expression of IL6ST by hindering its translation (Fig EV9M and N).

Overall, these results demonstrated that the increased efficiency of IL6ST translation induced by

the m^{6}A modification is mediated by YTHDF1 and YTHDF3.

In addition to IL-6/IL6ST/STAT3 signalling, the IL-15/IL15R and IL18/IL18R signalling which

dictate T cell response, regulate B cell homing, and activate NK cells (Wawrocki *et al*, 2016), may

also mediate the regulation of inflammatory responses by activating STAT3 (Mishra *et al*, 2014;

Yasuda *et al*, 2019). In our data, the m^{6}A peaks distributed in the 3'-UTR regions of *IL15RA* and

IL18R1, the crucial receptor molecules for IL-15/IL15R and IL18/IL18R signaling, also almost

disappeared when WTAP was depleted (Fig EV8B and C). The mRNA and protein abundance of

IL15Rα and IL18Rα was significantly reduced in $WTAP^{Δ1-77}$ THP-1 cells and BMDMs compared with the control cells (Figs 5M and N and EV9O), which suggested that WTAP-mediated m^6A modification can regulate the expression of IL15Rα and IL18Rα proteins. Although WTAP only affects the mRNA decay of *IL18R1* but not *IL15RA* (Figs 5O and EV9P), further reporter assays performed by replacing the putative m^6A-modified adenosine with a thymine in the 3' UTR of *IL15RA* and *IL18R1* mRNA confirmed that the protein abundances of *IL15RA* and *IL18R1* were promoted by WTAP-mediated m^6A (Fig 5P-R). Furthermore, the m^6A methylation of many proinflammatory factors, such as *RIPK2*, *IL1A*, *IL12B* and *CXCL11*, was significantly reduced in $WTAP^{Δ1-77}$ THP-1 cells (Fig EV8F-I), and reporter assays confirmed that WTAP promoted the protein output of these proinflammatory genes via m^6A modification (Fig EV9Q-S). Hence, our work in macrophages not only revealed the significance of WTAP for finetuning the activation of IL6/STAT3 signalling axis, but also implied that the protein synthesis of many proinflammatory genes can be regulated by WTAP via m^6A modification in immune homeostasis and disease occurrence, which is unique to WTAP compared to other m^6A regulators.

The phase separation of WTAP promotes METTL3 recruitment to efficiently modify inflammatory transcripts

By verifying the MeRIP-seq data, we found that the genes affected by WTAP in THP-1 cells did not completely overlap before and after LPS treatment (Fig EV10A). Transcripts with low m^6A abundance in $WTAP^{Δ1-77}$ THP-1 cells were involved in very few metabolic pathways in the resting state (Fig EV10B) but were mainly enriched in cytokine production and inflammatory signalling pathways after LPS treatment (Fig EV10C). Additionally, metagene profiles of the m^6A peak distribution showed that inflammatory stimulation increased the abundance of m^6A marks but decreased the number of m^6A peaks (Fig EV7E and F). These observations implied that the expression of WTAP activated by p65 may lead to more concentrated deposition of m^6A marks on inflammatory genes.

Recently, liquid-liquid phase separation (LLPS) has emerged as a widespread mechanism through which cells dynamically recruit and organize key signalling molecules. LLPS has also been revealed to play an important role in the formation of the m^6A "writer" complex and the regulation of the fate of m^6A-modified mRNAs (Chong & Forman-Kay, 2016; Han *et al*, 2022; Ries *et al*,

2019; Zhang *et al*, 2021a). An analysis of the disordered regions in the WTAP protein using the IUPred2A (https://iupred2a.elte.hu/) and PLAAC (http://plaac.wi.mit.edu/) tools revealed a low-complexity region of 16.5 kDa in the C-terminus of the WTAP coding sequence (CDS) (Fig 6A and B), which may form liquid droplets as a result of phase separation (Nott *et al*, 2015). We then further explored whether WTAP has an ability to undergo LLPS upon inflammatory stress and how it plays its roles in the progress of m^{6}A modification of those inflammatory transcripts. To this end, recombinantly expressed green fluorescent protein (GFP)-WTAP fusion proteins were bacterially expressed and purified (Fig EV11A). The GFP-WTAP formed spherical droplets with an aspect ratio close to 1, and an increase in the protein concentration and salt concentration increased the abundance of GFP droplets from barely detectable small foci to regular and large droplets (Figs 6C and EV11B). Through time-lapse microscopy, we found that the droplet size of GFP-WTAP increased over time (Fig EV11C), and captured the fusion between droplets within 40 s (Fig EV11D), implying the high dynamics and fluidity of the GFP-WTAP droplets. Conversely, droplet formation was substantially inhibited by 10% 1,6-hexanediol (Hex), a compound that putatively dissolves liquid-liquid phase-separated condensates (Fig EV11E).

To further test the ability of WTAP to undergo LLPS in intact cells, we constructed a C-terminal enhanced GFP (EGFP)-tagged WTAP vector and ectopically expressed it in 293T cells (Fig EV11F). We found that GFP-tagged WTAP formed discrete puncta in the nucleus (Fig 6D). In line with those *in vitro* results, fluorescence recovery after photobleaching (FRAP) assays in 293T cells showed that the fluorescence of GFP-tagged WTAP was efficiently and gradually recovered after bleaching (Fig 6E and F), indicating the potential phase separation capability of WTAP *in vivo*. Notably, endogenous WTAP condensates formed in HeLa cells, and the number of these droplets increased significantly after cell stimulation with TNF-α due to upregulated expression of WTAP (Figs 6G and H and EV11G), suggesting that upregulated WTAP induced by inflammatory stimuli is more prone to phase separation. In consistent with other study, the WTAP droplets colocalized with the nuclear speckle marker SC35 (Ping *et al.*, 2014) (Fig 6G and I). However, no difference in the number of SC35 droplets was found before and after inflammatory stimulation (Fig 6J), suggesting that endogenous WTAP droplets formed by self-occurring LLPS rather than the dynamics of the nuclear speckles, and the LLPS of WTAP may facilitate its colocalization with nuclear speckles.

To further identify the regions in WTAP that are needed for phase separation, we used the PONDR

tool (http://www.pondr.com) to predict 5 unfolded intrinsically disordered regions (IDRs) with

scores greater than 0.7 in the WTAP protein sequence (Fig EV11H). A series of truncated WTAP

mutants labelled with GFP were then constructed (Fig EV11H) and expressed in 293T cells (Fig

EV11I). Through confocal imaging, we found that WTAP failed to form droplets only when amino

acid residues 225-287 were deleted (Figs 6K and EV11J), suggesting that this sequence is

essential for the LLPS of WTAP. Biomolecular condensates formed through LLPS can maintain

locally elevated concentrations of resident proteins and/or RNAs (Banani et al, 2017). Hence, we

subsequently explored the biological function of the phase separation of WTAP. Previous studies

revealed that WTAP recruits METTL3 and other cofactors to nuclear speckles and that the RNA-

binding capacity of METTL3 is profoundly reduced in the absence of WTAP (Ping et $al.$, 2014).

Here, we found that mCherry-METTL3 (Fig EV11K) readily fused with phase-separated GFP-

WTAP and colocalized to all GFP-WTAP droplets (Fig EV11L), demonstrating that fusion with

WTAP enhanced the METTL3 LLPS in $vitro$. In $vivo$, METTL3 also formed liquid droplets with

the help of WTAP (Fig 6L-N), but deleting the LLPS sequence on WTAP significantly reduced the

ability of WTAP to recruit METTL3 through LLPS (Fig 6L) without affecting the direct binding

of METTL3 and WTAP (Fig EV11M), suggesting that WTAP LLPS might prime METTL3

condensation in cells. To confirm that the LLPS of WTAP promotes METTL3 to methylate

inflammatory mRNAs, RIP with antibodies against METTL3 followed by qRT–PCR was

performed and the results revealed that the amount of $IL6ST$, $IL15RA$ and $IL18R1$ mRNAs bound

by METTL3 was significantly increased after stimulation with LPS and was reduced by WTAP

depletion (Figs 6O and EV11N and O). These data further indicated that the LLPS of WTAP

facilitates the aggregation of METTL3 and the methylation of inflammatory mRNAs.

Consistently, the absence of phase transition capability observably inhibited WTAP-dependent

m^{6}A modification (Figs 6P and EV11P). Taken together, these data suggested that upregulated

WTAP undergoes phase separation to facilitate the assembly of the writer complex and the

localization of the writer complex to nuclear speckles. Since the nuclear speckles are active

regions for gene transcription and RNA processing and are rich in transcriptionally active

proinflammatory transcripts during inflammatory stress, WTAP-medicated LLPS may improve the

accessibility between the m^{6}A writer complex and inflammatory transcripts in nuclear speckles,

promoting the deposition of m^{6}A onto transcriptionally active inflammatory transcripts and the

activation of proinflammatory responses (Fig 6Q).

WTAP deficiency alleviates the disease progression of DSS-induced IBD in mice

All the aforementioned data suggest that WTAP is a true proinflammatory risk factor. To further

investigate the potential role and clinical relevance of upregulated WTAP in the progression of

inflammatory diseases, we first revealed that both WTAP and IL6ST were upregulated in SLE,

asthma, sepsis, RA, psoriasis and IBD patients (Figs EV1A-E and EV12A). Moreover, high

expression of WTAP was significantly positively correlated with the level of inflammation in

patients. For example, the upregulation of WTAP was significantly inhibited in psoriasis patients

treated with anti-IL17A therapy and Crohn's disease (CD) patients treated with anti-TNF therapy

(GSE137218/16879, Appendix Table S1), correspondingly, proinflammatory cytokine expression

was also significantly decreased in these patients after treatment (Fig 7A and B). We further

evaluated the expression of WTAP in the progression of IBD. An analysis of a single-cell dataset

of IBD patients (GSE125527, Fig 7C and Appendix Table S1) showed that the expression of

WTAP rather than METTL3 was significantly increased in some immune cell subsets in IBD

patients, particularly in monocytes (Fig 7D). The prevalence of WTAP upregulation in IBD patient

samples was then verified using more public datasets (GSE119600/10616, Fig EV12B and C and

Appendix Table S1). Moreover, using data from the IBDMDB database, an analysis of the

correlation between genes involved in m^{6}A methylation and genes that have been identified as

IBD risk genes or biomarkers showed that WTAP exhibited the highest correlation with the

expression of IBD genes among all the m^{6}A regulators (Fig EV12D). Observably, increased

expression of WTAP was accompanied by an increase in the global m^{6}A abundance in various

inflammatory diseases (Fig EV12E and F), which is consistent with our results in LPS-induced

macrophages. Consistent with this finding, similar tendencies were observed in mouse models of

imiquimod (IMQ)-induced psoriasis (Figs 7E and EV12G) and 3% dextran sodium sulfate (DSS)-

induced colitis (Figs 7F and EV12H). Thus, the high expression of WTAP accompanied by a high

level of m^{6}A modification and high expression levels of proinflammatory cytokines should be a

common characteristic of distinct inflammatory diseases, such as IBD.

To verify the biological correlation between WTAP and IBD signatures, we subjected *LyzM*-Cre$^+$

Wtap$^{Δ1-77}$ mice and their *Wtap$^{fl/fl}$* control littermates to DSS to induce acute colitis. The

administration of DSS in drinking water can cause the death of intestinal epithelial cells and thus

compromise gut barrier function and cause inflammation (Wirtz *et al*, 2017). We found that

compared with their *Wtap$^{fl/fl}$* control littermates, the *LyzM*-Cre$^+$ *Wtap$^{Δ1-77}$* mice treated with DSS

displayed attenuated colitis, accompanied by less weight loss (Fig 7G) and a lower disease activity

index (Fig 7H). Moreover, WTAP deficiency substantially attenuated the shortening of the colon

length in the DSS-challenged mice (Fig 7I and J). The low expression of *Il6*, *Il1a*, *Ccl2* and *Ccl8*

in DSS-challenged *LyzM*-Cre$^+$ *Wtap$^{Δ1-77}$* mice indicated attenuated colonic inflammation (Fig 7K).

A histopathological assessment revealed that the colonic mucosa of *LyzM*-Cre$^+$ *Wtap$^{Δ1-77}$* mice was

relatively intact, with less inflammatory cell infiltration after DSS treatment (Fig 7L).

Furthermore, *LyzM*-Cre$^+$ *Wtap$^{Δ1-77}$* mice exhibited later death onset and a lower death rate (Fig

7M). Taken together, these results suggested that WTAP deficiency attenuates the severity of DSS-

induced IBD, revealing a clinically relevant correlation between the WTAP levels and IBD

progression. Because the increased expression of WTAP is ubiquitous in many inflammatory

diseases, we hypothesize that WTAP is a novel and important risk factor that exerts a broad

spectrum of effects on inflammatory diseases and may serve as a novel potential therapeutic target

in the treatment of inflammatory diseases.

Discussion

WTAP is a smart adapter in the assembly of the m⁶A writer complex

Inflammation is vital for protecting the host against invading pathogens and for repairing tissue damage, which requires tight and concise control of pro- and anti-inflammatory gene expression. m^{6}A, the most prevalent internal mRNA modification, has emerged as an abundant modification throughout the transcriptome and has been recently linked to various inflammatory states, including autoimmunity, infection, metabolic diseases, and cancers. Although the roles played by m^{6}A in virus replication, host defense, immune cell activation and disease occurrence have been extensively investigated, basic knowledge of the relationship between the m^{6}A modification and inflammatory diseases has been largely unattained. Here, we found that WTAP, as a p65-controlled gene, is universally upregulated after distinct inflammatory stimuli. Increased WTAP protein is related to higher levels of m^{6}A modification and excessive inflammatory responses. Noteworthy, the transcription and expression of WTAP have been shown to be regulated in various biological processes. For example, at the transcriptional and posttranscriptional levels, HIF1α (Shao *et al*, 2023) and STAT3 (Liu *et al*, 2023) can transactivate WTAP gene expression by directly binding to its promoter region, and WTAP can also be upregulated by the epigenetic alteration of H3K4me3, resulting in the senescence of nucleus pulposus cells (NPCs) (Li *et al*, 2022a). At the translational and posttranslational levels, a recent study revealed that through epigenetic modifications, the pseudogene WTAPP1 can bind to WTAP mRNA to promote its translation by recruiting the EIF3 translation initiation complex, accelerating the progression of pancreatic ductal adenocarcinoma (PDAC) (Deng *et al*, 2021). And our previous study showed that WTAP undergoes degradation via the ubiquitination-proteasome pathway in virus-infected cells, leading to reductions in the abundance of m^{6}A marks and the subsequent protein output of *IRF3* and *IFNAR1* and providing the negative feedback to attenuate the intensity of IFN-I signalling (Ge *et al*, 2021). Using smart adapters without methyltransferase activity, such as WTAP, may provide more precise adjustment of m^{6}A "writer" complex activity, which is evolutionarily and physiologically beneficial to the dynamic regulation of phenotypic diversity and homeostasis in higher organisms, may be realized.

In addition to showing that WTAP is a p65-controlled gene, the basic mechanism by which WTAP coordinates the assembly dynamics of the writer complex to promote the m^{6}A modification of

many inflammatory genes was further revealed in this study. Although WTAP has been shown to

be critical for the localization of the m^{6}A "writer" complex to nuclear speckles, the dynamic

assembly of the "writer" complex is not fully unexplored. A previous study reported that METTL3

can also undergo LLPS (Han *et al.*, 2022), and METTL3 interacts with m^{6}A-METTL-associated

complex (MACOM) mainly through WTAP (Su *et al*, 2022). Here, we further found that when

WTAP is absent or cannot undergo LLPS, METTL3 is dispersedly distributed in the nucleus,

which indicates that the LLPS of WTAP is essential for the assembly of the writer complex,

providing dynamic m^{6}A regulation in distinct biological or physiological conditions. In this study,

we found that highly expressed WTAP spontaneously forms condensates through LLPS under

inflammatory stress, which may be important for the recruitment of more "writer" proteins to

nuclear speckles. Because nuclear speckles are sites with many transcriptionally active genes

(Faber *et al*, 2022), WTAP-mediated LLPS may increase the local "writer" concentrations and

improve the accessibility between the m^{6}A "writer" complex and inflammatory transcripts in

nuclear speckles, thereby promoting the m^{6}A modification of inflammatory genes and accelerating

of inflammatory response. In the past decade, biomolecular condensates formed by LLPS have

been widely reported to modulate many cellular functions by compartmentalizing specific proteins

and nucleic acids in subcellular environments with distinct properties (Shin & Brangwynne, 2017;

Vendruscolo & Fuxreiter, 2022). Our finding further expands the regulatory roles of phase

separation to the dynamic assembly of writer complex and the methylation of specific transcripts,

describing how cells utilize the composition and compartmentation of multivalent condensates to

affect the m^{6}A epitranscriptome.

577 The finetuning of the STAT3 signalling axis by targeting of WTAP LLPS may be an effective

578 therapeutic strategy for excessive inflammation

m^{6}A modification has been reported to regulate inflammatory responses and related diseases. For

example, KDM6B (Wu *et al.*, 2020), SOCS1 (Du *et al*, 2020), TRAF3 and TRAF6 (Zheng *et al*,

2017; Zong *et al.*, 2019) are dynamically regulated by m^{6}A modification in controlling

inflammatory responses. Through analyses of clinical data and mouse disease models, we clearly

found that increased expression of WTAP is associated with inflammatory diseases, such as sepsis,

SLE, asthma (Mo *et al*, 2022), RA, psoriasis (Mande *et al*, 2018) and IBD (Nie *et al*, 2022) (Fig

EV1). A recent study also suggested the relationship of WTAP to the anti-inflammatory effect of

Astragalus mongholicus polysaccharide (APS) (Long *et al*, 2022). In addition to showing the

relationship between increased WTAP and higher m^{6}A modification in inflammation, we

specifically showed that an increase in WTAP protein was accompanied by increases in the m^{6}A

abundances on *IL6ST*, *IL15RA*, *IL18R1*, and many other transcriptionally active proinflammatory

cytokines. High IL6ST, IL15RA and IL18R1 protein abundance promotes activation of the STAT3

signalling axis, which in turn contributes to the production of proinflammatory cytokines, such as

IL-6, CCL2 and CCL8 (Fig 6Q). Consistently, two recent studies individually noted that the

knockdown of FTO can significantly increase the overall m^{6}A abundance, STAT3 phosphorylation

and the secretion of proinflammatory cytokines and thereby positively regulates the inflammation

and IFN-I response (McFadden *et al*, 2022; Tang *et al.*, 2022). Moreover, the IL-6/STAT3

pathway is aberrantly hyperactivated in many cancers, and such hyperactivation is generally

associated with a poor clinical prognosis (Johnson *et al*, 2018). Recent studies have also shown

that high levels of m^{6}A modification enhance JAK1 translation in high-metastatic hepatocellular

carcinoma (HCC) cells, thereby promoting the activation of STAT3 (Liu *et al.*, 2023).

Interestingly, the expression of WTAP is also significantly elevated in many cancers, such as

pancreatic carcinoma (Deng *et al.*, 2021), osteosarcoma (Ren *et al*, 2022), HCC (Chen *et al*, 2019)

and nasopharyngeal carcinoma (Li *et al*, 2022b). This implies that the strategy of finetuning the

STAT3 signalling axis by WTAP may also play an important role in the progression of multiple

cancers. Because the STAT3 axis has been well characterized as a mechanism to accelerate the

production of a variety of cytokines and chemokines (Tanaka *et al*, 2014) and is highly associated

with inflammatory diseases, including cytokine storm syndromes, autoimmune diseases and

cancers (Gharibi *et al*, 2020; Matsuyama *et al*, 2020) , the novel NF-κB/WTAP/STAT3 axis

identified in this study indicated that WTAP is an ideal therapeutic target in the treatment of many

inflammatory diseases and cancers.

To date, effective therapeutic interventions for excessive inflammation by inhibiting STAT3

activity in the clinic remain to be further developed, largely due to the challenge of directly

targeting STAT3 (STAT3 is a transcription factor that lacks enzymatic activity), and in part due to

the complexity and diversity of the biological functions of STAT3 (Yu *et al*, 2014). Small-

molecule inhibitors targeting the NF-κB and STAT3 signalling pathways cause side effects because the long-term suppression of the NF-κB or STAT3 affects the homeostasis of the immune system as well as normal cell growth, differentiation, and survival (Siegel *et al.*, 2011; Zhang *et al.*, 2017). Here, we found that hyperinflammatory states in the body are often accompanied by high levels of m^6A methylation. Reducing the abundance of m^6A marks by STM2457 (Fig EV13A), a METTL3 inhibitor, significantly inhibited HKST-induced expression of proinflammatory genes (Fig EV13B). We also verified this effect in a mouse model of LPS-induced sepsis by showing that the m^6A modification levels of colon and lung tissues from mice were significantly reduced by STM2457 (Fig EV13C and D). A decrease in the m^6A modification level blocked the expression of inflammatory cytokines and activation of inflammatory signals induced by LPS (Fig EV13E and F). Similarly, STM2457 effectively alleviated lung inflammation and the pathological characteristics of lung injury (Fig EV13G). Based on these data combined with one study showing that STM2457 can effectively reduce the severity of DSS-induced IBD in mice (Yang *et al*, 2022), we believe that reversing the high abundance of m^6A modifications in disease states has broad prospects in the treatment of inflammatory diseases in the future. Because WTAP is more concentrated in the inflammatory context to affect the m^6A modification of nascent inflammatory genes, epigenetic therapies targeting WTAP to reprogram the m^6A landscape in cells may achieve more effective and safer therapeutic outcomes.

Elvitegravir (EVG), which was originally developed to treat HIV infection (Sato *et al*, 2006), was recently found to efficiently inhibit YAP oncogenic transcription by specifically disrupting SRC-1 LLPS (Zhu *et al*, 2021), demonstrating the feasibility of targeting LLPS to treat diseases. For the purpose of reducing hyperinflammation, one possible strategy is the development or identification of small-molecule drugs that can block WTAP LLPS to reduce the overall m^6A abundance, which might reverse the hyperinflammatory state of diseases. Moreover, interfering with WTAP LLPS during inflammatory state may reduce the damage to constitutively expressed genes and normal cells to successfully combat autoimmunity, infection, metabolic diseases, and cancers. In addition, given the critical role played by the NF-κB signalling pathway in cancer and various chronic diseases, NF-κB has become a potential target for pharmacological intervention, and numerous inhibitors of NF-κB may play important roles in the treatment of various inflammatory diseases in the future (Freitas & Fraga, 2018). Here, we found that the NF-κB p65 inhibitor PG490

significantly reduced the overall m^{6}A abundance by inhibiting the expression of WTAP (Fig

EV13H). Thus, for the development of novel small anti-inflammatory molecules, researchers may

need to consider the effects of these inhibitors on epitranscriptome.

In summary, we provide the first demonstration that high m^{6}A modification in a variety of

hyperinflammatory states is p65-dependent, due to the key component of the writer complex

WTAP is transcriptionally regulated by p65 and its overexpression can lead to higher m^{6}A

modification. We also discovered that upregulated WTAP undergoes phase separation, which

facilitates the aggregation of the writer complex and its localization to nuclear speckles and the

deposition of m^{6}A onto transcriptionally active transcripts, which results in the promotion of

proinflammatory responses. Based on the mouse model of inflammatory diseases generated in this

study and many online clinical data, we hypothesize that higher expression of WTAP should be a

risk factor in many hyperinflammatory states and inflammatory diseases. Interrupting the

assembly of the m^{6}A writer complex by targeting the phase separation of WTAP to reduce the

global m^{6}A level may be a potential therapeutic strategy for preventing excessive inflammation.

Material and Methods

Mice.

C57BL/6 *LyzM*-Cre⁺ *Wtap*ᶠˡ/ᶠˡ mice were generated by Gempharmatech Co., Ltd. using the CRISPR/Cas9 approach. All mouse lines were maintained at Sun Yat-sen University under specific pathogen-free (SPF) conditions in ventilated microisolator cages. The Institutional Animal Care and Use Committee (IACUC) of Sun Yat-sen University (Guangzhou, China) approved all the experimental protocols concerning the handling of mice. The study is compliant with all relevant ethical regulations regarding animals.

Cells.

The 293T, HeLa and THP-1 cell lines were purchased from ATCC. All these cells were cultured in endotoxin-free DMEM or RPMI-1640, supplemented with 10% FBS and 1% penicillin–streptomycin. PBMCs and BMDMs were isolated and cultured as described previously(Ge *et al.*, 2021).

In vitro recombinant protein expression and purification.

The expressing plasmids (pET-32a) encoding WTAP conjugated eGFP and METTL3 conjugated mCherry were transformed into BL21 *E.coli*. *E.coli* were cultured in LB with Ampicillin (50 μg/mL) at 37 °C for about 12 hr till OD600 = 0.6. After induction with 1 mM IPTG at 37 °C for 8 hr, the cultured *E.coli* was harvested by centrifugation at 4000 rpm, 4 °C, 10 min and resuspended in lysis buffer. Cells were lysed by sonication on ice and centrifuged (12,000 rpm, 30 min, 4 °C) to remove debris and collected the supernatant. The supernatant was purified by incubation with Ni-NTA agarose beads (QIAGEN) overnight at 4 °C. Then, Ni-NTA beads were washed with wash buffer (50 mM NaH2PO4, 300 mM NaCl, 20 mM imidazole, pH 7.8), and proteins were eluted with elution buffer (50 mM NaH2PO4, 300 mM NaCl, 300 mM imidazole, pH 7.8). The purified proteins were further dialyzed by using PD10 column (GE Healthcare), and concentrated using Amicon Ultra 30 K (Millipore) concentrators at 4 °C. Concentrated proteins were quantified by the BCA method (Thermo Fisher) and stored at -80 °C.

Generation of knockout THP-1 cells.

Knockout (KO) cells were constructed using the CRISPR/Cas9 system. Small guide RNAs (gRNAs) targeting the genome sequence of target genes were designed using an online gRNA design tool (by Zhang Feng laboratory, http://crispr.mit.edu/), and subcloned into the lentiCRISPR v2 vector. This vector was transfected into the HEK293T cells along with the following two lentiviral packing plasmids: psPAX2 and pVSV-G. The culture supernatants were collected at 48 and 60 hours after transfection and concentrated by ultracentrifugation before use for infection. The infection-positive (GFP$^+$) cells were selected by flow cytometry, monoclonal cells were confirmed by genomic sequencing and immunoblotting analysis with the corresponding antibody. The gRNA sequences used for generating the KO cells are listed in Table S2.

***In vivo* LPS challenge.**

For endotoxicity studies, age-matched female *Wtap*$^{fl/fl}$ and *LyzM-cre*$^+$ *Wtap*$^{fl/fl}$ mice (8 weeks old) were intraperitoneally injected with LPS (40 mg kg^{-1}). Mouse survival was monitored every 4 hr.

Female *Wtap*$^{fl/fl}$ and *LyzM-cre*$^+$ *Wtap*$^{A1-77}$ mice (8 weeks old) were intraperitoneally injected with LPS (10 mg kg^{-1}) or isodose PBS. After 12 hr, the mice were killed, blood was collected, and serum levels of IL-6 were measured by ELISA. Lung and colon tissues were collected, and RNA samples were extracted for qRT-PCR to determine the mRNA levels of inflammatory cytokines.

Female *Wtap*$^{fl/fl}$ and *LyzM-cre*$^+$ *Wtap*$^{A1-77}$ mice (8 weeks old) were intraperitoneally injected with LPS (10 mg kg^{-1}) or PBS. After 6 hr, the mice were killed, and lungs from control or LPS-stimulated mice were dissected, fixed in 4% paraformaldehyde (PFA) embedded in paraffin, sectioned, stained with hematoxylin and eosin (H&E) solution, and examined by light microscopy to detect histological changes.

Female *Wtap*$^{fl/fl}$ and *LyzM-cre*$^+$ *Wtap*$^{A1-77}$ mice (8 weeks old) were intraperitoneally injected with *P. aeruginosa* (ATCC27853, 2×10^8 CFU) or *L. monocytogenes* (ATCC19116, 1×10^8 CFU) or PBS. After 16 hr, the mice were killed, and the related phenotypes were measured as described above.

ELISA assays and quantitative RT–PCR (qRT–PCR).

The concentrations of IL-6, IL-1β and TNF-α in culture supernatants were measured using kits

from R&D Systems or Proteintech, according to the manufacturer's instructions.

Total RNA was extracted using TRIzol reagent (Invitrogen) and reversed-transcribed with a

PrimeScript[TM] RT reagent kit with gDNA Eraser (TaKaRa) according to the manufacturer's

instructions. And 2×Polarsignal® qPCR mix (MIKX) was used for quantitative real-time PCR

analysis. The data were normalized by the level of *GAPDH* or *ACTB* expression in each individual

sample, and primer sequences used are listed in Table S2.

Luciferase reporter gene assay.

For a promoter reporter gene assay, 293T cells were plated in 48-well plates and transiently

transfected with WTAP promoter reporter (pGL3 basic plasmid), with increasing doses of Flag-

tagged NF-κB p65, IRF3 or C/EBPβ and 7.5 ng of the Renilla luciferase reporter vector using

Lipofectamine 2000. 24 hrs post-transfection, luciferase activities were measured with a dual-

luciferase reporter assay system (Promega) according to the manufacturer's instructions. Reporter

gene activity was determined by normalization of Firefly luciferase activity to Renilla luciferase

activity.

For other reporter gene assays, wild-type or *WTAP* KO 293T cells were plated in 48-well plates

and transiently transfected with the indicated reporter vectors (psiCHECK[TM]-2) expressing wild-

type or mutated UTRs of *IL6ST* using Lipofectamine 2000. 24 hrs post-transfection, the luciferase

activities were measured as above. Reporter gene activity was determined by normalization of the

Renilla luciferase activity to Firefly luciferase activity.

Small interfering RNAs (siRNAs) and RNA interference.

The control nontargeting and WTAP-, YTHDF1-, YTHDF3-specific siRNAs were designed and

synthesized by RiboBio. THP-1 cells (8×10^5 cells/mL) were transiently transfected with siRNA

(final concentration of 20 μM) using Lipofectamine RNAiMAX following the manufacturer's

instructions. The sequences of siRNAs are listed in Table S2.

RNA decay assay.

WTAP^{Δ1-77} and WT THP-1 cells were seeded at a density of 8×10^5 cells/mL in 12-well plates.

After treatment with PMA (50 ng/mL, Sigma), the cells were treated with transcription inhibitor

Act D (5 μg/mL, Sigma) to block *de novo* RNA synthesis, and collected at different times. RNA

samples were extracted for qRT–PCR to determine the mRNA levels of the indicated genes.

Laser scanning confocal microscopy (LSCM).

Cells for fluorescence experiments were cultured in glass bottom culture dishes (Nest Scientific).

After treated with indicated stimulation, cells were fixed by 4% paraformaldehyde for 10 min,

followed with 3 times wash with PBS. Then, cells were permeabilized with methyl alcohol for

30 min at −20 °C and rinsed with PBS for 3 times. After blocking in 5% goat serum for 1 h at

room temperature, cells were incubated with primary antibodies at 4 °C, overnight. 1 × PBST

(PBS with 0.1% Tween20) was used to wash the cells for 3 times and subsequently incubated with

fluorescently labeled secondary antibodies at room temperature for 1 h. Confocal images were

obtained by Leica TCS SP8 STED 3X confocal microscope. The number of condensates were

counted and analyzed by ImageJ software.

Live-cell imaging.

At 24 hrs posttransfection incubation, cells were loaded into temperature- and CO2-controlled

live-cell imaging chamber of Leica TCS SP8 STED 3X confocal microscope. Cells were imaged

typically by use of 2 laser wavelengths (488 nm for eGFP imaging and 560 nm for mCherry

imaging).

In vitro phase separation assay.

The purified recombinant proteins were mixed at indicated concentration with LLPS buffer and

5% PEG8000 at 37 °C. The mixture was pipetted onto glass bottom dish and imaged by Leica

TCS SP8 STED 3X confocal microscope equipped with 100 × 1.40 NA oil objectives.

Fluorescence recovery after photobleaching (FRAP) assay.

FRAP assay was conducted by Leica TCS SP8 STED 3X confocal microscopy. 488- or 568-nm

laser beam was used to bleach the fluorescent protein at a region of interest (ROI), followed with

collecting time-lapse images. Fluorescence intensity of indicated ROI was measured and

normalized to the fluorescence intensity of pre-bleaching image by Leica AS Lite.

m⁶A dot blots.

Total RNA was extracted with TRIzol reagent. Equal amounts of RNA (300 ng) were denatured at

95 °C for 3 min. The samples were immediately chilled on ice, added to a positively charged nylon

membrane (PALL), and then cross-linked with a UV crosslinker. After blocking and incubating

with the m⁶A antibody (CST) overnight, the membrane was incubated with the secondary antibody

at room temperature for 1 hr. Signals were detected using a chemiluminescence imaging system.

Methylene blue in 0.3 M sodium acetate (pH 5.2) was used to indicate the amount of total RNA.

RNA immunoprecipitation assay.

RIP was conducted using a RIP assay Kit (MBL) following the manufacturer's instructions. In

brief, Protein A/G magnetic beads coated with 5 µg of specific antibody or normal IgG were

incubated with cell lysates at 4 °C overnight. Proteins were then extracted for immunoblot

analysis, and the co-precipitated RNAs were isolated by elution buffer and purified by TRIzol

reagent, and subsequently subjected to qRT–PCR analysis.

DNA/RNA pull-down assay.

Biotin-labeled DNA and RNA probes were synthesized by RiboBio. For DNA pull-down assay,

Flag-tagged NF-κB p65 expression constructs were transfected into 293T cells. 24 hrs post-

transfection, whole cell lysates were extracted from 293T cells using IP lysis buffer. Biotin-

coupled DNA-protein complex was pulled down by incubating whole cell lysates with high-

capacity streptavidin agarose beads (Thermo Fisher) according to the manufacturer's instructions.

The bound proteins were eluted and used for immunoblot analysis.

For RNA pull-down assay, biotin-coupled RNA complex was pulled down by incubating cell

lysates with high-capacity streptavidin agarose beads, bound proteins were then extracted for

immunoblot analysis.

m⁶A RNA-IP-qRT–PCR (MeRIP–qPCR).

An MeRIP assay was performed following a previously described procedure with minor

modifications (Ge *et al.*, 2021). Briefly, 200 μg of total RNA was fragmentated to approximately

150-200 nt in length, purified by magnetic beads, and then incubated with anti-m^{6}A antibody- or

IgG-conjugated protein A/G magnetic beads in 1 × IP buffer at 4 °C overnight.

Immunoprecipitated methylated RNAs were competitive eluted by free m^{6}A, and recovered with a

RNeasy kit (QIAGEN). One-tenth of the fragmented RNA was saved for use as the input control

and further analyzed by qRT–PCR with the MeRIP RNAs using primers for the targeted gene. The

related enrichment of m^{6}A in each sample was calculated by normalizing the number of

amplification cycles (Cq) of the m^{6}A-IP portion to the Cq of the corresponding input portion.

Quantification of the m^{6}A modification by LC-MS/MS.

200 ng extracted mRNA was digested into nucleosides by Nuclease P1 (1 U, NEB, M0660S) and

shrimp alkaline phosphatase (rSAP, 1 U, NEB, M0371S) in 50 μL RNase-free water at 37 °C

overnight. The mixture was diluted to 100 μL, 10 μL of which was injected into an LC–MS/MS

system consisting of a high-performance liquid chromatographer (Shimadzu) equipped with a

C18-T column (Weltech) and a Triple Quad 4500 (AB SCIEX) mass spectrometer in positive ion

mode by multiple-reaction monitoring. Mass transitions of m/z 268.0–136.0 (A), m/z 245.0–113.1

(U), m/z 244.0–112.1 (C), m/z 284.0–152.0 (G) and m/z 282.0–150.1 (m^{6}A) were monitored. A

concentration series of pure commercial nucleosides (MCE) was employed to generate standard

curves. The concentration of nucleosides in samples were obtained by fitting signal intensity to a

standard curve with certain ratios calculated subsequently.

RNA-seq and data analysis.

Whole-cell total RNA was isolated using TRIzol reagent and quantified using a NanoDrop 2000

spectrophotometer (Thermo). The cDNA library was constructed by Biomarker Technologies.

Sequencing was performed on an Illumina HiSeq 2500 platform. High-quality reads were mapped

to the human reference genome (hg19) using HISAT2. DESeq, an R package, was applied for

differential gene expression analysis. We filtered the differentially expressed genes based on a

false discovery rate (FDR) <0.05.

Methylated RNA immunoprecipitation sequencing (MeRIP-seq) and data analysis.

Total RNA was extracted by TRIzol reagent and mRNA was isolated using a Dynabeads mRNA

purification kit (Thermo Fisher). Then, cellar mRNA was fragmented using a fragmentation kit

(Thermo Fisher), and subsequent m^6A immunoprecipitation, MeRIP-seq and data analysis was

carried out as previously described with minor modifications (Zhang *et al*, 2021b).

Statistical analysis.

The data are represented as the mean ± s.d. unless otherwise indicated, and two-tailed Student's *t*-

test or Mann–Whitney U test with a confidence interval of 95% were performed for all statistical

analyses (survival curves were compared by the log-rank test) with GraphPad Prism 6 software.

Differences between two groups were considered statistically significant when the *P* value was

less than 0.05.

Data and code availability

Publicly available datasets, which were analyzed in this study including GSE19315, GSE198326,

GSE2411, GSE2638, GSE69063, GSE13887, GSE137268, GSE97779, GSE166388, GSE208303,

GSE125527, GSE16879, GSE193193, GSE179874, GSE119600, GSE10616, GSE137218,

GSE227851 and GSE189847 are available at the GEO database

(https://www.ncbi.nlm.nih.gov/geo/).

The validated IBDMDB data and relevant participants' information are available at the IBDMDB

database (http://ibdmdb.org).

All data generated for this paper have been deposited at the SRA under access number:

PRJNA943438.

Acknowledgments

This work was supported by projects from National Natural Science Foundation of China

(91942301, 31970852, 32200712), Guangdong Science and Technology Department

(2020B1212060031, 2023A1515010541), Ministry of Science and Technology of the People's

Republic of China (2018YFD0900502), "Decoding TCM" collaborative research project of

BUCM (90010060920009), Innovation Group Project of Southern Marine Science and

Engineering Guangdong Laboratory (Zhuhai) (311021006), National Key Research and

Development Project (2019YFC1710104) and Fundamental Research Funds for the Central

Universities (23yxqntd001).

Author Contributions

870 Y.G., S.Y. and A.X. designed the study, analyzed the data, and prepared the manuscript with input

871 from the other authors. Y.G., R.C., T.L., Y.C., J.H., Y.L., X.X. and G.X. performed the experiments

872 collaboratively. B.L., H.C. and G.L. performed the data analysis of MeRIP-seq. S.Y. and A.X.

873 conceived the study, supervised experiments, and wrote the paper. A.X. and S.Y. led the project

874 and finally approved the manuscript.

Declaration of Interests

877 The authors declare that they have no competing interests.

References

References

Banani SF, Lee HO, Hyman AA, Rosen MK (2017) Biomolecular condensates: organizers of cellular

biochemistry. *Nat Rev Mol Cell Biol* 18: 285-298

Banerjee S, Biehl A, Gadina M, Hasni S, Schwartz DM (2017) JAK-STAT signaling as a target for

inflammatory and autoimmune diseases: current and future prospects. *Drugs* 77: 521-546

Barbieri I, Kouzarides T (2020) Role of RNA modifications in cancer. *Nat Rev Cancer* 20: 303-322

Bradley JR (2008) TNF-mediated inflammatory disease. *J Pathol* 214: 149-160

Cao XT (2016) Self-regulation and cross-regulation of pattern-recognition receptor signalling in health

and disease. *Nat Rev Immunol* 16: 35-50

Carpenter S, Ricci EP, Mercier BC, Moore MJ, Fitzgerald KA (2014) Post-transcriptional regulation of

gene expression in innate immunity. *Nat Rev Immunol* 14: 361-376

Chen YH, Peng CH, Chen JR, Chen DY, Yang B, He B, Hu WD, Zhang YP, Liu H, Dai LF *et al* (2019)

WTAP facilitates progression of hepatocellular carcinoma via m^{6}A-HuR-dependent epigenetic

silencing of ETS1. *Mol Cancer* 18: 127

Chong PA, Forman-Kay JD (2016) Liquid-liquid phase separation in cellular signaling systems. *Curr*

Opin Struct Biol 41: 180-186

Cui L, Ma R, Cai JLY, Guo CY, Chen ZY, Yao LL, Wang YY, Fan R, Wang X, Shi YL (2022) RNA

modifications: importance in immune cell biology and related diseases. *Signal Transduct Target Ther*

7: 334

Deng JG, Zhang JL, Ye Y, Liu KJ, Zeng LX, Huang JY, Pan L, Li M, Bai RH, Zhuang LS *et al* (2021)

N^6-methyladenosine-mediated upregulation of WTAPP1 promotes WTAP translation and Wnt signaling

to facilitate pancreatic cancer progression. *Cancer Res* 81: 5268-5283

Du J, Liao W, Liu WC, Deb DK, He L, Hsu PJ, Nguyen T, Zhang LD, Bissonnette M, He C *et al*

(2020) N^6-adenosine methylation of Socs1 mRNA is required to sustain the negative feedback control

of macrophage activation. *Dev Cell* 55: 737-753.e737

Faber GP, Nadav-Eliyahu S, Shav-Tal Y (2022) Nuclear speckles - a driving force in gene expression. *J*

Cell Sci 135: jcs259594

Freitas RHCN, Fraga CAM (2018) NF-kappa B-IKK beta pathway as a target for drug development:

realities, challenges and perspectives. *Curr Drug Targets* 19: 1933-1942

Fu Y, Dominissini D, Rechavi G, He C (2014) Gene expression regulation mediated through reversible

m^{6}A RNA methylation. *Nat Rev Genet* 15: 293-306

Ge Y, Ling T, Wang Y, Jia X, Xie XM, Chen R, Chen SW, Yuan SC, Xu AL (2021) Degradation of

WTAP blocks antiviral responses by reducing the m^{6}A levels of IRF3 and IFNAR1 mRNA. *EMBO*

Rep 22: e52101

Gharibi T, Babaloo Z, Hosseini A, Abdollahpour-alitappeh M, Hashemi V, Marofi F, Nejati K,

Baradaran B (2020) Targeting STAT3 in cancer and autoimmune diseases. *Eur J Pharmacol* 878:

173107

Gong T, Liu L, Jiang W, Zhou RB (2020) DAMP-sensing receptors in sterile inflammation and

inflammatory diseases. *Nat Rev Immunol* 20: 95-112

Greten FR, Grivennikov SI (2019) Inflammation and cancer: triggers, mechanisms, and consequences.

Immunity 51: 27-41

Han D, Longhini AP, Zhang X, Hoang V, Wilson MZ, Kosik KS (2022) Dynamic assembly of the

mRNA m^{6}A methyltransferase complex is regulated by METTL3 phase separation. *PLoS biol* 20:

e3001535

Hirano T (2021) IL-6 in inflammation, autoimmunity and cancer. *Int Immunol* 33: 127-148

Huang Y, Su R, Sheng Y, Dong L, Dong Z, Xu HJ, Ni TF, Zhang ZS, Zhang T, Li CY *et al* (2019)
Small-molecule targeting of oncogenic FTO demethylase in acute myeloid leukemia. *Cancer Cell* 35:
677-691

Hunter CA, Jones SA (2015) IL-6 as a keystone cytokine in health and disease. *Nat Immunol* 16: 448-
457

Jiang XL, Liu BY, Nie Z, Duan LC, Xiong QX, Jin ZX, Yang CP, Chen YB (2021) The role of m^{6}A
modification in the biological functions and diseases. *Signal Transduct Target Ther* 6: 74

Johnson DE, O'Keefe RA, Grandis JR (2018) Targeting the IL-6/JAK/STAT3 signalling axis in cancer.
Nat Rev Clin Oncol 15: 234-248

Kang SJ, Tanaka T, Narazaki M, Kishimoto T (2019) Targeting interleukin-6 signaling in clinic.
Immunity 50: 1007-1023

Li GC, Ma L, He SJ, Luo RJ, Wang BJ, Zhang WF, Song Y, Liao ZW, Ke WC, Xiang Q *et al* (2022a)
WTAP-mediated m^{6}A modification of lncRNA NORAD promotes intervertebral disc degeneration. *Nat*
Commun 13: 1469

Li N, Rana TM (2022) Regulation of antiviral innate immunity by chemical modification of viral RNA.
Wiley Interdiscip Rev RNA 13: e1720

Li ZX, Zheng ZQ, Yang PY, Lin L, Zhou GQ, Lv JW, Zhang LL, Chen FP, Li YQ, Wu CF *et al* (2022b)
WTAP-mediated m^{6}A modification of lncRNA DIAPH1-AS1 enhances its stability to facilitate
nasopharyngeal carcinoma growth and metastasis. *Cell Death Differ* 29: 1137-1151

Liu BW, Cao JL, Wu BT, Hao KX, Wang XY, Chen X, Shen ZF (2023) METTL3 and STAT3 form a
positive feedback loop to promote cell metastasis in hepatocellular carcinoma. *Cell Commun Signal*
21: 121

Liu J, Qian C, Cao XT (2016) Post-translational modification control of innate immunity. *Immunity*
45: 15-30

Long HJ, Lin HY, Zheng P, Hou LJ, Zhang M, Lin SY, Yin K, Zhao GJ (2022) WTAP mediates the
anti-inflammatory effect of Astragalus mongholicus polysaccharide on THP-1 macrophages. *Front*
Pharmacol 13: 1023878

Luo JH, Xu T, Sun K (2021) N^6-methyladenosine RNA modification in inflammation: roles,
mechanisms, and applications. *Front Cell Dev Biol* 9: 670711

Mande P, Zirak B, Ko WC, Taravati K, Bride KL, Brodeur TY, Deng A, Dresser K, Jiang ZZ, Ettinger
R *et al* (2018) Fas ligand promotes an inducible TLR-dependent model of cutaneous lupus-like
inflammation. *J Clin Invest* 128: 2966-2978

Mapperley C, van de Lagemaat LN, Lawson H, Tavosanis A, Paris J, Campos J, Wotherspoon D,
Durko J, Sarapuu A, Choe J *et al* (2021) The mRNA m^{6}A reader YTHDF2 suppresses proinflammatory
pathways and sustains hematopoietic stem cell function. *J Exp Med* 218: e20200829

Matsuyama T, Kubli SP, Yoshinaga SK, Pfeffer K, Mak TW (2020) An aberrant STAT pathway is
central to COVID-19. *Cell Death Differ* 27: 3209-3225

McFadden MJT, Sacco MTA, Murphy KA, Park MS, Gokhale NSY, Somfleth KYM, Horner SM
(2022) FTO suppresses STAT3 activation and modulates proinflammatory interferon-stimulated gene
expression. *J Mol Biol* 434: 167247

Medzhitov R, Horng T (2009) Transcriptional control of the inflammatory response. *Nat Rev Immunol*
9: 692-703

Mishra A, Sullivan L, Caligiuri MA (2014) Molecular pathways: interleukin-15 signaling in health and

in cancer. *Clin Cancer Res* 20: 2044-2050

Mo BW, Li XM, Li SM, Xiao B, Yang J, Li HM (2022) m^{6}A echoes with DNA methylation: Coordinated DNA methylation and gene expression data analysis identified critical m^{6}A genes associated with asthma. *Gene* 828: 146457

Nie K, Yi J, Yang YY, Deng MZ, Yang Y, Wang TY, Chen XJ, Zhang ZY, Wang XY (2022) A broad m^{6}A modification landscape in inflammatory bowel disease. *Front Cell Dev Biol* 9: 782636

Nott TJ, Petsalaki E, Farber P, Jervis D, Fussner E, Plochowietz A, Craggs TD, Bazett-Jones DP, Pawson T, Forman-Kay JD *et al* (2015) Phase transition of a disordered nuage protein generates environmentally responsive membraneless organelles. *Mol Cell* 57: 936-947

O'Connell MA, Mannion NM, Keegan LP (2015) The epitranscriptome and innate immunity. *PLoS Genet* 11: e1005687

Ping XL, Sun BF, Wang L, Xiao W, Yang X, Wang WJ, Adhikari S, Shi Y, Lv Y, Chen YS *et al* (2014) Mammalian WTAP is a regulatory subunit of the RNA N^6-methyladenosine methyltransferase. *Cell Res* 24: 177-189

Ren ZP, Hu YC, Sun J, Kang YX, Li GS, Zhao HJ (2022) N^6-methyladenosine methyltransferase WTAP-stabilized FOXD2-AS1 promotes the osteosarcoma progression through m^{6}A/FOXM1 axis. *Bioengineered* 13: 7963-7973

Ries RJ, Zaccara S, Klein P, Olarerin-George A, Namkoong S, Pickering BF, Patil DP, Kwak H, Lee JH, Jaffrey SR (2019) m^{6}A enhances the phase separation potential of mRNA. *Nature* 571: 424-428

Sato M, Motomura T, Aramaki H, Matsuda T, Yamashita M, Ito Y, Kawakami H, Matsuzaki Y, Watanabe W, Yamataka K *et al* (2006) Novel HIV-1 integrase inhibitors derived from quinolone antibiotics. *J Med Chem* 49: 1506-1508

Scholler E, Weichmann F, Treiber T, Ringle S, Treiber N, Flatley A, Feederle R, Bruckmann A, Meister G (2018) Interactions, localization, and phosphorylation of the m^{6}A generating METTL3-METTL14-WTAP complex. *RNA* 24: 499-512

Shao YL, Li YQ, Li MY, Wang LL, Zhou HS, Liu DH, Yu L, Lin J, Gao XN (2023) HIF1 alpha-mediated transactivation of WTAP promotes AML cell proliferation via m^{6}A-dependent stabilization of KDM4B mRNA. *Leukemia* 37: 1254-1267

Shi HL, Wang X, Lu ZK, Zhao BXS, Ma HH, Hsu PJ, Liu C, He C (2017) YTHDF3 facilitates translation and decay of N^6-methyladenosine-modified RNA. *Cell Res* 27: 315-328

Shin Y, Brangwynne CP (2017) Liquid phase condensation in cell physiology and disease. *Science* 357: eaaf4382

Shulman Z, Stern-Ginossar N (2020) The RNA modification N^6-methyladenosine as a novel regulator of the immune system. *Nat Immunol* 21: 501-512

Siegel AM, Heimall J, Freeman AF, Hsu AP, Brittain E, Brenchley JM, Douek DC, Fahle GH, Cohen JI, Holland SM *et al* (2011) A critical role for STAT3 transcription factor signaling in the development and maintenance of human t cell memory. *Immunity* 35: 806-818

Stone JH, Tuckwell K, Dimonaco S, Klearman M, Aringer M, Blockmans D, Brouwer E, Cid MC, Dasgupta B, Rech J *et al* (2017) Trial of tocilizumab in giant-cell arteritis. *N Engl J Med* 377: 317-328

Su SC, Li SS, Deng T, Gao MS, Yin Y, Wu BX, Peng C, Liu JZ, Ma JB, Zhang KM (2022) Cryo-EM structures of human m^{6}A writer complexes. *Cell Res* 32: 982-994

Tanaka T, Narazaki M, Kishimoto T (2014) IL-6 in Inflammation, Immunity, and Disease. *Cold Spring Harb Perspect Biol* 6: a016295

Tang SY, Meng JY, Tan J, Liu XY, Zhou HX, Li N, Hou SP (2022) N^6-methyladenosine demethylase

FTO regulates inflammatory cytokine secretion and tight junctions in retinal pigment epithelium cells.
Clin Immunol 241: 109080

Triantafilou M, Triantafilou K (2002) Lipopolysaccharide recognition: CD14, TLRs and the LPS-
activation cluster. *Trends Immunol* 23: 301-304

Vendruscolo M, Fuxreiter M (2022) Protein condensation diseases: therapeutic opportunities. *Nat*
Commun 13: 5550

Wang JH, Yan SS, Lu HY, Wang SF, Xu DH (2019) METTL3 attenuates LPS-induced inflammatory
response in macrophages via NF-kappa B signaling pathway. *Mediators Inflamm* 2019: 3120391

Wawrocki S, Druszczynska M, Kowalewicz-Kulbat M, Rudnicka W (2016) Interleukin 18 (IL-18) as a
target for immune intervention. *Acta Biochim Pol* 63: 59-63

Wirtz S, Popp V, Kindermann M, Gerlach K, Weigmann B, Fichtner-Feigl S, Neurath MF (2017)
Chemically induced mouse models of acute and chronic intestinal inflammation. *Nat Protoc* 12: 1295-
1309

Wu CL, Chen WX, He JC, Jin SH, Liu YK, Yi Y, Gao ZX, Yang JY, Yang JH, Cui J *et al* (2020)
Interplay of m^{6}A and H3K27 trimethylation restrains inflammation during bacterial infection. *Sci Adv*
6: aba0647

Xu HJ, Lin CJ, Yang JH, Chen X, Chen YY, Chen JG, Guo AZ, Hu CM (2022) The role of N^6-
methyladenosine in inflammatory diseases. *Oxid Med Cell Longev* 2022: 9744771

Xu SL, Grande F, Garofalo A, Neamati N (2013) Discovery of a novel orally active small-molecule
gp130 inhibitor for the treatment of ovarian cancer. *Mol Cancer Ther* 12: 937-949

Yang LC, Wu GT, Wu Q, Peng LX, Yuan LW (2022) METTL3 overexpression aggravates LPS-induced
cellular inflammation in mouse intestinal epithelial cells and DSS-induced IBD in mice. *Cell Death*
Discov 8: 62

Yankova E, Blackaby W, Albertella M, Rak J, De Braekeleer E, Tsagkogeorga G, Pilka ES, Aspris D,
Leggate D, Hendrick AG *et al* (2021) Small-molecule inhibition of METTL3 as a strategy against
myeloid leukaemia. *Nature* 593: 597-601

Yasuda K, Nakanishi K, Tsutsui H (2019) Interleukin-18 in health and disease. *Int J Mol Sci* 20: 649

Yeung YT, Aziz F, Guerrero-Castilla A, Arguelles S (2018) Signaling pathways in inflammation and
anti-inflammatory therapies. *Curr Pharm Des* 24: 1449-1484

Yu H, Lee H, Herrmann A, Buettner R, Jove R (2014) Revisiting STAT3 signalling in cancer: new and
unexpected biological functions. *Nat Rev Cancer* 14: 736-746

Zaccara S, Ries RJ, Jaffrey SR (2019) Reading, writing and erasing mRNA methylation. *Nat Rev Mol*
Cell Biol 20: 608-624

Zhang JZ, Mehta S, Zhang J (2021a) Liquid-liquid phase separation: a principal organizer of the cell's
biochemical activity architecture. *Trends Pharmacol Sci* 42: 845-856

Zhang Q, Lenardo MJ, Baltimore D (2017) 30 years of NF-kappa B: A blossoming of relevance to
human pathobiology. *Cell* 168: 37-57

Zhang S, Liu F, Wu ZS, Xie JF, Yang Y, Qiu HB (2020) Contribution of m^{6}A subtype classification on
heterogeneity of sepsis. *Ann Transl Med* 8: 306

Zhang Z, Chen T, Chen HX, Xie YY, Chen LQ, Zhao YL, Liu BD, Jin LM, Zhang WT, Liu C *et al*
(2021b) Systematic calibration of epitranscriptomic maps using a synthetic modification-free RNA
library. *Nat Methods* 18: 1213-1222

Zhao YC, Shi YF, Shen HF, Xie WZ (2020) m^{6}A-binding proteins: the emerging crucial performers in
epigenetics. *J Hematol Oncol* 13: 35

Zheng QL, Hou J, Zhou Y, Li ZY, Cao XT (2017) The RNA helicase DDX46 inhibits innate immunity
by entrapping m^{6}A-demethylated antiviral transcripts in the nucleus. *Nat Immunol* 18: 1361-1361
Zhu GY, Xie JJ, Fu ZZ, Wang ML, Zhang QC, He H, Chen ZJ, Guo X, Zhu JD (2021) Pharmacological
inhibition of SRC-1 phase separation suppresses YAP oncogenic transcription activity. *Cell Res* 31:
1028-1031
Zhu SM, Lu Y (2020) Dexmedetomidine suppressed the biological behavior of HK-2 cells treated with
LPS by down-regulating ALKBH5. *Inflammation* 43: 2256-2263
Zong X, Zhao J, Wang H, Lu ZQ, Wang FQ, Du HH, Wang YZ (2019) Mettl3 deficiency sustains long-
chain fatty acid absorption through suppressing Traf6-dependent inflammation response. *J Immunol*
202: 567-578
1065

Figure legends

Figure 1. Multiple inflammatory stimuli markedly increased the expression of WTAP in macrophages.

A Heatmap showing a change in the mRNA abundance of m^{6}A-related genes in THP-1 cells stimulated with lipopolysaccharide (LPS) at the indicated time points.

B-D qRT-PCR showing the mRNA abundance of the METTL3/METTL14/WTAP heterotrimer in THP-1 cells (**B**), peripheral blood mononuclear cells (PBMCs) (**C**) and bone marrow-derived macrophages (BMDMs) (**D**) stimulated with LPS at the indicated time points.

E-G Immunoblots showing the protein abundance of the METTL3/METTL14/WTAP heterotrimer in THP-1 cells (**E**), PBMCs (**F**) and BMDMs (**G**) stimulated with LPS at the indicated time points.

H qRT-PCR showing the mRNA abundance of WTAP in THP-1 cells stimulated with CL097, Pam3CSK4, heat-killed *Salmonella typhimurium* (HKST) or heat-killed *Listeria monocytogenes* (HKLM) at the indicated time points.

I-K Immunoblots showing the protein abundance of the METTL3/METTL14/WTAP heterotrimer in THP-1 cells stimulated with Pam3CSK4 (**I**), HKST (**J**) or HKLM (**K**) at the indicated time points.

L qRT-PCR showing the mRNA abundance of WTAP in BMDMs stimulated with CL097, Pam3CSK4, HKST or HKLM at the indicated time points.

M Immunoblots showing the protein abundance of the METTL3/METTL14/WTAP heterotrimer in BMDMs stimulated with Pam3CSK4 at the indicated time points.

Data information: Data are presented as the mean ± s.d. in **B-D**, **H** and **L**, with individual measurements overlaid as dots, statistical analysis was performed using a two-tailed Student's *t*-test. Data are representative of three independent biological experiments in **E-G**, **I-K** and **M**.

Figure 2. WTAP is an NF-κB p65-stimulated gene.

A Schematic of the full-length, mutant and truncated human *WTAP* promoter.

B Schematic of the mouse *Wtap* promoter.

C, D Luciferase activity analyses in 293T cells transfected with a luciferase reporter for the *WTAP* promoter with WT (**C**), mutant or truncated (**D**) p65-binding motif, together with increasing doses of Flag-tagged NF-κB p65.

E-H qRT-PCR or immunoblots showing the expression of WTAP in THP-1 cells (**E**, **F**) or PBMCs (**G**, **H**) treated with PG490, followed by stimulation with LPS at the indicated time points.

I-L qRT-PCR or immunoblots showing the expression of WTAP in WT and *RELA*$^{-/-}$ THP-1 cells stimulated with LPS (**I**, **J**) or Pam3CSK4 (**K**, **L**) at different time points. m Schematic showing the biotin-labeled *WTAP* promoter probes. n DNA pull-down assays showing the effect of NF-κB p65 binding with *WTAP* promoter probes.

Data information: Data are presented as the mean ± s.d. in **C-E**, **G**, **L** and **K**, with individual measurements overlaid as dots, statistical analysis was performed using a two-tailed Student's *t*-test. Data are representative of three independent biological experiments in **F**, **H**, **J**, **L** and **N**.

Figure 3. WTAP positively regulates the expression of proinflammatory cytokines.

A KEGG analyses of down-regulated genes in *WTAP*$^{Δ1-77}$ THP-1 cells during LPS-induced inflammatory stress compared with WT cells.

B Heatmap showing a change in the mRNA abundance of pro-inflammatory cytokines in WT and *WTAP*$^{Δ1-77}$ THP-1 cells stimulated with LPS at the indicated time points.

C qRT–PCR showing the mRNA abundance of *IL6*, *CCL2*, *CCL8* and *CXCL8* in WT and *WTAP*$^{Δ1-77}$ THP-1 cells stimulated with LPS at the indicated time points.

D ELISAs were performed to detect IL-6 secretion in supernatants of WT and *WTAP*$^{Δ1-77}$ THP-1 cells stimulated with LPS, Pam3CSK4, HKST or HKLM at 3 and 6 hr.

E Heatmap showing a change of mRNA abundance of pro-inflammatory cytokines in WT and *Wtap*$^{Δ1-77}$ BMDMs stimulated with LPS at the indicated time points.

F-H qRT–PCR showing the mRNA abundance of *Il6* (**F**), *Ccl2* (**G**) and *Ccl8* (**H**) in the lung or colon tissues from *Wtap*$^{fl/fl}$ and *LyzM-cre*$^{+}$ *Wtap*$^{Δ1-77}$ mice that were intraperitoneally injected with LPS (10 mg kg^{-1}) for 12 hr. n = 10 mice per group.

I Heatmap showing a change of mRNA abundance of pro-inflammatory cytokines in the colon tissues from *Wtap*$^{fl/fl}$ and *LyzM-cre*$^{+}$ *Wtap*$^{Δ1-77}$ mice that were intraperitoneally injected with LPS for 12 hr.

J Hematoxylin & eosin staining (H&E) assays showing the lung injury of the *Wtap*$^{fl/fl}$ and *LyzM-cre*$^{+}$ *Wtap*$^{Δ1-77}$ mice that were intraperitoneally injected with LPS (10 mg kg^{-1}) for 6 hr. Scale bars, 50 μm. n = 4 mice per group.

K The survival of *Wtap*^fl/fl and *LyzM-cre*^+ *Wtap*^Δ1-77 mice was monitored for 32 hr following the

intraperitoneal injection of LPS (40 mg kg^{-1}). n = 10 mice per group.

L ELISAs were performed to detect IL-6 secretion in serum from *Wtap*^fl/fl and *LyzM-cre*^+ *Wtap*^Δ1-

^77 mice that were intraperitoneally injected with LPS (10 mg kg^{-1}) for 12 hr. n = 8 mice per group.

M qRT–PCR showing the mRNA abundance of *Il6* in the lung tissues from *Wtap*^fl/fl and *LyzM-cre*^+

Wtap^Δ1-77 mice that were intraperitoneally injected with *P. aeruginosa* or *L. monocytogenes* for 16

hr. n = 8 mice per group.

N ELISAs were performed to detect IL-6 secretion in serum from *Wtap*^fl/fl and *LyzM-cre*^+ *Wtap*^Δ1-

^77 mice as in that were intraperitoneally injected with *P. aeruginosa* or *L. monocytogenes* for 16 hr.

n = 3 mice per group.

Data information: Data are presented as the mean ± s.d. in **C**, **D**, **F-H**, and **L-N**, with individual

measurements overlaid as dots, statistical analysis was performed using a two-tailed Student's *t*-

test. Data are representative of three independent biological experiments in **J**.

1139

Figure 4. WTAP promotes the activation of STAT3 signaling through the m^{6}A modification

to accelerate inflammatory responses.

A, B LC–MS/MS quantification of m^{6}A abundance in mRNA extracted from WT and *WTAP*^Δ1-77

THP-1 cells (**A**) or *Wtap*^Δ1-77 BMDMs (**B**) with or without LPS treatment.

C Venn diagrams showing transcripts with decreased m^{6}A abundance in *WTAP*^Δ1-77 THP-1 cells

and *Wtap*^Δ1-77 BMDMs compared with WT cells stimulated with LPS for 4 hr.

D KEGG enrichment analysis of the overlapping transcripts presented in (**C**).

E-H Immunoblots showing total and phosphorylated (p-) STAT3, p65 and p38 levels in WT and

WTAP^Δ1-77 THP-1 cells (**E, F**) or BMDMs (**G, H**) stimulated with LPS or HKST at the indicated

time points.

I, J Immunoblots showing the total and phosphorylated (p-) STAT3 level in the lung (**I**) or colon

(**J**) tissues from *Wtap*^fl/fl and *LyzM-cre*^+ *Wtap*^Δ1-77 mice that had been intraperitoneally injected

with LPS (10 mg kg^{-1}) for 6 hr.

K qRT–PCR showing the expression of *IL6*, *CCL2*, *CCL8* and *CXCL8* in THP-1 cells treated with

SC144, followed by stimulation with LPS at the indicated time points.

L ELISAs were performed to measure IL-1β and TNF-α secretion in supernatants of WT and

WTAP$^{Δ1-77}$ THP-1 cells that were stimulated with LPS at 3 and 6 hr.

Data information: Data are presented as the mean ± s.d. in **A**, **B**, **K** and **L**, with individual

measurements overlaid as dots, statistical analysis was performed using a two-tailed Student's *t*-

test. Data are representative of three independent biological experiments in **E-J**.

1160

Figure 5. WTAP promotes the protein output of proinflammatory genes through m⁶A

modification.

A Immunoblots showing the expression of critical adapters in the IL-6/ STAT3 signaling pathway

in WT and *WTAP*$^{Δ1-77}$ THP-1 cells.

B Immunoblots showing the expression of IL6ST in WT and *WTAP*$^{Δ1-77}$ THP-1 cells stimulated

with LPS at the indicated time points.

C Representative flow cytometry data showing IL6ST-FITC fluorescence in cell surface of

peritoneal macrophages and BMDMs.

D Immunoblots showing the expression of WTAP and IL6ST in the lung or colon tissues from

Wtap$^{fl/fl}$ and *LyzM-cre*$^+$ *Wtap*$^{Δ1-77}$ mice that had been intraperitoneally injected with LPS (10 mg

kg^{-1}) for 12 hr.

E Immunoblots showing the total and phosphorylated (p-) STAT3, p65 and p38 levels in WT and

IL6ST$^{-/-}$ THP-1 cells stimulated with LPS at the indicated time points.

F ELISAs were performed to measure IL-6 secretion in the supernatants of WT and *IL6ST*$^{-/-}$ THP-

1 cells stimulated with LPS, Pam3CSK4, HKST or HKLM for 6 hr.

G Schematic representation showing the position of m⁶A motifs within the IL6ST transcripts.

F1/R1 represents detection site 1, which includes four "DRACH" elements, and F2/R2 represents

detection site 2, which includes three "DRACH" elements.

H The abundance of *IL6ST* transcripts (with detection sites 1 and 2) among mRNAs

immunoprecipitated with an anti-m⁶A antibody in WT and *WTAP*$^{Δ1-77}$ THP-1 cells stimulated with

LPS for 6 hr.

I WT or mutant (Mut) m⁶A consensus sequences (A-to-T mutation) on the *IL6ST*-3' UTR were

fused with the Renilla luciferase reporter in the psiCHECK^{TM}-2 vector.

J Relative luciferase activities in 293T cells after transfection with reporter vectors bearing *IL6ST*-

3' UTR with WT or Mut m⁶A sites. Renilla luciferase activity was measured and normalized to

firefly luciferase activity.

K Relative luciferase activities in WT and *WTAP* KO 293T cells after transfection with reporter

vectors bearing the *IL6ST*-3' UTR with WT m^{6}A sites. Renilla luciferase activity was measured

and normalized to firefly luciferase activity.

L qRT–PCR showing the proportion of *IL6ST* mRNA in polysome fractions from WT and

WTAP$^{Δ1-77}$ THP-1 cells stimulated with LPS for 6 hr.

M, N Immunoblots (**M**) and qRT–PCR (**N**) analyses showing the mRNA and protein abundance of

IL18Rα or IL15Rα in WT and *WTAP$^{Δ1-77}$* THP-1 cells stimulated with LPS at the indicated time

points.

O qRT–PCR showing the mRNA abundance of *IL18R1* in WT and *WTAP$^{Δ1-77}$* THP-1 cells treated

with Act D at the indicated time points.

P Relative luciferase activities in 293T cells after transfection with reporter vectors bearing

IL15RA- or *IL18R1*-3' UTR with WT or Mut m^{6}A sites. Renilla luciferase activity was measured

and normalized to firefly luciferase activity.

Q Relative luciferase activities in WT and *WTAP* KO 293T cells after transfection with reporter

vectors bearing the *IL15RA*- or *IL18R1*-3' UTR with WT m^{6}A sites.

R Relative luciferase activities of reporter vectors bearing *IL15RA*- or *IL18R1*-3' UTR with wild-

type m^{6}A sites after co-transfection with empty vector (EV) or Flag-tagged WTAP into *WTAP* KO

293T cells.

Data information: Data are representative of three independent biological experiments in **A-E** and

M. Data are presented as the mean ± s.d. in **F**, **H**, **J-L** and **N-R**, with individual measurements

overlaid as dots, statistical analysis was performed using a two-tailed Student's *t*-test.

1208

Figure 6. The phase separation of WTAP leads to METTL3 recruitment to modify

inflammatory transcripts efficiently.

A, B Prediction of the disordered regions and prion-like domains (PrD) of WTAP using the

IUPred2A (**A**) or PLAAC (**B**) tool.

C Images of GFP-WTAP droplets formation at room temperature with indicated GFP-WTAP

concentrations (350 mM NaCl).

D LSCM images of 293T cells expressing PEGFP-N1 or GFP-tagged WTAP constructs.

E FRAP assays of the GFP-tagged WTAP droplets before and after photobleaching.

F FRAP quantification of GFP-WTAP droplets over a period of 65 s.

G LSCM images of HeLa cells treated with TNF-α for 2 hr.

H Statistics of the number of WTAP droplets in (**g**).

I Quantitative line profile of colocalization along a white arrow of the image of (**g**).

J Statistics of the number of SC35 droplets in (**g**).

K LSCM images of 293T cells expressing full-length or truncated WTAP labeled with GFP.

L LSCM images of the mCherry-tagged METTL3 droplets (red) in 293T cells before and after co-transfection with GFP-tagged full-length or truncated WTAP.

M FRAP assays of the mCherry-tagged METTL3 droplets before and after photobleaching from (**l**).

N FRAP quantification of mCherry-METTL3 droplets over a period of 100 s.

O METTL3 was immunoprecipitated and RIP-qPCR was performed to assess the association of *IL6ST* transcripts with METTL3.

P LC–MS/MS quantification of m^{6}A abundance in mRNA extracted from 293T cells transfected with indicated plasmids.

Q Working model for WTAP facilitating inflammatory responses through m^{6}A modification and phase separation.

Data information: Data are representative of three independent biological experiments in **C-E**, **G**, and **K-M**. Data are presented as the mean ± s.d. in **H**, **I**, **O-P**, with individual measurements overlaid as dots, statistical analysis was performed using a two-tailed Student's *t*-test. Indicated scale bars are shown in **C-E**, **G**, and **K-M**.

Figure 7. WTAP deficiency alleviates the disease progression of DSS-induced IBD in mice.

A Heatmap showing the expression of WTAP and proinflammatory genes in samples from psoriasis patients before and after anti-IL-17A therapy.

B Heatmap showing the expression of WTAP and proinflammatory genes in samples from CD patients before and after anti-TNF therapy.

C Uniform manifold approximation and projection (UMAP) plots showing rectal tissue-derived

CD45[+] immune cells from all participants. The meaning of the cluster colors is indicated.

D Dot plots showing the expression of METTL14, METTL3, WTAP and IL6ST in major immune

cell groups.

E, F LC-MS/MS quantification of m^{6}A abundance in mRNA extracted from skin tissues in IMQ-

induced psoriasis (**E**) or colon tissues in DSS-induced colitis (**F**) of mice.

G Body weight changes of mice given 3% DSS in their drinking water were monitored daily. n =

7 mice per group.

H The disease activity index of *Wtap*$^{fl/fl}$ and *LyzM-cre*$^+$ *Wtap*$^{Δ1-77}$ mice was scored daily. n = 7

mice per group.

I, J Macroscopic appearances (**I**) and colon lengths (**J**) of *Wtap*$^{fl/fl}$ and *LyzM-cre*$^+$ *Wtap*$^{Δ1-77}$ mice

were recorded on day 6. n = 4 mice per group.

K qRT–PCR showing the mRNA abundance of *Il6*, *Il1α*, *Ccl2* and *Ccl8* in the colon tissues from

Wtap$^{fl/fl}$ and *LyzM-cre*$^+$ *Wtap*$^{Δ1-77}$ mice that had been given 3% DSS in their drinking water for 6

days. n = 10 mice per group.

L Histopathological changes in colon tissue were determined by H&E staining. n = 4 mice per

group.

M The survival of the *Wtap*$^{fl/fl}$ and *LyzM-cre*$^+$ *Wtap*$^{Δ1-77}$ mice after continuous feeding with 3%

DSS was monitored for 15 days. n = 10 mice per group.

Data information: Data are representative of three independent biological experiments in **I** and **L**.

Data are presented as the mean ± s.d. in **E, F, G, K,** with individual measurements overlaid as

dots, statistical analysis was performed using a two-tailed Student's *t*-test. Indicated scale bars are

shown in **I** and **L**.

1267

Expanded View Figure legends

Figure EV1. Hyperinflammation is accompanied by elevated WTAP level in many inflammatory diseases.

A-F Heatmap showing the mRNA abundance of m^{6}A-related genes in patients with SLE (A), asthma (B), sepsis (C), RA (D), psoriasis (E) and IBD (F) compared with healthy controls.

G-L Heatmap showing the mRNA abundance of m^{6}A-related genes in THP-1 cells (G) and human macrophages (H) stimulated with LPS, in lung tissues of mice with LPS-induced lung injury (I), in TNF-α-stimulated HMECs (J), in *Mycobacterium tuberculosis*-infected THP-1 cells (K) or in *S. typhimurium*-infected macrophages (L). The relative mRNA expression of *WTAP* is shown below.

Data are presented as the mean ± s.d. in A-L with individual measurements overlaid as dots, statistical analysis was performed using Mann–Whitney U test in A-F or two-tailed Student's *t*-test in G-L.

Figure EV2. WTAP is an NF-κB p65-stimulated gene.

A, B Luciferase activity analyses in 293T cells transfected with a luciferase reporter for the *WTAP* promoter, together with increasing doses of Flag-tagged C/EBPβ (A) or IRF3 (B).

C Luciferase activity analyses in 293T cells transfected with a luciferase reporter for mouse *Wtap* promoter, together with increasing doses of Flag-tagged NF-κB p65.

D, E qRT-PCR (D) or immunoblots (E) showing the expression of WTAP in THP-1 cells treated with PG490, followed by stimulation with Pam3CSK4 at the indicated time points.

F Immunoblots showing the efficiency of p65 knockout in THP-1 cells.

G, H qRT-PCR (G) or immunoblots (H) showing the expression of WTAP in WT and *RELA*$^{-/-}$ THP-1 cells that were stimulated with HKST or HKLM at the indicated time points.

I DNA pull-down assays showing the effect of C/EBPβ binding with *WTAP* promoter probes.

Data are presented as the mean ± s.d. in A-D and G with individual measurements overlaid as dots, statistical analysis was performed using a two-tailed Student's *t*-test. Data are representative of three independent biological experiments in A-C, E, F, H and I.

Figure EV3. WTAP KO cells express shorter but dysfunctional isoforms of WTAP.

A Schematic representation of the construction of *WTAP* KO THP-1 cells.

B Genotypes of sequenced cells. PCR products were cloned and sequenced.

C Open reading frame prediction.

D Immunoblots showing the expression of WTAP in WT and KO THP-1 cells.

E Schematic showing the *Wtap* conditional KO (cKO) strategy using the Cre-loxP system in

mouse myeloid cells.

F Genotypes of sequenced cells. PCR products were cloned and sequenced.

G Open reading frame prediction.

H Immunogen information for different brands of WTAP antibodies.

I Immunoblots showing the expression of WTAP in WT and KO BMDMs.

J Immunoblots showing the subcellular localization of wild-type and alternative WTAP proteins in

WT, $WTAP^{\Delta 1\text{-}20}$ and $WTAP^{\Delta 1\text{-}77}$ THP-1 cells.

K LC–MS/MS detection the m^6A abundance in mRNA extracted from *WTAP* KO 293T cells

transfected with PEGFP-N1, GFP-tagged WTAP, GFP-tagged $WTAP^{\Delta 1\text{-}20}$ or GFP-tagged $WTAP^{\Delta 1\text{-}77}$

^{77}.

L LC–MS/MS detection the m^6A abundance in mRNA extracted from WT, $WTAP^{\Delta 1\text{-}20}$ and $WTAP^{\Delta}$

$^{1\text{-}77}$ THP-1 cells.

M LC–MS/MS detection the m^6A abundance in mRNA extracted from WT and $Wtap^{\Delta 1\text{-}77}$

BMDMs.

Data are representative of three independent biological experiments in D, I and G. Data are

presented as the mean ± s.d. in K-M, with individual measurements overlaid as dots, statistical

analysis was performed using a two-tailed Student's *t*-test.

1319

Figure EV4. *Wtap* deficiency in myeloid cells did not affect the development or maturation of

major immune cell populations.

A The gating strategy for sorting out various immune cells.

B FACS analyses of major immune cell populations in spleens from $Wtap^{\text{fl/fl}}$ mice and $LyzM\text{-}Cre^{+}$

$Wtap^{\Delta 1\text{-}77}$ mice. n = 3 mice per group.

C Percentage of macrophages, neutrophils, DC and NK cells from (B). n = 3 mice per group.

Data are representative of three independent biological experiments in B. Data are presented as the mean ± s.d. in C, with individual measurements overlaid as dots, statistical analysis was performed using a two-tailed Student's *t*-test.

Figure EV5. WTAP positively regulates the expression of proinflammatory cytokines.

A KEGG analyses of up-regulated DEGs in *WTAP$^{\Delta 1-77}$* THP-1 cells during LPS-induced inflammatory stress compared with WT cells.

B qRT–PCR showing the mRNA abundance of *IL6*, *CCL2*, *CCL8* and *CXCL8* in WT and *WTAP$^{-/-}$* THP-1 cells that were stimulated with Pam3CSK4, HKST or HKLM at 6 hr.

C qRT–PCR and Immunoblots showing the ectopic expression of GFP-tagged WTAP in HeLa cells.

D qRT–PCR showing the mRNA abundance of *IL6* in HeLa cells transfected with GFP-tagged WTAP, followed by stimulation with TNF-α or HKST at different time points.

Data are presented as the mean ± s.d. in B-D, with individual measurements overlaid as dots, statistical analysis was performed using a two-tailed Student's *t*-test. Data are representative of three independent biological experiments in C.

Figure EV6. WTAP aggravates LPS-induced sepsis in mice.

A, B qRT–PCR showing the mRNA abundance of *Il1a*, *Il1b* and *Tnfa* in the lung (A) or colon (B) tissues from *Wtap$^{fl/fl}$* and *LyzM-cre^{+} Wtap$^{\Delta 1-77}$* mice that were intraperitoneally injected with LPS (10 mg kg^{-1}) for 12 hr. n = 10 mice per group.

C, D qRT–PCR showing the mRNA abundance of *Ccl2* (C) and *Ccl8* (D) in the lung tissues from *Wtap$^{fl/fl}$* and *LyzM-cre^{+} Wtap$^{\Delta 1-77}$* mice that were intraperitoneally injected with *P. aeruginosa* or *L. monocytogenes* for 16 hr. n = 8 mice per group.

E-G qRT–PCR showing the mRNA abundance of *Il6* (E), *Ccl2* (F) and *Ccl8* (G) in the colon tissues from *Wtap$^{fl/fl}$* and *LyzM-cre^{+} Wtap$^{fl/fl}$* mice that were intraperitoneally injected with *P. Aeruginosa* or *L. Monocytogenes* for 16 hr. n = 8 mice per group.

Data are presented as the mean ± s.d. in A-G, with individual measurements overlaid as dots, statistical analysis was performed using a two-tailed Student's *t*-test.

1355

Figure EV7. WTAP deficiency significantly reduces the m⁶A modification abundance of

inflammatory genes.

A LC–MS/MS detection the m⁶A abundance in mRNA extracted from 293T cells transfected with

empty vector or GFP-tagged WTAP.

B LC–MS/MS detection the m⁶A abundance in mRNA extracted from WTAP KO and WT 293T

cells.

C Sequence motif was identified within m⁶A peaks by HOMER analyses of WT and *WTAP^Δ1-77*

THP-1 cells with or without LPS treatment.

D Pie charts depicting the proportion of m⁶A peak distribution in the 5' UTR, CDS, and 3' UTR

regions across mRNA transcriptome.

E, F Metagene profiles of m⁶A peak distribution across the 5' UTR, CDS, and 3' UTR in WT and

WTAP^Δ1-77 THP-1 cells (E) or *Wtap^Δ1-77* BMDMs (F) with or without LPS treatment.

G, H EGG enrichment analysis of the transcripts with decreased m⁶A marks in *WTAP^Δ1-77* THP-1

cells (G) or *Wtap^Δ1-77* BMDMs (H) after treatment with LPS.

I qRT–PCR showing the expression of *TNFA*, *IL1B* and *LTA* in THP-1 cells treated with SC144,

followed by stimulation with LPS at the indicated time points.

J qRT–PCR showing the expression of *TNFA*, *IL1B* and *LTA* in WT and *WTAP^Δ1-77* THP-1 cells

that were stimulated with LPS at 3 and 6 hr.

K, L ELISAs were performed to measure IL-1β (K) and TNF-α (L) secretion in supernatants of

WT and *WTAP^Δ1-77* THP-1 cells that were stimulated with HKST or HKLM at 3 and 6 hr.

Data are presented as the mean ± s.d. in A, B and I-L, with individual measurements overlaid as

dots, statistical analysis was performed using a two-tailed Student's *t*-test.

1378

Figure EV8. WTAP deficiency decreased the m⁶A modification of many inflammatory genes.

A-I Integrative Genomics Viewer (IGV) browser tracks showing that m⁶A peaks were enriched in

the 3' UTR or CDS regions of *IL6ST* (**A**), *IL15RA* (**B**), *IL18R1* (**C**), *TYK2* (**D**), *JAK2* (**E**), *RIPK2*

(**F**), *IL1A* (**G**), *IL12B* (**H**), and *CXCL11* (**I**) transcripts.

J-L IGV browser tracks showing that m^{6}A peaks were enriched in the 3' UTR or CDS regions of *Il6st* (**J**), *Jak2* (**K**), and *Ripk2* (**L**) transcripts.

Figure EV9. WTAP promotes the protein output of proinflammatory genes through m^{6}A modification.

A Immunoblots showing the expression of critical adaptors in the IL-6/STAT3 signaling pathway in BMDMs from *Wtap*$^{fl/fl}$ or *LyzM*-Cre$^+$ *Wtap*$^{Δl-77}$ mice.

B, C Immunoblots showing the expression of WTAP and IL6ST in in the lung (B) or colon (C) tissues from wild-type mice that had been intraperitoneally injected with LPS or PBS for 12 hr.

D Relative luciferase activities of reporter vectors bearing *IL6ST*-3' UTR with wild-type m^{6}A sites after co-transfection with empty vector (EV) or Flag-tagged WTAP into *WTAP* KO 293T cells.

E qRT–PCR showing the mRNA abundance of *IL6ST* in WT and *WTAP*$^{Δl-77}$ THP-1 cells or BMDMs.

F qRT–PCR showing the mRNA abundance of *IL6ST* in WT and *WTAP*$^{Δl-77}$ THP-1 cells treated with Act D at the indicated time points.

G Schematic representation of the polysome profiling.

H qRT–PCR showing the proportion of *GAPDH* mRNA in polysome fractions from WT and *WTAP*$^{-/-}$ THP-1 cells treated with LPS for 6 hr.

I Immunoblots showing the expression of IL6ST in WT and *YTHDF1*$^{-/-}$ or *YTHDF3*$^{-/-}$ THP-1 cells.

J Schematic representation of the biotin-labeled probes of *IL6ST* transcripts.

K RNA pull-down analyses showing the interaction between different *IL6ST* RNA probes and the YTHDF1 or YTHDF3 protein.

L YTHDF1 or YTHDF3 were immunoprecipitated, and RNA immunoprecipitation (RIP)-qPCR was performed to assess the association of *IL6ST* with YTHDF1 or YTHDF3 in WT and *WTAP*$^{Δl-77}$ THP-1 cells.

M, N qRT–PCR showing the proportion of *IL6ST* mRNA in polysome fractions from WT and *YTHDF1*$^{-/-}$ (M) or *YTHDF3*$^{-/-}$ (N) THP-1 cells.

O qRT–PCR analyses showing the mRNA abundance of IL18Rα or IL15Rα in WT and $Wtap^{\Delta 1-77}$

BMDMs stimulated with LPS at the indicated time points.

P qRT–PCR showing the mRNA abundance of *IL15RA* in WT and $WTAP^{\Delta 1-77}$ THP-1 cells treated

with Act D at the indicated time points.

Q Relative luciferase activities in 293T cells after transfection with reporter vectors bearing *IL1A*-,

IL12B- or *CXCL11*-3' UTR with WT or Mut m^6A sites. Renilla luciferase activity was measured

and normalized to firefly luciferase activity.

R Relative luciferase activities in WT and *WTAP* KO 293T cells after transfection with reporter

vectors bearing the *IL1A*-, *IL12B*- or *CXCL11*-3' UTR with WT m^6A sites.

S Relative luciferase activities of reporter vectors bearing *IL1A*-, *IL12B*- or *CXCL11*-3' UTR with

wild-type m^6A sites after co-transfection with empty vector (EV) or Flag-tagged WTAP into

WTAP KO 293T cells.

Data are representative of three independent biological experiments in A-C, I and K. Data are

presented as the mean $\pm$ s.d. in D-F, H and L-S, with individual measurements overlaid as dots,

statistical analysis was performed using a two-tailed Student's *t*-test.

1425

Figure EV10. WTAP is deeply involved in the regulation of inflammatory response.

A Venn diagrams showing transcripts with decreased m^6A abundance in $WTAP^{\Delta 1-77}$ THP-1 cells

compared with WT cells stimulated with or without LPS.

B KEGG enrichment analysis of the 1816 transcripts with decreased m^6A abundance only at

resting state.

C KEGG enrichment analysis of the 770 transcripts with decreased m^6A abundance only after

treatment with LPS for 4 h.

1433

Figure EV11. The phase separation of WTAP leads to METTL3 recruitment to modify

inflammatory transcripts efficiently.

A Bacterial purified GFP-WTAP proteins were analyzed by SDS-PAGE and detected by

coomassie blue staining.

B Images of GFP-WTAP droplets formation at room temperature with indicated NaCl

concentrations (5 μM GFP-WTAP).

C EqDiameter (the diameter of a circle with the same area as the measured object) frequency

distribution of GFP-WTAP liquid droplets formed at the indicated timepoints (350 mM NaCl, 5

μM GFP-WTAP).

D Time-lapse micrographs of the fusion at room temperature of GFP-WTAP (5 μM) liquid

droplets formed at the indicated times.

E GFP-WTAP were treated with 5% Hex and then subjected to droplet formation assay *in vitro*

(350 mM NaCl, 5 μM GFP-WTAP).

F Immunoblots showing of the expression of GFP-tagged WTAP.

G qRT–PCR showing the mRNA abundance of *WTAP* in HeLa cells stimulated with TNF-α at the

indicated time points.

H Prediction of the unfolded intrinsically disordered regions (IDRs) of WTAP using the PONDR

tool, and a series of truncated WTAP labeled with GFP are as shown.

I, J Immunoblots (I) and representative confocal images (J) of 293T expressing truncated WTAP

labeled with GFP.

K Bacterial purified mCherry-METTL3 proteins were analyzed by SDS-PAGE and detected by

coomassie blue staining.

L Fluorescence of mCherry-METTL3 (10 μM) and GFP-WTAP (10 μM) mixed at room

temperature with 350 mM NaCl. Quantitative line profile of colocalization along a white arrow of

the image on the left.

M Coimmunoprecipitation and immunoblot analyses of extracts of 293T cells transfected with

various combinations of plasmids encoding GFP-tagged WTAP and mCherry-tagged METTL3.

N, O METTL3 was immunoprecipitated (N) and RIP-qPCR was performed to assess the

association of *IL15RA* or *IL18R1* transcripts with METTL3 (O).

P m^6A dot quantification of m^6A abundance in mRNA extracted from *WTAP* KO 293T cells

transfected with indicated plasmids.

Data are representative of three independent biological experiments in A-F, I-N and P. Data are

presented as the mean ± s.d. in G and O, with individual measurements overlaid as dots, statistical

analysis was performed using a two-tailed Student's *t*-test. Indicated scale bars are shown in B, D,

E, J and L.

1469

Figure EV12. The expression of WTAP and IL6ST is increased in patients with

inflammatory diseases.

A The mRNA abundance of *IL6ST* in patients with SLE, asthma, RA, psoriasis and IBD compared

with those in healthy controls.

B, C The mRNA abundance of *WTAP* and *IL6ST* in patients with IBD compared with healthy

controls.

D Spearman correlations between m^{6}A regulators and IBD core cytokines in IBDMDB patients.

E, F Spearman correlations between the expression of WTAP and the global m^{6}A abundance in

RA (E) patients, or in ovalbumin (OVA)-induced acute allergic asthma mice (F).

G, H m^{6}A dot blot quantification of m^{6}A abundance in mRNA extracted from skin tissues in IMQ-

induced psoriasis (G) or colon tissues in DSS-induced colitis (H) of mice.

Data are presented as the mean ± s.d. in A-C, with individual measurements overlaid as dots,

statistical analysis was performed using Mann–Whitney U test. Data are representative of three

independent biological experiments in G and H.

1484

Figure EV13. Reducing the level of m^{6}A modification can reverse the high inflammatory

state.

A m^{6}A dot blot quantification of m^{6}A levels in mRNA extracted from BMDMs that treated with

DMSO or STM2457, followed by stimulation with HKST for 6 hr. UN, unstimulated.

B Heatmap showing the mRNA expression of proinflammatory genes in BMDMs treated with

DMSO or STM2457, followed by stimulation with HKST for 6 hr. The number represents three

independent experiments.

C, D m^{6}A dot blot quantification of m^{6}A levels in mRNA extracted from colon (C) or lung (D)

tissues in LPS-induced sepsis of mice, followed by four consecutive days of intraperitoneal

injections of DMSO or STM2457. n = 3 mice per group.

E, F qRT–PCR showing the mRNA abundance of *Il1b*, *Tnfa*, *Il16*, *Ccl2* and *Ccl8* in the colon (E)

or lung (F) tissues from mice treated as above. n = 6 mice per group.

G H&E assays showing the lung injury of LPS-induced sepsis of mice, followed by four

consecutive days of intraperitoneal injections of DMSO or STM2457. Scale bars, 50 µm. n = 3

mice per group.

H m^{6}A dot blot quantification of m^{6}A levels in mRNA extracted from THP-1 cells that treated

with DMSO or PG490, followed by stimulation with LPS at the indicated time points.

Data are representative of three independent biological experiments in A, C, D and H. Data are

presented as the mean ± s.d. in B, E and F, with individual measurements overlaid as dots,

statistical analysis was performed using a two-tailed Student's *t*-test. Indicated scale bars are

shown in I.

1506

A GSE137218

Nonlesional Skin | Lesional psoriatic skin (After anti-IL-17A (secukinumab) treatment)
0 d | 4 d | 14 d | 42 d | 84 d

WTAP
CCL20
IL23A
IL17A
CXCL1
IL6
IL12B
IL22
CXCL10
CCL8
CCL18

B GSE16879

Anti-TNF Non-Responders — After / Before
Anti-TNF Responders — After / Before
HC

WTAP IL6ST IL6 CCL2 CCL8 TNF CCL18 CXCL10

C GSE125527

UMAP_1 / UMAP_2

CD4+ T cells
B cells
Plasma cells
CD8+ T cells
T cells
Monocytes
Progenitors

D

Healthy / UC

METTL14
METTL3
WTAP
IL6ST

CD4+ T cell | B cell | Plasma cell | CD8+ T cell | T cell | Monocyte | Progenitor

Percent Expressed: 10, 20, 30
Average Expression: 2, 0, -2

E Skin

m6A/A in mRNA

Vaseline / IMQ
P = 0.0164

F Colon

m6A/A in mRNA

Water / 3% DSS
P = 0.0113

G

Body Weight Changes (%)
0 1 2 3 4 5 6 7 days

P = 0.0025
P < 0.0001
P < 0.0001

*Wtap*fl/fl
*LyzM-cre+ Wtap*Δ1-77

H

Disease Activity Index
0 1 2 3 4 5 6 7 days

*Wtap*fl/fl
*LyzM-cre+ Wtap*Δ1-77

P = 0.0001
P = 0.0003
P < 0.0001
P < 0.0001

I

*Wtap*fl/fl : 1# 2# 3# 4#
*LyzM-cre+ Wtap*Δ1-77 : 1# 2# 3# 4#

J

Colon Length (cm)

*Wtap*fl/fl
*LyzM-cre+ Wtap*Δ1-77
P = 0.0007

K

Water / 3% DSS

*Wtap*fl/fl
*LyzM-cre+ Wtap*Δ1-77

L

Il6 — P = 0.0254
Il1a — P = 0.0171
Ccl2 — P = 0.0299
Ccl8 — P = 0.0009

Rel. mRNA Level

*Wtap*fl/fl
*LyzM-cre+ Wtap*Δ1-77

M Survival Curve

Percent survival (%)
6 8 10 12 14 16
Time after 3% DSS-administration (Days)

*Wtap*fl/fl
*LyzM-cre+ Wtap*Δ1-77
P = 0.0045

A

WTAP^{Δ1-77} vs WT

0 h_unique

1816   2633   770   4 h_unique
        (common)

m⁶A Level_Down   m⁶A Level_Down
(LPS-0 h)   (LPS-4 h)

B

KEGG Enrichment

Lysine degradation
Inositol phosphate metabolism
ErbB signaling pathway
Glycosaminoglycan biosynthesis
Chronic myeloid leukemia
Biotin metabolism

P-value
0.01
0.02
0.03
0.04

Gene count
0   5   10   15

C

KEGG Enrichment

NF−kappa B signaling pathway
Osteoclast differentiation
TNF signaling pathway
Cytokine−cytokine receptor interaction
NOD-like receptor signaling pathway
Notch signaling pathway
Toll-like receptor signaling pathway
C-type lectin receptor signaling pathway
IL17A signaling pathway
TGF-beta signaling pathway
Pathogenic Escherichia coli infection
Lipid and atherosclerosis
Lysine degradation
Shigellosis
Toxoplasmosis
MAPK signaling pathway

P-value
0.005
0.010
0.015
0.020
0.025

Gene count
0   10   20   30