Shifted levels of sleep and activity under darkness as mechanisms underlying ectoparasite resistance

Joshua B. Benoit[1], Oluwaseun M. Ajayi, Ashley Webster, Karl Grieshop, David Lewis, Hailie Talbott, Joy Bose, and Michal Polak

Department of Biological Sciences, University of Cincinnati, Cincinnati, OH 45211

[1]Author for correspondence: Joshua B. Benoit, Department of Biological Sciences, University of Cincinnati, OH 45221, USA.
E-mail address: joshua.benoit@uc.edu (J.B. Benoit).

**Abstract -** Parasites harm host fitness and are pervasive agents of natural selection. Host defensive traits in natural populations typically show genetic variation, which may be maintained when parasite resistance imposes fitness costs on the host coupled with variability in parasite prevalence in space and/or time. Previously we demonstrated significant evolutionary responses to artificial selection for increasing behavioral immunity to *Gamasodes queenslandicus* mites in replicate lines of *Drosophila melanogaster*. Here, we report transcriptional shifts in metabolic processes between selected and control fly lines based on RNA-seq analyses. We also show decreased starvation resistance and increased use of nutrient reserves in flies from mite-resistant lines. Additionally, mite-resistant lines exhibited increased behavioral activity, reduced sleep and elevated oxygen consumption under conditions of darkness. The link between resistance and sleep was confirmed in an independent panel of *D. melanogaster* genetic lines exhibiting variable sleep durations, showing a positive correlation between mite resistance and reduced sleep. Experimentally restraining the activity of artificially selected mite-resistant flies during exposure to parasites under dark conditions reduced their resistance advantage relative to control flies. The results suggest that ectoparasite resistance in this system involves increased dark-condition activity and metabolic gene expression at the expense of nutrient reserves and starvation resistance.

**Keywords:** host sleep, parasitism, resistance, costs, nutrient reserves, metabolic gene expression

**Introduction**

Parasites are ubiquitous, and parasite-mediated selection can represent a strong evolutionary pressure that influences many host phenotypic traits and ecological processes (1–4). Defensive adaptations against parasites are common in plant, animal, and microbial host populations, where genetic variation in resistance is an essential component of host capacity to respond evolutionarily to parasite pressure (5–9). *Drosophila* and the many parasites that target species within this genus represent excellent systems to examine mechanisms of resistance and the factors that promote and constrain their evolution (10–14). Specifically, associations between *Drosophila* and ectoparasitic mites (15–19) are exceptional models for studies of ectoparasitism, as many such systems that are naturally occurring are also experimentally tractable, and the behavioral traits flies deploy against mites parallel those seen in other animal hosts (20). The benefits ectoparasitic mites may derive from attaching to flies include increased nutrient intake and reproductive output, and improved dispersal to new habitats (21). The host is harmed through damage to the cuticle and resource extraction along with costly immune and repair responses (5, 15, 18, 22). The combined effects of mite-derived damage negatively influence the expression of male and female reproductive tissues, mating success, and lifespan (5, 15, 22). Other host behavioral and physiological changes that occur in response to parasitism or parasite exposure include adaptive shifts in male reproductive effort (23), anti-mite defensive behaviors (24), oviposition site preference (25), and rates of respiration (26).

*Drosophila melanogaster* and other fruit fly species have been noted to be parasitized by *Gamasodes* mites in Asia (Taiwan and Thailand) and Australia (17, 19, 22). The extended geographic distribution and host range of *Gamasodes* mites indicates that they are likely to be a pervasive selective force affecting many aspects of fly biology. The behaviors that flies use to avoid infestation by ectoparasitic mites include bursts of flight from the substrate, locomotor responses, and vigorous grooming once the mite has made contact with the fly (24, 27). Whereas strong selection is likely to act on these first-line forms of defense (27, 28), the molecular and physiological bases of these defensive traits are yet to be established.

In the present study of *D. melanogaster*, RNA-seq and functional studies revealed that metabolic activity increased in response to artificial selection for increased behavioral resistance to *Gamasodes* mites (27). The increased metabolism was associated with reductions in nutrient reserves and starvation resistance and was correlated with increased activity and reduced sleep during the dark. These results identify metabolic and activity changes as a potential causal factor underlying ectoparasite resistance. Resistance assays of an independent panel of *Drosophila* genetic lines with variable sleep supported a link between decreased sleep and enhanced parasite avoidance. The results suggest that increased activity and reduced sleep could be a critical factor in preventing ectoparasite attachment and that this energetically costly behavioral immunity trades off with nutrient reserves and starvation resistance.

**Results**

Ectoparasite resistance yielded transcriptional changes associated with metabolism

Previously we performed artificial selection for increasing resistance to *Gamasodes queenslandicus* mites in male *D. melanogaster* for 16 generations (27), which we continued in the present study for a total of 22 generations. The selection program resulted in significant evolutionary responses, yielding three replicate selection lines that were significantly more behaviorally resistant to ectoparasitic mites than their unselected counterparts (Fig. 1A). At the terminus of the 22 generations of selection, transcriptional analyses revealed that 108 genes were significantly differentially expressed (68 up-regulated and 40 down-regulated) in selection lines relative to control lines (three biological replicates of each of the three replicate lines per selection treatment) (Table S1). Gene ontology enrichment analyses revealed increased oxidoreductase activity in selection lines relative to controls along with increases in other factors, such as three genes involved in glycolysis processes (aldolase, acetyl-CoA synthetase, and phosphoglycerate kinase) (Fig. 1B, Table S1). These results suggest that there are potential differences in metabolism that occur in flies selected for mite resistance.

Starvation resistance decreased with increased metabolism under dark periods

Compared to control lines, the observed upregulation of metabolism-associated factors among resistant lines suggests that a trade-off exists between resistance and host nutrient reserve management. Consistent with this expectation, starvation resistance in ectoparasite-resistant lines was reduced compared to control lines (Fig. 2A). This apparent trade-off was corroborated by the observation of greater reductions in body lipid and protein reserves after 36 hours of starvation in resistant lines relative to controls (Fig. 2B, C). This difference in levels of nutrient stores suggests relatively increased lipolysis and proteolysis in the resistant lines. Thus, there are discernible metabolic phenotypes that co-evolved with ectoparasite resistance under artificial selection, identifying potential cost of ectoparasite resistance as a reduction in host survival resulting from accelerated reduction in body condition under nutrient deprivation.

To further probe this metabolic trade-off, we measured oxygen consumption of flies during a daily cycle of light and darkness. Surprisingly, oxygen consumption was not different between control and selected lines under light conditions (Fig. 2D). However, when oxygen consumption was measured during the dark, it was significantly higher in the resistant lines compared to controls (Fig. 2D). This result indicates that the metabolic costs of ectoparasite resistance primarily manifest under dark conditions.

Activity during darkness was greater in ectoparasite-resistant lines

As oxygen consumption was increased under dark conditions, we tested whether fly activity and sleep profiles were altered following selection throughout a daily cycle of light and dark conditions (Fig. 3A). As with oxygen consumption, activity and sleep levels were not significantly different during the light. However, resistant lines did show a significant increase in activity and a decrease in the duration of sleep bouts during the dark relative to controls (Fig. 3B, C) – two traits that are certainly related but not one in the same. Whereas activity was measured as the number of instances a fly crossed an infrared beam per unit time, a bout of sleep was defined as a minimum of 5 min of continuous inactivity. To further evaluate whether reduced sleep and increased activity profiles were associated with resistance to ectoparasitism, we exposed an independent panel of inbred fly lines that differed in activity and sleep levels to ectoparasitic mites (29). There was a

significant relationship between the level of sleep under dark conditions and the proportion of flies parasitized by mites (Fig. 3D). This result supports the conclusion that sleep and activity levels are causal factors affecting ectoparasite resistance in flies.

Movement restriction and assay under light conditions reduced ectoparasite resistance

To confirm the causal role of activity and sleep-like behavior in mediating ectoparasite resistance, we manipulated activity by restraining flies. We found that selected lines no longer showed improved resistance under dark conditions relative to controls when restrained (Fig. 4A), matching a previous study showing that flight restriction erodes mite resistance (27). Secondly, we performed mite parasitism assays under light and dark conditions, and showed that resistance divergence between selected and control lines was reduced in light relative to dark conditions (Fig. 4B). These results support the conclusion that increased activity under dark conditions and reduced sleep-like behavior are important factors mediating host response to selection for improved mite resistance.

**Discussion**

This study indicates that declining sleep and increased activity are critical mechanisms for preventing ectoparasitism in fruit flies. Following artificial selection for resistance (27), specific metabolic transcriptional pathways associated with increased oxidoreductase and monooxygenase activity along with a multitude of other factors such as specific genes underlying glycolysis (*e.g.,* aldolase, acetyl-CoA synthetase, and phosphoglycerate kinase) were up-regulated. These expressional changes are supported by increased metabolic activity at the expense of starvation resistance in flies resistant to mite parasitism. This finding matched increased activity and reduced sleep during dark conditions in selected versus control flies, which is likely a driving factor governing the evolved resistance to ectoparasitic mites. We saw experimental confirmation that altered sleep and activity profiles are critical for ectoparasite resistance when fly activity was prevented and an independent panel of fly lines of varying sleep levels was assayed.

Sleep and host-parasite dynamics have been studied in multiple systems, but these have predominantly involved endoparasites such as malaria (30, 31). In general, the response to internal parasites requires a robust immune response, where increased

sleep likely improves this response by allowing the increased expression of immune factors (30, 31). In contrast, host sleep (or lack thereof) could influence ectoparasite resistance differently, as behavioral resistance, rather than immune response, is of greater importance to avoid infestation. Here, we confirmed that among resistant lines, sleep decreased and activity increased during darkness, which is when sleep predominantly occurs in *D. melanogaster* (32, 33). Importantly, there could be differences that occur during light conditions, but differences may not have been apparent as *D. melanogaster* sleeps much less during the day (32, 33). These shifted daily patterns likely allow the flies to be more responsive during encounters with mites under dark conditions. Importantly, as the artificial selection was conducted under both light and dark conditions (spanning the diurnal transition) (27), this raises the possibility that a small change in sleep patterns or activity in the dark could have been a direct target of selection that yielded the increased resistance to mite infestation. The altered sleep and activity profiles are likely to be a general phenomenon that occurs among animals in the presence of parasites. For example, ectoparasitism has been noted in birds and bats to be associated with shifts in sleeping and rest behavior (34, 35). In primates, changes in sleep patterns have been suggested to reduce exposure to parasitic worms (36).

The present study is unique in providing integrated transcriptional, physiological and behavioral evidence for a causal link between reduced host sleep, increased activity, and improved resistance to ectoparasites. Recently, phototaxis effects have been noted during fly-mite interactions (37), where *Drosophila nigrospiracula* exposed to parasitic mites under light conditions had lower mite parasitism than those held in the dark with mites. Furthermore, flies will reside more in lighted areas after exposure to mites (37). As light exposure increases fruit fly activity and reduces sleep (32, 33), the studies on *D. nigrospiracula* provide additional support for our observations in *D. melanogaster* that activity and sleep are critical to fly-mite interactions.

Parasite resistance may be associated with distinct fitness costs, which include reduced stress tolerance and damaged fecundity identified in multiple animal systems (2–4) including other fly-mite systems (5, 27). The increased expression of metabolism-associated genes and altered starvation resistance in selection lines reported here suggest that mite resistance is likely to alter fly physiology in relation to metabolic output,

which we demonstrated through the assessment of lipid reserves during starvation, and altered oxygen consumption. Our artificial selection experiment may have allowed mite resistance to evolve despite these metabolic costs because the *ad libitum* food conditions reduced the strength of selection for nutrient management. Importantly, direct parasitism by mites induces a physiologically costly immune response (5, 22), and significantly reduces host egg production and lifespan (23, 38), thus avoiding parasitism is critical. Thus, the optimal degree of metabolically demanding mite resistance behaviors in nature is likely determined by the balance of parasite pressure versus resource limitation and nutrient management, which could vary temporally, geographically and taxonomically.

The finding that altered sleep patterns impact ectoparasite resistance represents a novel form of host resistance, similar to those involved in predator-prey dynamics. Indeed, many ectoparasites such as blood-feeding flies and mites have been classified as micropredators (39, 40), and differences in sleep have been associated with predation levels (41–43), such that animals with increased exposure to predators have reduced or lighter sleep (42, 44). Here, we showed that reduced sleep-like behavior in *D. melanogaster* was the target of selection for parasite resistance, providing evidence that sleep behavior can directly impact interactions with an ectoparasite. The phenotypes that confer ectoparasite resistance occur with a direct metabolic cost that could act as a trade-off constraining the evolution of resistance to the mites, especially when parasitism rates happen to be low and/or nutrients limiting.

**Acknowledgments**

The research was supported by National Science Foundation (NSF) grant 1654417 (to M.P. and J.B.B.). Partial funding for reusable equipment was provided indirectly by the United States Department of Agriculture 2018-67013 and the National Institute of Allergy and Infectious Diseases, award numbers R01AI148551 and R21AI166633 to J.B.B. We thank Rachel Black and Hannah Tran for their excellent assistance in media preparation.

**Materials and methods**

Flies, mites, and selection for resistance

The flies (*Drosophila melanogaster*) and mites (*Gamasodes queenslandicus*) were collected from Australia as described previously (27). The selection protocol was described previously (27), which resulted in a significant increase in resistance to mite parasitism after eight generations that continually increased to 16 generations of selection and remained high through 22 generations. The resistance levels were assessed every two to four generations (27, 45). Even though selection was only conducted on males, both sexes showed increased resistance (27). Furthermore, increased resistance to another mite species (*Macrocheles subbadius*) was noted (27), indicating that the defensive mechanisms that were selected are likely general to ectoparasitic mites. Lines of *D. melanogaster* with varying levels of sleep were acquired from the Bloomington Stock Center from short and long sleep lines that were cross-bred to increase the levels of sleep (29). Male flies were used in these analyses.

Mite infestation experiments

Mite infestation experiments were conducted according to previously developed methods (5, 22, 24, 38) that allow for determining infestation differences between fly lines. These were assays similar to those used for selection, where flies were exposed to mites in a jar containing media used to rear *G. queenslandicus* (22, 24, 38) . Each assay measured ectoparasitism in each fly line relative to its paired control line across a set of replicate infestation chambers. For each comparison, flies equal in size were aspirated into a chamber (total flies ranged between 40 and 50 flies). Minute wing clips at the tip of each wing were used to distinguish groups (≤ 3% of the wing), which have been previously shown to have no impact on fly susceptibility to mites (22, 24), and clipping was alternated between groups and across chambers. A recovery period of 24h was used after clipping the wing under $CO_2$. These jars were examined after 6-8 hours based on the relative number of mites in the media. Direct presence of mites and scars on flies from chambers in relation to the wing clipping that marks the specific lines was used to assess the prevalence of parasitism. These parasitic assays were conducted on the selected and control lines after 22 generations under either full light or in complete darkness (selection

line S3 was lost during COVID-19 restrictions as such only S1 and S2 were used in the light and dark comparative analyses) and for flies from the sleep inbred panel. For the sleep inbred panel, lines from the panel were intermixed during the assays to allow for random comparisons between each line.

For susceptibility tests under restraint, flies were transferred to shortened 200-µl micropipette tips, a confined space that severely limits fly movement (45, 46), and exposed to two mites, effectively mimicking periods of reduced activity. After two hours in the dark, the flies were removed and examined for attached mites or mite-induced scars. This assay was conducted on the selected and control lines after 20 generations of artificial selection.

RNA-seq analysis following selection

Flies were collected after 16 generations of selection for the RNA-seq studies. RNA was extracted from the flies as previously described (22) in three biological replicates per line. Illumina sequencing was conducted at Novogene on an Illumina NovaSeq Hiseq platform. Illumina datasets are available at the National Center for Biotechnology Information's Sequence Read Archive: Bioproject PRJNA999910. Differential expression of genes was determined based on methods previously described (22, 47, 48). Briefly, the quality and presence of adapters were determined with FASTQC (49), followed by a filtering/trimming procedure with Trimmomatic v.0.36 using default settings (50). At least 30 million reads were available for each sample following trimming. Reads were mapped to the transcript assembly release 6.10 from Flybase (51) using Kallisto v0.48.0 with recommended settings (52) summarized to the gene level using tximport v1.12.3 (53). Differential gene expression analysis was performed with DESeq2 (54) with the read counts generated from Kallisto. Filtering in DESeq2 was conducted to remove predicted transcripts with fewer two mapped reads. Selection-specific effects to identify overlapping differences among lines were determined by a nested analysis that included each line and replicates. An adjusted p-value of ≤ 0.01 was used to identify genes as differentially expressed. Specific enriched gene ontology (GO) categories were determined using g:Profiler with gene sets increased and decreased during selection analyzed separately (55) and treemaps to visualize the GO was generated with REVIGO (56).

Starvation assay, nutritional reserve levels, oxygen consumption.

Flies were collected after 17 and 18 generations of selection for these assays. Flies were transferred to 1% agar vials to provide water without nutrients (1 or 3 males per vial, no difference was noted in the duration of starvation between the two groups, and the results were combined), and the number of dead flies was noted every twelve hours until all flies were dead. Nutrient reserve levels were measured during starvation based on previously used methods (22, 57), which use spectrophotometric assays to establish the lipid and protein levels in relation to the dry mass of flies and relative to a protein or lipid standard. Differences in oxygen consumption between select and control lines were determined using a microrespirometer (57, 58). In brief, individual flies were placed in microrespirometers were positioned in a water bath (25°C) and were allowed 15 minutes to equilibrate before measurements. $CO_2$ production, measured through the movement of KOH, was measured every hour for 6 hours. Assuming that one mole of $O_2$ is consumed for every mole of $CO_2$ released, oxygen consumption was calculated using the distance traveled by the KOH and expressed as µl O2 mg dry mass$^{-1}$ h$^{-1}$.

Sleep and activity assessment

The rest-activity rhythms of the flies (control, selected, and sleep lines) were quantified with the aid of a Locomotor Activity Monitor 25 (LAM25) system (TriKinetics Inc., Waltham, MA, USA) and the DAMSystem3 Data Collection Software (TriKinetics). Flies were collected after 18-22 generations of selection and those from the sleep inbred panel (29) were used for these assays. Individuals were loaded into 25 x 150 mm clear glass tubes with access to fly media provided *ad libitum*. The glass tubes were then positioned horizontally in the LAM25 system, allowing the simultaneous recording of 32 individuals in an "8 x 4" horizontal by vertical matrix. Replicates were intermixed between trials to allow for randomization. The entire set-up was placed in a light-proof incubator supplied with its own lighting system at 24°C, 70-75% RH, under a 12hr:12hr L/D cycle. Following the acclimation of individuals for 48 hours, activity level was measured in 1 min bins (as the number of times an individual crosses an infrared beam in 1 min), and sleep levels were assessed as 5 min of continuous inactivity (29, 32). This allows for the determination of sleep bouts, duration of bouts, and total sleep. No differences were noted in the sleep

bouts or total sleep, but the duration of bouts were significantly different between control and selected lines. Although sleep is related to activity, there are distinct differences in the quantification of activity and sleep, and sleep levels are not directly an inverse of general activity levels.  Data collected with the DAMSystem3 was processed using the Rethomics platform in R with its associated packages including *behavr, ggetho*, *damr* and *sleepr* (59).

Statistical analyses

For all analyses, we used the R software (60) and plots were generated with *ggplot*. Survival analyses were determined with a Kaplan-Meier test and compared with a general Wilcoxon test. General linear models were used to examine the interactions between selection status and response variables where replicates for each selection and control lines are nested as a random effect within the selection and control treatments to establish that effects are significant among control and selected groups. Statistical outputs and replicate numbers are provided within the figure legends.

**Figure legends**

**Figure 1 - Selection increases resistance to mites and RNA-seq studies reveal altered metabolic-associated factors following selection for ectoparasite resistance.** A. Artificial selection for mite resistance resulted in significantly reduced susceptibility to mites in selected (S) versus control (C) lines after 22 generations. * denotes significance at $P < 0.05$ between control and selected lines. (F = 16.24, d.f. = 2,5, P = 0.001) Error bars represent the 95% confidence intervals. B. Gene ontology categories enriched in differentially expressed genes between S and C from a full analysis of all samples (nested within the replicate, revealing factors generally associated with metabolism, FDR, P < 0.05). There were three biological replicates of RNA-Seq data for each treatment (S and C) of the three artificial selection replicates.

**Figure 2 - Flies with increased resistance have higher metabolism during the dark conditions.** A. Death due to starvation was increased following selection in all three lines. Kaplan-Meier, generalized Wilcoxon test, Chi-square = 18.234, P < 0.001 for control compared to selected for each line, N = 150 individuals per line. B and C. Lipolysis (F = 11.84, d.f. = 2,5, P = 0.031) and protein (F = 11.08, d.f. = 2,5, P = 0.001) breakdown was increased following selection in all lines. N = 3 groups of three flies examined per line. D. Oxygen consumption was increased in the selected flies under darkness (scotophase) compared to light conditions (photophase) (F = 10.47, d.f. = 1,4, P = 0.009). N = 15 flies measured per line when selected and control lines are compared without light. * denotes significance at $P < 0.05$ between control and selected lines

**Figure 3 - Sleep and activity profiles are altered in selected lines.** A. Daily activity profile for control and selected fly lines, all lines combined. Shaded areas represent the 95% confidence intervals. N = 40-45 individuals per line. B. Sleep duration decreased during the dark cycle for lines selected for resistance (F = 19.291, d.f. = 1,4, P < 0.001). Other aspects of sleep, such as bouts or duration of bouts, are not significantly different between selected and control lines ( P > 0.05 in all cases). C. Activity is increased

during the night for lines selected for resistance (F = 15.98, d.f. = 1,4, P = 0.002).Each point represents the activity of a single fly. D. A significant interaction occurs between an independent panel of eight inbred lines with different sleep levels (29) and susceptibility to mites  (F = 3.021, d.f. = 7,40, P = 0.012). N = 6 for each line was assayed for parasitism. * denotes significance at P < 0.05 between control and selected lines

**Figure 4 - Factors that alter sleep status shift ectoparasite resistance.** A. Restraining flies reduces the resistance of selected lines compared to controls, (F = 28.66, d.f. = 1,4, P < 0.002).  B. The effect of selection on mite resistance is more profound when flies are exposed to mites under dark conditions. N = 4 parasitism assays per line (F = 11.839, d.f. = 1,3, P = 0.016). C. Summary of dynamics between sleep and activity in relation to resistance to ectoparasitic mites. This highlights that altered sleep and activity patterns are associated with resistance to parasitism.

**Supplemental Tables**

**Table S1** - Genes with significantly different $\log_2$ fold-change in expression (normalized read counts) between selection and control samples when all lines are included in a single nested analysis.

**References**

1. M. E. Scott, A. Dobson, The role of parasites in regulating host abundance. *Parasitology Today* **5**, 176–183 (1989).

2. P. W. Ewald, The evolution of virulence: a unifying link between parasitology and ecology. *J. Parasitol.* **81**, 659–669 (1995).

3. D. A. Windsor, Controversies in parasitology, Most of the species on Earth are parasites. *Controversies in parasitology, Most of the species on Earth are parasites. International Journal for Parasitology* **12**, 1939–1941 (1998).

4. P. S. Fitze, B. Tschirren, H. Richner, Life history and fitness consequences of ectoparasites. *J. Anim. Ecol.* **73**, 216–226 (2004).

5. L. T. Luong, M. Polak, Environment-dependent trade-offs between ectoparasite resistance and larval competitive ability in the *Drosophila–Macrocheles* system. *Heredity* **99**, 632–640 (2007).

6. H. J. Henter, S. Via, The potential for coevolution in a host-parasitoid system. I. Genetic variation within an aphid population in susceptibility to a parasitic wasp. *Evolution* (1995).

7. C. Combes, *The Art of Being a Parasite* (University of Chicago Press, 2020).

8. M. D. Rausher, Co-evolution and plant resistance to natural enemies. *Nature* **411**, 857–864 (2001).

9. E. Mazé-Guilmo, G. Loot, D. J. Páez, T. Lefèvre, S. Blanchet, Heritable variation in host tolerance and resistance inferred from a wild host–parasite system. *Proceedings of the Royal Society B: Biological Sciences* **281**, 20132567 (2014).

10. J. Jaenike, S. J. Perlman, Ecology and evolution of host-parasite associations: mycophagous *Drosophila* and their parasitic nematodes. *Am. Nat.* **160 Suppl 4**, S23–39 (2002).

11. B. Lemaitre, J. Hoffmann, The host defense of *Drosophila melanogaster*. *Annu. Rev. Immunol.* **25**, 697–743 (2007).

12. M. D. Fellowes, A. R. Kraaijeveld, H. C. Godfray, Trade-off associated with selection for increased ability to resist parasitoid attack in *Drosophila melanogaster*. *Proc. Biol. Sci.* **265**, 1553–1558 (1998).

13. C. Neyen, A. J. Bretscher, O. Binggeli, B. Lemaitre, Methods to study Drosophila immunity. *Methods* **68**, 116–128 (2014).

14. E. O. Campbell, L. T. Luong, Mite choice generates sex- and size-biased infection in Drosophila hydei. *Parasitology* **143**, 787–793 (2016).

15. M. Polak, T. A. Markow, Effect of ectoparasitic mites on sexual selection in a Sonoran desert fruit fly. *Evolution* **49**, 660–669 (1995).

16. R. B. Halliday, The Australian species of Macrocheles (Acarina : Macrochelidae). *Invertebr. Syst.* **14**, 273–326 (2000).

17. R. B. Halliday, D. E. Walter, M. Polak, A new species of *Gamasodes* Oudemans from Australia (Acari: Parasitidae). *Zootaxa* **1001**, 17 (2005).

18. A. Perez-Leanos, M. R. Loustalot-Laclette, N. Nazario-Yepiz, T. A. Markow, Ectoparasitic mites and their *Drosophila* hosts. *Fly* **11**, 10–18 (2017).

19. M.-Y. Yao, J.-J. Guo, P. Michal, T.-C. Yi, D.-C. Jin, A new species and new record of *Gamasodes* (Mesostigmata: Parasitidae) from China. *Systematic and Applied Acarology* **25**, 1299–1318 (2020).

20. B. L. Hart, L. A. Hart, How mammals stay healthy in nature: the evolution of behaviours to avoid parasites and pathogens. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **373** (2018).

21. D. E. Walter, H. C. Proctor, "Mites and Biological Diversity" in *Mites: Ecology, Evolution & Behaviour: Life at a Microscale*, D. E. Walter, H. C. Proctor, Eds. (Springer Netherlands, 2013), pp. 447–459.

22. J. B. Benoit, J. Bose, S. T. Bailey, M. Polak, Interactions with ectoparasitic mites induce host metabolic and immune responses in flies at the expense of reproduction-associated factors. *Parasitology* **147**, 1196–1205 (2020).

23. M. Polak, W. T. Starmer, Parasite–induced risk of mortality elevates reproductive effort in male *Drosophila*. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **265**, 2197–2201 (1998).

24. M. Polak, Heritability of resistance against ectoparasitism in the *Drosophila-- Macrocheles* system. *J. Evol. Biol.* **16**, 74–82 (2003).

25. C. J. Horn, M. K. Mierzejewski, M. E. Elahi, L. T. Luong, Extending the ecology of fear: Parasite-mediated sexual selection drives host response to parasites. *Physiol. Behav.* **224**, 113041 (2020).

26. T. Brophy, L. T. Luong, Ectoparasite-induced increase in *Drosophila* host metabolic rate. *Physiological Entomology* **46**, 1–7 (2021).

27. M. Polak, J. Bose, J. B. Benoit, H. Singh, Heritability and pre-adult survivorship costs of ectoparasite resistance in the naturally occurring *Drosophila-Gamasodes* mite system. *bioRxiv*, 2022.12.15.520609 (2022).

28. P. Schmid-Hempel, D. Ebert, On the evolutionary ecology of specific immune defence. *Trends Ecol. Evol.* **18**, 27–32 (2003).

29. Y. L. Serrano Negron, N. F. Hansen, S. T. Harbison, The sleep inbred panel, a collection of inbred *Drosophila melanogaster* with extreme long and short Sseep duration. *G3* **8**, 2865–2873 (2018).

30. L. Lungato, M. L. Gazarini, E. J. Paredes-Gamero, S. Tufik, V. D'Almeida, Paradoxical sleep deprivation impairs mouse survival after infection with malaria parasites. *Malar. J.* **14**, 183 (2015).

31. M. R. Opp, Sleeping to fuel the immune system: mammalian sleep and resistance to parasites. *BMC Evol. Biol.* **9**, 8 (2009).

32. C. Dubowy, A. Sehgal, Circadian rhythms and sleep in *Drosophila melanogaster*. *Genetics* **205**, 1373–1397 (2017).

33. O. T. Shafer, A. C. Keene, The regulation of *Drosophila* sleep. *Curr. Biol.* **31**, R38–R49 (2021).

34. P. Christe, H. Richner, A. Oppliger, Of great tits and fleas: sleep baby sleep. *Anim. Behav.* **52**, 1087–1092 (1996).

35. M. S. Giorgi, R. Arlettaz, P. Christe, P. Vogel, The energetic grooming costs imposed by a parasitic mite (*Spinturnix myoti*) upon its bat host (*Myotis myotis*). *Proc. Biol. Sci.* **268**, 2071–2075 (2001).

36. G. Hausfater, B. Jean Meade, Alternation of sleeping groves by yellow baboons (*Papio cynocephalus*) as a strategy for parasite avoidance. *Primates* **23**, 287–297 (1982).

37. C. J. Horn, J. A. Wasylenko, L. T. Luong, Scared of the dark? Phototaxis as behavioural immunity in a host–parasite system. *Biol. Lett.* **18**, 20210531 (2022).

38. M. Polak, Ectoparasitic effects on host survival and reproduction: The *Drosophila--Macrocheles* association. *Ecology* **77**, 1379–1389 (1996).

39. K. D. Lafferty, A. M. Kuris, Trophic strategies, animal diversity and body size. *Trends Ecol. Evol.* **17**, 507–513 (2002).

40. T. R. Raffel, L. B. Martin, J. R. Rohr, Parasites as predators: unifying natural enemy ecology. *Trends Ecol. Evol.* **23**, 610–618 (2008).

41. A. Acerbi, C. L. Nunn, Predation and the phasing of sleep: an evolutionary individual-based model. *Anim. Behav.* **81**, 801–811 (2011).

42. R. K. Tisdale, J. A. Lesku, G. J. L. Beckers, A. L. Vyssotski, N. C. Rattenborg, The low-down on sleeping down low: pigeons shift to lighter forms of sleep when sleeping near the ground. *J. Exp. Biol.* **221** (2018).

43. S. L. Lima, N. C. Rattenborg, J. A. Lesku, C. J. Amlaner, Sleeping under the risk of predation. *Anim. Behav.* **70**, 723–736 (2005).

44. J. A. Lesku, *et al.*, Predator-induced plasticity in sleep architecture in wild-caught Norway rats (*Rattus norvegicus*). *Behav. Brain Res.* **189**, 298–305 (2008).

45. L. T. Luong, B. D. Heath, M. Polak, Host inbreeding increases susceptibility to ectoparasitism. *J. Evol. Biol.* **20**, 79–86 (2007).

46. L. T. Luong, L. R. Penoni, C. J. Horn, M. Polak, Physical and physiological costs of ectoparasitic mites on host flight endurance. *Ecol. Entomol.* **40**, 518–524 (2015).

47. M. J. Scott, *et al.*, Genomic analyses of a livestock pest, the New World screwworm, find potential targets for genetic control programs. *Commun Biol* **3**, 424 (2020).

48. R. W. Hagan, *et al.*, Dehydration prompts increased activity and blood feeding by mosquitoes. *Sci. Rep.* **8**, 6804 (2018).

49. S. Andrews, Others, FastQC: a quality control tool for high throughput sequence data (2010).

50. A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

51. J. Thurmond, *et al.*, FlyBase 2.0: the next generation. *Nucleic Acids Res.* **47**, D759–D765 (2019).

52. N. L. Bray, H. Pimentel, P. Melsted, L. Pachter, Near-optimal probabilistic RNA-seq quantification. *Nat. Biotechnol.* **34**, 525–527 (2016).

53. C. Soneson, M. I. Love, M. D. Robinson, Differential analyses for RNA-seq: transcript-level estimates improve gene-level inferences. F1000Res 4: 1521 (2015).

54. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

55. J. Reimand, R. Kolde, T. Arak, gProfileR: Interface to the'g: Profiler'Toolkit. *R package version 0. 6* **7** (2018).

56. F. Supek, M. Bošnjak, N. Škunca, T. Šmuc, REVIGO summarizes and visualizes long lists of gene ontology terms. *PLoS One* **6**, e21800 (2011).

57. A. J. Rosendale, M. E. Dunlevy, M. D. McCue, J. B. Benoit, Progressive behavioural, physiological and transcriptomic shifts over the course of prolonged starvation in ticks. *Mol. Ecol.* **28**, 49–65 (2019).

58. A. M. Fieler, *et al.*, Larval thermal characteristics of multiple ixodid ticks. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* **257**, 110939 (2021).

59. Q. Geissmann, L. Garcia Rodriguez, E. J. Beckwith, G. F. Gilestro, Rethomics: An R framework to analyse high-throughput behavioural data. *PLoS One* **14**, e0209331

(2019).

60. R. Core Team, Others, R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing. *Available* (2013).

# Figure 1

**A. Resistance levels**

Proportion flies uninfected

0.700
0.600
0.500
0.400
0.300
0.200

\*    \*    \*

C  S    C  S    C  S
Line 1   Line 2   Line 3

**B. S vs. C (Nested analysis)**

oxidoreductase activity

heme binding

monooxygenase activity

iron ion binding

Figure 2

# Figure 3

# Figure 4