Precision-Cut Liver Slices as an *ex vivo* model to evaluate antifibrotic therapies for liver fibrosis and cirrhosis

Yongtao Wang[1,2,*], Ben Leaker[3,4,*], Guoliang Qiao[1,*], Mozhdeh Sojoodi[1], Ibrahim Ragab Eissa[1], Eliana T. Epstein[2], Jonathan Eddy[2], Oizoshimoshiofu Dimowo[2], Alan C. Mullen[2,5,6], Georg M. Lauer[2], Raymond T. Chung[2], Motaz Qadan[1], Michael Lanuti[7], Bryan C. Fuchs[1], Kenneth K. Tanabe[1]

[1]Division of Gastrointestinal and Oncologic Surgery, Massachusetts General Hospital and Harvard Medical School, Boston, MA, United States

[2]Liver Center, Division of Gastroenterology, Massachusetts General Hospital and Harvard Medical School, Boston, MA, United States

[3]Wellman Center for Photomedicine, Massachusetts General Hospital and Harvard Medical School, Boston, MA, United States

[4]Harvard-MIT program in Health Sciences and Technology, Massachusetts Institute of Technology, Boston, MA, United States

[5]Division of Gastroenterology, University of Massachusetts Chan Medical School, Worcester, MA, United States

[6]Broad Institute, Cambridge, MA, United States

[7]Division of Thoracic Surgery, Massachusetts General Hospital and Harvard Medical School, Boston, MA, United States

*Equal contribution

Correspondence:
Kenneth K. Tanabe
Division of Gastrointestinal and Oncologic Surgery
Massachusetts General Hospital
55 Fruit Street, Boston, MA 02114
Phone: 617-724-3868
ktanabe@partners.org

Abstract

Background: Precision-Cut Liver Slices (PCLS) are an *ex vivo* culture model developed to study hepatic drug metabolism. One of the main benefits of this model is that it retains the structure and cellular composition of the native liver. PCLS also represents a potential model system to study liver fibrosis in a setting that more closely approximates *in vivo* pathology than *in vitro* methods. The aim of this study was to assess whether responses to antifibrotic interventions can be detected and quantified with PCLS.

Methods: PCLS of 250 μm thickness were prepared from four different murine fibrotic liver models: choline-deficient, L-amino acid-defined, high-fat diet (CDAHFD), thioacetamide (TAA), diethylnitrosamine (DEN), and carbon tetrachloride (CCl_4). PCLS were treated with 5 μM Erlotinib for 72 hours. Histology and gene expression were then compared with *in vivo* murine experiments and TGF-β1 activated hepatic stellate cells (HSC). These types of PCLS characterization were also evaluated in PCLS from human cirrhotic liver.

Results: PCLS viability in culture was stable for 72 hours. Erlotinib treatment significantly inhibited the expression of profibrogenic genes *Il6*, *Col1a1* and *Timp1* in PCLS from CDAHFD-induced cirrhotic mice, and *Il6*, *Col1a1* and *Tgf-β* in PCLS from TAA-induced cirrhotic rats. Erlotinib treatment of PCLS from DEN-induced cirrhotic rats inhibited the expression of *Col1a1*, Timp1, *Tgf-β* and *Tnf-α*, which was consistent with the impact of erlotinib on *Col1a1* and *Tgf-β* expression in *in vivo* DEN-induced cirrhosis. Erlotinib treatment of PCLS from CCl_4-induced cirrhosis caused reduced expression of *Timp1*, *Col1a1* and *Tgf-β*, which was consistent with the effect of erlotinib in *in vivo* CCl_4-induced cirrhosis. In addition, in stellate cells at PCLS from normal mice, TGF-β1 treatment upregulated *Acta2* (*αSMA*), while treatment with erlotinib inhibited the expression of *Acta2*. Similar expression results were observed in TGF-β1 treated HSC cell lines. Expression of MMPs and TIMPs, key regulators of fibrosis progression and regression, were also significantly altered under erlotinib treatment in PCLS. Expression changes under erlotinib treatment were also corroborated with PCLS form human cirrhosis samples.

Conclusion: The responses to antifibrotic interventions can be detected and quantified with PCLS at the gene expression level. PCLS from murine cirrhotic livers after treatment with an antifibrotic

drug are consistent with those observed *in vivo* and *in vitro*. These PCLS characterizations were also observed in PCLS from human cirrhosis. PCLS is an excellent model to assess antifibrotic therapies that is aligned with the principles of Replacement, Reduction and Refinement (3Rs).

Introduction

Cirrhosis is the final stage of liver fibrosis caused by many forms of liver injury, such as chronic hepatitis B and C, chronic alcohol use disorder, and metabolic dysfunction-associated steatohepatitis (MASH, previously known as non-alcoholic steatohepatitis and NASH).[1, 2] In 2015, cirrhosis affected 2.8 million people and resulted in 1.3 million deaths globally.[3, 4] Cirrhosis also contributes to the vast majority of hepatocellular carcinoma (HCC) cases, which is the 6th most common cancer worldwide, and the 2nd leading cause of cancer-related death.[5] Considering the lack of successful treatment options and poor prognosis for cirrhosis,[6] new strategies and platforms to investigate antifibrotic therapies are urgently needed.

Many of the limitations of cellular and animal models of liver fibrosis can be overcome by using the relatively recently proposed precision cut liver slice (PCLS) model. This technique maintains thin slices of liver tissue in culture for several days, enabling drugs to be tested in an *ex vivo* setting that retains the architecture and cell populations of the liver. Compared to two-dimensional *in vitro* cell culture, PCLS better model the interactions and signal transduction pathways between different cell types.[7, 8] The PCLS model also offers the benefit of reduced animal experiments and time compared to *in vivo* models, and is in line with the principles of Replacement, Reduction and Refinement.[9]

PCLS has been utilized to study various liver conditions, such as normal liver function,[10] drug metabolism,[11] drug-induced liver injury and toxicity,[12-14] fatty liver,[15] early stages of liver fibrosis,[8, 15-20] and immunological responses.[21] A study using PCLS derived from rats subjected to bile duct ligation supported PCLS`s ability for molecular antifibrotic evaluation with transcriptomic characterization.[22] However, PCLS as a platform to evaluate antifibrotic therapies for cirrhosis, a more severe stage of liver disease, has not been reported.

Previous experiments from our lab have demonstrated the efficacy of a small-molecule EGF receptor (EGFR) inhibitor erlotinib as an antifibrotic agent in murine models of liver fibrosis.[23] Erlotinib inhibited the activation of hepatic stellate cells (HSCs), stopped the progression of cirrhosis, and prevented subsequent development of HCC. Here, erlotinib was used as an example antifibrotic therapy in PCLS. The aim of this study was to assess whether responses to antifibrotic interventions can be detected and quantified with PCLS.

Methods

Chemicals

Stock solution of 50 mM erlotinib (Tarceva[TM], Genentech) was prepared in DMSO.

Animal models for PCLS

All animal experiments were approved by the MGH Institutional Animal Care and Use Committee (IACUC). PCLS were prepared from four established murine models of cirrhosis: choline-deficient, L-amino acid-defined, high-fat diet (CDAHFD), thioacetamide (TAA), diethylnitrosamine (DEN), and carbon tetrachloride (CCl_4).

DEN-induced cirrhosis rat model

The DEN model of cirrhosis was prepared as in our previous publications.[23] Male Wistar rats received weekly intraperitoneal (ip) injections of DEN (Sigma) at 50 mg/kg for 18 weeks.

CCl_4-induced cirrhosis mouse model

The CCl_4 model of cirrhosis was prepared as in our previous publication.[23] Male C57Bl/6 mice were treated three times a week for 18 weeks with 0.1 ml of 40% CCl_4 (Sigma) in olive oil by oral gavage.

CDAHFD-induced cirrhosis mouse model

The CDAHFD model of cirrhosis was induced as in our previous publication.[24] Male C57Bl/6 mice received CDAHFD (L-amino acid diet with 60 kcal% fat with 0.1% methionine and no added choline. Research Diets Inc. A06071302) for 16 weeks.

TAA-induced cirrhosis rat model

The TAA model of cirrhosis was induced as in our previous publication.[25] Male Wistar rats received ip injections with 200 mg/kg TAA (Sigma) twice a week for 12 weeks.

PCLS as an *ex vivo* model of liver biology

PCLS from healthy or cirrhotic livers were performed as published.[7, 26, 27] Tissue samples were glued to the mounting stage of a 7000smz-2 vibratome instrument (Campden Instruments

Limited) and submersed in sterile Krebs Henseleit buffer (Sigma-Aldrich). Using 7550-1-C ceramic blades (Campden Instruments Limited), the tissue was cut into 250 μm slices at an advance speed of 0.1 mm/s, with 2.5 mm amplitude and 50Hz frequency. Slices from human and rat livers were trimmed to a uniform size with an 8 mm biopsy punch (Acuderm Inc.), and slices from mouse livers were trimmed with a 6 mm biopsy punch. PCLS were transferred into 8 μm-pore Transwell inserts and cultured in standard 6-well plates (Corning), with William's E medium (Sigma-Aldrich) containing 2.0 g/L glucose, 10% fetal bovine serum (FBS, Gibco), 2 mM L-glutamine supplement (Gibco), 100 U/mL penicillin and 100 μg/mL streptomycin (Lonza), at 37°C in a humidified atmosphere of 5% CO_2.

PCLS from cirrhotic animals were then treated with 5 μM erlotinib or vehicle control for 72 hours. PCLS from normal murine livers were treated with 10 ng/mL TGF-β1 for 24 hours, then treated with 10ng/mL TGF-β and 5 μM erlotinib for an additional 72 hours.

In vivo erlotinib experiments
DEN-induced cirrhosis rat model

As previously published,[23] male Wistar rats received weekly ip injections of 50 mg/kg DEN for 18 weeks. Starting from week 8, rats received daily ip injections with either 2 mg/kg erlotinib or vehicle control (n = 8 per group). All rats were sacrificed at week 19 after a one-week washout period to eliminate the acute effects of DEN. The livers from this previous *in vivo* experiment were used in this current study.

CCl4-induced cirrhosis mouse model

As previously published,[23] male A/J mice were treated three times a week for 18 weeks with 0.1 ml of 40% CCl4 in olive oil by oral gavage. Mice received daily ip injections of control or 2 mg/kg erlotinib from week 13 to 19 (n = 8 per group). Mice were sacrificed at week 19 after a one-week washout to eliminate acute the effects of CCl4. The livers from this previous *in vivo* experiment were used in this current study.

Cell culture

Cell culture was performed as described previously.[28, 29] Human HSC, LX2 and TWNT4 cells were purchased from American Type Culture Collection (ATCC). Cells were grown in

Dulbecco's modified Eagle's medium (DMEM, Cellgro) or Roswell Park Memorial Institute (RPMI) 1640 medium (HyClone) containing 15% FBS (Gibco), 1 mM sodium pyruvate (Gibco), 100 U/mL penicillin sodium and 100 µg/mL streptomycin sulfate (Lonza), at 37°C in a humidified atmosphere of 5% CO_2.

MTS

MTS assay was performed as described previously.[21] Individual PCLS were placed in a 24-well plate with 400 µL William's E medium and 80 µL MTS solution (Abcam) for each well. Plates were incubated at 37°C in standard culture conditions for 1 hour, then mixed briefly on a shaker. Supernatants were transferred to a 96-well plate and absorbance was measured using a plate reader (Molecular Devices) at OD = 490 nm.

Histological Hematoxylin & Eosin (H&E) and Sirius red staining

Initially, PCLS were fixed in formalin for 3 days at room temperature then paraffin-embedded and sectioned into 7 µm slices for H&E and Sirius red staining with standard protocols.[30] However, we found that PCLS from cirrhotic livers were fragile and easily degraded during sectioning and staining (**Fig. S3A**, left panel). The standard protocol was then modified. PCLS were fixed in formalin at 4°C overnight, embedded in paraffin, and then sectioned into 10 µm slices. The time for all steps were also reduced to half of the standard time. This modified protocol resulted in higher quality stains and retained the structure of the PCLS (**Fig. S3A**, right panel). Amount of collagen in Sirius red stained sections was scored according to the modified Ishak method[31] as described in **Supplementary Table 1**. Collagen proportionate area was morphometrically quantified with image processing software (ImageJ, NIH).

Quantitative RT-PCR

Quantitative RT-PCR was performed as described previously.[32, 33] Total RNA was isolated from rat liver tissues using TRIzol (Invitrogen) and subsequently treated with DNase I (Promega) according to the manufacturer's instructions. 1 µg of total RNA from each sample was used to synthesize cDNA (SuperScript III First-Strand Synthesis SuperMix for qRT-PCR, Invitrogen). Quantitative real-time PCR was performed using the 7900HT Fast Real-Time PCR System

(Thermo fisher) with commercial TaqMan primers (Thermo fisher) as shown in **Supplementary Table 2**.

Human samples

All human samples were harvested in accordance with the protocols approved by the Mass General Brigham Institutional Review Boards (IRBs).

Statistical analysis

All values were expressed as mean ± S.E.M.. Two-tailed Student's t tests were performed to compare data between control and one experimental group, and One-way ANOVA followed by post-hoc Tukey tests with 2-tailed distribution were performed to analyze data among groups of 3 or more. Graphs were prepared with GraphPad Prism v6.0c software. All experiments were independently repeated three times. Significance is represented by $*p < 0.05$, $**p < 0.01$, $***p < 0.001$ versus control.

Results

Stable viability of PCLS

PCLS with 250 μm thickness were prepared and further cultured with treatment (**Fig. 1A, S1**). To ensure minimal tissue degradation *ex vivo,* the period of stable viability was determined. MTS assays showed the viability of PCLS from mouse CDAHFD-induced, rat DEN-induced, rat TAA-induced, and mouse CCl_4-induced established cirrhosis were not significantly changed after 72 hours in culture (**Fig. 1B-1E, S2A, S2B**). This timepoint was used for the remaining experiments with cirrhotic PCLS. The DEN model was also used to investigate the lifespan of cirrhotic PCLS. The nominal decrease in viability after 5 days was not statistically significant (**Fig. 1C**). The CDAHFD and DEN models were further used to determine whether erlotinib had an impact on PCLS viability. There was no difference between vehicle and erlotinib treated slices with either model at 72 hours (**Fig. S2C, S2D**).

Antifibrotic treatment of PCLS from CDAHFD-induced mouse cirrhosis and TAA-induced rat cirrhosis

To first assess whether responses to antifibrotic interventions can be detected and quantified with PCLS, PCLS from CDAHFD-induced and TAA-induced cirrhosis were utilized. CDAHFD-induced cirrhosis[30] was confirmed with Sirius red staining (**Fig. S4A**), and PCLS from this CDAHFD-induced cirrhosis were prepared. Erlotinib treatment of PCLS slices for 72 hours significantly suppressed the expression of the profibrogenic genes *Il6*, *Col1a1* and *Timp1* (**Fig. 2A**), and suppressed *Acta2* expression with marginal significance ($p = 0.0777$). No significant effect was observed on the expression of *Tgf-β*. This short-term exposure of PCLS slices to erlotinib did not significantly reduce the amount of collagen measured with Sirius red staining (**Fig. 2B, 2C**). No change was evident on H&E stained morphology (**Fig. S5A**).

Cirrhosis was induced by TAA[34] and confirmed with Sirius staining (**Fig. S4B**). PCLS were prepared from these TAA-induced cirrhotic livers. Erlotinib treatment of PCLS slices for 72 hours significantly inhibited the expression of the profibrogenic genes *Il6*, *Tgf-β* and *Col1a1* in PCLS (**Fig. 2D**). No significant change in expression of *Timp1* and *Acta2* was observed. This short-term exposure to erlotinib did not significantly reduce the amount of collagen (**Fig. 2E, 2F**). No change was evident on H&E staining (**Fig. S5B**).

Taken together with the CDAHFD and TAA data above, acute responses to antifibrotic interventions can be detected and quantified with PCLS on the molecular gene expression level.

Antifibrotic treatment in PCLS from DEN-induced rat cirrhosis compared to *in vivo* DEN cirrhosis model

DEN-induced cirrhosis[23, 35, 36, 37, 38] was confirmed with Sirius staining (**Fig. S4C**), and PCLS from this DEN-induced cirrhosis were then prepared. Exposure of PCLS slices to erlotinib for 72 hours significantly suppressed the expression of the profibrogenic genes *Col1a1*, *Tgf-β*, *Il6* and *Timp1* (**Fig. 3A**). No significant difference in the amount of collagen (**Fig. 3B, 3C**) and H&E staining (**Fig. S5C**) were observed after erlotinib treatment of PCLS slices for 72 hours.

To compare the effect of erlotinib between *ex vivo* cirrhotic PCLS and an *in vivo* cirrhosis model, we analyzed livers obtained from DEN-induced cirrhotic rats that had been treated with erlotinib for 10 weeks (**Fig. S5E**). As observed in the short-term PCLS model, longer-term treatment rats with DEN-induced cirrhosis resulted in significant reduction in *Col1a1* and *Acta2* (**Fig. 3D**), and marginally reduced the expression of *Tgf-β* ($p = 0.0624$). No effect on the expression of *Il6* and *Timp1* was observed. As previously published, 10 weeks of erlotinib treatment in these rats also significantly reduced the amount of collagen (**Fig. 3E, 3F**).

Antifibrotic treatment in PCLS from CCl₄-induced mouse cirrhosis compared to *in vivo* CCl₄ cirrhosis model

CCl$_4$-induced cirrhosis[23] was confirmed with Sirius red staining (**Fig. S4D**) and PCLS from this CCl$_4$-induced cirrhosis were then prepared. Short-term exposure of PCLS slicers to erlotinib significantly suppressed the expression of the profibrogenic genes *Timp1*, *Col1a1* and *Tgf-β* (**Fig. 4A**), but did not affect the expression of *Il6* and *Acta2*. No significant difference in the amount of collagen (**Fig. 4B, 4C**) and H&E staining (**Fig. S5D**) were observed after erlotinib treatment of the PCLS for 72 hours.

To compare the effect of erlotinib between *ex vivo* cirrhotic PCLS and an *in vivo* cirrhosis model, we analyzed liver obtained from CCl$_4$-induced cirrhotic mice that had been treated with erlotinib for 6 weeks (**Fig. S5F**). Six weeks of erlotinib treatment in these mice significantly suppressed the expression of *Timp1*, *Col1a1*, *Tgf-β*, *Il6* and *Acta2* (**Fig. 4D**), and reduced the amount of collagen (**Fig. 4E, 4F**).

Taken together with the DEN and CCl₄ data above, these results demonstrate that erlotinib induces similar acute effects in PCLS that are observed *in vivo* in these two murine models of cirrhosis.

Antifibrotic treatment in stellate cells from PCLS compared to *in vitro* hepatic stellate cells

In PCLS from normal rats (**Fig. S5G**), stellate cells which expressing *Acta2* were only a small minority of cells. TGF-β1 treatment for 72 hours induced the expression of *Acta2* in stellate cells at PCLS, while treatment with erlotinib significantly inhibited the expression of *Acta2* (**Fig. 5A**). In stellate cell lines LX2 (**Fig. 5B**) and TWNT4 (**Fig. 5C**), TGF-β1 significantly induced the expression of *ACTA2*, while treatment with erlotinib for 72 hours suppressed *ACTA2* expression. Erlotinib treatment for 72 hours also reduced Tgf-β1-induced collagen deposition (**Fig. 5D**) on the morphological level. These results demonstrate that erlotinib induces similar effects in stellate cells at PCLS and *in vitro* HSC cell lines.

Besides, TGF-β1 treatment induced the expression of other profibrogenic genes *Il6*, *Tgf-β Timp1* and *Col1a1* in PCLS, while treatment with erlotinib significantly inhibited the expression of these genes (**Fig. 5E**). In stellate cell lines LX2 (**Fig. 5F**) and TWNT4 (**Fig. 5G**), TGF-β1 significantly induced the expression of *COL1A1*, while treatment with erlotinib suppressed *COL1A* expression. However, *TGF-β* and *IL6* did not respond to erlotinib treatment, and *TIMP1* did not respond to TGF-β1 and erlotinib treatment on these cells.

Effect of antifibrotic therapy on MMPs and TIMPs expression in cirrhotic PCLS

To explore other potential mechanisms of antifibrotic therapy of erlotinib in cirrhosis PCLS, we assessed expression of other regulatory pathways for fibrosis generation and degradation. In livers from CDAHFD-induced cirrhotic mice, exposure of PCLS slices to erlotinib for 72 hours significantly increased the expression of *Mmp2*, *Mmp3* and *Mmp8* (**Fig. 6A**), indicating activation of antifibrotic pathways. Erlotinib treatment also significantly decreased *Mmp9*, *Mmp13* (**Fig. 6B**), and *Timp1* (**Fig. 6C**) expression, reflecting suppression of profibrotic pathways. Decreased *Timp1* expression was also observed in PCLS from CDAHFD (**Fig. 2A**), DEN (**Fig. 3A**), and CCl₄-induced (**Fig. 4A**) cirrhotic livers.

Antifibrotic evaluation in human cirrhotic PCLS

The PCLS from diet-induced rodent cirrhotic model and cirrhotic patients were utilized to assess whether the types of PCLS characterizations in rodent cirrhosis PCLS were also observed in PCLS from human cirrhosis. As described above, PCLS from CDAHFD-induced murine cirrhosis was used to assess the effects of 72 hours exposure to erlotinib (**Fig. 2A-C**). We next tested the effects of 72 hours of erlotinib on PCLS from human liver cirrhosis (**Fig. S6**). Erlotinib treatment of PCLS did not change the liver morphology (**Fig. S5H**), but significantly inhibited the expression of the profibrogenic genes *COL1A1*, *TIMP1*, *IL6* and *TNF-α* (**Fig. 7A**). No significant effect was observed on the expression of *Tgf-β* and *Acta2*. No significant difference in collagen accumulation was observed after erlotinib treatment (**Fig. 7B, 7C**). Analysis using PCLS demonstrated that the effects of erlotinib as an antifibrotic therapy were similar in murine and human cirrhosis samples.

Discussion

New strategies to investigate antifibrotic therapies for cirrhosis are urgently needed. Compared with two-dimensional *in vitro* cell lines, PCLS are a better representation of the *in vivo* environment because they retain the structure and cellular composition.[7, 8] It therefore has the capacity to better capture the complex multi-cellular pathways involved in liver injury and the progression of cirrhosis. Compared to *in vivo* models, PCLS offer faster readouts, reduced animal numbers, and the ability to run multiple tests on highly similar samples by using serial sections.

In this study, we demonstrated that PCLS is a reliable *ex vivo* model to evaluate antifibrotic therapies across four established murine models of (CDAHFD, TAA, DEN, and CCl4-induced) cirrhosis, using erlotinib as an example drug. Expression analysis of PCLS after erlotinib treatment showed suppression in a variety of profibrogenic genes. We demonstrated the responses to antifibrotic interventions can be detected and quantified with PCLS at the molecular level. As expected, morphology of PCLS after erlotinib treatment did not significantly change. This is likely because the short time in culture (72 hours) does not provide enough time for significant remodeling to occur.[7] Our previous i*n vivo* experiments showing reduced collagen accumulation in DEN-induced cirrhosis in rats or CCl4-induced cirrhosis in mice used erlotinib treatment for longer-term weeks.[23] Our current study also confirmed that the viability of PCLS from cirrhotic liver stabilized during 72 hours in culture.

We directly compared PCLS with *in vivo* experiments using the same method of inducing cirrhosis, and showed that many of the acute responses to antifibrotic therapies in PCLS were consistent with *in vivo* results. Erlotinib treatment of PCLS from DEN-induced cirrhotic rats suppressed the expression of partial profibrogenic genes, which was consistent with the impact of erlotinib on these genes *in vivo*. Similar results were demonstrated with the CCl4 mouse model.

Stellate cells expressing *Acta2* are a small minority population within PCLS. We also compared the stellate cells at PCLS to common *in intro* HSC models. TGF-β1 activated LX2 and TWNT4 cell lines are well established models to evaluate antifibrotic therapies *in vitro*.[36, 39] Here we showed erlotinib treatment significantly suppressed expression of the TGF-β1 induced profibrogenic gene *Acta2* in stellate cells on normal liver PCLS. Erlotinib also significantly inhibited the expression of TGF-β1 activated *ACTA2* in LX2 and TWNT4 cells. As other profibrogenic genes *Il6*, *Tgf-β*, *Timp1* and *Col1a1* are secreted by hepatocytes, HSCs and other types of liver cells,[40-43] the decreased expression of these genes in PCLS after erlotinib treatment

might come from all these liver cells. However, erlotinib only significantly inhibited the expression of TGF-β1 activated *COL1A1* in LX2 and TWNT4 cells. *IL6*, *TIMP1* and *TGF-β* were not reduced after erlotinib treatment, which might also indicate a lower sensitivity for cell lines to respond to certain antifibrotic therapies.

We then investigated other potential mechanisms contributing to these molecular observations of fibrogenesis in PCLS. Matrix metalloproteinases (MMPs) are a family of enzymes that regulate the degradation of extracellular matrix proteins. MMP13 induces inflammation, cytokines and activation of myofibroblast to acquire the profibrotic effect. On the contrary, MMP2 and MMP8 induce clearance of collagen, and MMP3 inhibits activation of myofibroblast to foster the anti-fibrosis effect. Besides, lack of MMP9 results in reduced fibrosis. Tissue inhibitors of metalloproteinases (TIMPs) inhibit the MMPs.[44] Regulatory pathways on the fibrosis generation and degradation[22, 44] were significantly altered to exert the antifibrotic therapy of erlotinib in PCLS. After erlotinib treatment, antifibrotic pathways including MMP2, MMP3 and MMP8 were enhanced in PCLS from CDAHFD-induced cirrhosis, while profibrotic pathways including MMP9 and MMP13 were hampered. Besides, TIMP1 as the inhibitor of MMPs, was reduced after erlotinib treatment.

We then confirmed the antifibrotic effects of erlotinib using PCLS from human cirrhosis samples. The type of PCLS characterizations in rodent cirrhosis PCLS were also observed in PCLS from human cirrhosis, which addressed the prospect to further assess the effect of erlotinib in *in vivo* human cirrhotic liver. The majority of PCLS research to date has focused on murine samples[8, 10, 11, 13-19] and only a small fraction of studies used human tissue.[12, 20] PCLS from human samples may better predict the *in vivo* response to an antifibrotic therapy in humans, and also opens up the possibility of using a liver biopsy sample to test an individual patient's response to a variety of drugs. Human PCLS may also be used as an experimental platform to expedite basic research and drug development, by directly testing the effects of drugs in clinical cirrhosis samples rather than animal models.

In summary, the responses to antifibrotic interventions can be detected and quantified with PCLS on the molecular level. PCLS accurately capture the changes in expression that occur *in vitro* and *in vivo* during treatment with an antifibrotic therapy. These types of PCLS characterizations were also observed in PCLS from human cirrhosis. PCLS reduces animal numbers, in alignment with the principles of Replacement, Reduction and Refinement[9] by

enabling many slices to collected from one animal. PCLS is a promising platform for the future development of antifibrotic therapies for cirrhosis.

Acknowledgements

This study has been supported by R01DK104956, NIH.

Author contributions

KKT and YW designed the experiments. YW, BL and GQ conducted the experiments and performed data analysis. YW wrote the manuscript, and KKT and BL improved the manuscript. MS, IRE, KK, SCB, ETE, JE, ACM, GML, RTC, MQ, ML, BCF and RTC contributed to new reagents or analytical tools.

Conflict of interest

The authors declare that they have no competing interest.

Legends

Figure 1. Evaluation of viability in PCLS.

(A) Workflow for PCLS preparation and treatment. Viability analyzed with MTS assays for PCLS from (B) CDAHFD, (C) DEN, (D) CCl_4 and (E) TAA-induced cirrhosis rats. Experiments were independently repeated three times (n = 3). All values were expressed as the mean ± S.E.M with two-tailed Student's *t* tests or one-way ANOVA analysis.

Figure 2. Antifibrotic evaluation in PCLS from CDAHFD-induced mouse cirrhosis and TAA-induced rat cirrhosis.

(A) Quantitative RT-PCR analysis of profibrogenic genes. Amount of collagen (B) measured with Sirius red staining and (C) quantified with collagen proportionate area analysis in PCLS from CDAHFD-induced mouse cirrhosis after 5 μM erlotinib treatment for 72 hours. (D) Quantitative RT-PCR analysis of profibrogenic genes. Amount of collagen (E) measured with Sirius red staining and (F) quantified in PCLS from TAA-induced rat cirrhosis after 5 μM erlotinib treatment for 72 hours. Experiments were independently repeated three times (n = 4). All values were expressed as the mean ± S.E.M with two-tailed Student's *t* tests. Significance is represented by *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

Figure 3. Comparison of antifibrotic therapy in *ex vivo* PCLS from DEN-induced *rat* cirrhosis with *in vivo* DEN cirrhotic model.

For *ex vivo* PCLS analysis, (A) quantitative RT-PCR analysis of profibrogenic genes. Amount of collagen (B) measured with Sirius red staining and (C) quantified with collagen proportionate area analysis in PCLS after 5 μM erlotinib treatment for 72 hours. Experiments were independently repeated three times (n = 4). For *in vivo* analysis, (D) quantitative RT-PCR analysis of profibrogenic genes. Amount of collagen (E) measured with Sirius red staining and (F) quantified after daily ip injections of 2 mg/kg erlotinib for 10 weeks. All values were expressed as the mean ± S.E.M with two-tailed Student's *t* tests. Significance is represented by *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

Figure 4. Comparison of antifibrotic therapy in *ex vivo* PCLS from CCl_4-induced mouse cirrhosis with *in vivo* CCl_4 cirrhotic model.

For *ex vivo* PCLS analysis, (A) quantitative RT-PCR analysis of profibrogenic genes. Amount of collagen (B) measured with Sirius red staining and (C) quantified with collagen proportionate area analysis in PCLS after 5 µM erlotinib treatment for 72 hours. Experiments were independently repeated three times (n = 4). For *in vivo* analysis, (D) quantitative RT-PCR analysis of profibrogenic genes. Amount of collagen (E) measured with Sirius red staining and (F) quantified after daily ip injections of 2 mg/kg erlotinib for 6 weeks. All values were expressed as the mean ± S.E.M with two-tailed Student's *t* tests. Significance is represented by *$p < 0.05$, ***$p < 0.001$.

Figure 5. Comparison of antifibrotic evaluation in *ex vivo* PCLS with *in vitro* hepatic stellate cells.

Quantitative RT-PCR analysis of *Acta2* mRNA from (A) PCLS from normal rats, (B) LX2 and (C) TWNT4 cells. PCLS or HSC cells were activated with 10 ng/mL TGF-β1 for 24 hours, and further treated with 5 µM erlotinib for 72 hours. (D) Amount of collagen measured with Sirius red staining and quantified with collagen proportionate area analysis in PCLS from normal rats. Quantitative RT-PCR analysis of *Il6*, *Tgf-β*, *Timp1* and *Col1a1* mRNA from (E) PCLS from normal rats, (F) LX2 and (G) TWNT4 cells. These experiments were repeated three times (n = 4). All values were expressed as the mean ± S.E.M with one-way ANOVA analysis. Significance is represented by *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

Figure 6. Regulatory pathways of the antifibrotic evaluation of erlotinib in cirrhotic PCLS.

Quantitative RT-PCR analysis of the mRNA expression on (A) antifibrotic pathways, (B) profibrotic pathways, and (C) inhibitory pathways of MMPs in PCLS from CDAHFD-induced cirrhosis mice after erlotinib treatment for 72 hours. These experiments were repeated three times (n = 4). All values were expressed as the mean ± S.E.M with two-tailed Student's *t* tests. Significance is represented by *$p < 0.05$, ***$p < 0.001$.

Figure 7. The Antifibrotic evaluation in human cirrhotic PCLS.

(A) Quantitative RT-PCR analysis of mRNA expression of profibrogenic genes. Amount of collagen (B) measured with Sirius red staining and (C) quantified in PCLS from human cirrhosis treated with 5 µM erlotinib for 72 hours. These experiments were repeated three times (n = 4). All

values were expressed as the mean ± S.E.M with two-tailed Student's *t* tests. Significance is represented by **$p < 0.01$.

Legends for Supplementary Figures

Supplementary Figure 1. Schedule of PCLS preparation and erlotinib treatment.

Workflow of PCLS preparation and erlotinib treatment for (A) cirrhotic and (B) normal livers. (C) Summary table of PCLS preparation and erlotinib treatment.

Supplementary Figure 2. Viability evaluation in PCLS

(A) Representative images of PCLS from CDAHFD-induced cirrhosis rats on MTS assays, the cells alive were stained with purple color. (B) Representative images of PCLS from DEN-induced cirrhosis rats on MTS viability assays. MTS assays in PCLS from (C) CDAHFD and (D) DEN-induced rat cirrhosis after erlotinib treatment. These experiments were repeated three times (n = 3). All values were expressed as the mean ± S.E.M with two-tailed Student's *t* tests.

Supplementary Figure 3. Improvement of PCLS processing.

(A) Comparison of standard and modified protocols to proceed PCLS from TAA-induced rat cirrhosis.

Supplementary Figure 4. Confirmation of established murine cirrhosis.

Confirmation of established (A) CDAHFD, (B) TAA, (C) DEN and (D) CCl_4-induced murine cirrhosis.

Supplementary Figure 5. H&E staining.

Representative images for H&E staining in PCLS from (A) CDAHFD-induced mouse cirrhosis, (B) TAA-induced rat cirrhosis, (C) DEN-induced *rat* cirrhosis, and (D) CCl_4-induced *mouse* cirrhosis. (E) Representative images for H&E staining of PCLS from male healthy C57Bl/6 mice livers. (F) Representative images for H&E staining of human cirrhotic PCLS. Representative images for H&E staining of the samples from (G) *in vivo* DEN-induced rat cirrhotic models and (H) *in vivo* CCl_4-induced mouse cirrhotic models.

Supplementary Figure 6. Confirmation of established human cirrhosis.

Confirmation of human established cirrhosis with (A) H&E and (B) Sirius red staining. (C) H&E and (D) Sirius red staining of PCLS from human established cirrhosis.

Supplementary Tables

Supplementary Table 1. Modified *Ishak* Scoring[31]

Score	Description
0	No fibrosis
1	Fibrous expansion of some or most portal areas, with or without short fibrous septa
2	Fibrous expansion of most portal areas, with occasional portal to portal bridging, or with marked portal to portal bridging as well as portal areas to central bridging
3	Marked bridging with occasional nodules
4	Probable to definite cirrhosis

Supplementary Table 2. Oligonucleotides

Gene Symbol	Species Specificity	Company	Assay ID
18S	Human	Thermo fisher	Hs03003631_g1
IL6	Human	Thermo fisher	Hs00174131_m1
ACTA2	Human	Thermo fisher	Hs00426835_g1
TIMP1	Human	Thermo fisher	Hs01092512_g1
COL1A1	Human	Thermo fisher	Hs0016004_m1
TNF-α	Human	Thermo fisher	Hs00174128_m1
TGF-β	Human	Thermo fisher	Hs00998133_m1
Il6	Rat	Thermo fisher	Rn01410330_m1
Acta2	Rat	Thermo fisher	Rn01759928_g1
Timp1	Rat	Thermo fisher	Rn01430873_g1
Col1a1	Rat	Thermo fisher	Rn01463848_m1
Tgf-β	Rat	Thermo fisher	Rn00572010_m1
Tnf-α	Rat	Thermo fisher	Rn99999017_m1
Il6	Mouse	Thermo fisher	Mm00446190_m1
Acta2	Mouse	Thermo fisher	Mm00725412_s1
Timp1	Mouse	Thermo fisher	Mm01341361_m1

Col1a1	Mouse	Thermo fisher	Mm00801666_g1
Tgf-β	Mouse	Thermo fisher	Mm01178820_m1
Tnf-α	Mouse	Thermo fisher	Mm00443258_m1
Mmp2	Mouse	Thermo fisher	Mm00439498_m1
Mmp3	Mouse	Thermo fisher	Mm00440295_m1
Mmp8	Mouse	Thermo fisher	Mm00439509_m1
Mmp9	Mouse	Thermo fisher	Mm00442991_m1
Mmp13	Mouse	Thermo fisher	Mm00439491_m1
Timp2	Mouse	Thermo fisher	Mm00441825_m1

References

1.   Global, regional, and national age–sex specific all-cause and cause-specific mortality for 240 causes of death, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *The Lancet* **2015,** *385* (9963), 117-171.

2.   Castera, L.; Friedrich-Rust, M.; Loomba, R., Noninvasive Assessment of Liver Disease in Patients With Nonalcoholic Fatty Liver Disease. *Gastroenterology* **2019,** *156* (5), 1264-1281.e4.

3.   Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980-2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* **2016,** *388* (10053), 1459-1544.

4.   Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990-2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* **2016,** *388* (10053), 1545-1602.

5.   McGlynn, K. A.; Petrick, J. L.; London, W. T., Global epidemiology of hepatocellular carcinoma: an emphasis on demographic and regional variability. *Clin Liver Dis* **2015,** *19* (2), 223-38.

6.   Zhang, D. Y.; Friedman, S. L., Fibrosis-dependent mechanisms of hepatocarcinogenesis. *Hepatology* **2012,** *56* (2), 769-75.

7.   Paish, H. L.; Reed, L. H.; Brown, H.; Bryan, M. C.; Govaere, O.; Leslie, J.; Barksby, B. S.; Garcia Macia, M.; Watson, A.; Xu, X.; Zaki, M. Y. W.; Greaves, L.; Whitehall, J.; French, J.; White, S. A.; Manas, D. M.; Robinson, S. M.; Spoletini, G.; Griffiths, C.; Mann, D. A.; Borthwick, L. A.; Drinnan, M. J.; Mann, J.; Oakley, F., A Bioreactor Technology for Modeling Fibrosis in Human and Rodent Precision-Cut Liver Slices. *Hepatology* **2019,** *70* (4), 1377-1391.

8.   van de Bovenkamp, M.; Groothuis, G. M.; Meijer, D. K.; Olinga, P., Precision-cut fibrotic rat liver slices as a new model to test the effects of anti-fibrotic drugs in vitro. *J Hepatol* **2006,** *45* (5), 696-703.

9.   Hubrecht, R. C.; Carter, E., The 3Rs and Humane Experimental Technique: Implementing Change. *Animals (Basel)* **2019,** *9* (10).

10.   Gandolfi, A. J.; Wijeweera, J.; Brendel, K., Use of precision-cut liver slices as an in vitro tool for evaluating liver function. *Toxicol Pathol* **1996,** *24* (1), 58-61.

11.   Graaf, I. A.; Groothuis, G. M.; Olinga, P., Precision-cut tissue slices as a tool to predict metabolism of novel drugs. *Expert Opin Drug Metab Toxicol* **2007,** *3* (6), 879-98.

12.   Hadi, M.; Westra, I. M.; Starokozhko, V.; Dragovic, S.; Merema, M. T.; Groothuis, G. M., Human precision-cut liver slices as an ex vivo model to study idiosyncratic drug-induced liver injury. *Chem Res Toxicol* **2013,** *26* (5), 710-20.

13.   de Graaf, I. A.; Olinga, P.; de Jager, M. H.; Merema, M. T.; de Kanter, R.; van de Kerkhof, E. G.; Groothuis, G. M., Preparation and incubation of precision-cut liver and intestinal slices for application in drug metabolism and toxicity studies. *Nat Protoc* **2010,** *5* (9), 1540-51.

14.   Granitzny, A.; Knebel, J.; Schaudien, D.; Braun, A.; Steinberg, P.; Dasenbrock, C.; Hansen, T., Maintenance of high quality rat precision cut liver slices during culture to study hepatotoxic responses: Acetaminophen as a model compound. *Toxicol In Vitro* **2017,** *42*, 200-213.

15.   Thiele, G. M.; Duryee, M. J.; Thiele, G. E.; Tuma, D. J.; Klassen, L. W., Review: Precision Cut Liver Slices for the Evaluation of Fatty Liver and Fibrosis. *Curr Mol Pharmacol* **2017,** *10* (3), 249-254.

16.	van de Bovenkamp, M.; Groothuis, G. M.; Draaisma, A. L.; Merema, M. T.; Bezuijen, J. I.; van Gils, M. J.; Meijer, D. K.; Friedman, S. L.; Olinga, P., Precision-cut liver slices as a new model to study toxicity-induced hepatic stellate cell activation in a physiologic milieu. *Toxicol Sci* **2005,** *85* (1), 632-8.

17.	Dewyse, L.; De Smet, V.; Verhulst, S.; Eysackers, N.; Kunda, R.; Messaoudi, N.; Reynaert, H.; van Grunsven, L. A., Improved Precision-Cut Liver Slice Cultures for Testing Drug-Induced Liver Fibrosis. *Front Med (Lausanne)* **2022,** *9*, 862185.

18.	Westra, I. M.; Oosterhuis, D.; Groothuis, G. M.; Olinga, P., The effect of antifibrotic drugs in rat precision-cut fibrotic liver slices. *PLoS One* **2014,** *9* (4), e95462.

19.	Westra, I. M.; Oosterhuis, D.; Groothuis, G. M.; Olinga, P., Precision-cut liver slices as a model for the early onset of liver fibrosis to test antifibrotic drugs. *Toxicol Appl Pharmacol* **2014,** *274* (2), 328-38.

20.	Westra, I. M.; Mutsaers, H. A.; Luangmonkong, T.; Hadi, M.; Oosterhuis, D.; de Jong, K. P.; Groothuis, G. M.; Olinga, P., Human precision-cut liver slices as a model to test antifibrotic drugs in the early onset of liver fibrosis. *Toxicol In Vitro* **2016,** *35*, 77-85.

21.	Wu, X.; Roberto, J. B.; Knupp, A.; Kenerson, H. L.; Truong, C. D.; Yuen, S. Y.; Brempelis, K. J.; Tuefferd, M.; Chen, A.; Horton, H.; Yeung, R. S.; Crispe, I. N., Precision-cut human liver slice cultures as an immunological platform. *J Immunol Methods* **2018,** *455*, 71-79.

22.	Huang, X.; Cai, H.; Ammar, R.; Zhang, Y.; Wang, Y.; Ravi, K.; Thompson, J.; Jarai, G., Molecular characterization of a precision-cut rat liver slice model for the evaluation of antifibrotic compounds. *Am J Physiol Gastrointest Liver Physiol* **2019,** *316* (1), G15-G24.

23.	Fuchs, B. C.; Hoshida, Y.; Fujii, T.; Wei, L.; Yamada, S.; Lauwers, G. Y.; McGinn, C. M.; DePeralta, D. K.; Chen, X.; Kuroda, T.; Lanuti, M.; Schmitt, A. D.; Gupta, S.; Crenshaw, A.; Onofrio, R.; Taylor, B.; Winckler, W.; Bardeesy, N.; Caravan, P.; Golub, T. R.; Tanabe, K. K., Epidermal growth factor receptor inhibition attenuates liver fibrosis and development of hepatocellular carcinoma. *Hepatology* **2014,** *59* (4), 1577-90.

24.	Sojoodi, M.; Wang, Y.; Erstad, D. J.; Caravan, P.; Lanuti, M.; Qadan, M.; Fuchs, B. C.; Hoang, K.; Or, Y. S.; Jiang, L.; Tanabe, K. K., EDP-297, a novel and potent fxr agonist, exhibit robust anti-fibrotic effects with significant liver function in a rat model of non-alcoholic steatohepatitis. *Journal of Hepatology* **2020,** *73*.

25.	Chen, H. C.; Awale, S.; Wu, C. P.; Lee, H. H.; Wu, H. T., Co-cultured bone marrow mesenchymal stem cells repair thioacetamide-induced hepatocyte damage. *Cell Biol Int* **2020,** *44* (12), 2459-2472.

26.	Pearen, M. A.; Lim, H. K.; Gratte, F. D.; Fernandez-Rojo, M. A.; Nawaratna, S. K.; Gobert, G. N.; Olynyk, J. K.; Tirnitz-Parker, J. E. E.; Ramm, G. A., Murine Precision-Cut Liver Slices as an Ex Vivo Model of Liver Biology. *J Vis Exp* **2020,** (157).

27.	Leaker, B. D.; Wang, Y.; Tam, J.; Anderson, R. R., Analysis of culture and RNA isolation methods for precision-cut liver slices from cirrhotic rats. *bioRxiv* **2023**, 2023.07.18.549535.

28.	Yu, T.; Wang, Y.; Zhang, H.; Johnson, C. H.; Jiang, Y.; Li, X.; Wu, Z.; Liu, T.; Krausz, K. W.; Yu, A.; Gonzalez, F. J.; Huang, M.; Bi, H., Metabolomics reveals mycoplasma contamination interferes with the metabolism of PANC-1 cells. *Anal Bioanal Chem* **2016,** *408* (16), 4267-73.

29.	Wang, Y.; Yu, T.; Zhou, Y.; Wang, S.; Zhou, X.; Wang, L.; Ou, T.; Chen, Y.; Zhou, Y.; Zhang, H.; Wang, Y.; Fan, X.; Chen, P.; Gonzalez, F. J.; Yu, A.; Huang, P.; Huang, M.; Bi, H.,

Carnitine palmitoyltransferase 1C contributes to progressive cellular senescence. *Aging (Albany NY)* **2020,** *12* (8), 6733-6755.

30.   Sojoodi, M.; Erstad, D. J.; Barrett, S. C.; Salloum, S.; Zhu, S.; Qian, T.; Colon, S.; Gale, E. M.; Jordan, V. C.; Wang, Y.; Li, S.; Ataeinia, B.; Jalilifiroozinezhad, S.; Lanuti, M.; Zukerberg, L.; Caravan, P.; Hoshida, Y.; Chung, R. T.; Bhave, G.; Lauer, G. M.; Fuchs, B. C.; Tanabe, K. K., Peroxidasin Deficiency Re-programs Macrophages Toward Pro-fibrolysis Function and Promotes Collagen Resolution in Liver. *Cell Mol Gastroenterol Hepatol* **2022,** *13* (5), 1483-1509.

31.   Ishak, K. G., Chronic hepatitis: morphology and nomenclature. *Mod Pathol* **1994,** *7* (6), 690-713.

32.   Wang, Y.; Chen, Y.; Guan, L.; Zhang, H.; Huang, Y.; Johnson, C. H.; Wu, Z.; Gonzalez, F. J.; Yu, A.; Huang, P.; Wang, Y.; Yang, S.; Chen, P.; Fan, X.; Huang, M.; Bi, H., Carnitine palmitoyltransferase 1C regulates cancer cell senescence through mitochondria-associated metabolic reprograming. *Cell Death Differ* **2018,** *25* (4), 735-748.

33.   Yu, T.; Wang, Y. T.; Chen, P.; Li, Y. H.; Chen, Y. X.; Zeng, H.; Yu, A. M.; Huang, M.; Bi, H. C., Effects of nicotinamide N-methyltransferase on PANC-1 cells proliferation, metastatic potential and survival under metabolic stress. *Cell Physiol Biochem* **2015,** *35* (2), 710-21.

34.   Fukawa, A.; Baba, S.; Iwasawa, K.; Yamaguchi, M.; Hosono, A., Dietary alpha-Lactalbumin protects against thioacetamide-induced liver cirrhosis by maintaining gut-liver axis function in rats. *Biosci Biotechnol Biochem* **2020,** *84* (1), 171-177.

35.   Li, S.; Ghoshal, S.; Sojoodi, M.; Arora, G.; Masia, R.; Erstad, D. J.; Lanuti, M.; Hoshida, Y.; Baumert, T. F.; Tanabe, K. K.; Fuchs, B. C., Pioglitazone Reduces Hepatocellular Carcinoma Development in Two Rodent Models of Cirrhosis. *J Gastrointest Surg* **2019,** *23* (1), 101-111.

36.   Sojoodi, M.; Wei, L.; Erstad, D. J.; Yamada, S.; Fujii, T.; Hirschfield, H.; Kim, R. S.; Lauwers, G. Y.; Lanuti, M.; Hoshida, Y.; Tanabe, K. K.; Fuchs, B. C., Epigallocatechin Gallate Induces Hepatic Stellate Cell Senescence and Attenuates Development of Hepatocellular Carcinoma. *Cancer Prev Res (Phila)* **2020,** *13* (6), 497-508.

37.   Tolba, R.; Kraus, T.; Liedtke, C.; Schwarz, M.; Weiskirchen, R., Diethylnitrosamine (DEN)-induced carcinogenic liver injury in mice. *Lab Anim* **2015,** *49* (1 Suppl), 59-69.

38.   Heindryckx, F.; Colle, I.; Van Vlierberghe, H., Experimental mouse models for hepatocellular carcinoma research. *Int J Exp Pathol* **2009,** *90* (4), 367-86.

39.   Nakagawa, S.; Wei, L.; Song, W. M.; Higashi, T.; Ghoshal, S.; Kim, R. S.; Bian, C. B.; Yamada, S.; Sun, X.; Venkatesh, A.; Goossens, N.; Bain, G.; Lauwers, G. Y.; Koh, A. P.; El-Abtah, M.; Ahmad, N. B.; Hoshida, H.; Erstad, D. J.; Gunasekaran, G.; Lee, Y.; Yu, M. L.; Chuang, W. L.; Dai, C. Y.; Kobayashi, M.; Kumada, H.; Beppu, T.; Baba, H.; Mahajan, M.; Nair, V. D.; Lanuti, M.; Villanueva, A.; Sangiovanni, A.; Iavarone, M.; Colombo, M.; Llovet, J. M.; Subramanian, A.; Tager, A. M.; Friedman, S. L.; Baumert, T. F.; Schwarz, M. E.; Chung, R. T.; Tanabe, K. K.; Zhang, B.; Fuchs, B. C.; Hoshida, Y.; Precision Liver Cancer Prevention, C., Molecular Liver Cancer Prevention in Cirrhosis by Organ Transcriptome Analysis and Lysophosphatidic Acid Pathway Inhibition. *Cancer Cell* **2016,** *30* (6), 879-890.

40.   Schon, H.-T.; Weiskirchen, R., Immunomodulatory effects of transforming growth factor-β in the liver. *Hepatobiliary Surgery and Nutrition* **2014,** *3* (6), 386-406.

41.   Wang, H.; Lafdil, F.; Wang, L.; Yin, S.; Feng, D.; Gao, B., Tissue inhibitor of metalloproteinase 1 (TIMP-1) deficiency exacerbates carbon tetrachloride-induced liver injury

and fibrosis in mice: involvement of hepatocyte STAT3 in TIMP-1 production. *Cell Biosci* **2011,** *1* (1), 14.

42.	Cheng, N.; Kim, K. H.; Lau, L. F., Senescent hepatic stellate cells promote liver regeneration through IL-6 and ligands of CXCR2. *JCI Insight* **2022,** *7* (14).

43.	Li, J.; Zhang, J.; Zhang, B.; Chen, L.; Chen, G.; Zhu, D.; Chen, J.; Duan, L.; Duan, Y., rSjP40 Inhibited the Activity of Collagen Type I Promoter via Ets-1 in HSCs. *Front Cell Dev Biol* **2021,** *9*, 765616.

44.	Geervliet, E.; Bansal, R., Matrix Metalloproteinases as Potential Biomarkers and Therapeutic Targets in Liver Diseases. *Cells* **2020,** *9* (5).

Figure 1

A

Fresh cirrhosis liver → Vibrotome instrument → PCLS (250 µm) → Anti-cirrhosis/fibrosis treatment (3 days) → Analysis

Analysis:
- MTS proliferation
- H&E staining
- Sirius Red staining
- RT-qPCR

B PCLS from CDAHFD-induced cirrhosis

C PCLS from DEN-induced cirrhosis

D PCLS from CCl4-induced cirrhosis

E PCLS from TAA-induced cirrhosis

Figure 2

Ex vivo PCLS from CDAHFD-induced mouse cirrhotic livers

A

Il6

Col1a1

Timp1

Tgf-β

Acta2

B CDAHFD + Vehicle CDAHFD + Erlotinib

C

Ex vivo PCLS from TAA-induced rat cirrhotic livers

D

Il6

Col1a1

Tgfβ

Timp1

Acta2

E TAA + Vehicle TAA + Erlotinib

F

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Supplementary Figure 1

A

B

C

Model types	Model induction	Erlotinib treatment
ex vivo PCLS from 4 murine cirrhotic models	CDAHFD for 16 weeks	72 hours
	200 mg/kg thioacetamide for 12 weeks	
	50 mg/kg DEN for 18 weeks	
	40% CCl4 for 18 weeks	
ex vivo PCLS from murine normal livers	Tgf-β for 96 hours	72 hours
in vitro human HSC (LX2, TWNT4)		
in vivo murine cirrhotic models	50 mg/kg DEN for 18 weeks	10 weeks
	40% CCl4 for 18 weeks	6 weeks
ex vivo PCLS from human cirrhotic liver		72 hours

Supplementary Figure 2

Supplementary Figure 3

A

 PCLS from thioacetamide-induced cirrhosis

Supplementary Figure 4

Supplementary Figure 5

Ex vivo PCLS from rodent cirrhotic livers

A CDAHFD + Vehicle   CDAHFD + Erlotinib

B TAA + Vehicle   TAA + Erlotinib

C DEN + Vehicle   DEN + Erlotinib

D CCl4 + Vehicle   CCl4 + Erlotinib

In vivo PCLS from rodent cirrhotic livers

E DEN + Vehicle   DEN + Erlotinib

F CCl4 + Vehicle   CCl4 + Erlotinib

Ex vivo PCLS from rodent normal livers

G Mock   TGF-β1 + Vehicle   TGF-β1 + Erlotinib

Ex vivo PCLS from human cirrhotic livers

H Vehicle   Erlotinib

Supplementary Figure 6

| Human established cirrhosis | PCLS from human established cirrhosis |

A H&E

B Sirius Red

C H&E

D Sirius Red