Different complex regulatory phenotypes underlie hybrid male sterility in divergent rodent crosses

Kelsie E. Hunnicutt[*,1], Colin Callahan[2], Sara Keeble[2], Emily C. Moore[1,2], Jeffrey M. Good[2], and Erica L. Larson[*,1]

[1] University of Denver, Department of Biological Sciences, Denver, CO, 80208
[2] Division of Biological Sciences, University of Montana, Missoula, MT, 59812
*Corresponding authors

ORCID: 0000-0002-9674-0630 (KEH)
ORCID: 0000-0002-0385-1121 (CC)
ORCID: 0000-0002-7318-3122 (SK)
ORCID: 0000-0001-6166-3367 (ECM)
ORCID: 0000-0003-0707-5374 (JMG)
ORCID: 0000-0003-3006-645X (ELL)

Corresponding Authors:
Kelsie Hunnicutt
Department of Biological Sciences
University of Denver
Denver, CO 80208
Email: Kelsie.Hunnicutt@du.edu

Erica Larson
Department of Biological Sciences
University of Denver
Denver, CO 80208
Email: Erica.Larson@du.edu

Running title: Rodent hybrid male sterility

Key words: speciation, hybrid male sterility, gene expression, fluorescence activated cell sorting, meiotic sex chromosome inactivation, *Phodopus*

1 **ABSTRACT**
2
Hybrid incompatibilities are a critical component of species barriers and may arise due

to negative interactions between divergent regulatory elements in parental species. We used a

comparative approach to identify common themes in the regulatory phenotypes associated

with hybrid male sterility in two divergent rodent crosses, dwarf hamsters and house mice. We

investigated three potential characteristic regulatory phenotypes in hybrids including the

propensity towards over or underexpression relative to parental species, the influence of

developmental stage on the extent of misexpression, and the role of the sex chromosomes on

misexpression phenotypes. In contrast to near pervasive overexpression in hybrid house mice,

we found that misexpression in hybrid dwarf hamsters was dependent on developmental

stage. In both house mouse and dwarf hamster hybrids, however, misexpression increased

with the progression of spermatogenesis, although to varying extents and with potentially

different consequences. In both systems, we detected sex-chromosome specific

overexpression in stages of spermatogenesis where inactivated X chromosome expression

was expected, but the hybrid overexpression phenotypes were fundamentally different.

Importantly, misexpression phenotypes support the presence of multiple histological blocks to

spermatogenesis in dwarf hamster hybrids, including a potential role of meiotic stalling early in

spermatogenesis. Collectively, we demonstrate that while there are some similarities in hybrid

regulatory phenotypes of house mice and dwarf hamsters, there are also clear differences that

point towards unique mechanisms underlying hybrid male sterility in each system. Our results

highlight the potential of comparative approaches in helping to understand the importance of

disrupted gene regulation in speciation.

INTRODUCTION

The evolution of postzygotic reproductive barriers, such as hybrid inviability and

sterility, is an important part of the speciation process, and identifying the genetic architecture

of hybrid incompatibilities has been a common goal uniting speciation research (Coughlan and

Matute 2020). While identifying the genetic basis of hybrid dysfunction remains difficult in many

systems, downstream regulatory phenotypes can provide insight into the underlying

mechanisms of speciation (Mack and Nachman 2017). An outstanding question surrounding

the role of disrupted gene regulation and speciation is whether the combination of two

divergent genomes in hybrids results in expression perturbations that are consistent or

repeatable across species. At the broadest level, it is unclear whether hybrids tend to have

over or underexpression of genes relative to parental species (Ortíz-Barrientos et al. 2007), and

whether misexpression is modulated by cis or trans regulatory elements (Wittkopp et al. 2004;

McManus et al. 2010; Oka et al. 2014; Mack et al. 2016; Mugal et al. 2020; Kopania, Larson, et

al. 2022). We might predict that misexpression in hybrids would be biased towards

overexpression if hybrid incompatibilities disrupt gene repression (Meiklejohn et al. 2014;

Barreto et al. 2015; Larson et al. 2017). Alternatively, expression in hybrids may tend towards

underexpression if transcription factor binding with promoter or enhancer elements is impaired

(Oka et al. 2014; Guerrero et al. 2016) or if divergence stimulates epigenetic silencing (Paun et

al. 2007; Shivaprasad et al. 2012; Lafon-Placette and Köhler 2015; Zhu et al. 2017). At a finer

scale, misexpression may depend on developmental stage: for example, if there is greater

pleiotropy earlier in development (Ortíz-Barrientos et al. 2007; Cutter and Bundus 2020). In

particular, we might expect sterile hybrids to have more misexpression during later stages of

gametogenesis when genes are evolving rapidly and are potentially under less regulatory

constraint (Kopania, Larson, et al. 2022; Murat et al. 2023). Finally, the role of sex chromosome

regulation in inviable or sterile hybrids encompasses both of these larger questions. Sex

chromosomes may be prone to asymmetry in their expression divergence (Oka and Shiroishi

2014; Civetta 2016) and be regulated differently across different stages of development

(Presgraves 2008; Larson et al. 2018), particularly in reproductive tissues, and thus may play a

central role in hybrid dysregulation relative to autosomes.

Disruption of sex chromosome regulation is thought to be a potentially widespread

regulatory phenotype in sterile hybrids (Lifschytz and Lindsley 1972; Larson et al. 2018), in part

because X chromosome repression may be crucial to normal spermatogenesis in diverse taxa

(McKee and Handel 1993; Landeen et al. 2016; Taxiarchi et al. 2019; Rappaport et al. 2021;

Viera et al. 2021; Murat et al. 2023). Furthermore, misregulation of the X chromosome is

associated with hybrid sterility in several species pairs (Davis et al. 2015; Bredemeyer et al.

2021; Sánchez-Ramírez et al. 2021), although it has been best studied in house mice. In fertile

male mice, the X chromosome is silenced early in meiosis through meiotic sex chromosome

inactivation (MSCI; McKee and Handel 1993; Handel 2004) and is again repressed in

postmeiotic sperm development (*i.e.,* postmeiotic sex chromosome repression or PSCR;

Namekawa et al. 2006). In contrast, the X chromosome is not properly inactivated and is

overexpressed in sterile hybrid mice (Good et al. 2010; Bhattacharyya et al. 2013; Campbell et

al. 2013; Turner et al. 2014; Larson et al. 2017; Larson et al. 2022). Disrupted MSCI in house

mice is associated with divergence at *Prdm9*, a gene that is a major contributor to hybrid male

sterility (Mihola et al. 2009; Davies et al. 2016). However, misexpression of the X chromosome

in sterile hybrids could result from mechanisms other than *Prdm9*-associated disrupted MSCI,

such as mispairing of the sex chromosomes due to divergence in their region of homology

known as the pseudoautosomal region (PAR; Burgoyne 1982; Ellis and Goodfellow 1989;

Raudsepp and Chowdhary 2015). In sum, the ubiquity of X chromosome repression and the

growing body of evidence linking disrupted MSCI to hybrid sterility in mammals suggest that

disrupted sex chromosome regulation may be a common regulatory phenotype in sterile hybrid

males.

Here, we characterized disruption of gene regulation associated with hybrid male

sterility in two rodent crosses, dwarf hamsters and house mice, which span ~35 million years

of divergence (Swanson et al. 2019). The regulatory phenotypes of hybrid male sterility have

been thoroughly studied in house mice (Good et al. 2010; Bhattacharyya et al. 2013; Campbell

et al. 2013; Turner et al. 2014; Larson et al. 2017; Hunnicutt et al. 2022; Larson et al. 2022). We

contrast these with an analogous cross between two sister species of dwarf hamster,

Campbell's dwarf hamster (*Phodopus campbelli*) and the Siberian dwarf hamster (*P.*

sungorus), and their sterile F1 hybrid male offspring. These species diverged only ~0.8-1.0

million years ago (Neumann et al. 2006), and they are not thought to interbreed in the wild due

to geographic separation (Ishishita et al. 2015). Crosses between female *P. sungorus* and male

P. campbelli produce sterile hybrid males that, similar to mice, have a range of sterility

phenotypes, suggesting multiple histological blocks to spermatogenesis (Ishishita et al. 2015;

Bikchurina et al. 2018). Hybrids from the reciprocal cross are usually inviable due to abnormal

growth in utero (Brekke and Good 2014). Intriguingly, the species origin of the X chromosome

is the primary genetic factor controlling hybrid inviability (Brekke et al. 2021). The frequently

observed asynapsis of the sex chromosomes in hybrid dwarf hamsters provide additional

reason to think that X chromosome-specific misregulation may be observed in sterile hybrid

dwarf hamsters (Ishishita et al. 2015; Bikchurina et al. 2018). The combination of these

abnormal histological phenotypes and the interactions between diverse reproductive barriers

make dwarf hamsters an important comparison to mice for investigating what regulatory

phenotypes may be associated with reproductive isolation between species.

Regulatory phenotypes associated with hybrid sterility have historically been difficult to

assess because the cellular diversity of reproductive tissues (*e.g.,* testes; Ramm and Schärer

2014) confounds inferences of gene expression (Good et al. 2010; Montgomery and Mank

2016; Hunnicutt et al. 2022), thus requiring molecular approaches that permit greater spatial

and developmental resolution. To overcome these difficulties, we used Fluorescence Activated

Cell Sorting (FACS) to isolate and sequence cell populations across the developmental timeline

of spermatogenesis for each species pair and their F1 hybrids, including stages that span the

different sex chromosome regulatory states. We used both datasets to address three main

questions about the regulatory phenotypes observed in sterile hybrids: (1) does misexpression

tend towards up- or downregulation in hybrids compared to parents? (2) are there similar

patterns of disrupted expression across stages of development? and (3) is there a clear

differentiation between autosomes and sex chromosomes in regulatory phenotype? And if so,

is sex chromosome specific misexpression consistent with either disrupted MSCI and/or

disrupted PAR regulation? Collectively, we demonstrate the power of cell-specific approaches

for untangling the regulatory phenotypes associated with the evolution of hybrid male sterility

and for identifying common themes in the mechanistic basis of hybrid incompatibilities across

divergent taxa.

RESULTS

Impaired sperm production in hybrid male dwarf hamsters

We first established the extent of hybrid male sterility in dwarf hamsters by comparing

reproductive phenotypes for *P. campbelli*, *P. sungorus*, and F1 hybrid males (Figure 1; Table

S1). *Phodopus sungorus* males had smaller testes and seminal vesicles than *P. campbelli*

males (Dunn's Test relative testes weight p < 0.001; relative seminal vesicle weight p = 0.0061;

Figure 1), perhaps reflecting inbreeding depression on this laboratory colony. However, these

species did not differ in normalized sperm counts (p = 0.071) or in sperm motility (p = 0.12).

The F1 hybrid males exhibited extreme reproductive defects relative to *P. campbelli* (Figure 1).

Hybrid males had smaller testes (p < 0.001) and seminal vesicles (p < 0.001) than male *P.*

campbelli hamsters, and importantly, produced almost no mature spermatozoa. In the one

instance where a hybrid male produced a single mature spermatozoon, it was non-motile,

indicating severe reproductive impairment in hybrid males. Overall, our results confirmed

previous reports of reduced fertility in hybrid male dwarf hamsters (Ishishita et al. 2015;

Bikchurina et al. 2018).

Figure 1. Hybrid dwarf hamsters exhibit multiple sterility phenotypes. We assessed paired

testes weight and paired seminal vesicle weight (SV), both normalized by body weight, sperm

count, and proportion of motile sperm for *P. sungorus*, *P. campbelli*, and F1 hybrids. Whiskers

extend to either the largest or smallest value or no further than 1.5 times the interquartile range,

and *** indicates p < 0.001, ** indicates p < 0.01, and n.s. indicates non-significant difference

between means at p > 0.05 using a post-hoc Dunn's test with FDR correction. Upwards-

pointing triangles (▲) indicate *P. campbelli*, downwards-pointing triangles (▼) indicate *P.*

sungorus, and crosses (✖) indicate F1 hybrids.

Cell-specific gene expression across spermatogenesis

To characterize cell-specific gene expression, we used FACS to isolate enriched cell

populations from each fertile parent species and their sterile F1 hybrids across four stages of

spermatogenesis. The four targeted populations included: spermatogonia (mitotic precursor

cells), leptotene/zygotene spermatocytes (meiotic cells before MSCI), diplotene spermatocytes

(meiotic cells after MSCI), and round spermatids (postmeiotic cells). We were unable to isolate

round spermatids from the F1 hybrids, which was consistent with the lack of mature

spermatozoa present in the cauda epididymis extractions (Figure 1). We sequenced RNA from

each cell population for *P. campbelli* (spermatogonia n = 4, leptotene/zygotene n = 5, diplotene

n = 5, and round spermatids n = 4; Table S2), *P. sungorus* (spermatogonia n = 4,

leptotene/zygotene n = 4, diplotene n = 5, and round spermatids n = 3), and F1 hybrid males (

P. campbelli♀ x *P. sungorus*♂; spermatogonia n = 4, leptotene/zygotene n = 4, diplotene n =

4). We compared our hamster expression data to an analogous cell-specific RNASeq dataset

from two species of house mice, *Mus musculus musculus* and *M. m. domesticus*, and their

sterile F1 hybrids (Larson et al. 2017).

We used two approaches to evaluate the purity of spermatogonia, leptotene/zygotene

spermatocytes, diplotene spermatocytes, and round spermatids isolated from males from both

parental species: (1) quantifying the relative expression of a panel of cell population marker

genes and (2) characterizing the expression patterns of the sex chromosomes across

development. We found that our candidate marker genes had the highest expression in their

expected cell population for all stages except leptotene/zygotene (Figures S4 and 5), indicating

high purity of spermatogonia, diplotene spermatocytes, and round spermatids.

Leptotene/zygotene markers did not have the highest expression in leptotene/zygotene

samples (except for *Ccnb1ip1*), potentially indicating lower purity of this cell population.

Nonetheless, the patterns of X chromosome expression in fertile parents was consistent with

expectations across this developmental timeline: the X chromosome had active expression in

spermatogonia and leptotene/zygotene cells, was inactivated in diplotene cells consistent with

MSCI, and was partially inactivated in round spermatids, consistent with PSCR (Figure 2;

Namekawa et al. 2006), indicating successful isolation of these cell populations. When we

examined overall expression differences among dwarf hamsters, we found that samples

separated primarily by cell population, then by species within a cell population, with a lack of

clear separation of spermatogonia and leptotene/zygotene samples but distinct populations of

diplotene and round spermatids (Figure 3). Hybrids showed intermediate expression level

relative to parent species for all cell populations (Figures S2 and 3). The expression profiles of

hybrid diplotene cells ranged from clustering with parental leptotene/zygotene to parental

diplotene cells, which contrasted with what we observed in house mice where hybrid diplotene

cells clustered distinctly with parental diplotene cells (Figure 3). Overall, our results indicate

that we successfully isolated cell populations in parental dwarf hamsters that span the full

extent of spermatogenesis.

Figure 2. Overexpression of X-linked genes in diplotene spermatocytes in both house mouse and dwarf hamster hybrids. Heatmap of X-linked gene expression in house mice (upper panel) and dwarf hamsters (lower panel) plotted as normalized FPKM values that are hierarchically clustered using Euclidean distance. Each column represents a different individual, each row represents a gene, and darker colors indicate higher expression. The heatmap was generated with the R package ComplexHeatmap v.2.12.0 (Gu et al. 2016). Note, hybrid dwarf hamsters do not produce mature spermatozoa, and accordingly, we were unable to isolate round spermatids.

Figure 3. Hybrid gene expression profiles cluster by parental spermatogenic cell

population in house mice but not dwarf hamsters. Multidimensional scaling (MDS) plots of

distances among house mouse (upper panels) and dwarf hamster (lower panels) samples for

expressed autosomal (left) and X-linked (right) genes. Distances are calculated as the root-

mean-square deviation (Euclidean distance) of log2 fold changes among genes that distinguish

each sample. Each species and F1 hybrid is indicated by a symbol, and samples are colored

by cell population.

Disrupted transcriptional activation early in spermatogenesis in dwarf hamsters

We sought to characterize which regulatory phenotypes were associated with sterile

hybrids in both house mice and dwarf hamsters. We first investigated whether misexpression

tends towards up- or downregulation in hybrids compared to parents [measured as the log fold

change (logFC) in expression between hybrids and parents sharing the same X chromosome

for differentially expressed (DE) genes]. Misexpression in hybrid house mice was biased

towards upregulation across spermatogenesis in sterile hybrids (average logFC of

spermatogonia autosomal DE genes: +1.17/ (X^2): p < 0.001; diplotene: +1.13/ (X^2): p < 0.001;

Figure 4), as was X-linked misexpression (spermatogonia average logFC: +1.63/ (X^2): p < 0.001;

leptotene/zygotene: +1.20 (X^2): p < 0.001; diplotene: +2.54/ (X^2): p < 0.001), with only the

exception of autosomal leptotene/zygotene DE genes (leptotene/zygotene: +0.49/ (X^2): p =

0.408). In contrast, we found that the direction of misexpression in dwarf hamster hybrids was

stage-specific. Differentially expressed autosomal genes in dwarf hamster hybrids were

overwhelmingly downregulated in both early stages of spermatogenesis especially in

comparison to house mice (average logFC of spermatogonia autosomal DE genes: -2.63/ (X^2):

p < 0.001; leptotene/zygotene: -1.73/ (X^2): p < 0.001; Figure 4) but upregulated in diplotene

(average logFC = +2.15/ (X^2): p < 0.001; Figure 4). When comparing both X-linked and

autosomal expression in dwarf hamsters, we found similar patterns: almost all differentially

expressed X-linked genes in the first two stages of spermatogenesis were exclusively

downregulated (spermatogonia average logFC = -6.50/ (X^2): p < 0.001; leptotene/zygotene: -

5.44/ (X^2): p < 0.001), and differential X-linked expression in diplotene was biased towards

upregulation (average logFC: 5.57/ (X^2): p < 0.001). These patterns held regardless of which

parent species was used as a contrast for dwarf hamsters (Figure S6). When hybrid house

mice were compared with the parent species with a different X chromosome (*M. m.*

domesticus), the direction of misexpression was positive for autosomal and X-linked

spermatogonia and diplotene, as in the comparisons with *M. m. musculus*, but negative for

autosomal and X-linked leptotene/zygotene.

Figure 4. House mice and dwarf hamster hybrids have opposite patterns of disrupted regulation early in spermatogenesis. logFC of differentially expressed genes between hybrids and their parent species with the same X chromosome (*i.e.,* hybrids versus *M. m. musculus* mice; hybrid versus *P. campbelli* dwarf hamsters) for each stage. Results are displayed for autosomes (left) and the X chromosome (right). *** indicates p < 0.001 for pairwise comparisons from Wilcoxon signed-rank tests after FDR correction. Whiskers extend to either the largest or smallest value or no further than 1.5 times the interquartile range. The number of up and downregulated differentially expressed genes in hybrids are listed next to arrows indicating direction of differential expression.

Second, we investigated how developmental stage influenced the extent of hybrid misexpression. For both dwarf hamsters and house mice, misregulation in hybrids increased with the progression of spermatogenesis, though to a greater extent in dwarf hamsters (Figure S9). In dwarf hamsters, there were substantially more differentially expressed genes in diplotene spermatocytes than earlier stages of spermatogenesis (spermatogonia = 1,386, leptotene/zygotene = 966, and diplotene = 12,015 genes). In mice, there was also a trend

towards misexpression late, although it was not as dramatic as in dwarf hamsters

(spermatogonia = 2,360, leptotene/zygotene = 2,434, diplotene = 2,673, and round spermatids

= 4,181 genes). We also found a much greater genome-wide disruption of expression in

diplotene cells of dwarf hamsters - nearly half the transcriptome was misexpressed (Figures

S7-S9), indicating more widespread regulatory disruption than was observed in house mouse

hybrids.

To determine if the differences we observed in the extent of misregulation between

dwarf hamsters and house mice was due to greater expression variability across our dwarf

hamster samples, we calculated the Biological Coefficient of Variation (BCV), a measurement

of inter-replicate variability, for each species and hybrid across the first three cell populations.

Inter-replicate variability was higher in dwarf hamster hybrids relative to house mouse hybrids

(dwarf hamsters BCV = 0.69; house mice = 0.18). However, the extent of the variability

observed in hybrids relative to the inter-replicate variability of parental species differed

between house mice and dwarf hamsters: dwarf hamster hybrid variability was less than dwarf

hamster parental samples (*P. campbelli* = 0.49; *P. sungorus* = 0.40), but hybrid variability was

similar to parental samples for house mice (*M. m. musculus* = 0.19; *M. m. domesticus* = 0.22).

The greater inter-replicate variability in dwarf hamster hybrids relative to parent species

suggests that the increased misexpression we see in hybrids cannot be explained by greater

inter-replicate variability in our dwarf hamster samples. Thus, while there were hints of

disruption early in spermatogenesis in both mouse and dwarf hamster hybrids, misexpression

was amplified across the developmental timeline in both species.

Third, we characterized whether there was a clear difference between the autosomes

and sex chromosomes in regulatory phenotype. Across all stages of spermatogenesis in house

mice, autosomal expression is relatively unperturbed in contrast to greatly disrupted sex

chromosome expression (Figure 4). Surprisingly, misregulation in dwarf hamsters early in

spermatogenesis was not sex chromosome-specific, as the sex chromosomes in dwarf

hamsters showed no over-enrichment for differentially expressed genes early in

spermatogenesis between parent species and hybrids (spermatogonia: $p = 0.995$;

leptotene/zygotene: $p = 0.30$; Figure S7). However, we note that the magnitude of

misregulation was greater for sex chromosomes than autosomes in dwarf hamsters for both

spermatogonia and leptotene/zygotene (Figure 4; discussed above). Only one autosome

showed either over- or under-enrichment for differentially expressed genes in the first two cell

populations: over-enrichment of chromosome 11 in spermatogonia (hypergeometric test; $p =$

0.018) and leptotene/zygotene ($p = 0.017$; Figure S7). However, the X chromosome was

enriched for DE genes in diplotene cells in hybrid dwarf hamsters (observed X-linked DE genes

$= 585$; expected X-linked DE genes $= 247$; $p < 0.001$; Figure S7), consistent with sex

chromosome-specific misexpression in this cell population. In mice, the X chromosome was

over-enriched for differentially expressed genes between hybrids and parents for all stages

except leptotene/zygotene (spermatogonia $p = 0.001$; leptotene/zygotene $p = 0.436$; diplotene

$p < 0.001$; round spermatids $p < 0.001$; Figure S8), suggesting a difference in the extent of the

role for sex chromosome-specific disruption between systems.

Misexpression in diplotene appears unrelated to disrupted MSCI in dwarf hamsters

We next asked whether similar disrupted regulatory processes resulted in sex

chromosome-specific misexpression in both systems. In sterile hybrid house mice, X-linked

genes were upregulated during diplotene relative to autosomal genes (Wilcoxon signed rank

test: $p < 0.001$; Figure 2), consistent with disrupted MSCI (Good et al. 2010; Bhattacharyya et

al. 2013; Campbell et al. 2013; Turner and Harr 2014; Larson et al. 2017; Larson et al. 2022). In

sterile dwarf hamster hybrids, we also found that X-linked genes were upregulated during

diplotene relative to autosomal genes (Wilcoxon signed rank test: $p < 0.001$), but the extent of

295 X chromosome overexpression was greater than in hybrid house mice (average logFC X-linked

296 genes dwarf hamster: 5.57; house mice: 2.54; Wilcoxon signed rank test: $p < 0.001$; Figure 2).

297 There was also more variability in the relative extent of overexpression of X-linked genes in

298 hybrid dwarf hamsters during diplotene compared to the mean expression of X-linked genes of

299 parental dwarf hamsters (relative overexpression = 18.5 +/- 6.8) compared to hybrid house

300 mice (relative overexpression = 1.75 +/- 0.09; Figures 2 and S10). Strikingly, some hybrid male

301 dwarf hamsters had an almost completely silenced X chromosome, while others had an almost

302 completely transcriptionally-activated X chromosome (Figure 2).

303  Despite overexpression of the X chromosome during diplotene in hybrid dwarf

304 hamsters, the overall regulatory phenotype, including the identity and the extent of

305 misexpression of overexpressed genes, appeared to fundamentally differ between house

306 mouse and dwarf hamster hybrids (Figures 2 and 5). We established these differences in X-

307 linked overexpression using two approaches. First, we tested which parental cell types had the

308 highest expression correlation with hybrid diplotene cell types for both X-linked and autosomal

309 genes. In mice, the expression profile of X-linked genes in hybrids during diplotene was most

310 positively correlated with the expression profile of X-linked parental round spermatid genes,

311 consistent with disrupted MSCI (spermatogonia *(r)* = -0.25, $p < 0.001$; leptotene/zygotene *(r)* = -

312 0.073, $p = 0.035$; round spermatid *(r)* = 0.26, $p < 0.001$; Figures 2 and 5A). Autosomal genes

313 showed no positive correlations with either spermatogonia (*r* = -0.26; $p < 0.001$),

314 leptotene/zygotene (*r* = -0.040; $p < 0.001$), or round spermatids (*r* = -0.060; $p < 0.001$; Figure

315 5). If the X-linked overexpression phenotype in dwarf hamsters was consistent with disrupted

316 MSCI, then we would also expect X-linked and autosomal expression profiles in hybrid

317 diplotene to follow the same patterns. In contrast to this prediction, hybrid dwarf hamster

318 diplotene expression profiles for both X-linked and autosomal genes had a positive correlation

319 with parental leptotene/zygotene (autosomal *(r)* = 0.13, $p < 0.001$; X-linked *(r)* = 0.22, $p < 0.001$)

and spermatogonia (autosomal *(r)* = 0.027, p < 0.001; X-linked *(r)* = 0.091, p = 0.017) and a

negative correlation with round spermatids (autosomal *(r)* = -0.30, p < 0.001; X-linked *(r)* = -

0.31, p < 0.001; Figure 5). These striking differences in the strength and direction of expression

profile correlations suggest that the regulatory mechanisms underlying the overexpression

phenotype of X-linked genes in sterile hybrids differed between house mice and dwarf

hamsters.

This difference in pattern was further supported when we compared which sets of

genes were overexpressed in hybrid diplotene. For this approach, we characterized which

stages of spermatogenesis genes were active in and characteristic of, then identified which X-

linked genes in hybrid diplotene were overexpressed (defined as genes with normalized

expression in the top 10% of X-linked genes), and finally compared which parental stages

overexpressed X-linked genes were characteristic of in both systems. Similar to our correlation

analysis, we found that in hybrid house mice, the genes that were overexpressed in diplotene

most closely resembled genes that characterize round spermatids in parental mice (56.6% of

genes), but that in hybrid dwarf hamsters, overexpressed diplotene genes resembled

spermatogonia and leptotene/zygotene specific genes (45.6% and 44.6% respectively; Figure

5C). Because of the overwhelming similarity of the hamster hybrid diplotene samples to

parental spermatogonia and leptotene/zygotene samples, we next performed gene ontology

(GO) enrichment analyses on the set of differentially expressed genes in hybrid diplotene

samples relative to *P. campbelli* diplotene samples to determine which biological processes

could be potentially contributing to this pattern. Three Biological Process GO terms were

significant after FDR correction: "negative regulation of cell proliferation", "positive regulation

of gene expression", and "positive regulation of cell migration" (Figure 5D and S11). The first

two GO terms were consistent with the patterns we observed in our hybrid dwarf hamster

samples: "negative regulation of cell proliferation" could explain the similarity of our dwarf

hamster hybrid diplotene samples to the earlier spermatogenic cell types, and "positive

regulation of gene expression" was consistent with the dramatic overexpression phenotype of

both autosomal and X-linked genes during this stage. Collectively, these results suggest that

the X-linked overexpression phenotype in sterile hybrid dwarf hamsters is inconsistent with

disrupted MSCI and is possibly related to a stalling of spermatogenesis between

leptotene/zygotene and diplotene during Prophase I.

Figure 5. Spermatogenesis in hybrid dwarf hamsters appears to stall after leptotene/

zygotene. A. We calculated the Pearson's correlation coefficient *(r)* between mean hybrid

diplotene expression and the mean expression for each parental cell type for both house mice

and **B.** dwarf hamsters. Correlation coefficients were calculated for both autosomal (left panels)

and X-linked (right panels) genes. We then generated bootstrap values by randomly sampling

the expression matrices for 1000 replicates. All values of *r* were significantly different from zero

(p < 0.05) after FDR correction. **C.** Classification of X-linked overexpressed genes in diplotene

cells of hybrid mice and dwarf hamsters by parental stage in which genes are induced.

Samples are colored by sample type (red = spermatogonia, yellow = leptotene/zygotene, green

= diplotene, and blue = round spermatids). **D.** The top 10 Biological Process GO terms (ranked

by FDR) for differentially expressed genes between hybrid dwarf hamsters and *P. campbelli* are

listed. Significant GO terms are colored while non-significant GO terms (FDR > 0.05) are gray.

Size of each point corresponds to the number of genes belonging to each GO term, and terms

are plotted by the fold enrichment of the GO term in the dataset relative to the provided gene

backgrounds.

PAR expression was not disrupted in sterile hybrid dwarf hamsters

370   Hybrid sterility in dwarf hamsters may be correlated with asynapsis of the X and Y

chromosomes because of divergence in the pseudoautosomal region (PAR) which prevents

proper chromosome pairing (Bikchurina et al. 2018). The PAR is the only portion of the sex

chromosomes that is able to synapse during routine spermatogenesis, and PAR genes are

assumed to escape silencing by MSCI (Raudsepp and Chowdhary 2015). However, because

XY asynapsis is common in dwarf hamster hybrids, Bikchurina et al. (2018) hypothesized that

MSCI may extend to the PAR of hybrid male dwarf hamsters, resulting in the silencing of PAR

genes in hybrids that may be critical to meiosis (Figure 6A). To test this hypothesis, we

compared the expression of genes located in the dwarf hamster PAR (Moore et al. 2022)

between parental dwarf hamster species and hybrid offspring across the timeline of

spermatogenesis. Specifically, we hypothesized that if XY asynapsis results in an extension of

MSCI to the PAR in hybrid dwarf hamsters, then hybrids should have similar PAR gene

expression to parents early in meiosis before homologous chromosome synapse during

pachytene. This should be followed by the silencing of PAR genes in hybrids, but not parent

species, during diplotene. If XY asynapsis does not alter the regulation of the PAR in hybrids,

then we may see two possible patterns. First, if all PAR genes are critical to the later stages of

spermatogenesis, then PAR genes in both hybrids and parents should be uniformly expressed

in diplotene. Alternatively, if PAR genes are not critical to the later stages of spermatogenesis,

then hybrids and parent species should have similar PAR gene expression, and not all PAR

genes may be expressed during diplotene.

We did not find evidence supporting PAR-wide silencing in dwarf hamster hybrids

during diplotene suggesting that MSCI is not extended to the PAR in dwarf hamster hybrids

because of XY asynapsis (Figure 6B). Furthermore, we also do not see PAR-wide expression of

genes during diplotene in hybrids or parents, indicating that not all PAR genes are critical to

the progression of spermatogenesis in dwarf hamsters. In general, most PAR gene expression

followed similar trends between hybrids and parent species. Three PAR genes were

differentially expressed between hybrids and *P. campbelli* during diplotene (*Ndrg2*,

Psun_G000022883, and Psun_G000022886; Table S3), but these genes were still expressed in

hybrids. Further, an association between PAR misregulation during the early stages of

spermatogenesis and hybrid male sterility also seems unlikely as only one gene, *Gprin1*,

showed consistent differential expression in early meiosis between both parent species and

hybrids (Table S3). Thus, based on the current annotation of the PAR in *P. sungorus*, we

currently find no direct evidence linking improper silencing of PAR genes to hybrid male sterility

in dwarf hamsters.

Figure 6. PAR gene expression is not disrupted during spermatogenesis in hybrid dwarf

hamsters. A. Hypothesized types of PAR gene regulation across spermatogenesis. If MSCI

extends to the entire X chromosome, then PAR genes would show some level of expression

early in spermatogenesis which would then drop to zero in diplotene when MSCI occurs. If the

PAR escapes silencing by MSCI and if PAR genes are critical to spermatogenesis, then we

would expect PAR genes to be uniformly expressed in diplotene when the rest of the X

chromosome is silenced. Finally, if the PAR escapes silencing by MSCI but all PAR genes are

not critical to spermatogenesis, then we would expect some PAR genes to be expressed and

some to be not expressed in diplotene. **B.** Observed patterns of PAR gene expression (as

mean normalized RPKM across individuals) in parental species and hybrid dwarf hamsters for

all PAR genes (indicated by line type and color) across spermatogenesis (SP = spermatogonia,

LZ = leptotene/zygotene·, and DIP = diplotene).

DISCUSSION

We used a comparative approach to understand common gene expression phenotypes

associated with hybrid male sterility in two divergent rodent crosses. We characterized

asymmetry of misexpression in hybrids, how misexpression changed over developmental

timelines, and how the X chromosome and autosomes differed in both of these aspects. We

found that while there were similarities in hybrid regulatory phenotypes in house mice and

dwarf hamsters, there were also differences in the timing and chromosomal distribution of

disrupted gene expression that point towards different underlying mechanisms behind hybrid

male sterility.

Asymmetry and developmental timing of misexpression in hybrids

We first investigated patterns of gene misexpression in sterile male hybrids. Studies of

misexpression in sterile or inviable hybrids have often focused on whether hybrid expression is

biased towards over or underexpression, with the hypothesis that expression may be biased

towards overexpression if hybrid incompatibilities disrupt repressive gene regulatory elements

(Meiklejohn et al. 2014; Barreto et al. 2015; Larson et al. 2017). In house mice, there is strong

support for overexpression of both autosomal and X-linked genes in sterile F1 hybrids (Mack et

al. 2016; Larson et al. 2017; Hunnicutt et al. 2022; Larson et al. 2022), and further work has

shown that F1 hybrid expression patterns depend on both autosomal background and sex

chromosome mismatch (Kopania, Watson, et al. 2022). Surprisingly, we found that in dwarf

hamster hybrids, there was nearly uniform downregulation of differentially expressed genes in

mitotic cells and early meiotic cells, suggesting that a loss of regulatory repression is not an

inevitable outcome of hybrid genomes. Asymmetric patterns of misexpression have been

found in many hybrids, including underexpression in sterile *Drosophila* hybrids (Michalak and

Noor 2003; Haerty and Singh 2006; Llopart 2012) and sterile introgression lines of tomato

(Guerrero et al. 2016) and *Drosophila* (Meiklejohn et al. 2014), but overexpression has also

been found in other sterile hybrids (Llopart 2012; Davis et al. 2015). The variation we and

others have found in hybrid regulatory phenotypes suggests that the mechanisms of disrupted

expression are complex, even within groups with relatively shallow divergence times (such as

rodents), and we need more data from diverse hybrid sterility systems to begin to understand

common drivers of asymmetric hybrid misexpression.

The downregulation we observed in early spermatogenesis in hybrid dwarf hamsters

could be due to impaired transcription factor binding with promoter or enhancer elements (Oka

et al. 2014; Guerrero et al. 2016) or disrupted epigenetic silencing. Disruption of epigenetic

regulation of gene expression has been increasingly linked to hybrid dysfunction in plants

(Shivaprasad et al. 2012; Lafon-Placette and Köhler 2015; Zhu et al. 2017), especially

polyploids (Paun et al. 2007), and may also contribute to hybrid male sterility in *Drosophila*

(Bayes and Malik 2009) and cattle x yak hybrids (Luo et al. 2022). At least one known

chromatin difference has been documented between parental dwarf hamster species on the X

chromosome (Gamperl et al. 1977; Haaf et al. 1987). However, it is unknown if genome-wide

epigenetic regulatory mechanisms are disrupted in hybrid dwarf hamster spermatogenesis, and

further work is needed to distinguish between potential mechanisms underlying the observed

genome-wide downregulation.

Spermatogenesis as a developmental process may be sensitive to disruption (*i.e.,*

"faster male" evolution; Wu and Davis 1993), but it remains an open question whether specific

stages of spermatogenesis, or developmental processes more broadly, may be more prone to

the accumulation of hybrid incompatibilities. In general, earlier developmental stages are

thought to be under greater pleiotropic constraint and less prone to disruption (Cutter and

Bundus 2020). With the progression of mouse spermatogenesis, pleiotropy decreases (as

approximated by increases in tissue specificity; Murat et al. 2023) and the rate of protein-

coding evolution increases (Larson et al. 2016; Kopania, Larson, et al. 2022; Murat et al. 2023),

which may make the later stages of spermatogenesis more prone to accumulating hybrid

incompatibilities. Indeed, we found fewer differentially expressed genes in sterile hybrids of

both dwarf hamsters and house mice during the early stages of spermatogenesis than in later

stages (Figures S7 and 8). Generally, there might be some tolerance for misregulation early in

spermatogenesis, since misregulation of some genes during early stages does not always

completely halt the progression of sperm development (Oka et al. 2010; Ishishita et al. 2015;

Mipam et al. 2023). However, the patterns we find in dwarf hamsters suggests that

spermatogenesis may be disrupted between zygotene and diplotene cell stages from early

misexpression. Dwarf hamster hybrid diplotene cell populations had X-linked and autosomal

gene expression profiles which more closely resemble parental leptotene/zygotene cell

populations than either diplotene or postmeiotic cell populations. Furthermore, differentially

expressed genes in hybrids during diplotene were enriched for genes associated with the GO

term for "negative regulation of cell proliferation" suggesting that there may be a regulatory

stalling of meiosis occurring in hybrid dwarf hamsters during early spermatogenesis. It's

unclear what underlying genomic mechanisms could result in this stalling, but it is possible that

this disruption could potentially act as a major contributor to hybrid sterility in this system.

Ultimately, we find that spermatogenesis is a complex and rapidly evolving developmental

program that may provide many potential avenues across its timeline for the evolution of hybrid

incompatibilities.

Abnormal sex chromosome expression in dwarf hamster hybrids is inconsistent with

disrupted MSCI

The sex chromosomes play a central role in speciation, an observation which has been

supported by both Haldane's rule (Haldane 1922) and the large X-effect on hybrid male sterility

(Coyne and Orr 1989). Misregulation of the X chromosome may contribute to hybrid sterility in

several species pairs (Davis et al. 2015; Morgan et al. 2020; Sánchez-Ramírez et al. 2021). The

X chromosome is transcriptionally repressed during routine spermatogenesis in many

organisms including eutherian mammals (McKee and Handel 1993), monotremes (Murat et al.

2023), *Drosophila* (Landeen et al. 2016), grasshoppers (Viera et al. 2021), mosquitos (Taxiarchi

et al. 2019), and nematodes (Rappaport et al. 2021). Because of the ubiquity of X chromosome

repression during spermatogenesis, disruption of transcriptional repression could be a

widespread regulatory phenotype in sterile hybrids (Lifschytz and Lindsley 1972; Larson et al.

2018). In sterile hybrid mice, disrupted X repression (disrupted MSCI) leads to the

overexpression of the normally silenced X chromosome, but not the autosomes, during

diplotene (Good et al. 2010; Bhattacharyya et al. 2013; Campbell et al. 2013; Turner and Harr

2014; Larson et al. 2017; Larson et al. 2022). We also found overexpression of the X

chromosome during diplotene in sterile hybrid dwarf hamsters, but in a manner inconsistent

with disrupted MSCI. Unlike in house mouse hybrids, both the X and autosomes are

overexpressed in dwarf hamster hybrid diplotene cells and the extent of X overexpression was

greater than what was observed in house mice, and importantly, more variable. In fact, some

dwarf hamster hybrids had wildly overexpressed X chromosomes while others appeared to

have properly silenced X chromosomes. Our expression correlation and gene set analyses of

hybrid diplotene cells provide additional evidence that the genes overexpressed in hybrid

hamster diplotene are different than those overexpressed in house mouse hybrids, sharing

more similarity to the earlier meiotic cell types than downstream postmeiotic cell types. Overall,

our results indicate fundamentally different patterns of X-linked overexpression in both

systems, with X-linked overexpression in dwarf hamster hybrids being inconsistent with

disrupted MSCI.

Much of what we know about the genomic architecture and the role of sex

chromosome misregulation in hybrid male sterility in mammals comes from decades of work

that have shown a major gene, *Prdm9*, and its X chromosome modulator, *Hstx2*, may be

responsible for most F1 hybrid male sterility in house mice (Forejt et al. 1991; Trachtulec et al.

1997; Mihola et al. 2009; Lustyk et al. 2019; Forejt et al. 2021). *Prdm9* directs the location of

double strand breaks during meiotic recombination (Mihola et al. 2009; Oliver et al. 2009;

Smagulova et al. 2016). In hybrid mice, divergence at *Prdm9* binding sites leads to asymmetric

double-stranded breaks and results in autosomal asynapsis, triggering Meiotic Silencing of

Unsynapsed Chromatin, shutting down transcription on asynapsed autosomes using the same

cellular machinery as MSCI (Turner 2015), and eventually meiotic arrest and cell death

(Bhattacharyya et al. 2013; Forejt et al. 2021). This process is associated with the disruption of

MSCI and a characteristic overexpression of the X chromosome during meiosis (Good et al.

2010; Bhattacharyya et al. 2013; Campbell et al. 2013; Turner and Harr 2014; Larson et al.

2017; Larson et al. 2022), but whether disrupted MSCI directly contributes to hybrid male

sterility or is simply a downstream consequence of *Prdm9* divergence is still uncertain (Forejt et

al. 2021).

Whether we should have expected patterns of disrupted sex chromosome expression

in sterile hybrid hamsters to be the same as house mice is unclear. The sex chromosomes in

pachytene cells of hybrid dwarf hamsters display normal yH2AFX staining (Ishishita et al. 2015;

Bikchurina et al. 2018), a key marker in MSCI (Abe et al. 2022), which may indicate that the

hybrid sex chromosomes are properly silenced. Additionally, autosomal asynapsis is rarely

observed in hybrid dwarf hamsters, and asynapsis is almost exclusive to the sex chromosomes

(Ishishita et al. 2015; Bikchurina et al. 2018). This contrasts *Prdm9*-mediated sterility in house

mice, where hybrid autosomes are often asynapsed and decorated with yH2AFX

(Bhattacharyya et al. 2013; Forejt et al. 2021). Mechanisms other than *Prdm9* may also disrupt

MSCI and result in sterility, such as macrosatellite copy number divergence (Bredemeyer et al.

2021) and X-autosome translocations that impair synapsis (Homolka et al. 2007), although

there is no evidence for X-autosome translocations between the two species of dwarf hamsters

(Moore et al. 2022). Thus, while MSCI may be a major target for the accumulation of

reproductive barriers between species in many mammalian systems, either through *Prdm9*

divergence or alternative mechanisms, our results suggest that sterility in dwarf hamsters has a

more composite regulatory basis.

Another mechanism often proposed to underlie mammalian male hybrid sterility,

especially in rodents, is divergence in the PAR between parental species. The PAR is the only

portion of the sex chromosomes which can synapse during spermatogenesis, and it is still

unclear if the regulation of the PAR is uniformly detached from MSCI across divergent

mammalian sex chromosome systems (Raudsepp and Chowdhary 2015). We find no evidence

that PAR-specific misregulation is associated with hybrid sterility in dwarf hamsters. PAR

genes are not silenced in hybrids or parents, a pattern that is inconsistent with MSCI that has

extended to the PAR due to XY asynapsis, and further, expression of PAR genes in hybrids

differs little from parental PAR expression. While we find no evidence that PAR misregulation

per se is associated with hybrid sterility in this system, we cannot rule out the possibility that

structural and sequence divergence between the PARs of *P. sungorus* and *P. campbelli* may

be associated with hybrid sterility. Structural and sequence divergence in the PAR has been

hypothesized to activate the meiotic spindle checkpoint by interfering with proper pairing of

sex chromosomes (Burgoyne et al. 2009; Dumont 2017). The PAR evolves rapidly in rodents

(White, Ikeda, et al. 2012; Raudsepp and Chowdhary 2015; Morgan et al. 2019), and this

elevated divergence may underlie sex chromosome asynapsis and apoptosis in several hybrid

mouse crosses (Matsuda et al. 1991; Oka et al. 2010; White, Stubbings, et al. 2012; Dumont

2017). Furthermore, divergence in the mouse PAR has been implicated in spermatogenic

defects in crosses where *Prdm9*-divergence is minimal, such as between closely related

subspecies (Dumont 2017) or in mice with genetically-modified *Prdm9* alleles (Davies et al.

2021). Meiosis is likely tolerant to some degree of divergence in the PAR (Morgan et al. 2019),

but exact limits are currently unknown. At this time, thorough analysis of structural and

sequence divergence between the PARs in dwarf hamsters is challenging as the PAR is

notoriously difficult to assemble (but see Kasahara et al. 2022), and there are annotation gaps

in the current assembly of the PAR in dwarf hamsters. In sum, we find no clear pattern of

regulatory disruption of PAR genes in sterile hybrid dwarf hamsters, though this result may

change pending further refinement of the PAR annotation.

Conclusions

Cell-specific approaches for quantifying regulatory phenotypes are powerful tools for

providing insight into the underlying mechanisms behind hybrid dysfunction (Hunnicutt et al.

2022), especially in systems where it remains difficult to interrogate the underlying genomic

architecture of these traits. Using a contrast of dwarf hamster and house mouse hybrids, we

have shown that transgressive overexpression is not an inevitable outcome of hybridization,

that hybrid incompatibilities are likely to arise during multiple stages of spermatogenesis, and

that disrupted sex chromosome silencing does not appear to play an equal role in sterility

between these two systems. Both the regulatory phenotypes we observed here and

histological evidence from other studies (Ishishita et al. 2015; Bikchurina et al. 2018) suggest

that several reproductive barriers are acting during spermatogenesis in dwarf hamster hybrids.

It has become increasingly apparent as more study systems are investigated that the genetic

basis of postzygotic species barriers are often complex and polymorphic (Cutter 2012;

Coughlan and Matute 2020), and implementing approaches which account for the

developmental complexities of hybrid dysfunction, as we have done here, will allow us to make

further advances in understanding the processes of speciation.

MATERIALS AND METHODS

Hamster crosses and male reproductive phenotypes

We used wild-derived colonies of two sister species of dwarf hamster, *P. sungorus* and

P. campbelli, established by Kathy Wynne-Edwards (Scribner and Wynne-Edwards 1994) and

housed at the University of Montana. Both species were maintained as outbred colonies, but

inbreeding levels of these closed colonies are still high (Brekke et al. 2018). We used males

from both parent species and male F1 hybrid offspring from crosses of female *P. campbelli*

with male *P. sungorus*. We weaned males in same-sex sibling groups between 17 and 21 dpp

and housed them individually at 45 dpp. We euthanized reproductively mature males using

carbon dioxide followed by cervical dislocation between 59 - 200 dpp (Table S1). All animal

use was approved by the University of Montana (IACUC protocols 050-16JGDBS & 035-

19JGDBS).

We measured several fertility metrics for parent species and hybrid males including

paired testes weight, paired seminal vesicle weight, normalized sperm counts, and sperm

motility (Good et al. 2008). Paired testes weight and paired seminal vesicle weight were

correlated with body weight (paired testes weight Pearson's $r(29)$ = 0.47, p = 0.007; paired

seminal vesicle weight Pearson's $r(23)$ = 0.56, p = 0.003), so we standardized both metrics

relative to body weight. We calculated sperm count by isolating sperm from caudal

epididymides diced in 1 ml of Dulbecco's PBS (Sigma) and incubated at 37°C for 10 minutes.

We quantified sperm motility (proportion of motile sperm in a 5 µl suspension) and sperm count

(number of sperm with head and tail in a heat shocked 5 µl suspension) across a fixed area on

a Makler counting chamber. We performed statistical comparisons of fertility phenotypes in R

v.4.3.1, and we used the FSA package v.0.9.4 for the Kruskal-Wallis and Dunn's tests (Ogle

and Ogle 2017).

Isolation of enriched cell populations from hamster testes

To investigate the regulatory dynamics of the sex chromosomes during

spermatogenesis, we isolated four spermatogenic cell populations from whole testes using

FACS: spermatogonia, leptotene/zygotene spermatocytes, diplotene spermatocytes, and

round spermatids. Briefly, we disassociated a single testis per male following a published

protocol originally developed for house mice (Getun et al. 2011) with modifications

(github.com/goodest-goodlab/good-protocols/tree/main/protocols/FACS, last accessed June

16, 2021). We doubled the volumes of all reagents to account for the increased mass of testes

in dwarf hamsters relative to house mice. We isolated cell populations based on size,

granularity, and fluorescence on a FACSAria IIu cell sorter (BD Biosciences) at the University of

Montana Center for Environmental Health Sciences Fluorescence Cytometry Core. For each

sorted cell population, we extracted RNA using RNeasy kits (Qiagen) following protocols for

Purification of Total RNA from Animal Cells. We quantified sample RNA quantity and quality

(RNA integrity number > 7) on a Tapestation 2200 (Agilent) at the University of Montana

genomics core. RNA libraries were prepared by Novogene and sequenced on Illumina

NovaSeq 6000s (paired end, 150 bp). Six samples (distributed across different species and cell

types) had low RNA concentrations, and for these samples we used Novogene's low input

RNA library preparation (Table S2). MDS plots indicated no severe library batch effects

between samples from different library preparations (Figure S1) so we included all samples

from both libraries in subsequent analyses.

Read processing and mapping

We sequenced RNA from each cell population for three to five individuals of each

parent species and F1 hybrids generating an average of ~27.5 million read pairs per individual

(Table S2). We trimmed reads using Trimmomatic v.0.39 (Bolger et al. 2014) to remove low

quality bases from the first and last 5 bp of each read and bases with an average Phred score

of less than 15 across a 4 bp sliding window and only retained reads of at least 36 bp. We next

used an approach (based on the modtools pipeline) which maps reads from each sample to

pseudogenomes for both parent species (described below) to obtain a merged output

alignment file in order to alleviate reference bias associated with mapping hybrids to only a

single reference genome (Holt et al. 2013; Huang et al. 2014). For this approach, we mapped

reads for each individual to both a *P. sungorus* pseudogenome and a *P. campbelli*

pseudogenome with Hisat v.2.2.0 (Kim et al. 2019) with default settings and retaining at most

100 distinct, primary alignments. We generated the *P. sungorus* pseudogenome by mapping

RNASeq reads from a male *P. sungorus* individual (30.6 million total read pairs; NCBI SRA:

SRR17223284; Moore et al. 2022) to the *P. sungorus* reference genome with bwa-mem v.2.2.1

(Vasimuddin et al. 2019), and the *P. campbelli* pseudogenome by mapping female *P. campbelli*

whole genome sequencing reads (average coverage: 33x; NCBI SRA: SRR17223279; Moore et

al. 2022) to the *P. sungorus* reference genome. Because our *P. sungorus* pseudogenome was

based on a male hamster and the reference genome on a female hamster, we excluded reads

mapping to the PAR because sequences mapping to this region could have originated from

either the X or Y chromosomes and interfered with subsequent variant calling. Following

mapping, we used GATK v.4.2.5.0 HaplotypeCaller (-ERC GVCF) to call SNPs then performed

genotyping with genotypeGVCFs. We hard-filtered our SNPs (--mask-extension 5 "QD < 2.0"

"FS > 60.0" "MQ < 40.0" "QUAL < 30.0" "DP < 10" "DP > 150") and restricted SNPs to biallelic

loci. Finally, we incorporated filtered SNPs back into the *P. sungorus* reference genome with

FastaAlternateReferenceMaker to create the *P. sungorus* and *P. campbelli* pseudoreferences.

For our RNASeq data, we appended query hit indexes to resulting alignment files using

hisat2Tophat.py (https://github.com/goodest-goodlab/pseudo-it/tree/master/helper-

scripts/hisat2Tophat.py, last accessed March 8th, 2022) to maintain compatibility with the

modtools pipeline. We used our vcfs (above) to generate a mod-file for both species with

vcf2mod from Lapels v.1.1.1 to convert alignments to the *P. sungorus* reference genome, and

Suspenders 0.2.6 to merge alignments while retaining the highest quality alignment per read

(Holt et al. 2013; Huang et al. 2014). We used featureCounts v.2.0.1 (Liao et al. 2014) to

estimate counts of read pairs that aligned to the same chromosome (-B and -C) and retained

only singly-mapped reads. Summaries of properly mapped reads for each sample can be

found in Table S2.

We sought to compare the gene expression phenotypes observed in dwarf hamsters to

those previously documented in house mice using published RNASeq data for the same four

spermatogenic cell types of two subspecies of house mouse and their sterile F1 hybrids

(Larson et al. 2017; Hunnicutt et al. 2022). These studies examined two subspecies of house

mice, *Mus musculus musculus* (intra-subspecific F1 males between wild-derived inbred strains

PWK/PhJ♀ and CZECHII/EiJ♂) and *M. m. domesticus* (intra-subspecific F1 males between

wild-derived inbred strains WSB/EiJ♀ and LEWES/EiJ♂) and their sterile (PWK♀ x LEWES♂)

F1 hybrids for disrupted gene expression across spermatogenesis following the same FACS

protocols implemented in this study. For all comparisons between house mice and dwarf

hamsters, we used read count files generated previously for house mice (Hunnicutt et al. 2022)

and performed all subsequent analyses in parallel for both systems.

Gene expression pre-processing

Following read processing and mapping, we conducted all analyses in R v.4.3.1. We

classified genes as "expressed" if genes had a minimum of one Fragment Per Kilobase of exon

per Million mapped reads (FPKM) in at least three samples, resulting in 21,077 expressed

genes across the whole dwarf hamster dataset and 21,212 expressed genes across the house

mouse dataset. We also identified sets of genes "induced" in a given cell-type defined as

genes with a median expression in a given cell population (normalized FPKM) greater than two

times its median expression across all other sorted cell populations (following Kousathanas et

al. 2014). We calculated normalized FPKM values by adjusting the sum of squares to equal one

using the R package vegan v.2.6-4 (Oksanen et al. 2013). We conducted expression analyses

using edgeR v.3.42.4 (Robinson et al. 2010) and normalized the data using the scaling factor

method (Anders and Huber 2010).

We assessed cell population purity using a panel of marker genes specific to the four

cell populations targeted by our FACS protocol and present in only a single copy in the *P.*

sungorus annotation. Spermatogonia markers included *Dmrt1* (Raymond et al. 2000) and *Hells*

(Green et al. 2018). Leptotene/zygotene markers included *Ccnb1ip1* and *Adad2* (Hermann et al.

2018). Diplotene was characterized by *Aurka* and *Tank* expression (Murat et al. 2023) and

round spermatids by *Cabyr* and *Acrv1* expression (Green et al. 2018). To estimate relative

purity, we quantified mean marker gene expression across replicates for a given cell population

for both parent species. A cell population was considered pure if it had higher marker gene

expression than other populations isolated by our FACS protocol and if X-linked gene

expression matched the expected regulatory dynamics (*i.e.,* active vs. silenced (MSCI) vs.

repressed (PSCR); Handel 2004; Namekawa et al. 2006). We examined expression patterns

across cell populations for all genes, autosomal genes, and X-linked genes using MDS plots

generated with the plotMDS function in limma v.3.56.2 (Ritchie et al. 2015) and heatmaps using

ComplexHeatmap v.2.16.0 (Gu et al. 2016). MDS plots used the top 500 genes with the largest

fold change difference between samples.

Differential gene expression analysis

We assessed differential gene expression by contrasting hybrids and each parent

species for all cell populations. We fit the expression data for dwarf hamsters and house mice

separately with negative binomial generalized linear models with Cox-Reid tagwise dispersion

estimates and adjusted P-values to a false discovery rate (FDR) of 5% (Benjamini and

Hochberg 1995). We quantified the biological coefficient of variation (BCV), a metric

representing the variation in gene expression among replicates (McCarthy et al. 2012), for each

dataset. Additionally, we calculated the BCV of just parental males or hybrid samples for each

species for the first three cell populations to examine whether dwarf hamster hybrids exhibited

more variability in expression than house mouse hybrids and parental dwarf hamsters. For our

differential expression analyses, we contrasted expression between hybrids and parents so

that a positive log fold-change (logFC) indicated overexpression in sterile males. Unless

otherwise specified, results discussed in the main text contrast hybrid offspring to the parent

with the same X chromosome (*P. campbelli* for dwarf hamster F1 hybrids and *M. m. musculus*

for house mouse F1 hybrids). Contrasts between F1 hybrids and *P. sungorus* or *M. m.*

domesticus are provided in supplementary material (Figures S6, 9, and 10).

We tested for significant differences in the number of under and overexpressed DE

genes within a stage for both house mouse and hamster hybrids using X^2 tests with chisq.tests

in R and used FDR correction for multiple comparisons. We also tested for differences in the

magnitude of misexpression between mouse and hamster hybrids for each stage by

comparing the distributions of the logFC of DE genes between hybrids and parent species

between mice and hamsters using Wilcoxon signed-rank tests and FDR correction. To

characterize hybrid diplotene expression in both house mice and dwarf hamsters, we used two

approaches. First, we calculated Pearson's correlation coefficient *(r)* between average

normalized hybrid diplotene expression and the average normalized expression in each

parental cell type. We corrected p-values for each correlation with FDR. We generated

bootstrap values for each correlation coefficient by randomly sampling the expression matrices

with replacement for each sample type for 1000 replicates. Second, we compared the gene

sets that escaped MSCI in each species with gene sets that characterize stage-specific

expression in parent species. For this analysis, we defined sets of overexpressed X-linked

diplotene genes as genes with expression (normalized FPKM) in hybrids that was in the top

10% of X-linked genes in parental diplotene samples (*i.e.,* genes that normally escape MSCI).

We then compared these sets of overexpressed hybrid diplotene genes to genes "induced" in

each parental stage for each species. We used DAVID v.2021 (Sherman et al. 2022) to perform

gene ontology (GO) analysis to identify GO terms overrepresented in differentially expressed

genes in hybrid samples across each cell population. We only included the longest genes

associated with mouse orthologs in our GO analysis, and for our background gene lists, we

used genes that were "expressed" in both hybrid and parent species in a given stage.

Biological Process GO terms with FDR below 0.05 were retained as significant. Finally, to test

whether specific chromosomes were enriched for differentially expressed genes for a given

stage, we performed hypergeometric tests on the number of differentially expressed genes on

a given chromosome with phyper and adjusted P-values to an FDR of 5%.

Characterizing the behavior of PAR genes in dwarf hamsters

We sought to characterize the regulatory behavior of PAR genes in dwarf hamster

parent species and hybrids to determine if (1) PAR genes are normally silenced in parent

species and (2) if PAR genes were overexpressed in hybrids (consistent with an extension of

761 MSCI to the PAR). The PAR on the *P. sungorus* X chromosome is on the distal arm of the X

762 chromosome from around 115,350,000-119,112,095 bp (Moore et al. 2022). There are 15

763 annotated *P. sungorus* genes in this region (Table S3), which is comparable to the latest PAR

764 assembly in C57BL/6J house mice (Kasahara et al. 2022). Six of these are orthologous to

765 annotated genes in mice (*Tppp2*, *Gprin1*, *Ndrg2*, *Kcnip4*, *Ndrg2*, and *Hs6st3*), but they are not

766 located in the mouse PAR (Kasahara et al. 2022). Only seven of the annotated genes in the

767 PAR were expressed in more than three replicates across all samples (Psun_G000022875,

768 Psun_G000022880, Psun_G000022883, Psun_G000022886, *Tppp2*, *Gprin1*, and *Ndrg2*). For

769 these genes, we assessed whether these genes were consistently expressed or silenced in

770 parent species at any cell stage and whether any genes were differentially expressed between

771 hybrids and either parent species at any cell stage.

Acknowledgements:

This work was supported by a National Science Foundation Graduate Research Fellowship to KEH (DGE-2034612) and the following grants awarded to KEH: Sigma Xi Grant-in-aid of Research, American Society of Mammalogists Grant-in-Aid of Research Grant, and a Society for the Study of Evolution R.C. Lewontin Early Award. Additionally, this work was supported by grants from the Eunice Kennedy Shriver National Institute of Child Health and Human Development of the National Institutes of Health (R01-HD073439, R01-HD094787) to JMG, and the National Science Foundation to ELL (DEB-2012041). We would like to thank members of the Larson, Velotta, and Good Labs for feedback on this project, Ivan Kovanda for support using the University of Denver Research Data Analysis Cluster, and Pamela K. Shaw and the UM Fluorescence Cytometry Core supported by an Institutional Development Award from the NIGMS (P30GM103338). Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation or the National Institutes of Health.

Author contributions:

KEH, JMG, and ELL conceived of the study. CC, SK, ELL, and KEH conducted lab work. KEH conducted the analyses. KEH and ELL wrote the manuscript with input from ECM, CC, SK, and JMG.

Data accessibility:

The data reported in this paper are available through the National Center for Biotechnology Information under accession number PRJNA1024468. The code used for the analyses is available from GitHub (https://github.com/KelsieHunnicutt/dwarf_hamster_hybrids).

LITERATURE CITED

Abe H, Yeh Y-H, Munakata Y, Ishiguro K-I, Andreassen PR, Namekawa SH. 2022. Active DNA

damage response signaling initiates and maintains meiotic sex chromosome inactivation.

Nat. Commun. 13:7212.

Anders S, Huber W. 2010. Differential expression analysis for sequence count data. Nature

Precedings:1–1.

Barreto FS, Pereira RJ, Burton RS. 2015. Hybrid dysfunction and physiological compensation

in gene expression. Mol. Biol. Evol. 32:613–622.

Bayes JJ, Malik HS. 2009. Altered heterochromatin binding by a hybrid sterility protein in

Drosophila sibling species. Science 326:1538–1541.

Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: a practical and powerful

approach to multiple testing. J. R. Stat. Soc. 57:289–300.

Bhattacharyya T, Gregorova S, Mihola O, Anger M, Sebestova J, Denny P, Simecek P, Forejt J.

2013. Mechanistic basis of infertility of mouse intersubspecific hybrids. Proc. Natl. Acad.

Sci. U. S. A. 110:E468–E477.

Bikchurina TI, Tishakova KV, Kizilova EA, Romanenko SA, Serdyukova NA, Torgasheva AA,

Borodin PM. 2018. Chromosome synapsis and recombination in male-sterile and female-

fertile interspecies hybrids of the dwarf hamsters (Phodopus, Cricetidae). Genes 9:227.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence

data. Bioinformatics 30:2114–2120.

Bredemeyer KR, Seabury CM, Stickney MJ, McCarrey JR, vonHoldt BM, Murphy WJ. 2021.

Rapid macrosatellite evolution promotes X-linked hybrid male sterility in a feline

interspecies cross. Mol. Biol. Evol. 38:5588–5609.

Brekke TD, Good JM. 2014. Parent-of-origin growth effects and the evolution of hybrid

inviability in dwarf hamsters. Evolution 68:3134–3148.

Brekke TD, Moore EC, Campbell-Staton SC, Callahan CM, Cheviron ZA, Good JM. 2021. X

chromosome-dependent disruption of placental regulatory networks in hybrid dwarf

hamsters. Genetics 218:iyab043.

Brekke TD, Steele KA, Mulley JF. 2018. Inbred or outbred? Genetic diversity in laboratory

rodent colonies. G3 8:679–686.

Burgoyne PS. 1982. Genetic homology and crossing over in the X and Y chromosomes of

mammals. Hum. Genet. 61:85–90.

Burgoyne PS, Mahadevaiah SK, Turner JMA. 2009. The consequences of asynapsis for

mammalian meiosis. Nat. Rev. Genet. 10:207–216.

Campbell P, Good JM, Nachman MW. 2013. Meiotic sex chromosome inactivation is disrupted

in sterile hybrid male house mice. Genetics 193:819–828.

Civetta A. 2016. Misregulation of gene expression and sterility in interspecies hybrids: causal

links and alternative hypotheses. J. Mol. Evol. 82:176–182.

Coughlan JM, Matute DR. 2020. The importance of intrinsic postzygotic barriers throughout

the speciation process. Philos. Trans. R. Soc. Lond. B Biol. Sci. 375:20190533.

Coyne JA, Orr HA. 1989. Two rules of speciation. In: Otte D, Endler JA, editors. Speciation and

its consequences. Sunderland, MA: Sinauer Associates. p. 180–207.

Cutter AD. 2012. The polymorphic prelude to Bateson-Dobzhansky-Muller incompatibilities.

Trends Ecol. Evol. 27:209–218.

Cutter AD, Bundus JD. 2020. Speciation and the developmental alarm clock. Elife 9:e56276.

Davies B, Gupta Hinch A, Cebrian-Serrano A, Alghadban S, Becker PW, Biggs D, Hernandez-

Pliego P, Preece C, Moralli D, Zhang G, et al. 2021. Altering the binding properties of

PRDM9 partially restores fertility across the species boundary. Mol. Biol. Evol. 38:5555–

5562.

Davies B, Hatton E, Altemose N, Hussin JG, Pratto F, Zhang G, Hinch AG, Moralli D, Biggs D,

Diaz R, et al. 2016. Re-engineering the zinc fingers of PRDM9 reverses hybrid sterility in

mice. Nature 530:171–176.

Davis BW, Seabury CM, Brashear WA, Li G, Roelke-Parker M, Murphy WJ. 2015. Mechanisms

underlying mammalian hybrid sterility in two feline interspecies models. Mol. Biol. Evol.

32:2534–2546.

Dumont BL. 2017. Meiotic consequences of genetic divergence across the murine

pseudoautosomal region. Genetics 205:1089–1100.

Ellis N, Goodfellow PN. 1989. The mammalian pseudoautosomal region. Trends Genet. 5:406–

410.

Forejt J, Jansa P, Parvanov E. 2021. Hybrid sterility genes in mice (Mus musculus): a peculiar

case of PRDM9 incompatibility. Trends Genet. 37:1095–1108.

Forejt J, Vincek V, Klein J, Lehrach H, Loudová-Micková M. 1991. Genetic mapping of the t-

complex region on mouse chromosome 17 including the Hybrid sterility-1 gene. Mamm.

Genome 1:84–91.

Gamperl R, Vistorin G, Rosenkranz W. 1977. New observations on the karyotype of the

860   Djungarian hamster, Phodopus sungorus. Experientia 33:1020–1021.

861   Getun IV, Torres B, Bois PRJ. 2011. Flow cytometry purification of mouse meiotic cells. J. Vis.

862   Exp.:e2602.

863   Good JM, Giger T, Dean MD, Nachman MW. 2010. Widespread over-expression of the X

864   chromosome in sterile F1 hybrid mice. PLoS Genet. 6:e1001148.

865   Good JM, Handel MA, Nachman MW. 2008. Asymmetry and polymorphism of hybrid male

866   sterility during the early stages of speciation in house mice. Evolution 62:50–65.

867   Green CD, Ma Q, Manske GL, Shami AN, Zheng X, Marini S, Moritz L, Sultan C, Gurczynski SJ,

868   Moore BB, et al. 2018. A comprehensive roadmap of murine spermatogenesis defined by

869   single-cell RNA-Seq. Dev. Cell 46:651–667.e10.

870   Guerrero RF, Posto AL, Moyle LC, Hahn MW. 2016. Genome-wide patterns of regulatory

871   divergence revealed by introgression lines. Evolution 70:696–706.

872   Gu Z, Eils R, Schlesner M. 2016. Complex heatmaps reveal patterns and correlations in

873   multidimensional genomic data. Bioinformatics 32:2847–2849.

874   Haaf T, Weis H, Schmid M. 1987. A comparative cytogenetic study on the mitotic and meiotic

875   chromosomes in hamster species of the genus Phodopus (Rodentia, Cricetinae).

876   Zeitschrift für Säugetierkunde 52:281–290.

877   Haerty W, Singh RS. 2006. Gene regulation divergence is a major contributor to the evolution

878   of Dobzhansky-Muller incompatibilies between species of Drosophila. Mol. Biol. Evol.

23:1707–1714.

Haldane JBS. 1922. Sex ratio and unisexual sterility in hybrid animals. J. Genet. 12:101–109.

Handel MA. 2004. The XY body: a specialized meiotic chromatin domain. Exp. Cell Res.

296:57–63.

Hermann BP, Cheng K, Singh A, Roa-De La Cruz L, Mutoji KN, Chen I-C, Gildersleeve H, Lehle

JD, Mayo M, Westernströer B, et al. 2018. The mammalian spermatogenesis single-cell

transcriptome, from spermatogonial stem cells to spermatids. Cell Rep. 25:1650–1667.e8.

Holt J, Huang S, McMillan L, Wang W. 2013. Read annotation pipeline for high-throughput

sequencing data. In: Proceedings of the International Conference on Bioinformatics,

Computational Biology and Biomedical Informatics - BCB'13. New York, New York, USA:

ACM Press. p. 605–612.

Homolka D, Ivanek R, Capkova J, Jansa P, Forejt J. 2007. Chromosomal rearrangement

interferes with meiotic X chromosome inactivation. Genome Res. 17:1431–1437.

Huang S, Holt J, Kao C-Y, McMillan L, Wang W. 2014. A novel multi-alignment pipeline for

high-throughput sequencing data. Database 2014:bau057.

Hunnicutt KE, Good JM, Larson EL. 2022. Unraveling patterns of disrupted gene expression

across a complex tissue. Evolution 76:275–291.

Ishishita S, Tsuboi K, Ohishi N, Tsuchiya K, Matsuda Y. 2015. Abnormal pairing of X and Y sex

chromosomes during meiosis I in interspecific hybrids of Phodopus campbelli and *P.*

sungorus. Sci. Rep. 5:9435.

Kasahara T, Mekada K, Abe K, Ashworth A, Kato T. 2022. Complete sequencing of the mouse

pseudoautosomal region, the most rapidly evolving "chromosome."

bioRxiv:2022.03.26.485930.

Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019. Graph-based genome alignment and

genotyping with HISAT2 and HISAT-genotype. Nat. Biotechnol. 37:907–915.

Kopania EEK, Larson EL, Callahan C, Keeble S, Good JM. 2022. Molecular evolution across

mouse spermatogenesis. Mol. Biol. Evol. 39:msac023.

Kopania EEK, Watson EM, Rathje CC, Skinner BM, Ellis PJI, Larson EL, Good JM. 2022. The

contribution of sex chromosome conflict to disrupted spermatogenesis in hybrid house

mice. Genetics 222:iyac151.

Kousathanas A, Halligan DL, Keightley PD. 2014. Faster-X adaptive protein evolution in house

mice. Genetics 196:1131–1143.

Lafon-Placette C, Köhler C. 2015. Epigenetic mechanisms of postzygotic reproductive isolation

in plants. Curr. Opin. Plant Biol. 23:39–44.

Landeen EL, Muirhead CA, Wright L, Meiklejohn CD, Presgraves DC. 2016. Sex chromosome-

wide transcriptional suppression and compensatory cis-regulatory evolution mediate gene

expression in the Drosophila male germline. PLoS Biol. 14:e1002499.

Larson EL, Keeble S, Vanderpool D, Dean MD, Good JM. 2017. The composite regulatory

basis of the large X-effect in mouse speciation. Mol. Biol. Evol. 34:282–295.

Larson EL, Kopania EEK, Good JM. 2018. Spermatogenesis and the evolution of mammalian

sex chromosomes. Trends Genet. 34:722–732.

Larson EL, Kopania EEK, Hunnicutt KE, Vanderpool D, Keeble S, Good JM. 2022. Stage-

specific disruption of X chromosome expression during spermatogenesis in sterile house

mouse hybrids. G3 12:jkab407.

Larson EL, Vanderpool D, Keeble S, Zhou M, Sarver BAJ, Smith AD, Dean MD, Good JM.

2016. Contrasting levels of molecular evolution on the mouse X chromosome. Genetics

203:1841–1857.

Liao Y, Smyth GK, Shi W. 2014. featureCounts: an efficient general purpose program for

assigning sequence reads to genomic features. Bioinformatics 30:923–930.

Lifschytz E, Lindsley DL. 1972. The role of X-chromosome inactivation during

spermatogenesis. Proc. Natl. Acad. Sci. U. S. A. 69:182–186.

Llopart A. 2012. The rapid evolution of X-linked male-biased gene expression and the large-X

effect in Drosophila yakuba, D. santomea, and their hybrids. Mol. Biol. Evol. 29:3873–

3886.

Luo H, Mipam T, Wu S, Xu C, Yi C, Zhao W, Chai Z, Chen X, Wu Z, Wang J, et al. 2022. DNA

methylome of primary spermatocyte reveals epigenetic dysregulation associated with male

sterility of cattleyak. Theriogenology 191:153–167.

Lustyk D, Kinský S, Ullrich KK, Yancoskie M, Kašíková L, Gergelits V, Sedlacek R, Chan YF,

Odenthal-Hesse L, Forejt J, et al. 2019. Genomic structure of Hstx2 modifier of Prdm9-

dependent hybrid male sterility in mice. Genetics 213:1047–1063.

Mack KL, Campbell P, Nachman MW. 2016. Gene regulation and speciation in house mice.

Genome Res. 26:451–461.

Mack KL, Nachman MW. 2017. Gene regulation and speciation. Trends Genet. 33:68–80.

Matsuda Y, Hirobe T, Chapman VM. 1991. Genetic basis of X-Y chromosome dissociation and

male sterility in interspecific hybrids. Proc. Natl. Acad. Sci. U. S. A. 88:4850–4854.

McCarthy DJ, Chen Y, Smyth GK. 2012. Differential expression analysis of multifactor RNA-

Seq experiments with respect to biological variation. Nucleic Acids Res. 40:4288–4297.

McKee BD, Handel MA. 1993. Sex chromosomes, recombination, and chromatin conformation.

Chromosoma 102:71–80.

McManus CJ, Coolon JD, Duff MO, Eipper-Mains J, Graveley BR, Wittkopp PJ. 2010.

Regulatory divergence in Drosophila revealed by mRNA-seq. Genome Res. 20:816–825.

Meiklejohn CD, Coolon JD, Hartl DL, Wittkopp PJ. 2014. The roles of cis- and trans-regulation

in the evolution of regulatory incompatibilities and sexually dimorphic gene expression.

Genome Res. 24:84–95.

Michalak P, Noor MAF. 2003. Genome-wide patterns of expression in Drosophila pure species

and hybrid males. Mol. Biol. Evol. 20:1070–1076.

Mihola O, Trachtulec Z, Vlcek C, Schimenti JC, Forejt J. 2009. A mouse speciation gene

encodes a meiotic histone H3 methyltransferase. Science 323:373–375.

Mipam T, Chen X, Zhao W, Zhang P, Chai Z, Yue B, Luo H, Wang J, Wang H, Wu Z, et al.

2023. Single-cell transcriptome analysis and in vitro differentiation of testicular cells reveal

novel insights into male sterility of the interspecific hybrid cattle-yak. BMC Genomics

24:149.

Montgomery SH, Mank JE. 2016. Inferring regulatory change from gene expression: the

confounding effects of tissue scaling. Mol. Ecol. 25:5114–5128.

Moore EC, Thomas GWC, Mortimer S, Kopania EEK, Hunnicutt KE, Clare-Salzler ZJ, Larson

EL, Good JM. 2022. The evolution of widespread recombination suppression on the dwarf

hamster (Phodopus) X chromosome. Genome Biol. Evol. 14:evac080.

Morgan AP, Bell TA, Crowley JJ, Pardo-Manuel de Villena F. 2019. Instability of the

pseudoautosomal boundary in house mice. Genetics 212:469–487.

Morgan K, Harr B, White MA, Payseur BA, Turner LM. 2020. Disrupted gene networks in

subfertile hybrid house mice. Mol. Biol. Evol. 37:1547–1562.

Mugal CF, Wang M, Backström N, Wheatcroft D, Ålund M, Sémon M, McFarlane SE, Dutoit L,

Qvarnström A, Ellegren H. 2020. Tissue-specific patterns of regulatory changes underlying

gene expression differences among Ficedula flycatchers and their naturally occurring F1

hybrids. Genome Res. 30:1727–1739.

Murat F, Mbengue N, Winge SB, Trefzer T, Leushkin E, Sepp M, Cardoso-Moreira M, Schmidt

J, Schneider C, Mößinger K, et al. 2023. The molecular evolution of spermatogenesis

across mammals. Nature 613:308–316.

Namekawa SH, Park PJ, Zhang L-F, Shima JE, McCarrey JR, Griswold MD, Lee JT. 2006.

Postmeiotic sex chromatin in the male germline of mice. Curr. Biol. 16:660–667.

Neumann K, Michaux J, Lebedev V, Yigit N, Colak E, Ivanova N, Poltoraus A, Surov A, Markov

G, Maak S, et al. 2006. Molecular phylogeny of the Cricetinae subfamily based on the

mitochondrial cytochrome b and 12S rRNA genes and the nuclear vWF gene. Mol.

Phylogenet. Evol. 39:135–148.

Ogle D, Ogle MD. 2017. Package "FSA." CRAN Repos:1–206.

Oka A, Mita A, Takada Y, Koseki H, Shiroishi T. 2010. Reproductive isolation in hybrid mice

due to spermatogenesis defects at three meiotic stages. Genetics 186:339–351.

Oka A, Shiroishi T. 2014. Regulatory divergence of X-linked genes and hybrid male sterility in

mice. Genes Genet. Syst. 89:99–108.

Oka A, Takada T, Fujisawa H, Shiroishi T. 2014. Evolutionarily diverged regulation of X-

chromosomal genes as a primal event in mouse reproductive isolation. PLoS Genet.

10:e1004301.

Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'hara RB, Simpson GL, Solymos

P, Stevens MHH, Wagner H, et al. 2013. Package "vegan." Community ecology package,

version 2:1–295.

Oliver PL, Goodstadt L, Bayes JJ, Birtle Z, Roach KC, Phadnis N, Beatson SA, Lunter G, Malik

HS, Ponting CP. 2009. Accelerated evolution of the Prdm9 speciation gene across diverse

metazoan taxa. PLoS Genet. 5:e1000753.

Ortíz-Barrientos D, Counterman BA, Noor MAF. 2007. Gene expression divergence and the

origin of hybrid dysfunctions. Genetica 129:71–81.

Paun O, Fay MF, Soltis DE, Chase MW. 2007. Genetic and epigenetic alterations after

hybridization and genome doubling. Taxon 56:649–656.

Presgraves DC. 2008. Sex chromosomes and speciation in Drosophila. Trends Genet. 24:336–

343.

Ramm SA, Schärer L. 2014. The evolutionary ecology of testicular function: size isn't

everything. Biol. Rev. Camb. Philos. Soc. 89:874–888.

Rappaport Y, Achache H, Falk R, Murik O, Ram O, Tzur YB. 2021. Bisection of the X

chromosome disrupts the initiation of chromosome silencing during meiosis in

Caenorhabditis elegans. Nat. Commun. 12:4802.

Raudsepp T, Chowdhary BP. 2015. The eutherian pseudoautosomal region. Cytogenet.

Genome Res. 147:81–94.

Raymond CS, Murphy MW, O'Sullivan MG, Bardwell VJ, Zarkower D. 2000. Dmrt1, a gene

related to worm and fly sexual regulators, is required for mammalian testis differentiation.

Genes Dev. 14:2587–2595.

Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK. 2015. limma powers

differential expression analyses for RNA-sequencing and microarray studies. Nucleic Acids

Res. 43:e47.

Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a Bioconductor package for differential

expression analysis of digital gene expression data. Bioinformatics 26:139–140.

Sánchez-Ramírez S, Weiss JG, Thomas CG, Cutter AD. 2021. Widespread misregulation of

inter-species hybrid transcriptomes due to sex-specific and sex-chromosome regulatory

evolution. PLoS Genet. 17:e1009409.

Scribner SJ, Wynne-Edwards KE. 1994. Disruption of body temperature and behavior rhythms

during reproduction in dwarf hamsters (Phodopus). Physiol. Behav. 55:361–369.

Sherman BT, Hao M, Qiu J, Jiao X, Baseler MW, Lane HC, Imamichi T, Chang W. 2022. DAVID:

a web server for functional enrichment analysis and functional annotation of gene lists

(2021 update). Nucleic Acids Res. 50:W216–W221.

Shivaprasad PV, Dunn RM, Santos BA, Bassett A, Baulcombe DC. 2012. Extraordinary

transgressive phenotypes of hybrid tomato are influenced by epigenetics and small

silencing RNAs. EMBO J. 31:257–266.

Smagulova F, Brick K, Pu Y, Camerini-Otero RD, Petukhova GV. 2016. The evolutionary

turnover of recombination hot spots contributes to speciation in mice. Genes Dev. 30:266–

280.

Swanson MT, Oliveros CH, Esselstyn JA. 2019. A phylogenomic rodent tree reveals the

repeated evolution of masseter architectures. Proc. Biol. Sci. 286:20190672.

Taxiarchi C, Kranjc N, Kriezis A, Kyrou K, Bernardini F, Russell S, Nolan T, Crisanti A, Galizi R.

2019. High-resolution transcriptional profiling of Anopheles gambiae spermatogenesis

reveals mechanisms of sex chromosome regulation. Sci. Rep. 9:14841.

Trachtulec Z, Mnuková-Fajdelová M, Hamvas RM, Gregorová S, Mayer WE, Lehrach HR,

Vincek V, Forejt J, Klein J. 1997. Isolation of candidate hybrid sterility 1 genes by cDNA

selection in a 1.1 megabase pair region on mouse chromosome 17. Mamm. Genome

8:312–316.

Turner JMA. 2015. Meiotic silencing in mammals. Annu. Rev. Genet. 49:395–412.

Turner LM, Harr B. 2014. Genome-wide mapping in a house mouse hybrid zone reveals hybrid

sterility loci and Dobzhansky-Muller interactions. Elife 3:e02504.

Turner LM, White MA, Tautz D, Payseur BA. 2014. Genomic networks of hybrid sterility. PLoS

Genet. 10:e1004162.

Vasimuddin M, Misra S, Li H, Aluru S. 2019. Efficient architecture-aware acceleration of BWA-

MEM for multicore systems. In: 2019 IEEE International Parallel and Distributed Processing

Symposium (IPDPS). IEEE. p. 314–324.

Viera A, Parra MT, Arévalo S, García de la Vega C, Santos JL, Page J. 2021. X chromosome

inactivation during grasshopper spermatogenesis. Genes 12:1844.

White MA, Ikeda A, Payseur BA. 2012. A pronounced evolutionary shift of the

pseudoautosomal region boundary in house mice. Mamm. Genome 23:454–466.

White MA, Stubbings M, Dumont BL, Payseur BA. 2012. Genetics and evolution of hybrid male

sterility in house mice. Genetics 191:917–934.

Wittkopp PJ, Haerum BK, Clark AG. 2004. Evolutionary changes in cis and trans gene

regulation. Nature 430:85–88.

Wu CI, Davis AW. 1993. Evolution of postmating reproductive isolation: the composite nature

of Haldane's rule and its genetic bases. Am. Nat. 142:187–212.

Zhu W, Hu B, Becker C, Doğan ES, Berendzen KW, Weigel D, Liu C. 2017. Altered chromatin

compaction and histone methylation drive non-additive gene expression in an interspecific

Arabidopsis hybrid. Genome Biol. 18:157.