Different complex regulatory phenotypes underlie hybrid male sterility in divergent rodent crosses

Kelsie E. Hunnicutt[*,1], Colin Callahan[2], Sara Keeble[2], Emily C. Moore[1,2], Jeffrey M. Good[2], and Erica L. Larson[*,1]

[1] University of Denver, Department of Biological Sciences, Denver, CO, 80208
[2] Division of Biological Sciences, University of Montana, Missoula, MT, 59812
*Corresponding authors

ORCID: 0000-0002-9674-0630 (KEH)
ORCID: 0000-0002-0385-1121 (CC)
ORCID: 0000-0002-7318-3122 (SK)
ORCID: 0000-0001-6166-3367 (ECM)
ORCID: 0000-0003-0707-5374 (JMG)
ORCID: 0000-0003-3006-645X (ELL)

Corresponding Authors:
Kelsie Hunnicutt
Department of Biological Sciences
University of Denver
Denver, CO 80208
Email: Kelsie.Hunnicutt@du.edu

Erica Larson
Department of Biological Sciences
University of Denver
Denver, CO 80208
Email: Erica.Larson@du.edu

Running title: Rodent hybrid male sterility

Key words: speciation, hybrid male sterility, gene expression, fluorescence activated cell sorting, meiotic sex chromosome inactivation, *Phodopus*

ABSTRACT

Hybrid incompatibilities are a critical component of species barriers and may arise due to negative interactions between divergent regulatory elements in parental species. We used a comparative approach to identify common themes in the regulatory phenotypes associated with hybrid male sterility in two divergent rodent crosses, dwarf hamsters and house mice. We investigated three potential characteristic gene expression phenotypes in hybrids including the propensity of transgressive differentially expressed genes towards over or underexpression, the influence of developmental stage on patterns of misexpression, and the role of the sex chromosomes on misexpression phenotypes. In contrast to near pervasive overexpression in hybrid house mice, we found that misexpression in hybrid dwarf hamsters was dependent on developmental stage. In both house mouse and dwarf hamster hybrids, however, misexpression increased with the progression of spermatogenesis, although to varying extents and with potentially different consequences. In both systems, we detected sex-chromosome specific overexpression in stages of spermatogenesis where inactivated X chromosome expression was expected, but the hybrid overexpression phenotypes were fundamentally different. Importantly, misexpression phenotypes support the presence of multiple developmental blocks to spermatogenesis in dwarf hamster hybrids, including a potential role of meiotic stalling or breakdown early in spermatogenesis. Collectively, we demonstrate that while there are some similarities in hybrid expression phenotypes of house mice and dwarf hamsters, there are also clear differences that point towards unique mechanisms underlying hybrid male sterility. Our results highlight the potential of comparative approaches in helping to understand the causes and consequences of disrupted gene expression in speciation.

INTRODUCTION

The evolution of postzygotic reproductive barriers, such as hybrid inviability and

sterility, is an important part of the speciation process, and identifying the genetic architecture

of hybrid incompatibilities has been a common goal uniting speciation research (Coughlan and

Matute 2020). While identifying the genetic basis of hybrid dysfunction remains difficult in many

systems, downstream regulatory phenotypes can provide insight into the underlying

mechanisms of speciation (Mack and Nachman 2017). An outstanding question surrounding

the role of disrupted gene regulation and speciation is whether the combination of two

divergent genomes in hybrids results in gene expression perturbations that are consistent or

repeatable across species. At the broadest level, it is unclear whether gene expression in

hybrids tends to be intermediate or transgressive (outside the range of parental gene

expression), whether transgressive gene expression is biased towards over or underexpression

(Ortíz-Barrientos *et al.* 2007), and whether transgressive misexpression tends to be modulated

by *cis* or *trans* regulatory elements (Wittkopp *et al.* 2004; McManus *et al.* 2010; Oka *et al.* 2014;

Mack *et al.* 2016; Mugal *et al.* 2020; Kopania *et al.* 2022a). By investigating trends in the

magnitude and direction of transgressive expression across different hybrid systems, we can

begin to understand the evolutionary forces shaping regulatory-based hybrid incompatibilities.

For example, if transgressive misexpression in hybrids tends towards overexpression, this may

mean that genes with disrupted regulation in hybrids tend to be genes that are normally

repressed in parental lineages (Meiklejohn *et al.* 2014; Barreto *et al.* 2015; Larson *et al.* 2017).

Alternatively, transgressive misexpression in hybrids may tend towards underexpression if

regulatory divergence between parental lineages results in impaired transcription factor binding

with promoter or enhancer elements (Oka *et al.* 2014; Guerrero *et al.* 2016) or if divergence

stimulates epigenetic silencing (Paun *et al.* 2007; Shivaprasad *et al.* 2012; Lafon-Placette and

Köhler 2015; Brekke *et al.* 2016; Zhu *et al.* 2017). At a finer scale, transgressive misexpression

patterns may depend on developmental stage: for example, if there is greater pleiotropy earlier

in development (Ortíz-Barrientos *et al.* 2007; Cutter and Bundus 2020). In particular, we might

expect sterile hybrids to have more transgressive misexpression during later stages of

gametogenesis when genes are evolving rapidly and are potentially under less regulatory

constraint (Kopania *et al.* 2022a; Murat *et al.* 2023). Finally, the role of sex chromosome

regulation in inviable or sterile hybrids encompasses both larger questions. Sex chromosomes

may be prone to asymmetry in their expression divergence (Oka and Shiroishi 2014; Civetta

2016) and be regulated differently across stages of development (Presgraves 2008; Larson *et*

al. 2018), particularly in reproductive tissues, and thus may play a central role in hybrid

dysregulation relative to autosomes.

Disruption of sex chromosome regulation is thought to be a potentially widespread

regulatory phenotype in sterile hybrids (Lifschytz and Lindsley 1972; Larson *et al.* 2018), in part

because X chromosome repression may be crucial to normal spermatogenesis in diverse taxa

(McKee and Handel 1993; Landeen *et al.* 2016; Taxiarchi *et al.* 2019; Rappaport *et al.* 2021;

Viera *et al.* 2021; Murat *et al.* 2023). Furthermore, misregulation of the X chromosome is

associated with hybrid sterility in several species pairs (Davis *et al.* 2015; Sánchez-Ramírez *et*

al. 2021; Bredemeyer *et al.* 2021), although it has been best studied in house mice. In fertile

male mice, the X chromosome is silenced just prior to the Diplotene stage of meiosis through

meiotic sex chromosome inactivation (MSCI; McKee and Handel 1993; Handel 2004) and is

again repressed in postmeiotic sperm development (*i.e.,* postmeiotic sex chromosome

repression or PSCR; Namekawa *et al.* 2006). In contrast, the X chromosome is not properly

inactivated and is overexpressed in sterile hybrid mice (Good *et al.* 2010; Bhattacharyya *et al.*

2013; Campbell *et al.* 2013; Turner *et al.* 2014; Larson *et al.* 2017, 2022). Disrupted MSCI in

house mice is associated with divergence at *Prdm9*, a gene that is a major contributor to

hybrid male sterility (Mihola *et al.* 2009; Davies *et al.* 2016). However, misexpression of the X

chromosome in sterile hybrids could result from mechanisms other than *Prdm9*-associated

disrupted MSCI, such as mispairing of the sex chromosomes due to divergence in their region

of homology known as the pseudoautosomal region (PAR; Burgoyne 1982; Ellis and

Goodfellow 1989; Raudsepp and Chowdhary 2015). In sum, the ubiquity of X chromosome

repression and the growing body of evidence linking disrupted MSCI to hybrid sterility in

mammals suggest that disrupted sex chromosome regulation may be a common regulatory

phenotype in sterile hybrid males.

Here, we characterized disruption of gene expression associated with hybrid male

sterility in two rodent crosses, dwarf hamsters and house mice, which span ~35 million years

of divergence (Swanson *et al.* 2019). The regulatory phenotypes of hybrid male sterility have

been thoroughly studied in house mice (Good *et al.* 2010; Bhattacharyya *et al.* 2013; Campbell

et al. 2013; Turner *et al.* 2014; Larson *et al.* 2017, 2022; Hunnicutt *et al.* 2022). We contrast

these with an analogous cross between two sister species of dwarf hamster, Campbell's dwarf

hamster (*Phodopus campbelli*) and the Siberian dwarf hamster (*P. sungorus*), and their sterile

F1 hybrid male offspring. These species diverged only ~0.8-1.0 million years ago (Neumann *et*

al. 2006), and they are not thought to interbreed in the wild due to geographic separation

(Ishishita *et al.* 2015). Crosses between female *P. sungorus* and male *P. campbelli* produce

sterile hybrid males that, similar to mice, have a range of sterility phenotypes, suggesting

multiple developmental blocks to spermatogenesis (Ishishita *et al.* 2015; Bikchurina *et al.*

2018). Hybrids from the reciprocal cross are usually inviable due to abnormal growth *in utero*

(Brekke and Good 2014), and the species origin of the X chromosome is the primary genetic

factor controlling hybrid inviability (Brekke *et al.* 2021). Additionally, sex chromosome

asynapsis during spermatogenesis is common in hybrid dwarf hamsters, providing further

reason to think that X chromosome-specific misregulation may be observed in sterile hybrid

dwarf hamsters (Ishishita *et al.* 2015; Bikchurina *et al.* 2018). Both the abnormal spermatogenic

phenotypes observed in dwarf hamster hybrids and the potential regulatory interactions that

may result from the involvement of the X chromosome in multiple reproductive barriers make

dwarf hamsters an important comparison to mice for investigating what regulatory phenotypes

may be repeatedly associated with the evolution of postzygotic reproductive isolation.

Expression phenotypes associated with hybrid sterility have historically been difficult to

assess because of the cellular diversity of reproductive tissues (*e.g.,* testes; Ramm and

Schärer 2014) and because hybrids may differ from parents in both tissue composition and

developmental timing (reviewed in Montgomery and Mank 2016; Hunnicutt *et al.* 2022). To

overcome these difficulties, we used Fluorescence Activated Cell Sorting (FACS) to isolate and

sequence cell populations across the developmental timeline of spermatogenesis for each

species pair and their F1 hybrids, including stages that span the different sex chromosome

regulatory states. Our developmental timeline spans stages where we expect fertile parents to

have a transcriptionally active X chromosome (spermatogonia and leptotene/zygotene

spermatocytes) and an inactive X chromosome (diplotene spermatocytes and round

spermatids). We used both datasets to address three main questions about the transgressive

gene expression phenotypes observed in sterile hybrids: (1) within transgressive differentially

expressed genes, does misexpression tend towards up- or downregulation in hybrids

compared to parents? (2) are there similar patterns of disrupted transgressive expression

across stages of development? and (3) are there clear differences between autosomes and sex

chromosomes in expression phenotypes? And if so, is sex chromosome-specific transgressive

misexpression consistent with either disrupted MSCI and/or disrupted PAR regulation?

Collectively, we demonstrate the power of cell type-specific approaches for untangling the

expression phenotypes associated with the evolution of hybrid male sterility and for identifying

common themes in the mechanistic basis of hybrid incompatibilities across divergent taxa.

MATERIALS AND METHODS

Hamster crosses and male reproductive phenotypes

We used wild-derived colonies of two sister species of dwarf hamster, *P. sungorus* and

P. campbelli, established by Kathy Wynne-Edwards (Scribner and Wynne-Edwards 1994) and

housed at the University of Montana. Both species were maintained as closed colonies with a

breeding scheme to minimize inbreeding. Nonetheless, inbreeding levels of these closed

colonies are still high as indicated by very low nucleotide diversity (Brekke *et al.* 2018). We

used males from both parent species and male F1 hybrid offspring from crosses of female *P.*

campbelli with male *P. sungorus*. We weaned males in same-sex sibling groups between 17 -

21 dpp and housed them individually at 45 dpp. We euthanized reproductively mature males

using carbon dioxide followed by cervical dislocation between 59 - 200 dpp (Table S1 in File

S1). All animal use was approved by the University of Montana (IACUC protocols 050-

16JGDBS & 035-19JGDBS).

We measured several fertility metrics for parent species and hybrid males including

paired testes weight, paired seminal vesicle weight, normalized sperm counts, and sperm

motility (Good *et al.* 2008). Paired testes weight and paired seminal vesicle weight were

correlated with body weight (paired testes weight Pearson's $r(29) = 0.47$, $p = 0.007$; paired

seminal vesicle weight Pearson's $r(23) = 0.56$, $p = 0.003$), so we standardized both metrics

relative to body weight. We calculated sperm count by isolating sperm from caudal

epididymides diced in 1 ml of Dulbecco's PBS (Sigma) and incubated at 37°C for 10 minutes.

We quantified sperm motility (proportion of motile sperm in a 5 µl suspension) and sperm count

(number of sperm with head and tail in a heat shocked 5 µl suspension) across a fixed area on

a Makler counting chamber. We performed statistical comparisons of fertility phenotypes in R

v.4.3.1, and we used the FSA package v.0.9.4 for the Kruskal-Wallis and Dunn's tests (Ogle

and Ogle 2017).

Isolation of enriched cell populations from hamster testes

To investigate the regulatory dynamics of the sex chromosomes during

spermatogenesis, we isolated four spermatogenic cell populations from whole testes using

FACS. These cell populations span a developmental timeline of spermatogenesis from mitosis

(spermatogonia), meiosis prior to X inactivation (leptotene/zygotene spermatocytes), meiosis,

after X inactivation (diplotene spermatocytes), and post-meiosis (round spermatids). Briefly, we

disassociated a single testis per male following a published protocol originally developed for

house mice (Getun *et al.* 2011) with modifications (github.com/goodest-goodlab/good-

protocols/tree/main/protocols/FACS, last accessed June 16, 2021). We doubled the volumes

of all reagents to account for the increased mass of testes in dwarf hamsters relative to house

mice. We isolated cell populations based on size, granularity, and fluorescence on a FACSAria

Ilu cell sorter (BD Biosciences) at the University of Montana Center for Environmental Health

Sciences Fluorescence Cytometry Core. For each sorted cell population, we extracted RNA

using RNeasy kits (Qiagen) following protocols for Purification of Total RNA from Animal Cells.

We quantified sample RNA quantity and quality (requiring an RNA integrity number > 7) on a

Tapestation 2200 (Agilent) at the University of Montana genomics core. RNA libraries were

prepared by Novogene and sequenced on Illumina NovaSeq 6000s (paired end, 150 bp). Six

samples (distributed across different species and cell types) had low RNA concentrations, and

for these samples we used Novogene's low input RNA library preparation (Table S2 in File S1).

MDS plots indicated no severe library batch effects between samples from different library

preparations (Figure S1 in File S1), so we included all samples from both libraries in

subsequent analyses.

Read processing and mapping

We sequenced RNA from each cell population for three to five individuals of each

parent species and F1 hybrids generating an average of ~27.5 million read pairs per individual

(Table S2 in File S1). We trimmed reads using Trimmomatic v.0.39 (Bolger *et al.* 2014) to

remove low quality bases from the first and last 5 bp of each read and bases with an average

Phred score of less than 15 across a 4 bp sliding window and only retained reads of at least 36

bp. We next used an approach (based on the modtools pipeline) which maps reads from each

sample to pseudogenomes for both parent species (described below) to obtain a merged

output alignment file in order to alleviate reference bias associated with mapping hybrids to

only a single reference genome (Holt *et al.* 2013; Huang *et al.* 2014). For this approach, we

mapped reads for each individual to both a *P. sungorus* pseudogenome and a *P. campbelli*

pseudogenome with Hisat v.2.2.0 (Kim *et al.* 2019) with default settings and retaining at most

100 distinct, primary alignments, although multi-mapped reads were removed downstream

(described below). We generated the *P. sungorus* pseudogenome by mapping RNASeq reads

from a male *P. sungorus* individual (30.6 million total read pairs; NCBI SRA: SRR17223284;

Moore *et al.* 2022) to the *P. sungorus* reference genome (GCA_023856395.1) with bwa-mem

v.2.2.1 (Vasimuddin *et al.* 2019), and the *P. campbelli* pseudogenome by mapping female *P.*

campbelli whole genome sequencing reads (average coverage: 33x; NCBI SRA: SRR17223279;

Moore *et al.* 2022) to the *P. sungorus* reference genome. Because our *P. sungorus*

pseudogenome was based on a male hamster and the reference genome on a female hamster,

we excluded reads mapping to the PAR because sequences mapping to this region could have

originated from either the X or Y chromosomes and interfered with subsequent variant calling.

Following mapping, we used GATK v.4.2.5.0 HaplotypeCaller (-ERC GVCF) to call SNPs then

performed genotyping with genotypeGVCFs. We hard-filtered our SNPs (--mask-extension 5

"QD < 2.0" "FS > 60.0" "MQ < 40.0" "QUAL < 30.0" "DP < 10" "DP > 150") and restricted

SNPs to biallelic loci. Finally, we incorporated filtered SNPs back into the *P. sungorus*

reference genome with FastaAlternateReferenceMaker to create the *P. sungorus* and *P.*

campbelli pseudoreferences. For our RNASeq data, we appended query hit indexes to

resulting alignment files using hisat2Tophat.py (https://github.com/goodest-goodlab/pseudo-

it/tree/master/helper-scripts/hisat2Tophat.py, last accessed March 8th, 2022) to maintain

compatibility with the modtools pipeline. We used our VCFs (above) to generate a mod-file for

both species with vcf2mod from Lapels v.1.1.1 to convert alignments to the *P. sungorus*

reference genome, and Suspenders v.0.2.6 to merge alignments while retaining the highest

quality alignment per read (Holt *et al.* 2013; Huang *et al.* 2014). We used featureCounts v.2.0.1

(Liao *et al.* 2014) to estimate counts of read pairs that aligned to the same chromosome (-B

and -C) and retained only singly-mapped reads. Summaries of properly mapped reads for each

sample can be found in Table S2 in File S1.

We sought to compare the gene expression phenotypes observed in dwarf hamsters to

those previously documented in house mice using published RNASeq data for the same four

spermatogenic cell types of two subspecies of house mouse and their sterile F1 hybrids

(Larson *et al.* 2017; Hunnicutt *et al.* 2022). These studies examined two subspecies of house

mice, *Mus musculus musculus* (intra-subspecific F1 males between wild-derived inbred strains

PWK/PhJ♀ and CZECHII/EiJ♂) and *M. m. domesticus* (intra-subspecific F1 males between

wild-derived inbred strains WSB/EiJ♀ and LEWES/EiJ♂) and their sterile (PWK♀ x LEWES♂)

F1 hybrids for disrupted gene expression across spermatogenesis following the same FACS

protocols implemented in this study. For all comparisons between house mice and dwarf

hamsters, we used read count files generated previously for house mice (Hunnicutt *et al.* 2022)

and performed all subsequent analyses in parallel for both systems.

Estimating nucleotide diversity and divergence within and between crosses and strains

We next estimated nucleotide diversity within and divergence between each strain or

cross for parent species and hybrids (π and d_{XY}). We used the bam files generated above from

mapping each strain or cross to its respective reference genome (either GCA_023856395.1 for

dwarf hamsters or GRCm38.p6 for house mice) to call SNPs as described above but with the

addition of a step to split reads at intronic regions using the GATK function SplitNCigarReads.

We generated two VCFs, one for dwarf hamster crosses and the other for house mouse

crosses, which we processed and filtered separately. If an individual mouse or hamster was

sequenced for more than one cell type, we randomly chose one of the represented cell types

to be included in the analysis. We hard-filtered our SNPs as above and restricted SNPs to

biallelic loci. Because our SNPs were called from RNASeq data and thus may be susceptible to

allelic imbalances or coverage differences between samples, we next investigated how the

inclusion of SNPs with different levels of missing data impacted estimation of π and d_{XY}. We

used vcftools v.0.1.17 (Danecek *et al.* 2011) to filter SNPs allowing between 0% and 90%

missing data (--max_missing; Figure S3 in File S1), and filtered SNPs with a depth lower than 5

and higher than 60 (~2-3 higher than average coverage) to eliminate multi-mapped reads.

Finally, we used pixy v.1.2.10.beta2 (Korunes and Samuk 2021) on our filtered VCFs to

estimate π and d_{XY}. Patterns of nucleotide diversity across strains and crosses were

qualitatively similar across missing data thresholds, so we present results corresponding to

10% missing data in the main text. However, we note that for all comparisons, estimates of π

and d_{XY} decreased with more stringent missing data thresholds regardless of strain or cross

(Figure S3 in File S1).

Gene expression pre-processing

Following read processing and mapping, we conducted all analyses in R v.4.3.1. We

classified genes as "expressed" if genes had a minimum of one Fragment Per Kilobase of exon

per Million mapped reads (FPKM) in at least three samples, resulting in 21,077 expressed

genes across the dwarf hamster dataset and 21,212 expressed genes across the house mouse

dataset. We also identified sets of genes "induced" in a given cell type defined as genes with a

median expression in a given cell population (normalized FPKM) greater than two times its

median expression across all other sorted cell populations (following Kousathanas *et al.* 2014).

We calculated normalized FPKM values by adjusting the sum of squares to equal one using the

R package vegan v.2.6-4 (Oksanen *et al.* 2013). We conducted expression analyses using

edgeR v.3.42.4 (Robinson *et al.* 2010) and normalized the data using the scaling factor method

(Anders and Huber 2010).

We qualitatively assessed cell population purity both by visual inspection during cell

sorting and following sequencing by assessing the expression of a panel of marker genes

specific to the four cell populations targeted by our FACS protocol and present in only a single

copy in the *P. sungorus* annotation. Spermatogonia markers included *Dmrt1* (Raymond *et al.*

2000) and *Hells* (Green *et al.* 2018). Leptotene/zygotene markers included *Ccnb1ip1* and

Adad2 (Hermann *et al.* 2018). Diplotene was characterized by *Aurka* and *Tank* expression

(Murat *et al.* 2023) and round spermatids by *Cabyr* and *Acrv1* expression (Green *et al.* 2018).

To estimate relative purity, we compared mean marker gene expression across replicates for a

given cell population for both parent species. A cell population was considered qualitatively

pure if it had higher marker gene expression than other populations isolated by our FACS

protocol and if X-linked gene expression matched the expected regulatory dynamics (*i.e.,*

active vs. silenced (MSCI) vs. repressed (PSCR); Handel 2004; Namekawa *et al.* 2006). We

examined expression patterns across cell populations for all genes, autosomal genes, and X-

linked genes using MDS plots generated with the plotMDS function in limma v.3.56.2 (Ritchie

et al. 2015) and heatmaps using ComplexHeatmap v.2.16.0 (Gu *et al.* 2016). MDS plots used

the top 500 genes with the largest fold change difference between samples.

Differential gene expression analysis

We assessed differential gene expression by contrasting hybrids and each parent

species for all cell populations. We fit the expression data for dwarf hamsters and house mice

separately with negative binomial generalized linear models with Cox-Reid tagwise dispersion

estimates and adjusted P-values to a false discovery rate (FDR) of 5% (Benjamini and

Hochberg 1995). We quantified the biological coefficient of variation (BCV), a metric

representing the variation in gene expression among replicates (McCarthy *et al.* 2012), for each

dataset. Additionally, we calculated the BCV of just parental males or hybrid samples for each

species for the first three cell populations to examine whether dwarf hamster hybrids exhibited

more variability in expression than house mouse hybrids and parental dwarf hamsters. For our

differential expression analyses, we contrasted expression between hybrids and each parent

so that a positive log fold-change (logFC) indicated overexpression in sterile males and

implemented a logFC cutoff of 1.25. We then categorized differentially expressed (DE) genes

into one of four categories: DE relative to only one parent species, DE relative to both parent

species but with intermediate expression (intermediate), and DE relative to both parent species

but outside of the range of either parent species (transgressive; Figure S2 in File S1). Unless

otherwise specified, results discussed in the main text are restricted to transgressive DE genes

and are presented with the logFC from the contrast of the hybrid offspring to the parent with

the same X chromosome (*P. campbelli* for dwarf hamster F1 hybrids and *M. m. musculus* for

house mouse F1 hybrids). Transgressive DE genes have similar logFC values regardless of

which parent is used as the contrast, but figures depicting the logFC between F1 hybrids and

P. sungorus or *M. m. domesticus* are provided in supplementary material. We also assessed

differential expression between parent species/strains (parental DE) by contrasting expression

between parents sharing the same X chromosome as hybrids (*i.e., M. m. musculus* and *P.*

campbelli) with parents with the alternate X chromosome (*i.e., M. m. domesticus* and *P.*

sungorus) so that a positive log fold-change indicated overexpression in parents with the same

X chromosome as hybrids and implemented a logFC cutoff of 1.25.

We tested for significant differences in the number of under and overexpressed

transgressive DE genes within a stage for both house mouse and hamster hybrids using X^2

tests with chisq.tests in R and used FDR correction for multiple comparisons. We also tested

for differences in the magnitude of misexpression between mouse and hamster hybrids for

each stage by comparing the distributions of logFC of transgressive DE genes between hybrids

and parent species using Wilcoxon signed-rank tests and FDR correction. To characterize

hybrid diplotene expression in both house mice and dwarf hamsters, we used two approaches.

First, we calculated Pearson's correlation coefficient *(r)* between average normalized hybrid

diplotene expression and the average normalized expression in each parental cell type. We

corrected p-values for each correlation with FDR. We generated bootstrap values for each

correlation coefficient by randomly sampling the expression matrices with replacement for

each sample type for 1000 replicates. Second, we compared the gene sets that escaped MSCI

in each species with gene sets that characterize stage-specific expression in parent species.

For this analysis, we defined sets of overexpressed X-linked diplotene genes as genes with

expression (normalized FPKM) in hybrids that was in the top 10% of X-linked genes in parental

diplotene samples (*i.e.,* genes that normally escape MSCI). We then compared these sets of

overexpressed hybrid diplotene genes to genes "induced" in each parental stage for each

species. We used gProfiler2 v.0.2.3 (Kolberg *et al.* 2020) in R to perform gene ontology (GO)

analysis to identify GO terms overrepresented in hybrid transgressive DE genes sets for each

cell population. We only included *P. sungorus* genes associated with mouse orthologs in our

GO analysis (as established by Moore *et al.* 2022), and for our background gene lists, we used

P. sungorus genes associated with mouse orthologs that were "expressed" in hybrids and both

parent species in a given stage. We retained only Biological Process GO terms with an FDR

below 0.05 and ran gProfiler2 both with and without the highlight option, a two-stage algorithm

for reducing resulting GO terms by grouping significant terms into sub-ontologies and then

identifying the gene sets that give rise to other significant functions (Kolberg *et al.* 2020). To

test whether specific chromosomes were enriched or depleted for transgressive DE genes for a

given stage, we performed hypergeometric tests on the number of transgressive DE genes on

a given chromosome with phyper and adjusted P-values to an FDR of 5%. We also assessed

overlap in specific transgressive DE genes within stages between house mice and dwarf

hamsters. We estimated whether overlap was more or less than expected by chance and

whether overlapping genes were preferentially located on the X chromosome using

hypergeometric tests and performed GO enrichment on overlapping genes. For all

hypergeometric tests, we defined the background sets of genes as those with non-zero logFC

values in both differential expression comparisons between hybrids and each parent for a

given spermatogenic stage.

Characterizing the behavior of PAR genes in dwarf hamsters

339   We sought to characterize the regulatory behavior of PAR genes in dwarf hamster

parent species and hybrids to determine if (1) PAR genes are normally silenced in parent

species (consistent with an extension of MSCI to the PAR) and (2) if PAR genes were

overexpressed in hybrids (consistent with disrupted MSCI in the PAR of hybrids). The PAR on

the *P. sungorus* X chromosome is on the distal arm of the X chromosome from around

115,350,000-119,112,095 bp (Moore *et al.* 2022). There are 15 annotated *P. sungorus* genes in

this region (Table S3 in File S1), which is comparable to the latest PAR assembly in C57BL/6J

house mice (Kasahara *et al.* 2022). Six of these are orthologous to annotated genes in mice

(*Tppp2*, *Gprin1*, *Ndrg2*, *Kcnip4*, *Ndrg2*, and *Hs6st3*), but they are not located in the mouse PAR

(Kasahara *et al.* 2022). Only seven of the annotated genes in the PAR were expressed in more

than three replicates across all samples (Psun_G000022875, Psun_G000022880,

Psun_G000022883, Psun_G000022886, *Tppp2*, *Gprin1*, and *Ndrg2*). For these genes, we

assessed whether these genes were consistently expressed or silenced in parent species in

any cell population and whether any genes were differentially expressed between hybrids and

either species.

RESULTS

Impaired sperm production in hybrid male dwarf hamsters

We first established the extent of hybrid male sterility in dwarf hamsters by comparing

reproductive phenotypes for *P. campbelli*, *P. sungorus*, and F1 hybrid males (Figure 1; Table

S1 in File S1). *Phodopus sungorus* males had smaller testes and seminal vesicles than *P.*

campbelli males (Dunn's Test relative testes weight p < 0.001; relative seminal vesicle weight p

= 0.0061; Figure 1). These differences are qualitatively consistent across independent

laboratory colonies (Ishishita *et al.* 2015; Bikchurina *et al.* 2018) and likely reflect species-

specific differences between *P. campbelli* and *P. sungorus*. *Phodopus sungorus* males also

had lower nucleotide diversity (π = 0.00013) than *P. campbelli* males (π = 0.00045) and both

house mouse parental crosses between fully inbred mouse strains (*M. m. musculus* π =

0.00029; *M. m. domesticus* π 0.00017; Figure S3 in File S1), which could contribute to

depression of male fertility within this highly inbred laboratory colony (Brekke *et al.* 2018).

Nucleotide divergence between parental dwarf hamster species was elevated relative to house

mice (d_{xy} = 0.0020 vs. 0.0010), consistent with reported older divergence time estimates for

dwarf hamsters (Neumann *et al.* 2006). However, these dwarf hamster species did not differ in

normalized sperm counts (p = 0.071) or in sperm motility (p = 0.12). The F1 hybrid males

exhibited extreme reproductive defects relative to *P. campbelli* (Figure 1). Hybrid males had

smaller testes (p < 0.001) and seminal vesicles (p < 0.001) than male *P. campbelli* hamsters,

and importantly, produced almost no mature spermatozoa. In the one instance where a hybrid

male produced a single mature spermatozoon, it was non-motile, indicating severe

reproductive impairment in hybrid males. Overall, our results confirmed previous reports of

reduced fertility in hybrid male dwarf hamsters (Ishishita *et al.* 2015; Bikchurina *et al.* 2018).

Figure 1. Evidence of some hybrid male sterility in dwarf hamsters. We assessed paired

testes weight and paired seminal vesicle weight (SV; both normalized by body weight), sperm

count, and proportion of motile sperm for *P. sungorus*, *P. campbelli*, and F1 hybrids. Whiskers

extend to either the largest or smallest value or no further than 1.5 times the interquartile range,

and *** indicates p < 0.001, ** indicates p < 0.01, and n.s. indicates non-significant difference

between means at p > 0.05 using a post-hoc Dunn's test with FDR correction. Upwards-

pointing triangles (▲) indicate *P. campbelli*, downwards-pointing triangles (▼) indicate *P.*

sungorus, and crosses (✖) indicate F1 hybrids.

Cell type-specific gene expression across spermatogenesis

To characterize cell type-specific gene expression, we used FACS to isolate enriched

cell populations from each fertile parent species and their sterile F1 hybrids across four stages

of spermatogenesis. The four targeted populations included: spermatogonia (mitotic precursor

cells), leptotene/zygotene spermatocytes (meiotic cells before MSCI), diplotene spermatocytes

(meiotic cells after MSCI), and round spermatids (postmeiotic cells). We were unable to isolate

round spermatids from the F1 hybrids, which was consistent with the lack of mature

spermatozoa present in the cauda epididymis extractions (Figure 1). We sequenced RNA from

each cell population for *P. campbelli* (spermatogonia n = 4, leptotene/zygotene n = 5, diplotene

n = 5, and round spermatids n = 4; Table S2 in File S1), *P. sungorus* (spermatogonia n = 4,

leptotene/zygotene n = 4, diplotene n = 5, and round spermatids n = 3), and F1 hybrid males (

P. campbelli ♀ x *P. sungorus* ♂; spermatogonia n = 4, leptotene/zygotene n = 4, diplotene n =

4). We compared our hamster expression data to an analogous cell type-specific RNASeq

dataset from two species of house mice, *Mus musculus musculus* and *M. m. domesticus*, and

their sterile F1 hybrids (n = 3 for all cell populations in each cross; Larson *et al.* 2017).

We used two approaches to qualitatively evaluate the purity of spermatogonia,

leptotene/zygotene spermatocytes, diplotene spermatocytes, and round spermatids isolated

from males from both parental species: (1) we quantified the relative expression of a panel of

cell population marker genes, and (2) we characterized the expression patterns of the sex

chromosomes across development. We found that our candidate marker genes had the

highest expression in their expected cell population for all stages except leptotene/zygotene

(Figures S4 and S5 in File S1), indicating high purity of spermatogonia, diplotene

spermatocytes, and round spermatids. Leptotene/zygotene markers did not have the highest

expression in leptotene/zygotene samples (except for *Ccnb1ip1*), potentially indicating lower

purity of this cell population. Nonetheless, the patterns of X chromosome expression in fertile

parents were consistent with expectations across this developmental timeline: the X

chromosome had active expression in spermatogonia and leptotene/zygotene cells, was

inactivated in diplotene cells consistent with MSCI, and was partially inactivated in round

spermatids, consistent with PSCR (Figure 2; Namekawa *et al.* 2006), indicating successful

isolation of these cell populations.

When we examined overall expression differences within dwarf hamsters and within

house mice, we found that samples clustered primarily by cell population on MDS1 and 2

(Figure 3), then by cross/strain when cell populations were examined separately (Figures S6

and S7 in File S1). Within all cell populations across both systems, hybrids showed

intermediate overall expression patterns to parent species (Figures S6 and S7 in File S1).

However, in dwarf hamsters but not house mice, spermatogonia and leptotene/zygotene

samples overlap rather than forming distinct clusters (Figure 3). Further, the expression profiles

of hybrid diplotene cell populations ranged from clustering with parental leptotene/zygotene to

parental diplotene cell populations, which contrasted with what we observed in house mice

where hybrid diplotene cell populations clustered distinctly with parental diplotene cell

populations (Figure 3). Overall, our results indicate that we successfully isolated cell

populations in dwarf hamsters that span key stages of spermatogenesis.

Figure 2. Overexpression of X-linked genes in diplotene spermatocytes in both house mouse and dwarf hamster hybrids. Heatmap of X-linked gene expression in house mice (upper panel) and dwarf hamsters (lower panel) plotted as normalized FPKM values that are hierarchically clustered using Euclidean distance. Each column represents a different individual, each row represents a gene, and darker colors indicate higher expression. The heatmap was generated with the R package ComplexHeatmap v.2.12.0 (Gu et al. 2016). Note, hybrid dwarf hamsters do not produce mature spermatozoa, and accordingly, we were unable to isolate round spermatids.

Figure 3. Hybrid gene expression profiles cluster by parental spermatogenic cell

population in house mice but not dwarf hamsters. Multidimensional scaling (MDS) plots of

distances among house mouse (upper panels) and dwarf hamster (lower panels) samples for

expressed autosomal (left) and X-linked (right) genes. Distances are calculated as the root-

mean-square deviation (Euclidean distance) of log2 fold changes among the top 500 genes

that distinguish each sample. Each strain or cross is indicated by a symbol, and samples are

colored by cell population.

Disrupted transcription early in spermatogenesis in dwarf hamsters

We sought to characterize which expression phenotypes were associated with sterile

hybrids in both house mice and dwarf hamsters. We first investigated whether differential gene

expression in hybrids tended towards intermediate or transgressive expression, and then

within transgressive DE genes, whether misexpression tends towards up- or downregulation in

hybrids compared to parents. Differential gene expression in mouse hybrids had a slight bias

towards transgressive expression except in round spermatids (percentage of transgressive DE

genes: SP: 72.0%, LZ: 62.7%, DIP: 60.8%, and RS: 24.9%; Figures 4b and S2 in File S1), while

almost all differential expression in dwarf hamster hybrids was transgressive (percentage of

transgressive DE genes: SP: 99.2%, LZ: 98.0%, and DIP: 97.1). For transgressive DE genes,

autosomal misexpression in hybrid house mice was biased towards upregulation across

spermatogenesis (mean logFC of spermatogonia autosomal transgressive DE genes: +1.90/ X^2:

p < 0.001; leptotene/zygotene: +1.33/ X^2: p < 0.001; diplotene: +0.47/ X^2: p < 0.001; Figures 4c

and S8 in File S1), as was X-linked misexpression (spermatogonia mean logFC: +1.98 / X^2: p <

0.001; leptotene/zygotene: +2.50/ X^2: p < 0.001; diplotene: +2.78/ X^2: p < 0.001). In contrast,

we found that the direction of misexpression in dwarf hamster hybrids was cell type-specific.

Autosomal transgressive DE genes in dwarf hamster hybrids were overwhelmingly

downregulated in both early stages of spermatogenesis, especially in comparison to house

mice (mean logFC of spermatogonia autosomal transgressive DE genes: -4.38/ X^2: p < 0.001;

leptotene/zygotene: -4.67/ X^2: p < 0.001; Figure 4d) but upregulated in diplotene (average

logFC = +4.67/ X^2: p < 0.001; Figures 4d and S8 in File S1). When comparing both X-linked and

autosomal expression in dwarf hamsters, we found similar patterns: almost all X-linked

transgressive DE genes in the first two stages of spermatogenesis were exclusively

downregulated (spermatogonia mean logFC = -6.57/ X^2: p < 0.001; leptotene/zygotene: -6.73/

X^2: p < 0.001), but misexpression was biased towards upregulation in diplotene (mean logFC:

5.70/ X^2: p < 0.001).

Figure 4. House mice and dwarf hamster hybrids have opposite patterns of disrupted

regulation early in spermatogenesis. a) Up- and down-regulated DE genes between parent

species for house mice (left; *M. m. musculus* vs. *M. m. domesticus*) and dwarf hamsters (right;

P. sungorus vs. *P. campbelli*). b) Counts of DE genes between hybrids and one parent species

(two lightest shades of gray where Parent 1 was either *M. m. musculus or P. sungorus* and

parent 2 was either *M. m. domesticus* or *P. campbelli)* or between hybrids and both parents

(two darker shades of gray) that showed either intermediate or transgressive expression. c)

Transgressive DE genes in house mouse and dwarf hamster hybrids for autosomal genes

where the logFC represents hybrid expression relative to *M. m. musculus* or *P. campbelli,*

respectively, and d) transgressive DE gene expression for X-linked genes*.* Results are

displayed for autosomes (left) and the X chromosome (right). *** indicates p < 0.001 for

pairwise comparisons from Wilcoxon signed-rank tests after FDR correction. Whiskers extend

to either the largest or smallest value or no further than 1.5 times the interquartile range. The

number of up and downregulated transgressive DE genes in hybrids are listed next to arrows

indicating direction of differential expression.

Second, we investigated how developmental stage influenced the extent of hybrid

misexpression. In both systems, the number of DE genes between parent species increased

with the progression of spermatogenesis, consistent with less constraint on gene expression

levels as spermatogenesis progresses (Figure 4a; Kopania *et al.* 2022a; Murat *et al.* 2023).

Similarly, transgressive misexpression in both hybrid dwarf hamsters and house mice

increased with the progression of spermatogenesis, though to a greater extent in dwarf

hamsters (Figure 4b). While both parental DE genes and transgressive hybrid DE genes

increased with the progression of spermatogenesis, parental DE genes initially exceeded the

number of transgressive hybrid DE genes in hybrids in spermatogonia and leptotene/zygotene

spermatocytes. However, in diplotene spermatocytes, transgressive hybrid differential

expression surpassed parental differential expression in both house mice and dwarf hamsters.

We also found a much greater genome-wide disruption of expression in diplotene cell

populations of hybrid dwarf hamsters than in hybrid house mice (Figures 4c, 4d, and S8-S10 in

File S1), indicating more widespread regulatory disruption. We then compared transgressive

DE genes between house mice and dwarf hamster hybrids across all three stages and found

that the number of shared transgressive DE genes did not differ from the number expected by

chance in early spermatogenesis (spermatogonia = 3 genes, hypergeometric test p = 0.98;

leptotene/zygotene = 2 genes, p = 0.59; Table S4 in File S1). However, the number of shared

transgressive DE genes did exceed the number expected by chance in diplotene cell

populations (n = 68; p < 0.001), and these shared genes were preferentially located on the X

chromosome (61/68; p < 0.001). Additionally, these sets of shared transgressive DE genes

between house mice and dwarf hamsters were not significantly enriched for any GO terms,

though many play known roles in spermatogenesis, the apoptotic process, and cell

differentiation (Table S5 in File S1) and may be promising candidates for future functional

analysis. Ultimately, we found that there are similar trends in the patterns of transgressive

expression across stages in both systems, and although few specific genes had disrupted

expression in both dwarf hamster and house mouse hybrids, genes with similar patterns of

disrupted expression tend to be X-linked and disrupted in later stages of spermatogenesis.

To determine if the differences we observed in the extent of misexpression between

dwarf hamsters and house mice was due to greater expression variability across our dwarf

hamster samples, we calculated the BCV, a measurement of inter-replicate variability, for each

species and hybrid across the first three cell populations. Inter-replicate variability was higher

in dwarf hamster hybrids relative to house mouse hybrids (dwarf hamster BCV = 0.69; house

mice = 0.18). Additionally, nucleotide diversity within dwarf hamster hybrids was higher (π =

0.0012) than within house mouse hybrids (π = 0.00052; Figure S3 in File S1). However, the

extent of the expression variability observed in hybrids relative to the inter-replicate variability

of parental species differed between house mice and dwarf hamsters: dwarf hamster hybrid

variability was more than dwarf hamster parental samples (*P. campbelli* = 0.49; *P. sungorus* =

0.40), but hybrid variability was similar to parental samples for house mice (*M. m. musculus* =

0.19; *M. m. domesticus* = 0.22). The greater inter-replicate expression variability in dwarf

hamster hybrids relative to parent species suggests that the increased misexpression we see

in hybrids cannot be explained by greater inter-replicate variability in our dwarf hamster

samples alone, and it may also reflect the greater nucleotide diversity present among dwarf

hamster hybrids.

Third, we characterized whether there was a clear difference between the autosomes

and sex chromosomes in expression phenotype by testing whether the X chromosome was

enriched for transgressive DE genes in each stage for both systems. Across all stages of

spermatogenesis in house mice, the X chromosome was enriched for transgressive DE genes

between hybrids and parents (spermatogonia $p < 0.001$; leptotene/zygotene $p < 0.001$;

diplotene $p < 0.001$; round spermatids $p < 0.0036$; Figure S9 in File S1). In contrast,

misexpression in dwarf hamsters was not uniformly sex chromosome-specific across all

stages, as the sex chromosomes in dwarf hamsters showed no enrichment for transgressive

DE genes early in spermatogenesis (spermatogonia: $p = 0.21$; Figure S10 in File S1) despite X

chromosome enrichment in both leptotene/zygotene ($p = 0.0064$) and round spermatids ($p <$

0.001). However, we note that the magnitude of misexpression was greater for sex

chromosomes than autosomes in dwarf hamsters across all stages (Figures 4c and 4d;

discussed above). Only two autosomes were enriched for transgressive DE genes in dwarf

hamsters in any cell population: one scaffold on chromosome 5 in spermatogonia

(JAJQIY010003390.1; hypergeometric test; $p = 0.003$) and chromosome 11 in

leptotene/zygotene ($p = 0.017$; Figure S10 in File S1). Together, the subtle differences in the

distribution of transgressive DE genes across autosomes and the X chromosome between

dwarf hamster hybrids and house mouse hybrids suggest a difference in the extent of the role

for sex chromosome-specific disruption between systems.

Misexpression in diplotene appears to be unrelated to disrupted MSCI in dwarf hamsters

We next asked whether expression patterns indicated similar disrupted regulatory

processes resulting in sex chromosome-specific misexpression in both systems. In sterile

hybrid house mice, the mean logFC of transgressive X-linked DE genes during diplotene was

higher than the mean logFC of transgressive autosomal DE genes (X logFC = +2.78 in contrast

to autosomal logFC = +0.47; Figures 4c and 4d), consistent with disrupted MSCI (Good *et al.*

2010; Bhattacharyya *et al.* 2013; Campbell *et al.* 2013; Turner and Harr 2014; Larson *et al.*

2017, 2022). In sterile dwarf hamster hybrids, we also found elevated mean logFC of

transgressive X-linked DE genes relative to autosomal genes (X logFC = +5.70 in contrast to

autosomal logFC = +3.69), but the extent of X chromosome overexpression, as measured by

logFC of X-linked transgressive DE genes, was greater than in hybrid house mice (5.7/3.69 or

~1.5x higher; Figures 2, 4c, and 4d). There was also more variability in the extent of

overexpression of X-linked genes in hybrid dwarf hamsters compared to normal parental X-

linked expression during diplotene (relative overexpression = 18.5 +/- 6.8) than for

overexpression of X-linked genes in hybrid house mice compared to normal parental X-linked

expression in diplotene (relative overexpression = 1.75 +/- 0.09; Figures 2 and S11 in File S1).

Strikingly, some hybrid male dwarf hamsters had an almost completely silenced X

chromosome, while others had an almost completely transcriptionally-activated X chromosome

(Figure 2).

Despite overexpression of the X chromosome during diplotene in hybrid dwarf

hamsters, the overall expression phenotype, including the identity and the extent of

misexpression of overexpressed genes, appeared to fundamentally differ between house

mouse and dwarf hamster hybrids (Figures 2 and 5a-5c). We established these differences in

X-linked overexpression using two approaches. First, we tested which parental cell types had

the highest expression correlation with hybrid diplotene cell types for both X-linked and

autosomal genes. In mice, the expression profile of X-linked genes in hybrids during diplotene

was most positively correlated with the expression profile of X-linked parental round spermatid

genes, consistent with disrupted MSCI (spermatogonia *(r)* = -0.25, p < 0.001;

leptotene/zygotene *(r)* = -0.073, p = 0.035; round spermatid *(r)* = 0.26, p < 0.001; Figures 2 and

5a). Autosomal diplotene genes showed no positive correlations with either spermatogonia *(r* =

-0.26; p < 0.001), leptotene/zygotene (r = -0.032; p < 0.001), or round spermatids (r = -0.060; p

< 0.001; Figure 5a). If the X-linked overexpression phenotype in dwarf hamsters was consistent

with disrupted MSCI, then we would also expect X-linked and autosomal expression profiles in

hybrid diplotene to follow the same patterns. In contrast to this prediction, hybrid dwarf

hamster diplotene expression profiles for both X-linked and autosomal genes had a positive

correlation with parental leptotene/zygotene (autosomal *(r)* = 0.14, p < 0.001; X-linked *(r)* =

0.22, p < 0.001) and spermatogonia (autosomal *(r)* = 0.027, p < 0.001; X-linked *(r)* = 0.091, p =

0.016) and a negative correlation with round spermatids (autosomal *(r)* = -0.30, p < 0.001; X-

linked *(r)* = -0.31, p < 0.001; Figure 5b). These striking differences in the strength and direction

of expression profile correlations suggest that the regulatory mechanisms underlying the

overexpression phenotype of X-linked genes in sterile hybrids differed between house mice

and dwarf hamsters.

This difference in pattern was further supported when we compared which sets of

genes were overexpressed in hybrid diplotene in dwarf hamsters and house mice. For this

approach, we looked at parental gene expression patterns to characterize which stages of

spermatogenesis all genes were normally active during and characteristic of (*i.e.,* "induced";

see Methods). Using this information, we then identified which X-linked genes were

overexpressed in hybrid diplotene (defined as genes with normalized expression in the top

10% of X-linked genes) and assessed which parental stages the overexpressed X-linked genes

were characteristic of in both systems. As in our correlation analysis, we found that in hybrid

house mice, the genes that were overexpressed in diplotene most closely resembled genes

that are normally active in round spermatids in parental mice (51.6% of genes), but that in

hybrid dwarf hamsters, overexpressed diplotene genes resembled spermatogonia- and

leptotene/zygotene-specific genes (47.6% and 41.6% respectively; Figure 5c). Because of the

dissimilarity in the genes that were overexpressed during diplotene in both systems, we next

performed gene ontology (GO) enrichment analyses on the set of transgressive DE genes in

hybrids to determine which biological processes could be potentially contributing to this

pattern. In contrast to house mice hybrids where transgressive DE genes were enriched for no

biological processes, the transgressive DE genes in dwarf hamster were enriched for several

biological processes including cell junction/extracellular matrix organization, system

development, inflammatory response, and apoptotic process which together point to

widespread disruption of numerous processes necessary for normal male fertility (highlighted

terms presented in Figure 5d and the full lists in Tables S6-S8 in File S1). Collectively, these

results suggest that the X-linked overexpression phenotype in sterile hybrid dwarf hamsters is

inconsistent with disrupted MSCI and is possibly related to a stalling and breakdown of

spermatogenesis between leptotene/zygotene and diplotene during Prophase I.

Figure 5. Spermatogenesis in hybrid dwarf hamsters appears to stall after leptotene/ zygotene. a) We calculated the Pearson's correlation coefficient *(r)* between mean hybrid diplotene expression and the mean expression for each parental cell type for both house mice and b) dwarf hamsters. Correlation coefficients were calculated for both autosomal (left panels) and X-linked (right panels) genes. We then generated bootstrap values by randomly sampling the expression matrices for 1000 replicates. All correlation coefficients were significantly different from zero (p < 0.05) after FDR correction. c) Classification of X-linked overexpressed genes in diplotene cell populations of hybrid mice and dwarf hamsters by parental stage in which genes are induced. Genes are colored by cell type (red = spermatogonia, yellow = leptotene/zygotene, green = diplotene, and blue = round spermatids). d) Select enriched

Biological Process GO terms (ranked by FDR) for transgressive DE genes between hybrid

dwarf hamsters and *P. campbelli* in diplotene spermatocytes. Included terms shown are the

result of the highlight function within gProfiler2 which collapses GO terms in a two-step

clustering algorithm (Kolberg *et al.* 2020). Point size corresponds to the number of genes

belonging to each GO term, and terms are plotted by the fold enrichment of the GO term in the

dataset relative to the provided gene backgrounds.

PAR expression was not disrupted in sterile hybrid dwarf hamsters

Finally, we tested the hypothesis that hybrid sterility in dwarf hamsters may be

correlated with asynapsis of the X and Y chromosomes because of divergence in the

pseudoautosomal region (PAR) which prevents proper chromosome pairing (Bikchurina *et al.*

2018). The PAR is the only portion of the sex chromosomes that is able to synapse during

routine spermatogenesis, and PAR genes are assumed to escape silencing by MSCI

(Raudsepp and Chowdhary 2015). However, because XY asynapsis is common in dwarf

hamster hybrids, Bikchurina et al. (2018) hypothesized that MSCI may extend to the PAR of

hybrid male dwarf hamsters, resulting in the silencing of PAR genes in hybrids that may be

critical to meiosis (Figure 6a). To test this hypothesis, we compared the expression of genes

located in the dwarf hamster PAR (Moore et al. 2022) between parental dwarf hamster species

and hybrid offspring across the timeline of spermatogenesis. Specifically, we hypothesized that

if XY asynapsis results in an extension of MSCI to the PAR in hybrid dwarf hamsters, then

hybrids should have similar PAR gene expression to parents early in meiosis before

homologous chromosome synapse during pachytene. This should be followed by the silencing

of PAR genes in hybrids, but not parent species, during diplotene. If XY asynapsis does not

alter the regulation of the PAR in hybrids, then we may see two possible patterns. First, if all

PAR genes are critical to the later stages of spermatogenesis, then PAR genes in both hybrids

and parents should be uniformly expressed in diplotene. Alternatively, if PAR genes are not

critical to the later stages of spermatogenesis, then hybrids and parent species should have

similar PAR gene expression, and not all PAR genes may be expressed during diplotene.

Figure 6. PAR gene expression is not disrupted during spermatogenesis in hybrid dwarf

hamsters. a) Hypothesized types of PAR gene expression across spermatogenesis. If MSCI

extends to the entire X chromosome, then PAR genes would show some level of expression

early in spermatogenesis which would then drop to zero in diplotene when MSCI occurs. If the

PAR escapes silencing by MSCI and if PAR genes are critical to spermatogenesis, then we

would expect PAR genes to be uniformly expressed in diplotene when the rest of the X

chromosome is silenced. Finally, if the PAR escapes silencing by MSCI but all PAR genes are

not critical to spermatogenesis, then we would expect some PAR genes to be expressed and

some to be not expressed in diplotene. b) Observed patterns of PAR gene expression (as mean

normalized RPKM across individuals) in parental species and hybrid dwarf hamsters for all PAR

genes (indicated by line type and color) across spermatogenesis (SP = spermatogonia, LZ =

leptotene/zygotene, and DIP = diplotene).

We did not find evidence supporting PAR-wide silencing in dwarf hamster hybrids

during diplotene suggesting that MSCI is not extended to the PAR in dwarf hamster hybrids

because of XY asynapsis (Figure 6b). Furthermore, we do not see PAR-wide expression of

genes during diplotene in hybrids or parents, indicating that not all PAR genes are critical to

the progression of spermatogenesis in dwarf hamsters. In general, most PAR gene expression

followed similar trends between hybrids and parent species. Two PAR genes were differentially

expressed between hybrids and *P. campbelli* during diplotene (*Ndrg2* and Psun_G000022883;

Table S3 in File S1), but these genes were still expressed in hybrids. Further, an association

between PAR misregulation during the early stages of spermatogenesis and hybrid male

sterility also seems unlikely as only one gene, *Gprin1*, showed transgressive differential

expression in early meiosis between both parent species and hybrids (Table S3 in File S1).

Thus, based on the current annotation of the PAR in *P. sungorus*, we currently find no direct

evidence linking improper silencing of PAR genes to hybrid male sterility in dwarf hamsters.

DISCUSSION

We used a comparative approach to understand common gene expression phenotypes

associated with hybrid male sterility in two divergent rodent crosses. We characterized the

asymmetry in the expression patterns of transgressive genes, how misexpression changed

over developmental timelines, and how the X chromosome and autosomes differed in both of

these aspects. We found that while there were similarities in hybrid expression phenotypes in

house mice and dwarf hamsters, there were also differences in the timing and chromosomal

distribution of disrupted gene expression that point towards different underlying mechanisms

behind hybrid male sterility.

Asymmetry and developmental timing of misexpression in hybrids

We first investigated patterns of transgressive gene misexpression in sterile male

hybrids. Studies of transgressive misexpression in sterile or inviable hybrids have often

focused on whether hybrid expression is biased towards over or underexpression, with the

hypothesis that expression may be biased towards overexpression if hybrid incompatibilities

disrupt repressive gene regulatory elements (Meiklejohn *et al.* 2014; Barreto *et al.* 2015; Larson

et al. 2017). In house mice, there is strong support for overexpression of both autosomal and

X-linked genes in sterile F1 hybrids (Mack *et al.* 2016; Larson *et al.* 2017, 2022; Hunnicutt *et al.*

2022). Surprisingly, we found that in dwarf hamster hybrids, there was nearly uniform

downregulation of transgressive DE genes in mitotic and early meiotic cell populations,

suggesting that a loss of regulatory repression is not an inevitable outcome of hybrid genomes.

Hybrid house mice expression is also more similar to the parent with the same X chromosome,

M. m. musculus, than to the parent with a different X chromosome, *M. m. domesticus* (Figure

4b and S2 in File S1; Larson *et al.* 2017). Further work in house mice has shown that F1 hybrid

expression patterns depend on both autosomal background and sex chromosome mismatch

(Kopania *et al.* 2022b). In contrast, hybrid dwarf hamsters showed similar levels of

misexpression in both the *P. campbelli* and *P. sungorus* comparisons. Determining what

factors shape the misexpression of parental alleles is a fruitful area of future research.

Asymmetric patterns of misexpression have been found in many hybrids, including

underexpression in sterile *Drosophila* hybrids (Michalak and Noor 2003; Haerty and Singh

2006; Llopart 2012) and sterile introgression lines of tomato (Guerrero *et al.* 2016) and

Drosophila (Meiklejohn *et al.* 2014), but overexpression has also been found in other sterile

hybrids (Llopart 2012; Davis *et al.* 2015). In many of these studies, patterns of misexpression

may be complicated by differences in cell composition or differences in the developmental

timeline of sterile hybrids and their parents (Good *et al.* 2010; Wei *et al.* 2014; Montgomery and

Mank 2016; Kerwin and Sweigart 2020; Hunnicutt *et al.* 2022). The variation we and others

have found in hybrid expression phenotypes suggests that the mechanisms of disrupted

expression are complex, even within groups with relatively shallow divergence times, such as

rodents, and we need more data from diverse hybrid sterility systems to begin to understand

common drivers of transgressive hybrid misexpression.

The downregulation we observed in early spermatogenesis in hybrid dwarf hamsters

could be due to impaired transcription factor binding with promoter or enhancer elements (Oka

et al. 2014; Guerrero *et al.* 2016) or disrupted epigenetic silencing. Disruption of epigenetic

regulation of gene expression has been increasingly linked to hybrid dysfunction in plants

(Shivaprasad *et al.* 2012; Lafon-Placette and Köhler 2015; Zhu *et al.* 2017), especially

polyploids (Paun *et al.* 2007), and may also contribute to hybrid male sterility in *Drosophila*

(Bayes and Malik 2009) and cattle x yak hybrids (Luo *et al.* 2022). At least one known

chromatin difference, an expansion of the heterochromatin-enriched Xp arm of the X

chromosome, has been documented between parental dwarf hamster species (Gamperl *et al.*

1977; Haaf *et al.* 1987). However, it is unknown what the functional consequences of this

chromatin state divergence or other diverged epigenetic regulatory mechanisms, such as

methylation, may be in hybrid dwarf hamster spermatogenesis, and further work is needed to

distinguish between potential mechanisms underlying the observed genome-wide

downregulation.

Spermatogenesis as a developmental process may be sensitive to disruption (Lifschytz

and Lindsley 1972; Wu and Davis 1993), but it remains an open question whether specific

stages of spermatogenesis, or developmental processes more broadly, may be more prone to

the accumulation of hybrid incompatibilities. In general, earlier developmental stages are

thought to be under greater pleiotropic constraint and less prone to disruption (Cutter and

Bundus 2020). With the progression of mouse spermatogenesis, pleiotropy decreases (as

approximated by increases in tissue specificity; Murat *et al.* 2023) and the rate of protein-

coding evolution increases (Larson *et al.* 2016; Kopania *et al.* 2022a; Murat *et al.* 2023), which

may make the later stages of spermatogenesis more prone to accumulating hybrid

incompatibilities. Indeed, we found fewer DE genes both between parent species and in sterile

hybrids for both dwarf hamsters and house mice during the early stages of spermatogenesis

than in later stages (Figures S9 and S10), and hybrid misexpression greatly exceeds parental

expression divergence in late spermatogenesis. When examining transgressive DE genes

shared between analogous cell types in dwarf hamster and house mouse hybrids, we found

that there were similar or fewer shared genes than expected by chance during early

spermatogenesis but more shared genes than expected by chance, especially on the X

chromosome, in later spermatogenesis, suggesting that disrupted expression of shared genes

of large effect during early spermatogenesis is unlikely to be responsible for the repeated

evolution of hybrid male sterility in these species.

Despite general similarities in patterns of misexpression across spermatogenesis in

hybrids, studies in house mice suggest that early spermatogenesis may be tolerant of some

misregulation as low levels of gene misexpression in early meiotic stages does not always

correlate with a complete cessation of sperm development (Oka *et al.* 2010; Ishishita *et al.*

2015; Mipam *et al.* 2023). However, the patterns we find in dwarf hamsters suggests that

spermatogenesis may be disrupted between zygotene and diplotene cell stages from early

misexpression. Dwarf hamster hybrid diplotene cell populations had X-linked and autosomal

gene expression profiles which more closely resemble parental leptotene/zygotene cell

populations than either diplotene or postmeiotic cell populations. Furthermore, transgressive

DE genes in hybrids during diplotene were enriched for genes associated with cell

differentiation, proliferation, and programmed cell death, suggesting misexpression during this

stage could be a consequence of a stalling or breakdown of early meiosis in hybrid dwarf

hamsters. It's unclear what underlying genomic mechanisms could result in this breakdown,

but it is possible that this disruption could potentially act as a major contributor to hybrid

sterility in this system. Ultimately, we find that spermatogenesis is a complex and rapidly

evolving developmental program that may provide many potential avenues across its timeline

for the evolution of hybrid incompatibilities.

Abnormal sex chromosome expression patterns differ between dwarf hamster and house

mouse hybrids

The sex chromosomes play a central role in speciation, an observation which has been

supported by both Haldane's rule (Haldane 1922) and the large X-effect on hybrid male sterility

(Coyne and Orr 1989). Misregulation of the X chromosome may contribute to hybrid sterility in

several species pairs (Davis *et al.* 2015; Morgan *et al.* 2020; Sánchez-Ramírez *et al.* 2021). The

X chromosome is transcriptionally repressed during routine spermatogenesis in many

organisms including eutherian mammals (McKee and Handel 1993), monotremes (Murat *et al.*

2023), *Drosophila* (Landeen *et al.* 2016), grasshoppers (Viera *et al.* 2021), mosquitos (Taxiarchi

et al. 2019), and nematodes (Rappaport *et al.* 2021). Because of the ubiquity of X chromosome

repression during spermatogenesis, disruption of transcriptional repression could be a

widespread regulatory phenotype in sterile hybrids (Lifschytz and Lindsley 1972; Larson *et al.*

2018). In sterile hybrid mice, disrupted X repression (disrupted MSCI) leads to the

overexpression of the normally silenced X chromosome during diplotene (Good *et al.* 2010;

Bhattacharyya *et al.* 2013; Campbell *et al.* 2013; Turner and Harr 2014; Larson *et al.* 2017,

2022). We also found overexpression of the X chromosome during diplotene in sterile hybrid

dwarf hamsters, but in a manner inconsistent with sterile hybrid house mice. Both the X

chromosome and autosomes are overexpressed in dwarf hamster hybrid diplotene cell

populations to a greater extent on average than was observed in house mice, and importantly,

X-linked overexpression was more variable in dwarf hamster hybrids than house mice hybrids.

In fact, some dwarf hamster hybrids had wildly overexpressed X chromosomes while others

appeared to have properly silenced X chromosomes. Our expression correlation and gene set

analyses of hybrid diplotene cell populations provide additional evidence that the genes

overexpressed in hybrid hamster diplotene are different than those overexpressed in house

mouse hybrids, sharing more similarity to the earlier meiotic cell types than downstream

postmeiotic cell types. Overall, our results indicate fundamentally different patterns of X-linked

overexpression in both systems, with X-linked overexpression in dwarf hamster hybrids being

inconsistent with disrupted MSCI patterns observed in house mouse hybrids.

Much of what we know about the genomic architecture and the role of sex

chromosome misregulation in hybrid male sterility in mammals comes from decades of work

that have shown a major gene, *Prdm9*, and its X chromosome modulator, *Hstx2*, may be

responsible for most F1 hybrid male sterility in house mice (Forejt *et al.* 1991, 2021; Trachtulec

et al. 1997; Mihola *et al.* 2009; Lustyk *et al.* 2019). *Prdm9* directs the location of double strand

breaks during meiotic recombination (Mihola *et al.* 2009; Oliver *et al.* 2009; Smagulova *et al.*

2016). In hybrid mice, divergence at *Prdm9* binding sites leads to asymmetric double-stranded

breaks and results in autosomal asynapsis, triggering Meiotic Silencing of Unsynapsed

Chromatin, shutting down transcription on asynapsed autosomes using the same cellular

machinery as MSCI (Turner 2015), and eventually meiotic arrest and cell death (Bhattacharyya

et al. 2013; Forejt *et al.* 2021). This process is associated with the disruption of MSCI and a

characteristic overexpression of the X chromosome during meiosis (Good *et al.* 2010;

Bhattacharyya *et al.* 2013; Campbell *et al.* 2013; Turner and Harr 2014; Larson *et al.* 2017,

2022), but whether disrupted MSCI directly contributes to hybrid male sterility or is simply a

downstream consequence of *Prdm9* divergence is still uncertain (Forejt *et al.* 2021).

Whether we should have expected patterns of disrupted sex chromosome expression

in sterile hybrid hamsters to be the same as house mice is unclear. The sex chromosomes in

pachytene cells of hybrid dwarf hamsters display normal γH2AFX staining (Ishishita *et al.* 2015;

Bikchurina *et al.* 2018), a key marker in MSCI (Abe *et al.* 2022), which may indicate that the

hybrid sex chromosomes are properly silenced. Additionally, autosomal asynapsis is rarely

observed in hybrid dwarf hamsters, and asynapsis is almost exclusive to the sex chromosomes

(Ishishita *et al.* 2015; Bikchurina *et al.* 2018). This contrasts *Prdm9*-mediated sterility in house

mice, where hybrid autosomes are often asynapsed and decorated with γH2AFX

(Bhattacharyya *et al.* 2013; Forejt *et al.* 2021). Mechanisms other than *Prdm9* may also disrupt

MSCI and result in sterility, such as macrosatellite copy number divergence (Bredemeyer *et al.*

2021) and X-autosome translocations that impair synapsis (Homolka *et al.* 2007), although

there is no evidence for X-autosome translocations between these two species of dwarf

hamsters (Moore *et al.* 2022). Thus, while MSCI may be a major target for the accumulation of

reproductive barriers between species in many mammalian systems, either through *Prdm9*

divergence or alternative mechanisms, our results suggest that sterility in dwarf hamsters has a

more composite regulatory basis.

Another mechanism often proposed to underlie mammalian male hybrid sterility,

especially in rodents, is divergence in the PAR between parental species. The PAR is the only

portion of the sex chromosomes which can synapse during spermatogenesis, and it is still

unclear if the regulation of the PAR is uniformly detached from MSCI across divergent

mammalian sex chromosome systems (Raudsepp and Chowdhary 2015). We find no evidence

that PAR-specific misregulation is associated with hybrid sterility in dwarf hamsters. PAR

genes are not silenced in hybrids or parents, a pattern that is inconsistent with MSCI that has

extended to the PAR due to XY asynapsis, and further, expression of PAR genes in hybrids

differs little from parental PAR expression. While we find no evidence that PAR misregulation

per se is associated with hybrid sterility in this system, we cannot rule out the possibility that

structural and sequence divergence between the PARs of *P. sungorus* and *P. campbelli* may

be associated with hybrid sterility. Structural and sequence divergence in the PAR has been

hypothesized to activate the meiotic spindle checkpoint by interfering with proper pairing of

sex chromosomes (Burgoyne *et al.* 2009; Dumont 2017). The PAR evolves rapidly in rodents

(White *et al.* 2012b; Raudsepp and Chowdhary 2015; Morgan *et al.* 2019), and this elevated

divergence may underlie sex chromosome asynapsis and apoptosis in several hybrid mouse

crosses (Matsuda *et al.* 1991; Oka *et al.* 2010; White *et al.* 2012a; Dumont 2017). Furthermore,

divergence in the mouse PAR has been implicated in spermatogenic defects in crosses where

Prdm9-divergence is minimal, such as between closely related subspecies (Dumont 2017) or in

mice with genetically-modified *Prdm9* alleles (Davies *et al.* 2021). Meiosis is likely tolerant to

some degree of divergence in the PAR (Morgan *et al.* 2019), but exact limits are currently

unknown. At this time, thorough analysis of structural and sequence divergence between the

PARs in dwarf hamsters is challenging as the PAR is notoriously difficult to assemble (but see

Kasahara *et al.* 2022), and there are annotation gaps in the current assembly of the PAR in

dwarf hamsters. In sum, we find no clear pattern of regulatory disruption of PAR genes in

sterile hybrid dwarf hamsters, though this result may change pending further refinement of the

PAR annotation.

Conclusions

Cell-specific approaches for quantifying expression phenotypes are powerful tools for

providing insight into the underlying mechanisms behind hybrid dysfunction (Hunnicutt *et al.*

2022), especially in systems where it remains difficult to interrogate the underlying genomic

architecture of these traits. Using a contrast of dwarf hamster and house mouse hybrids, we

have shown that transgressive overexpression is not an inevitable outcome of hybridization,

that misexpression resulting from hybrid incompatibilities may be likely to arise in differing

stages of spermatogenesis, and that disrupted sex chromosome silencing does not appear to

play an equal role in sterility between these two systems. Both the expression phenotypes we

observed here and histological evidence from other studies (Ishishita *et al.* 2015; Bikchurina *et*

al. 2018) suggest that several reproductive barriers are acting during spermatogenesis in dwarf

hamster hybrids. It has become increasingly apparent as more study systems are investigated

that the genetic basis of postzygotic species barriers are often complex and polymorphic

(Cutter 2012; Coughlan and Matute 2020), and implementing approaches which account for

the developmental complexities of hybrid dysfunction, as we have done here, will allow us to

make further advances in understanding the processes of speciation.

Acknowledgements:

This work was supported by a National Science Foundation Graduate Research Fellowship to

KEH (DGE-2034612) and the following grants awarded to KEH: Sigma Xi Grant-in-aid of

Research, American Society of Mammalogists Grant-in-Aid of Research Grant, and a Society

for the Study of Evolution R.C. Lewontin Early Award. Additionally, this work was supported by

grants from the Eunice Kennedy Shriver National Institute of Child Health and Human

Development of the National Institutes of Health (R01-HD073439, R01-HD094787) to JMG, and

the National Science Foundation to ELL (DEB-2012041). We would like to thank members of

the Larson and Good labs, as well as Jonathan Velotta and members of the Velotta lab, for

feedback on this project, Gabrielle Welsh for technical assistance, Ivan Kovanda for support

using the University of Denver Research Data Analysis Cluster, and Pamela K. Shaw and the

UM Fluorescence Cytometry Core supported by an Institutional Development Award from the

Author contributions:

KEH, JMG, and ELL conceived of the study. CC, SK, ELL, and KEH conducted lab work. KEH

conducted the analyses. KEH and ELL wrote the manuscript with input from ECM, CC, SK, and

JMG.

Data accessibility:

Hamster sequence data are available through the NCBI SRA under accession number

PRJNA1024468, and mouse sequence data are available under SRA accession PRJNA296926.

File S1 contains all supplemental figures. Code used for the analyses and fertility phenotypic

data are available from GitHub (https://github.com/KelsieHunnicutt/dwarf_hamster_hybrids).

LITERATURE CITED

907
908

Abe H., Y.-H. Yeh, Y. Munakata, K.-I. Ishiguro, P. R. Andreassen, *et al.*, 2022 Active DNA

damage response signaling initiates and maintains meiotic sex chromosome

inactivation. Nat. Commun. 13: 7212.

Anders S., and W. Huber, 2010 Differential expression analysis for sequence count data.

Nature Precedings 1–1.

Barreto F. S., R. J. Pereira, and R. S. Burton, 2015 Hybrid dysfunction and physiological

compensation in gene expression. Mol. Biol. Evol. 32: 613–622.

Bayes J. J., and H. S. Malik, 2009 Altered heterochromatin binding by a hybrid sterility protein

in Drosophila sibling species. Science 326: 1538–1541.

Benjamini Y., and Y. Hochberg, 1995 Controlling the false discovery rate: a practical and

powerful approach to multiple testing. J. R. Stat. Soc. 57: 289–300.

Bhattacharyya T., S. Gregorova, O. Mihola, M. Anger, J. Sebestova, *et al.*, 2013 Mechanistic

basis of infertility of mouse intersubspecific hybrids. Proc. Natl. Acad. Sci. U. S. A. 110:

E468–E477.

Bikchurina T. I., K. V. Tishakova, E. A. Kizilova, S. A. Romanenko, N. A. Serdyukova, *et al.*,

2018 Chromosome synapsis and recombination in male-sterile and female-fertile

interspecies hybrids of the dwarf hamsters (Phodopus, Cricetidae). Genes 9: 227.

Bolger A. M., M. Lohse, and B. Usadel, 2014 Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics 30: 2114–2120.

Bredemeyer K. R., C. M. Seabury, M. J. Stickney, J. R. McCarrey, B. M. vonHoldt, *et al.*, 2021

929   Rapid macrosatellite evolution promotes X-linked hybrid male sterility in a feline

930   interspecies cross. Mol. Biol. Evol. 38: 5588–5609.

931   Brekke T. D., and J. M. Good, 2014 Parent-of-origin growth effects and the evolution of hybrid

932   inviability in dwarf hamsters. Evolution 68: 3134–3148.

933   Brekke T. D., L. A. Henry, and J. M. Good, 2016 Genomic imprinting, disrupted placental

934   expression, and speciation. Evolution 70: 2690–2703.

935   Brekke T. D., K. A. Steele, and J. F. Mulley, 2018 Inbred or outbred? Genetic diversity in

936   laboratory rodent colonies. G3 8: 679–686.

937   Brekke T. D., E. C. Moore, S. C. Campbell-Staton, C. M. Callahan, Z. A. Cheviron, *et al.*, 2021 X

938   chromosome-dependent disruption of placental regulatory networks in hybrid dwarf

939   hamsters. Genetics 218: iyab043.

940   Burgoyne P. S., 1982 Genetic homology and crossing over in the X and Y chromosomes of

941   mammals. Hum. Genet. 61: 85–90.

942   Burgoyne P. S., S. K. Mahadevaiah, and J. M. A. Turner, 2009 The consequences of asynapsis

943   for mammalian meiosis. Nat. Rev. Genet. 10: 207–216.

944   Campbell P., J. M. Good, and M. W. Nachman, 2013 Meiotic sex chromosome inactivation is

945   disrupted in sterile hybrid male house mice. Genetics 193: 819–828.

946   Civetta A., 2016 Misregulation of gene expression and sterility in interspecies hybrids: causal

947   links and alternative hypotheses. J. Mol. Evol. 82: 176–182.

948   Coughlan J. M., and D. R. Matute, 2020 The importance of intrinsic postzygotic barriers

949   throughout the speciation process. Philos. Trans. R. Soc. Lond. B Biol. Sci. 375:

20190533.

Coyne J. A., and H. A. Orr, 1989 Two rules of speciation, pp. 180–207 in *Speciation and its*

consequences, edited by Otte D., Endler J. A. Sinauer Associates, Sunderland, MA.

Cutter A. D., 2012 The polymorphic prelude to Bateson-Dobzhansky-Muller incompatibilities.

Trends Ecol. Evol. 27: 209–218.

Cutter A. D., and J. D. Bundus, 2020 Speciation and the developmental alarm clock. Elife 9:

e56276.

Danecek P., A. Auton, G. Abecasis, C. A. Albers, E. Banks, *et al.*, 2011 The variant call format

and VCFtools. Bioinformatics 27: 2156–2158.

Davies B., E. Hatton, N. Altemose, J. G. Hussin, F. Pratto, *et al.*, 2016 Re-engineering the zinc

fingers of PRDM9 reverses hybrid sterility in mice. Nature 530: 171–176.

Davies B., A. Gupta Hinch, A. Cebrian-Serrano, S. Alghadban, P. W. Becker, *et al.*, 2021

Altering the binding properties of PRDM9 partially restores fertility across the species

boundary. Mol. Biol. Evol. 38: 5555–5562.

Davis B. W., C. M. Seabury, W. A. Brashear, G. Li, M. Roelke-Parker, *et al.*, 2015 Mechanisms

underlying mammalian hybrid sterility in two feline interspecies models. Mol. Biol. Evol.

32: 2534–2546.

Dumont B. L., 2017 Meiotic consequences of genetic divergence across the murine

pseudoautosomal region. Genetics 205: 1089–1100.

Ellis N., and P. N. Goodfellow, 1989 The mammalian pseudoautosomal region. Trends Genet.

5: 406–410.

Forejt J., V. Vincek, J. Klein, H. Lehrach, and M. Loudová-Micková, 1991 Genetic mapping of

the t-complex region on mouse chromosome 17 including the Hybrid sterility-1 gene.

Mamm. Genome 1: 84–91.

Forejt J., P. Jansa, and E. Parvanov, 2021 Hybrid sterility genes in mice (Mus musculus): a

peculiar case of PRDM9 incompatibility. Trends Genet. 37: 1095–1108.

Gamperl R., G. Vistorin, and W. Rosenkranz, 1977 New observations on the karyotype of the

Djungarian hamster, Phodopus sungorus. Experientia 33: 1020–1021.

Getun I. V., B. Torres, and P. R. J. Bois, 2011 Flow cytometry purification of mouse meiotic

cells. J. Vis. Exp. e2602.

Good J. M., M. A. Handel, and M. W. Nachman, 2008 Asymmetry and polymorphism of hybrid

male sterility during the early stages of speciation in house mice. Evolution 62: 50–65.

Good J. M., T. Giger, M. D. Dean, and M. W. Nachman, 2010 Widespread over-expression of

the X chromosome in sterile F1 hybrid mice. PLoS Genet. 6: e1001148.

Green C. D., Q. Ma, G. L. Manske, A. N. Shami, X. Zheng, *et al.*, 2018 A comprehensive

roadmap of murine spermatogenesis defined by single-cell RNA-Seq. Dev. Cell 46:

651-667.e10.

Gu Z., R. Eils, and M. Schlesner, 2016 Complex heatmaps reveal patterns and correlations in

multidimensional genomic data. Bioinformatics 32: 2847–2849.

Guerrero R. F., A. L. Posto, L. C. Moyle, and M. W. Hahn, 2016 Genome-wide patterns of

regulatory divergence revealed by introgression lines. Evolution 70: 696–706.

Haaf T., H. Weis, and M. Schmid, 1987 A comparative cytogenetic study on the mitotic and

meiotic chromosomes in hamster species of the genus Phodopus (Rodentia,

Cricetinae). Zeitschrift für Säugetierkunde 52: 281–290.

Haerty W., and R. S. Singh, 2006 Gene regulation divergence is a major contributor to the

evolution of Dobzhansky-Muller incompatibilities between species of Drosophila. Mol.

Biol. Evol. 23: 1707–1714.

Haldane J. B. S., 1922 Sex ratio and unisexual sterility in hybrid animals. J. Genet. 12: 101–

109.

Handel M. A., 2004 The XY body: a specialized meiotic chromatin domain. Exp. Cell Res. 296:

57–63.

Hermann B. P., K. Cheng, A. Singh, L. Roa-De La Cruz, K. N. Mutoji, *et al.*, 2018 The

mammalian spermatogenesis single-cell transcriptome, from spermatogonial stem cells

to spermatids. Cell Rep. 25: 1650-1667.e8.

Holt J., S. Huang, L. McMillan, and W. Wang, 2013 Read annotation pipeline for high-

throughput sequencing data, pp. 605–612 in *Proceedings of the International*

Conference on Bioinformatics, Computational Biology and Biomedical Informatics -

BCB'13, ACM Press, New York, New York, USA.

Homolka D., R. Ivanek, J. Capkova, P. Jansa, and J. Forejt, 2007 Chromosomal rearrangement

interferes with meiotic X chromosome inactivation. Genome Res. 17: 1431–1437.

Huang S., J. Holt, C.-Y. Kao, L. McMillan, and W. Wang, 2014 A novel multi-alignment pipeline

for high-throughput sequencing data. Database 2014: bau057.

Hunnicutt K. E., J. M. Good, and E. L. Larson, 2022 Unraveling patterns of disrupted gene

expression across a complex tissue. Evolution 76: 275–291.

Ishishita S., K. Tsuboi, N. Ohishi, K. Tsuchiya, and Y. Matsuda, 2015 Abnormal pairing of X and

Y sex chromosomes during meiosis I in interspecific hybrids of Phodopus campbelli

and P. sungorus. Sci. Rep. 5: 9435.

Kasahara T., K. Mekada, K. Abe, A. Ashworth, and T. Kato, 2022 Complete sequencing of the

mouse pseudoautosomal region, the most rapidly evolving 'chromosome.' bioRxiv

2022.03.26.485930.

Kerwin R. E., and A. L. Sweigart, 2020 Rampant Misexpression in a Mimulus (Monkeyflower)

Introgression Line Caused by Hybrid Sterility, Not Regulatory Divergence. Mol. Biol.

Evol. 37: 2084–2098.

Kim D., J. M. Paggi, C. Park, C. Bennett, and S. L. Salzberg, 2019 Graph-based genome

alignment and genotyping with HISAT2 and HISAT-genotype. Nat. Biotechnol. 37: 907–

915.

Kolberg L., U. Raudvere, I. Kuzmin, J. Vilo, and H. Peterson, 2020 gprofiler2 -- an R package

for gene list functional enrichment analysis and namespace conversion toolset

g:Profiler. F1000Res. 9. https://doi.org/10.12688/f1000research.24956.2

Kopania E. E. K., E. L. Larson, C. Callahan, S. Keeble, and J. M. Good, 2022a Molecular

evolution across mouse spermatogenesis. Mol. Biol. Evol. 39: msac023.

Kopania E. E. K., E. M. Watson, C. C. Rathje, B. M. Skinner, P. J. I. Ellis, *et al.*, 2022b The

contribution of sex chromosome conflict to disrupted spermatogenesis in hybrid house

mice. Genetics 222: iyac151.

Korunes K. L., and K. Samuk, 2021 pixy: Unbiased estimation of nucleotide diversity and

divergence in the presence of missing data. Mol. Ecol. Resour. 21: 1359–1368.

Kousathanas A., D. L. Halligan, and P. D. Keightley, 2014 Faster-X adaptive protein evolution in

house mice. Genetics 196: 1131–1143.

Lafon-Placette C., and C. Köhler, 2015 Epigenetic mechanisms of postzygotic reproductive

isolation in plants. Curr. Opin. Plant Biol. 23: 39–44.

Landeen E. L., C. A. Muirhead, L. Wright, C. D. Meiklejohn, and D. C. Presgraves, 2016 Sex

chromosome-wide transcriptional suppression and compensatory cis-regulatory

evolution mediate gene expression in the Drosophila male germline. PLoS Biol. 14:

e1002499.

Larson E. L., D. Vanderpool, S. Keeble, M. Zhou, B. A. J. Sarver, *et al.*, 2016 Contrasting levels

of molecular evolution on the mouse X chromosome. Genetics 203: 1841–1857.

Larson E. L., S. Keeble, D. Vanderpool, M. D. Dean, and J. M. Good, 2017 The composite

regulatory basis of the large X-effect in mouse speciation. Mol. Biol. Evol. 34: 282–295.

Larson E. L., E. E. K. Kopania, and J. M. Good, 2018 Spermatogenesis and the evolution of

mammalian sex chromosomes. Trends Genet. 34: 722–732.

Larson E. L., E. E. K. Kopania, K. E. Hunnicutt, D. Vanderpool, S. Keeble, *et al.*, 2022 Stage-

specific disruption of X chromosome expression during spermatogenesis in sterile

house mouse hybrids. G3 12: jkab407.

Liao Y., G. K. Smyth, and W. Shi, 2014 featureCounts: an efficient general purpose program for

assigning sequence reads to genomic features. Bioinformatics 30: 923–930.

Lifschytz E., and D. L. Lindsley, 1972 The role of X-chromosome inactivation during
spermatogenesis. Proc. Natl. Acad. Sci. U. S. A. 69: 182–186.

Llopart A., 2012 The rapid evolution of X-linked male-biased gene expression and the large-X
effect in Drosophila yakuba, D. santomea, and their hybrids. Mol. Biol. Evol. 29: 3873–
3886.

Luo H., T. Mipam, S. Wu, C. Xu, C. Yi, *et al.*, 2022 DNA methylome of primary spermatocyte
reveals epigenetic dysregulation associated with male sterility of cattleyak.
Theriogenology 191: 153–167.

Lustyk D., S. Kinský, K. K. Ullrich, M. Yancoskie, L. Kašíková, *et al.*, 2019 Genomic structure of
Hstx2 modifier of Prdm9-dependent hybrid male sterility in mice. Genetics 213: 1047–
1063.

Mack K. L., P. Campbell, and M. W. Nachman, 2016 Gene regulation and speciation in house
mice. Genome Res. 26: 451–461.

Mack K. L., and M. W. Nachman, 2017 Gene regulation and speciation. Trends Genet. 33: 68–
80.

Matsuda Y., T. Hirobe, and V. M. Chapman, 1991 Genetic basis of X-Y chromosome
dissociation and male sterility in interspecific hybrids. Proc. Natl. Acad. Sci. U. S. A. 88:
4850–4854.

McCarthy D. J., Y. Chen, and G. K. Smyth, 2012 Differential expression analysis of multifactor
RNA-Seq experiments with respect to biological variation. Nucleic Acids Res. 40: 4288–
4297.

McKee B. D., and M. A. Handel, 1993 Sex chromosomes, recombination, and chromatin

conformation. Chromosoma 102: 71–80.

McManus C. J., J. D. Coolon, M. O. Duff, J. Eipper-Mains, B. R. Graveley, *et al.*, 2010

Regulatory divergence in Drosophila revealed by mRNA-seq. Genome Res. 20: 816–

825.

Meiklejohn C. D., J. D. Coolon, D. L. Hartl, and P. J. Wittkopp, 2014 The roles of cis- and trans-

regulation in the evolution of regulatory incompatibilities and sexually dimorphic gene

expression. Genome Res. 24: 84–95.

Michalak P., and M. A. F. Noor, 2003 Genome-wide patterns of expression in Drosophila pure

species and hybrid males. Mol. Biol. Evol. 20: 1070–1076.

Mihola O., Z. Trachtulec, C. Vlcek, J. C. Schimenti, and J. Forejt, 2009 A mouse speciation

gene encodes a meiotic histone H3 methyltransferase. Science 323: 373–375.

Mipam T., X. Chen, W. Zhao, P. Zhang, Z. Chai, *et al.*, 2023 Single-cell transcriptome analysis

and in vitro differentiation of testicular cells reveal novel insights into male sterility of the

interspecific hybrid cattle-yak. BMC Genomics 24: 149.

Montgomery S. H., and J. E. Mank, 2016 Inferring regulatory change from gene expression: the

confounding effects of tissue scaling. Mol. Ecol. 25: 5114–5128.

Moore E. C., G. W. C. Thomas, S. Mortimer, E. E. K. Kopania, K. E. Hunnicutt, *et al.*, 2022 The

evolution of widespread recombination suppression on the dwarf hamster (Phodopus) X

chromosome. Genome Biol. Evol. 14: evac080.

Morgan A. P., T. A. Bell, J. J. Crowley, and F. Pardo-Manuel de Villena, 2019 Instability of the

pseudoautosomal boundary in house mice. Genetics 212: 469–487.

Morgan K., B. Harr, M. A. White, B. A. Payseur, and L. M. Turner, 2020 Disrupted gene

networks in subfertile hybrid house mice. Mol. Biol. Evol. 37: 1547–1562.

Mugal C. F., M. Wang, N. Backström, D. Wheatcroft, M. Ålund, *et al.*, 2020 Tissue-specific

patterns of regulatory changes underlying gene expression differences among *Ficedula*

flycatchers and their naturally occurring F1 hybrids. Genome Res. 30: 1727–1739.

Murat F., N. Mbengue, S. B. Winge, T. Trefzer, E. Leushkin, *et al.*, 2023 The molecular evolution

of spermatogenesis across mammals. Nature 613: 308–316.

Namekawa S. H., P. J. Park, L.-F. Zhang, J. E. Shima, J. R. McCarrey, *et al.*, 2006 Postmeiotic

sex chromatin in the male germline of mice. Curr. Biol. 16: 660–667.

Neumann K., J. Michaux, V. Lebedev, N. Yigit, E. Colak, *et al.*, 2006 Molecular phylogeny of the

Cricetinae subfamily based on the mitochondrial cytochrome b and 12S rRNA genes

and the nuclear vWF gene. Mol. Phylogenet. Evol. 39: 135–148.

Ogle D., and M. D. Ogle, 2017 Package 'FSA.' CRAN Repos 1–206.

Oka A., A. Mita, Y. Takada, H. Koseki, and T. Shiroishi, 2010 Reproductive isolation in hybrid

mice due to spermatogenesis defects at three meiotic stages. Genetics 186: 339–351.

Oka A., and T. Shiroishi, 2014 Regulatory divergence of X-linked genes and hybrid male

sterility in mice. Genes Genet. Syst. 89: 99–108.

Oka A., T. Takada, H. Fujisawa, and T. Shiroishi, 2014 Evolutionarily diverged regulation of X-

chromosomal genes as a primal event in mouse reproductive isolation. PLoS Genet. 10:

e1004301.

Oksanen J., F. G. Blanchet, R. Kindt, P. Legendre, P. R. Minchin, *et al.*, 2013 Package 'vegan.'

Community ecology package, version 2: 1–295.

Oliver P. L., L. Goodstadt, J. J. Bayes, Z. Birtle, K. C. Roach, *et al.*, 2009 Accelerated evolution

of the Prdm9 speciation gene across diverse metazoan taxa. PLoS Genet. 5: e1000753.

Ortíz-Barrientos D., B. A. Counterman, and M. A. F. Noor, 2007 Gene expression divergence

and the origin of hybrid dysfunctions. Genetica 129: 71–81.

Paun O., M. F. Fay, D. E. Soltis, and M. W. Chase, 2007 Genetic and epigenetic alterations

after hybridization and genome doubling. Taxon 56: 649–656.

Presgraves D. C., 2008 Sex chromosomes and speciation in Drosophila. Trends Genet. 24:

336–343.

Ramm S. A., and L. Schärer, 2014 The evolutionary ecology of testicular function: size isn't

everything. Biol. Rev. Camb. Philos. Soc. 89: 874–888.

Rappaport Y., H. Achache, R. Falk, O. Murik, O. Ram, *et al.*, 2021 Bisection of the X

chromosome disrupts the initiation of chromosome silencing during meiosis in

Caenorhabditis elegans. Nat. Commun. 12: 4802.

Raudsepp T., and B. P. Chowdhary, 2015 The eutherian pseudoautosomal region. Cytogenet.

Genome Res. 147: 81–94.

Raymond C. S., M. W. Murphy, M. G. O'Sullivan, V. J. Bardwell, and D. Zarkower, 2000 Dmrt1,

a gene related to worm and fly sexual regulators, is required for mammalian testis

differentiation. Genes Dev. 14: 2587–2595.

Ritchie M. E., B. Phipson, D. Wu, Y. Hu, C. W. Law, *et al.*, 2015 limma powers differential

expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Res.

43: e47.

Robinson M. D., D. J. McCarthy, and G. K. Smyth, 2010 edgeR: a Bioconductor package for

differential expression analysis of digital gene expression data. Bioinformatics 26: 139–

140.

Sánchez-Ramírez S., J. G. Weiss, C. G. Thomas, and A. D. Cutter, 2021 Widespread

misregulation of inter-species hybrid transcriptomes due to sex-specific and sex-

chromosome regulatory evolution. PLoS Genet. 17: e1009409.

Scribner S. J., and K. E. Wynne-Edwards, 1994 Disruption of body temperature and behavior

rhythms during reproduction in dwarf hamsters (Phodopus). Physiol. Behav. 55: 361–

369.

Shivaprasad P. V., R. M. Dunn, B. A. Santos, A. Bassett, and D. C. Baulcombe, 2012

Extraordinary transgressive phenotypes of hybrid tomato are influenced by epigenetics

and small silencing RNAs. EMBO J. 31: 257–266.

Smagulova F., K. Brick, Y. Pu, R. D. Camerini-Otero, and G. V. Petukhova, 2016 The

evolutionary turnover of recombination hot spots contributes to speciation in mice.

Genes Dev. 30: 266–280.

Swanson M. T., C. H. Oliveros, and J. A. Esselstyn, 2019 A phylogenomic rodent tree reveals

the repeated evolution of masseter architectures. Proc. Biol. Sci. 286: 20190672.

Taxiarchi C., N. Kranjc, A. Kriezis, K. Kyrou, F. Bernardini, *et al.*, 2019 High-resolution

transcriptional profiling of Anopheles gambiae spermatogenesis reveals mechanisms of

sex chromosome regulation. Sci. Rep. 9: 14841.

Trachtulec Z., M. Mnuková-Fajdelová, R. M. Hamvas, S. Gregorová, W. E. Mayer, *et al.*, 1997

Isolation of candidate hybrid sterility 1 genes by cDNA selection in a 1.1 megabase pair

region on mouse chromosome 17. Mamm. Genome 8: 312–316.

Turner L. M., M. A. White, D. Tautz, and B. A. Payseur, 2014 Genomic networks of hybrid

sterility. PLoS Genet. 10: e1004162.

Turner L. M., and B. Harr, 2014 Genome-wide mapping in a house mouse hybrid zone reveals

hybrid sterility loci and Dobzhansky-Muller interactions. Elife 3: e02504.

Turner J. M. A., 2015 Meiotic silencing in mammals. Annu. Rev. Genet. 49: 395–412.

Vasimuddin M., S. Misra, H. Li, and S. Aluru, 2019 Efficient architecture-aware acceleration of

BWA-MEM for multicore systems, pp. 314–324 in *2019 IEEE International Parallel and*

Distributed Processing Symposium (IPDPS), IEEE.

Viera A., M. T. Parra, S. Arévalo, C. García de la Vega, J. L. Santos, *et al.*, 2021 X chromosome

inactivation during grasshopper spermatogenesis. Genes 12: 1844.

Wei K. H.-C., A. G. Clark, and D. A. Barbash, 2014 Limited gene misregulation is exacerbated

by allele-specific upregulation in lethal hybrids between Drosophila melanogaster and

Drosophila simulans. Mol. Biol. Evol. 31: 1767–1778.

White M. A., M. Stubbings, B. L. Dumont, and B. A. Payseur, 2012a Genetics and evolution of

hybrid male sterility in house mice. Genetics 191: 917–934.

White M. A., A. Ikeda, and B. A. Payseur, 2012b A pronounced evolutionary shift of the

pseudoautosomal region boundary in house mice. Mamm. Genome 23: 454–466.

Wittkopp P. J., B. K. Haerum, and A. G. Clark, 2004 Evolutionary changes in cis and trans gene

regulation. Nature 430: 85–88.

Wu C. I., and A. W. Davis, 1993 Evolution of postmating reproductive isolation: the composite

nature of Haldane's rule and its genetic bases. Am. Nat. 142: 187–212.

Zhu W., B. Hu, C. Becker, E. S. Doğan, K. W. Berendzen, *et al.*, 2017 Altered chromatin

compaction and histone methylation drive non-additive gene expression in an

interspecific Arabidopsis hybrid. Genome Biol. 18: 157.