Different complex regulatory phenotypes underlie hybrid male sterility in divergent rodent crosses

Kelsie E. Hunnicutt[*,1], Colin Callahan[2], Sara Keeble[2], Emily C. Moore[1,2], Jeffrey M. Good[2], and Erica L. Larson[*,1]

[1] University of Denver, Department of Biological Sciences, Denver, CO, 80208
[2] Division of Biological Sciences, University of Montana, Missoula, MT, 59812
*Corresponding authors

ORCID: 0000-0002-9674-0630 (KEH)
ORCID: 0000-0002-0385-1121 (CC)
ORCID: 0000-0002-7318-3122 (SK)
ORCID: 0000-0001-6166-3367 (ECM)
ORCID: 0000-0003-0707-5374 (JMG)
ORCID: 0000-0003-3006-645X (ELL)

Corresponding Authors:
Kelsie Hunnicutt
Department of Biological Sciences
University of Denver
Denver, CO 80208
Email: Kelsie.Hunnicutt@du.edu

Erica Larson
Department of Biological Sciences
University of Denver
Denver, CO 80208
Email: Erica.Larson@du.edu

Running title: Rodent hybrid male sterility

Key words: speciation, hybrid male sterility, gene expression, fluorescence activated cell sorting, meiotic sex chromosome inactivation, *Phodopus*

ABSTRACT

Hybrid incompatibilities are a critical component of species barriers and may arise due to negative interactions between divergent regulatory elements in parental species. We used a comparative approach to identify common themes in the regulatory phenotypes associated with hybrid male sterility in two divergent rodent crosses, dwarf hamsters and house mice. We investigated three potential characteristic gene expression phenotypes in hybrids including the propensity of transgressive differentially expressed genes in hybrids towards over or underexpression relative to parental species, the influence of developmental stage on the extent of misexpression, and the role of the sex chromosomes on misexpression phenotypes. In contrast to near pervasive overexpression in hybrid house mice, we found that misexpression in hybrid dwarf hamsters was dependent on developmental stage. In both house mouse and dwarf hamster hybrids, however, transgressive misexpression increased with the progression of spermatogenesis, although to varying extents and with potentially different consequences. In both systems, we detected sex-chromosome specific overexpression in stages of spermatogenesis where inactivated X chromosome expression was expected, but the hybrid overexpression phenotypes were fundamentally different. Importantly, misexpression phenotypes support the presence of multiple developmental blocks to spermatogenesis in dwarf hamster hybrids, including a potential role of meiotic stalling or breakdown early in spermatogenesis. Collectively, we demonstrate that while there are some similarities in hybrid expression phenotypes of house mice and dwarf hamsters, there are also clear differences that point towards unique mechanisms underlying hybrid male sterility in each system. Our results highlight the potential of comparative approaches in helping to understand the causes and consequences of disrupted gene expression in speciation.

INTRODUCTION

The evolution of postzygotic reproductive barriers, such as hybrid inviability and

sterility, is an important part of the speciation process, and identifying the genetic architecture

of hybrid incompatibilities has been a common goal uniting speciation research (Coughlan and

Matute 2020). While identifying the genetic basis of hybrid dysfunction remains difficult in many

systems, downstream regulatory phenotypes can provide insight into the underlying

mechanisms of speciation (Mack and Nachman 2017). An outstanding question surrounding

the role of disrupted gene regulation and speciation is whether the combination of two

divergent genomes in hybrids results in gene expression perturbations that are consistent or

repeatable across species. At the broadest level, it is unclear whether gene expression in

hybrids tends to be intermediate or transgressive (outside the range of parental gene

expression), whether transgressive gene expression is biased towards over or underexpression

(Ortíz-Barrientos *et al.* 2007), and whether misexpression tends to be modulated by *cis* or *trans*

regulatory elements (Wittkopp *et al.* 2004; McManus *et al.* 2010; Oka *et al.* 2014; Mack *et al.*

2016; Mugal *et al.* 2020; Kopania *et al.* 2022a). By investigating trends in the magnitude and

direction of transgressive expression across different hybrid systems, we can begin to

understand the evolutionary forces shaping regulatory-based hybrid incompatibilities. For

example, if expression in hybrids tends towards overexpression, this may mean that genes

contributing to hybrid incompatibilities are normally repressed in parental lineages (Meiklejohn

et al. 2014; Barreto *et al.* 2015; Larson *et al.* 2017). Alternatively, expression in hybrids may

tend towards underexpression if regulatory divergence between parental lineages results in

impaired transcription factor binding with promoter or enhancer elements (Oka *et al.* 2014;

Guerrero *et al.* 2016) or if divergence stimulates epigenetic silencing (Paun *et al.* 2007;

Shivaprasad *et al.* 2012; Lafon-Placette and Köhler 2015; Brekke *et al.* 2016; Zhu *et al.* 2017).

At a finer scale, misexpression patterns may depend on developmental stage: for example, if

there is greater pleiotropy earlier in development (Ortíz-Barrientos *et al.* 2007; Cutter and

Bundus 2020). In particular, we might expect sterile hybrids to have more misexpression during

later stages of gametogenesis when genes are evolving rapidly and are potentially under less

regulatory constraint (Kopania *et al.* 2022a; Murat *et al.* 2023). Finally, the role of sex

chromosome regulation in inviable or sterile hybrids encompasses both of these larger

questions. Sex chromosomes may be prone to asymmetry in their expression divergence (Oka

and Shiroishi 2014; Civetta 2016) and be regulated differently across stages of development

(Presgraves 2008; Larson *et al.* 2018), particularly in reproductive tissues, and thus may play a

central role in hybrid dysregulation relative to autosomes.

Disruption of sex chromosome regulation is thought to be a potentially widespread

regulatory phenotype in sterile hybrids (Lifschytz and Lindsley 1972; Larson *et al.* 2018), in part

because X chromosome repression may be crucial to normal spermatogenesis in diverse taxa

(McKee and Handel 1993; Landeen *et al.* 2016; Taxiarchi *et al.* 2019; Rappaport *et al.* 2021;

Viera *et al.* 2021; Murat *et al.* 2023). Furthermore, misregulation of the X chromosome is

associated with hybrid sterility in several species pairs (Davis *et al.* 2015; Sánchez-Ramírez *et

al.* 2021; Bredemeyer *et al.* 2021), although it has been best studied in house mice. In fertile

male mice, the X chromosome is silenced early in meiosis through meiotic sex chromosome

inactivation (MSCI; McKee and Handel 1993; Handel 2004) and is again repressed in

postmeiotic sperm development (*i.e.,* postmeiotic sex chromosome repression or PSCR;

Namekawa *et al.* 2006). In contrast, the X chromosome is not properly inactivated and is

overexpressed in sterile hybrid mice (Good *et al.* 2010; Bhattacharyya *et al.* 2013; Campbell *et

al.* 2013; Turner *et al.* 2014; Larson *et al.* 2017, 2022). Disrupted MSCI in house mice is

associated with divergence at *Prdm9*, a gene that is a major contributor to hybrid male sterility

(Mihola *et al.* 2009; Davies *et al.* 2016). However, misexpression of the X chromosome in sterile

hybrids could result from mechanisms other than *Prdm9*-associated disrupted MSCI, such as

mispairing of the sex chromosomes due to divergence in their region of homology known as

the pseudoautosomal region (PAR; Burgoyne 1982; Ellis and Goodfellow 1989; Raudsepp and

Chowdhary 2015). In sum, the ubiquity of X chromosome repression and the growing body of

evidence linking disrupted MSCI to hybrid sterility in mammals suggest that disrupted sex

chromosome regulation may be a common regulatory phenotype in sterile hybrid males.

Here, we characterized disruption of gene expression associated with hybrid male

sterility in two rodent crosses, dwarf hamsters and house mice, which span ~35 million years

of divergence (Swanson *et al.* 2019). The regulatory phenotypes of hybrid male sterility have

been thoroughly studied in house mice (Good *et al.* 2010; Bhattacharyya *et al.* 2013; Campbell

et al. 2013; Turner *et al.* 2014; Larson *et al.* 2017, 2022; Hunnicutt *et al.* 2022). We contrast

these with an analogous cross between two sister species of dwarf hamster, Campbell's dwarf

hamster (*Phodopus campbelli*) and the Siberian dwarf hamster (*P. sungorus*), and their sterile

F1 hybrid male offspring. These species diverged only ~0.8-1.0 million years ago (Neumann *et*

al. 2006), and they are not thought to interbreed in the wild due to geographic separation

(Ishishita *et al.* 2015). Crosses between female *P. sungorus* and male *P. campbelli* produce

sterile hybrid males that, similar to mice, have a range of sterility phenotypes, suggesting

multiple developmental blocks to spermatogenesis (Ishishita *et al.* 2015; Bikchurina *et al.*

2018). Hybrids from the reciprocal cross are usually inviable due to abnormal growth *in utero*

(Brekke and Good 2014), and the species origin of the X chromosome is the primary genetic

factor controlling hybrid inviability (Brekke *et al.* 2021). Additionally, sex chromosome

asynapsis during spermatogenesis is common in hybrid dwarf hamsters, providing further

reason to think that X chromosome-specific misregulation may be observed in sterile hybrid

dwarf hamsters (Ishishita *et al.* 2015; Bikchurina *et al.* 2018). Both the abnormal spermatogenic

phenotypes observed in dwarf hamster hybrids and the potential regulatory interactions that

may result from the involvement of the X chromosome in multiple reproductive barriers make

dwarf hamsters an important comparison to mice for investigating what regulatory phenotypes

may be repeatedly associated with the evolution of postzygotic reproductive isolation.

Expression phenotypes associated with hybrid sterility have historically been difficult to

assess because of the cellular diversity of reproductive tissues (*e.g.,* testes; Good *et al.* 2010;

Ramm and Schärer 2014; Montgomery and Mank 2016; Hunnicutt *et al.* 2022), thus requiring

molecular approaches that permit greater spatial and developmental resolution into expression

evolution. To overcome these difficulties, we used Fluorescence Activated Cell Sorting (FACS)

to isolate and sequence cell populations across the developmental timeline of

spermatogenesis for each species pair and their F1 hybrids, including stages that span the

different sex chromosome regulatory states. We used both datasets to address three main

questions about the gene expression phenotypes observed in sterile hybrids: (1) within

transgressive differentially expressed genes, does misexpression tend towards up- or

downregulation in hybrids compared to parents? (2) are there similar patterns of disrupted

expression across stages of development? and (3) are there clear differences between

autosomes and sex chromosomes in regulatory phenotypes? And if so, is sex chromosome-

specific misexpression consistent with either disrupted MSCI and/or disrupted PAR regulation?

Collectively, we demonstrate the power of cell type-specific approaches for untangling the

expression phenotypes associated with the evolution of hybrid male sterility and for identifying

common themes in the mechanistic basis of hybrid incompatibilities across divergent taxa.

MATERIALS AND METHODS

Hamster crosses and male reproductive phenotypes

We used wild-derived colonies of two sister species of dwarf hamster, *P. sungorus* and

P. campbelli, established by Kathy Wynne-Edwards (Scribner and Wynne-Edwards 1994) and

housed at the University of Montana. Both species were maintained as closed colonies with a

breeding scheme to minimize inbreeding. Nonetheless, inbreeding levels of these closed

colonies are still high as indicated by very low nucleotide diversity (Brekke *et al.* 2018). We

used males from both parent species and male F1 hybrid offspring from crosses of female *P.*

campbelli with male *P. sungorus*. We weaned males in same-sex sibling groups between 17 -

21 dpp and housed them individually at 45 dpp. We euthanized reproductively mature males

using carbon dioxide followed by cervical dislocation between 59 - 200 dpp (Table S1). All

animal use was approved by the University of Montana (IACUC protocols 050-16JGDBS &

035-19JGDBS).

We measured several fertility metrics for parent species and hybrid males including

paired testes weight, paired seminal vesicle weight, normalized sperm counts, and sperm

motility (Good *et al.* 2008). Paired testes weight and paired seminal vesicle weight were

correlated with body weight (paired testes weight Pearson's $r(29) = 0.47$, p = 0.007; paired

seminal vesicle weight Pearson's $r(23) = 0.56$, p = 0.003), so we standardized both metrics

relative to body weight. We calculated sperm count by isolating sperm from caudal

epididymides diced in 1 ml of Dulbecco's PBS (Sigma) and incubated at 37°C for 10 minutes.

We quantified sperm motility (proportion of motile sperm in a 5 µl suspension) and sperm count

(number of sperm with head and tail in a heat shocked 5 µl suspension) across a fixed area on

a Makler counting chamber. We performed statistical comparisons of fertility phenotypes in R

v.4.3.1, and we used the FSA package v.0.9.4 for the Kruskal-Wallis and Dunn's tests (Ogle

and Ogle 2017).

Isolation of enriched cell populations from hamster testes

To investigate the regulatory dynamics of the sex chromosomes during

spermatogenesis, we isolated four spermatogenic cell populations from whole testes using

FACS: spermatogonia, leptotene/zygotene spermatocytes, diplotene spermatocytes, and

round spermatids. Briefly, we disassociated a single testis per male following a published

protocol originally developed for house mice (Getun *et al.* 2011) with modifications

(github.com/goodest-goodlab/good-protocols/tree/main/protocols/FACS, last accessed June

16, 2021). We doubled the volumes of all reagents to account for the increased mass of testes

in dwarf hamsters relative to house mice. We isolated cell populations based on size,

granularity, and fluorescence on a FACSAria IIu cell sorter (BD Biosciences) at the University of

Montana Center for Environmental Health Sciences Fluorescence Cytometry Core. For each

sorted cell population, we extracted RNA using RNeasy kits (Qiagen) following protocols for

Purification of Total RNA from Animal Cells. We quantified sample RNA quantity and quality

(requiring an RNA integrity number > 7) on a Tapestation 2200 (Agilent) at the University of

Montana genomics core. RNA libraries were prepared by Novogene and sequenced on Illumina

NovaSeq 6000s (paired end, 150 bp). Six samples (distributed across different species and cell

types) had low RNA concentrations, and for these samples we used Novogene's low input

RNA library preparation (Table S2). MDS plots indicated no severe library batch effects

between samples from different library preparations (Figure S1), so we included all samples

from both libraries in subsequent analyses.

Read processing and mapping

We sequenced RNA from each cell population for three to five individuals of each

parent species and F1 hybrids generating an average of ~27.5 million read pairs per individual

(Table S2). We trimmed reads using Trimmomatic v.0.39 (Bolger *et al.* 2014) to remove low

quality bases from the first and last 5 bp of each read and bases with an average Phred score

of less than 15 across a 4 bp sliding window and only retained reads of at least 36 bp. We next

used an approach (based on the modtools pipeline) which maps reads from each sample to

pseudogenomes for both parent species (described below) to obtain a merged output

alignment file in order to alleviate reference bias associated with mapping hybrids to only a

single reference genome (Holt *et al.* 2013; Huang *et al.* 2014). For this approach, we mapped

reads for each individual to both a *P. sungorus* pseudogenome and a *P. campbelli*

pseudogenome with Hisat v.2.2.0 (Kim *et al.* 2019) with default settings and retaining at most

100 distinct, primary alignments, although multi-mapped reads were removed downstream

(described below). We generated the *P. sungorus* pseudogenome by mapping RNASeq reads

from a male *P. sungorus* individual (30.6 million total read pairs; NCBI SRA: SRR17223284;

Moore *et al.* 2022) to the *P. sungorus* reference genome (GCA_023856395.1) with bwa-mem

v.2.2.1 (Vasimuddin *et al.* 2019), and the *P. campbelli* pseudogenome by mapping female *P.*

campbelli whole genome sequencing reads (average coverage: 33x; NCBI SRA: SRR17223279;

Moore *et al.* 2022) to the *P. sungorus* reference genome. Because our *P. sungorus*

pseudogenome was based on a male hamster and the reference genome on a female hamster,

we excluded reads mapping to the PAR because sequences mapping to this region could have

originated from either the X or Y chromosomes and interfered with subsequent variant calling.

Following mapping, we used GATK v.4.2.5.0 HaplotypeCaller (-ERC GVCF) to call SNPs then

performed genotyping with genotypeGVCFs. We hard-filtered our SNPs (--mask-extension 5

"QD < 2.0" "FS > 60.0" "MQ < 40.0" "QUAL < 30.0" "DP < 10" "DP > 150") and restricted

SNPs to biallelic loci. Finally, we incorporated filtered SNPs back into the *P. sungorus*

reference genome with FastaAlternateReferenceMaker to create the *P. sungorus* and *P.*

campbelli pseudoreferences. For our RNASeq data, we appended query hit indexes to

resulting alignment files using hisat2Tophat.py (https://github.com/goodest-goodlab/pseudo-

it/tree/master/helper-scripts/hisat2Tophat.py, last accessed March 8th, 2022) to maintain

compatibility with the modtools pipeline. We used our VCFs (above) to generate a mod-file for

both species with vcf2mod from Lapels v.1.1.1 to convert alignments to the *P. sungorus*

reference genome, and Suspenders 0.2.6 to merge alignments while retaining the highest

quality alignment per read (Holt *et al.* 2013; Huang *et al.* 2014). We used featureCounts v.2.0.1

(Liao *et al.* 2014) to estimate counts of read pairs that aligned to the same chromosome (-B

and -C) and retained only singly-mapped reads. Summaries of properly mapped reads for each

sample can be found in Table S2.

We sought to compare the gene expression phenotypes observed in dwarf hamsters to

those previously documented in house mice using published RNASeq data for the same four

spermatogenic cell types of two subspecies of house mouse and their sterile F1 hybrids

(Larson *et al.* 2017; Hunnicutt *et al.* 2022). These studies examined two subspecies of house

mice, *Mus musculus musculus* (intra-subspecific F1 males between wild-derived inbred strains

PWK/PhJ♀ and CZECHII/EiJ♂) and *M. m. domesticus* (intra-subspecific F1 males between

wild-derived inbred strains WSB/EiJ♀ and LEWES/EiJ♂) and their sterile (PWK♀ x LEWES♂)

F1 hybrids for disrupted gene expression across spermatogenesis following the same FACS

protocols implemented in this study. For all comparisons between house mice and dwarf

hamsters, we used read count files generated previously for house mice (Hunnicutt *et al.* 2022)

and performed all subsequent analyses in parallel for both systems.

Estimating nucleotide diversity and divergence within and between crosses and strains
217
We next estimated nucleotide diversity within and divergence between each strain or

cross for parent species and hybrids (π and d_{XY}). We used the bam files generated above from

mapping each strain or cross to its respective reference genome (either GCA_023856395.1 for

dwarf hamsters or GRCm38.p6 for house mice) to call SNPs as described above but with the

addition of a step to split reads at intronic regions using the GATK function SplitNCigarReads.

We generated two VCFs, one for dwarf hamster crosses and the other for house mouse

crosses, which we processed and filtered separately. If an individual mouse or hamster was

sequenced for more than one cell type, we randomly chose one of the represented cell types

to be included in the analysis. We hard-filtered our SNPs as above and restricted SNPs to

biallelic loci. Because our SNPs were called from RNASeq data and thus may be susceptible to

allelic imbalances or coverage differences between samples, we next investigated how the

inclusion of SNPs with different levels of missing data impacted estimation of π and d_{XY}. We

used vcftools v.0.1.17 (Danecek *et al.* 2011) to filter SNPs allowing between 0% and 90%

missing data (--max_missing; Supplemental Figure S3), and filtered SNPs with a depth lower

than 5 and higher than 60 (~2-3 higher than average coverage) to eliminate multi-mapped

reads. Finally, we used pixy v.1.2.10.beta2 (Korunes and Samuk 2021) on our filtered VCFs to

estimate π and d_{XY}. Patterns of nucleotide diversity across strains and crosses were

qualitatively similar across missing data thresholds, so we present results corresponding to

10% missing data in the main text. However, we note that for all comparisons, estimates of π

and d_{XY} decreased with more stringent missing data thresholds regardless of strain or cross

(Supplemental Figure S3).

Gene expression pre-processing

Following read processing and mapping, we conducted all analyses in R v.4.3.1. We

classified genes as "expressed" if genes had a minimum of one Fragment Per Kilobase of exon

per Million mapped reads (FPKM) in at least three samples, resulting in 21,077 expressed

genes across the dwarf hamster dataset and 21,212 expressed genes across the house mouse

dataset. We also identified sets of genes "induced" in a given cell-type defined as genes with a

median expression in a given cell population (normalized FPKM) greater than two times its

median expression across all other sorted cell populations (following Kousathanas *et al.* 2014).

We calculated normalized FPKM values by adjusting the sum of squares to equal one using the

R package vegan v.2.6-4 (Oksanen *et al.* 2013). We conducted expression analyses using

edgeR v.3.42.4 (Robinson *et al.* 2010) and normalized the data using the scaling factor method

(Anders and Huber 2010).

We qualitatively assessed cell population purity both by visual inspection during cell

sorting and following sequencing by assessing the expression of a panel of marker genes

specific to the four cell populations targeted by our FACS protocol and present in only a single

copy in the *P. sungorus* annotation. Spermatogonia markers included *Dmrt1* (Raymond *et al.*

2000) and *Hells* (Green *et al.* 2018). Leptotene/zygotene markers included *Ccnb1ip1* and

Adad2 (Hermann *et al.* 2018). Diplotene was characterized by *Aurka* and *Tank* expression

(Murat *et al.* 2023) and round spermatids by *Cabyr* and *Acrv1* expression (Green *et al.* 2018).

To estimate relative purity, we compared mean marker gene expression across replicates for a

given cell population for both parent species. A cell population was considered qualitatively

pure if it had higher marker gene expression than other populations isolated by our FACS

protocol and if X-linked gene expression matched the expected regulatory dynamics (*i.e.,*

active vs. silenced (MSCI) vs. repressed (PSCR); Handel 2004; Namekawa *et al.* 2006). We

examined expression patterns across cell populations for all genes, autosomal genes, and X-

linked genes using MDS plots generated with the plotMDS function in limma v.3.56.2 (Ritchie

et al. 2015) and heatmaps using ComplexHeatmap v.2.16.0 (Gu *et al.* 2016). MDS plots used

the top 500 genes with the largest fold change difference between samples.

Differential gene expression analysis

We assessed differential gene expression by contrasting hybrids and each parent

species for all cell populations. We fit the expression data for dwarf hamsters and house mice

separately with negative binomial generalized linear models with Cox-Reid tagwise dispersion

estimates and adjusted P-values to a false discovery rate (FDR) of 5% (Benjamini and

Hochberg 1995). We quantified the biological coefficient of variation (BCV), a metric

representing the variation in gene expression among replicates (McCarthy *et al.* 2012), for each

dataset. Additionally, we calculated the BCV of just parental males or hybrid samples for each

species for the first three cell populations to examine whether dwarf hamster hybrids exhibited

more variability in expression than house mouse hybrids and parental dwarf hamsters. For our

differential expression analyses, we contrasted expression between hybrids and each parent

so that a positive log fold-change (logFC) indicated overexpression in sterile males and

implemented a logFC cutoff of 1.25. We then categorized differentially expressed (DE) genes

into one of four categories: DE relative to only one parent species, DE relative to both parent

species but with intermediate expression (intermediate), and DE relative to both parent species

but outside of the range of either parent species (transgressive; Figure S2). Unless otherwise

specified, results discussed in the main text are restricted to transgressive DE genes and are

presented with the logFC from the contrast of the hybrid offspring to the parent with the same

X chromosome (*P. campbelli* for dwarf hamster F1 hybrids and *M. m. musculus* for house

mouse F1 hybrids). Transgressive DE genes have similar logFC values regardless of which

parent is used as the contrast, but figures depicting the logFC between F1 hybrids and *P.*

sungorus or *M. m. domesticus* are provided in supplementary material.

We tested for significant differences in the number of under and overexpressed

transgressive DE genes within a stage for both house mouse and hamster hybrids using X^2

tests with chisq.tests in R and used FDR correction for multiple comparisons. We also tested

for differences in the magnitude of misexpression between mouse and hamster hybrids for

each stage by comparing the distributions of logFC of transgressive DE genes between hybrids

and parent species using Wilcoxon signed-rank tests and FDR correction. To characterize

hybrid diplotene expression in both house mice and dwarf hamsters, we used two approaches.

First, we calculated Pearson's correlation coefficient *(r)* between average normalized hybrid

diplotene expression and the average normalized expression in each parental cell type. We

corrected p-values for each correlation with FDR. We generated bootstrap values for each

correlation coefficient by randomly sampling the expression matrices with replacement for

each sample type for 1000 replicates. Second, we compared the gene sets that escaped MSCI

in each species with gene sets that characterize stage-specific expression in parent species.

For this analysis, we defined sets of overexpressed X-linked diplotene genes as genes with

expression (normalized FPKM) in hybrids that was in the top 10% of X-linked genes in parental

diplotene samples (*i.e.,* genes that normally escape MSCI). We then compared these sets of

overexpressed hybrid diplotene genes to genes "induced" in each parental stage for each

species. We used gProfiler2 v.0.2.3 (Kolberg *et al.* 2020) in R to perform gene ontology (GO)

analysis to identify GO terms overrepresented in hybrid DE genes sets for each cell population.

We only included *P. sungorus* genes associated with mouse orthologs in our GO analysis (as

established by Moore *et al.* 2022), and for our background gene lists, we used *P. sungorus*

genes associated with mouse orthologs that were "expressed" in hybrids and both parent

species in a given stage. We retained only Biological Process GO terms with an FDR below

0.05 and ran gProfiler2 both with and without the highlight option, a two-stage algorithm for

reducing resulting GO terms by grouping significant terms into sub-ontologies and then

identifying the gene sets that give rise to other significant functions (Kolberg *et al.* 2020). To

test whether specific chromosomes were enriched or depleted for transgressive DE genes for a

given stage, we performed hypergeometric tests on the number of DE genes on a given

chromosome with phyper and adjusted P-values to an FDR of 5%. We also assessed overlap

in specific DE genes within stages between house mice and dwarf hamsters. We estimated

whether overlap was more or less than expected by chance using hypergeometric tests and

performed GO enrichment on overlapping genes.

Characterizing the behavior of PAR genes in dwarf hamsters

325  We sought to characterize the regulatory behavior of PAR genes in dwarf hamster

326 parent species and hybrids to determine if (1) PAR genes are normally silenced in parent

327 species (consistent with an extension of MSCI to the PAR) and (2) if PAR genes were

328 overexpressed in hybrids (consistent with disrupted MSCI in the PAR of hybrids). The PAR on

329 the *P. sungorus* X chromosome is on the distal arm of the X chromosome from around

330 115,350,000-119,112,095 bp (Moore *et al.* 2022). There are 15 annotated *P. sungorus* genes in

331 this region (Table S3), which is comparable to the latest PAR assembly in C57BL/6J house

332 mice (Kasahara *et al.* 2022). Six of these are orthologous to annotated genes in mice (*Tppp2*,

333 *Gprin1*, *Ndrg2*, *Kcnip4*, *Ndrg2*, and *Hs6st3*), but they are not located in the mouse PAR

334 (Kasahara *et al.* 2022). Only seven of the annotated genes in the PAR were expressed in more

335 than three replicates across all samples (Psun_G000022875, Psun_G000022880,

336 Psun_G000022883, Psun_G000022886, *Tppp2*, *Gprin1*, and *Ndrg2*). For these genes, we

337 assessed whether these genes were consistently expressed or silenced in parent species at

338 any cell stage and whether any genes were differentially expressed between hybrids and either

339 species.

340 **RESULTS**

341 **Impaired sperm production in hybrid male dwarf hamsters**

342  We first established the extent of hybrid male sterility in dwarf hamsters by comparing

343 reproductive phenotypes for *P. campbelli*, *P. sungorus*, and F1 hybrid males (Figure 1; Table

344 S1). *Phodopus sungorus* males had smaller testes and seminal vesicles than *P. campbelli*

345 males (Dunn's Test relative testes weight $p < 0.001$; relative seminal vesicle weight $p = 0.0061$;

346 Figure 1). These differences are qualitatively consistent across independent laboratory colonies

(Ishishita *et al.* 2015; Bikchurina *et al.* 2018) and likely reflect species-specific differences

between *P. campbelli* and *P. sungorus*. *Phodopus sungorus* males also had lower nucleotide

diversity (π = 0.00013) than *P. campbelli* males (π = 0.00045) and both house mouse parental

crosses between fully inbred mouse strains (*M. m. musculus* π = 0.00029; *M. m. domesticus* π

0.00017; Figure S3), which could contribute to depression of male fertility within this highly

inbred laboratory colony (Brekke *et al.* 2018). Nucleotide divergence between parental dwarf

hamster species was elevated relative to house mice (d_{xy} = 0.0020 vs. 0.0010), consistent with

reported older divergence time estimates for dwarf hamsters (Neumann *et al.* 2006). However,

these dwarf hamster species did not differ in normalized sperm counts (p = 0.071) or in sperm

motility (p = 0.12). The F1 hybrid males exhibited extreme reproductive defects relative to *P.*

campbelli (Figure 1). Hybrid males had smaller testes (p < 0.001) and seminal vesicles (p <

0.001) than male *P. campbelli* hamsters, and importantly, produced almost no mature

spermatozoa. In the one instance where a hybrid male produced a single mature

spermatozoon, it was non-motile, indicating severe reproductive impairment in hybrid males.

Overall, our results confirmed previous reports of reduced fertility in hybrid male dwarf

hamsters (Ishishita *et al.* 2015; Bikchurina *et al.* 2018).

Figure 1. Evidence of some hybrid male sterility in dwarf hamsters. We assessed paired

testes weight and paired seminal vesicle weight (SV; both normalized by body weight), sperm

count, and proportion of motile sperm for *P. sungorus*, *P. campbelli*, and F1 hybrids. Whiskers

extend to either the largest or smallest value or no further than 1.5 times the interquartile range,

and *** indicates p < 0.001, ** indicates p < 0.01, and n.s. indicates non-significant difference

between means at p > 0.05 using a post-hoc Dunn's test with FDR correction. Upwards-

pointing triangles (▲) indicate *P. campbelli*, downwards-pointing triangles (▼) indicate *P.*

sungorus, and crosses (✖) indicate F1 hybrids.

Cell-specific gene expression across spermatogenesis

To characterize cell-specific gene expression, we used FACS to isolate enriched cell

populations from each fertile parent species and their sterile F1 hybrids across four stages of

spermatogenesis. The four targeted populations included: spermatogonia (mitotic precursor

cells), leptotene/zygotene spermatocytes (meiotic cells before MSCI), diplotene spermatocytes

(meiotic cells after MSCI), and round spermatids (postmeiotic cells). We were unable to isolate

round spermatids from the F1 hybrids, which was consistent with the lack of mature

spermatozoa present in the cauda epididymis extractions (Figure 1). We sequenced RNA from

each cell population for *P. campbelli* (spermatogonia n = 4, leptotene/zygotene n = 5, diplotene

n = 5, and round spermatids n = 4; Table S2), *P. sungorus* (spermatogonia n = 4,

leptotene/zygotene n = 4, diplotene n = 5, and round spermatids n = 3), and F1 hybrid males (

P. campbelli ♀ x *P. sungorus* ♂; spermatogonia n = 4, leptotene/zygotene n = 4, diplotene n =

4). We compared our hamster expression data to an analogous cell-specific RNASeq dataset

from two species of house mice, *Mus musculus musculus* and *M. m. domesticus*, and their

sterile F1 hybrids (n = 3 for all cell populations in each cross; Larson *et al.* 2017).

We used two approaches to qualitatively evaluate the purity of spermatogonia,

leptotene/zygotene spermatocytes, diplotene spermatocytes, and round spermatids isolated

from males from both parental species: (1) we quantified the relative expression of a panel of

cell population marker genes, and (2) we characterized the expression patterns of the sex

chromosomes across development. We found that our candidate marker genes had the

highest expression in their expected cell population for all stages except leptotene/zygotene

(Figures S4 and S5), indicating high purity of spermatogonia, diplotene spermatocytes, and

round spermatids. Leptotene/zygotene markers did not have the highest expression in

leptotene/zygotene samples (except for *Ccnb1ip1*), potentially indicating lower purity of this

cell population. Nonetheless, the patterns of X chromosome expression in fertile parents were

consistent with expectations across this developmental timeline: the X chromosome had active

expression in spermatogonia and leptotene/zygotene cells, was inactivated in diplotene cells

consistent with MSCI, and was partially inactivated in round spermatids, consistent with PSCR

(Figure 2; Namekawa *et al.* 2006), indicating successful isolation of these cell populations.

When we examined overall expression differences within dwarf hamsters and within

house mice, we found that samples clustered primarily by cell population on MDS1 and 2

(Figure 3), then by cross/strain when cell populations were examined separately (Figures S6

and S7). Within all cell populations across both systems, hybrids showed intermediate overall

expression patterns to parent species (Figures S6 and S7). However, in dwarf hamsters but not

house mice, spermatogonia and leptotene/zygotene samples overlap rather than forming

distinct clusters (Figure 3). Further, the expression profiles of hybrid diplotene cells ranged

from clustering with parental leptotene/zygotene to parental diplotene cells, which contrasted

with what we observed in house mice where hybrid diplotene cells clustered distinctly with

parental diplotene cells (Figure 3). Overall, our results indicate that we successfully isolated cell

populations in dwarf hamsters that span key stages of spermatogenesis.

Figure 2. Overexpression of X-linked genes in diplotene spermatocytes in both house mouse and dwarf hamster hybrids. Heatmap of X-linked gene expression in house mice (upper panel) and dwarf hamsters (lower panel) plotted as normalized FPKM values that are hierarchically clustered using Euclidean distance. Each column represents a different individual, each row represents a gene, and darker colors indicate higher expression. The heatmap was generated with the R package ComplexHeatmap v.2.12.0 (Gu et al. 2016). Note, hybrid dwarf hamsters do not produce mature spermatozoa, and accordingly, we were unable to isolate round spermatids.

Figure 3. Hybrid gene expression profiles cluster by parental spermatogenic cell

population in house mice but not dwarf hamsters. Multidimensional scaling (MDS) plots of

distances among house mouse (upper panels) and dwarf hamster (lower panels) samples for

expressed autosomal (left) and X-linked (right) genes. Distances are calculated as the root-

mean-square deviation (Euclidean distance) of log2 fold changes among the top 500 genes

that distinguish each sample. Each strain or cross is indicated by a symbol, and samples are

colored by cell population.

Disrupted transcription early in spermatogenesis in dwarf hamsters

We sought to characterize which expression phenotypes were associated with sterile

hybrids in both house mice and dwarf hamsters. We first investigated whether differential gene

expression in hybrids tended towards intermediate or transgressive expression, and then

within transgressive DE genes, whether misexpression tends towards up- or downregulation in

hybrids compared to parents. Differential gene expression in mouse hybrids had a slight bias

towards transgressive expression except in round spermatids (percentage of transgressive DE

genes: SP: 72.0%, LZ: 62.7%, DIP: 60.8%, and RS: 24.9%; Figures 4B and S2), while almost

all differential expression in dwarf hamster hybrids was transgressive (percentage of

transgressive DE genes: SP: 99.2%, LZ: 98.0%, and DIP: 97.1). For transgressive DE genes,

autosomal misexpression in hybrid house mice was biased towards upregulation across

spermatogenesis (mean logFC of spermatogonia autosomal DE genes: +1.90/ X^2: p < 0.001;

leptotene/zygotene: +1.33/ X^2: p < 0.001; diplotene: +0.47/ X^2: p < 0.001; Figures 4C and S8),

as was X-linked misexpression (spermatogonia mean logFC: +1.98 / X^2: p < 0.001;

leptotene/zygotene: +2.50/ X^2: p < 0.001; diplotene: +2.78/ X^2: p < 0.001). In contrast, we found

that the direction of misexpression in dwarf hamster hybrids was stage-specific. Autosomal DE

genes in dwarf hamster hybrids were overwhelmingly downregulated in both early stages of

spermatogenesis, especially in comparison to house mice (mean logFC of spermatogonia

autosomal DE genes: -4.38/ X^2: p < 0.001; leptotene/zygotene: -4.67/ X^2: p < 0.001; Figure 4D)

but upregulated in diplotene (average logFC = +4.67/ X^2: p < 0.001; Figures 4D and S8). When

comparing both X-linked and autosomal expression in dwarf hamsters, we found similar

patterns: almost all X-linked DE genes in the first two stages of spermatogenesis were

exclusively downregulated (spermatogonia mean logFC = -6.57/ X^2: p < 0.001;

leptotene/zygotene: -6.73/ X^2: p < 0.001), but misexpression was biased towards upregulation

in diplotene (mean logFC: 5.70/ X^2: p < 0.001).

Figure 4. House mice and dwarf hamster hybrids have opposite patterns of disrupted regulation early in spermatogenesis. A. Up- and down-regulated DE genes between parent species for house mice (left; *M. m. musculus* vs. *M. m. domesticus*) and dwarf hamsters (right; *P. sungorus* vs. *P. campbelli*). **B**. Counts of DE genes between hybrids and one parent species (two lightest shades of gray where Parent 1 was either *M. m. musculus or P. sungorus* and parent 2 was either *M. m. domesticus* or *P. campbelli)* or between hybrids and both parents (two darker shades of gray) that showed either intermediate or transgressive expression. **C.** Transgressive DE genes in house mouse and dwarf hamster hybrids for autosomal genes where the logFC represents hybrid expression relative to *M. m. musculus* or *P. campbelli,* respectively, and **D.** transgressive DE gene expression for X-linked genes*.* Results are displayed for autosomes (left) and the X chromosome (right). *** indicates p < 0.001 for

pairwise comparisons from Wilcoxon signed-rank tests after FDR correction. Whiskers extend

to either the largest or smallest value or no further than 1.5 times the interquartile range. The

number of up and downregulated DE genes in hybrids are listed next to arrows indicating

direction of differential expression.

Second, we investigated how developmental stage influenced the extent of hybrid

misexpression. In both systems, the number of DE genes between parent species increased

with the progression of spermatogenesis, consistent with less constraint on gene expression

levels as spermatogenesis progresses (Figure 4A; Kopania *et al.* 2022a; Murat *et al.* 2023).

Similarly, transgressive misexpression in both hybrid dwarf hamsters and house mice

increased with the progression of spermatogenesis, though to a greater extent in dwarf

hamsters (Figure 4B). While both parental DE genes and transgressive hybrid DE genes

increased with the progression of spermatogenesis, parental DE genes initially exceeded the

number of transgressive hybrid DE genes in hybrids in spermatogonia and leptotene/zygotene

spermatocytes. However, in diplotene spermatocytes, transgressive hybrid differential

expression surpassed parental differential expression in both house mice and dwarf hamsters.

We also found a much greater genome-wide disruption of expression in diplotene cells of

hybrid dwarf hamsters than in hybrid house mice (Figures 4C-D and S8-S10), indicating more

widespread regulatory disruption. We then compared transgressive DE genes between house

mice and dwarf hamster hybrids across all three stages and found that the number of shared

DE genes did not differ from the number expected by chance (spermatogonia = 3 genes,

hypergeometric test $p = 0.98$; leptotene/zygotene = 2 genes, $p = 0.59$; and diplotene = 78

genes, $p = 0.18$; Table S4). These sets of shared DE genes between house mice and dwarf

hamsters were not significantly enriched for any GO terms, although many play known roles in

spermatogenesis, the apoptotic process, and cell differentiation (Table S5) and may be

promising candidates for future functional analysis. Ultimately, we found that, although few

specific genes have similarly disrupted expression in dwarf hamster and house mouse hybrids,

there are similar trends in the patterns of transgressive expression across stages in both

systems.

To determine if the differences we observed in the extent of misexpression between

dwarf hamsters and house mice was due to greater expression variability across our dwarf

hamster samples, we calculated the BCV, a measurement of inter-replicate variability, for each

species and hybrid across the first three cell populations. Inter-replicate variability was higher

in dwarf hamster hybrids relative to house mouse hybrids (dwarf hamster BCV = 0.69; house

mice = 0.18). Additionally, nucleotide diversity within dwarf hamster hybrids was higher (π =

0.0012) than within house mouse hybrids (π = 0.00052; Figure S3). However, the extent of the

expression variability observed in hybrids relative to the inter-replicate variability of parental

species differed between house mice and dwarf hamsters: dwarf hamster hybrid variability was

more than dwarf hamster parental samples (*P. campbelli* = 0.49; *P. sungorus* = 0.40), but

hybrid variability was similar to parental samples for house mice (*M. m. musculus* = 0.19; *M. m.*

domesticus = 0.22). The greater inter-replicate expression variability in dwarf hamster hybrids

relative to parent species suggests that the increased misexpression we see in hybrids cannot

be explained by greater inter-replicate variability in our dwarf hamster samples alone, and it

may also reflect the greater nucleotide diversity present among dwarf hamster hybrids.

Third, we characterized whether there was a clear difference between the autosomes

and sex chromosomes in expression phenotype by testing whether the X chromosome was

enriched for transgressive DE genes in each stage for both systems. Across all stages of

spermatogenesis in house mice, the X chromosome was enriched for transgressive DE genes

between hybrids and parents (spermatogonia $p < 0.001$; leptotene/zygotene $p < 0.001$;

diplotene $p < 0.001$; round spermatids $p < 0.0036$; Figure S9). In contrast, misexpression in

dwarf hamsters was not uniformly sex chromosome-specific across all stages, as the sex

chromosomes in dwarf hamsters showed no enrichment for transgressive DE genes early in

spermatogenesis (spermatogonia: p = 0.21; Figure S10) despite X chromosome enrichment in

both leptotene/zygotene (p = 0.0064) and round spermatids (p < 0.001). However, we note that

the magnitude of misexpression was greater for sex chromosomes than autosomes in dwarf

hamsters across all stages (Figures 4C and 4D; discussed above). Only two autosomes were

enriched for transgressive DE genes in dwarf hamsters in any cell population: one scaffold on

chromosome 5 in spermatogonia (JAJQIY010003390.1; hypergeometric test; p = 0.003) and

chromosome 11 in leptotene/zygotene (p = 0.017; Figure S10). Together, the subtle differences

in the distribution of transgressive DE genes across autosomes and the X chromosome

between dwarf hamster hybrids and house mouse hybrids suggest a difference in the extent of

the role for sex chromosome-specific disruption between systems.

Misexpression in diplotene appears to be unrelated to disrupted MSCI in dwarf hamsters

We next asked whether expression patterns indicated similar disrupted regulatory

processes resulting in sex chromosome-specific misexpression in both systems. In sterile

hybrid house mice, the mean logFC of transgressive X-linked DE genes during diplotene was

higher than the mean logFC of transgressive autosomal DE genes (+0.47 in contrast to +2.78;

Figure 4C and D), consistent with disrupted MSCI (Good *et al.* 2010; Bhattacharyya *et al.* 2013;

Campbell *et al.* 2013; Turner and Harr 2014; Larson *et al.* 2017, 2022). In sterile dwarf hamster

hybrids, we also found elevated mean logFC of transgressive DE X-linked genes relative to

autosomal genes (+3.69 in contrast to +5.70), but the extent of X chromosome overexpression

was greater than in hybrid house mice (Figures 2, 4C, and 4D). There was also more variability

in the relative extent of overexpression of X-linked genes in hybrid dwarf hamsters during

diplotene compared to the mean expression of X-linked genes of parental dwarf hamsters

(relative overexpression = 18.5 +/- 6.8) compared to hybrid house mice (relative overexpression

= 1.75 +/- 0.09; Figures 2 and S11). Strikingly, some hybrid male dwarf hamsters had an

almost completely silenced X chromosome, while others had an almost completely

transcriptionally-activated X chromosome (Figure 2).

Despite overexpression of the X chromosome during diplotene in hybrid dwarf

hamsters, the overall expression phenotype, including the identity and the extent of

misexpression of overexpressed genes, appeared to fundamentally differ between house

mouse and dwarf hamster hybrids (Figures 2 and 5A-5C). We established these differences in

X-linked overexpression using two approaches. First, we tested which parental cell types had

the highest expression correlation with hybrid diplotene cell types for both X-linked and

autosomal genes. In mice, the expression profile of X-linked genes in hybrids during diplotene

was most positively correlated with the expression profile of X-linked parental round spermatid

genes, consistent with disrupted MSCI (spermatogonia *(r)* = -0.25, p < 0.001;

leptotene/zygotene *(r)* = -0.073, p = 0.035; round spermatid *(r)* = 0.26, p < 0.001; Figures 2 and

5A). Autosomal diplotene genes showed no positive correlations with either spermatogonia (r =

-0.26; p < 0.001), leptotene/zygotene (r = -0.032; p < 0.001), or round spermatids (r = -0.060; p

< 0.001; Figure 5A). If the X-linked overexpression phenotype in dwarf hamsters was

consistent with disrupted MSCI, then we would also expect X-linked and autosomal expression

profiles in hybrid diplotene to follow the same patterns. In contrast to this prediction, hybrid

dwarf hamster diplotene expression profiles for both X-linked and autosomal genes had a

positive correlation with parental leptotene/zygotene (autosomal *(r)* = 0.14, p < 0.001; X-linked

(r) = 0.22, p < 0.001) and spermatogonia (autosomal *(r)* = 0.027, p < 0.001; X-linked *(r)* = 0.091,

p = 0.016) and a negative correlation with round spermatids (autosomal *(r)* = -0.30, p < 0.001;

X-linked *(r)* = -0.31, p < 0.001; Figure 5B). These striking differences in the strength and

direction of expression profile correlations suggest that the regulatory mechanisms underlying

the overexpression phenotype of X-linked genes in sterile hybrids differed between house mice

and dwarf hamsters.

This difference in pattern was further supported when we compared which sets of

genes were overexpressed in hybrid diplotene in dwarf hamsters and house mice. For this

approach, we looked at parental gene expression patterns to characterize which stages of

spermatogenesis all genes were normally active during and characteristic of (*i.e.,* "induced";

see Methods). Using this information, we then identified which X-linked genes were

overexpressed in hybrid diplotene (defined as genes with normalized expression in the top

10% of X-linked genes) and assessed which parental stages the overexpressed X-linked genes

were characteristic of in both systems. As in our correlation analysis, we found that in hybrid

house mice, the genes that were overexpressed in diplotene most closely resembled genes

that are normally active in round spermatids in parental mice (51.6% of genes), but that in

hybrid dwarf hamsters, overexpressed diplotene genes resembled spermatogonia- and

leptotene/zygotene-specific genes (47.6% and 41.6% respectively; Figure 5C). Because of the

dissimilarity in the genes that were overexpressed during diplotene in both systems, we next

performed gene ontology (GO) enrichment analyses on the set of transgressive DE genes in

hybrids to determine which biological processes could be potentially contributing to this

pattern. In contrast to house mice hybrids where transgressive DE genes were enriched for no

biological processes, the transgressive DE genes in dwarf hamster were enriched for several

biological processes including cell junction/extracellular matrix organization, system

development, inflammatory response, and apoptotic process which together point to

widespread disruption of numerous processes necessary for normal male fertility (highlighted

terms presented in Figure 5D and the full lists in Tables S6-S8). Collectively, these results

suggest that the X-linked overexpression phenotype in sterile hybrid dwarf hamsters is

inconsistent with disrupted MSCI and is possibly related to a stalling and breakdown of

spermatogenesis between leptotene/zygotene and diplotene during Prophase I.

Figure 5. Spermatogenesis in hybrid dwarf hamsters appears to stall after leptotene/

zygotene. A. We calculated the Pearson's correlation coefficient *(r)* between mean hybrid

diplotene expression and the mean expression for each parental cell type for both house mice

and **B.** dwarf hamsters. Correlation coefficients were calculated for both autosomal (left panels)

and X-linked (right panels) genes. We then generated bootstrap values by randomly sampling

the expression matrices for 1000 replicates. All correlation coefficients were significantly

different from zero ($p < 0.05$) after FDR correction. **C.** Classification of X-linked overexpressed

genes in diplotene cells of hybrid mice and dwarf hamsters by parental stage in which genes

are induced. Genes are colored by cell type (red = spermatogonia, yellow =

leptotene/zygotene, green = diplotene, and blue = round spermatids). **D.** Select enriched

Biological Process GO terms (ranked by FDR) for transgressive DE genes between hybrid

dwarf hamsters and *P. campbelli* in diplotene spermatocytes*.* Included terms shown are the

result of the highlight function within gProfiler2 which collapses GO terms in a two-step

clustering algorithm (Kolberg *et al.* 2020). Point size corresponds to the number of genes

belonging to each GO term, and terms are plotted by the fold enrichment of the GO term in the

dataset relative to the provided gene backgrounds.

PAR expression was not disrupted in sterile hybrid dwarf hamsters

Finally, we tested the hypothesis that hybrid sterility in dwarf hamsters may be

correlated with asynapsis of the X and Y chromosomes because of divergence in the

pseudoautosomal region (PAR) which prevents proper chromosome pairing (Bikchurina *et al.*

2018). The PAR is the only portion of the sex chromosomes that is able to synapse during

routine spermatogenesis, and PAR genes are assumed to escape silencing by MSCI

(Raudsepp and Chowdhary 2015). However, because XY asynapsis is common in dwarf

hamster hybrids, Bikchurina et al. (2018) hypothesized that MSCI may extend to the PAR of

hybrid male dwarf hamsters, resulting in the silencing of PAR genes in hybrids that may be

critical to meiosis (Figure 6A). To test this hypothesis, we compared the expression of genes

located in the dwarf hamster PAR (Moore et al. 2022) between parental dwarf hamster species

and hybrid offspring across the timeline of spermatogenesis. Specifically, we hypothesized that

if XY asynapsis results in an extension of MSCI to the PAR in hybrid dwarf hamsters, then

hybrids should have similar PAR gene expression to parents early in meiosis before

homologous chromosome synapse during pachytene. This should be followed by the silencing

of PAR genes in hybrids, but not parent species, during diplotene. If XY asynapsis does not

alter the regulation of the PAR in hybrids, then we may see two possible patterns. First, if all

PAR genes are critical to the later stages of spermatogenesis, then PAR genes in both hybrids

and parents should be uniformly expressed in diplotene. Alternatively, if PAR genes are not

critical to the later stages of spermatogenesis, then hybrids and parent species should have

similar PAR gene expression, and not all PAR genes may be expressed during diplotene.

636   We did not find evidence supporting PAR-wide silencing in dwarf hamster hybrids

during diplotene suggesting that MSCI is not extended to the PAR in dwarf hamster hybrids

because of XY asynapsis (Figure 6B). Furthermore, we do not see PAR-wide expression of

genes during diplotene in hybrids or parents, indicating that not all PAR genes are critical to

the progression of spermatogenesis in dwarf hamsters. In general, most PAR gene expression

followed similar trends between hybrids and parent species. Two PAR genes were differentially

expressed between hybrids and *P. campbelli* during diplotene (*Ndrg2* and Psun_G000022883;

Table S3), but these genes were still expressed in hybrids. Further, an association between

PAR misregulation during the early stages of spermatogenesis and hybrid male sterility also

seems unlikely as only one gene, *Gprin1*, showed transgressive differential expression in early

meiosis between both parent species and hybrids (Table S3). Thus, based on the current

annotation of the PAR in *P. sungorus*, we currently find no direct evidence linking improper

silencing of PAR genes to hybrid male sterility in dwarf hamsters.

Figure 6. PAR gene expression is not disrupted during spermatogenesis in hybrid dwarf

hamsters. A. Hypothesized types of PAR gene expression across spermatogenesis. If MSCI

extends to the entire X chromosome, then PAR genes would show some level of expression

early in spermatogenesis which would then drop to zero in diplotene when MSCI occurs. If the

PAR escapes silencing by MSCI and if PAR genes are critical to spermatogenesis, then we

would expect PAR genes to be uniformly expressed in diplotene when the rest of the X

chromosome is silenced. Finally, if the PAR escapes silencing by MSCI but all PAR genes are

not critical to spermatogenesis, then we would expect some PAR genes to be expressed and

some to be not expressed in diplotene. **B.** Observed patterns of PAR gene expression (as

mean normalized RPKM across individuals) in parental species and hybrid dwarf hamsters for

all PAR genes (indicated by line type and color) across spermatogenesis (SP = spermatogonia,

LZ = leptotene/zygotene, and DIP = diplotene).

DISCUSSION

We used a comparative approach to understand common gene expression phenotypes

associated with hybrid male sterility in two divergent rodent crosses. We characterized the

asymmetry in the expression patterns of transgressive genes, how misexpression changed

over developmental timelines, and how the X chromosome and autosomes differed in both of

these aspects. We found that while there were similarities in hybrid expression phenotypes in

house mice and dwarf hamsters, there were also differences in the timing and chromosomal

distribution of disrupted gene expression that point towards different underlying mechanisms

behind hybrid male sterility.

Asymmetry and developmental timing of misexpression in hybrids

We first investigated patterns of gene misexpression in sterile male hybrids. Studies of

transgressive misexpression in sterile or inviable hybrids have often focused on whether hybrid

expression is biased towards over or underexpression, with the hypothesis that expression

may be biased towards overexpression if hybrid incompatibilities disrupt repressive gene

regulatory elements (Meiklejohn *et al.* 2014; Barreto *et al.* 2015; Larson *et al.* 2017). In house

mice, there is strong support for overexpression of both autosomal and X-linked genes in

sterile F1 hybrids (Mack *et al.* 2016; Larson *et al.* 2017, 2022; Hunnicutt *et al.* 2022), and further

work has shown that F1 hybrid expression patterns depend on both autosomal background

and sex chromosome mismatch (Kopania *et al.* 2022b). Surprisingly, we found that in dwarf

hamster hybrids, there was nearly uniform downregulation of transgressive DE genes in mitotic

cells and early meiotic cells, suggesting that a loss of regulatory repression is not an inevitable

outcome of hybrid genomes. Asymmetric patterns of misexpression have been found in many

hybrids, including underexpression in sterile *Drosophila* hybrids (Michalak and Noor 2003;

Haerty and Singh 2006; Llopart 2012) and sterile introgression lines of tomato (Guerrero *et al.*

2016) and *Drosophila* (Meiklejohn *et al.* 2014), but overexpression has also been found in other

sterile hybrids (Llopart 2012; Davis *et al.* 2015). The variation we and others have found in

hybrid expression phenotypes suggests that the mechanisms of disrupted expression are

complex, even within groups with relatively shallow divergence times (such as rodents), and we

need more data from diverse hybrid sterility systems to begin to understand common drivers of

transgressive hybrid misexpression.

The downregulation we observed in early spermatogenesis in hybrid dwarf hamsters

could be due to impaired transcription factor binding with promoter or enhancer elements (Oka

et al. 2014; Guerrero *et al.* 2016) or disrupted epigenetic silencing. Disruption of epigenetic

regulation of gene expression has been increasingly linked to hybrid dysfunction in plants

(Shivaprasad *et al.* 2012; Lafon-Placette and Köhler 2015; Zhu *et al.* 2017), especially

polyploids (Paun *et al.* 2007), and may also contribute to hybrid male sterility in *Drosophila*

(Bayes and Malik 2009) and cattle x yak hybrids (Luo *et al.* 2022). At least one known

chromatin difference, an expansion of the heterochromatin-enriched Xp arm of the X

chromosome, has been documented between parental dwarf hamster species (Gamperl *et al.*

1977; Haaf *et al.* 1987). However, it is unknown what the functional consequences of this

chromatin state divergence or other diverged epigenetic regulatory mechanisms, such as

methylation, may be in hybrid dwarf hamster spermatogenesis, and further work is needed to

distinguish between potential mechanisms underlying the observed genome-wide

downregulation.

Spermatogenesis as a developmental process may be sensitive to disruption (Lifschytz

and Lindsley 1972; Wu and Davis 1993), but it remains an open question whether specific

stages of spermatogenesis, or developmental processes more broadly, may be more prone to

the accumulation of hybrid incompatibilities. In general, earlier developmental stages are

thought to be under greater pleiotropic constraint and less prone to disruption (Cutter and

Bundus 2020). With the progression of mouse spermatogenesis, pleiotropy decreases (as

approximated by increases in tissue specificity; Murat *et al.* 2023) and the rate of protein-

coding evolution increases (Larson *et al.* 2016; Kopania *et al.* 2022a; Murat *et al.* 2023), which

may make the later stages of spermatogenesis more prone to accumulating hybrid

incompatibilities. Indeed, we found fewer DE genes both between parent species and in sterile

hybrids for both dwarf hamsters and house mice during the early stages of spermatogenesis

than in later stages (Figures S9 and S10), and hybrid misexpression greatly exceeds parental

expression divergence in late spermatogenesis. Despite general similarity in patterns of

misexpression across spermatogenesis in hybrids, we found that few DE genes were shared

between dwarf hamster and house mouse hybrids, suggesting that shared genes of large

effect are unlikely to be responsible for the repeated evolution of hybrid male sterility. The low

levels of disrupted gene expression early in spermatogenesis that we and others have shown

suggests that there might be some tolerance for misregulation early in spermatogenesis, since

misregulation of some genes during early stages does not always completely halt the

progression of sperm development (Oka *et al.* 2010; Ishishita *et al.* 2015; Mipam *et al.* 2023).

However, the patterns we find in dwarf hamsters suggests that spermatogenesis may be

disrupted between zygotene and diplotene cell stages from early misexpression. Dwarf

hamster hybrid diplotene cell populations had X-linked and autosomal gene expression profiles

which more closely resemble parental leptotene/zygotene cell populations than either diplotene

or postmeiotic cell populations. Furthermore, DE genes in hybrids during diplotene were

enriched for genes associated with cell differentiation, proliferation, and programmed cell

death, suggesting misexpression during this stage could be a consequence of a stalling or

breakdown of early meiosis in hybrid dwarf hamsters. It's unclear what underlying genomic

mechanisms could result in this breakdown, but it is possible that this disruption could

potentially act as a major contributor to hybrid sterility in this system. Ultimately, we find that

spermatogenesis is a complex and rapidly evolving developmental program that may provide

many potential avenues across its timeline for the evolution of hybrid incompatibilities.

Abnormal sex chromosome expression in dwarf hamster hybrids is inconsistent with

disrupted MSCI

The sex chromosomes play a central role in speciation, an observation which has been

supported by both Haldane's rule (Haldane 1922) and the large X-effect on hybrid male sterility

(Coyne and Orr 1989). Misregulation of the X chromosome may contribute to hybrid sterility in

several species pairs (Davis *et al.* 2015; Morgan *et al.* 2020; Sánchez-Ramírez *et al.* 2021). The

X chromosome is transcriptionally repressed during routine spermatogenesis in many

organisms including eutherian mammals (McKee and Handel 1993), monotremes (Murat *et al.*

2023), *Drosophila* (Landeen *et al.* 2016), grasshoppers (Viera *et al.* 2021), mosquitos (Taxiarchi

et al. 2019), and nematodes (Rappaport *et al.* 2021). Because of the ubiquity of X chromosome

repression during spermatogenesis, disruption of transcriptional repression could be a

widespread regulatory phenotype in sterile hybrids (Lifschytz and Lindsley 1972; Larson *et al.*

2018). In sterile hybrid mice, disrupted X repression (disrupted MSCI) leads to the

overexpression of the normally silenced X chromosome during diplotene (Good *et al.* 2010;

Bhattacharyya *et al.* 2013; Campbell *et al.* 2013; Turner and Harr 2014; Larson *et al.* 2017,

2022). We also found overexpression of the X chromosome during diplotene in sterile hybrid

dwarf hamsters, but in a manner inconsistent with disrupted MSCI. Both the X chromosome

and autosomes are overexpressed in dwarf hamster hybrid diplotene cells to a greater extent

on average than was observed in house mice, and importantly, X-linked overexpression was

more variable in dwarf hamster hybrids than house mice hybrids. In fact, some dwarf hamster

hybrids had wildly overexpressed X chromosomes while others appeared to have properly

silenced X chromosomes. Our expression correlation and gene set analyses of hybrid diplotene

cells provide additional evidence that the genes overexpressed in hybrid hamster diplotene are

different than those overexpressed in house mouse hybrids, sharing more similarity to the

earlier meiotic cell types than downstream postmeiotic cell types. Overall, our results indicate

fundamentally different patterns of X-linked overexpression in both systems, with X-linked

overexpression in dwarf hamster hybrids being inconsistent with disrupted MSCI.

Much of what we know about the genomic architecture and the role of sex

chromosome misregulation in hybrid male sterility in mammals comes from decades of work

that have shown a major gene, *Prdm9*, and its X chromosome modulator, *Hstx2*, may be

responsible for most F1 hybrid male sterility in house mice (Forejt *et al.* 1991, 2021; Trachtulec

et al. 1997; Mihola *et al.* 2009; Lustyk *et al.* 2019). *Prdm9* directs the location of double strand

breaks during meiotic recombination (Mihola *et al.* 2009; Oliver *et al.* 2009; Smagulova *et al.*

2016). In hybrid mice, divergence at *Prdm9* binding sites leads to asymmetric double-stranded

breaks and results in autosomal asynapsis, triggering Meiotic Silencing of Unsynapsed

Chromatin, shutting down transcription on asynapsed autosomes using the same cellular

machinery as MSCI (Turner 2015), and eventually meiotic arrest and cell death (Bhattacharyya

et al. 2013; Forejt *et al.* 2021). This process is associated with the disruption of MSCI and a

characteristic overexpression of the X chromosome during meiosis (Good *et al.* 2010;

Bhattacharyya *et al.* 2013; Campbell *et al.* 2013; Turner and Harr 2014; Larson *et al.* 2017,

2022), but whether disrupted MSCI directly contributes to hybrid male sterility or is simply a

downstream consequence of *Prdm9* divergence is still uncertain (Forejt *et al.* 2021).

Whether we should have expected patterns of disrupted sex chromosome expression

in sterile hybrid hamsters to be the same as house mice is unclear. The sex chromosomes in

pachytene cells of hybrid dwarf hamsters display normal γH2AFX staining (Ishishita *et al.* 2015;

Bikchurina *et al.* 2018), a key marker in MSCI (Abe *et al.* 2022), which may indicate that the

hybrid sex chromosomes are properly silenced. Additionally, autosomal asynapsis is rarely

observed in hybrid dwarf hamsters, and asynapsis is almost exclusive to the sex chromosomes

(Ishishita *et al.* 2015; Bikchurina *et al.* 2018). This contrasts *Prdm9*-mediated sterility in house

mice, where hybrid autosomes are often asynapsed and decorated with yH2AFX

(Bhattacharyya *et al.* 2013; Forejt *et al.* 2021). Mechanisms other than *Prdm9* may also disrupt

MSCI and result in sterility, such as macrosatellite copy number divergence (Bredemeyer *et al.*

2021) and X-autosome translocations that impair synapsis (Homolka *et al.* 2007), although

there is no evidence for X-autosome translocations between these two species of dwarf

hamsters (Moore *et al.* 2022). Thus, while MSCI may be a major target for the accumulation of

reproductive barriers between species in many mammalian systems, either through *Prdm9*

divergence or alternative mechanisms, our results suggest that sterility in dwarf hamsters has a

more composite regulatory basis.

801   Another mechanism often proposed to underlie mammalian male hybrid sterility,

especially in rodents, is divergence in the PAR between parental species. The PAR is the only

portion of the sex chromosomes which can synapse during spermatogenesis, and it is still

unclear if the regulation of the PAR is uniformly detached from MSCI across divergent

mammalian sex chromosome systems (Raudsepp and Chowdhary 2015). We find no evidence

that PAR-specific misregulation is associated with hybrid sterility in dwarf hamsters. PAR

genes are not silenced in hybrids or parents, a pattern that is inconsistent with MSCI that has

extended to the PAR due to XY asynapsis, and further, expression of PAR genes in hybrids

differs little from parental PAR expression. While we find no evidence that PAR misregulation

per se is associated with hybrid sterility in this system, we cannot rule out the possibility that

structural and sequence divergence between the PARs of *P. sungorus* and *P. campbelli* may

be associated with hybrid sterility. Structural and sequence divergence in the PAR has been

hypothesized to activate the meiotic spindle checkpoint by interfering with proper pairing of

sex chromosomes (Burgoyne *et al.* 2009; Dumont 2017). The PAR evolves rapidly in rodents

(White *et al.* 2012b; Raudsepp and Chowdhary 2015; Morgan *et al.* 2019), and this elevated

divergence may underlie sex chromosome asynapsis and apoptosis in several hybrid mouse

crosses (Matsuda *et al.* 1991; Oka *et al.* 2010; White *et al.* 2012a; Dumont 2017). Furthermore,

divergence in the mouse PAR has been implicated in spermatogenic defects in crosses where

Prdm9-divergence is minimal, such as between closely related subspecies (Dumont 2017) or in

mice with genetically-modified *Prdm9* alleles (Davies *et al.* 2021). Meiosis is likely tolerant to

some degree of divergence in the PAR (Morgan *et al.* 2019), but exact limits are currently

unknown. At this time, thorough analysis of structural and sequence divergence between the

PARs in dwarf hamsters is challenging as the PAR is notoriously difficult to assemble (but see

Kasahara *et al.* 2022), and there are annotation gaps in the current assembly of the PAR in

dwarf hamsters. In sum, we find no clear pattern of regulatory disruption of PAR genes in

sterile hybrid dwarf hamsters, though this result may change pending further refinement of the

PAR annotation.

Conclusions

Cell-specific approaches for quantifying expression phenotypes are powerful tools for

providing insight into the underlying mechanisms behind hybrid dysfunction (Hunnicutt *et al.*

2022), especially in systems where it remains difficult to interrogate the underlying genomic

architecture of these traits. Using a contrast of dwarf hamster and house mouse hybrids, we

have shown that transgressive overexpression is not an inevitable outcome of hybridization,

that hybrid incompatibilities are likely to arise during multiple stages of spermatogenesis, and

that disrupted sex chromosome silencing does not appear to play an equal role in sterility

between these two systems. Both the expression phenotypes we observed here and

histological evidence from other studies (Ishishita *et al.* 2015; Bikchurina *et al.* 2018) suggest

that several reproductive barriers are acting during spermatogenesis in dwarf hamster hybrids.

It has become increasingly apparent as more study systems are investigated that the genetic

basis of postzygotic species barriers are often complex and polymorphic (Cutter 2012;

Coughlan and Matute 2020), and implementing approaches which account for the

developmental complexities of hybrid dysfunction, as we have done here, will allow us to make

further advances in understanding the processes of speciation.

Acknowledgements:

This work was supported by a National Science Foundation Graduate Research Fellowship to

KEH (DGE-2034612) and the following grants awarded to KEH: Sigma Xi Grant-in-aid of

Research, American Society of Mammalogists Grant-in-Aid of Research Grant, and a Society

for the Study of Evolution R.C. Lewontin Early Award. Additionally, this work was supported by

grants from the Eunice Kennedy Shriver National Institute of Child Health and Human

Development of the National Institutes of Health (R01-HD073439, R01-HD094787) to JMG, and

the National Science Foundation to ELL (DEB-2012041). We would like to thank members of

the Larson and Good labs, as well as Jonathan Velotta and members of the Velotta lab, for

feedback on this project, Gabrielle Welsh for technical assistance, Ivan Kovanda for support

using the University of Denver Research Data Analysis Cluster, and Pamela K. Shaw and the

UM Fluorescence Cytometry Core supported by an Institutional Development Award from the

NIGMS (P30GM103338). Any opinions, findings, and conclusions or recommendations

expressed in this material are those of the author(s) and do not necessarily reflect the views of

the National Science Foundation or the National Institutes of Health.

Author contributions:

KEH, JMG, and ELL conceived of the study. CC, SK, ELL, and KEH conducted lab work. KEH

conducted the analyses. KEH and ELL wrote the manuscript with input from ECM, CC, SK, and

JMG.

Data accessibility:

Hamster sequence data are available through the NCBI SRA under accession number

PRJNA1024468, and mouse sequence data are available under SRA accession PRJNA296926.

File S1 contains all supplemental figures. Code used for the analyses and fertility phenotypic

data are available from GitHub (https://github.com/KelsieHunnicutt/dwarf_hamster_hybrids).

LITERATURE CITED

870
871

Abe H., Y.-H. Yeh, Y. Munakata, K.-I. Ishiguro, P. R. Andreassen, *et al.*, 2022 Active DNA

damage response signaling initiates and maintains meiotic sex chromosome

inactivation. Nat. Commun. 13: 7212.

Anders S., and W. Huber, 2010 Differential expression analysis for sequence count data.

Nature Precedings 1–1.

Barreto F. S., R. J. Pereira, and R. S. Burton, 2015 Hybrid dysfunction and physiological

compensation in gene expression. Mol. Biol. Evol. 32: 613–622.

Bayes J. J., and H. S. Malik, 2009 Altered heterochromatin binding by a hybrid sterility protein

in Drosophila sibling species. Science 326: 1538–1541.

Benjamini Y., and Y. Hochberg, 1995 Controlling the false discovery rate: a practical and

powerful approach to multiple testing. J. R. Stat. Soc. 57: 289–300.

Bhattacharyya T., S. Gregorova, O. Mihola, M. Anger, J. Sebestova, *et al.*, 2013 Mechanistic

basis of infertility of mouse intersubspecific hybrids. Proc. Natl. Acad. Sci. U. S. A. 110:

E468–E477.

Bikchurina T. I., K. V. Tishakova, E. A. Kizilova, S. A. Romanenko, N. A. Serdyukova, *et al.*,

2018 Chromosome synapsis and recombination in male-sterile and female-fertile

interspecies hybrids of the dwarf hamsters (Phodopus, Cricetidae). Genes 9: 227.

Bolger A. M., M. Lohse, and B. Usadel, 2014 Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics 30: 2114–2120.

Bredemeyer K. R., C. M. Seabury, M. J. Stickney, J. R. McCarrey, B. M. vonHoldt, *et al.*, 2021

Rapid macrosatellite evolution promotes X-linked hybrid male sterility in a feline

893   interspecies cross. Mol. Biol. Evol. 38: 5588–5609.

894   Brekke T. D., and J. M. Good, 2014 Parent-of-origin growth effects and the evolution of hybrid

895   inviability in dwarf hamsters. Evolution 68: 3134–3148.

896   Brekke T. D., L. A. Henry, and J. M. Good, 2016 Genomic imprinting, disrupted placental

897   expression, and speciation. Evolution 70: 2690–2703.

898   Brekke T. D., K. A. Steele, and J. F. Mulley, 2018 Inbred or outbred? Genetic diversity in

899   laboratory rodent colonies. G3 8: 679–686.

900   Brekke T. D., E. C. Moore, S. C. Campbell-Staton, C. M. Callahan, Z. A. Cheviron, *et al.*, 2021 X

901   chromosome-dependent disruption of placental regulatory networks in hybrid dwarf

902   hamsters. Genetics 218: iyab043.

903   Burgoyne P. S., 1982 Genetic homology and crossing over in the X and Y chromosomes of

904   mammals. Hum. Genet. 61: 85–90.

905   Burgoyne P. S., S. K. Mahadevaiah, and J. M. A. Turner, 2009 The consequences of asynapsis

906   for mammalian meiosis. Nat. Rev. Genet. 10: 207–216.

907   Campbell P., J. M. Good, and M. W. Nachman, 2013 Meiotic sex chromosome inactivation is

908   disrupted in sterile hybrid male house mice. Genetics 193: 819–828.

909   Civetta A., 2016 Misregulation of gene expression and sterility in interspecies hybrids: causal

910   links and alternative hypotheses. J. Mol. Evol. 82: 176–182.

911   Coughlan J. M., and D. R. Matute, 2020 The importance of intrinsic postzygotic barriers

912   throughout the speciation process. Philos. Trans. R. Soc. Lond. B Biol. Sci. 375:

20190533.

Coyne J. A., and H. A. Orr, 1989 Two rules of speciation, pp. 180–207 in *Speciation and its*

consequences, edited by Otte D., Endler J. A. Sinauer Associates, Sunderland, MA.

Cutter A. D., 2012 The polymorphic prelude to Bateson-Dobzhansky-Muller incompatibilities.

Trends Ecol. Evol. 27: 209–218.

Cutter A. D., and J. D. Bundus, 2020 Speciation and the developmental alarm clock. Elife 9:

e56276.

Danecek P., A. Auton, G. Abecasis, C. A. Albers, E. Banks, *et al.*, 2011 The variant call format

and VCFtools. Bioinformatics 27: 2156–2158.

Davies B., E. Hatton, N. Altemose, J. G. Hussin, F. Pratto, *et al.*, 2016 Re-engineering the zinc

fingers of PRDM9 reverses hybrid sterility in mice. Nature 530: 171–176.

Davies B., A. Gupta Hinch, A. Cebrian-Serrano, S. Alghadban, P. W. Becker, *et al.*, 2021

Altering the binding properties of PRDM9 partially restores fertility across the species

boundary. Mol. Biol. Evol. 38: 5555–5562.

Davis B. W., C. M. Seabury, W. A. Brashear, G. Li, M. Roelke-Parker, *et al.*, 2015 Mechanisms

underlying mammalian hybrid sterility in two feline interspecies models. Mol. Biol. Evol.

32: 2534–2546.

Dumont B. L., 2017 Meiotic consequences of genetic divergence across the murine

pseudoautosomal region. Genetics 205: 1089–1100.

Ellis N., and P. N. Goodfellow, 1989 The mammalian pseudoautosomal region. Trends Genet.

5: 406–410.

Forejt J., V. Vincek, J. Klein, H. Lehrach, and M. Loudová-Micková, 1991 Genetic mapping of

the t-complex region on mouse chromosome 17 including the Hybrid sterility-1 gene.

Mamm. Genome 1: 84–91.

Forejt J., P. Jansa, and E. Parvanov, 2021 Hybrid sterility genes in mice (Mus musculus): a

peculiar case of PRDM9 incompatibility. Trends Genet. 37: 1095–1108.

Gamperl R., G. Vistorin, and W. Rosenkranz, 1977 New observations on the karyotype of the

Djungarian hamster, Phodopus sungorus. Experientia 33: 1020–1021.

Getun I. V., B. Torres, and P. R. J. Bois, 2011 Flow cytometry purification of mouse meiotic

cells. J. Vis. Exp. e2602.

Good J. M., M. A. Handel, and M. W. Nachman, 2008 Asymmetry and polymorphism of hybrid

male sterility during the early stages of speciation in house mice. Evolution 62: 50–65.

Good J. M., T. Giger, M. D. Dean, and M. W. Nachman, 2010 Widespread over-expression of

the X chromosome in sterile F1 hybrid mice. PLoS Genet. 6: e1001148.

Green C. D., Q. Ma, G. L. Manske, A. N. Shami, X. Zheng, *et al.*, 2018 A comprehensive

roadmap of murine spermatogenesis defined by single-cell RNA-Seq. Dev. Cell 46:

651-667.e10.

Gu Z., R. Eils, and M. Schlesner, 2016 Complex heatmaps reveal patterns and correlations in

multidimensional genomic data. Bioinformatics 32: 2847–2849.

Guerrero R. F., A. L. Posto, L. C. Moyle, and M. W. Hahn, 2016 Genome-wide patterns of

regulatory divergence revealed by introgression lines. Evolution 70: 696–706.

Haaf T., H. Weis, and M. Schmid, 1987 A comparative cytogenetic study on the mitotic and

meiotic chromosomes in hamster species of the genus Phodopus (Rodentia,

Cricetinae). Zeitschrift für Säugetierkunde 52: 281–290.

Haerty W., and R. S. Singh, 2006 Gene regulation divergence is a major contributor to the

evolution of Dobzhansky-Muller incompatibilities between species of Drosophila. Mol.

Biol. Evol. 23: 1707–1714.

Haldane J. B. S., 1922 Sex ratio and unisexual sterility in hybrid animals. J. Genet. 12: 101–

109.

Handel M. A., 2004 The XY body: a specialized meiotic chromatin domain. Exp. Cell Res. 296:

57–63.

Hermann B. P., K. Cheng, A. Singh, L. Roa-De La Cruz, K. N. Mutoji, *et al.*, 2018 The

mammalian spermatogenesis single-cell transcriptome, from spermatogonial stem cells

to spermatids. Cell Rep. 25: 1650-1667.e8.

Holt J., S. Huang, L. McMillan, and W. Wang, 2013 Read annotation pipeline for high-

throughput sequencing data, pp. 605–612 in *Proceedings of the International*

Conference on Bioinformatics, Computational Biology and Biomedical Informatics -

BCB'13, ACM Press, New York, New York, USA.

Homolka D., R. Ivanek, J. Capkova, P. Jansa, and J. Forejt, 2007 Chromosomal rearrangement

interferes with meiotic X chromosome inactivation. Genome Res. 17: 1431–1437.

Huang S., J. Holt, C.-Y. Kao, L. McMillan, and W. Wang, 2014 A novel multi-alignment pipeline

for high-throughput sequencing data. Database 2014: bau057.

Hunnicutt K. E., J. M. Good, and E. L. Larson, 2022 Unraveling patterns of disrupted gene

expression across a complex tissue. Evolution 76: 275–291.

Ishishita S., K. Tsuboi, N. Ohishi, K. Tsuchiya, and Y. Matsuda, 2015 Abnormal pairing of X and

Y sex chromosomes during meiosis I in interspecific hybrids of Phodopus campbelli

and P. sungorus. Sci. Rep. 5: 9435.

Kasahara T., K. Mekada, K. Abe, A. Ashworth, and T. Kato, 2022 Complete sequencing of the

mouse pseudoautosomal region, the most rapidly evolving 'chromosome.' bioRxiv

2022.03.26.485930.

Kim D., J. M. Paggi, C. Park, C. Bennett, and S. L. Salzberg, 2019 Graph-based genome

alignment and genotyping with HISAT2 and HISAT-genotype. Nat. Biotechnol. 37: 907–

915.

Kolberg L., U. Raudvere, I. Kuzmin, J. Vilo, and H. Peterson, 2020 gprofiler2 -- an R package

for gene list functional enrichment analysis and namespace conversion toolset

g:Profiler. F1000Res. 9. https://doi.org/10.12688/f1000research.24956.2

Kopania E. E. K., E. L. Larson, C. Callahan, S. Keeble, and J. M. Good, 2022a Molecular

evolution across mouse spermatogenesis. Mol. Biol. Evol. 39: msac023.

Kopania E. E. K., E. M. Watson, C. C. Rathje, B. M. Skinner, P. J. I. Ellis, *et al.*, 2022b The

contribution of sex chromosome conflict to disrupted spermatogenesis in hybrid house

mice. Genetics 222: iyac151.

Korunes K. L., and K. Samuk, 2021 pixy: Unbiased estimation of nucleotide diversity and

divergence in the presence of missing data. Mol. Ecol. Resour. 21: 1359–1368.

Kousathanas A., D. L. Halligan, and P. D. Keightley, 2014 Faster-X adaptive protein evolution in

house mice. Genetics 196: 1131–1143.

Lafon-Placette C., and C. Köhler, 2015 Epigenetic mechanisms of postzygotic reproductive

isolation in plants. Curr. Opin. Plant Biol. 23: 39–44.

Landeen E. L., C. A. Muirhead, L. Wright, C. D. Meiklejohn, and D. C. Presgraves, 2016 Sex

chromosome-wide transcriptional suppression and compensatory cis-regulatory

evolution mediate gene expression in the Drosophila male germline. PLoS Biol. 14:

e1002499.

Larson E. L., D. Vanderpool, S. Keeble, M. Zhou, B. A. J. Sarver, *et al.*, 2016 Contrasting levels

of molecular evolution on the mouse X chromosome. Genetics 203: 1841–1857.

Larson E. L., S. Keeble, D. Vanderpool, M. D. Dean, and J. M. Good, 2017 The composite

regulatory basis of the large X-effect in mouse speciation. Mol. Biol. Evol. 34: 282–295.

Larson E. L., E. E. K. Kopania, and J. M. Good, 2018 Spermatogenesis and the evolution of

mammalian sex chromosomes. Trends Genet. 34: 722–732.

Larson E. L., E. E. K. Kopania, K. E. Hunnicutt, D. Vanderpool, S. Keeble, *et al.*, 2022 Stage-

specific disruption of X chromosome expression during spermatogenesis in sterile

house mouse hybrids. G3 12: jkab407.

Liao Y., G. K. Smyth, and W. Shi, 2014 featureCounts: an efficient general purpose program for

assigning sequence reads to genomic features. Bioinformatics 30: 923–930.

Lifschytz E., and D. L. Lindsley, 1972 The role of X-chromosome inactivation during

spermatogenesis. Proc. Natl. Acad. Sci. U. S. A. 69: 182–186.

Llopart A., 2012 The rapid evolution of X-linked male-biased gene expression and the large-X

effect in Drosophila yakuba, D. santomea, and their hybrids. Mol. Biol. Evol. 29: 3873–

3886.

Luo H., T. Mipam, S. Wu, C. Xu, C. Yi, *et al.*, 2022 DNA methylome of primary spermatocyte

reveals epigenetic dysregulation associated with male sterility of cattleyak.

Theriogenology 191: 153–167.

Lustyk D., S. Kinský, K. K. Ullrich, M. Yancoskie, L. Kašíková, *et al.*, 2019 Genomic structure of

Hstx2 modifier of Prdm9-dependent hybrid male sterility in mice. Genetics 213: 1047–

1063.

Mack K. L., P. Campbell, and M. W. Nachman, 2016 Gene regulation and speciation in house

mice. Genome Res. 26: 451–461.

Mack K. L., and M. W. Nachman, 2017 Gene regulation and speciation. Trends Genet. 33: 68–

80.

Matsuda Y., T. Hirobe, and V. M. Chapman, 1991 Genetic basis of X-Y chromosome

dissociation and male sterility in interspecific hybrids. Proc. Natl. Acad. Sci. U. S. A. 88:

4850–4854.

McCarthy D. J., Y. Chen, and G. K. Smyth, 2012 Differential expression analysis of multifactor

RNA-Seq experiments with respect to biological variation. Nucleic Acids Res. 40: 4288–

4297.

McKee B. D., and M. A. Handel, 1993 Sex chromosomes, recombination, and chromatin

conformation. Chromosoma 102: 71–80.

McManus C. J., J. D. Coolon, M. O. Duff, J. Eipper-Mains, B. R. Graveley, *et al.*, 2010

Regulatory divergence in Drosophila revealed by mRNA-seq. Genome Res. 20: 816–

825.

Meiklejohn C. D., J. D. Coolon, D. L. Hartl, and P. J. Wittkopp, 2014 The roles of cis- and trans-

regulation in the evolution of regulatory incompatibilities and sexually dimorphic gene

expression. Genome Res. 24: 84–95.

Michalak P., and M. A. F. Noor, 2003 Genome-wide patterns of expression in Drosophila pure

species and hybrid males. Mol. Biol. Evol. 20: 1070–1076.

Mihola O., Z. Trachtulec, C. Vlcek, J. C. Schimenti, and J. Forejt, 2009 A mouse speciation

gene encodes a meiotic histone H3 methyltransferase. Science 323: 373–375.

Mipam T., X. Chen, W. Zhao, P. Zhang, Z. Chai, *et al.*, 2023 Single-cell transcriptome analysis

and in vitro differentiation of testicular cells reveal novel insights into male sterility of the

interspecific hybrid cattle-yak. BMC Genomics 24: 149.

Montgomery S. H., and J. E. Mank, 2016 Inferring regulatory change from gene expression: the

confounding effects of tissue scaling. Mol. Ecol. 25: 5114–5128.

Moore E. C., G. W. C. Thomas, S. Mortimer, E. E. K. Kopania, K. E. Hunnicutt, *et al.*, 2022 The

evolution of widespread recombination suppression on the dwarf hamster (Phodopus) X

chromosome. Genome Biol. Evol. 14: evac080.

Morgan A. P., T. A. Bell, J. J. Crowley, and F. Pardo-Manuel de Villena, 2019 Instability of the

pseudoautosomal boundary in house mice. Genetics 212: 469–487.

Morgan K., B. Harr, M. A. White, B. A. Payseur, and L. M. Turner, 2020 Disrupted gene

networks in subfertile hybrid house mice. Mol. Biol. Evol. 37: 1547–1562.

Mugal C. F., M. Wang, N. Backström, D. Wheatcroft, M. Ålund, *et al.*, 2020 Tissue-specific

patterns of regulatory changes underlying gene expression differences among *Ficedula*

flycatchers and their naturally occurring F1 hybrids. Genome Res. 30: 1727–1739.

Murat F., N. Mbengue, S. B. Winge, T. Trefzer, E. Leushkin, *et al.*, 2023 The molecular evolution

of spermatogenesis across mammals. Nature 613: 308–316.

Namekawa S. H., P. J. Park, L.-F. Zhang, J. E. Shima, J. R. McCarrey, *et al.*, 2006 Postmeiotic

sex chromatin in the male germline of mice. Curr. Biol. 16: 660–667.

Neumann K., J. Michaux, V. Lebedev, N. Yigit, E. Colak, *et al.*, 2006 Molecular phylogeny of the

Cricetinae subfamily based on the mitochondrial cytochrome b and 12S rRNA genes

and the nuclear vWF gene. Mol. Phylogenet. Evol. 39: 135–148.

Ogle D., and M. D. Ogle, 2017 Package 'FSA.' CRAN Repos 1–206.

Oka A., A. Mita, Y. Takada, H. Koseki, and T. Shiroishi, 2010 Reproductive isolation in hybrid

mice due to spermatogenesis defects at three meiotic stages. Genetics 186: 339–351.

Oka A., and T. Shiroishi, 2014 Regulatory divergence of X-linked genes and hybrid male

sterility in mice. Genes Genet. Syst. 89: 99–108.

Oka A., T. Takada, H. Fujisawa, and T. Shiroishi, 2014 Evolutionarily diverged regulation of X-

chromosomal genes as a primal event in mouse reproductive isolation. PLoS Genet. 10:

e1004301.

Oksanen J., F. G. Blanchet, R. Kindt, P. Legendre, P. R. Minchin, *et al.*, 2013 Package 'vegan.'

Community ecology package, version 2: 1–295.

Oliver P. L., L. Goodstadt, J. J. Bayes, Z. Birtle, K. C. Roach, *et al.*, 2009 Accelerated evolution

of the Prdm9 speciation gene across diverse metazoan taxa. PLoS Genet. 5: e1000753.

Ortíz-Barrientos D., B. A. Counterman, and M. A. F. Noor, 2007 Gene expression divergence

and the origin of hybrid dysfunctions. Genetica 129: 71–81.

Paun O., M. F. Fay, D. E. Soltis, and M. W. Chase, 2007 Genetic and epigenetic alterations

after hybridization and genome doubling. Taxon 56: 649–656.

Presgraves D. C., 2008 Sex chromosomes and speciation in Drosophila. Trends Genet. 24:

336–343.

Ramm S. A., and L. Schärer, 2014 The evolutionary ecology of testicular function: size isn't

everything. Biol. Rev. Camb. Philos. Soc. 89: 874–888.

Rappaport Y., H. Achache, R. Falk, O. Murik, O. Ram, *et al.*, 2021 Bisection of the X

chromosome disrupts the initiation of chromosome silencing during meiosis in

Caenorhabditis elegans. Nat. Commun. 12: 4802.

Raudsepp T., and B. P. Chowdhary, 2015 The eutherian pseudoautosomal region. Cytogenet.

Genome Res. 147: 81–94.

Raymond C. S., M. W. Murphy, M. G. O'Sullivan, V. J. Bardwell, and D. Zarkower, 2000 Dmrt1,

a gene related to worm and fly sexual regulators, is required for mammalian testis

differentiation. Genes Dev. 14: 2587–2595.

Ritchie M. E., B. Phipson, D. Wu, Y. Hu, C. W. Law, *et al.*, 2015 limma powers differential

expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Res.

43: e47.

Robinson M. D., D. J. McCarthy, and G. K. Smyth, 2010 edgeR: a Bioconductor package for

differential expression analysis of digital gene expression data. Bioinformatics 26: 139–

140.

Sánchez-Ramírez S., J. G. Weiss, C. G. Thomas, and A. D. Cutter, 2021 Widespread

misregulation of inter-species hybrid transcriptomes due to sex-specific and sex-

chromosome regulatory evolution. PLoS Genet. 17: e1009409.

Scribner S. J., and K. E. Wynne-Edwards, 1994 Disruption of body temperature and behavior

rhythms during reproduction in dwarf hamsters (Phodopus). Physiol. Behav. 55: 361–

369.

Shivaprasad P. V., R. M. Dunn, B. A. Santos, A. Bassett, and D. C. Baulcombe, 2012

Extraordinary transgressive phenotypes of hybrid tomato are influenced by epigenetics

and small silencing RNAs. EMBO J. 31: 257–266.

Smagulova F., K. Brick, Y. Pu, R. D. Camerini-Otero, and G. V. Petukhova, 2016 The

evolutionary turnover of recombination hot spots contributes to speciation in mice.

Genes Dev. 30: 266–280.

Swanson M. T., C. H. Oliveros, and J. A. Esselstyn, 2019 A phylogenomic rodent tree reveals

the repeated evolution of masseter architectures. Proc. Biol. Sci. 286: 20190672.

Taxiarchi C., N. Kranjc, A. Kriezis, K. Kyrou, F. Bernardini, *et al.*, 2019 High-resolution

transcriptional profiling of Anopheles gambiae spermatogenesis reveals mechanisms of

sex chromosome regulation. Sci. Rep. 9: 14841.

Trachtulec Z., M. Mnuková-Fajdelová, R. M. Hamvas, S. Gregorová, W. E. Mayer, *et al.*, 1997

Isolation of candidate hybrid sterility 1 genes by cDNA selection in a 1.1 megabase pair

region on mouse chromosome 17. Mamm. Genome 8: 312–316.

Turner L. M., M. A. White, D. Tautz, and B. A. Payseur, 2014 Genomic networks of hybrid

sterility. PLoS Genet. 10: e1004162.

Turner L. M., and B. Harr, 2014 Genome-wide mapping in a house mouse hybrid zone reveals

hybrid sterility loci and Dobzhansky-Muller interactions. Elife 3: e02504.

Turner J. M. A., 2015 Meiotic silencing in mammals. Annu. Rev. Genet. 49: 395–412.

Vasimuddin M., S. Misra, H. Li, and S. Aluru, 2019 Efficient architecture-aware acceleration of

BWA-MEM for multicore systems, pp. 314–324 in *2019 IEEE International Parallel and*

Distributed Processing Symposium (IPDPS), IEEE.

Viera A., M. T. Parra, S. Arévalo, C. García de la Vega, J. L. Santos, *et al.*, 2021 X chromosome

inactivation during grasshopper spermatogenesis. Genes 12: 1844.

White M. A., M. Stubbings, B. L. Dumont, and B. A. Payseur, 2012a Genetics and evolution of

hybrid male sterility in house mice. Genetics 191: 917–934.

White M. A., A. Ikeda, and B. A. Payseur, 2012b A pronounced evolutionary shift of the

pseudoautosomal region boundary in house mice. Mamm. Genome 23: 454–466.

Wittkopp P. J., B. K. Haerum, and A. G. Clark, 2004 Evolutionary changes in cis and trans gene

regulation. Nature 430: 85–88.

Wu C. I., and A. W. Davis, 1993 Evolution of postmating reproductive isolation: the composite

nature of Haldane's rule and its genetic bases. Am. Nat. 142: 187–212.

Zhu W., B. Hu, C. Becker, E. S. Doğan, K. W. Berendzen, *et al.*, 2017 Altered chromatin

compaction and histone methylation drive non-additive gene expression in an

interspecific Arabidopsis hybrid. Genome Biol. 18: 157.