Title: Decoding Heterogenous Single-cell Perturbation Responses

One sentence summary: We present a method to quantify diverse perturbation responses and discover
novel biological insights in single-cell perturbation datasets.

Bicna Song[1,2], Dingyu Liu[3,4], Weiwei Dai[5,6], Natalie McMyn[5], Qingyang Wang[7], Dapeng Yang[3], Adam
Krejci[8], Anatoly Vasilyev[8], Nicole Untermoser[8], Anke Loregger[8], Dongyuan Song[9], Breanna Williams[3],
Bess Rosen[3,10], Xiaolong Cheng[1,2], Lumen Chao[1,2], Hanuman T. Kale[3], Hao Zhang[5], Yarui Diao[11],
Tilmann Bürckstümmer[8], Jenet M. Siliciano[5], Jingyi Jessica Li[7,9,12-14,], Robert Siliciano[5,6], Danwei
Huangfu[3], Wei Li[1,2,#]

1 Center for Genetic Medicine Research, Children's National Hospital, Washington DC, USA
2 Department of Genomics and Precision Medicine, George Washington University, Washington DC,
USA
3 Developmental Biology Program, Sloan Kettering Institute, New York City, NY, USA
4 Louis V. Gerstner Jr. Graduate School of Biomedical Sciences, Memorial Sloan Kettering Cancer
Center, New York City, NY, USA.
5 Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, MD, USA
6 Howard Hughes Medical Institute, Johns Hopkins University School of Medicine, Baltimore, MD,
USA.
7 Department of Statistics and Data Science, University of California, Los Angeles, CA, USA
8 Myllia Biotechnology GmbH. Am Kanal 27, 1110 Vienna Austria.
9 Bioinformatics Interdepartmental Ph.D. Program, University of California, Los Angeles, CA, USA
10 Weill Cornell Graduate School of Medical Sciences, Weill Cornell Medicine, New York, NY, USA.
11 Department of Cell Biology, Duke University Medical Center, Durham, NC, USA
12 Department of Human Genetics, University of California, Los Angeles, CA, USA
13 Department of Biostatistics, University of California, Los Angeles, CA, USA
14 Department of Computational Medicine, University of California, Los Angeles, CA, USA

Correspondences should be addressed. wli2@childrensnational.org;

Keywords: Perturb-seq, CRISPR-based genetic perturbations, single-cell RNA-seq, computational
model

Abstract

Understanding diverse responses of individual cells to the same perturbation is central to many biological and biomedical problems. Current methods, however, do not precisely quantify the strength of perturbation responses and, more importantly, reveal new biological insights from heterogeneity in responses. Here we introduce the perturbation-response score (PS), based on constrained quadratic optimization, to quantify diverse perturbation responses at a single-cell level. Applied to single-cell transcriptomes of large-scale genetic perturbation datasets (e.g., Perturb-seq), PS outperforms existing methods for quantifying partial gene perturbation responses. In addition, PS presents two major advances. First, PS enables large-scale, single-cell-resolution dosage analysis of perturbation, without the need to titrate perturbation strength. By analyzing the dose-response patterns of over 2,000 essential genes in Perturb-seq, we identify two distinct patterns, depending on whether a moderate reduction in their expression induces strong downstream expression alterations. Second, PS identifies intrinsic and extrinsic biological determinants of perturbation responses. We demonstrate the application of PS in contexts such as T cell stimulation, latent HIV-1 expression, and pancreatic cell differentiation. Notably, PS unveiled a previously unrecognized, cell-type-specific role of coiled-coil domain containing 6 (CCDC6) in guiding liver and pancreatic lineage decisions, where CCDC6 knockouts drive the endoderm cell differentiation towards liver lineage, rather than pancreatic lineage. The PS approach provides an innovative method for dose-to-function analysis and will enable new biological discoveries from single-cell perturbation datasets.

Introduction

Perturbation is essential for understanding the functions of the mammalian genome that encodes protein-coding genes and non-coding elements (e.g., enhancers). Single-cell profiling of cells undergoing genetic, chemical, environmental or mechanical perturbations is commonly used to examine perturbation responses at the single-cell level. Recently, high-throughput approaches of perturbation have been developed using single-cell RNA-seq (scRNA-seq) readout, including multiplexing of perturbations and single-cell CRISPR screen (e.g., Perturb-seq, CROP-seq)[1–7]. This concept has been extended to study changes in single-cell chromatin accessibility[8,9], spatial transcriptomics[10] upon perturbations or perturbation combinations[11–13], and other phenomena.

Understanding how perturbations lead to different responses within cells is critical to understanding the fundamental biology behind perturbation. Technical factors, including single-cell assays used to profile the response, and the on-target/off-target effects of perturbations, are known drivers that lead to differences of single-cell profiles in the data[14–16]. In Perturb-seq experiments that use CRISPR/Cas9 for knockouts, both in-frame deletions[16] and chromosomal losses[17] contribute to different expression profiles and clustering patterns of single cells.

Perhaps more interestingly, the heterogeneity perturbation responses may be driven by underlying biological factors (**Fig. 1a**). These factors may be either cell-intrinsic (e.g., the activities of other coding-and non-coding genomic elements) or cell-extrinsic (e.g., cell states or types, environment factors), all of which define the context of perturbation response. For example, combined expressions of transcription factors (TFs) are critical for many cellular state conversions. Therefore, to properly decode the functions of these TFs via perturbation, one must consider the effect of the cell state and the activities of other companion TFs. For this reason, defining the heterogeneity of perturbation response and identifying

82 factors that contribute to these outcomes is important for understanding how cells respond to
83 perturbation.
84
Unfortunately, computational frameworks are currently lacking to decode the diverse outcomes of
perturbations. For technical factors, mixscape is the only method to detect and mitigate confounding
variations (e.g., incomplete knockouts from CRISPR/Cas9)[16]. However, its performance has not been
rigorously benchmarked, especially when partial gene functions are perturbed using techniques like
CRISPR interference (CRISPRi). More importantly, no methods have been developed to reveal new
biological insights from the heterogenous perturbation outcomes, including studying how partial gene
perturbations affect a phenotype of interest (i.e., "dosage" analysis), and discovering biological
determinants that govern differential perturbation responses.
93
Here we present a computational framework, the perturbation-response score (PS), to quantify
heterogenous perturbation outcomes in single-cell transcriptomics datasets. The PS, estimated from
constrained quadratic optimization, quantifies the strength of the perturbation outcome for a single cell.
We performed comprehensive benchmark studies that demonstrated the outstanding performance of PS
over existing methods, including simulated datasets, genome-scale Perturb-seq, and published Perturb-
seq datasets that cover various CRISPR-based technologies. More importantly, PS analysis presents two
major conceptual advances for analyzing single-cell perturbation data: it enables analysis of the dose of
perturbation, and identification of novel biological determinants that govern the heterogeneity of
perturbation responses. First, we analyzed essential gene Perturb-seq and found two patterns of dose
response, based on whether moderate perturbation leads to strong expression changes of downstream
genes. Second, we identified intrinsic and extrinsic biological factors governing critical gene functions
in latent HIV-1 expression and pancreatic/liver development. Based on PS analysis results, we identified
and confirmed a novel function of CCDC6, wherein perturbation drives duodenum cell differentiation
towards liver commitment. Collectively, PS analysis provides a powerful tool to decode heterogenous
perturbation outcomes from single-cell assays.
109
Results
Using PS to detect heterogenous perturbation outcomes within and across datasets.
Perturbing the same gene (or non-coding elements) may result in different phenotypic changes or
transcriptional outcomes (**Fig. 1a**), depending on technical factors (e.g., perturbation efficiency) and
biological factors (e.g., cell type, cell state, activities of cofactors). Unfortunately, existing methods can
detect only technical factors[16], while biological factors remain unexplored. To bridge this gap, we built a
computational framework to quantify perturbation outcomes in single-cell datasets using scRNA-seq as
readout. Corresponding assays include single-cell CRISPR screens (e.g., Perturb-seq), or simply
multiplex scRNA-seq profiling of various perturbations (e.g., sci-Plex; **Fig. 1b, c**). We define the
perturbation-response score (PS) to quantify the strength of perturbation, where PS=0 indicates no
perturbation effect (e.g., effects corresponding to unperturbed, wild-type gene functions) and PS=1
indicates the maximum perturbation effect observed within a dataset; for example, effects that
correspond to homozygous knockouts on both alleles of a gene. We utilize the expressions of multiple
downstream targets of a perturbed gene to infer the (unknown) values of PS (**Fig. 1b**). For example, if
one cell has dramatic expression changes on the known downstream target genes, then its value of PS
should be higher than cells with weak expression changes of these genes.
126

We built a computational model, based on a constrained quadratic optimization, to automatically
identify the downstream targets of perturbed genes and calculate PS (**Fig. 1c**). This model, named
"scMAGeCK-PS", is based on our previously published scMAGeCK algorithm[15] and consists of three
steps. First, scMAGeCK-PS identifies differentially expressed genes (DEGs) upon perturbation (e.g.,
perturbing the function of gene X), by comparing the transcriptome profiles between perturbed cells and
unperturbed cells. These DEGs are served as "signature" target genes of X. Second, scMAGeCK-PS
used a previously developed scMAGeCK model to estimate the average effect of perturbation on these
target genes, which can be estimated from the first step. Third, scMAGeCK-PS uses a constraint
optimization procedure to find the value of PS that minimizes the sum of mean squared errors between
predicted and measured expression changes of all downstream targets (see Methods). The constraints are
established such that any PS is non-negative for cells with X perturbed, and is exactly zero in cells
without perturbation. Such constraints can be established based on the prior information of
perturbations; for example, the expression matrix of single-guide RNAs (sgRNAs).
140
PS outperforms mixscape in quantifying partial perturbations.
mixscape[16] is currently the only method to detect and remove technical factors that affect perturbation
outcomes, especially incomplete gene knockouts that are generated from CRISPR/Cas9. However, the
performance of mixscape on partial gene perturbations has not been fully evaluated. Here we compare
PS with mixscape using multiple benchmark datasets. We first used synthetic datasets to evaluate the
performances of different methods, because finding a real scRNA-seq dataset that contains ground truth
(*i.e.*, accurate measurements of loss-of-function upon perturbation) is challenging. For synthetic data
generation, we used scDesign3[18] to simulate the single-cell transcriptomic responses upon perturbing the
50% and 100% functions of Nelfb, based on a real scRNA-seq dataset that deletes Nelfb in mouse T
cells[19] (**Supplementary Fig. S1a;** see Methods). We specified different numbers of DEGs (from 10 to
500) and simulated their expression changes upon 50% or 100% perturbations of Nelfb functions. In all
the cases, PS correctly estimated partial perturbation, where the median PSs range from 0.32-0.34 for
50% perturbation, and greater than 0.8 for 100% perturbation, respectively (**Fig. 1d-e; Supplementary
Fig. S1b-e**). In contrast, mixscape uniformly assigned the posterior probability of perturbation to 1 in all
cases, an indication that mixscape is not suited to analyze the outcome of partial gene perturbations (**Fig.
1d-e; Supplementary Fig. S1b-e)**, possibly due to the bimodal statistic model it uses, which only
considers 100% knockout effects[16].
158
We next evaluated different methods using real single-cell perturbation datasets. We chose CRISPRi-
based Perturb-seq datasets (**Fig. 1f-i**) because the CRISPRi system directly modulates the expression
levels of perturbed genes, and the perturbation efficiency can be accessed using single-cell
transcriptomic data. We use two published K562 CROP-seq datasets:[20] in the first, only 1 gRNA is
expressed within each cell (*i.e.,* low multiplicity of infection or MOI), and in the second multiple
gRNAs are expressed (*i.e.,* high MOI). We examine cells where the transcription starting sites (TSS) of
highly expressed protein-coding genes are targeted (23 and 342 genes, respectively; **Fig. 1g-i,
Supplementary Fig. S1f-g**). If the TSS of gene X is perturbed, we first removed the expression of X
from expression matrix, and used the rest of gene expressions to measure the perturbation efficiency of
X. The scores of different methods were then compared with the expression of X, producing a direct
measurement of perturbation efficiency (**Fig. 1f**). In over 40% of these genes (10 out of 23 for low MOI,
139 out of 342 for high MOI), PS has a strong negative correlation with the expression of X (**Fig. 1g**),
defined as Pearson correlation coefficient < -0.1 and p value <0.01. In contrast, mixscape scores
correlate with X expression in none of the genes (for low MOI dataset; **Fig. 1g),** or in less than 5% of all

the genes (for high MOI dataset; **Supplementary Fig. S1f-g**). PS detects a much greater number of cells
that have a strong perturbation effect (PS or mixscape score >0.5; **Fig. 1h**), whose scores are strongly
negatively correlated with gene expression (**Fig. 1i**). We also tested both methods in another CRISPRi-
based Perturb-seq dataset, where sgRNAs with mismatches reduce efficiency, leading to partial
perturbation effects[21] (**Supplementary Fig. S1h-i**). PS has a high sensitivity and a good balance
between sensitivity and specificity, evidenced by the higher Pearson correlation coefficients
(**Supplementary Fig. S1h**) and areas under the receiver-operating characteristic (ROC) curve (AUC)
values (**Supplementary Fig. S1i**).
181
To further benchmark methods in terms of a phenotype of interest, we designed and performed a
genome-scale CRISPRi Perturb-seq on both unstimulated and stimulated Jurkat, a T lymphocyte cell
model (**Fig. 2a**), and evaluated the performances of different methods in identifying known regulators of
T cell activation. We designed Perturb-seq library that contains sgRNAs targeting the TSS of 18,595
genes (4-6 guides per gene) and used a TAP-seq-based[22] multiplex primer panel to detect the
expressions of 374 genes with high sensitivity (see **Supplementary Table S1** and Methods). We
obtained high-quality scRNA-seq data on over 586,000 single cells after quality control, and the UMAP
clustering of Perturb-seq datasets clearly demonstrated the differences between stimulated and non-
stimulated cells (**Fig. 2b**). Next, we ran PS or mixscape to calculate the scores of all perturbations at a
single-cell level; and for each perturbed gene, we calculated its overall perturbation-response score, by
adding the scores of all cells that express a corresponding sgRNA targeting that gene. Because our
system focuses on T cell stimulation, perturbing a gene that reaches a highest (and lowest) cumulative
score should have the strongest (and no) effect on T cell stimulation, respectively. For an independent
evaluation, we extracted 385 (and 1297) positive (and negative) hits whose perturbation impairs (or does
not impair) the stimulation of T cells from a published genome-scale CRISPR screen[23]. Both Perturb-seq
and pooled CRISPR screen identified many known positive regulators of T cell activation, such as
components of the T cell receptor complex (e.g., CD3D) and proximal signaling components (e.g., LCK;
Fig. 2c). For many positive genes, cells with higher values of PS or mixscape score are skewed towards
non-stimulating state, consistent with their negative selections in pooled CRISPR screens using T cell
stimulation as readout (**Fig. 2c; Supplementary Fig. S2**). However, when comparing the ROC score,
PS reaches a higher AUC score than mixscape (**Fig. 2d**), indicating its better performance in accurately
separating positive from negative hits.
204
Finally, we tested different methods on a published ECCITE-seq, which simultaneously measures
single-cell transcriptomes, surface proteins, and perturbations[16]. PDL1 protein expression was used as an
independent metric of evaluation (**Fig. 2e**), because PDL1 is a well-studied gene whose protein
expression is well understood. Among 25 perturbed genes in the ECCITE-seq perturbation library, 17
are known to regulate PDL1 expression (**Fig. 2f**). We compared PS with mixscape in terms of predicting
changes in PDL1 expression (**Fig. 2f; Supplementary Fig. S3**). In addition, the expression of the
perturbed gene is included in the comparison as a naïve method. In 19 out of 25 genes (76%), PS
outperformed mixscape and perturbed gene expression in predicting PDL1 expression (**Fig. 2e**),
including 12 out of 17 (71%) known PDL1 regulators. Notably, for genes whose perturbations led to
strong transcriptomic changes (*e.g.,* IFNGR1, IFNGR2, JAK2, STAT1), both PS and mixscape work
well, reaching AUC > 0.8 (**Fig. 2f**). For other genes whose perturbation only leads to moderate or weak
expression changes, as described previously[16], PS outperforms mixscape, including those that are
confirmed to be PLD1 regulators (*i.e.*, genes marked in red in **Fig. 2f**).
218

Analyzing dose-dependent effects of perturbation.

Traditionally, dosage analysis requires a careful, time-consuming adjustment of perturbation strength, including changing drug concentrations or designing sgRNA sequences to achieve various editing efficiencies[21,24]. Since the quantifying partial gene perturbation by PS is highly accurate (**Fig. 1-2**), we can use PS to perform dose-response analysis of perturbation, without the need to titrate the strength of perturbation. By examining ECCITE-seq data in which PDL1 expression was measured directly (**Fig. 2e**), we found correlations between PDL1 expression and the PS of known PDL1 regulators (**Fig. 3a**). The PSs of positive PDL1 regulators (e.g., IFNGR1/2, STAT1; **Fig. 3b**) are negatively correlated with PDL1 expression, while the scores of negative regulators (e.g., CUL3, BRD4) are negatively correlated (**Supplementary Fig. S3; Fig. 3c**). One example is CUL3, which is known to destabilize and degrade PDL1 protein expression[25]. Consequently, higher CUL3 PSs, indicating higher CUL3 functional perturbation, correspond to higher PDL1 protein expressions (**Fig. 3a**). Compared with mixscape, PS more accurately predicts the quantitative changes in PDL1 expression, evidenced by stronger Pearson correlations between the two (**Supplementary Fig. S3**).

We further investigated the relationships between perturbation efficiency and the strength of perturbation responses, which is measured by PS (**Fig. 3d**). In particular, we are interested in genes that show one of two different patterns of PSs upon perturbation: "buffered" distribution, where genes have high PSs only when stronger perturbation efficiency is achieved; and "sensitive" distribution, where the PSs are high, even with moderate or weak perturbation efficiency. Both "buffered" or "sensitive" terms have been coined previously to describe the effects of transcription factor dosages to chromatin accessibility[26]. CRISPRi-based Perturb-seq datasets are used, as the efficiencies of CRISPR inhibition can be directly evaluated by examining perturbed gene expressions (**Fig. 2e**).

We calculated PS for every gene in a published essential-wide Perturb-seq[27], which uses CRISPRi to inhibit the expressions of 2,285 common essential genes. We classified genes based on their PS quantiles that correspond to around 50% perturbation efficiency (**Fig. 3e**): a gene is classified as "buffered" if its median PS is smaller than 0.1; or "sensitive", if its 95% quantile is greater than 0.75. Among over 2,000 essential genes, we classified 613 genes as either buffered or sensitive. The majority are buffered (529 out of 613), indicating high robustness to perturbation, possibly due to their essential roles in cellular functions that require compensations on expression reductions. Many buffered genes belong to essential protein complexes, including proteosomes (e.g., PSMA3; **Fig. 3f**) and ribosomal subunits (e.g., RPL4; **Supplementary Fig. S4a**). 30% of the genes (185 out of 613) belong to "sensitive" category, showing strong transcriptome responses even with moderate or weak efficiencies upon perturbing gene expression (**Supplementary Fig. S4b-c**). Many of the sensitive genes are also displaying buffering effect, a demonstration of complex, heterogenous responses of cells undergoing the same perturbation of essential genes. Notably, 50% reduction of HSPA5 and GATA1 expression achieved near-maximal transcriptional response (and the associated growth defect), as in previous studies[22].

We further examined possible mechanisms by which buffered genes resist perturbation, especially those that belong to the same functional protein complex. Interestingly, perturbing one member of the protein complex usually leads to the expression up-regulation of other members of the complex, indicating a possible mechanism for compensation. For example, perturbing proteosome subunits led to a strong expression reduction of the perturbed gene (*e.g.,* PSMA5; blue squares in **Fig. 3g**) and concurrent up-regulation of other members of the proteosomes (*e.g.,* PSMB7, PSMD2). Perturbing many other protein

complexes, including ribosomal subunit, mediator, and RNA polymerases, also leads to similar up-
regulation of some members of the same functional unit (**Supplementary Fig. S5a-c**), indicating that
compensation occurs by up-regulation of other submits of the same molecular machine. To confirm our
findings on a different cellular system, we examined the effects of perturbing proteosomes in our
genome-scale Perturb-seq dataset (**Fig. 2a**). The TAP-seq approach used in this dataset provides a
sensitive and accurate measurement of gene expression changes upon perturbation[27]. Indeed, perturbing
members of the proteosome subunits leads to the up-regulation of other proteosomes (**Supplementary**
Fig. S5d), consistent with the known transcriptional feedback loop that is observed between proteosome
genes[28]. Overall, the widespread existence of such compensatory effect may explain the perturbation-
expression phenotype of buffered genes, where a strong perturbation efficiency is needed to achieve
strong expression changes.
276
PS reveals intrinsic and extrinsic biological factors that regulate gene functions in latent HIV
expression.
We next perform Perturb-seq experiment and use PS to investigate the functions of key genes regulating
latent HIV-1 expression. We used a Jurkat HIV cell model that we previously established for pooled
CRISPR screening[29], where cells stably express Cas9 and are latently infected with HIV-GFP viral
vector. We designed a Perturb-seq library that targets 10 protein-coding genes (**Supplementary Table**
S2), which are either (1) known factors in HIV-1 virus expression and T cell activation (e.g., BIRC2), or
(2) top hits from genome-scale CRISPR screens that we previously performed (e.g., BRD4)[29]. We
performed Perturb-seq experiments in three different conditions, including stimulated Jurkat (by PMA/I)
followed by GFP expression sorting (GFP+ or GFP-), and unstimulated cells (**Fig. 4a**). The single-cell
transcriptomes were profiled via the 10X Genomics Chromium platform, and expressed guide RNAs can
be captured directly. After quality controls, we received 7,063-8,811 single cells per sample, where the
mean reads per cell (and median genes expressed per cell) in each sample is at least 69,888 (and 4,744),
respectively (**Supplementary Fig. S6a**). Guide RNAs were detected in over 96% of the cells, and over
85% of these cells could be assigned a unique guide RNA (**Supplementary Table S3**). The
transcriptome profiles of cells are primarily clustered by cell states (stimulated vs. unstimulated),
indicating that the primary sources of expression variation are coming from cell states (**Fig. 4b**).
294
We investigated gene functions using our PS framework. Among all perturbed genes, the PS of BRD4
(bromodomain containing 4) demonstrates a strong cell state-specific pattern, where a subset of cells
with BRD4 perturbation has strong BRD4 PSs (named "BRD4-PS+ cells") than other BRD4-perturbed
cells (or "BRD4-PS- cells; **Fig. 4c**). BRD4-PS+ cells overexpress genes that are involved in known
functions of BRD4[30,31] including NF-kB/TNF-alpha signaling, hypoxia and apoptosis (**Fig. 4c-d**,
Supplementary Fig. S6b-d). We examined whether the differences in BRD4 PS reflects the degree of
BRD4 functional perturbation. We first checked the expressions of BRD4 "signature" genes from
another published study[32]. Compared with BRD4-PS- cells, BRD4-PS+ cells have a much lower
expressions of these signature genes (**Supplementary Fig. S6e**), indicating a stronger functional BRD4
perturbation. In addition, BRD4 has been shown to inhibit HIV transcription and activation in many
studies, including our previous CRISPR screens[29,33], consistent with the fact that HIV-GFP is one of the
strongest up-regulated genes in BRD4-PS+ cells (**Supplementary Fig. S6f**). Furthermore, BRD4-PS+
cells have a stronger GFP expression (**Fig. 4d**) than other cells, confirming a stronger BRD4 functional
perturbation in these cells.
309

To build a quantitative perturbation-expression relationship, we recalculated BRD4 PS without using
HIV-GFP expression and examined how the scores are associated with a phenotype of interest (i.e.,
latent HIV-GFP expression) in different conditions (**Fig. 4e**). BRD4 PS correlation with HIV-GFP
expression is cell-state dependent: in stimulated T cells (PMA/I treatment), a linear, positive correlation
is observed regardless of the GFP expression. In contrast, a nonlinear relationship exists in unstimulated
T cells (DMSO), where stronger BRD4 PS (>0.5) leads to a sharp increase in HIV-GFP expression (**Fig.
4e**).
317
Another gene, cyclin T1 (CCNT1), also displays heterogeneity in PS distribution: cells with CCNT1
perturbation have a high PS distribution only in stimulated cells (**Fig. 4f**). This is different from CCNT1
gene expression or guide distribution, which do not show such pattern differences between cell states
(**Supplementary Fig. S7a**). Confirming our findings, the number of DEGs (cells with CCNT1
perturbation vs. cells expressing non-targeting guides) is over 100 in stimulated cells, but only 1 in non-
stimulated cells (adjusted p value <0.001; **Supplementary Fig. S7b**). In particular, HIV-GFP is the
strongest DEG in cells with CCNT1 perturbation, consistent with the known role of CCNT1 in
activating HIV transcription.
326
CCNT1 is a key subunit of P-TEFb (positive transcription elongation factor b)/CDK9 complex that
drives RNA transcription, including the transcription of HIV. The transcription elongation control of P-
TEFb/CDK9 is a complicated process that is regulated by multiple mechanisms, including various T cell
signaling pathways (e.g., NF-kB signaling), translation control, and epigenetic modification (reviewed in
[34]). The activities of these factors are different in different states of T cells (e.g., NF-kB;
Supplementary Fig. S7c), which may explain the differences of CCNT1 PSs. Despite the strong cell
state dependency of CCNT1 PS, PS shows weak correlation with HIV-GFP within one cell state
(**Supplementary Fig. S7d**), which is different from BRD4 PS (**Fig. 4e**).
335
To further confirm our finding that different cellular states affect the transcriptomic responses of
CCNT1 perturbation, we stimulated Jurkat cells using a different agonist (TNF-alpha). To measure the
downstream effect of CCNT1 perturbation, we sorted cells by expression of HIV-GFP, which is the
strongest down-regulated gene upon CCNT1 knockout (**Supplementary Fig. S7b**), and whose
expression is known to be regulated by CCNT1[35,36]. Indeed, with the presence of TNF-alpha, CCNT1
knockout leads to a strong reduction in HIV-GFP expression (over 50% reduction), while such reduction
is much smaller (<5% reduction) in cellular states without TNF-alpha stimulation (**Fig. 4g**).
Collectively, these results demonstrated that PS is a powerful computational framework for investigating
cofactors (cell states, other genes) that drive transcriptomic responses upon gene perturbation.
345
PS enables identification of novel cell-type dependent gene functions in regulating pancreatic cell
differentiation from multiplex single-cell transcriptomics.
Besides Perturb-seq, multiplexing cells with different perturbations are also used to measure single-cell
responses to perturbation[2,19]. A mixture of cells from different perturbations can be sequenced at the
same time, and the identity of cells can be established using various methods including cell hashing[37],
the expressions of pre-defined barcodes[38], or a combination of random barcodes[39]. We therefore tested
our PS framework on pooled single-cell transcriptomics of different perturbations to study the functions
of lineage regulators during human pancreatic differentiation. By using an established in vitro human
embryonic stem cell (hESC) pancreatic differentiation system, we generated cells corresponding to early
stage (definitive endoderm, DE) and middle stage (pancreatic progenitor, PP) pancreas development. To

test the performance of PS framework and uncover the functions of unknown regulators, we picked ten clonal hESC lines with the homozygous knockout of four genes (**Supplementary Table S4**), including two known pancreatic lineage regulators (HHEX, FOXA1) and two uncharacterized candidate regulators from previous genetic screens (OTUD5, CCDC6)[40,41]. These clones are then labelled with different LARRY (Lineage and RNA recovery) DNA barcodes[38], pooled together and differentiated into DE and PP stages using established protocols[40]. Finally, the single-cell expressions of these cells were profiled via 10X genomics Chromium platform (**Fig. 5a**). The clone information of each cell was identified from LARRY barcodes. Among 26,286 single cells that passed the quality control measurements, over 97% (25,694/26,286) of the cells had at least one barcode detected, and over 80% (20,678/25,694) were identified as singlets and retained for downstream analysis. UMAP clustering revealed different known cell types during pancreatic differentiation, based on the expression markers of known cell types (**Fig. 5b; Supplementary Fig. S8**), including DE, PP, liver/duodenum progenitor (LV/DUO), endocrine precursor (EP), and cells in transition stages (e.g., DE in transition, PP in transition).

We next applied the PS framework to the pooled single-cell RNA-seq datasets containing different knockout clones. Among all knockout genes, HHEX PS is high in cells whose type is between two different differentiated cell types (PP and LP/DP; **Fig. 5c; Supplementary Fig. S8**), consistent with the known function of HHEX as a key determinant of cell fate decision, whose deletion drives DE cell differentiation towards LP/DP, rather than PP[40]. Indeed, *HHEX* knockout led to a much fewer percentage of cells that are annotated as PP (**Fig. 5d**). The PS of FOXA1, another key transcription factor during PP differentiation, is strong in DE and PP cell types, consistent with the specific expression pattern of FOXA1 in DE/PP cell types (**Supplementary Fig. S9a-c).**

As in our previous genome-wide CRISPR screens, CCDC6 is one of the top hits whose perturbation hinders PP differentiation[40,42]. However, the exact function of CCDC6 during pancreatic differentiation is largely unknown. CCDC6 may have different functions at different cell types, evidenced by the few overlaps of DEGs between different cell types (**Supplementary Fig. S9d-f**). To investigate these different functions, we calculated PSs from the DEGs from four major cell types in the dataset (DE in transition, DE, PP/PP in transition, and LV/DUO). An unbiased clustering on these CCDC6 PSs demonstrated two distinct distributions across cell types (**Fig. 5e**), where scores calculated from late-stage cell types including PP/PP in transition/LV/DUO ("pattern 1") are distributed differently from scores calculated from early-stage cell types including DE in transition/DE ("pattern 2"; **Fig. 5f; Supplementary Fig. S10a-b**), implying different behaviors of CCDC6 perturbation at different cell types. Indeed, functional analysis on DEGs leading to both patterns have distinct enrichment terms. In early-stage cell types, DEG genes are enriched in the targets of stem cell transcription factors (e.g., SOX2, POU5F1, NANOG) and cell cycle regulation (**Supplementary Fig. S10c-e**), consistent with the known function of *CCDC6* as a cell cycle regulator[43,44]. In contrast, DEGs in late-stage cell types are primarily the targets of HNF4A, a key transcription factor that drives LP/DP differentiation (**Fig. 5g; Supplementary Fig. S10f**). The expressions of these transcription factors (SOX2, HNF4A) are among the up-regulated genes in both programs, respectively (**Supplementary Fig. S9d-e**). Furthermore, compared with wild-type cells, *CCDC6* knockout cells have a much lower percentage of PP cells and a higher percentage of LP/DP cells (**Fig. 5h**). Collectively, these results imply that *CCDC6* has different functions for early vs. late-stage cell types. Especially in late-stage cell types, *CCDC6* knockout drives cell differentiation towards LV/DUO cell types rather than PP cell types.

To further validate the prediction results of CCDC6, we performed flow cytometry analysis to evaluate
the effects of *CCDC6* knockout on the composition of late-stage cell types (PP/LV/DUO). We examined
the percentage of HNF4A+ cells, a marker for LV population, and PDX1+ cells, a marker for PP
population. Indeed, both clones of *CCDC6* knockout greatly reduced PDX1+ population and increased
HNF4A+ population in three biological replicates (**Fig. 5i; Supplementary Fig. S11**), confirming our
finding on the enrichment of CCDC6 PS in LP/DP populations (**Fig. 5f-g**).
407
Discussion
Understanding cellular responses to perturbations is a central task in modern biology, from studying
tumor heterogeneity to developing personalized medicine. These perturbations may be genetic (e.g.,
knocking out genes or non-coding elements), chemical (e.g., drug treatments), mechanical (e.g.,
pressure) or environmental (e.g., temperature changes). Single-cell genomics profiles of perturbations
are commonly used to investigate the mechanisms of perturbations. Many technologies, including
Perturb-seq and sci-Plex, provide a high-content readout of the results of systematically perturbing many
genes or non-coding elements. Despite rapid technological advancements, a major bottleneck is the lack
of a computational model to fully unlock the potential of high-content perturbation, especially for
discovering novel biological insights from the data. Here we introduce the PS framework to model the
heterogenous transcriptomic responses of perturbations and to enable novel biological discovery from
modeling perturbation heterogeneity.
420
Partial gene perturbation is common in perturbation experiments. Partial perturbations may come from
dose-controlled drug treatment, gene editing technology that does not fully knockout gene function (e.g.,
RNA- or CRISPR-interference, epigenome editors), or from CRISPR/Cas9 that generates random DNA
editing outcomes. We demonstrated the outstanding performance of our PS method over existing
methods in quantifying partial gene perturbation. Specifically, partial perturbation identification enables
the analysis of dose-dependent effect, which is demonstrated in this study using various datasets.
427
More importantly, PS enables novel biological investigations, including analysis of perturbation dosage
without the need to titrate perturbation strength and identification of cell-intrinsic and extrinsic
biological factors that regulate perturbation responses. In the latter case, the PS, ranging between 0 and
1, no longer represents the quantity of partial perturbation, but instead represents the strength of the
perturbation outcome. Therefore, PS becomes a convenient tool to identify cell context that determines
perturbation outcome. We demonstrated the application of PS in various biological problems, including
T cell activation, essential gene function, latent HIV-1 virus expression, and pancreatic cell
differentiation. Importantly, our PS model leads the discovery of novel CCDC6 functions that are cell
type dependent, whose role as a regulator during pancreatic and liver cell fate decision is experimentally
validated.
438
Partial perturbations of gene functions contribute to the complexity of many biological processes. For
example, "haploinsufficient" genes are able to cause disease phenotypes when 50% of their functions
are disrupted, while "haplosufficient" genes will require a nearly complete gene knockout. However, we
currently lack a method to investigate the phenotypes of partial gene perturbations or to efficiently
perform dosage analysis at a large scale. Current approaches, such as introducing mismatches to guide
RNAs to modulate the effects of CRISPRi[21] or Cas13[28], require a complex design of a specific CRISPR
system. Here we demonstrated that both CRISPR knockout (e.g., **Fig. 2f, Fig. 4e**) and CRISPRi
naturally introduce partial perturbation effects, which can be used to study the dose effect of partial gene

perturbations on downstream gene expressions or a phenotype of interest. Our PS framework is
versatile, enabling the dosage analysis using various perturbation methods (e.g., CRISPRi or CRISPR
knockout) and assays (e.g., Perturb-seq or multiplex scRNA-seq).
450
Results from genetic perturbations (e.g., via CRISPR/Cas9) are informative for drug development, and
confirmations from genetic perturbation experiments are usually required to demonstrate the feasibility
of candidate drug targets. However, titrating pharmaceutical interventions are easy (e.g., by using
different doses of drugs), while it is much more difficult to precisely control the degree of genetic
perturbations. Our PS framework provides a convenient alternative to dose-dependent perturbations,
especially genetic perturbations, and their associations with phenotypic changes, which will be
informative in designing drugs. For example, BRD4 is the primary target of bromodomain inhibitors
(BETi), many of which have been proposed as candidates of latency reversing agents (LRAs) to
reactivate latent HIV-1 expression. The distribution of BRD4 PSs (**Fig. 3**) reveals that stronger
perturbation effects are needed to induce the desired phenotype, in this case, the expression of HIV-GFP
(**Fig. 3**). Since BRD4 is an essential gene, a strong BRD4 perturbation may lead to unexpected toxicity,
thereby limiting the efficacy of BETi. Indeed, our previous study[29] demonstrated that 10-1000x higher
doses of JQ1, a commonly used BETi, are needed to induce latent HIV-1 expression at a similar level
with other potent LRAs. Our results further warrant the development of synergistic drug combinations to
mitigate the narrow therapeutic window of BETi, which is currently tested in many studies.
466
Our PS analysis provides a general framework to analyze several major sources that contribute to the
heterogeneity of perturbation responses: the strength of perturbation *per se* (e.g., **Fig. 1i, 3d;** BRD4 in
Fig. 4c), compensations to perturbation especially on essential genes (e.g., proteosomes; **Fig. 3g**), and
cell type/state specificity (e.g., T cell states in **Fig. 4**; differentiation cell types in **Fig. 5**). Importantly,
cell type/state is linked to perturbation responses in three distinct ways: cell type/state may change as a
result of perturbation (e.g., CCDC6 and HHEX in Fig. 5); cell type/state serving a critical context to
define perturbation responses (e.g., T cell states in response to CCNT1 perturbation in **Fig. 4f-g**); and
cell type/state as a confounding factor that drives perturbation responses (e.g., BRD4 perturbation
heterogeneity in unstimulated T cells in **Fig. 4c**). Compared with other methods, PS is currently the only
method to analyze heterogeneity of perturbation responses from all these aspects.
477
Confounding factors are the major sources of variation when analyzing single-cell perturbation effects.
These confounding factors can be modeled explicitly (e.g., using generalized linear models) if
confounding source is known; or be detected and corrected using mathematical or statistical approaches
including matrix factorization (e.g., using GSFA[45]) or independent component analysis (e.g., using
CINEMA-OT[46]). In contrast, PS does not explicitly model confounding factors. Instead, PS scores can
be used in combination with methods that remove confounding sources of variation, or to detect these
confounding factors that contribute to the heterogeneity in perturbation responses (e.g., **Fig. 4c**).
Importantly, many confounding factors defined in previous methods[16,46] are not always confounding;
instead, they can be used to discover novel biological insights, as are shown in this study (e.g.,
perturbation efficiency, cell type/state). The orthogonal algorithmic design of PS compared with existing
methods also allows the combination of PS with these methods to simultaneously remove confounding
factors and measure the strength of perturbation responses.
490
One limitation of PS is its power in detecting drastic changes in cell types or states. For example, even
moderate perturbations on essential gene functions affect cellular viability[47,48]. In this case, single-cell

profiling only captures surviving cells that are resistant to essential gene perturbations in various
mechanisms (e.g., expression compensation in **Fig. 3**), and largely misses dead cells due to essential
gene dysfunction. Consequently, due to this "survival bias", PS probably only reflects the perturbation
responses in a fraction of cells, rather than the full spectrum of perturbations. To overcome this
limitation, PS can combine with recently developed prediction methods that predict the responses of
perturbations, even if cells between perturbed/non-perturbed states are unevenly distributed[49].
499
500 **Methods**
501
502 **The Perturbation-response Score (PS) framework**
503 Estimating PS proceeds in three steps, as illustrated in Figure 1c: target gene identification (Step 1),
504 average perturbation effect estimation using a previously published scMAGeCK (Step 2), and PS
505 estimation using constrained optimization (Step 3).
506
Step 1: target gene identification. We first performed differential expression analysis between cells with
certain perturbation (e.g., knocking out gene X) and negative control cells. In most cases, negative
control cells are cells that express non-targeting guide RNAs (in Perturb-seq), or wild-type cells (in
pooled scRNA-seq). In Perturb-seq with high MOI condition, these cells may come from cells that do
not have a particular perturbation. We used Wilcoxon rank sum test (implemented in Seurat) to identify
and rank differentially expressed genes. Top genes were then selected as potential target genes of the
specific perturbation. The maximum and minimum numbers of top genes can be specified by the user.
Alternatively, users can provide the list of target genes for each perturbation, based on prior knowledge,
therefore skipping the differential expression analysis in this step.
516
Step 2: average perturbation effect estimation. We used the linear regression module in scMAGeCK
(scMAGeCK-LR) to estimate the average perturbation effect. scMAGeCK-LR takes the expressions of
all target genes (identified in Step 1) in all cells as input and outputs a β score, which is conceptually
similar to log fold change. There are two advantages of using β score, instead of simply using the log
fold changes in Step 1. First, scMAGeCK-LR naturally supports datasets from high MOI Perturb-seq,
where one cell may express multiple guides targeting different genes. Second, scMAGeCK-LR is able to
estimate average perturbation effects of multiple perturbations (e.g., genome-scale perturbations) in one
step, while a naïve DEG analysis can only calculate LFC for each perturbation.
525
The mathematical model of scMAGeCK-LR is described as follows. Let Y be the log-transformed, $M*N$
expression matrix of M single cells and N target genes. These genes are the union of all target genes for
all K perturbations, extracted from Step 1. Let D be the $M*K$ binary cell identity matrix of M single cells
and K perturbations, where $d_{jX} = 1$ if single cell j contains sgRNAs targeting gene X ($j =$
$1,2,...,M; X = 1,2,...,K$), and $d_{jX} = 0$ otherwise. D can be obtained from the detected guide RNA
expression matrix from Perturb-seq or from the prior sample information from pooled scRNA-seq. The
effect of target gene knockout on all expressed genes is indicated as a β score in a matrix B with size
$K*N$, where $\beta_{XA} > 0$ (< 0) indicates gene X is positively (or negatively) selected on gene A expression,
respectively. In other words, gene X knockout increases (or decreases) gene A expression if $\beta_{XA} > 0$ ($<$
0), respectively.
536
537 The log-transformed expression matrix Y is modeled as follows:
538

$$Y = Y_0 + D \times B + \epsilon, \qquad\qquad \text{Eq (1)}$$

where Y_0 is the basal expression level of all genes in an unperturbed state, and ϵ is a noise term following a Gaussian distribution with zero means. Y_0 can be estimated from negative control cells (e.g., wild-type cells or cells expressing non-targeting guides), or be modeled using the expressions of neighboring negative control cells (e.g., the approach used by mixscape[16]). The value of B can be estimated using ridge regression:

$$B = (D^T D + \lambda I)^{-1} D^T Y, \qquad\qquad \text{Eq (2)}$$

where I is the identity matrix, and λ is a small positive value (default 0.01).

Step 3: PS estimation using constrained optimization. We revise Eq (1) to incorporate PS. Here, the log-transformed expression matrix Y is modelled as follows:

$$Y = Y_0 + \Psi \times B + \epsilon, \qquad\qquad \text{Eq (3)}$$

Where Ψ is the non-negative, raw PS matrix with the same size as D in Step 2 (*M*K)*. Each element ψ_{jx} in Ψ indicates the raw PS of cell j of perturbing gene X. Here, B is the β score matrix which is estimated in Step 2. We find the value of Ψ to minimize the squared error of predicted and observed expressions of all genes within all cells, subject to constraints and regularization terms:

$$\min \sum\nolimits_{ji}\left(y_{ji} - y_{ji}^0 - \sum\nolimits_k \psi_{jk}\beta_{ki}\right)^2 + \lambda \sum\nolimits_{jk} |\psi_{jk}|, \qquad\qquad \text{Eq (4)}$$

subject to the following constraints:

$$\begin{cases} 0 \le \psi_{jk} \le U, & if\ d_{jk} = 1 \\ \quad \psi_{jk} = 0 & if\ d_{jk} = 0 \end{cases}.$$

Here, U is a positive value indicating the upper bound of raw Ψ values, and d_{ik} is the value of the binary cell identity matrix in Step 2. $1 \le j \le M$ is the index of single cells, $1 \le i \le N$ is the index of target genes, and $1 \le k \le K$ is the index of perturbations.

Because we are imposing non-negative constraints to Ψ, the absolute operator can be removed from the objective function in Eq (4) and can be rewritten as

$$\min \sum\nolimits_{ji}\left(y_{ji} - y_{ji}^0 - \sum\nolimits_k \psi_{jk}\beta_{ki}\right)^2 + \lambda \sum\nolimits_{jk} \psi_{jk}. \qquad\qquad \text{Eq (4)}$$

This becomes a constrained quadratic optimization problem where the best solution can be easily achieved using methods like Newton's method. The final, normalized PS is to scale values of ψ_{ik} to [0,1]:

$$PS_{ik} = \psi_{ik}/U.$$

We implemented this framework as part of the scMAGeCK pipeline[18]. The PS source code, documentation and tutorials can be found on Github: https://github.com/davidliwei/PS

Simulated datasets

The eight simulated datasets are generated by the simulator scDesign3[46] with modifications for Perturb-
seq. The simulation utilizes scDesign3's parametric model to capture the characteristics of the user-
inputted reference data, specify the desired ground truth, and simulate synthetic cells via sampling from
the model (to be detailed in **Steps 1-4** below). The reference data is the real scRNA-seq dataset with the
gene Nelfb perturbed in some mouse T cells[47]; the cells with Nelfb perturbed are referred to as *knockout*
cells, and the cells with Nelfb unperturbed serve as the negative control and are referred to as *wild-type*
cells. Based on the same reference data, the eight simulated datasets are generated under eight different
settings. Each setting corresponds to a combination of two simulation parameters' values: the number of
Nelfb's downstream genes (i.e., the genes whose expression levels are affected by Nelfb's knockout;
with candidate values 0, 10, 200, and 500) and the perturbation efficiency (with candidate values 50%
and 100%). The candidate downstream genes of Nelfb are the top differentially expressed (DE) genes
identified from the bulk RNA-seq data of the same biological sample (from the second sheet in the
Excel file from Wu et al.'s Supplementary Data 1[48]). Thus, we have 4×2=8 simulated datasets in total.
595
Before running the simulation, we pre-process the scRNA-seq dataset and the bulk DE gene rank list.
1. First, we perform the same quality control as in the dataset's original publication[49]. Specifically,
cells are retained only if their numbers of detected genes are between 1,000 and 5,000, and their
UMI counts have less than 12% mitochondrial counts.
2. Second, we impute and amplify the gene-by-cell count matrix of the wild-type mouse cells to
enhance the perturbation effects in the simulated data. Specifically, we first impute the wild-type
count matrix using scImpute[50] (default version 0.0.9) to reduce the sparsity. Then we multiply
the imputed count matrix by an amplification factor of 10 to increase the range of gene
expression levels.
3. Third, we construct a gene-by-cell count matrix by combining the wild-type cells in the post-
imputation-and-amplification wild-type count matrix and the knockout cells in the knockout
count matrix. By the end of this step, the dimension of this combined matrix is $(P+1)\times N$, with
rows corresponding to $P+1$ genes (Nelfb and P other genes) and columns corresponding to N
cells, which consist of N^{wt} wild-type cells and N^{ko} knockout cells.
4. Fourth, we extract the row corresponding to Nelfb as a vector, which contains Nelfb's counts in
all cells (an N-dimensional vector denoted as C, where C_j is Nelfb's count in cell j), and we
denote the remaining P rows as a $P\times N$ matrix $\mathbf{Y}$, where Y_{ij} is gene i's count in cell j.
5. Fifth, using $\mathbf{Y}$, we refine the list of bulk DE genes by excluding the DE genes that correspond to
zero rows in $\mathbf{Y}$ or do not correspond to any rows in $\mathbf{Y}$.
6. Lastly, to reduce the computation time for data simulation, we use the scran package[51] to select
3,000 highly variable genes in $\mathbf{Y}$. We only keep the union of these 3,000 highly variable genes
and the refined bulk DE genes as the rows in $\mathbf{Y}$. The number of the kept genes is 3,390, so the
dimension of $\mathbf{Y}$ is 3,390×N.
619
Additionally, we know which cells have Nelfb perturbed; thus, we have another N-dimensional binary
vector denoted as K, where K_j indicates whether the j-th cell has Nelfb perturbed or not; that is, $K_j = 0$
means the j-th cell is a wild-type cell, and $K_j = 1$ means the j-th cell is a knockout cell. K and C are used
as two covariate vectors, and $\mathbf{Y}$ is used as the reference count matrix for scDesign3. Finally, we modify
scDesign3 by using $\mathbf{Y}$, C, K, the refined DE genes, the number of Nelfb's downstream genes, and the
perturbation efficiency to simulate data in the following four steps:
626

<u>Step 1: modeling each gene's marginal distribution independently.</u> For each gene i, if it is a downstream
gene of Nelfb, we assume that Y_{ij}, conditional on C_j, follows a zero-inflated negative binomial (ZINB)
distribution with the mean parameter μ_{ij}, the dispersion parameter ϕ_i, and the zero-inflation probability
parameter ν_{ij}. Otherwise, if gene i is not a downstream gene of Nelfb, we assume that Y_{ij} follows a
ZINB distribution with the mean parameter μ_i, the dispersion parameter ϕ_i, and the zero-inflation
probability parameter ν_i. This marginal distribution for each gene is specified by a generalized additive
model for location, scale, and shape (GAMLSS). Without loss of generality, we define the first D genes
in $\mathbf{Y}$ to be the top D DE genes in the refined DE gene list ($D \in \{0, 10, 200, 500\}$); we treat these top D
DE genes as the D downstream genes of Nelfb. Then we modify scDesign3's original code
implementation so Nelfb's downstream genes and non-downstream genes have different marginal
distributions: a downstream gene's marginal distribution in each cell j depends on C_j, Nelfb's count in
cell j; a non-downstream gene's marginal distribution in each cell j is irrelevant to C_j.
639
640 For Nelfb's downstream gene $i = 1, \dots, D$:

$$
\begin{cases}
Y_{ij} \mid C_j \sim \text{ZINB}(\mu_{ij}, \phi_i, \nu_{ij}) \\
\log(\mu_{ij}) = \alpha_i + \beta_i \times C_j \\
\log(\phi_i) = \omega_i \\
\text{logit}(\nu_{ij}) = \gamma_i + \eta_i \times C_j
\end{cases}
.
$$

641
642
643 For Nelfb's non-downstream gene $i = D + 1, \dots, P$:

$$
\begin{cases}
Y_{ij} \sim \text{ZINB}(\mu_i, \phi_i, \nu_i) \\
\log(\mu_i) = \alpha_i \\
\log(\phi_i) = \omega_i \\
\text{logit}(\nu_i) = \gamma_i
\end{cases}
.
$$

644
645
646 After parameter estimation by the R package gamlss, the fitted distribution of $Y_{ij} \mid C_j$, for $i = 1, \dots, D$, is
647 denoted as $\text{ZINB}(\hat{\mu}_{ij}, \hat{\phi}_i, \hat{\nu}_{ij})$ with the CDF $\hat{F}_{ij}$; the fitted distribution of Y_{ij}, for $i = D + 1, \dots, P$, is
648 denoted as $\text{ZINB}(\hat{\mu}_i, \hat{\phi}_i, \hat{\nu}_i)$ with the CDF $\hat{F}_i$. The other parameters including $\alpha_i, \beta_i, \gamma_i$, and η_i are
649 estimated as $\hat{\alpha}_i, \hat{\beta}_i, \hat{\gamma}_i$, and $\hat{\eta}_i$ for each i respectively.
650
<u>Step 2: modeling genes' joint distribution using the Gaussian copula.</u> To approximate the pairwise gene-
gene correlations in the reference dataset, scDesign3 utilizes a multivariate statistical technique, the
Gaussian copula. Given each gene's marginal distribution fitted in Step 1, scDesign3 approximates the
multivariate joint distribution of the P genes in cell j as

$\left(\Phi^{-1}(\hat{F}_{1j}(Y_{1j})), \dots, \Phi^{-1}(\hat{F}_{Dj}(Y_{Dj})), \Phi^{-1}(\hat{F}_{D+1}(Y_{(D+1)j})), \dots, \Phi^{-1}(\hat{F}_P(Y_{Pj})) \right) \sim N\left(\mathbf{0}, \widehat{\mathbf{R}}(K_j) \right)$,

where $\Phi^{-1}(\cdot)$ denotes the inverse of the cumulative distribution function (CDF) of the standard
Gaussian distribution, $\mathbf{0}$ is the P-dimensional zero vector, and $\widehat{\mathbf{R}}(K_j)$ is the estimated $P \times P$ gene-gene
correlation matrix of the Gaussian copula conditional on the value of K_j. Specifically, since K_j is binary,
we have two estimated gene-gene correlation matrices, one for the wild-type cells ($K_j = 0$) and the other
for the knockout cells ($K_j = 1$). For $\hat{F}_{1j}(Y_{1j}), \dots, \hat{F}_{Dj}(Y_{Dj}), \hat{F}_{D+1}(Y_{(D+1)j}), \dots, \hat{F}_P(Y_{Pj})$, a technique called
"distributional transform" is used to make the CDFs continuous; see Sun et al.[52] for a detailed
explanation.
663

<u>Step 3: modifying the fitted parameters.</u> Since we want to generate synthetic datasets with two
perturbation efficiencies, for each downstream gene $i = 1, \dots, D$, we modify the mean parameters for all
downstream genes in the knockout cells to reflect the user-specified perturbation efficiency. Without
loss of generality, we assume the first $N^{\text{ko}} = \sum_{j=1}^{N} I(K_j = 1)$ of the N cells as the knockout cells. Then,
we update the mean parameters for Nelfb's D downstream genes in the N^{ko} knockout cells (i.e., $\hat{\mu}_{ij}$ for
$i \in \{1, \dots, D\}, j \in \{1, \dots, N^{\text{ko}}\}$) based on the user-specified perturbation efficiency as follows.
670
For the 50% perturbation efficiency: We randomly sample N^{ko} values from $\{C_j, j \in \{N^{\text{ko}} + 1, \dots, N\}\}$
(i.e., Nelfb's counts in the wild-type cells) and multiply the sampled C_j values by 0.5 to represent the
50% perturbation efficiency. We store these sampled and scaled values by $C^* = \left(C_1^*, \dots, C_{N^{\text{ko}}}^*\right)^T$ as
Nelfb's counts in the N^{ko} synthetic knockout cells to be simulated. Then, we modify the mean
parameters for the D downstream genes in the N^{ko} synthetic knockout cells (for $i \in \{1, \dots, D\}, j \in$
$\{1, \dots, N^{\text{ko}}\}$) as

677
$$\hat{\mu}_{ij} = \hat{\alpha}_i + \hat{\beta}_i \cdot C_j^* .$$

678
679   For the 100% perturbation efficiency: C^* becomes a zero vector with length N^{ko}, and we modify $\hat{\mu}_{ij}$ for
680     $i \in \{1, \dots, D\}, j \in \{1, \dots, N^{\text{ko}}\}$ in the same way as above.
681
682   We do not change any estimated mean parameters for the D downstream genes in the N^{wt} wild-type
683   cells, any estimated mean parameters for the non-downstream $(P - D)$ genes in all N cells, any
684   estimated dispersion parameters, or any estimated zero-inflation probability parameters.
685
686   Moreover, we use S to denote an N-dimensional vector representing Nelfb's counts in the N synthetic
687   cells, with the counts in the first N^{ko} synthetic knockout cells set above based on the perturbation
688   efficiency, and the counts in the last N^{wt} synthetic wild-type cells same as those in the real N^{wt} wild-
689   type cells. That is, $S_j = C_j^*$ for $j \in \{1, \dots, N^{\text{ko}}\}$, and $S_j = C_j$ for $j \in \{N^{\text{ko}} + 1, \dots, N\}$.
690
691   <u>Step 4: generating synthetic data with the fitted model and modified parameters.</u> First, we independently
692   sample N^{wt} Gaussian vectors of length P from the estimated P-dimensional multivariate Gaussian
693   distribution $N\left(0, \hat{R}(K_j = 0)\right)$ and N^{ko} Gaussian vectors of length P from the estimated P-dimensional
694   multivariate Gaussian distribution $N\left(0, \hat{R}(K_j = 1)\right)$. Together, we stack these $N = N^{\text{wt}} + N^{\text{ko}}$ vectors
695     $(\tilde{Z}_{11}, \dots, \tilde{Z}_{P1})^T, \dots, (\tilde{Z}_{1N}, \dots, \tilde{Z}_{PN})^T$ by row into a $P{\times}N$ Gaussian matrix $\tilde{Z}$.
696
697   Given the parameter estimates (modified or not) from Step 3, we convert the $P{\times}N$ Gaussian
698   matrix $\tilde{Z}$ into a $P{\times}N$ ZINB count matrix $\tilde{Y}$ as

699
$$\tilde{Y} = \begin{bmatrix} \tilde{Y}_1 = (\hat{F}_{11}^{-1}(\tilde{Z}_{11} | \hat{\mu}_{11}, \hat{\phi}_1, \hat{v}_{11}), \dots, \hat{F}_{1N}^{-1}(\tilde{Z}_{1N} | \hat{\mu}_{1N}, \hat{\phi}_1, \hat{v}_{1N})) \\ \vdots \\ \tilde{Y}_D = (\hat{F}_D^{-1}(\tilde{Z}_{D1} | \hat{\mu}_{D1}, \hat{\phi}_D, \hat{v}_{D1}), \dots, \hat{F}_{DN}^{-1}(\tilde{Z}_{DN} | \hat{\mu}_{DN}, \hat{\phi}_D, \hat{v}_{DN})) \\ \tilde{Y}_{D+1} = (\hat{F}_{D+1}^{-1}(\tilde{Z}_{(D+1)1} | \hat{\mu}_{D+1}, \hat{\phi}_{D+1}, \hat{v}_{D+1}), \dots, \hat{F}_{D+1}^{-1}(\tilde{Z}_{(D+1)N} | \hat{\mu}_{D+1}, \hat{\phi}_{D+1}, \hat{v}_{D+1})) \\ \vdots \\ \tilde{Y}_P = (\hat{F}_P^{-1}(\tilde{Z}_{P1} | \hat{\mu}_P, \hat{\phi}_P, \hat{v}_P), \dots, \hat{F}_P^{-1}(\tilde{Z}_{PN} | \hat{\mu}_P, \hat{\phi}_P, \hat{v}_P)) \end{bmatrix} .$$

700
701 Lastly, we combine $\widetilde{Y}$ with S by row into a $(P+1) \times N$ matrix, obtaining the final $(P+1) \times N$ synthetic

702 count matrix ($\begin{bmatrix} \widetilde{Y} \\ S \end{bmatrix}$).

703
Genome-scale Perturb-seq on Jurkat cells
Perturb-seq. We performed genome-scale Perturb-seq on Jurkat E6 cell line expressing dCas9-KRAB as
our model to study. We transduced them with a genome-wide CRISPRi CROP-seq library at a high
MOI. After infection, we split the cells into two populations, including untreated cells and activated cells
(cells treated with anti-TCR and anti-CD28 antibodies for approximately 24 hours to stimulate TCR
signaling). Cells were then labelled with cell hashing antibodies. Multiple labels were used for the
activated population to help with cell multiplet detection. Cells were loaded on 16 channels of a 10x
Chromium X instrument. We loaded 115 000 cells per channel, and the expected recovery rate was 60
000 cells per channel, including 24% multiples. Samples were pooled unequally before they were loaded
on the ChromiumX: 10% untreated cells, 90% treated cells. A sequencing library was prepared using
3'Chemistry with a targeted primer panel: custom multiplex PCR step to enrich for specific transcripts.
Libraries were sequenced on NovaSeq S4 PE100 in asymetric read mode (R1: 28 cycles; R2: 172
cycles), with PhiX concentration of 1%. The expected coverage is around 9 000 ~ 10 000 input reads per
cell.

718
719 Hash oligos.

oligo	condition
CGGCTCGTGCTGCGTCGTCTCAAGTCCAGAAACTCCGTGTATCCT	untreated
CTCCCTGGTGTTCAATACCCGATGTGGTGGGCAGAATGTGGCTGG	activated
TTACCCGCAGGAAGACGTATACCCCTCGTGCCAGGCGACCAATGC	activated
TGTCTACGTCGGACCGCAAGAAGTGAGTCAGAGGCTGCACGCTGT	activated
CCCCACCAGGTTGCTTTGTCGGACGAGCCCGCACAGCGCTAGGAT	activated
GTGATCCGCGCAGGCACACATACCGACTCAGATGGGTTGTCCAGG	activated
GCAGCCGGCGTCGTACGAGGCACAGCGGAGACTAGATGAGGCCCC	activated

720
sgRNA library design. The genome-wide CRISPRi sgRNA library was designed to target the
transcription start site (TSS) coordinates, calculated from publicly available FANTOM CAGE peaks
data. In total, 18 595 genes were targeted, with 4 sgRNAs per gene. On top of that, we designed another
CRISPRi library targeting 3220 genes with 4 sgRNAs per gene. This library was designed using Jurkat-
specific TSS, which were calculated from public Jurkat CAGE-seq datasets. Both libraries were
combined into a final library targeting 3220 genes with 8 sgRNA/gene and 15 375 genes with 4
sgRNA/gene.

728
729 Targeted primer panel. The primer panel for targeted transcriptomic readout consisted of 374 target
730 genes from several categories:

Source	Number of targets (genes)
bulk RNA-seq DEG, 50 top up/downregulated	100
T-cell related genes	51
KEGG TCR signaling pathway	35
RNA-binding proteins	27

Replogle*: unfolded protein response	51
Replogle: proteasome	41
Replogle: NFkB	20
Replogle: cell cycle	29
Replogle: targets of nonsense-mediated decay	5
Controls (cell cycle, mitochondrial, Cas9)	15
Total number of targets	374

* "Replogle" prefix means that this category of targets was derived from the published genome-scale
Perturb-seq dataset[50].

<u>Data preprocessing.</u> For each of the 16 channels, all 3 kinds of sequencing libraries (mRNA, sgRNA,
cell hashing) were indexed using the same Illumina index sequence. We obtained high-quality scRNA-
seq data of over 586,000 single cells after quality control, with a median 13 guides detected per cell. We
obtained an average of 400 cells per gene perturbation. STAR and STAR solo 2.7.10a were used to map
transcriptomic reads against a custom gtf annotation, which was based on gencode.v34.annotation.gtf
(hg38). Reads that did not map to the transcriptome reference were then mapped & counted using STAR
solo against a custom fasta reference with guide sequences and a fasta reference with hash label
sequences. STAR Solo output transcriptome matrices were first filtered using an approach similar to 10x
cellranger EmptyDrops filtering, which retained cells with at least 10 % UMI count of the 99th percentile
UMI counts of the top expected cells number. Then, an initial Seurat object was created from those
filtered transcriptome matrices using the CreateSeuratObject function with following parameters:
min.cells = 5; min.features = 10; all other parameters at default values. Outlier cells were filtered out by
mithochondrial and mRNA content (percent.mt, nCount_RNA). In order to detect cell multiplets and
determine cell population (untreated or activated), cell labels (also known as hashes) were called using
the MULTI-seq approach (deMULTIplex::classifyCells in R). Only cells with exactly 1 known label
were kept. Then, sgRNA calling was conducted using a binomial test, with total sgRNA UMI counts
used to derive background frequencies. A threshold of 0.05 on Benjamini-Hochberg corrected p-values
(per channel) was use to generate the final calls. The sgRNA assays are sparse matrices containing 1,
where the respective cell is considered to be carrying the respective sgRNA and 0 elsewhere. Following
that, all results from steps above from 16 channels were merged together, and merged counts were
normalized using NormalizeData and scaled using ScaleData. Cell cycle scoring was performed via
CellCycleScoring, PCA was calculated using RunPCA, and UMAP was calculated on first 30 principal
components using RunUMAP in Seurat.

HIV latency Perturb-seq
We used a previously established cell line model of HIV latency[40]. In this model, Jurkat cells were
infected with an HIV vector with GFP tied to the LTR promoter, resulting in a positive GFP signal as a
measurement of viral transcription reactivation and HIV latency reversal. These cells, which already
express Cas9, were transduced with a lenti-sgRNA library. The lenti-sgRNA library (MilliporeSigma;
LV14, U6-gRNA-10x:EF1a-Puro-2a-BFP) was designed to target 10 genes, with 3 gRNAs per gene. In
addition to non-targeting controls, the library contained five positive regulators (NFKB1, CCNT1,
PRKCA, TLR1, MAP3K14) and five negative regulators (NFKBIA, NELFE, HDAC2, BRD4, BIRC2)
of HIV transcription. Transduction was carried out on 850,000 cells at an MOI of 0.3 using 8ug/ml
polybrene in 2 ml of RPMI containing 10% FBS and 1% penicillin–streptomycin. The media was
replaced 24 hours later with fresh media without polybrene. Two days after transduction, the cells were

selected for using 1.5 ug/ml puromycin for 5 days. After selection, the cells were split evenly into three groups. One-third of the cells were kept in culture with no drug added, and two-thirds of the cells were stimulated with PMA/I (50ng/ml PMA in combination with 1 µM Ionomycin). After 16 hours, the stimulated cells were sorted into GFP+ and GFP- populations. All three samples were then analyzed following the 10x Genomics single-cell sequencing protocol. Sequencing data, encompassing gene expression and CRISPR guide capture libraries, underwent demultiplexing and processing using Cell Ranger (version 6.1.2). The resulting feature-barcode matrices from three samples were then merged, and subsequent analysis was carried out utilizing the Seurat R package (version 4.3.1). To ensure data quality, cells were excluded if the number of expressed genes was greater than 7,500 or fewer than 200. Additionally, cells were removed if the percentage of mitochondrial reads exceeded 15%. Single cells harboring more than one detected sgRNA sequence, attributable to either multiple sgRNA transductions or the presence of multiple cells in a single-cell droplet, were also excluded from the analysis. Following quality control measures, merged counts underwent normalization and scaling. PCA was computed based on the top 2,000 highly variable genes. Subsequently, clustering and UMAP embeddings were performed using default parameters. To gain further insights into the biological significance of the obtained clusters, enrichment analysis was conducted utilizing Enrichr (PMID: 27141961).

Pancreatic differentiation clones and pooled single-cell RNA-seq

<u>Culture of hESC.</u> Generation of KO hESCs was described in published studies, including HHEX KO H1 and HUES8 cell lines[53], FOXA1 KO HUES8 cell lines[54], OTUD5 KO HUES8 cell lines, and CCDC6 KO H1 cell lines[41]. Cells were regularly confirmed to be mycoplasma-free by the Memorial Sloan Kettering Cancer Center (MSKCC) Antibody & Bioresource Core Facility. KO and WT hESCs were maintained in Essential 8 (E8) medium (Thermo Fisher Scientific, A1517001) on vitronectin (Thermo Fisher Scientific, A14700) pre-coated plates at 37 °C with 5% CO2. The Rho-associated protein kinase (ROCK) inhibitor Y-276325 (5 µM; Selleck Chemicals, S1049) was added to the E8 medium the first day after passaging or thawing of hESCs.

<u>hESC-directed pancreatic differentiation.</u> hESCs were seeded at a density of 2.3×10^5 cells/cm^2 on vitronectin-coated plates in E8 medium with 10 µM Y-27632. After 24 hours, cells were washed with PBS and differentiated to DE (stage 1), primitive gut tube (stage 2), PP1 (stage 3) and PP2 (stage 4) stages following previously described 4-stage protocol[40]. In brief, stage 1 (3 d): S1/2 medium supplemented with 100 ng ml^{-1} Activin A (Bon Opus Biosciences) and 5 µM CHIR99021 (04-0004-10, Stemgent) for 1 d. S1/2 medium supplemented with 100 ng ml^{-1}Activin A for the next 2 d. Stage 2 (2 d): S1/2 medium supplemented with 50 ng ml^{-1} KGF (AF-100-19, PeproTech) and 0.25 mM vitamin C (VitC) (Sigma-Aldrich, A4544). Stage 3 (2 d): S3/4 medium supplemented with 50 ng ml^{-1} KGF, 0.25 mM VitC and 1 µM retinoic acid (R2625, MilliporeSigma). Stage 4 (4 d): S3/4 medium supplemented with 50 ng ml^{-1} KGF, 0.1 µM retinoic acid, 200 nM LDN (Stemgent, 04-0019), 0.25 µM SANT-1 (Sigma, S4572), 0.25 mM VitC and 200 nM TPB (EMD Millipore, 565740). The base differentiation medium formulations used in each stage were as follows. S1/2 medium: 500 ml MCDB 131 (15-100-CV, Cellgro) supplemented with 2 ml 45% glucose (G7528, MilliporeSigma), 0.75 g sodium bicarbonate (S5761, MilliporeSigma), 2.5 g BSA (68700, Proliant), 5 ml GlutaMAX (35050079, Invitrogen). S3/4 medium: 500 ml MCDB 131 supplemented with 0.52 ml 45% glucose, 0.875 g sodium bicarbonate, 10 g BSA, 2.5 ml ITS-X, 5 ml GlutaMAX.

<u>Cell infection with LARRY barcode virus.</u> Individual LARRY barcode constructs were cloned from the LARRY barcode library (Addgene:140024) and transfected to 293T cells to generate lentivirus. Next,

817 each KO and WT hESC clone was infected with a unique LARRY barcode at low MOI. One week after
818 lentiviral infection, the barcoded cells, which expressed GFP, were sorted out and cultured in E8
819 medium as described in previous section.
820
<u>Pooled single-cell RNA-seq.</u> One day before differentiation, each of 10 hESC barcoded clones were
counted, mixed at the same cell number ratio, and then seeded at a density of 2.3×10^5 cells/cm^2 onto a
12-well cell culture plate. At DE and PP2 stages, pooled differentiating cells were dissociated into single
cell suspension by TrypLE Select for 5 min at 37 °C. Cells were then stored in BAMBANKER™
freezing medium for future experiments. For scRNA-seq, frozen cells were thawed and sorted to collect
live GFP+ cells. Cellular suspensions were then loaded on a Chromium Controller following the
manufacturer's instructions (10x Genomics Chromium Single Cell 3′ Reagent Kit v3.1 User Guide).
cDNA libraries and targeted LARRY barcode libraries were generated separately using 10ul cDNA
each. cDNA libraries were made under manufacturer's instructions and targeted LARRY barcode
libraries were amplified using specific primers (F: CTACACGACGCTCTTCCGATCT; R:
GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTtaaccgttgctaggagagaccataT).
832
<u>Data analysis.</u> The sequencing data which included transcriptome and LARRY barcode libraries,
underwent demultiplexing and processing via Cell Ranger (version 6.1.2). Subsequent analysis was
conducted using the Seurat R package (version 4.3.1). Quality control measures were implemented to
ensure robust data analysis. Cells were excluded if the number of expressed genes exceeded 7,000 or fell
below 200. Additionally, cells were removed if the percentage of mitochondrial read exceeded 20%.
Singlet cells were defined by considering the highest feature barcode count, ensuring it was at least
twice as large as the second highest feature barcode count. Single cells containing more than one
detected barcode sequence were excluded from the dataset. This process resulted in a final set of 20,678
cells for downstream analysis. After quality control measures, the count matrix underwent normalization
and scaling. PCA was performed using the top 2,000 highly variable genes. Subsequently, clustering and
UMAP embeddings were generated using default parameters to elucidate the underlying structure and
relationships within the dataset.
845
<u>Flow cytometry.</u> Cells were dissociated using TrypLE Select and resuspended in FACS buffer (5% FBS
in PBS). Live/Dead Fixable Violet cell stain (Invitrogen, L34955) was used to discriminate dead cells
from live cells. Permeabilization/fixation was performed at room temperature for 1 h. Antibody staining
was performed in permeabilization buffer. Antibodies for this study include HNF4A, Novus Biologicals,
NBP2-67679, 1:200; PDX1, R&D Systems, AF2419, 1:500, Donkey anti-Rabbit IgG (H+L) Highly
Cross-Adsorbed Secondary Antibody, Thermo Fisher Scientific, 1:500; Donkey anti-goat IgG (H+L)
Highly Cross-Adsorbed Secondary Antibody, Thermo Fisher Scientific, 1:500. Cells were then analysed
using BD LSRFortessa. Flow cytometry analysis and figures were generated using FlowJo v.10.
854

Acknowledgements
The authors thank all members from the Li and Huangfu laboratory for comments and discussions. The
authors thank Jake P. Taylor-King for discussions. This study is supported by NIH R01 HG010753,
HL168174 (to W.L., B.S., L.C.), District of Columbia Center for AIDS (DC-CFAR) Research
Transitioning Investigator Award (AI117970, to W.L.), and startup support from the Center for Genetic
Medicine Research at Children's National Hospital. D.H is supported by NIH UM1 HG012654, U01
HG012051. J.J. Li is supported by National Science Foundation DBI-1846216 and DMS-2113754, NIH
R35 GM140888, Johnson and Johnson WiSTEM2D Award, Sloan Research Fellowship, UCLA David

Geffen School of Medicine W.M. Keck Foundation Junior Faculty Award, and Chan-Zuckerberg
Initiative Single-Cell Biology Data Insights [Silicon Valley Community Foundation Grant Number:
2022-249355]. W.D. and R.F.S. is supported by the Howard Hughes Medical Institute.

Author contributions

W.L. conceived the project. W.L. and B.S. developed the method. W.L., B.S., and D.L. designed and
performed the experiments and analyzed the data. W.D., N.M. and B.S. performed and analyzed HIV
Perturb-seq under the supervision of J.M.S., R.S. and W.L. B.S., Q.W. and D.S. performed synthetic
experiments under the supervision of W.L. and J.J.L. D.L., D.Y., B.W. and B.R. generated pancreatic
differentiation dataset and performed validations under the supervision of D.H. A.K., A.V., N.U., and
A.L. generated and analyzed genome-scale Perturb-seq under the supervision of T.B. X.C., L.C. and
Y.D. performed the analysis and interpretation of the results. W.L., and B.S. wrote the manuscript with
input from all the authors. W.L. T.B., J.J.L., R.S. and D.H. supervised the study.

Competing interests

878   T.B. is a co-founder and Managing Director of Myllia Biotechnology. A.K., A.V., N.U. and A.L. are
879   employees of Myllia Biotechnology. Other authors declare that they have no competing interest.

Data and materials availability

882   The Perturb-seq scRNA-seq data have been deposited to Gene Expression Omnibus (GEO) under the
883   accession number GSE247601. The source code of the PS method, and the documentation and demos
884   are available on GitHub: (https://github.com/davidliwei/PS).

References

1   High-content CRISPR screening. *Nat Rev Methods Primers* 2022;**2**:.
https://doi.org/10.1038/s43586-022-00098-7.
2   Srivatsan SR, McFaline-Figueroa JL, Ramani V, Saunders L, Cao J, Packer J, *et al.* Massively
multiplex chemical transcriptomics at single-cell resolution. *Science* 2020;**367**:45–51.
3   Dixit A, Parnas O, Li B, Chen J, Fulco CP, Jerby-Arnon L, *et al.* Perturb-Seq: Dissecting
Molecular Circuits with Scalable Single-Cell RNA Profiling of Pooled Genetic Screens. *Cell*
2016;**167**:1853-1866.e17.
4   Adamson B, Norman TM, Jost M, Cho MY, Nuñez JK, Chen Y, *et al.* A Multiplexed Single-Cell
CRISPR Screening Platform Enables Systematic Dissection of the Unfolded Protein Response. *Cell*
2016;**167**:1867-1882.e21.
5   Jaitin DA, Weiner A, Yofe I, Lara-Astiaso D, Keren-Shaul H, David E, *et al.* Dissecting Immune
Circuits by Linking CRISPR-Pooled Screens with Single-Cell RNA-Seq. *Cell* 2016;**167**:1883-
1896.e15.
6   Datlinger P, Rendeiro AF, Schmidl C, Krausgruber T, Traxler P, Klughammer J, *et al.* Pooled
CRISPR screening with single-cell transcriptome readout. *Nat Methods* 2017;**14**:297–301.

7   Xie S, Duan J, Li B, Zhou P, Hon GC. Multiplexed Engineering and Analysis of Combinatorial
Enhancer Activity in Single Cells. *Mol Cell* 2017;**66**:285-299.e5.
8   Rubin AJ, Parker KR, Satpathy AT, Qi Y, Wu B, Ong AJ, *et al.* Coupled Single-Cell CRISPR
Screening and Epigenomic Profiling Reveals Causal Gene Regulatory Networks. *Cell*
2019;**176**:361-376.e17.
9   Liscovitch-Brauer N, Montalbano A, Deng J, Méndez-Mancilla A, Wessels H-H, Moss NG, *et al.*
Profiling the genetic determinants of chromatin accessibility with scalable single-cell CRISPR
screens. *Nat Biotechnol* 2021;**39**:1270–7.
10   Dhainaut M, Rose SA, Akturk G, Wroblewska A, Nielsen SR, Park ES, *et al.* Spatial CRISPR
genomics identifies regulators of the tumor microenvironment. *Cell* 2022;**185**:1223-1239.e20.
11   Norman TM, Horlbeck MA, Replogle JM, Ge AY, Xu A, Jost M, *et al.* Exploring genetic
interaction manifolds constructed from rich single-cell phenotypes. *Science* 2019;**365**:786–93.
12   Replogle JM, Norman TM, Xu A, Hussmann JA, Chen J, Cogan JZ, *et al.* Combinatorial single-cell
CRISPR screens by direct guide RNA capture and targeted sequencing. *Nat Biotechnol*
2020;**38**:954–61.
13   Wessels H-H, Méndez-Mancilla A, Hao Y, Papalexi E, Mauck WM 3rd, Lu L, *et al.* Efficient
combinatorial targeting of RNA transcripts in single cells with Cas13 RNA Perturb-seq. *Nat*
Methods 2023;**20**:86–94.
14   Hill AJ, McFaline-Figueroa JL, Starita LM, Gasperini MJ, Matreyek KA, Packer J, *et al.* On the
design of CRISPR-based single-cell molecular screens. *Nat Methods* 2018;**15**:271–4.
15   Yang L, Zhu Y, Yu H, Cheng X, Chen S, Chu Y, *et al.* scMAGeCK links genotypes with multiple
phenotypes in single-cell CRISPR screens. *Genome Biol* 2020;**21**:19.
16   Papalexi E, Mimitou EP, Butler AW, Foster S, Bracken B, Mauck WM 3rd, *et al.* Characterizing
the molecular regulation of inhibitory immune checkpoints with multimodal single-cell screens.
Nat Genet 2021;**53**:322–31.
17   Tsuchida CA, Brandes N, Bueno R, Trinidad M, Mazumder T, Yu B, *et al.* Mitigation of
chromosome loss in clinical CRISPR-Cas9-engineered T cells. *Cell* 2023;**186**:4567-4582.e20.
18   Song D, Wang Q, Yan G, Liu T, Sun T, Li JJ. scDesign3 generates realistic in silico data for
multimodal single-cell and spatial omics. *Nat Biotechnol* 2023. https://doi.org/10.1038/s41587-023-
01772-1.
19   Wu B, Zhang X, Chiang H-C, Pan H, Yuan B, Mitra P, *et al.* RNA polymerase II pausing factor
NELF in CD8+ T cells promotes antitumor immunity. *Nat Commun* 2022;**13**:2155.
20   Gasperini M, Hill AJ, McFaline-Figueroa JL, Martin B, Kim S, Zhang MD, *et al.* A Genome-wide
Framework for Mapping Gene Regulation via Cellular Genetic Screens. *Cell* 2019;**176**:1516.
21   Jost M, Santos DA, Saunders RA, Horlbeck MA, Hawkins JS, Scaria SM, *et al.* Titrating gene
expression using libraries of systematically attenuated CRISPR guide RNAs. *Nat Biotechnol*
2020;**38**:355–64.
22   Schraivogel D, Gschwind AR, Milbank JH, Leonce DR, Jakob P, Mathur L, *et al.* Targeted
Perturb-seq enables genome-scale genetic screens in single cells. *Nat Methods* 2020;**17**:629–35.
23   Shifrut E, Carnevale J, Tobin V, Roth TL, Woo JM, Bui CT, *et al.* Genome-wide CRISPR screens
in primary human T cells reveal key regulators of immune function. *Cell* 2018;**175**:1958-1971.e15.
24   Wessels H-H, Stirn A, Méndez-Mancilla A, Kim EJ, Hart SK, Knowles DA, *et al.* Prediction of on-
target and off-target activity of CRISPR-Cas13d guide RNAs using deep learning. *Nat Biotechnol*
2023. https://doi.org/10.1038/s41587-023-01830-8.
25   Zhang J, Bu X, Wang H, Zhu Y, Geng Y, Nihira NT, *et al.* Cyclin D–CDK4 kinase destabilizes
PD-L1 via cullin 3–SPOP to control cancer immune surveillance. *Nature* 2018;**553**:91–5.

954 26 Naqvi S, Kim S, Hoskens H, Matthews HS, Spritz RA, Klein OD, *et al.* Precise modulation of
955   transcription factor levels identifies features underlying dosage sensitivity. *Nat Genet* 2023;**55**:841–
956   51.

957 27 Replogle JM, Saunders RA, Pogson AN, Hussmann JA, Lenail A, Guna A, *et al.* Mapping
958   information-rich genotype-phenotype landscapes with genome-scale Perturb-seq. *Cell*
959   2022;**185**:2559-2575.e28.

960 28 Radhakrishnan SK, Lee CS, Young P, Beskow A, Chan JY, Deshaies RJ. Transcription factor Nrf1
961   mediates the proteasome recovery pathway after proteasome inhibition in mammalian cells. *Mol*
962   *Cell* 2010;**38**:17–28.

963 29 Dai W, Wu F, McMyn N, Song B, Walker-Sperling VE, Varriale J, *et al.* Genome-wide CRISPR
964   screens identify combinations of candidate latency reversing agents for targeting the latent HIV-1
965   reservoir. *Sci Transl Med* 2022;**14**:eabh3351.

966 30 Yin M, Guo Y, Hu R, Cai WL, Li Y, Pei S, *et al.* Potent BRD4 inhibitor suppresses cancer cell-
967   macrophage interaction. *Nat Commun* 2020;**11**:1833.

968 31 Tan Y-F, Wang M, Chen Z-Y, Wang L, Liu X-H. Inhibition of BRD4 prevents proliferation and
969   epithelial-mesenchymal transition in renal cell carcinoma via NLRP3 inflammasome-induced
970   pyroptosis. *Cell Death Dis* 2020;**11**:239.

971 32 Shu S, Wu H-J, Ge JY, Zeid R, Harris IS, Jovanović B, *et al.* Synthetic Lethal and Resistance
972   Interactions with BET Bromodomain Inhibitors in Triple-Negative Breast Cancer. *Mol Cell*
973   2020;**78**:1096-1113.e8.

974 33 Li Z, Guo J, Wu Y, Zhou Q. The BET bromodomain inhibitor JQ1 activates HIV latency through
975   antagonizing Brd4 inhibition of Tat-transactivation. *Nucleic Acids Res* 2013;**41**:277–87.

976 34 Mbonye U, Kizito F, Karn J. New insights into transcription elongation control of HIV-1 latency
977   and rebound. *Trends Immunol* 2023;**44**:60–71.

978 35 Wei P, Garber ME, Fang S-M, Fischer WH, Jones KA. A novel CDK9-associated C-type cyclin
979   interacts directly with HIV-1 tat and mediates its high-affinity, loop-specific binding to TAR RNA.
980   *Cell* 1998;**92**:451–62.

981 36 Peng J, Zhu Y, Milton JT, Price DH. Identification of multiple cyclin subunits of human P-TEFb.
982   *Genes Dev* 1998;**12**:755–62.

983 37 Stoeckius M, Zheng S, Houck-Loomis B, Hao S, Yeung BZ, Mauck WM 3rd, *et al.* Cell Hashing
984   with barcoded antibodies enables multiplexing and doublet detection for single cell genomics.
985   *Genome Biol* 2018;**19**:224.

986 38 Weinreb C, Rodriguez-Fraticelli A, Camargo FD, Klein AM. Lineage tracing on transcriptional
987   landscapes links state to fate during differentiation. *Science* 2020;**367**:eaaw3381.

988 39 Vitak SA, Torkenczy KA, Rosenkrantz JL, Fields AJ, Christiansen L, Wong MH, *et al.* Sequencing
989   thousands of single-cell genomes with combinatorial indexing. *Nat Methods* 2017;**14**:302–8.

990 40 Yang D, Cho H, Tayyebi Z, Shukla A, Luo R, Dixon G, *et al.* CRISPR screening uncovers a central
991   requirement for HHEX in pancreatic lineage commitment and plasticity restriction. *Nat Cell Biol*
992   2022;**24**:1064–76.

993 41 Rosen BP, Li QV, Cho H, Liu D, Yang D, Graff S, *et al.* Parallel genome-scale CRISPR screens
994   distinguish pluripotency and self-renewal. *BioRxivorg* 2023.
995   https://doi.org/10.1101/2023.05.03.539283.

996 42 Li QV, Dixon G, Verma N, Rosen BP, Gordillo M, Luo R, *et al.* Genome-scale screens identify
997   JNK-JUN signaling as a barrier for pluripotency exit and endoderm differentiation. *Nat Genet*
998   2019;**51**:999–1010.

43   Thanasopoulou A, Stravopodis DJ, Dimas KS, Schwaller J, Anastasiadou E. Loss of CCDC6
affects cell cycle through impaired intra-S-phase checkpoint control. *PLoS One* 2012;**7**:e31007.

44   Morra F, Luise C, Merolla F, Poser I, Visconti R, Ilardi G, *et al.* FBXW7 and USP7 regulate
CCDC6 turnover during the cell cycle and affect cancer drugs susceptibility in NSCLC. *Oncotarget*
2015;**6**:12697–709.

45   Zhou Y, Luo K, Liang L, Chen M, He X. A new Bayesian factor analysis method improves
detection of genes and biological processes affected by perturbations in single-cell CRISPR
screening. *Nat Methods* 2023. https://doi.org/10.1038/s41592-023-02017-4.

46   Dong M, Wang B, Wei J, de O Fonseca AH, Perry CJ, Frey A, *et al.* Causal identification of single-
cell experimental perturbation effects with CINEMA-OT. *Nat Methods* 2023;**20**:1769–79.

47   Hart T, Brown KR, Sircoulomb F, Rottapel R, Moffat J. Measuring error rates in genomic
perturbation screens: gold standards for human functional genomics. *Mol Syst Biol* 2014;**10**:733.

48   Morgens DW, Deans RM, Li A, Bassik MC. Systematic comparison of CRISPR/Cas9 and RNAi
screens for essential genes. *Nat Biotechnol* 2016;**34**:634–6.

49   Bunne C, Stark SG, Gut G, Del Castillo JS, Levesque M, Lehmann K-V, *et al.* Learning single-cell
perturbation responses using neural optimal transport. *Nat Methods* 2023;**20**:1759–68.

50   Li WV, Li JJ. An accurate and robust imputation method scImpute for single-cell RNA-seq data.
Nat Commun 2018;**9**:997.

51   Lun ATL, Bach K, Marioni JC. Pooling across cells to normalize single-cell RNA sequencing data
with many zero counts. *Genome Biol* 2016;**17**:75.

52   Sun T, Song D, Li WV, Li JJ. scDesign2: a transparent simulator that generates high-fidelity
single-cell gene expression count data with gene correlations captured. *Genome Biol* 2021;**22**:163.

53   Hao Y, Hao S, Andersen-Nissen E, Mauck WM 3rd, Zheng S, Butler A, *et al.* Integrated analysis of
multimodal single-cell data. *Cell* 2021;**184**:3573-3587.e29.

54   Lee K, Cho H, Rickert RW, Li QV, Pulecio J, Leslie CS, *et al.* FOXA2 is required for enhancer
priming during pancreatic differentiation. *Cell Rep* 2019;**28**:382-393.e7.

1025

1026

1027

1028

1029

 Figures

Figure 1. The Perturbation-response Score (PS) framework and benchmark. a, Overview of
different technical and biological factors that contribute to heterogenous perturbation outcomes from
single-cell perturbation datasets. **b,** Using downstream gene expressions to infer the value of PSs. **C,**
Overview of the scMAGeCK-PS that estimates PS value. **d-e,** Benchmark results of both PS and
mixscape using simulated datasets, where 50% (**d**) and 100% € gene perturbation effects are simulated
using scDesign3. Here, the expressions of 200 differentially expressed genes (DEGs) from bulk RNA-
seq (Nelf knockout vs. wild-type) are simulated, and ground truth efficiency value is indicated in red
color. **f,** Benchmark pipeline using real CRISPRi-based Perturb-seq datasets, where the perturbation
efficiency can be evaluated directly via gene expression. **g-h,** Benchmark results of mixscape and
scMAGeCK-PS using a published Perturb-seq dataset, by counting the numbers of cells or genes with
strong perturbation effects. A gene is considered to have strong perturbation effect, if a strong negative
correlation (Pearson correlation coefficient < -0.1) is observed between PS and the expression of that
gene across all perturbed cells. A cell is considered to be strongly perturbed, if its predicted efficiency
score (by scMAGeCK-PS or mixscape) within one cell is greater than 0.5. The Perturb-seq experiment

1046 is performed with low MOI condition, where most cells have only 1 expressed guide. **i,** An
1047 representative estimation results of scMAGeCK-PS and their correlations of ACTB expression.
1048
1049
1050
1051

1052
Figure 2. Additional benchmark results using genome-scale Perturb-seq and ECCITE-seq. a,
Benchmark procedure using a genome-scale Perturb-seq and a published, pooled T cell CRISPR screen.
b, The distribution of unstimulated and stimulated Jurkat cells along the UMAP plot. **c,** The correlation
of predicted scores by scMAGeCK-PS and mixscape. **d,** the Receiver-Operating Characteristic (ROC)
curve of both methods in separating positive and negative hits. **e-f,** Benchmark using a published
ECCITE-seq where PDL1 protein expression is used as gold standard (**e**), and the performance of
different methods in terms of predicting PLD1 protein expression (**f**).
1060

Figure 3. Dose-dependent responses of perturbations. a, The correlation between a gene's PS and a
phenotype of interest indicates positive (or negative) regulations. **b-c,** The correlation between PDL1
protein expression and the PS of CUL3 (**b**) and STAT1 (**c**). CUL3 is a known negative regulator of
PDL1, while STAT1 is a known positive regulator. **d,** The classification of buffered or sensitive genes,
based on perturbed gene expression and PS. **e,** The classification of buffered or sensitive genes from
published Perturb-seq datasets focusing essential genes in K562[26]. **f,** The perturbation-expression plot of
PSMA3, a buffered gene. **g,** The log fold changes of mark gene expressions (columns) upon perturbing
proteasome genes (rows) from the essential gene Perturb-seq dataset.
1071

Figure 4. Perturb-seq on HIV latency. a, The experimental design of Perturb-seq. **b,** The UMAP plot of single-cell transcriptome profiles. Cells are colored by three different conditions. **c,** The distribution of BRD4 PS. **d,** The expression of HIV-GFP. **e,** The correlations between HIV-GFP expression and BRD4 PS that does not use HIV-GFP as target gene. **f,** The distribution of CCNT1 PS. **g,** The protein expression of HIV-GFP in response to CCNT1 knockout in different cell states (TNF-alpha vs non-stimulated).

Figure 5. Pooled scRNA-seq on pancreatic differentiation. a, Experimental design of multiplexing scRNA-seq on the knockout clones of different genes. **b,** The UMAP plot of single-cell transcriptome profiles, colored by different clusters (left) or clones (right). **c,** The PS distribution of HHEX. **d,** The percentage of cells in PP/LV/DUO cell types from different clones. **e,** The correlations of CCDC6 PSs calculated from different HHEX cell types. The Pearson Correlation Coefficient (PCC) is calculated from all cells with CCDC6 knockouts and is shown as numbers on the heatmap. **f,** Two different distribution patterns of CCDC6 PSs. **g,** The top enriched GO terms of DEGs from PP/PP in transition. Enrichr was used to perform enrichment analysis. **h,** The percentage of cells in PP/LV/DUO cell types from CCDC6 clones. **i,** The percentage of cells with PDX1+ (a PP marker) or HNF4A+ (a LV marker) by flow cytometry sorting. The data is based on two CCDC6 knockouts (KO1, KO2) and one wild-type (WT) control. Three independent replicates are performed for each condition. The multiple comparison-adjusted p value is calculated by one-way ANOVA test. *p<0.05, **p<0.01.

Supplementary Figure Legends

Supplementary Figure S1. Benchmark different methods using simulated and real datasets. a, Steps to generate simulated datasets using scDesign3 from a real scRNA-seq dataset that knocks out Nelfb gene. **b-e,** The score distribution of scMAGeCK-PS and mixscape using different DEGs and different values of true efficiencies. **f-g,** Similar with Figure 1f-g, but using a published high MOI Perturb-seq dataset in the same study. **h-i,** Benchmark results of different methods on another published CRISPRi-based Perturb-seq, where mismatches are introduced into guides to attenuate perturbation effects. The Pearson correlation coefficients (PCCs) between the predicted scores of each method and the expressions of perturbed genes are reported for every perturbed gene (**h**), and between the predicted scores and predicted sgRNA activities (**i**), using the prediction methods provided in the original study[21].

Supplementary Figure S2. A genome-scale Perturb-seq. a-b, The distribution of scMAGeCK-PS and mixscape predicted scores of top hits including CD247 (**a**) and LCK (**b**) in the pooled screen. **c,** The correlation between PSs and perturbed gene expression.

Supplementary Figure S3. Predictions of PDL1 protein expression from a published ECCITE-seq dataset. The ROC curve, the correlations between scMAGeCK-PS results with PDL1 protein expression, and the correlations between mixscape results with PDL1 protein expression are reported for each gene. The correlations are separated by classifications of each single cell: NP (non-perturbed), defined as mixscape score <=0.5; and KO (knockout), defined as mixscape score >0.5. For a fair comparison, we used mixscape classification results to plot PSs (mid panel).

Supplementary Figure S4. Buffered genes and sensitive genes. a, RPL4, a buffered gene. **b-c,** HSPA5 and GATA1, two sensitive genes. **d,** A gene (BRD4) whose expression has no correlation with PS.

Supplementary Figure S5. The log fold changes of gene expressions upon perturbing genes within the same protein complex, including ribosomal subunits (**a**), RNA polymerase (**b**) and mediator complex (**c**) in essential gene Perturb-seq. **(d)** The log fold changes of proteosome gene expressions (columns) upon perturbing proteasome genes (rows) from the genome-scale Perturb-seq.

Supplementary Figure S6. HIV Perturb-seq. a, The number of genes (nFeature_RNA), UMI counts (nCount_RNA) and the fraction of mitochondrial RNAs in three different conditions. **b,** Clustering results. **c,** Enriched Gene Ontology (GO) terms of cluster 8. **d,** The distribution of BRD4-targeting gRNAs. **e,** The expression distribution of BRD4 signature genes in cluster 8 vs other clusters. Only cells express BRD4-targeting gRNAs are included. **f,** Differential expression results between BRD4 PS+ cells vs BRD4 PS- cells.

Supplementary Figure S7. HIV Perturb-seq. a, The expressions of CCNT1 (left) and CCNT1-targeting gRNAs (right). **b,** Differential expression results between CCNT1-targeting cells and non-targeting control cells in two different cell states. **c,** The expressions of NFKB1. **d,** The quantitative perturbation-expression relationship between GFP and CCNT1 PS, similar with Figure 4e.

Supplementary Figure S8. Cell type assignment based on known expression markers of different cell types in pancreatic differentiation scRNA-seq.

Supplementary Figure S9. DEG analysis. a-b, The distribution of FOXA1 PSs across two different clones. **c,** The expression pattern of FOXA1. **d-e,** The DEG analysis results of CCDC6 knockout clones vs. wild-type clones in different cell types. **f,** The overlap of statistically significant DEGs in DE and LV/DUO cell types.

Supplementary Figure S10. Different CCDC6 functions. a-b, The two patterns of CCDC6 PSs in LV/DUO (**a**) and DE in transition (**b**) cell types. **c-f,** Additional enriched terms using Enrichr on DEGs of CCDC6 knockout.

Supplementary Figure S11. Flow cytometry analysis of PDX1 and HNF4A expression upon CCDC6 knockout. One representative plots of three biological replicates are shown.

Supplementary Table S1. Genome-scale Perturb-seq library design.

Supplementary Table S2. HIV Perturb-seq library design.

Supplementary Table S3. Sequencing summary of HIV Perturb-seq.

Supplementary Table S4. Genotype summary of 10-clone scRNA-seq pancreatic differentiation dataset.