Fortune may favor the flexible: Environment-dependent behavioral shifts in invasive

coquí frogs

Katharina M Soto[1,2], Devin Edmonds[3], Andrea L Colton[3], Faith O Hardin[1], Eva K Fischer[1,4,5*]

[1] Department of Evolution, Ecology, and Behavior; School of Integrative Biology; University of

Illinois Urbana-Champaign

[2] Program of Evolution, Ecology, and Conservation; School of Integrative Biology; University of

Illinois Urbana-Champaign

[3] Illinois Natural History Survey; Prairie Research Institute; University of Illinois Urbana-

Champaign

[4] Carl R. Woese Institute for Genomic Biology; University of Illinois Urbana-Champaign

[5] Department of Neurobiology, Physiology, and Behavior; University of California Davis

*Corresponding author:

Eva K Fischer

Department of Evolution, Ecology, and Behavior

School of Integrative Biology

University of Illinois Urbana-Champaign

505 S Goodwin Ave

Urbana, IL 61801

Abstract

Biological invasions are a major driver of global biodiversity loss, impacting endemic species, ecosystems, and economies. While the influence of life history traits on invasive success is well-established, the role of behavior in the invasive potential of animals is less studied. The common coquí frog, *Eleutherodactylus coqui*, is a highly successful invader in Hawaii. We build on previous research characterizing changes in physiology and morphology to explore behavioral variation across the invasive range of coquí in Hawaii. We hypothesized that behavioral traits contributing to invasion expansion would be reflected in behavioral differences at the invasive center versus invasive edge. To investigate whether differences in the field represent local adaptation or behavioral plasticity, we additionally evaluated behavior following acclimation to a shared laboratory environment. While we identified only subtle behavioral variation among populations in the field, we found that individuals from all populations became less bold, active, and exploratory in the laboratory, converging on a similar behavioral phenotype. Alongside previous work, our results suggest that coquí adjust their behavior to local environmental conditions across their invasive range, and that behavioral flexibility may contribute to invasive success.

Keywords: behavior, invasive species, behavioral plasticity, anuran, physiology

Introduction

Biological invasions are a major cause of global biodiversity loss, impacting endemic species, ecosystems, and economies (Kumschick et al. 2012; Lennox et al. 2015). While the impacts of invasive species have been thoroughly examined, we are still in the process of understanding the attributes that contribute to invasive species' success. Research examining invasive animals has demonstrated the importance of life history traits, such as high reproductive rates (Alcaraz et al. 2005), rapid population growth, dispersal capabilities, and a generalist diet (Sakai et al. 2001; Alex Perkins et al. 2013). The role of behavior remains less studied, but has been implicated in the spread of some invasive species (Sol and Lefebvre 2000; Weis 2010; Hudina and Hock 2012; Damas-Moreira et al. 2019). For instance, Candler & Bernal (2015) found that introduced cane toads (*Rhinella marina*) displayed reduced neophobia and increased exploration of novel prey and objects compared to toads from native populations (Candler and Bernal 2015). Similarly, Damas-Moreira et al. (2019) found that invasive wall lizards (*Podarcis sicula*) were more bold, exploratory, and neophilic compared to non-invasive wall lizards (*Podarcis virescens*) (Damas-Moreira et al. 2019). These studies highlight how certain behavioral traits and/or collections of behaviors ("behavioral syndromes") can drive invasive success by facilitating range expansion and the exploitation of new resources. Furthermore, behavioral plasticity may be particularly crucial during the initial stages of invasion when adaptation can be slow due to low genetic diversity resulting from founder effects and bottlenecks during colonization (Tsutsui et al. 2000; Peacock et al. 2009; Wright et al. 2010). Yet to succeed, invaders must adjust rapidly to novel environments, prey, predators, and competitors (Chapple et al. 2012), and behavioral plasticity may help bridge this gap.

Behavioral and physiological changes may be closely linked to invasion potential as behavior (behavioral plasticity) may help overcome novel physiological challenges, and physiological adjustments may in turn facilitate coordination of favorable behavior. Indeed, colonization of novel environments often presents physiological challenges, and behavior may

be an important coping mechanism. High metabolic rates, often associated with a 'fast pace of

life'—marked by rapid growth, early reproduction, and a generalist diet—are traits already

recognized as key contributors to invasive potential (Lagos et al. 2017). Additionally, changes in

metabolic rates have been correlated with changes in boldness, exploration, and activity (Biro

and Stamps 2010; Myles-Gonzalez et al. 2015; Baškiera and Gvoždík 2022), behaviors that

have been identified as relatively greater in invasive species as compared to native organisms

(Candler and Bernal 2015; Damas-Moreira et al. 2019). These relationships may be particularly

important in ectotherms that directly regulate body temperature through their behavior; however,

connections between behavior and physiology during biological invasions have only been

examined in a few systems (Myles-Gonzalez et al. 2015; Mathot et al. 2019; Mowery et al.

2021).

The common coquí (*Eleutherodactylus coqui*) is an excellent model for investigating

mechanisms that contribute to invasion success. These small, nocturnal frogs are endemic to

Puerto Rico, where they are an integral part of the island's ecosystem and culture (reviewed in

Westrick et al. 2022). However, their introduction to other regions has garnered them a

reputation as one of the worst invasive species in the world (Lowe et al. 2000). Coquí have

successfully colonized Hawaii, Guam, and various Pacific and Caribbean islands (Kraus and

Campbell 2002; Beard and Pitt 2005), and have been introduced to Costa Rica (Barrantes-

Madrigal et al. 2019). Their invasion of Hawaii extends across the archipelago, including the

islands of Hawaii, Maui, Oahu, and Kauai (Beard et al. 2009), though the Kauai population is

believed to have been exterminated (Beard and Pitt 2012). Their loud and incessant calls have

made them an unmistakable presence in the Hawaiian nightscape, impacting property values,

tourism, and the ornamental plant trade, and prompting campaigns to prevent their spread

(Beard and Pitt 2005).

Since their introduction to Hawaii in the 1980s, coquí populations have exploded to

densities three-fold those in their native range, reaching up to 91,000 frogs per hectare (Beard

et al. 2009). Unlike most frogs, coquí are direct developing, meaning they complete

metamorphosis within the egg, skipping a free-living tadpole stage. This life history allows for

rapid reproduction and independence from standing freshwater (reviewed in Westrick et al.

2022). This reproductive strategy, combined with a voracious, generalist diet, has fostered their

rapid population growth.

On the Big Island of Hawaii where populations are largest, coquí are continuing to

expand, including to higher elevations (Marchetti et al. 2023; Marchetti et al. 2024). Generally,

this combination of factors has led to a tradeoff between escaping high competition at low

elevations and coping with colder and more variable environmental conditions further uphill.

Previous studies have documented the invasive spread of coquí in Hawaii (Marchetti et al.

2023; Marchetti et al. 2024), characterized the impacts of coquí on local plants, animals, and

economies (Kraus and Campbell 2002; Beard and Pitt 2005; Beard et al. 2008; Smith et al.

2018), and explored physiological changes associated with the colonization of distinct habitats

(Beard et al. 2008; Rollins-Smith et al. 2015; Haggerty 2016; O'Neill et al. 2018; Marchetti et al.

2023; Marchetti et al. 2024). Behavior remains relatively understudied, aside from Marchetti and

Beard (2021) who demonstrated that Hawaiian coquí maintain avoidance responses to native

Puerto Rican predators ~20 generations after invasion. Like other invasive species, Hawaiian

coquí exhibit reduced genetic diversity as compared to their native counterparts due to founder

effects (Tsutsui et al. 2000; Peacock et al. 2009). Nonetheless, their populations have thrived,

establishing a presence in diverse habitats and ecosystems across the island. This versatility in

the face of low genetic diversity suggests a role for plasticity in their invasive potential (Wright et

al. 2010). Indeed, the above studies document a combination of population differences and

plasticity in various traits.

Our goal was to investigate how behavior might contribute to the invasive expansion and

success of *E. coqui* in Hawaii. We adopted a stepwise approach, involving field and laboratory

experiments. First, we characterized activity, exploration, and boldness across elevational and

density gradients in Hawaii to understand to what extent behavior varies across environmental

conditions at the invasive center verses edge (Chuang and Peterson 2016). We considered

high-density, low-elevation sites representative of the invasive center, and low-density sites at

both high and low elevations as invasive edges. We tested the alternative predictions that (1)

more active and bold coquí would be found at invasive edges as these individuals would be

more likely to push range boundaries, or (2) more bold, active coquí would be found at the

invasive center as these individuals would be better equipped to deal with the competitive

challenges of high densities. By targeting high- and low-density sites at different elevations, we

additionally asked whether the effects of density were modulated by more challenging

environmental conditions.

Second, we asked whether behavioral traits changed when we introduced coquí to a

novel, laboratory environment, specifically measuring behavior in the same individuals in the

field and in the laboratory. This allowed us to distinguish environmentally-mediated behavioral

plasticity from stable behavioral differences that would indicate developmental influences or

local adaptation to different densities and/or elevations. We also measured resting metabolic

rate in the laboratory to test for an association between metabolic rate and behavior, thereby

exploring a potential mechanism by which behavioral changes are coordinated and linked with

physiological adaptation to high elevations. Taken together, our findings contribute a behavioral

perspective to complementary studies characterizing life history and metabolic traits in Hawaiian

coquí, adding an additional dimension to our understanding of this small frog's invasive

success.

Methods

All frog husbandry and experimental methods were approved by the University of Illinois

Urbana-Champaign Animal Care and Use Committee (Protocol # 20147). Frog collection and

export were approved by the State of Hawaii Department of Land and Natural Resources

(Permit # H0422-04; EX-22-11). Given the invasive nature of coquí in Hawaii, our collection

posed no conservation concerns.

Field Collection

We collected 60 male coquí on the Big Island of Hawaii. We collected frogs from four site types:

high-density/low-elevation (N=16), low-density/low-elevation (N=14), high-density/high-elevation

(N=15), and low-density/high-elevation (N=15). Collection sites were at Lava Tree State

Monument Park (HL, 195m, [19°28'58"N 154°54'09"W]), 'ŌlaʻA Forest Reserve (LH, 870m,

[19°27'10"N 155°11'14"W]), Watershed Forest Reserve (LH, 860m, [19°25'02"N 155°09'45"W]),

Upper (HH, 850m, [19°34'28"N 155°11'27"W]) and Lower (HL, 250m, [19°37'46"N

155°05'37"W]) Stainback Highway, Homestead Road (LL, 200m, [19°50'33"N 155°06'50"W]),

and Hilo Watershed Reserve (HH, 720m, [19°50'36"N 155°11'08"W]).

We defined low-elevation as <300m and high-elevation as >700m. We classified

population density based on recent surveys (Marchetti et al. 2024), and used call surveys during

collection as a rough confirmation of density differences, considering higher sound pressure is

associated higher male coquí density (Benevides et al. 2019). We performed acoustic

recordings at each collection site and our field station using a shotgun microphone (Sennheiser

ME 66/K6) and a handheld recorder (Zoom H1 Handy Recorder). Intensity analysis of recorded

calls was conducted using Audacity v. 3.0.2 (Audacity Team 2021). **L**ow-density sites had an

average frequency of 500 Hz at 40.7 dB and high-density sites had an average frequency of

2,639 Hz at 55.6 dB**,** lending support to our initial classifications based on Marchetti et al. 2024.

Frogs were hand-captured between 19:00 and 00:00 hours and individually placed in

500 mL plastic cups with ventilation holes and damp paper towels for moisture and humidity.

Frogs were transported from each collection site to our field station in Leilani Estates, Hawaii

County, HI, and held for 24 hours before behavioral testing the following night. We conducted all

field behavioral assays in this manner to minimize the acute effects of temperature and humidity

differences between sites on behavior, make the timing of trials uniform, allow frogs to recover

from capture stress, and due to limitations in transporting our behavioral arena and recording

equipment.

Field Behavioral Assays

We conducted field behavioral trials between May and June 2022, from 19:00 to 03:00 hours.

Average temperature was 22.9 °C (range: 19–25) and humidity was 76.7% (range: 70–85). We

used two behavioral arenas constructed from ½" PVC pipes and white mesh fabric, with

dimensions measuring 83 x 58 x 48 cm. The arenas were illuminated by red light lamps and

infrared illuminators (Univivi IR Illuminator), and we recorded behavior using infrared

camcorders (Sony DCR-SR85). Arenas were placed ~8 meters apart on opposite sides of the

field station to prevent trial interference. We constructed emergence chambers using a ½" PVC

pipe held vertical by a plastic cup base and a retractable cardboard cover.

To examine behavior, we used two tests: (1) an emergence test to quantify latency to

emerge from a shelter (a commonly used metric of boldness, see Magnhagen et al. 2014;

Myles-Gonzalez et al. 2015; Yuen et al. 2017), and (2) an open field test to quantify activity and

exploration. For the emergence test, each frog was placed in the emergence chamber and

given a 3-minute acclimation period. After the acclimation period, the cover of the chamber was

removed from outside the arena. Each frog was given 10 minutes to emerge from the chamber.

Individuals that did not emerge after 10 minutes were gently removed from the PVC pipe without

direct handling of the frog. To ensure this experience did not bias subsequent behavior, we

compared activity and exploration between frogs that emerged on their own versus those that

were removed and found no differences (exploration: $F_{1,52}= 0.04$, p= 0.838; activity: $F_{1,52}= 0.06$,

p= 0.801). Latency to emerge was recorded as the moment when the majority of the frog's body

had exited the PVC pipe. Immediately following the emergence test, we performed an open field

test for 15 minutes. During the open field test, we quantified exploration as the proportion of

areas visited, and activity as the number of times a frog crossed lines between areas

superimposed during analysis (see below). We expected a positive correlation between activity

and exploration; however, these two variables are not perfectly correlated and can vary between

individuals or populations, as frogs that move the same amount can do so over smaller or larger

areas (e.g., 10 line crossings back and forth between 4 areas versus 10 line crossings across

10 areas).

Laboratory Transfer and Husbandry

Within four days of capture, all frogs were shipped to the University of Illinois Urbana-

Champaign. For transport, each frog was placed in a 500 ml plastic cup with a moist paper

towel and dead leaves. Cups were packed into Styrofoam boxes with insulation and padding

material and shipped via climate-controlled UPS overnight. All frogs arrived alive and in good

body condition. Upon arrival, frogs were individually housed in tanks measuring 30 x 20 x 15

cm, with ~ 2-3 cm of soil and sphagnum moss as a substrate. Additionally, each frog was

provided a ~20cm long PVC pipe as a shelter. We maintained humidity >70% and temperature

69–74 °F on a 12L:12D light cycle with a shifted dark period (lights off at noon and on at

midnight). We fed frogs live brown crickets (*Acheta domesticus*) gut loaded with vitamins three

times weekly. Four individuals died over the course of the study due to escape and subsequent

desiccation.

Laboratory Behavioral Assays

We conducted laboratory trials (N=56) between September and November 2022 after frogs

were confirmed healthy and had acclimated to the laboratory for 14–18 weeks. All trials were

conducted between 13:00 and 17:00 hours, one to four hours into the frogs' wake cycle after

lights went off. To replicate the conditions of field capture and acclimation, we held individuals in

500 mL plastic cups lined with moist paper towel for 24 hours prior to laboratory behavioral

trials. Trials were conducted in an unoccupied animal room with average temperature 21.7 °C

(range: 18–24) and average humidity 74.8% (range: 50–98). We used the same arena as in the

field assays; however, we used curtains to dampen sound reflections off the walls and maintain

humidity in the arena. Behavioral assays in the laboratory closely followed the field protocol,

consisting of a 13-minute emergence test (3 minutes acclimation and 10 minutes to emerge)

immediately followed by a 15-minute open field test. Ambient sound recordings collected during

behavioral trials in the field were used to replicate natural background noise, including the

presence of other coquí.

Physiological Assays

We conducted resting metabolic rate (RMR) assays (N=55) in the laboratory between February

and March of 2023, 12 weeks after laboratory behavioral assays. Given that resting metabolic

rate (RMR) can be influenced by numerous factors, including diet, temperature, age, sex, and

environmental conditions (particularly in ectotherms), prior studies have highlighted that a

common garden approach is the preferred method for controlling these variables (Biro and

Stamps 2010; Burton et al. 2011).

We used an intermittent flow-through respirometry system to measure RMR. We placed

frogs in 127 mL chambers made from ½" diameter PVC plastic pipe and PVC end caps. We

configured a flow-through system with an open circuit and incurrent flow measurement using a

PP2 Dual Pump, FB8 Flow Measurement System, RH-300 Water Vapor Analyzer, Fox Box, and

UL2 Universal Interface (all from Sable Systems International). The incurrent airstream was

scrubbed of CO_2 and water vapor. After allowing frogs to acclimate in the chamber for 20

minutes, we sampled CO_2 production and O_2 consumption for each individual every 5 seconds

over a 40-minute window. We switched from a chamber containing a frog to a control chamber

between individuals to establish a baseline measurement of ambient CO_2 and O_2, and to

account for measurement drift. Measurements were recorded following a 2-day fast to eliminate

the energy expenditure associated with digestion (Naya et al. 2009). To calculate RMR, we first

matched each frog to the control measurement taken directly after the trial. For three

individuals, we substituted controls from the following trial on the same day due to problems

with their paired control. We deleted 5 minutes from the start of each trial to remove any error

associated with switching tubing between the control and frog. We also deleted 7.3 minutes

(determined from visually inspecting plots) from the end of each trial to remove the lag between

sequential CO_2, O_2, and H_2O sensors in our system. After trimming data, we calculated the

fractional concentration of CO_2 and O_2, and H_2O adjusted for barometric pressure.

To calculate RMR, we first calculated the rate of O_2 consumption (VO_2) and CO_2

emission (VCO_2) using Lighton (2008) equations 10.1:

$$VO_2 = FR * {O_{2\,diff}} \Big/ {1 - O_2}$$

261   and 10.8:

$$VCO_2 = \frac{FR * (CO_{2\,diff} - (CO_2 * (1 - H_2O)) * \frac{(FR * O_{2\,diff})}{(1 - O_2)})}{1 - (CO_2 * (1 - H_2O))}$$

where FR = flow rate, $O_{2\,diff}$ is the average O_2 during the control minus the O_2 produced by a

frog, $CO_{2\,diff}$ is the CO_2 produced by a frog minus the average CO_2 during the control, and CO_2,

O_2, and H_2O are the average values during control. We then calculated RMR as the lowest 3-

minute average VO_2 and VCO_2 during each trial. Next, we calculated the respiratory quotient

(RQ) by dividing the lowest 3-minute average VCO_2 by the lowest 3-minute average VO_2. Finally,

we calculated mass corrected RMR by dividing each frog's RMR by its mass (g), hereafter

referred to as RMR. Data manipulation was performed using the packages 'zoo' and 'tidyverse'

(Wickham et al. 2019; Zeileis et al. 2023) in R.

Statistical Analysis

We used Behavioral Observation Research Interactive Software (BORIS) version 8.13 (Friard

and Gamba 2016) to code exploration and activity from videos. We analyzed data in R version

4.2.4 (R Core Team 2022) in Rstudio v.4.2.2 (RStudio Team 2021). We analyzed response

variables (i.e., behavior and RMR) using raw data for those variables with a normal distribution.

We applied a square root transformation to activity because it was not normally distributed. We

lost field video recordings for five frogs from low-density/low-elevation (LL) and eight frogs from

high-density/low-elevation sites (HL) due to a hard drive failure. The analysis of field behavior

(i.e., exploration and activity) is based on the remaining dataset (N=49). Field emergence data

was recorded live during trials, and all available data (N=60) were analyzed. We used power

analysis with the 'pwr' package (Champley 2020) in R to assess the consequences of this data

loss. Even with reduced sample sizes, we maintained high power (94%) to detect large effects

and moderate power (64%) to detect medium effects. We used the R packaged 'effectsize'

(Ben-Shachar et al. 2020) to estimate eta-squared (η^2) effect sizes.

To examine behavioral differences in the field, we used the 'lm' function from the 'stats'

package to construct linear models that included behavior (activity or exploration) as the

response variable, and density, elevation, and their interaction as predictor variables. We used

similar models to test for behavioral and RMR differences in the laboratory. For linear models,

statistical significance was called at $p \leq 0.05$ and a trend toward statistical significance at $p \leq$

0.10. We interpreted effect sizes: $\eta^2 > 0.01$ as a small effect, $\eta^2 > 0.06$ as a medium effect, and

$\eta^2 > 0.14$ as a large effect,

For emergence, not all frogs performed the behavior during the trial period, making a

direct test of latency to emerge problematic. To address this issue, we ran Cox Proportional

Hazard Models using the R packages 'survival' (Therneau 2021) and 'survminer' (Kassambara

et al. 2020) to test for differences in the probability of emergence based on density, elevation,

and their interaction.

To characterize behavioral differences between the field and the laboratory, we

constructed linear mixed models using the 'lmer' function in the 'lme4' package (Bates et al.

2011) with frog ID as a random effect and trial (field vs laboratory), density, elevation, and their

interactions as fixed effects. We used Type III ANOVAs to generate test statistics and p-values

for main effects and interactions. To prevent overfitting, we first ran models with all possible two-

and three-way interactions and removed interactions when non-significant. Where appropriate,

we used the 'emmeans' package (Lenth et al. 2023) for pairwise post-hoc comparisons. We

calculated Kendall's tau correlation coefficients and test statistics between behaviors and RMR

using the cor.test function from the 'Hmisc'. We interpreted correlation coefficients following the

criteria ≈0.1 as a small effect, ≈0.3 as a moderate effect, and values >0.5 as a strong effect

(Garamszegi et al. 2013).

Finally, we again used linear models to test whether correlations between behaviors, and

between RMR and behavior, depended on density and elevation at collection sites. We ran the

latter tests using measures of behavior in the laboratory as these behavioral assays occurred

closer in time and under the same conditions as RMR measurements.

Results

Field Behavioral Assays

For exploration, we found statistically significant main effects of density ($F_{1,37}$= 4.91, p= 0.033),

elevation ($F_{1,37}$= 5.38, p= 0.026), and their interaction (i.e., site type; $F_{1,37}$= 6.26, p= 0.017)

(Table 1, Figure 1a). While the effect sizes of density (η^2<0.01) and elevation (η^2=0.01) were

small, the effect of the interaction was large (η^2=0.14). For activity, we found a trend toward

statistical significance of density ($F_{1,36}$= 3.34, p= 0.072) and the interaction between density and

elevation ($F_{1,36}$= 3.38, p= 0.074) (Table 1, Figure 1b). The effect sizes of density (η^2<0.01) and

elevation (η^2<0.01) were small, but the effect of the interaction was moderate (η^2=0.09). For

emergence, there was also a trend toward significance for the effect of elevation (X^2= 3.29, p=

0.070), with low elevation frogs emerging more quickly (Table 1, Figure 1c). However, pairwise

comparisons were not significant in Tukey corrected, post hoc comparisons for any behavior

(exploration, activity, or emergence).

Laboratory Behavioral Assays

After acclimation to the laboratory, we found no differences in exploration or activity based on

density, elevation, or their interaction (Table 1, Figure 2a,b). The effect of elevation on

emergence time had a trend toward significance in the lab as in the field (X^2= 3.08, p= 0.079)

(Table 1), though we note very few frogs emerged at all (5/54 frogs). Effect sizes were small to

negligible for all models.

Comparison of Field and Laboratory Behaviors

Comparing behavior for the same individuals in field and laboratory assays, we observed

significant main effects of trial (field vs lab) on exploration ($F_{1,43}$= 5.41, p= 0.025) and activity

($F_{1,47}$= 7.54, p= 0.008) (Figure 2b; Table 2). No two-way interactions were significant, but the

three-way interaction between trial, density, and elevation was significant for exploration ($F_{1,43}$=

5.12, p= 0.029) and approaching significance for activity ($F_{1,43}$= 3.80, p= 0.057), and the effect

sizes were moderate (exploration: η^2=0.11; activity: η^2=0.07). These interactions were driven by

the fact that behavioral shifts from the field to the lab were greater for some site types than

others (Table 3, Figure 2a,b). Specifically, frogs from high-density/low-elevation sites showed

the largest shift in behavior (exploration: t_{51}= 2.79, p= 0.0046; activity: t_{46}= 2.46, p= 0.018).

The effect of trial on emergence probability showed a non-significant trend in the full

model (X^2= 3.01, p= 0.083) (Table 2) and a significant effect (X^2= 18.48, p<0.0001) when all

non-significant two- and three-way interactions were removed (Figure 2c). In general, frogs

explored less, moved less, and were less likely to emerge in the laboratory as compared to the

field (Figure 2).

Activity and exploration were strongly correlated in both the field (tau= 0.82, z= 7.25,

p<0.001) and the lab (tau= 0.71, z= 7.38, p<0.001), with the strength of this relationship was

significantly different based on trial context ($F_{1,92}$= 4.9, p= 0.029) (Figure 3). Neither exploration

nor activity was correlated with time to emerge in the field (exploration: tau= -0.06; activity: tau=

0.004) or the lab (exploration: tau= 0.04; activity: tau= 0.03).

Due to known effects of temperature and humidity on behavior, and variation in these

metrics between the lab and the field, we tested for effects of temperature and humidity on

behavior within and across contexts. We found no significant effects of either temperature or

humidity on any behavior.

Physiological Assays

We found no main effect of density ($F_{1,50}$= 0.74, p= 0.393) or elevation ($F_{1,50}$= 0.32, p= 0.571) on

RMR after acclimation to the laboratory, but a non-significant trend in their interaction ($F_{1,50}$=

3.00, p= 0.089) (Figure 4a). The effect sizes of density (η^2=0.01) and elevation (η^2<0.01) were

small, but the effect of the interaction was moderate (η^2=0.06). However, pairwise differences

between site types were not significant in Tukey corrected, post hoc comparisons. While there

were no substantial differences in RMR based on population of origin, RMR moderately

predicted exploration (tau= 0.27, $F_{1,50}$= 12.03, p= 0.001) (Figure 4b) and activity (tau= 0.25,

$F_{1,50}$= 7.53, p= 0.008) (Figure 4c), but not propensity to emerge (tau= -0.02, $F_{1,50}$= 0.20, p=

0.66).

Discussion

During their expansion across Hawaii, coquí have experienced changes in ecological

conditions, notably extreme increases in population densities and climatic fluctuations

associated with colonizing higher elevations. Previous research has characterized the

consequences of these shifting conditions on ecosystem dynamics, morphology, and physiology

(Beard et al. 2008; Rollins-Smith et al. 2015; Haggerty 2016; O'Neill et al. 2018; Marchetti et al.

2023; Marchetti et al. 2024), but behavior remains relatively understudied in this context (but

see Marchetti and Beard 2021). To address this gap, we conducted a series of experiments,

examining variations in behavior across environmental and experimental contexts. We found

that behavior varied based on interactions between density and elevation across the invasive

range. However, these field-based differences disappeared after acclimation to the laboratory,

suggesting behavioral variation in the field is primarily driven by behavioral plasticity rather than

local adaptation. Additionally, we found that physiology predicted behavior in the laboratory.

Below, we discuss the implications and limitations of our findings.

Our initial prediction was that behavior would vary with density if certain individuals were

more likely to expand outward from high-density population centers toward low-density

population edges, and that this effect might be modulated by the challenges associated with

moving up in elevation. In the field, we found a trend toward faster emergence time at high

densities, in line with the prediction that individuals living at high densities are bolder, perhaps

conferring an advantage in acquiring food and mates in a competitive environment. For

exploration and activity, we found a significant interaction between density and elevation on

exploration, and a trend toward the interaction of density and elevation influencing activity.

Importantly, effect sizes indicated that neither variable alone predominated, but that the effects

of density differ across elevations. For example, interactions could arise if density dependent

differences associated with competitive abilities are modulated by the physiological constraints

imposed by higher elevations. We note that our sample sizes for exploration and activity were

relatively small. Nonetheless, statistical analyses indicated sufficient power to detect medium to

large effects, suggesting that—if additional behavioral differences between populations do

exist—they are subtle. Furthermore, previous studies in amphibians have primarily compared

invasive and native populations, where behavioral differences might be more pronounced. We

suggest this is an interesting avenue for future comparisons in coqui. In sum, while the subtle

effects we identify provide only weak evidence for behavioral differences associated with

invasive range expansion in Hawaiian coqui, they underscore the potentially complex

interactions of multiple factors influencing behavior.

Although we found little evidence for population differences in the field, by comparing

behavior of the same individuals in the field and after acclimation to the laboratory, we found

strong evidence of behavioral changes in a novel environment. Following acclimation to the

laboratory, frogs from all populations exhibited reduced activity, exploration, and boldness. We

propose two potential explanations for these behavioral shifts. First, decreased activity and

exploration may signal a shift towards cryptic behavior, which is a key factor in the success of

invasive species, enabling them to evade predation, reduce competition, and thrive in new

environments (Jarić et al. 2019). Alternatively, this behavioral shift may result from a reduced

need or motivation to move and explore under laboratory conditions where there is an absence

of resource competition and mating opportunities. Although we cannot distinguish these

alternatives, uniform behavior in a novel environment indicates a generalized strategy across

individuals from different site types, suggesting environmentally induced behavioral

responses—rather than local adaptation—drive behavioral variation among Hawaiian coquí.

Behavioral plasticity may enhance invasion success by enabling rapid responses to new

environmental conditions, even in the absence of genetic variation (Wright et al. 2010). Indeed,

another study in Hawaiian coquí found that behavioral plasticity in microhabitat use may

mediate colonization of higher elevation sites (Marchetti et al. 2024), providing an explanation

for a lack of physiological changes among coquí from different elevations in some studies

(Haggerty 2016; Marchetti et al. 2024), and underscoring the potential for a combination of

genetic and environmental influences underlying traits across coquí's native and invasive range.

In addition to single behaviors, we were interested in correlations among our behavioral

metrics. Chapple et al. (2012) suggest that collections of correlated behaviors could enhance

invasive success and be classified as an 'invasion syndrome' (Chapple et al. 2012). For

example, correlations between aggression and activity in invasive crayfish (*Pacifastasus*

leniusculus) at higher densities facilitate their dominance over native species in the communities

they invade (Pintor et al. 2009). We found that exploration and activity were strongly correlated

in both the lab and the field, although there was a significant decrease in the strength of this

relationship in the laboratory. We expected an overall positive relationship between exploration

and activity, but note that variation in the magnitude of this relationship is possible, as evidenced

by our findings. Though additional work is needed, the strength of this relationship could reflect

the commonly recovered activity:exploration behavioral syndrome (for a review in frogs see

Kelleher et al. 2018). Neither exploration nor activity were correlated with time to emerge, our

measure of boldness. However, because emergence probability was overall low, especially in

the laboratory, it is difficult to draw conclusions. Additionally, commonly used measures of

boldness are not always correlated with one another (Carter et al. 2013; Yuen et al. 2017), and

here too, additional work is needed. In sum, our findings here provide starting points for future

work characterizing additional behaviors and the same behaviors in alternative assays.

In addition to correlations among behaviors, correlations between behavior and

physiology also play a role in invasive success (Candler and Bernal 2015; Damas-Moreira et al.

2019). Coquí in Puerto Rico and Hawaii show physiological variation across elevational

gradients (Chaparro 2023; Marchetti et al. 2023; Marchetti et al. 2024) and in response to

manipulations in the lab (Haggerty 2016; Marchetti et al. 2023). Therefore, we were interested in

whether and how such physiological differences might be associated with behavioral variation.

We measured resting metabolic rate (RMR) after acclimation to the laboratory to ask whether

we could still detect differences based on population of origin and whether RMR was associated

with behavior. We found a trend toward an interaction of density and collection site elevation on

RMR following acclimation to laboratory conditions. Notably, these effects mirrored behavioral

findings, where the effects of density were modulated by elevation. As we did not measure RMR

in the field, we cannot say whether RMR differed among site types in the field. However, we

also found a positive association between RMR and exploration and activity in the laboratory.

Taken together with those of others, our findings here demonstrate associations between

metabolic rate and behavior, and suggest that both change plastically in response to

environmental conditions across the invasive range of coqui in Hawaii.

Conclusions

Our findings highlight the relationships between behavioral traits, physiological factors, and

environmentally induced changes in an impressive invader. Collectively, our findings underscore

the ability of coquí to thrive in diverse ecological conditions and suggest that behavioral and

physiological plasticity play a role in their success as invaders. This study contributes to the

growing body of research on the role of behavior in invasive potential.

Acknowledgements

We thank the members of the Fischer Laboratory for their help with frog care, input on data

analysis, and feedback on previous versions of the manuscript. For their assistance with

equipment and technical support, we thank Dr. Jen Moss and Reed Crawford, as well as the

Illinois Natural History Survey and the University of Illinois Urbana Champaign (UIUC) College

of Agricultural, Consumer, and Environmental Sciences for providing technical equipment

funding. We thank Michael R. Britton for advice on metabolic rate data collection and analysis.

We thank Jack Marchetti, Drs. Karen H. Beard and Susannah S. French, and residents and

employees of the Hawaii Division of Wildlife for their assistance with navigating the permitting

process and identifying collection sites. Our work would not be possible without the support of

the Division of Animal Resources at UIUC.

Funding

This work was supported by the United States Department of Agriculture National Institute of

Food and Agriculture Hatch project 1026333 (ILLU-875-984 to K.M.S); a University of Illinois

Graduate College Master's Fellowship (to K.M.S); a University of Illinois Graduate College

Travel Award (to K.M.S); Illinois State Toll Highway Authority funding (to D.E and A.L.C); and

University of Illinois Laboratory Start-up funds (to E.K.F).

Data Availability

Data analyzed in the current manuscript are included as supplemental materials.

Author Contributions

Conceptualization: KMS and EKF. Methodology: KMS, DE, ALC, and FOH. Formal analysis:

KMS and EKF. Investigation: KMS, DE, ALC, and FOH. Data curation: KMS and EKF. Writing –

Original draft: KMS and EKF. Writing – Review & Editing: KMS, DE, ALC, FOH, and EKF.

Visualization: KMS and EKF. Project administration: EKF. Funding acquisition: KMS and EKF.

References

Alcaraz C, Vila-Gispert A, García-Berthou E. 2005. Profiling invasive fish species: The
importance of phylogeny and human use. Divers Distrib. 11(4). doi:10.1111/j.1366-
9516.2005.00170.x.

Alex Perkins T, Phillips BL, Baskett ML, Hastings A. 2013. Evolution of dispersal and life history
interact to drive accelerating spread of an invasive species. Ecol Lett. 16(8).
doi:10.1111/ele.12136.

Baškiera S, Gvoždík L. 2022. Individual Variation in Thermal Reaction Norms Reveals
Metabolic-Behavioral Relationships in an Ectotherm. Front Ecol Evol. 10.
doi:10.3389/fevo.2022.850941.

Bates D, Mächler M, Zurich E, Bolker BM, Walker SC. 2011. Fitting linear mixed-effects models
using lme4. J Stat Softw.

Beard KH, Al-Chokhachy R, Tuttle NC, O'Neill EM. 2008. Population density estimates and
growth rates of *Eleutherodactylus coqui* in Hawaii. J Herpetol. 42(4). doi:10.1670/07-
314R1.1.

Beard KH, Pitt WC. 2005. Potential consequences of the coqui frog invasion in Hawaii. Divers
Distrib. 11(5). doi:10.1111/j.1366-9516.2005.00178.x.

Beard KH, Pitt WC. 2012. *Eleutherodactylus coqui* Thomas (Caribbean tree frog). In: A
Handbook of Global Freshwater Invasive Species.

Beard KH, Price EA, Pitt WC. 2009. Biology and impacts of pacific island invasive species. 5.
Eleutherodactylus coqui, the coqui frog (anura: Leptodactylidae). Pacific Sci. 63(3).
doi:10.2984/049.063.0301.

Ben-Shachar M, Lüdecke D, Makowski D. 2020. effectsize: Estimation of Effect Size Indices
and Standardized Parameters. J Open Source Softw. 5(56). doi:10.21105/joss.02815.

Benevides FL, Mautz WJ, Jacobsen CJ, Hara AH. 2019. Estimating density of calling male
Eleutherodactylus coqui in Hawaii from audio recordings of the nighttime frog chorus.

Bioacoustics. 28(2). doi:10.1080/09524622.2017.1399467.

Biro PA, Stamps JA. 2010. Do consistent individual differences in metabolic rate promote

consistent individual differences in behavior? Trends Ecol Evol. 25(11).

doi:10.1016/j.tree.2010.08.003.

Burton T, Killen SS, Armstrong JD, Metcalfe NB. 2011. What causes intraspecific variation in

resting metabolic rate and what are its ecological consequences? Proc R Soc B Biol Sci.

278(1724). doi:10.1098/rspb.2011.1778.

Candler S, Bernal XE. 2015. Differences in neophobia between cane toads from introduced and

native populations. Behav Ecol. 26(1). doi:10.1093/beheco/aru162.

Carter AJ, Feeney WE, Marshall HH, Cowlishaw G, Heinsohn R. 2013. Animal personality:

What are behavioural ecologists measuring? Biol Rev. 88(2). doi:10.1111/brv.12007.

Champley S. 2020. pwr: Basic Functions for Power Analysis. R Packag version 13-0.

Chaparro R. 2023. Characterizing Metabolic Responses of *Eleutherodactylus* Frogs in Puerto

Rico to Different Thermal Treatments: Implications for Conservation and Management.

North Carolina State University.

Chapple DG, Simmonds SM, Wong BBM. 2012. Can behavioral and personality traits influence

the success of unintentional species introductions? Trends Ecol Evol. 27(1).

doi:10.1016/j.tree.2011.09.010.

Chuang A, Peterson CR. 2016. Expanding population edges: Theories, traits, and trade-offs.

Glob Chang Biol. 22(2). doi:10.1111/gcb.13107.

Damas-Moreira I, Riley JL, Harris DJ, Whiting MJ. 2019. Can behaviour explain invasion

success? A comparison between sympatric invasive and native lizards. Anim Behav. 151.

doi:10.1016/j.anbehav.2019.03.008.

Garamszegi LZ, Markó G, Herczeg G. 2013. A meta-analysis of correlated behaviors with

implications for behavioral syndromes: Relationships between particular behavioral traits.

Behav Ecol. 24(5). doi:10.1093/beheco/art033.

Haggerty J. 2016. Thermal tolerance of the common coquí frog (*Eleutherodactylus coqui*) in

East Hawaii along an elevation gradient. University of Hawaii at Hilo.

Hudina S, Hock K. 2012. Behavioural determinants of agonistic success in invasive crayfish.

Behav Processes. 91(1). doi:10.1016/j.beproc.2012.05.011.

Jarić I, Heger T, Castro Monzon F, Jeschke JM, Kowarik I, McConkey KR, Pyšek P, Sagouis A,

Essl F. 2019. Crypticity in Biological Invasions. Trends Ecol Evol. 34(4).

doi:10.1016/j.tree.2018.12.008.

Kassambara A, Kosinski M, Biecek P. 2020. survminer: Drawing Survival Curves using

"ggplot2". R package version 0.4.8. https://cran.r-project.org/web/packages/survminer/.

Kelleher SR, Silla AJ, Byrne PG. 2018. Animal personality and behavioral syndromes in

amphibians: a review of the evidence, experimental approaches, and implications for

conservation. Behav Ecol Sociobiol. 72(5). doi:10.1007/s00265-018-2493-7.

Kraus F, Campbell EW. 2002. Human-mediated escalation of a formerly eradicable problem:

The invasion of Caribbean frogs in the Hawaiian Islands. Biol Invasions. 4(3).

doi:10.1023/A:1020909205908.

Kumschick S, Bacher S, Dawson W, Heikkilä J, Sendek A, Pluess T, Robinson T, Kühn I. 2012.

A conceptual framework for prioritization of invasive alien species for management

according to their impact. NeoBiota. 15. doi:10.3897/neobiota.15.3323.

Lagos ME, White CR, Marshall DJ. 2017. Do invasive species live faster? Mass-specific

metabolic rate depends on growth form and invasion status. Funct Ecol. 31(11).

doi:10.1111/1365-2435.12913.

Lennox R, Choi K, Harrison PM, Paterson JE, Peat TB, Ward TD, Cooke SJ. 2015. Improving

science-based invasive species management with physiological knowledge, concepts, and

tools. Biol Invasions. 17(8). doi:10.1007/s10530-015-0884-5.

Lenth RV, Bolker B, Buerkner P, Giné-Vázquez L, Herve M, Jung M, Love J, Miguez F, Riebl H,

Singmann H. 2023. Package "emmeans." Am Stat. 34(4).

Magnhagen C, Wacker S, Forsgren E, Myhre LC, Espy E, Amundsen T. 2014. Context

consistency and seasonal variation in boldness of male two-spotted gobies. PLoS One.

9(3). doi:10.1371/journal.pone.0093354.

Marchetti JR, Beard KH. 2021. Predator-Prey Reunion: Non-native Coquí Frogs Avoid Their

Native Predators. Ichthyol Herpetol. 109(3). doi:10.1643/h2020146.

Marchetti JR, Beard KH, Virgin EE, Lewis EL, Hess SC, Ki KC, Sermersheim LO, Furtado AP,

French SS. 2023. Invasive frogs show persistent physiological differences to elevation and

acclimate to colder temperatures. J Therm Biol. 114. doi:10.1016/j.jtherbio.2023.103590.

Marchetti JR, French SS, Virgin EE, Lewis EL, Ki KC, Sermersheim LO, Brusch GA, Beard KH.

2024. Invading nonnative frogs use different microhabitats and change physiology along

an elevation gradient. J Exp Zool Part A Ecol Integr Physiol. 341(1):73–85.

doi:10.1002/jez.2762.

Mathot KJ, Dingemanse NJ, Nakagawa S. 2019. The covariance between metabolic rate and

behaviour varies across behaviours and thermal types: meta-analytic insights. Biol Rev.

94(3). doi:10.1111/brv.12491.

Mowery MA, Vink C, Mason AC, Andrade MCB. 2021. Behavioural, morphological, and life

history shifts during invasive spread. Biol Invasions. 23(11). doi:10.1007/s10530-021-

02593-6.

Myles-Gonzalez E, Burness G, Yavno S, Rooke A, Fox MG. 2015. To boldly go where no goby

has gone before: Boldness, dispersal tendency, and metabolism at the invasion front.

Behav Ecol. 26(4). doi:10.1093/beheco/arv050.

O'Neill EM, Beard KH, Fox CW. 2018. Body Size and Life History Traits in Native and

Introduced Populations of Coqui Frogs. Copeia. 106(1). doi:10.1643/CE-17-642.

Peacock MM, Beard KH, O'Neill EM, Kirchoff VS, Peters MB. 2009. Strong founder effects and

low genetic diversity in introduced populations of Coqui frogs. Mol Ecol. 18(17).

doi:10.1111/j.1365-294X.2009.04308.x.

Pintor LM, Sih A, Kerby JL. 2009. Behavioral correlations provide a mechanism for explaining

high invader densities and increased impacts on native prey. Ecology. 90(3).

doi:10.1890/08-0552.1.

R-Core-Team. 2022. R: A language and environment for statistical computing. Vienna, Austria,

http://www.R-project.org

Rollins-Smith LA, Reinert LK, Burrowes PA. 2015. Coqui frogs persist with the deadly chytrid

fungus despite a lack of defensive antimicrobial peptides. Dis Aquat Organ. 113(1).

doi:10.3354/dao02823.

RStudio-Team. 2021. RStudio: Integrated Development Environment for R. RStudio, PBC,

Boston, MA, http://www.rstudio.com

Sakai AK, Allendorf FW, Holt JS, Lodge DM, Molofsky J, With KA, Baughman S, Cabin RJ,

Cohen JE, Ellstrand NC, et al. 2001. The population biology of invasive species. Annu Rev

Ecol Syst. 32. doi:10.1146/annurev.ecolsys.32.081501.114037.

Smith RL, Beard KH, Koons DN. 2018. Invasive coqui frogs are associated with greater

abundances of nonnative birds in Hawaii, USA. Condor. 120(1). doi:10.1650/CONDOR-

17-109.1.

Sol D, Lefebvre L. 2000. Behavioural flexibility predicts invasion success in birds introduced to

New Zealand. Oikos. 90(3). doi:10.1034/j.1600-0706.2000.900317.x.

Therneau TM. 2021. survival: A Package for Survival Analysis in R. R Packag version 238.

Tsutsui ND, Suarez A V., Holway DA, Case TJ. 2000. Reduced genetic variation and the

success of an invasive species. Proc Natl Acad Sci U S A. 97(11).

doi:10.1073/pnas.100110397.

Weis JS. 2010. The role of behavior in the success of invasive crustaceans. Mar Freshw Behav

Physiol. 43(2). doi:10.1080/10236244.2010.480838.

Westrick SE, Laslo M, Fischer EK. 2022. The big potential of the small frog *Eleutherodactylus*

coqui. Elife. 11:e73401. doi:10.7554/eLife.

Wright TF, Eberhard JR, Hobson EA, Avery ML, Russello MA. 2010. Behavioral flexibility and

species invasions: The adaptive flexibility hypothesis. Ethol Ecol Evol. 22(4).

doi:10.1080/03949370.2010.505580.

Yuen CH, Schoepf I, Schradin C, Pillay N. 2017. Boldness: are open field and startle tests

measuring the same personality trait? Anim Behav. 128.

doi:10.1016/j.anbehav.2017.04.009.

TABLES

640   **Table 1.** Model outputs for behavioral and resting metabolic rate (RMR) differences in the field and the laboratory.

	FIELD						LAB							
	exploration		activity		emergence		exploration		activity		emergence		RMR	
	$F_{1,37}$	p	$F_{1,36}$	p	X^2	p	$F_{1,52}$	p	$F_{1,52}$	p	X^2	p	$F_{1,50}$	p
density	4.91	0.033	3.43	0.072	0.02	0.891	0.00	0.989	0.08	0.772	0.06	0.812	0.74	0.393
elevation	5.38	0.026	2.83	0.101	3.29	0.070	1.51	0.225	0.57	0.453	3.08	0.079	0.32	0.571
density*elevation	6.26	0.017	3.38	0.074	2.19	0.140	0.03	0.874	0.07	0.794	1.28	0.259	3.01	0.090

643   **Table 2.** Model outputs for comparison of behavior in the field versus the laboratory.

	exploration		activity		emergence	
	F	p	F	p	X^2	p
trial	5.41	0.023	7.54	0.008	3.01	0.083
density	0.22	0.640	0.27	0.061	0.00	0.941
elevation	0.02	0.890	0.00	0.954	2.17	0.141
trial*density	0.06	0.813	0.50	0.483	0.06	0.813
trial*elevation	3.52	0.067	1.26	0.268	0.00	0.999
density*elevation	3.46	0.069	2.15	0.148	1.50	0.220
trial*density*elevation	5.11	0.029	3.80	0.057	0.00	0.999

645   **Table 3.** Post hoc comparison of within site type behavioral differences in the lab versus field.

	FIELD vs LAB			
	exploration		activity	
	t	p	t	p
high density / low elevation	2.97	0.005	2.46	0.016
high density / high elevation	-1.18	0.245	-0.61	0.549
low density / low elevation	1.07	0.291	1.25	0.219
low density / high elevation	1.36	0.178	1.99	0.051

Figure 1. Behavior across high- and low- density (D) and elevational (E) populations in the field. Key findings include (a) a significant interaction of density and elevation on exploration, (b) a trend toward statistical significance on activity, and (c) no statistically significant differences on probability to emerge (boldness). Complete statistical results are in Table 1.

Figure 2. Behavioral changes from the field to the laboratory. We observed no behavioral differences based on population of origin density or elevation after acclimation to the laboratory. However, we found overall decreased (a) exploration and (b) activity, and (c) a lower propensity to emerge in the laboratory as compared to the field. For exploration and activity, this shift was particularly pronounced in high density, low elevation populations. Lines in (a) and (b) connect field and laboratory observations for the same individuals.

Figure 3. Activity and exploration are strongly correlated. While we expected a positive correlation between activity and exploration

due to assay design, these metrics are not perfectly correlated, and we tested for variation between populations or origin and

between field and laboratory trials. While we found strong, significant correlations under both conditions, the strength of the

relationship was significantly different in the field (tau=0.82; grey line) and the laboratory (tau=0.71; black line). Activity (# line

crosses; x-axis) is shown on a square root adjusted scale to improve separation and visualization of points at the low end.

Figure 4. Resting metabolic rate (RMR) and behavior. (a) The interaction between population of origin density and elevation on RMR

trended toward statistical significance (Table 1) but all post hoc pairwise comparisons between site types were non-significant. There

were moderate, significant correlations between RMR and (b) exploration (tau=0.27) and (c) activity (tau=0.25) (behavioral data from

laboratory trials only). RMR is represented as the volume of CO_2 produced per minute per gram of wet mass (uL CO_2/min/g).