Warm temperature suppresses plant systemic acquired resistance by intercepting the *N*-hydroxypipecolic acid immune pathway

Authors: Alyssa Shields[1], Lingya Yao[2], Jong Hum Kim[3,4,5], Wasan Mudher Abo Al-Timmen[1,6], Sha Li[7], Eric J. R. Marchetta[1], Vanessa Shivnauth[1], Tao Chen[7], Sheng Yang He[3,4], Xiufang Xin[2], and Christian Danve M. Castroverde[1,*]

Affiliations:

[1]Department of Biology, Wilfrid Laurier University, Waterloo, Ontario, Canada N2L 3C5

[2]National Key Laboratory of Plant Molecular Genetics, CAS Center for Excellence in Molecular Plant Sciences, Institute of Plant Physiology and Ecology, Chinese Academy of Sciences, Shanghai 200032, China

[3]Department of Biology, Duke University, Durham, North Carolina, USA 27708

[4]Howard Hughes Medical Institute, Duke University, Durham, North Carolina, USA 27708

[5]Department of Life Sciences, Pohang University of Science and Technology, Pohang University of Science and Technology, Pohang 37673, Republic of Korea

[6]Department of Biology, College of Science, University of Babylon, Babylon, Iraq

[7]State Key Laboratory of Agriculture Microbiology, Huazhong Agricultural University, Wuhan, China

*Corresponding Author: dcastroverde@wlu.ca (ORCID: 0000-0002-9982-8451)

Running Title: Temperature regulation of plant SAR

Abstract

Climate warming influences disease development by targeting critical components of the plant immune system, including pattern-triggered immunity (PTI), effector-triggered immunity (ETI) and production of the central defence hormone salicylic acid (SA) at the primary pathogen infection site. However, it is not clear if and/or how temperature impacts systemic immunity. Here we show that pathogen-triggered systemic acquired resistance (SAR) in *Arabidopsis thaliana* is suppressed at elevated temperature. This was accompanied by global downregulation of SAR-induced genes at elevated temperature. Abolished SAR under warmer conditions was associated with reduced biosynthesis of the SAR metabolite *N*-hydroxypipecolic acid (NHP) in *Arabidopsis* and other plant species, as demonstrated by downregulation of NHP biosynthetic genes (*ALD1* and *FMO1*) and NHP precursor pipecolic acid (Pip) levels. Although multiple SAR signals have been shown previously, exogenous Pip was sufficient to restore disease protection at elevated temperature, indicating that heat-mediated SAR suppression is due to Pip-NHP downregulation. Along with *ALD1* and *FMO1*, systemic expression of the SA biosynthetic gene *ICS1* was also suppressed at warm temperature. Finally, we define a transcriptional network controlling thermosensitive NHP pathway via the master transcription factors CBP60g and SARD1. Our findings demonstrate that warm temperatures impact not only local but also systemic immunity by impinging on the NHP pathway, providing a roadmap towards engineering climate-resilient plant immune systems.

Keywords: Climate change, N-hydroxypipecolic acid, plant hormone, plant immunity, systemic acquired resistance, temperature

Main Text

Warming global temperatures due to climate change pose serious threats to the natural environment and human civilization (Altizer et al., 2016; Deutsch et al., 2018; Cavicchioli et al., 2019; Delgado-Baquerizo et al., 2020). In agricultural systems, increasing average temperatures are predicted to significantly reduce yields of the world's major crops (Zhao et al., 2017; Chaloner et al., 2021; Singh et al., 2023). As many crop varieties are bred to be grown outside the native growth ranges of their wild relatives, it is important to understand the impact

of elevated temperatures on plant growth, development, immunity and phenology. Together with

other environmental factors, temperature profoundly affects various aspects of plant physiology,

ranging from growth and development (Quint et al., 2016; Lippman et al., 2018; Hayes et al.,

2020; Vu et al., 2021; Castroverde and Dina, 2021) to stress responses and immunity

(Velasquez et al., 2018; Kim et al., 2021; Castroverde and Dina, 2021; Singh et al., 2023).

Environmentally compromised immune signaling is associated with increased plant disease

development, as postulated as the plant disease triangle (Stevens, 1960; Colhoun, 1973;

Roussin-Léveillée et al., 2023).

Elevated temperatures target various components of the plant immune system (Velasquez

et al., 2018; Cheng et al., 2019; Cohen and Leach, 2020; Kim et al., 2021), which encompass

both cell surface (transmembrane) and intracellular immune receptors (Jones and Dangl, 2006;

Zhou and Zhang, 2020; Kim and Castroverde, 2020; Ngou et al., 2022). Cell surface immune

receptors recognize apoplastic immunogenic molecules, typically conserved pathogen-

associated molecular patterns (PAMPs; Bigeard et al., 2015; Gust et al., 2017; DeFalco and

Zipfel, 2021), while intracellular nucleotide-binding leucine-rich receptors (NLRs) perceive

pathogen effectors and/or effector-induced host modifications (Cui et al., 2015; Jones et al.,

2016; Saur et al., 2021). There is evidence that warm temperatures lead to reduced abundance,

membrane localization or signaling of cell surface receptors (De Jong et al., 2002; Janda et al.,

2019; Hilleary et al., 2020) and perturbed nuclear localization or activation of certain NLRs (Zhu

et al., 2010; Mang et al., 2012; Cheng et al., 2013; Venkatesh and Kang, 2019).

Recent studies show that both PRR signaling and NLR signaling activate convergent

downstream signaling, including increased cytoplasmic calcium influx, oxidative burst,

phosphorylation cascades and production of defence hormone salicylic acid (SA), leading to

overlapping defence gene expression and metabolism at the primary infection site (Pieterse et

al., 2009; Ngou et al., 2021; Yuan et al., 2021). Previous research has shown that elevated

temperature suppresses pathogen-induced biosynthesis of the central defence hormone

salicylic acid (SA) (Malamy et al., 1992; Huot et al., 2017; Li et al., 2021; Kim et al., 2022; Rossi

et al., 2023), which aligns with temperature modulation of other plant hormones (Franklin et al.,

2011; Sun et al., 2012; Ibañez et al., 2018; Martínez et al., 2018; Ferrero et al., 2019; Bruessow

et al., 2021; Castroverde and Dina, 2021). Apart from resistance at the local infection site, SA is

also involved in systemic immunity known as systemic acquired resistance (SAR) (Fu and

Dong, 2013; Zhang and Li, 2019; Kachroo and Kachroo, 2020; Lim et al., 2020; Peng et al.,

2021). Primary infection and immune activation lead to SAR, which primes non-infected

systemic tissues against secondary infections (Cameron et al., 1994; Maldonado et al., 2002;

Fu and Dong, 2013; Yu et al., 2013; Vlot et al., 2021; Zeier, 2021). SAR is widely conserved

across the plant kingdom, leading to broad-spectrum host defences by rapidly generating

mobile signals at the infection site and transporting them throughout the plant (Carella et al.,

2016; Chen et al., 2018; Hartmann et al., 2018; Kachroo and Kachroo, 2020; Shine et al., 2022).

A major SAR signal is the immune-activating metabolite *N*-hydroxypipecolic acid (NHP)

(Chen et al., 2018; Hartmann and Zeier, 2018; Hartmann et al., 2018). NHP has been shown to

induce SAR, and blocking its biosynthesis results in SAR abolishment (Song et al., 2004;

Mishina and Zeier, 2006; Jing et al., 2011; Navarova et al., 2012; Huang et al., 2020). NHP

biosynthesis follows three consecutive enzymatic reactions (Hartmann and Zeier, 2018),

involving the aminotransferase AGD2-LIKE DEFENSE RESPONSE PROTEIN 1 (ALD1; Zeier,

2013; Bernsdorff et al., 2016), the reductase SAR-DEFICIENT 4 (SARD4; Ding et al., 2016;

Hartmann et al., 2017) and *N*-hydroxylase FLAVIN-DEPENDENT MONOOXYGENASE 1

(FMO1), which converts pipecolic acid (Pip) to *N*-hydroxypipecolic acid (NHP; Chen et al., 2018;

Hartmann et al., 2018). *ALD1*, *SARD4* and *FMO1* gene expression and NHP levels are induced

systemically following pathogenic attack at a primary site (Song et al., 2004; Mishina and Zeier,

2006; Ding et al., 2016; Bernsdorff et al., 2016; Hartmann et al., 2017; Yildiz et al., 2021).

Currently, the impact of elevated temperature on plant systemic immunity is

underexplored. Although previous studies have shown that increased temperatures have a

negative effect on several aspects of the plant immune system that occur in the local/primary

sites of infection (like PTI, ETI and SA production), it is unknown if and how elevated

temperature affect NHP-mediated plant immunity and the SAR response. In our study, we show

that elevated temperature suppresses SAR-associated NHP biosynthetic gene expression and

Pip production. Because it has been shown that the NHP immune pathway is redundantly

controlled by the master transcription factors CALMODULIN-BINDING PROTEIN 60-LIKE G

(CBP60g) and SAR-DEFICIENT 1 (SARD1) (Sun et al., 2015; Sun et al., 2018; Huang et al.,

2020), we further found that warm temperature-suppressed plant systemic immunity is restored

by constitutive *CBP60g/SARD1* gene expression or exogenous Pip application. Collectively, our

study indicates that CBP60g and SARD1 control the temperature-vulnerability of *Arabidopsis*

systemic immunity by regulating Pip/NHP biosynthesis.

Results

Warm temperature affects systemic acquired resistance in *Arabidopsis* plants

To determine if SAR is affected by temperature, lower leaves of *Arabidopsis* Col-0 plants were inoculated with the virulent bacterial pathogen *Pst* DC3000 as the primary challenge. Two days after primary infection, upper systemic leaves were inoculated with the pathogen *Pst* DC3000 as the secondary challenge. As shown in Figure 1a, primary pathogen infection expectedly lowered the systemic bacterial counts after secondary pathogen challenge at 23°C but not at 28°C in Col-0 plants. This indicates that *Arabidopsis* SAR due to local *Pst* DC3000 infection is effective at normal but not at elevated temperature, suggesting that temperature regulates plant systemic immunity.

Fig. 1 *Arabidopsis* SAR is suppressed at elevated temperature.

(A, B) Lower leaves of four-week-old *Arabidopsis* Col-0 plants were infiltrated with 0.25 mM $MgCl_2$ (mock) and *Pst*
DC3000 (OD600 = 0.02) in (A) or *Pst* DC3000/AvrRpt2 (OD600 = 0.02) in (B). Plants were then incubated at either
23°C or 28°C. Two days after primary local inoculation, upper systemic leaves were infiltrated with *Pst* DC3000
(OD600 = 0.001), and plants were incubated again at their respective temperatures (23°C or 28°C). Bacterial
numbers (upper panels) and symptom expression photos (lower panels) were taken at 3 days post-inoculation (dpi)
of systemic tissues. Data show the mean log CFU *Pst* DC3000/mL (± S.D.) and individual points (n=16 from 4
independent experiments) analyzed with two-way ANOVA and Tukey's Multiple Comparisons test. Statistical
differences of means are denoted by different letters. (C) Transcriptome analysis using SAR+ genes from Hartmann
et al. (2018) interfaced with temperature-regulated genes from Kim et al. (2022). The number of SAR+ genes that are
downregulated and upregulated at elevated temperature are shown (fold change cutoff > 2).

In addition to virulent pathogens, avirulent pathogens that induce local ETI can also

induce SAR (Cameron et al., 1994; Zeier, 2021). We therefore tested whether SAR activated by

the avirulent strain *Pst* DC3000/AvrRpt2, which activates RPS2-dependent immunity in Col-0

plants (Kunkel et al., 1993; Bent et al., 1994), is also impacted by warm conditions. As

expected, SAR was induced at normal temperature (23°C) after primary *Pst* DC3000/AvrRpt2

infection (Figure 1b). Strikingly, systemic *Pst* DC3000/AvrRpt2 levels remained similar

compared to mock treatment at elevated temperature (28°C), indicating that even ETI-induced

SAR protection is also negatively affected in *Arabidopsis* Col-0 plants at higher temperature.

To further understand global SAR immune signaling, we analyzed a previously generated

Arabidopsis SAR transcriptome after virulent *P. syringae* infection (Hartmann et al., 2018). We

interfaced the SAR-induced (SAR+) genes from that study with our previously published

temperature-regulated transcriptome (Kim et al., 2022) and categorized *Arabidopsis* SAR+

genes into downregulated or upregulated genes at elevated temperature. As shown in Figure

1c, 1037 SAR+ genes were downregulated, while 151 SAR+ genes were upregulated at

elevated temperatures. This demonstrates that a significant majority of temperature-regulated

SAR genes exhibit downregulated expression at elevated temperature, consistent with loss of

systemic protection against secondary infection at 28°C (Figures 1a-b). Overall, our collective

results indicate the *Arabidopsis* SAR induced by both virulent and avirulent pathogens are

negatively impacted by warm temperatures.

The *N*-hydroxypipecolic acid pathway is downregulated by elevated temperature

To determine the mechanism of how temperature regulates systemic immunity, we

measured expression levels of genes required for biosynthesis of NHP, which is an immune-

activating metabolite crucial for SAR (Hartmann et al., 2018; Chen et al., 2018). As shown in Figure 2a, transcript levels of *ALD1* and *FMO1* in local (primary) leaves of *Arabidopsis* Col-0 plants at 1 day post-infection with *Pst* DC3000 were lower at 28°C than at 23°C. In agreement, pathogen-induced levels of the NHP precursor metabolite Pip were also lower at the warmer temperature (Figure 2b). Similar trends in warm temperature-suppression of NHP biosynthetic gene expression was observed in tomato and rapeseed plants (Figures 2c-d). Finally, as shown in Figure 2e, systemic (uninfected) leaves of *Arabidopsis* plants at 2 dpi exhibited lower *ALD1* and *FMO1* gene expression levels at 28°C compared to those at 23°C. Taken together, these results indicate that elevated temperature impacts NHP biosynthesis in various plants species, which is associated with loss of SAR at higher temperatures.

Fig. 2 Plant NHP biosynthetic gene expression and Pip production are suppressed at elevated temperature.

(A-B) Leaves of four-week-old *Arabidopsis* Col-0 plants were infiltrated with 0.25 mM MgCl₂ (mock) or *Pst* DC3000 (OD600 = 0.001). Plants in were then incubated at either 23°C or 28°C. (A) *ALD1* and *FMO1* transcript levels and (B) Pip levels of pathogen-inoculated tissues were measured at 1 day post-inoculation (dpi). (C) Leaves of four-week-old tomato cultivar Castlemart plants were infiltrated with 0.25 mM MgCl₂ (mock) or *Pst* DC3000 (OD600 = 0.001). *SlALD1* transcript levels of pathogen-inoculated tissues were measured at 1 dpi. (D) Leaves of 4- to 5-week-old rapeseed cultivar Westar plants were infiltrated with 0.25 mM MgCl₂ (mock) or *Pst* DC3000 (OD600 = 0.0001). *BnaALD1* and *BnaFMO1* transcript levels of pathogen-inoculated tissues were measured at 1 dpi. (E) Lower leaves of four-week-old *Arabidopsis* Col-0 plants were infiltrated with 0.25 mM MgCl₂ (mock) or *Pst* DC3000 (OD600 = 0.02). Plants in were then incubated at either 23°C or 28°C. *ALD1* and *FMO1* transcript levels in upper systemic tissues were measured at 2 dpi. Data show the means (± S.D.) and individual points (n=3 to 4) analyzed with two-way ANOVA and Tukey's Multiple Comparisons test. Statistical differences of means are denoted by different letters. Experiments were performed at least two times with reproducible results.

Because SA is functionally linked and mutually amplified with Pip and NHP during plant

systemic immunity (Hartmann and Zeier, 2019; Huang et al., 2020; Zeier, 2021; Shields et al.,

2022), we also measured SA biosynthetic gene expression in upper, systemic leaves after

primary pathogen challenge. As shown in Figures 3a and 3b, systemic expression of the SA

biosynthetic gene *ICS1* (Wildermuth et al., 2001) was induced at 2 days after virulent *Pst*

DC3000 or avirulent *Pst* DC3000/AvrRpt2 inoculation at 23°C. However, *ICS1* transcript levels

in systemic tissues were comparable between mock and pathogen treatments at 28°C (Figures

3a-b), suggesting that elevated temperature downregulates SA accumulation systemically.

Altogether, these results demonstrate that the SA biosynthetic pathway is suppressed at higher

temperature similarly as NHP. This suggests that mutual amplification of these two important

SAR-associated metabolites is negatively affected when temperatures increase.

Fig. 3 Systemic SA biosynthetic gene expression is suppressed at elevated temperature.

Lower leaves of four-week-old *Arabidopsis* Col-0 plants were infiltrated with 0.25 mM MgCl$_2$ (mock), *Pst* DC3000 (OD600 = 0.02) or *Pst* DC3000 AvrRpt2 (OD600=0.02). Plants were then incubated at either 23°C or 28°C. (A-B) *ICS1* transcript levels were measured in systemic tissues using RT-qPCR at 2 days post-inoculation (dpi) with *Pst* DC3000 or *Pst* DC3000 AvrRpt2, respectively. (C-D) *CBP60g* and *SARD1* transcript levels were measured in systemic tissues at 2 dpi with *Pst* DC3000. Data show the means (± S.D.) and individual points (n=4) analyzed with two-way ANOVA and Tukey's Multiple Comparisons test. Statistical differences of means are denoted by different letters. Experiments were performed two times with reproducible results.

Local or systemic application of exogenous Pip restores *Arabidopsis* immune priming at elevated temperature

Having observed that Pip levels were suppressed at elevated temperature, we hypothesized that exogenous supplementation with Pip may restore disease protection at 28°C if Pip production is the rate-limiting step at elevated temperature. We infiltrated leaves of *Arabidopsis* plants with mock or 1 mM Pip and then infected the same leaves with *Pst* DC3000. As shown in Figure 4a, pathogen levels were reduced after Pip treatment compared to mock treatment at 23°C, expectedly indicating that Pip is sufficient to induce immune priming. Remarkably, Pip-induced disease protection was maintained at 28°C (Figure 4a), which confirms that Pip-NHP production is the rate-limiting step in SAR at elevated temperature.

Fig. 4 Exogenous Pip treatment restores *Arabidopsis* immune priming at warm temperature.

Four-week-old *Arabidopsis* Col-0 plants were treated with mock or 1mM Pip solution by leaf-infiltration (A) or root-drenching (B). Plants were then incubated at either 23°C or 28°C. Two days after Pip treatment, leaves were infiltrated with *Pst* DC3000 (OD600 = 0.001), and plants were incubated again at their respective temperatures (23°C or 28°C). Bacterial numbers were quantified at 3 days post-inoculation (dpi). Data show the mean log CFU *Pst* DC3000/mL (± S.D.) and individual points (n=12 from 3 independent experiments in A; n=24 from 6 independent

To determine if elevated temperature also affects systemic Pip-mediated disease

protection, we irrigated *Arabidopsis* roots with mock or 1 mM Pip two days before infiltrating the

leaves with *Pst* DC3000. Similar to the results with Pip-induced disease protection by leaf

infiltration, we also observed reduced pathogen levels after Pip root irrigation at both 23°C and

28°C (Figure 4b). These results indicate that the temperature-suppression of the Pip-NHP

pathway is governed at the level of biosynthesis and not systemic transport.

***CBP60g* and *SARD1* control temperature-vulnerability of the NHP-SAR pathway**

In addition to chemical supplementation, we next investigated if we could restore

systemic immune responses genetically at warm temperatures. We recently showed that

CBP60g and its functionally redundant paralog SARD1 control the temperature-sensitivity of

plant basal resistance and pathogen-induced SA biosynthesis (Kim et al., 2022). We therefore

investigated whether CBP60g and SARD1 also control temperature-sensitive systemic defence

responses. As shown in Figure 3c-d, *CBP60g* and *SARD1* transcript levels are induced

systemically after local *Pst* DC3000 challenge at 23°C, but this systemic induction is lost at

28°C. These suggest that the loss of systemic immunity at elevated temperature could be due to

significantly decreased *CBP60g* and *SARD1* gene expression.

To show a causative role for CBP60g/SARD1 downregulation with NHP immune

pathway suppression, we used plants constitutively expressing *CBP60g* (*35S::CBP60g*) or

SARD1 (*35S::SARD1*). As shown in Figures a-c, *35S::CBP60g* lines restored NHP biosynthetic

gene expression (*ALD1, FMO1*) and Pip levels at 28°C in contrast to the wild-type Col-0 plants.

Consistent with this, constitutive expression of the functionally redundant *SARD1* gene in

35S::SARD1 plants also led to restored expression of *ALD1* and *FMO1* at warm temperature

(Figure 5d-e). Taken together, a major mechanism by which higher temperatures target plant

systemic immunity and NHP biosynthesis is through the expression of the master immune

transcription factor genes *CBP60g* and *SARD1*.

Fig. 5 CBP60g and SARD1 control the temperature-vulnerability of NHP-mediated immunity.

Leaves of four-week-old *Arabidopsis* Col-0 and *35S::CBP60g* (A-C) or *35S::SARD1* plants (D-E) were infiltrated with 0.25 mM $MgCl_2$ (mock) or *Pst* DC3000 (OD600 = 0.001). Plants were then incubated at either 23°C or 28°C. (A) *ALD1* and (B) *FMO1* transcript levels were measured using RT-qPCR at 1 day post-inoculation (dpi) of pathogen-inoculated tissues of Col-0 and *35S::CBP60g* plants. (C) Pip levels were measured at 1 dpi in the same tissues. (D) *ALD1* and (E) *FMO1* transcript levels were measured at 1 day post-inoculation (dpi) of pathogen-inoculated tissues of Col-0 and *35S::SARD1* plants. Data show the means (± S.D.) and individual points (n=4 in A-C; n=3 in D-E), with experiments performed three times with reproducible results. Statistical analyses were conducted using two-way ANOVA with Tukey's Multiple Comparisons test, with statistically significant differences of means denoted by different letters.

Discussion

In this study, we showed that pathogen-induced SAR and the SAR-activating NHP pathway in *Arabidopsis* plants are sensitive to elevated temperatures. Local infection with both virulent or ETI-activating avirulent pathogens triggers SAR at normal temperatures but not at elevated temperatures. We showed that SAR regulation by temperature is caused by temperature-sensitive NHP pathway in local and systemic tissues. SAR-induced NHP biosynthetic gene expression and/or Pip levels are downregulated at elevated temperature. Even though multiple SAR signals have been proposed previously (Fu and Dong, 2013;

Návarová, et al., 2012; Hartmann et al., 2018; Wang, et al., 2018; Vlot et al., 2021; Zeier, 2021),

we show here that exogenous supplementation with the NHP precursor Pip (locally and

systemically) was sufficient to restore SAR at elevated temperature. This demonstrates a

causative relationship between temperature-suppressed Pip-NHP pathway and SAR.

Because NHP primes systemic SA biosynthesis and immunity (Yildiz et al., 2021; Zeier,

2021), we also observed that systemic SA biosynthetic gene expression is downregulated by

elevated temperature. SA induction is also important for SAR establishment since SA induction

deficient *sid2* and SA-insensitive *npr1* mutants also have abolished SAR (Fu and Dong, 2013;

Huang et al., 2020; Peng et al., 2021). Temperature-suppressed systemic SA pathway agrees

with our previous study showing a negative impact of higher temperatures on local pathogen-

induced SA biosynthesis and basal resistance (Huot et al., 2017; Kim et al., 2022). However, we

found that Pip supplementation alone can rescue SAR at elevated temperature, suggesting that

Pip-NHP signaling is sufficient in inducing SAR.

We previously showed that CBP60g and SARD1 controls the temperature-vulnerability of

local pathogen-induced SA biosynthesis (Kim et al., 2022). Functionally redundant paralogs

CBP60g and SARD1 are master transcription factors that directly target the promoters of

numerous immunity-related genes (Wang et al., 2009; Zhang et al., 2010; Wang et al., 2011;

Sun et al., 2015, Sun et al., 2018), including those important for SA biosynthesis (like *ICS1*) and

NHP biosynthesis (like *ALD1* and *FMO1*). In this study, we further demonstrate that CBP60g

and SARD1 control the temperature-regulation of NHP biosynthesis and systemic immunity.

Expression of *CBP60g* and *SARD1* in systemic tissues are downregulated at elevated

temperature, which is associated with suppressed NHP pathway. Restoring systemic *CBP60g*

or *SARD1* expression using constitutively expressing *35S::CBP60g* or *35S::SARD1* plants

rescues NHP biosynthetic gene expression and Pip levels under warming conditions.

Overall, changing climatic factors like elevated temperature have a broad and significant

impact on the plant immune system, not only at local sites of infection (PTI, ETI, SA) but also on

systemic immune priming in distal sites via the central SAR metabolite NHP. In this study, we

determined that elevated temperature regulates SAR by influencing NHP pathway. We further

demonstrated that this temperature-regulation of NHP-mediated SAR is also controlled by the

temperature-sensitive master regulators CBP60g and SARD1, reminiscent of their roles in local

immune responses (Kim et al., 2022). Not only does temperature affect the plant's ability to

directly defend against pathogen attacks but also the plant's immune preparedness for future

infections. Our discoveries advance our understanding of how the plant immune landscape is regulated by a changing environment. This foundational mechanistic knowledge of the plant disease triangle is important to inform strategies in mitigating the negative impacts of warming temperatures on plant health and to provide a molecular roadmap towards engineering climate-resilient plants.

Materials and Methods

Plant materials and growth conditions

Arabidopsis thaliana Columbia-0 (Col-0), *35S::CBP60g* (Wan et al., 2012) and *35S::SARD1* seeds (Kim et al., 2022) were surface-sterilized with 70% ethanol for 10 minutes and rinsed with autoclaved water. Seeds were suspended in sterile 0.1% agarose and incubated in cold conditions (4°C) for three days. Seeds were sown into pots containing autoclaved soil mix – one-part Promix-PGX soil (Plant Products, Ancaster, Ontario), one-part Turface (Turface Athletics, Buffalo Grove, IL), and one-part Vermiculite Prolite (Therm-O-Rock, New Eagle, PA) – supplemented with 100mL of Miracle-Gro solution (The Scotts Company, Mississauga, ON). Tomato (*Solanum lycopersicum*) cultivar Castlemart plants were grown as described previously (Shivnauth et al., 2023). Rapeseed (*Brassica napus*) cultivar Westar plants were grown as described previously (Kim et al., 2022). Plants were grown in environmentally controlled chambers at 23°C with 60% relative humidity and 12h light [$100 \pm 20\ \mu mol\ m^{-2}\ s^{-1}$]/12 dark cycle.

Plant systemic immunity and SAR assays

Four-week-old *Arabidopsis* plants were covered with plastic domes to increase humidity and open the stomata 24 hours before infiltration, Plants were infiltrated with 0.25 mM MgCl$_2$ (mock), *Pst* DC3000 (OD600=0.02) or *Pst* DC3000/AvrRpt2 (OD600=0.02) in their lower leaves (Xin et al., 2013; Xin et al., 2018). Following infiltration, plants were further incubated in environmentally growth chambers at either 23°C or 28°C with identical relative humidity (60%) and lighting conditions (12h light/12 dark). For SAR disease assays, upper systemic leaves were or infiltrated with *Pst* DC3000 (OD600 = 0.001). Bacterial levels were quantified 3 days after systemic infiltration based on a previously published protocol (Huot et al., 2017; Kim et al.,

2022). Briefly, in planta bacterial extracts were plated on rifampicin-containing LM media and

log colony forming units (CFUs) cm^{-2} were calculated.

Pip treatment and protection assay

Four-week-old *Arabidopsis* plants were infiltrated with mock solution or 1 mM Pip (Sigma)

in their lower leaves. In parallel to direct Pip infiltration into leaves, plants were irrigated with 1

mL of mock solution or 1 mM Pip (Sigma) for root inoculations. Following infiltration or irrigation,

plants were further incubated in environmentally growth chambers at either 23°C or 28°C with

identical relative humidity (60%) and lighting conditions (12h light/12 dark). After two days, the

Pip-treated leaves (for local Pip infiltration) or upper systemic leaves (for root irrigation) were

further infiltrated with *Pst* DC3000 (OD600 = 0.001). Bacterial levels were quantified 3 days

after systemic infiltration as stated in the previous section.

Gene expression analyses

Pathogen-infected leaves and upper systemic leaves at either 23°C or 28°C were

harvested at 1 or 2 days after local infection, respectively. Tissues were flash-frozen in liquid

nitrogen and stored at -80°C before total RNA extraction. Gene expression levels were

quantified based on a previously published protocol (Huot et al., 2017; Kim et al., 2022) with

slight modifications. RNA was extracted from flash-frozen plant tissues using the Qiagen Plant

RNeasy Mini Kit (Qiagen, Toronto, ON) or TRIzol Reagent (Aidlab Biotech, China) according to

the manufacturer's protocol. Resulting cDNA was synthesized using qScript cDNA super mix

(Quantabio) or M-MLV reverse transcriptase (RT, Vazyme Biotech, China) based on

manufacturers' recommendations. Real-time quantitative polymerase chain reaction (qPCR)

was performed using PowerTrack SYBR Green master mix (Life Technologies) or iTaq™

Universal SYBR® Green Supermix (Bio-Rad, USA) with approximately 1.5-10 ng of template

cDNA. Equivalently diluted mRNA without the qScript cDNA mix were used as negative

controls. The resulting qPCR mixes were run using the Applied Biosystems QuantStudio3

platform (Life Technologies) or CFX96 Real-Time PCR Detection System (Bio-Rad, USA). The

individual Ct values were determined for target genes and the internal control gene: *PP2AA3* for

Arabidopsis; *SlACT2* for tomato and *BnaGDI1* for rapeseed (Huot et al., 2017; Kim et al., 2022;

Shivnauth et al., 2023). Gene expression values were reported as $2^{-\Delta Ct}$, where ΔCt is Ct_{target}

$_{gene}$–Ct$_{internal\ control\ gene}$. qPCR was carried out with three technical replicates for each biological sample. Primers used for qPCR are shown in Supplementary Table 1.

Pip metabolite extraction and quantification

Pathogen-infected leaves at 23°C or 28°C were harvested at 1 day after infection. Pip extraction and quantification were performed based on previous reports with slight modifications (Yao et al., 2023). Approximately 100 mg leaf tissue was frozen and ground in liquid nitrogen, and Pip was extracted at 4℃ for 1h in 600 μL ice-cold extraction buffer (80% methanol in water, 0.1 g/L butylated hydroxytoluene). The extraction step was repeated twice, and a 1.2-mL supernatant was speed-dried in a vacuum centrifugal concentrator (Beijing JM Technology). The pellet was resuspended in 240 μL 30% methanol solution and diluted 10 times for quantification. Pip level was quantified using the AB SCIEX QTARP 5500 LC/MS/MS system. Selected ion monitoring (SIM) was conducted in the positive ES channel for Pip (m/z 130.0>84.0), which was done using a 20V collision energy and a 50V declustering potential. The instrument control and data acquisition were performed using Analyst 1.6.3 software (AB SCIEX), and data processing was performed using MultiQuant 3.0.2 software (AB SCIEX). Pip was separated with an ACQUITY UPLC BEH Amide column (1.7 μm, 2.0 x 100 mm, Waters) using the method in Supplementary Table 2. Pip levels were quantified by calculating the area of each individual peak and comparing it to standard curves. Reported Pip concentration was normalized by sample fresh weight (FW) in gram.

Reference List

1. Altizer, S., Ostfeld, R.S., Johnson, P.T., Kutz, S., Harvell, C.D. (2013) Climate change and infectious diseases: from evidence to a predictive framework. *Science.* 341(6145):514-9.
2. Bent, A.F., Kunkel, B.N., Dahlbeck, D., Brown, K.L., Schmidt, R., Giraudat, J., Leung, J., Staskawicz, B.J. (1994). RPS2 of Arabidopsis thaliana: a leucine-rich repeat class of plant disease resistance genes. *Science.* 265(5180):1856-60.
3. Bernsdorff, F., Döring, A.-C., Gruner, K., Schuck, S., Bräutigam, A., Zeier, J. (2016). Pipecolic Acid Orchestrates Plant Systemic Acquired Resistance and Defence Priming via Salicylic Acid-Dependent and -Independent Pathways. *The Plant Cell.* 28(1): 102-129.

4. Bigeard J., Colcombet J., Hirt H. (2015). Signaling Mechanisms in Pattern-Triggered
Immunity (PTI). *Mol. Plant.* 8, 521–539.
5. Bruessow, F. et al. (2021) Natural variation in temperaturemodulated immunity uncovers
transcription factor bHLH059 as a thermoresponsive regulator in Arabidopsis thaliana. *PLoS*
Genet. 17, e1009290
6. Cameron, R.K., Dixon, R.A., Lamb, C.J. (1994) Biologically induced systemic acquired
resistance in Arabidopsis thaliana. *Plant J.* 5: 715-725.
7. Carella, P., Merl-Pham, J., Wilson, D.C., Dey, S., Hauck, S.M., Vlot, A.C., Cameron, R.K.
(2016). Comparative Proteomics Analysis of Phloem Exudates Collected during the
Induction of Systemic Acquired Resistance. *Plant Physiol.* 171(2):1495-510.
8. Castroverde, C. D. M., Dina, D. (2021). Temperature regulation of plant hormone signaling
during stress and development. *J. Exp. Bot.* 72, 7436–7458.
9. Cavicchioli, R., Ripple, W.J., Timmis, K.N. et al. (2019). Scientists' warning to humanity:
microorganisms and climate change. *Nat. Rev. Microbiol.* 17, 569–586.
10. Chaloner, T.M. et al. (2021) Plant pathogen infection risk tracks global crop yields under
climate change. *Nat. Clim. Chang.* 11, 710–715
11. Chen, Y.C., Holmes, E.C., Rajniak, J., Kim, J.G., Tang, S., Fischer, C.R., Mudgett, M.B.,
Sattely, E.S. (2018). N-hydroxy-pipecolic acid is a mobile metabolite that induces systemic
disease resistance in Arabidopsis. *Proc. Natl. Acad. Sci.* A. 115(21):E4920-E4929.
12. Cheng, C., Gao, X., Feng, B. Sheen, J., Shan, L., He, P. (2013). Plant immune response to
pathogens differs with changing temperatures. *Nat Commun* 4, 2530.
13. Cheng, Y.T., Zhang, L., He, S.Y. (2019). Plant-Microbe Interactions Facing Environmental
Challenge. *Cell Host Microbe.* 26(2):183-192.
14. Cohen, S.P., Leach, J.E. (2020). High temperature-induced plant disease susceptibility:
more than the sum of its parts. *Curr. Opin. Plant Biol.* 56, 235–241.
15. Colhoun, J. (1973). Effects of environmental factors on plant disease. *Annu. Rev.*
Phytopathol. 11, 343–364.
16. Cui, H., Tsuda, K., Parker, J.E. (2015). Effector-triggered immunity: from pathogen
perception to robust defense. *Annu. Rev. Plant Biol.* 66:487-511.
17. De Jong, C.F., Takken, F.L., Cai, X., de Wit, P.J., Joosten, M.H. (2002) Attenuation of Cf-
mediated defense responses at elevated temperatures correlates with a decrease in elicitor-
binding sites. *Mol. Plant Microbe Interact.* 15(10):1040-9.
18. DeFalco, T. A., Zipfel, C. (2021). Molecular mechanisms of early plant pattern-triggered
immune signaling. *Mol. Cell* 81, 3449–3467.

19. Delgado-Baquerizo, M., Guerra, C.A., Cano-Díaz, C. et al. (2020) The proportion of soil-
borne pathogens increases with warming at the global scale. *Nat. Clim. Chang.* 10, 550–
554.

20. Deutsch, C.A. et al. (2018) Increase in crop losses to insect pests in a warming climate.
Science 361, 916–919.

21. Ding, P., Rekhter, D., Ding, Y., Feussner, K., Busta, L., Haroth, S., Xu, S., Li, X., Jetter, R.,
Feussner, I., Zhang, Y. (2016). Characterization of a pipecolic acid biosynthesis pathway
required for systemic acquired resistance. *Plant Cell.* 28(10):2603-2615.

22. Ferrero, V., Viola, I.L., Ariel, F.D., Gonzalez, D.H. (2019). Class I TCP transcription factors
target the gibberellin biosynthesis gene GA20ox1 and the growth-promoting genes HBI1
and PRE6 during thermomorphogenic growth in Arabidopsis. *Plant Cell Physiol.* 60, 1633–
1645.

23. Franklin, K.A., Lee, S.H., Patel, D. et al. (2011). Phytochrome-interacting factor 4 (PIF4)
regulates auxin biosynthesis at high temperature. *Proc. Natl. Acad. Sci.* 108, 20231–20235.

24. Fu, Z. Q., Dong, X. (2013). Systemic acquired resistance: turning local infection into global
defense. *Annu. Rev. Plant Biol.* 64, 839–863.

25. Gust, A.A., Pruitt, R., Murnberger, T. (2017). Sensing danger: key to activating plant
immunity. *Trends Plant Sci.* 22: 779-791.

26. Hartmann, M., Zeier, J. (2018). L-lysine metabolism to N-hydroxypipecolic acid: an integral
immune-activating pathway in plants. *Plant J.* 96: 5-21.

27. Hartmann, M., Zeier, J. (2019). N-hydroxypipecolic acid and salicylic acid: a metabolic duo
for systemic acquired resistance. *Curr. Opin. Plant Biol.* 50, 44–57.

28. Hartmann, M., Kim, D., Bernsdorff, F., Ajami-Rashidi, Z., Scholten, N., Schreiber, S., Zeier,
T., Schuck, S., Reichel-Deland, V., Zeier, J. (2017). Biochemical principles and functional
aspects of pipecolic acid biosynthesis in plant immunity. *Plant Physiol.* 174(1), 124–153.

29. Hartmann, M., Zeier, T., Bernsdorff, F., Reichel-Deland, V., Kim, D., Hohmann, M.,
Schloten, N., Schuck, S., Bräutigam, A., Hölzel, T., Ganter. C., Zeier, J. (2018). Flavin
Monooxygenase-Generated N-Hydroxypipecolic Acid Is a Critical Element of Plant Systemic
Immunity. *Cell* 173(2), 456-469.e16.

30. Hayes, S., Schachtschabel, J., Mishkind, M., Munnik, T., Arisz, S.A. (2020). Hot topic:
Thermosensing in plants. *Plant Cell Environ.* 44(7): 2018–2033.

31. Hilleary R, Paez-Valencia J, Vens C, Toyota M, Palmgren M, Gilroy S. (2020). Tonoplast-
localized Ca2+ pumps regulate Ca2+ signals during pattern-triggered immunity in
Arabidopsis thaliana. *Proc. Natl. Acad. Sci.* 117(31):18849-18857.

32. Huang, W., Wang, Y.., Li, X., Zhang, Y. (2020). Biosynthesis and Regulation of Salicylic
Acid and N-Hydroxypipecolic Acid in Plant Immunity. *Mol. Plant* 13: 31-41.
33. Huot, B., Castroverde, C. D. M., Velásquez, A. C., Hubbard, E., Pulman, J. A., Yao, J.,
Childs, K.L., Tsuda, K., Montgomery, L.B., He, S. Y. (2017). Dual impact of elevated
temperature on plant defence and bacterial virulence in Arabidopsis. *Nat. Commun. 8*(1),
1808.
34. Ibañez, C., Delker, C., Martinez, C., et al. (2018). Brassinosteroids dominate hormonal
regulation of plant thermomorphogenesis via BZR1. *Curr. Biol.* 28, 303–310.e3.
35. Janda, M., Lamparova, L., Zubıkova, A., Burketova, L., Martinec, J., Krckova, Z. (2019).
Temporary heat stress suppresses PAMP-triggered immunity and resistance to bacteria in
Arabidopsis thaliana. *Mol. Plant Pathol*, 20:1005-1012.
36. Jing, B., Xu, S., Xu, M., Li, Y., Li, S., Ding, J., Zhang, Y. (2011). Brush and spray: a high-
throughput systemic acquired resistance assay suitable for large-scale genetic screening.
Plant Physiol. 157:973–980.
37. Jones, J. D. G., Dangl, J. L. (2006). The plant immune system. *Nature* 444, 323–329.
38. Jones, J.D., Vance, R.E., Dangl, J.L. (2016). Intracellular innate immune surveillance
devices in plants and animals. *Science.* 354(6316):aaf6395.
39. Kachroo A, Kachroo P. (2020). Mobile signals in systemic acquired resistance. *Curr. Opin.*
Plant Biol. 58:41-47.
40. Kim, J.H. Castroverde, C.D.M. (2020). Diversity, Function and Regulation of Cell Surface
and Intracellular Immune Receptors in Solanaceae. *Plants* 9(4):434.
41. Kim, J.H., Hilleary, R., Seroka, A., He, S.Y. (2021). Crops of the future: building a climate-
resilient plant immune system. *Curr. Opin. Plant Biol.* 60, 101997.
42. Kim, J.H., Castroverde, C.D.M., Huang, S. et al. (2022). Increasing the resilience of plant
immunity to a warming climate. *Nature* 607(7918):339-344.
43. Kunkel, B.N., Bent, A.F., Dahlbeck, D., Innes, R.W., Staskawicz, B.J. (1993). RPS2, an
Arabidopsis disease resistance locus specifying recognition of Pseudomonas syringae
strains expressing the avirulence gene avrRpt2. *Plant Cell.* 5(8):865-75.
44. Li, Z., Liu, H., Ding, Z., Yan, J., Yu, H., Pan, R., Hu, J., Guan, Y., Hua, J. (2020). Low
temperature enhances plant immunity via salicylic acid pathway genes that are repressed by
ethylene. *Plant Physiol.* 182, 626–639.
45. Lim, G.H., Liu, H., Yu, K., Liu, R., Shine, M.B., Fernandez, J., Burch-Smith, T., Mobley, J.K.,
McLetchie, N., Kachroo, A., Kachroo, P. (2020). The plant cuticle regulates apoplastic
transport of salicylic acid during systemic acquired resistance. *Sci. Adv.* 6: eaaz0478.

46. Lippmann, R., Babben, S., Menger, A., Delker, C., Quint, M. (2019). Development of wild

and cultivated plants under global warming conditions. *Curr. Biol.* 29, R1326–R1338.

47. Malamy, J., Hennig, J., Klessig, D.F. (1992). Temperature-dependent induction of salicylic

acid and its conjugates during the resistance response to tobacco mosaic virus infection.

The Plant Cell 4, 359–366.

48. Maldonado, A.M., Doerner, P., Dixon, R.A., Lamb, C.J., Cameron, R.K. (2002). A putative

lipid transfer protein involved in systemic resistance signalling in Arabidopsis. *Nature.*

419(6905):399-403.

49. Mang, H.G., Qian, W., Zhu, Y., Qian, J., Kang, H.G., Klessig, D.F., Hua, J. (2012). Abscisic

acid deficiency antagonizes high-temperature inhibition of disease resistance through

enhancing nuclear accumulation of resistance proteins SNC1 and RPS4 in Arabidopsis. *The*

Plant Cell 24, 1271–1284.

50. Martínez, C., Espinosa-Ruíz, A., de Lucas, M., Bernardo-García, S., Franco-Zorrilla, J.M.,

Prat, S. (2018.) PIF4-induced BR synthesis is critical to diurnal and thermomorphogenic

growth. *EMBO J.* 37, e99552.

51. Mishina, T. E., Zeier, J. (2006). The Arabidopsis flavin-dependent monooxygenase FMO1 is

an essential component of biologically induced systemic acquired resistance. *Plant Physiol.*

141(4), 1666–1675.

52. Návarová, H., Bernsdorff, F., Döring, A. C., Zeier, J. (2012). Pipecolic acid, an endogenous

mediator of defence amplification and priming, is a critical regulator of inducible plant

immunity. *Plant Cell*, 24(12), 5123–5141.

53. Ngou, B.P.M., Ahn, H.K., Ding, P., Jones, J.D.G. (2021). Mutual potentiation of plant

immunity by cell-surface and intracellular receptors. *Nature.* 592(7852):110-115.

54. Ngou, B.P.M., Ding, P., Jones, J.D.G. (2022). Thirty years of resistance: Zig-zag through the

plant immune system. *Plant Cell.* 34(5):1447-1478.

55. Peng, Y. et al. (2021) Salicylic acid: biosynthesis and signalling. *Annu. Rev. Plant Biol.* 72,

761–791

56. Pieterse, C. M. J., Leon-Reyes, A., Van Der Ent, S., Van Wees, S. C. M. (2009).

Networking by small-molecule hormones in plant immunity. *Nat. Chem. Biol.* 5(5):308-16.

57. Quint, M., Delker, C., Franklin, K. et al. (2016). Molecular and genetic control of plant

thermomorphogenesis. *Nat. Plants* 2, 15190.

58. Rossi, C.A.M., Marchetta, E.J.R., Kim, J.H., Castroverde, C.D.M. (2023). Molecular

regulation of the salicylic acid hormone pathway in plants under changing environmental

conditions. *Trends Biochem. Sci.* 48(8):699-712.

59. Roussin-Léveillée, C., Rossi, C.A.M., Castroverde, C. D. M. Moffett, P. 2023. The Plant

disease triangle facing climate change. *Trends Plant Sci.* (under review).

60. Saur, I.M.L., Panstruga, R., Schulze-Lefert, P. (2021). NOD-like receptor-mediated plant

immunity: from structure to cell death. *Nat. Rev. Immunol.* 21(5):305-318.

61. Shields, A., Shivnauth, V., Castroverde, C.D.M. (2022) Salicylic acid and N-hydroxypipecolic

acid at the fulcrum of the plant immunity-growth equilibrium. *Front. Plant Sci.* 13, 841688

62. Shine, M.B., Zhang, K., Liu, H., Lim, G.H., Xia, F., Yu, K., Hunt, A.G., Kachroo, A., Kachroo,

P. (2022). Phased small RNA-mediated systemic signaling in plants. *Sci. Adv.*

8(25):eabm8791.

63. Shivnauth, V., Pretheepkumar, S., Marchetta, E.J.R., Rossi, C.A.M., Amani, K.,

Castroverde, C.D.M. (2023). Structural diversity and stress regulation of the plant immunity-

associated CALMODULIN-BINDING PROTEIN 60 (CBP60) family of transcription factors in

Solanum lycopersicum (tomato). Funct. Integr. Genomics 23(3):236.

64. Singh, B.K., Delgado-Baquerizo, M., Egidi, E. et al. (2023). Climate change impacts on plant

pathogens, food security and paths forward. Nat. Rev. Microbiol. 21(10):640-656.

65. Song, J.T., Lu, H., McDowell, J.M., Greenberg, J.T. (2004) A key role for *ALD1* in activation

of local and systemic defences in Arabidopsis. *Plant J.* 40: 200-212.

66. Stevens, R.B. (1960). Pages 357-429 in: *Plant Pathology, an Advanced Treatise*, Vol. 3.

J.G. Horsfall and A.E. Dimond, eds. Academic Press, NY.

67. Sun, J., Qi, L., Li, Y., Chu, J., Li, C. (2012). PIF4-mediated activation of YUCCA8 expression

integrates temperature into the auxin pathway in regulating Arabidopsis hypocotyl growth.

PLoS Genet. 8, e1002594.

68. Sun, T., Zhang, Y., Li, Y., Zhang, Q., Ding, Y., Zhang, Y. (2015) ChIP-seq reveals broad

roles of SARD1 and CBP60g in regulating plant immunity. *Nat. Commun.* 6:10159.

69. Sun, T., Busta, L., Zhang, Q., Ding, P., Jetter, R., Zhang, Y. (2018) TGACG-BINDING

FACTOR 1 (TGA1) and TGA4 regulate salicylic acid and pipecolic acid biosynthesis by

modulating the expression of SYSTEMIC ACQUIRED RESISTANCE DEFICIENT 1

(SARD1) and CALMODULIN-BINDING PROTEIN 60g (CBP60g). *New Phytol.* 217:344–354.

70. Velásquez, A. C., Castroverde, C. D. M., He, S. Y. (2018). Plant-Pathogen Warfare under

Changing Climate Conditions. *Curr. Biol.* 28(10):R619-R634.

71. Venkatesh, J., Kang, B-C. (2019). Current views on temperature-modulated *R* gene-

mediated plant defense responses and tradeoffs between plant growth and immunity. *Curr.*

Opin. Plant Biol. 50: 9-17.

72. Vlot, A.C., Sales, J.H., Lenk, M., Bauer, K., Brambilla, A., Sommer, A., Chen, Y., Wenigand,
M., Nayem, S. (2021). Systemic Propagation of Immunity in Plants. *New Phytol.* 229:1234–
1250.

73. Vu, L.D., Xu, X., Zhu, T., et al. (2021). The membrane-localized protein kinase
MAP4K4/TOT3 regulates thermomorphogenesis. *Nat. Commun.* 12(1):2842.

74. Wan, D., Li, R., Zou, B., Zhang, X., Cong, J., Wang, R., Xia, Y., Li, G. (2012). Calmodulin-
binding protein CBP60g is a positive regulator of both disease resistance and drought
tolerance in Arabidopsis. *Plant Cell Rep.* 31(7):1269–1281.

75. Wang, L., Tsuda, K., Sato, M., Cohen, J.D., Katagiri, F., Glazebrook, J. (2009). Arabidopsis
CaM binding protein CBP60g contributes to MAMP-induced SA accumulation and is
involved in disease resistance against Pseudomonas syringae. *PLoS Pathog.* 5:e1000301.

76. Wang, L., Tsuda, K., Truman, W., Sato, M., Nguyen, L.V., Katagiri, F., Glazebrook, J.
(2011). CBP60g and SARD1 play partially redundant critical roles in salicylic acid signaling.
Plant J. 67:1029–1041.

77. Wang, C., Liu, R., Lim, G. H., De Lorenzo, L., Yu, K., Zhang, K., Hunt, A.G., Kachroo, A.,
Kachroo, P. (2018). Pipecolic acid confers systemic immunity by regulating free radicals.
Sci. Adv. 4(5), 4509.

78. Wildermuth, M. C., Dewdney, J., Wu, G., Ausubel, F. M. (2001). Isochorismate synthase is
required to synthesize salicylic acid for plant defence. *Nature*, 414(6863), 562–565.

79. Xin, X. F., Kvitko, B., He, S. Y. (2018). Pseudomonas syringae: What it takes to be a
pathogen. *Nat. Rev. Microbiol.* 16, 316–328.

80. Xin, X.F., He, S.Y. (2013). Pseudomonas syringae pv. tomato DC3000: A Model Pathogen
for Probing Disease Susceptibility and Hormone Signaling in Plants. *Annu. Rev.*
Phytopathol. 51: 473-498.

81. Yao, L., Jiang, Z., Wang, Y., Hu, Y., Hao, G., Zhong, W., Wan, S., Xin, X.F. (2023). High air
humidity dampens salicylic acid pathway and NPR1 function to promote plant disease.
EMBO J. 20:e113499.

82. Yildiz, I., Mantz, M., Hartmann, M., Zeier, T., Kessel, J., Thurow, C., Gatz, C., Petzsch, P.,
Köhrer, K., Zeier, J. (2021). The mobile SAR signal N-hydroxypipecolic acid induces NPR1-
dependent transcriptional reprogramming and immune priming, *Plant Physiol.* 186(3):1679-
1705.

83. Yu, K., Soares, J. M., Mandal, M. K., Wang, C., Chanda, B., Gifford, A. N., Fowler, J.S.,
Navarre, D., Kachroo, A., Kachroo, P. (2013). A Feedback Regulatory Loop between G3P

and Lipid Transfer Proteins DIR1 and AZI1 Mediates Azelaic-Acid-Induced Systemic

Immunity. *Cell Rep. 3*(4), 1266–1278.

84. Yuan, M., Jiang, Z., Bi, G., Nomura, K., Liu, M., Wang, Y., Cai, B., Zhou, J.M., He, S.Y., Xin,

X.F. (2021). Pattern-recognition receptors are required for NLR-mediated plant immunity.

Nature. 592(7852):105-109.

85. Zeier, J. (2013). New insights into the regulation of plant immunity by amino acid metabolic

pathways. *Plant Cell Environ.* 36: 2085-2103.

86. Zeier, J. (2021). Metabolic regulation of systemic acquired resistance. *Curr. Opin. Plant Biol.*

62:102050.

87. Zhao, C., Liu, B., Piao, S., et al. (2017). Temperature increase reduces global yields of

major crops in four independent estimates. *Proc. Natl. Acad. Sci*. 114, 9326–9331.

88. Zhang, Y., Xu, S., Ding, P., Wang, D., Cheng, Y.T., He, J., Gao, M., Xu, F., Li, Y., Zhu, Z.,

Li, X., Zhang, Y. (2010) Control of salicylic acid synthesis and systemic acquired resistance

by two members of a plant-specific family of transcription factors. *Proc. Natl. Acad. Sci*.

107:18220–18225.

89. Zhang, Y., Li, X. (2019). Salicylic acid: biosynthesis, perception, and contributions to plant

immunity. *Curr. Opin. Plant Biol.* 50, 29–36.

90. Zhou, J.M., Zhang, Y. (2020). Plant Immunity: Danger Perception and Signaling. *Cell*. 181:

978-989.

91. Zhu, Y., Qian, W., Hua, J. (2010). Temperature modulates plant defense responses through

NB-LRR proteins. *PLoS Pathog*. 6, e1000844.

Acknowledgements

We thank the following colleagues for sharing biological materials: G. Howe for Castlemart
tomato, B. Staskawicz for *Pst* DC3000/AvrRpt2, and G. Li and D. Wan for *35S::CBP60g*. We
are grateful to the Laurier Faculty of Science (Max Pottier and Gena Braun), Michigan State
University (MSU) Growth Chamber Facility and Duke Phytotron for instrumentation support.
This work was supported by research funding from the NSERC Discovery Grant, Canada
Foundation for Innovation, Ontario Research Fund and Wilfrid Laurier University Faculty of
Science start-up funds. Our Laurier Biology research is located on the shared traditional territory
of the Neutral, Anishinaabe, and Haudenosaunee peoples.

Authors' contributions

C.D.M.C conceptualized and supervised the study. A.S. performed most of the experiments. L.Y.Y. performed the Pip metabolite measurements. J.H.K. performed gene expression experiments of *35S::SARD1* plants. W.M.A.A.T. and E.M. conducted experimental replicates of the SAR assays. V.S. conducted temperature experiments of tomato plants, extracted RNA and synthesized cDNA. S.L. and T.C. performed gene expression analyses of rapeseed plants. L.Y.Y., J.H.K. and S.L. were supervised by X.F.X., S.Y.H. and T.C., respectively. Everyone analyzed the data. A.S. and C.D.M.C. wrote the paper with input from all authors.