Anticipatory smooth pursuit eye movements scale with the probability of visual motion: the role of target speed and acceleration

Vanessa Carneiro Morita[1], David Souto[2], Guillaume S. Masson[1] and Anna Montagnini[*,1]

1 Institut de Neurosciences de la Timone, CNRS & Aix-Marseille Université, Marseille, France

2 School of Psychology and Vision Sciences, University of Leicester, Leicester, United Kingdom.

* Corresponding author

Acknowledgements: This work was funded by the *Agence Nationale de la Recherche* (ANR-PREDICTEYE 18-CE37--0019 to GSM, AM) and the *Fondation pour la Recherche Médicale* (Equipe FRM 2018 to GSM) and by Aix-Marseille Université (PhD Doctoral extension funding to VCM). We would like to thank Mr Alexis Ulian for helping in the data collection of the Experiment 2A as part of his Master internship.

Abstract

Sensory-motor systems can extract statistical regularities in dynamic uncertain environments, enabling quicker responses and anticipatory behavior for expected events. Anticipatory smooth pursuit eye movements (aSP) have been observed in primates when the temporal and kinematic properties of a forthcoming visual moving target are fully or partially predictable. To investigate the nature of the internal model of target kinematics underlying aSP, we tested the effect of varying the target kinematics and its predictability. Participants tracked a small visual target in a constant direction with either constant, accelerating or decelerating speed. Across experimental blocks, we manipulated the probability of each kinematic condition varying either speed or acceleration across trials; with either one kinematic condition (providing certainty) or with a mixture of conditions with a fixed probability within a block. We show that aSP is robustly modulated by target kinematics. With constant-velocity targets, aSP velocity scales linearly with target velocity in blocked sessions, and matches the probability-weighted average in the mixture sessions. Predictable target acceleration does also have an influence on aSP, suggesting that the internal model of motion which drives anticipation contains some information about the changing target kinematics, beyond the initial target speed. However, there is a large variability across participants in the precision and consistency with which this information is taken into account in order to control anticipatory behavior.

Introduction

Smooth pursuit eye movements allow us to maintain the image of a moving object of interest steady on the retinas. To do so, tracking eye movements rely primarily on the neural representation of the moving target's speed and direction (Carl & Gellman, 1987; Lisberger & Westbrook,; Tychsen & Lisberger, 1986). When the target moves at constant speed across the visual field, the eyes start accelerating in the same direction as the target motion with a short latency (~100-130 ms in humans; Carl & Gellman, 1987). In the optimal speed range for human pursuit (i.e. below 20-30°/s), the eyes typically reach a steady state velocity close to the target's velocity within ~300 ms from visual motion onset. Steady-state smooth tracking, in close coordination with the so-called catch-up saccades can dynamically maintain a good alignment between the fovea and the target retinal image position (Carl & Gellman, 1987; Orban De Xivry & Lefèvre, 2007). Even though natural objects rarely move at a constant velocity, only a few studies have investigated eye tracking behavior for accelerating (or decelerating) targets. Those studies reported that humans are indeed able to track visible targets with smoothly-varying speed (e.g. Bennett & Benguigui, 2013; Kreyenmeier et al., 2022), but with weak accuracy. Like perceptual discrimination judgments, tracking eye movements discriminate poorly between target accelerations as compared to target speeds in humans (e.g., Watamaniuk & Heinen, 2003; Kowler & McKee, 1987) and macaque monkeys (Lisberger & Movshon, 1999).

Most of the early work on pursuit eye movements and visual speed/acceleration processing were concerned with the online control. However, our dynamic environment is often, at least in part, predictable, because either the motion is produced by our own body (e.g., Gauthier et al., 1988; Landelle et al., 2016), or it can be inferred from general prior knowledge, past experience or perceptual cues about an object's motion (Kowler et al., 2019). For instance, when a target's velocity changes in a periodic way (e.g., sinusoidal motion), ocular tracking can rapidly take advantage of this predictable motion such that, after only a few cycles, the fovea is aligned to the target position with nearly no lag (Kowler & Steinman, 1979a). Moreover, motion predictability allows anticipation of the target motion onset, or target reappearance after a transient occlusion (e.g. Dodge et al., 1930; for a review, see Kowler et al., 2019 and Fukushima et al., 2013). These anticipatory smooth pursuit eye movements (aSP) are thought to help in quickly reducing the retinal position and velocity errors during the early phase of visually-guided pursuit (e.g. Kao & Morrow, 1994).

Large efforts have been devoted to understanding what kind of signals drive anticipatory responses (Kowler et al., 2019). The underlying computational mechanisms of aSP are however still not fully understood. Empirically, target motion predictability can be manipulated across different time scales (e.g. across or within trials in a standard visuomotor experiment) and repetition schedules. Our group, and others have previously shown that aSP amplitude is proportional to the probability of a given target motion direction in a direction-biased task (Damasse et al., 2018; Montagnini et al., 2010; Rubinstein et al., 2024; Santos & Kowler, 2017;

Wu et al., 2021). Expectation for a given target speed also modulates aSP:
anticipatory eye velocity increases with speed predictability, in a random versus
fully predictable speed design (e.g., Heinen et al., 2005; Jarrett & Barnes, 2002; Kao
& Morrow, 1994). aSP also depends on the recent trial history of target speed (Maus
et al. 2015). However, it has not been tested whether aSP exhibits the same linear
dependency upon speed probability as it does for direction probability.

Predictive aSP has been extensively investigated for simple target trajectories
of constant speed and/or direction (e.g., among many others, in Barnes & Asselman,
1991; Heinen et al., 2005; Kowler & Steinman, 1979b). However, much less is
known about aSP for accelerating targets and the contribution of acceleration
signals to the internal representation of motion raises several questions. First, it is
still not known if acceleration is taken into account when preparing for anticipatory
movements, or if the latter are based on simple estimates approximating the target
motion profile (e.g. instantaneous at some critical moments, or time-averaged speed,
Bennett, Orban De Xivry, et al., 2010). Second, most of the previous studies have
investigated acceleration-based predictive behavior only on a short time scale,
namely during the transient occlusion of a moving target with an accelerating
profile (Bennett et al., 2007, 2010; Bennett & Barnes, 2006).

From these previous studies, target acceleration does not appear to be fully
integrated in predictive pursuit. For instance, when an accelerating target is
occluded after a short presentation of accelerating motion (~ 200 ms), eye velocity
during the blanking period reduces, and its predictive reacceleration prior to the

target reappearance is not influenced by the target acceleration (Bennett et al., 2007). However, if the target visibility window before the occlusion is long enough (> 500 ms), eye velocity reduces during the blanking period, but recovers in an acceleration-scaled manner prior to the target reappearance (Bennett, Orban De Xivry, et al., 2010). Moreover, target acceleration is not appropriately taken into account neither in a task where participants are asked to track a target and, after a period of occlusion, to predict the target position, or to predict when the target is going to reach a certain position (Bennett & Benguigui, 2013; Kreyenmeier et al., 2022). However, one of the very few studies addressing the effect of acceleration predictability on a longer time scale (blocked design) showed that the amplitude of aSP observed before the target reappearance in an occlusion paradigm scales with target acceleration (Bennett & Barnes, 2006). Here, we aimed at better understanding how target motion acceleration shapes aSP when the predictability of the kinematic profile is manipulated on a relatively long timescale, namely across blocks of several tenths of trials.

In the present study we tested the following hypotheses: 1) Anticipatory smooth pursuit eye velocity scales linearly with target speed probability (similar to direction probability, Damasse et al. 2018, Santos et al. 2017), across experimental blocks with fixed probability; 2) Anticipatory eye velocity is modulated by target acceleration and 3) it scales with the probability of the accelerating motion. To test these hypotheses, we analyzed anticipatory oculomotor behavior for targets moving in a fully predictable direction but with different speeds and accelerations. We manipulated the probability of each kinematic condition across blocks. We showed

that target speed probability influences aSP similarly to motion direction probability, with a linear dependence of aSP velocity upon target speed probability. We also showed that aSP can be driven by predictable accelerating targets, in a way that accounts for the expected change in target velocity and for its probability, even though with large inter-individual variability.

Methods

Participants

Twenty-nine healthy adult volunteers signed an informed consent to participate in the experiments presented in this study. The experimental protocol was approved by the Ethics Committee *Comité de Protection des Personnes OUEST III* (CPP reference: PredictEye-2018-A02608-47), in full respect of the Helsinki declaration guidelines. Three of the authors (AM, GM, DS) participated in Experiment 1A (n = 3), two of the authors (AM, VCM) participated in Experiment 1B-2A (n = 13) and one of the authors (VCM) participated in Experiment 2B (n = 5) and in Experiment 3 (n = 8). In Experiment 1A, anticipatory eye movements and initial pursuit were recorded with high precision by using the scleral search coil technique (Robinson, 1963) in a small participant sample. A preliminary version of the results from Experiment 1A was presented previously at the VSS conference (Souto et al., 2008). The core finding of this experiment motivated Experiments 1B-2A-2B, that were run with a larger sample of participants, a smaller subset of probability conditions, and using a less invasive technique (video eye tracking). Experiment 3 was designed to test the effect of acceleration on anticipatory eye

150 speed in fully predictable blocks, again using a non-invasive eye recording
151 technique.

Stimuli and procedure

153   In all experiments, participants were instructed to visually track a small
154 moving target by smoothly pursuing it with their eyes, as accurately as they could,
155 while their eye movements were recorded. We used different materials across
156 experiments.

Experiment 1A

The detailed methods are described elsewhere (Wallace et al. 2005). Briefly,
a PC running the REX package controlled both stimulus presentation and data
acquisition. Stimuli were generated with an SGI Fuel workstation (ABC Corp., New
York, USA, no longer available) and back-projected onto a large translucent screen
(80° x 60°, viewing distance: 1m) using a Barco 908s video-projector (1280", 1024
pixels at 76 Hz). Oculomotor recordings were collected using the scleral search coil
technique (Collewijn et al. 1975).

This experiment probed aSP in different target speed probability contexts
(**Figure 1a**). Each trial started with a white fixation point, located at the center of
the screen for a random duration between 300 and 450 ms, on a black uniform
background (luminance <1 cd/m^2). If the participant fixated accurately (i.e. within
a 2°-side, square electronic window) during the last 200 ms, the fixation target was
extinguished and followed by a fixed-duration, 300-ms empty screen. At the end of

this gap, the target (a white, gaussian-windowed circle, 0.2° std, maximum luminance 45 cd/m^2) appeared at the center of the screen and started moving horizontally to the right, for a fixed period of 500 ms. The target speed was either 5.5 °/s (low speed, LS) or 16.5 °/s (high speed, HS). In each experimental block, a different target speed-probability condition defined the proportion of trials at high speed (P(HS)=0, 0.1, 0.25, 0.5, 0.75, 0.9 and 1). The complementary proportion of trials had a low-speed target motion (P(LS)=1-P(HS)). Participants completed 500 trials per block, except for the P(HS)=0 and P(HS)=1 conditions where only 250 trials were completed. One or two blocks were completed in a day, with the constraint of not exceeding a total of one hour of duration.

Figure 1. Experimental designs. (a). Each trial started with a fixation point displayed at the center of the screen for a random period, followed by a gap of 300 ms. The target then appeared at the center of the screen and started moving. **(b). Experiment 1.** In Experiment 1A, the target moved horizontally to the right at one of the two constant speeds (5.5 or 16.5 °/s, in blue). The probability of a high-speed (P(HS), v = 16.5 °/s) vs a low-speed

(P(LS) = 1-P(HS), v = 5.5 °/s) trial was varied between experimental blocks with 7 different values of P(HS) (0, 0.1, 0.25, 0.5, 0.75, 0.9, 1). In Experiment 1B the target always moved in a fixed direction, chosen between one of the four diagonals (counterbalanced between participants). We displayed two different constant target speeds (v11c and v33c, of 11 and 33 °/s, respectively, in blue). Conditions were designed with a constant target displacement, and we manipulated the probability of the target kinematics in each experimental block with the probability of v11c spanning the values (0, 0.3, 0.7, 1) and P(v33c) = 1-P(v11c). **(c) Experiment 2.** Experiment 2A followed a design similar to Exp 1B, but this time the target kinematics included acceleration conditions. The initial speed was kept the same as in Exp 1B, but the target with initial speed at 11°/s accelerated (v11a, a = 22 °/s^2, in green) and the target with initial speed at 33 °/s decelerated (v33d, a = -22°/s^2, in pink). In Experiment 2B, target kinematics were the same as in Exp 2A, except the time duration of the target motion was kept constant at 600 ms. **(d)** Experiment 3. We displayed three different initial speeds (v11, v22, v33 of 11, 22 and 33 °/s, respectively) combined with three different acceleration values (a, c and d, of 22, 0 and -22 °/s^2, in green, blue and pink respectively) in fully predictable blocks. Target motion duration was kept constant and lasted 500ms. **(e) Model fitting of eye velocity profiles in individual trials.** The bottom panel shows an example of the eye velocity trace (blue curve) and of the ANEMO model fit (red curve, Pasturel et al., 2018) in an individual trial. The selected model fitted an exponential function to the anticipatory phase, and a sigmoid to the initial visually-guided pursuit phase. From this model we extracted the maximum of the anticipatory velocity (aSPv). The model fitting procedure for Experiments 1B, 2A, 2B and 3 was similar to Experiment 1A.

Experiments 1B, 2 and 3

Experiment 1B-2A

The same group of participants took part in Experiments 1B and 2A. Stimuli were presented using the Psychtoolbox (Brainard, 1997) package for MATLAB. A Display++ monitor (CRS Ltd., Rochester, UK) with a refresh rate of 120 Hz was placed at 57 cm distance in front of the participant. Eye movements were recorded using an Eyelink1000, an infrared video-based eye tracker (SR Research Ltd., Ottawa, Canada). This experiment probed aSP with different target constant and varying speed conditions, while manipulating the probability of each condition.

With respect to Experiment 1A, a larger pool of participants was tested with a different set of target speed values and a smaller set of speed probabilities and fewer trials per condition. Because several studies have reported anisotropies for saccadic and smooth pursuit eye movements across different directions (e.g., Grasse & Lisberger, 1992; Ke et al., 2013; Rottach et al., 1996; Takeichi et al., 2003), we decided to investigate whether the results observed in Experiment 1A can be generalized across target motion directions. Accordingly, in Experiment 1B-2A the target moved along one of the four diagonal directions, counterbalanced between participants.

Figure 1a shows the experimental design. Each trial started with a white fixation dot in the center of the screen for a random interval between 300 ms and 600 ms. This fixation period was followed by a 300 ms gap. At the end of this period, the target (a white circle of 0.6° diameter, maximum luminance 45 cd/m^2 on a black background with luminance <1 cd/m^2) appeared at the center of the screen and started moving in one of the four diagonal directions (always the same for one participant) with different target kinematics conditions: the target speed was either constant (v11c and v33c; 11 and 33 °/s respectively), uniformly accelerating (v11a; starting from 11 °/s, a = 22 °/s^2), or decelerating (v33d; starting from 33 °/s, a = -22 °/s^2). In Experiments 1B and 2A, target motion duration was adapted to the target kinematic properties in order to achieve a similar spatial displacement on the screen across conditions: in practice target movement lasted 1 and 0.52 s for v11c, and v33c, respectively, 0.87 s for v11a, and 0.72 s for v33d.

Participants first completed 4 blocks of 100 trials with a single target kinematics condition (v11c, v33c, v11a, v33d, presented in randomized order across participants). Notice that these conditions can be considered as a probability of 1 for each kinematic value. Next, the probability of the different target kinematics was manipulated across blocks of 200 trials each. For blocks with constant speed, we used v11c and v33c, with P(v33c) being equal to either 0.3 or 0.7 and P(v11c)=1-P(v33c). The two probability levels were presented in random order across participants. For blocks with accelerating speed, P(v33d) could be either 0.3 or 0.7 (P(v11a)=1-P(v33d)). Again, these two probability conditions were randomly interleaved.

Experiment 2B

Having different motion durations across conditions might have introduced some confounds affecting anticipatory behavior. First, estimates of target acceleration can be impaired if the target motion is presented too briefly (Bennett, De Xivry, et al., 2010; Bennett et al., 2007). Second, if anticipatory behavior relies on the estimate of mean target velocity, rather than on its accelerating dynamics (Brouwer et al., 2002; Gottsdanker et al., 1961; Schmerler, 1976), then the duration of the visual motion epoch might influence the mean velocity estimate for targets with accelerating speed. Therefore, we ran an additional control experiment (Exp 2B) with the same design as Exp 2A but with one main difference: target motion duration was held constant (600ms), resulting in different target final positions across conditions.

Experiment 3

While Experiments 1 and 2 were designed to probe the effects of probability manipulation for both constant and accelerating target motion, we did not directly address whether and how the accelerating motion of predictable accelerating targets drive anticipatory eye movements. Most of the literature on smooth pursuit focus on constant speed targets, and the very few studies manipulating acceleration focuses on the effects of this manipulation on the visually-guided pursuit (Brostek et al., 2017; Krauzlis & Lisberger, 1994) or short-term predictive tracking (e.g. blanking paradigm, Bennett et al., 2007, 2010; Bennett & Barnes, 2006; Bennett & Benguigui, 2013). Interestingly, Bennett & Barnes (2006) showed that for a fixed initial target speed and when accelerating target motion is presented in a blocked design, anticipatory eye movements scale with target acceleration. In order to better understand the relationship between target acceleration and anticipatory eye movements, we ran Experiment 3 (**Figure 1d**), with a *fully-crossed* design, in which the target could take one out of three possible initial speeds (v11, v22, and v33 of 11, 22, and 33 °/s, respectively) and one out of three possible acceleration values (of 22, 0 and -22 °/s^2, labeled as "a" for *accelerating,* "c" for *constant* and "d" for *decelerating,* respectively). This design yielded in nine blocks of fully-predictable target motion. In all blocks, the target motion duration was held constant to 500 ms. Participants completed 100 trials in each block with v11 and v22 as initial speeds, and 120 trials in each block with v33 as initial speed. The small difference in the number of trials was introduced to compensate for the higher number of excluded trials for the data analysis, due to a higher occurrence of saccades around target

257 motion onset, a phenomenon already observed in the previous experiments when
258 the fastest speed (v33) was presented.

Eye-movements recording and preprocessing

For Experiment 1A, the analog voltage measure collected with the scleral coil
technique and reflecting the right-eye rotation was low-pass filtered (DC-130 Hz)
and digitized with 16-bit resolution at 1000 samples per second (to obtain the eye's
horizontal and vertical position). For Experiments 1B, 2A, 2B and 3, the right-eye
horizontal and vertical position was recorded with an infrared video-based eye
tracker, EyeLink 1000 (SR Research), at 1 kHz.

For all sets of recorded eye movements, position data was converted in an
ASCII format. After conversion, the ANEMO toolbox (Pasturel et al., 2018) and
custom-made python scripts were used to pre-process the data. Position data was
low-pass filtered (acausal second-order Butterworth low-pass filters, 30 Hz cut-off),
numerically differentiated to get the eye velocity in degrees per second and then
de-saccaded using ANEMO's implementations. In practice, saccade detection was
implemented by jointly applying the absolute eye velocity threshold criterion
(30°/s, by default in the Eyelink system) and the relative velocity threshold (akin to
the method proposed by Engbert & Kliegl, 2003). The epochs corresponding to
detected saccades were removed from analysis (given not-a-number values). Trials
with more than 40% of missing data points in the [-100,200] ms window around
the target onset for Exp 1A and more than 70% of missing data points in the
[-100,100] ms window around the target onset for Exp 1B, 2A,B and 3, or with

more than 60% of missing data points overall (for all experiments), were automatically excluded from the pre-processing pipeline. ANEMO was used to fit a piecewise linear and non-linear model to individual trials' eye velocity to extract the relevant oculomotor parameters (see **Figure 1e** for an illustration of the model fitting and extracted parameters). The model comprised a linear baseline phase (flat linear regression), and two non-linear phases: an anticipatory phase modeled by an exponential function and a visually-guided phase modeled by a sigmoid function. Importantly, we selected this model after comparing its performance quantitatively (using AIC, BIC and RMSE indicators of goodness of fit) with a more standard piecewise linear model (implemented in ANEMO): in all experiments our model (linear + non-linear model) performed better than the linear one in the large majority of trials. In Experiment 1A, the fit was performed in the time window of -300 to 350 ms relatively to the target motion onset. In Experiments 1B, 2A,B and 3, the fit was performed in the time window of -300 to 300 ms relatively to the target motion onset. The relevant model-fit parameter for this study was only the maximum velocity of the anticipatory phase (aSPv), calculated as the maximum value of the exponential function (i.e. its value at the offset of the anticipatory phase). Other oculomotor parameters are automatically extracted by the ANEMO toolbox, but we do not report them in the present study.

Eye velocity traces and model fits were visually inspected to exclude the remaining aberrant trials and those with extremely poor fits. Overall, 12% of the trials were excluded on average for Experiment 1A (median: 10%, max: 18%), 16%

for Experiment 1B-2A (median: 14 %, max: 32%), 13 % for Experiment 2B (median: 13 %, max: 15 %), and 26 % for Experiment 3 (median: 25 %, max: 33 %).

Data analysis

In all experiments, when analyzing the effects of the probability bias, the first 10 trials of each block were excluded from analysis. This was done in the aim of excluding large fluctuations related to learning the stimuli statistics and focus on average values rather than on their rate of change. For control, we also repeated all analyses excluding the first 50 trials: the results did not change significantly.

Linear Mixed Effects regression Models (*nlme* package for R) were used to evaluate the effects of the target kinematics (different constant speed and acceleration conditions) and of the kinematic probability on the anticipatory smooth pursuit velocity (aSPv) estimates. Because not including a true random effect can increase the false positive rate (Barr et al., 2013), participants were treated as a random effect and all fixed effects were allowed to vary with it. Since this approach usually leads to models that don't converge because of the high number of parameters and the correlations between them, when needed, we used the *buildmer* package for R (Voeten, 2020) to find the maximal model (i.e., the model including the most of variables) which still converges for the dependent variable. After finding the models, we fitted the data and result tables were exported with the *stargazer* package for R (Hlavac, 2022).

For Experiment 1A, the models included only speed probability as a fixed

effect for the oculomotor anticipatory velocity (aSPv). We chose to use the

probability of the highest speed, P(HS), as the independent variable in the model 1:

(1)   aSPv ~ 1 + P(HS) + (1 + P(HS) | participant)

The variable before the ~ symbol is the dependent variable, and the variables

after it are the independent variables (also called fixed effects). The 1 corresponds

to the model intercept. For the variables within the parentheses, each one before the

| symbol is allowed to vary for each level of the variable after it (also called random

effect). In other words, for each participant, the model will return a different best-fit

value for the intercept and the slope of the linear dependence upon probability. For

the analysis of the probability-mixtures of Experiment 1B % (v11c vs v33c) and

Experiment 2 (v11a vs v33d), we also added the axis (horizontal/vertical) as an

interaction factor with the probability, given that the target moved along one of the

diagonals. For Experiment 2, we included an interaction with the experiment

(2A/2B). The final models, as well as the statistics tables are presented in the

supplementary material.

It is known that recent trial-history can affect the behavior in the present trial

(sequential effect), and this could be seen as a confound in our analyses addressing

the effect of the target motion probability across several trials. Therefore, we re-ran

the regression models for Experiments 1A and 1B adding the target velocity at trial

N-1 (Tv_{N-1}) as an interaction variable:

(2) aSPv ~ 1 + P(HS)*Tv_{N-1} + (1 + P(HS) + Tv_{N-1} | participant)

In Experiment 3, considering that each target kinematic condition could be uniquely identified as a combination of an initial velocity and acceleration values, we ran a second linear mixed-effects regression model: this time the main independent variables were the target initial velocity ($v0$: 11, 22, or 33°/s) and the target acceleration (*accel*: 0, +22 and -22°/s^2), as well as their interaction, all treated as parametric variables:

(2)   *aSPv ~ 1 + v0*accel + (1 + v0 + accel | participant)*

In order to test the actual role of target acceleration this model was tested against the model including only v0 as fixed effect using the *bayestestR* package for R (Makowski et al., 2019). Pairwise comparisons between aSPv corresponding to the target kinematics (*Tk*) conditions of Experiment 3 were based on a categorical LMM, initially defined as:

(3)   aSPv ~ 1 + Tk + (1 + Tk| participant)

Contrasts between the different conditions were performed using the *emmeans* package for R (Lenth, 2017) and p-values were adjusted for multiple comparisons using the Benjamini & Hochberg method for controlling the false discovery rate.

We then tested the hypothesis that anticipatory eye velocity is driven by an estimate of the mean target velocity across a finite *temporal window of integration* (TWI) starting at target motion onset (time 0) and ending at time TWI^{end}. In order to do so we fitted, for the pooled participants' data, the following linear regression:

(4)   aSPv = slope*Tk + intercept

and extracted the individual best-fit value for the parameters *slope$_i$* and

intercept$_i$. Here, Tk included v11c, v22c, and v33c. We then used the same relation

(5) for the six conditions with accelerating targets, indexed by the suffix i, replacing

Tk by the target speed estimate (TSE$_i$), which approximates the accelerating

kinematics:

(5)   *aSPv$_i$ = slope*TSE$_i$ + intercept*

Solving with respect to TSE$_i$ we have:

(6)   TSE$_i$ = *(aSPv$_i$ - intercept)/ slope*

By imposing the equality between the estimated target speed TSE$_i$ and the mean of

the accelerating/decelerating target speed over a finite TWI, between 0 and TWI^{end},

we obtain:

(7) $(aSPv_i - intercept)/\ slope = 1/\text{TWI}^{end} \times \int_{0}^{TWIend} (v0 + a * t)\,dt$

where a is the acceleration value and v0 the initial target speed for accelerating

conditions. Finally, by solving Equation 8 with respect to TWI^{end} we could estimate,

for each acceleration condition, the final point in time of the temporal window of

integration,

(8)   TWI^{end} = 2/a × [(aSPv$_i$ - intercept)/ slope - V0]

In order to have a robust estimate of the variability of such temporal window of

integration across the population, we used the bootstrapping technique (n = 1000,

Efron, 1979) and extracted the 95% confidence interval around the mean estimated

TWI^{end}.

Results

Effects of target speed probability on anticipatory Smooth Eye Movements

Using a state-of-the-art eye movement recording technique (scleral search

coil), we first investigated the effects of target speed probability upon anticipatory

smooth eye movements. In Experiment 1A, three participants had to smoothly

pursue a target which moved rightwards along the horizontal axis with two

different possible speeds (referred to as high-speed, HS, and low-speed, LS)

randomly interleaved across trials but with a given probability of occurrence (P(HS)

and P(LS) = 1-P(HS)) within a block. **Figure 2a** shows the trial-averaged eye velocity

curves for one participant, sorted by probability and target speed conditions. Each

probability condition is represented by a different color. The two different target

speed profiles are illustrated by the horizontal dotted lines; time zero corresponds to

target motion onset. Participants were able to track the two target motions with

high accuracy, as shown by the convergence of eye velocity to the target speed

during steady-state pursuit. Since target motion direction was always rightward, we

observed a strong anticipatory response for all speed probability conditions, as

evidenced by the non-zero eye velocity at the usual pursuit latency in humans

(~100 ms). However these anticipatory responses in the predicted direction were

also modulated by target speed probability. We analyzed such modulation by

considering the relationship between amplitude of anticipatory responses and

P(HS), i.e., the probability of the highest speed (16.5 °/s). As illustrated in **Figure**

2a, higher values of P(HS) drove stronger anticipatory pursuit, regardless of the

actual target speed (16.5 ou 5.5 °/s). **Figure 2b** plots the relationship between mean

anticipatory eye velocity (aSPv) and P(HS), for the 3 participants. The gray curves

are the linear relationships estimated from the Linear Mixed Effects Model (LMM).

We found a clear linear dependency of anticipatory response upon the probability of

target speed, in the direction of target motion. The LMM statistical analysis (with

P(HS) as a fixed effect) shows that aSPv significantly increased with higher

probability (P(HS) effect: beta = 3.48, 95% CI = (3.14,3.84), $p < .001$). Overall,

anticipatory responses were stronger by ~200 %, rising from 2.5 to 7.5 °/s when

P(HS) increased from 0 to 1.

Figure 2. Experiment 1A: Dependence of anticipatory eye velocity upon target speed probability. **(a).** Average eye velocity as a function of time (+/- 95% confidence interval) for one representative participant. Trials are grouped by probability of the higher speed (HS). Each color corresponds to one probability condition. The time zero corresponds to the target onset, and the dashed lines indicate the two possible target speeds. **(b).** Anticipatory eye velocity for the group of participants. Each box plot (median in yellow, box limits corresponding to the 25% and 75% quartiles, and whiskers corresponding to 1.5 times the interquartile range, IQR) corresponds to one probability condition. The gray lines show the linear mixed model fit for each participant.

418   In a second version of this Experiment (Exp 1B), our objective was to

419 replicate the speed-probability dependency of anticipatory pursuit on a larger group

420 of 13 participants, with a less invasive eye tracking technique. We used two new

speeds (11 and 33°/s) and a reduced set of probability conditions in the mixture blocks (P(v33c) = 0, 0.3, 0.7 and 1). By using an oblique target motion direction, our second objective was to generalize this speed probability dependency across visual motion directions, by having the target moving along one of the 4 diagonal directions (**Figure 1b**). To directly compare the results of Experiment 1A and 1B, we first report the effects of target speed probability observed when varying P(v33c) – and therefore P(v11c) = 1-P(v33c) – across blocks with constant speed (v11c,v33c) mixtures. **Figure 3a** shows an example of the trial-averaged eye velocity as a function of time for one participant, sorted according to the target velocity profiles (dotted lines) and P(v33c) values. Horizontal dotted lines depict the target horizontal and vertical velocity components (i.e. 7.77 and 23.3 °/s for 11 and 33 °/s radial speed). A comparison with **Figure 2a** shows a behavior similar to Exp 1A where target motion was horizontal. Final pursuit velocity matches target velocity, especially for the lowest target speed (v11c). The fact that steady-state eye velocity gain remained lower than 1 for the fastest target speed (v33c) is consistent with previous studies (Carl & Gellman, 1987; Dodge, 1930). As in Exp 1A, target motion direction remained constant within a block, thus a robust anticipatory pursuit response was always observed. However, its amplitude increased when P(v33c) increased. Such dependency is illustrated in **Figure 3b**, where horizontal and vertical components of anticipatory eye velocity (aSPv) are plotted against P(v33c). Both components increased linearly with the probability of the highest speed. A symmetric relationship was observed with P(v11c). We ran the LMM statistical models for the anticipatory response (aSPv), including the effect of P(v33c), as for

Exp 1A. We added the effect of the eye velocity axis (horizontal or vertical) and its

interaction to test whether aSPv was differently modulated along the horizontal and

vertical dimensions. aSPv increased significantly for higher probability of P(v33c)

(**Figure 3b**, P(v33c) effect: beta = 2.74, 95% CI = (1.82, 3.66), p<.001). We did

not find a significant difference (given a criterion of alpha <0.01) between axes

(main axis effect: beta = -0.74, 95% CI = (-1.46, -0.016), p=0.046), but the effect

of P(v33c) was significantly smaller in the vertical axis (P(v33c)*axis effect: beta =

-0.67, 95% CI = (-0.98, -0.36), p<.0001). Overall, the two experiments strongly

support the fact that anticipatory eye velocity scales with the probability of target

speed, in a similar way across different motion directions in the plane, although

slightly less robustly along the vertical direction.

Figure 3. Experiment 1B: Dependence of anticipatory eye velocity upon target constant-speed probability. **(a).** Average eye velocity for trials grouped by probability of v33c and by target velocity for the (v11c,v33c) mix. The left panel shows the horizontal eye velocity, while the right panel shows the vertical eye velocity for a representative participant. Each color corresponds to the different probabilities of v33c. **(b).** Amplitude of anticipatory pursuit is plotted against P(v33c), along the horizontal (left panel) and vertical (right panel) axes. Data represented in the same way as Figure 2.

Short and long-time scale factors affecting speed expectation and eye movement anticipation

Recent trial history, that is the stimulus properties (e.g. the target speed) observed in the previous trial, or across the few previous trials, can modulate anticipatory eye movements (e.g. Heinen et al., 2005; Kowler & McKee, 1987; Kowler & Steinman, 1979; Maus et al., 2015). Importantly, several studies have also shown that both short-term factors related to one or few previous trials, and longer-term factors, related to global statistical estimates can coexist and interact to control perception and visuomotor behavior (e.g. Chopin & Mamassian, 2012; Maus et al., 2015, Damasse et al., 2018; Falmagne et al., 1975; Kowler, 1984; Pasturel et al., 2020; Wu et al., 2021). In order to quantify the effects of the previous trial's speed on the anticipatory eye velocity in the present study, as well as its interaction with the block's speed probability, we ran a new LMM, now including both the speed-probability and the target speed at the trial N-1 (Tv_{N-1}). Note that our study was not designed specifically to study sequential effects (i.e., by presenting all possible combinations of N-1, N-2, N-3,… trials), and therefore we limit our analysis of short-term effects only to the effect of the trial N-1. When the previous trial was a low-speed trial, aSPv decreased when compared to a previous high-speed trial

(Exp1A: beta = -0.50, 95% CI = (-0.66, -0.34), p<.001; Exp1B: beta = -0.61, 95%

CI = (-0.88, -0.34), p<.001). For both Experiment 1A and 1B, we found that the

main effect of target speed probability upon aSPv remained significant (Exp1A,

Tv_{N-1}=HS: beta = 4.35, 95% CI = (3.98, 4.72), p<.001; Exp1B, Tv_{N-1}=v33: beta =

1.38, 95% CI = (0.44, 2.32), p<0.01). The interaction between the previous trial's

speed and the block's speed probability was also significant, although with a

different sign: for Experiment 1A the probability effect was reduced when the

previous trial was low-speed compared to high speed (beta = -2.74, 95% CI =

(-3.10, -2.38), p<.001); in contrast, for Experiment 1B the probability effect

increased when the previous trial was at v11c compared to v33c (beta = 0.82, 95%

CI = (0.38, 1.26), p<.001). These results are illustrated in the left panels of **Figure**

4a,b. In the right panels of **Figure 4a** and **4b** we plotted, for Experiment 1A and 1B

respectively, the difference between the mean aSPv for trials following a high-speed

(in red) or a low-speed (in blue) trials and the mean aSPv across all trials in a

probability block. This illustration allows us to immediately capture how the N-1

trial's effect is modulated across the probability values: a high-speed previous trial

has a larger excitatory impact on subsequent anticipatory velocity when high-speed

trials are less frequent. The symmetric interaction is observed for a low-speed

previous trial, namely its inhibitory effect is stronger in blocks with a low

probability of low-speed trials.

Figure 4. Effect of previous trial's speed versus block's probability. **(a) Experiment 1A.** The left panel shows the probability effect on the aSPv given that the trial N-1 was a high-speed trial (HS, red) or a low-speed trial (LS, blue). The right panel shows the difference between the mean aSPv for trials with N-1 = HS (red), or N-1 = LS (blue) and the mean aSPv in the whole block. Each trace corresponds to one participant, while dots correspond to the average between participants. **(b)** Same analysis for Exp 1B. On the right panel, individual traces were substituted by the 95% confidence interval across participants for the sake of clarity.

Effects of the accelerating target probability on anticipatory eye movements

Following the same reasoning as for constant speed mixture blocks, we tested the effect of the probability of accelerating target kinematics on aSP. In Experiments 2A and 2B, we compared different probabilistic mixtures of trials of accelerating or decelerating target motion. Participants ran blocks of 4 different probability pairs

(P(v33d), P(v11a)): (0,1), (0.3, 0.7), (0.7, 0.3) and (1,0). In Figure 5, we present the

results relative to P(v33d) values. Figure 5a shows horizontal and vertical

trial-averaged eye velocities for one participant and each available combination of

P(v33d) and target kinematic conditions. Again, each color depicts one probability

condition and time zero indicates the target movement onset. We can see clear

anticipatory responses, with stronger anticipation occurring for higher probabilities

of the highest initial velocity and decelerating motion. After the anticipatory phase,

eye velocity traces corresponding to vacc or vdec trials separate and converge to the

target's velocity which keeps changing in time according to the acceleration

condition. Notice that the anticipation seen with P(v33d) $=0$ (i.e., P(v11a) $=1$) was

particularly small but still significant in participant 6, as in all others. Figure 5b

plots the horizontal and vertical aSPv, as a function of P(v33d), for all participants.

There is an increase in the amplitude of anticipatory pursuit as P(v33d) increases, as

confirmed by the LMM statistical analysis (P(v33d): beta = 1.88, 95% CI = (1.11,

2.65), p < .001). Consistently with the previous analysis, we did not find any

significant difference between axes (axis main effect: beta = -0.46, 95% CI =

(-1.26, 0.34), p $=0.26$) and the interaction between axis and probability did not

survive the model selection procedure, suggesting it was not significant. We found

that the aSPv was slightly higher in Exp 2B (constant target duration) than in Exp

2A (constant target displacement) (experiment effect: beta = 1.68, 95% CI = (1.41,

1.95), p < .001), but, again, the interaction between experiment and probability was

excluded from the selected model. Overall, in the probability-mixture blocks with

accelerating targets, we observed a robust probability-dependent anticipation,

similar to the mixture blocks with constant speed and this regardless of the motion

direction. The significant main effect of the experimental design (difference between

Exp 2A and 2B) suggests that the temporal regularity across trials generally favors

anticipatory behavior; however, the lack of interaction between the experiment

design and the probability effect argues against a critical role of the motion

presentation duration (at least in the tested range, namely above 500ms) on the

integration of information about the target acceleration.

Figure 5. Experiment 2A,B. Effect of the probability of accelerating target kinematics upon anticipatory eye movements (mixture blocks). **(a).** Average eye velocity across time for a representative participant. Trials are grouped according to the probability of v33d (P(v33d)) and sorted by the different target kinematic conditions for each (P(v33d), P(v11a)) mixture. Left and and right panels show horizontal and vertical eye velocity profiles, respectively. The dashed lines show the target velocity. **(b).** Mean anticipatory eye

velocity as a function of the probability of v33d, group results. The left panel shows the data for Exp 2A, while the right panel shows the data for Exp 2B. Horizontal data is shown in red, and vertical data is shown in blue. Data represented in the same way as Figure 2.

Effects of different target kinematics on anticipatory Smooth Eye Movements

Lastly, we questioned in Experiment 3 how the anticipation for accelerating target motion compares to that observed for constant target speeds. We compared three conditions where oblique target motion had a constant speed (v11c, v22c and v33c, corresponding to radial 11, 22 and 33 °/s, respectively) to conditions in which the target started at 11, 22 or 33 °/s and either accelerated uniformly (v11a, v22a, v33a, acceleration = 22 °/s^2) or decelerated uniformly (v11d, v22d, v33d, acceleration = -22 °/s^2). The different target kinematic conditions were presented in a block design and motion direction was fixed for each participant, leading to full predictability of both target's trajectory and kinematics (P=1). **Figure 6** shows eye velocity profiles recorded in one participant, for Exp 3, for each target kinematic condition illustrated by the dotted lines. Each row of **Figure 6** corresponds to one initial target speed value, while acceleration values are shown in different colors (accelerating motion in green, constant motion in blue, and decelerating motion in pink). Predictable constant speed targets drove strong anticipatory pursuits that were scaled according to target speed (blue curves, from top to bottom). Moreover, accelerating conditions also resulted in clear anticipatory pursuit responses. As clearly seen with the 22 and 33°/s initial speed conditions, green and red curves are,

548 respectively, above and below the blue ones, illustrating that anticipatory responses

549 were modulated by both initial speed and target acceleration.

Figure 6. Experiment 3: Effect of target kinematics on aSP (fully predictable blocks). **(a).** Average eye velocity over time with trials grouped by target speed (rows: v11, v22, and v33 from top to bottom) and target acceleration values (color coded: accelerating in green,

constant in blue, decelerating in pink) for one participant of Exp 3. The dashed lines show the corresponding target speed profile for each acceleration value.

Figure 7 illustrates the amplitude of anticipatory pursuit responses (aSPv) for
the different conditions across all participants. Note that, for each initial speed,
there is a tendency for aSPv to increase as the acceleration increases. To statistically
test the effect of acceleration on the anticipatory eye velocity, we ran a parametric
linear mixed-effects regression model including both the acceleration and the initial
target speed as independent variables.

We found that a model including both the initial speed and acceleration was
significantly better than a model including only the initial speed (speed-only model
BIC = 43521.6; full model BIC = 43308.97; Bayes Factor = 1.48e+46). The initial
speed significantly modulated the aSPv (v0: beta = 0.24, 95% CI = (0.15, 0.34),
$p < .001$), while the acceleration alone had a significant but smaller positive effect
on the aSPv (accel: beta = 0.017, 95% CI = (0.004, 0.030), $p = 0.009$), indicating
that indeed aSPv increases with target acceleration. We also found a significant
positive interaction between initial speed and acceleration (v0*accel: beta = 0.001 ,
95% CI = (0.0004, 0.002), $p = .004$), indicating that the effect of acceleration is
stronger for high initial speeds when compared to low initial speeds.

Anticipatory Eye Velocity

Figure 7. Experiment 3: Group effect of target kinematics on the aSPv. aSPv is grouped by initial velocity (v11, v22 and v33, from left to right, respectively) and acceleration values are shown in different colors (decelerating target motion in pink, constant target motion in blue and accelerating target motion in green). Data are represented in the same way as Figure 2.

Temporal window of estimation of the mean target speed for accelerating motion

Previous studies investigating predictive smooth pursuit during the transient disappearance of the target have compared different possible schemes of temporal integration to build an internal model of complex target motion. Such alternative internal models could either take into account only the last sample of observed target velocity before target blanking, or an average of velocity across a finite time window, or use both the last velocity sample and its rate of change. Their results suggest that the rate of speed change (i.e. acceleration) was only taken into account if target displacement properties were estimated during a sufficiently long interval

(Bennet et al. 2007). Performance was in any case inaccurate due to a lack of sufficient extrapolation of accelerating motion (Bennet & Benguigui 2013). Together with the results of Bennet and Barnes (2006), our results suggest that the target's acceleration is integrated, though with a small weight and large variability, in the internal model of visual motion that drives anticipatory smooth eye movements across a block of trials where the target motion is highly predictable (as illustrated in the previous section). However, such an effect does not imply, *per se,* that a representation of target acceleration is accessible to the visuo-motor system. For instance, an estimate of target speed across a finite temporal window could be used as a proxy for target motion and drive anticipation. Thus, we simulated a similar but more realistic version of the simplest, initial speed based, internal model of target motion. We reasoned that, rather than an instantaneous estimate of target velocity (in our case the target velocity at time 0, or target motion onset) the internal model of motion stored in memory would take into account an estimate of the mean target velocity computed over a finite time-window around target motion initiation. This alternative model could accommodate the observation that expectancy-based anticipatory velocity differs, but only weakly, for two targets with the same initial velocity and a different acceleration. Note that we can already speculate that such hypothetical temporal window of integration should be shorter than 500ms. If this was not the case, aSPv should not differ between target kinematic conditions that lead to the same mean target speed across the 500ms window of motion presentation. In Experiment 3, two pairs of conditions fit to this requirement, namely (v11a, v22d) and (v22a, v33d). For both pairs, aSPv is significantly higher

for the second than for the first target kinematic condition (v11a-v22d = -1.503, SE

= 0.139, p<.0001; v22a-v33d = -0.443, SE = 0.132, p<.001).

Figure 8 illustrates the rationale and the results of our model-based analysis of the

temporal window of velocity integration. We have shown in the previous sections

that a linear regression describes the relationship between target speed and

anticipatory eye velocity for predictable, constant target speeds (schematically

illustrated in **Figure 8a**, upper panel, blue line). We assumed that 1) the same linear

regression applies to the accelerating conditions as well, to describe the relationship

between anticipatory eye velocity and the target speed estimate (TSE)

approximating the accelerating kinematics; 2) such an estimate, TSE, would

correspond to the mean target speed computed over a finite temporal window of

integration (TWI), starting at time 0 (target motion onset) and ending at time

TWI^{end}. Thus, in order to have a reliable estimate of TWI^{end} for all kinematic

conditions across participants, we first estimated the linear regression between aSPv

and target speed at the group level, by pooling the data of all participants together.

We then computed the mean TSE knowing the aSPv values corresponding to

accelerating conditions and inverting the above-mentioned linear relation (as

illustrated in the top panel of **Figure 8a**). From the TSE value, we finally inferred

the mean TWI^{end} (as schematized in **Figure 8a,** bottom panel) and its variability

using a bootstrapping procedure (Efron, 1979, see **Figure 8b**; for details of the

calculations please refer to the figure caption and the Methods section)**.** Note that

the estimated TWI^{end} for accelerating conditions is between 0 and 500 ms, although

the distribution is very broad (v11a, mean = 0.15, 95% CI = (-0.04, 0.34); v22a,

mean = 0.18, 95% CI = (-0.03, 0.33); v33a, mean = 0.19, 95% CI = (-0.14,

0.49)). Remarkably, the TWI^{end} distribution for decelerating conditions displays

smaller values, largely overlapping with 0, but also includes a wide range of

unrealistic (negative) values (v11d, mean = -0.01, 95% CI = (-0.17, 0.10); v22d,

mean = -0.09, 95% CI = (-0.23, 0.07); v33d, mean = 0.004, 95% CI = (-0.33,

0.32)).

Figure 8. Testing the hypothesis of a finite temporal window of integration (TWI) to extract a target speed estimate (TSE) for accelerating conditions. (**a**) Schematic

illustration of the procedure to compute the temporal window (the number of datapoints is reduced, for the sake of clarity: for details refer to the Methods section). **Upper panel**: At the group level and for the constant speed blocks, we calculated the linear regression between aSPv and target speed (blue straight line). On the basis of this regression, for accelerating blocks (pink or green dots), we performed the inverse operation to calculate the estimated target speed (TSE) that would have elicited the observed aSPv. **Lower panel**: The TWI is estimated as the temporal window over which the mean of the accelerating target speed equals TSE, as exemplified for two particular accelerating conditions (v11a and v33d). **(b)** Distribution of the bootstrapped estimates of TWI for the accelerating conditions. Each panel shows the TWI distributions for one initial speed (v11, v22 and v33 from top to bottom) and for the accelerating (pink) vs decelerating (green) target motion. The smooth line depicts a gaussian fit of the histograms.

Overall, these results cannot exclude the possibility that visual accelerating

motion is approximated by the mean target speed estimate across a finite time

window. However, the large variability of the inferred TWI and the presence of

incoherent results, like negative TWI^{end} (especially for deceleration targets) impose a

strong caution in the interpretation. We will further discuss this finding in the

Discussion section.

634 Discussion

We had two main objectives with the present study. First, we wanted to analyze the

effects of a parametric change of the target speed probability upon human

anticipatory smooth eye movements. We have done this by re-analyzing previously

collected data (see Souto et al., 2008) and by replicating and generalizing those

results in a larger group of participants and different conditions, including targets

with different motion direction, speed and acceleration profiles. We used target

motion trajectories along both horizontal and oblique directions but kept target

motion direction constant within blocks in all experiments. Second, we compared different fully predictable target kinematics, namely constant or accelerating target speeds and analyzed their effect on anticipatory eye movements. We found that anticipatory responses were strongly modulated by both constant and accelerating target speeds. We report a linear scaling of anticipatory smooth pursuit with target speed (or acceleration) probability, similar to what we previously reported for a probability bias in direction and distinct from the short-term trial history effects. We also investigated, through statistical and model-based analyses, whether the internal model of visual motion that drives anticipatory smooth pursuit would integrate information about the target's accelerating profile. Our results, although very variable across participants, provide evidence that humans can integrate some information about acceleration and use it to anticipate the forthcoming target motion. Whether this integration is grounded on an internal, noisy representation of motion acceleration, or on an approximation of the target mean velocity over an extended temporal window remains to be further investigated.

A linear dependence between anticipatory pursuit and the probability of target kinematic cues

In two separate experiments, we showed that the velocity of anticipatory pursuit is modulated by the constant-speed probability of visual moving targets, regardless of its fully predictable direction. These results are consistent with previous reports (as reviewed in Kowler et al., 2019) that showed that anticipatory smooth velocity is modulated by the predictability of target speed, as well as by

short-term trial-sequence effects. Our results are however novel in several aspects. First, we demonstrate a parametric, linear relationship between the amplitude of the anticipatory phase and a broad range of target speed probability. Such a linear relationship is observed also over a large interval of target speeds (from 5 to above 30°/s) and is similar for targets moving along either the horizontal (**Figure 2**) or the oblique axes (**Figure 3**). In addition, similar to a previous study of our group (Damasse et al., 2018) we have provided evidence that short-term effects driven by the previous trial's speed can coexist and interact with the long-term effects that were the main objective in the present study. In particular, experiencing a high (low) speed trial yields an increase (decrease) of aSPv in the following trial with respect to the block's average (**Figure 4**), but this effect is strongly modulated by the context, namely by the probability of high (or low) speed trials. Moreover, we show that higher order kinematic cues such as acceleration can also modulate eye velocity during anticipatory pursuit, again regardless of the (predictable) target motion direction and such an effect is also modulated by the probability of these higher-order kinematic cues (**Figure 5**).

Overall, these novel results extend our previous findings of a linear dependence of the anticipatory eye velocity upon the target direction probability (Damasse et al., 2018; Santos & Kowler, 2017) and further demonstrate that the statistical regularities of different motion properties are efficiently stored in memory and used to drive anticipatory visuomotor control across a timescale of several seconds to several minutes and more. Our results argue for a probabilistic coding of target velocity (direction and speed), and possibly of target acceleration as well. Yet,

future work is needed to elucidate whether these aspects of target trajectories are encoded together or separately and by which neuronal populations and computational processes. It is important to notice that the present and most of the previous results demonstrating some degree of sensorimotor adaptation to the statistical regularities in the environment do not imply that probabilities are explicitly learnt and/or represented as abstract concepts in the brain. The very nature of probabilistic coding in the brain is at the heart of important lines of research, and the relation between probabilistic coding and probability-based behavior is not trivial at all. In any case, anticipatory smooth pursuit eye movements appear as an effective behavioral measure to elucidate how direction and kinematic parameters are encoded, together with their uncertainty, to control eye movements.

Is there an internal representation of accelerating target kinematics?

In this study, we also addressed the question of *how* information about accelerating motion could be integrated in the internal model of motion that drives anticipatory eye movements. To do so, we analyzed how predictable accelerating targets affect anticipatory eye velocity compared to constant-speed targets. Target motion conditions were designed such that constant speeds can directly be compared with accelerating conditions: each value of initial speed (11, 22 or 33 °/s) was paired with different acceleration values. The linear mixed-effects regression model indicated that anticipatory eye velocity was not simply scaled based on the instantaneous initial target speed, but also on its acceleration. Moreover, the effect

of acceleration increased with the initial speed (see **Figure 7**, and statistical analyses reported in the text). We performed an additional model-based analysis to tease apart the possibility that the internal model driving anticipatory smooth pursuit relies, rather than on an exact representation of target acceleration, on a simple approximation of it, namely the estimation of mean target speed across a finite temporal window close to motion onset. Leveraging on the linear relation between predictable target speed and anticipatory eye velocity, this analysis allowed us to simulate the size of the temporal window of integration and its variability across the group of participants tested in this study (**Figure 8**). Overall, the estimated temporal window of integration was strongly variable across participants and in the case of decelerating targets these estimates were often unrealistic (e.g. negative temporal windows). Interestingly, the distribution of TWI estimates with accelerating targets was clearly different from the one obtained with decelerating targets, and centered on positive and plausible values (i.e. around two-hundred milliseconds after target motion onset).

Whether and how the acceleration of moving targets is represented and used by the primate visual tracking system is still unclear (Lisberger & Movshon, 1999), even though target acceleration is a key component of many models of visual target-driven smooth pursuit eye movements (Brostek et al., 2017; Goldreich et al., 1992; Krauzlis & Lisberger, 1994). In humans, several previous studies have attempted to demonstrate a role for acceleration and whether such high-order motion cues can be learned through the history of target motion, on both the short (a few hundreds milliseconds) and medium to long (seconds to minutes and more)

timescales. Bennett et al. (2007) showed that when smoothly pursuing an accelerating target which undergoes an occlusion after a short exposition (200 ms) in a random-presentation condition, human participants are not able to adaptively use the acceleration information. Instead, participants seem to store the estimate of a constant velocity and use saccades to compensate for the displacement error between the eye position and the location where the target reappears. Those authors found, however, that after a longer exposure (500-800 ms, comparable to our visual motion duration), smooth pursuit and saccades discriminate between the different acceleration profiles. Still, prediction of the target position at the end of the occlusion was not accurate. Using again the transient target-occlusion paradigm, Bennett & Barnes (2006) probed predictive smooth pursuit of accelerating targets in blocked vs random presentation conditions. They reported two interesting and complementary results: in a blocked-design paradigm (thus with highly predictable motion over a long timescale), anticipatory eye velocity occurring (1) before the target motion onset and (2) before the end of target blanking was scaled to the target acceleration. However, increasing uncertainty about target acceleration, by mixing trials, had canceled such dependency and anticipatory eye movements were no longer distinguishable between acceleration conditions. Other studies came to the same conclusion: on a short timescale acceleration information can be used to somehow control online tracking eye (and hand) movements but not to build robust predictive motor responses to moving targets, or related perceptual judgements (Kreyenmeier et al. 2022). On the other hand, predictability over a longer timescale seems to favor the integration of acceleration information for visuomotor control

(Bennett & Barnes, 2006). Our results tend to corroborate the latter claim, with the

caveat of a large inter-individual variability observed in the anticipatory behavior

with predictable accelerating motion.

Early psychophysical studies have shown that the mean speed estimated over

the stimulus motion duration influences the perceptual discrimination of

acceleration (Brouwer et al., 2002; Gottsdanker et al., 1961; Schmerler, 1976).

Watamaniuk & Heinen (2003) showed that this is also the case when judging and

tracking an accelerated moving target. In addition, the duration of the temporal

window during which the target kinematic information is acquired seems to

influence the accuracy of acceleration estimation (Bennett et al., 2007). In the

present study, our model-based analysis of the temporal window of integration

highlighted a large inter-individual variability, as well as a dependence on the

acceleration sign, in the timescale that would be relevant to estimate the motion of

accelerating targets. Overall, several contextual factors seem to influence the

encoding and processing of visual motion acceleration. The precise nature -and the

mere existence!- of an explicit representation of visual acceleration in the brain

remains to be elucidated.

We also need a more complete understanding of how speed and acceleration

cues can be integrated through learning sensorimotor contingencies in specific tasks.

A very peculiar example is the vertical tracking of a target that changes speed by

following the gravity acceleration (Zago et al., 2010). What is being learned in an

experimental session (e.g. probabilities of occurrence) versus the entire lifespan (e.g.

the law of gravity, or friction, see Souto & Kerzel, 2013), and how this drives anticipatory pursuit responses questions the complex interactions between predictive and sensory information for an optimal tracking behavior.

Finally, in addition, in alternative or in parallel to the internal model of the retinal target speed and acceleration, reinforcement learning processes could play an important role in adapting anticipatory eye movements to predictable motion properties. Any combination of retinal position, velocity or acceleration errors could be estimated and eventually minimized over trials, akin to a *cost function*, to improve target visibility and tracking performance. Thus, we can speculate that participants could simply learn by trial and error and adapt their anticipation behavior to rapidly minimize the difference between the eyes and target position and velocity, as well as its change over time. Again, this sort of cost-minimization process remains to be thoroughly tested by future model-based experiments.

Neuronal bases of predictive tracking and processing of different kinematic properties

Electrophysiological studies in the non-human primates have provided evidence that a small subpart of the Frontal Eye Fields (FEFsem, slightly ventral compared to the saccadic FEF) is implicated in the control of predictive smooth pursuit (e.g. Fukushima et al., 2002; MacAvoy et al., 1991, see Kowler, 2019 for a review). Darlington et al. (2018) showed that FEFsem firing rate is modulated, before visual motion onset, by the expectations about the target speed. In addition, the speed-context modulation of neuronal activity continues throughout the

visually-guided phase of smooth pursuit, and it is stronger when the visual stimuli

are less reliable (i.e. at lower contrast), in agreement with Bayesian integration of

prior beliefs and sensory evidence. Such integration was also apparent in the

oculomotor recordings, with the monkeys' smooth pursuit eye velocity more

strongly modulated by the speed context for low-contrast targets. Unfortunately, the

authors could not compare the FEF preparatory activity with anticipatory eye

velocity, nor did they analyze the smooth pursuit latency dependence on motion

expectancy, thereby limiting the possibility to draw some correspondence with our

results. A second prefrontal oculomotor field, the Supplementary Eye Fields (SEF) is

also involved in the control of predictive smooth pursuit (Heinen & Liu, 1997). For

instance, de Hemptinne et al. (2008) showed that the activity of a population of SEF

neurons encoded the target direction expectations, as neurons became more active

after the presentation of a cue indicating deterministically a target motion in the

neuron's preferred direction. The evidence for the neural substrates of predictive

pursuit is much sparser in humans: Gagnon et al. (2006) have applied transcranial

magnetic stimulation (TMS) pulses to the human FEFsem and SEF regions during

visual tracking of sinusoidal target motion. They have reported an enhancement of

predictive pursuit when TMS was applied to FEFsem at different epochs, but only in

some specific conditions when TMS was applied to SEF. Several questions remain

yet unanswered. First, the respective role of FEFsem and SEF in predictive eye

movement is still debated. Second, how the different variables of target motion

trajectories are encoded and learned is yet to be investigated. Thanks to its fast,

block-designed protocol mixing different target motion cues, the present study may

inspire future neurophysiological studies in non-human and human primates,

focusing on the joint analysis of anticipatory responses and preparatory neural

activities in these two prefrontal areas.

Our results call for a reevaluation of the role of higher-order motion cues

(acceleration/deceleration) in the control and learning of predictive pursuit

behavior. There is very little evidence that the primate nervous system encodes

visual acceleration explicitly, in the visual or in the oculomotor systems. Lisberger &

Movshon (1999) measured MT single neurons' responses to image acceleration, but

did not find evidence that those neurons' activity varied with acceleration. They

found, however, that the simulated nonlinear readout of a population of MT

neurons was correlated to image acceleration (though not to its deceleration).

Similarly, Price et al. (2005) found speed tuning in MT single neurons, but not an

acceleration or deceleration tuning. However, Schlack et al. (2007) showed that a

linear classifier can extract acceleration signals from the MT population response,

given that the MT neurons' tuning to speed depended on the acceleration and

deceleration contexts of the task. Note, however, that these earlier studies focused

mainly on primate area MT while other parietal (MST) and frontal (FEF) cortical

areas might contribute to represent complex target motion trajectories and

higher-order kinematics. Future work shall elucidate how position, velocity and

acceleration cues are jointly or independently encoded across the visuo-oculomotor

distributed network, to represent and learn target trajectories for the efficient

control of action.

Conclusion

In this study, we showed that when the target speed is predictable, human participants show a linear dependence of anticipatory eye velocity upon the speed probability that is comparable to the dependence found for target direction probability. Moreover, participants also show anticipatory responses adjusted to accelerating target kinematics, and to their probability across-trials. Overall, this study contributes to the broad existing literature about the sensory and cognitive control of eye movements by better characterizing the role of predictive information about the target kinematics.

Data availability statement

The data and analysis scripts are available (DOI 10.17605/OSF.IO/SYD3T).

References

Barnes, G. R., & Asselman, P. T. (1991). The mechanism of prediction in human smooth pursuit eye movements. *The Journal of Physiology, 439*(1), 439–461. https://doi.org/10.1113/jphysiol.1991.sp018675

Barr, D. J., Levy, R., Scheepers, C., & Tily, H. J. (2013). Random effects structure for confirmatory hypothesis testing: Keep it maximal. *Journal of Memory and Language, 68*(3), 255–278. https://doi.org/10.1016/j.jml.2012.11.001

Bennett, S. J., & Barnes, G. R. (2006). Smooth ocular pursuit during the transient disappearance of an accelerating visual target: The role of reflexive and voluntary control. *Experimental Brain Research, 175*(1), 1–10. https://doi.org/10.1007/s00221-006-0533-4

Bennett, S. J., & Benguigui, N. (2013). Is Acceleration Used for Ocular Pursuit and Spatial Estimation during Prediction Motion? *PLoS ONE, 8*(5), e63382. https://doi.org/10.1371/journal.pone.0063382

Bennett, S. J., de Xivry, J.-J. O., Barnes, G. R., & Lefèvre, P. (2007). Target Acceleration Can Be Extracted and Represented Within the Predictive Drive to Ocular Pursuit. *Journal of Neurophysiology, 98*(3), 1405–1414. https://doi.org/10.1152/jn.00132.2007

Bennett, S. J., Orban De Xivry, J.-J., Lefèvre, P., & Barnes, G. R. (2010). Oculomotor prediction of accelerative target motion during occlusion: Long-term and short-term effects. *Experimental Brain Research, 204*(4), 493–504. https://doi.org/10.1007/s00221-010-2313-4

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision, 10*(4), 433–436.

Brostek, L., Eggert, T., & Glasauer, S. (2017). Gain Control in Predictive Smooth Pursuit Eye Movements: Evidence for an Acceleration-Based Predictive Mechanism. *Eneuro, 4*(3), ENEURO.0343-16.2017. https://doi.org/10.1523/ENEURO.0343-16.2017

Brouwer, A.-M., Brenner, E., & Smeets, J. B. J. (2002). Perception of acceleration with short presentation times: Can acceleration be used in interception?

Perception & Psychophysics, 64(7), 1160–1168. https://doi.org/10.3758/BF03194764

Carl, J. R., & Gellman, R. S. (1987). Human smooth pursuit: Stimulus-dependent responses. *Journal of Neurophysiology, 57*(5), 1446–1463. https://doi.org/10.1152/jn.1987.57.5.1446

Chopin, A., & Mamassian, P. (2012). Predictive Properties of Visual Adaptation. *Current Biology, 22*(7), 622–626. https://doi.org/10.1016/j.cub.2012.02.021

Collewijn, H., Van Der Mark, F., & Jansen, T. C. (1975). Precise recording of human eye movements. *Vision Research, 15*(3), 447-IN5. https://doi.org/10.1016/0042-6989(75)90098-X

Damasse, J., Perrinet, L. U., & Montagnini, A. (2018). *Reinforcement effects in anticipatory smooth eye movements. 18*(2018), 1–18.

Darlington, T. R., Beck, J. M., & Lisberger, S. G. (2018). Neural implementation of Bayesian inference in a sensorimotor behavior. *Nature Neuroscience, 21*(10), 1442–1451. https://doi.org/10.1038/s41593-018-0233-y

De Hemptinne, C., Lefèvre, P., & Missal, M. (2008). Neuronal Bases of Directional Expectation and Anticipatory Pursuit. *The Journal of Neuroscience, 28*(17), 4298–4310. https://doi.org/10.1523/JNEUROSCI.5678-07.2008

Dodge, R. (1930). OPTIC NYSTAGMUS: III. CHARACTERISTICS OF THE SLOW PHASE. *Archives of Neurology & Psychiatry, 24*(1), 21. https://doi.org/10.1001/archneurpsyc.1930.02220130024002

Efron, B. (1979). Bootstrap Methods: Another Look at the Jackknife. *The Annals of Statistics, 7*(1). https://doi.org/10.1214/aos/1176344552

Engbert, R., & Kliegl, R. (2003). Microsaccades uncover the orientation of covert attention. *Vision Research, 43*(9), 1035–1045. https://doi.org/10.1016/S0042-6989(03)00084-1

Falmagne, J.-C., Cohen, S. P., & Dwivedi, A. (1975). Two-choice reactions as an ordered memory scanning process. *Attention and Performance V*, 296–344.

Fukushima, K., Fukushima, J., Warabi, T., & Barnes, G. R. (2013). Cognitive processes involved in smooth pursuit eye movements: Behavioral evidence,

neural substrate and clinical correlation. *Frontiers in Systems Neuroscience, 7.* https://doi.org/10.3389/fnsys.2013.00004

Fukushima, K., Yamanobe, T., Shinmei, Y., Fukushima, J., Kurkin, S., & Peterson, B. W. (2002). Coding of smooth eye movements in three-dimensional space by frontal cortex. *Nature, 419*(6903), 157–162.

Gagnon, D., Paus, T., Grosbras, M.-H., Pike, G. B., & O'Driscoll, G. A. (2006). Transcranial magnetic stimulation of frontal oculomotor regions during smooth pursuit. *Journal of Neuroscience, 26*(2), 458–466.

Gauthier, G. M., Vercher, J. L., Mussa Ivaldi, F., & Marchetti, E. (1988). Oculo-manual tracking of visual targets: Control learning, coordination control and coordination model. *Experimental Brain Research, 73*(1), 127–137. https://doi.org/10.1007/BF00279667

Goldreich, D., Krauzlis, R. J., & Lisberger, S. G. (1992). Effect of changing feedback delay on spontaneous oscillations in smooth pursuit eye movements of monkeys. *Journal of Neurophysiology, 67*(3), 625–638. https://doi.org/10.1152/jn.1992.67.3.625

Gottsdanker, R., Frick, J. W., & Lockard, R. B. (1961). IDENTIFYING THE ACCELERATION OF VISUAL TARGETS. *British Journal of Psychology, 52*(1), 31–42. https://doi.org/10.1111/j.2044-8295.1961.tb00765.x

Grasse, K. L., & Lisberger, S. G. (1992). Analysis of a naturally occurring asymmetry in vertical smooth pursuit eye movements in a monkey. *Journal of Neurophysiology, 67*(1), 164–179. https://doi.org/10.1152/jn.1992.67.1.164

Heinen, S. J., Badler, J. B., & Ting, W. (2005). Timing and velocity randomization similarly affect anticipatory pursuit. *Journal of Vision, 5*(6), 1–1. https://doi.org/10.1167/5.6.1

Heinen, S. J., & Liu, M. (1997). Single-neuron activity in the dorsomedial frontal cortex during smooth-pursuit eye movements to predictable target motion. *Visual Neuroscience, 14*(5), 853–865. https://doi.org/10.1017/S0952523800011597

Hlavac, M. (2022). *stargazer: Beautiful LATEX, HTML and ASCII tables from R*

statistical output [R].

Jarrett, C. B., & Barnes, G. (2002). Volitional scaling of anticipatory ocular pursuit velocity using precues. *Cognitive Brain Research*, *14*(3), 383–388. https://doi.org/10.1016/S0926-6410(02)00140-4

Kao, G. W., & Morrow, M. J. (1994). The relationship of anticipatory smooth eye movement to smooth pursuit initiation. *Vision Research*, *34*(22), 3027–3036. https://doi.org/10.1016/0042-6989(94)90276-3

Ke, S. R., Lam, J., Pai, D. K., & Spering, M. (2013). Directional Asymmetries in Human Smooth Pursuit Eye Movements. *Investigative Ophthalmology & Visual Science*, *54*(6), 4409. https://doi.org/10.1167/iovs.12-11369

Kowler, E., Martins, A. J., & Pavel, M. (1984). The effect of expectations on slow oculomotor control—IV. Anticipatory smooth eye movements depend on prior target motions. *Vision Research*, *24*(3), 197–210.

Kowler, E., & McKee, S. P. (1987). Sensitivity of smooth eye movement to small differences in target velocity. *Vision Research*, *27*(6), 993–1015. https://doi.org/10.1016/0042-6989(87)90014-9

Kowler, E., Rubinstein, J. F., Santos, E. M., & Wang, J. (2019). Predictive Smooth Pursuit Eye Movements. *Annual Review of Vision Science*, *5*(1), 223–246. https://doi.org/10.1146/annurev-vision-091718-014901

Kowler, E., & Steinman, R. M. (1979a). The effect of expectations on slow oculomotor control—I. Periodic target steps. *Vision Research*, *19*(6), 619–632. https://doi.org/10.1016/0042-6989(79)90238-4

Kowler, E., & Steinman, R. M. (1979b). The effect of expectations on slow oculomotor control—II. Single target displacements. *Vision Research*, *19*(6), 633–646. https://doi.org/10.1016/0042-6989(79)90239-6

Krauzlis, R. J., & Lisberger, S. G. (1994). A model of visually-guided smooth pursuit eye movements based on behavioral observations. *Journal of Computational Neuroscience*, *1*(4), 265–283.

Kreyenmeier, P., Kämmer, L., Fooken, J., & Spering, M. (2022). Humans Can Track But Fail to Predict Accelerating Objects. *Eneuro*, *9*(5),

ENEURO.0185-22.2022. https://doi.org/10.1523/ENEURO.0185-22.2022

Landelle, C., Montagnini, A., Madelain, L., & Danion, F. (2016). Eye tracking a self-moved target with complex hand-target dynamics. *Journal of Neurophysiology*, *116*(4), 1859–1870. https://doi.org/10.1152/jn.00007.2016

Lenth, R. (2017). *Emmeans* [R]. https://github.com/rvlenth/emmeans

Lisberger, S. G., & Movshon, J. A. (1999). Visual motion analysis for pursuit eye movements in area MT of macaque monkeys. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *19*(6), 2224–2246. https://doi.org/10.1523/JNEUROSCI.19-06-02224.1999

Lisberger, S., & Westbrook, L. (1985). Properties of visual inputs that initiate horizontal smooth pursuit eye movements in monkeys. *The Journal of Neuroscience*, *5*(6), 1662–1673. https://doi.org/10.1523/JNEUROSCI.05-06-01662.1985

MacAvoy, M. G., Gottlieb, J. P., & Bruce, C. J. (1991). Smooth-Pursuit Eye Movement Representation in the Primate Frontal Eye Field. *Cerebral Cortex*, *1*(1), 95–102. https://doi.org/10.1093/cercor/1.1.95

Makowski, D., Ben-Shachar, M., & Lüdecke, D. (2019). bayestestR: Describing Effects and their Uncertainty, Existence and Significance within the Bayesian Framework. *Journal of Open Source Software*, *4*(40), 1541. https://doi.org/10.21105/joss.01541

Maus, G. W., Potapchuk, E., Watamaniuk, S. N. J., & Heinen, S. J. (2015). Different time scales of motion integration for anticipatory smooth pursuit and perceptual adaptation. *Journal of Vision*, *15*(2), 16–16. https://doi.org/10.1167/15.2.16

Montagnini, A., Souto, D., & Masson, G. (2010). Anticipatory eye-movements under uncertainty: A window onto the internal representation of a visuomotor prior. *Journal of Vision, 10*(7), 554–554. https://doi.org/10.1167/10.7.554

Orban De Xivry, J.-J., & Lefèvre, P. (2007). Saccades and pursuit: Two outcomes of a single sensorimotor process: Saccades and smooth pursuit eye movements. *The Journal of Physiology*, *584*(1), 11–23.

https://doi.org/10.1113/jphysiol.2007.139881

Pasturel, C., Montagnini, A., & Perrinet, L. U. (2018). ANEMO: Quantitative tools for the ANalysis of Eye MOvements. *Grenoble Workshop on Models and Analysis of Eye Movements, Grenoble, France.* https://laurentperrinet.github.io/publication/pasturel-18-anemo

Pasturel, C., Montagnini, A., & Perrinet, L. U. (2020). Humans adapt their anticipatory eye movements to the volatility of visual motion properties. *PLOS Computational Biology*, *16*(4), e1007438. https://doi.org/10.1371/journal.pcbi.1007438

Price, N. S. C., Ono, S., Mustari, M. J., & Ibbotson, M. R. (2005). Comparing Acceleration and Speed Tuning in Macaque MT: Physiology and Modeling. *Journal of Neurophysiology*, *94*(5), 3451–3464. https://doi.org/10.1152/jn.00564.2005

Robinson, D. A. (1963). A Method of Measuring Eye Movemnent Using a Scieral Search Coil in a Magnetic Field. *IEEE Transactions on Bio-Medical Electronics*, *10*(4), 137–145. https://doi.org/10.1109/TBMEL.1963.4322822

Rottach, K. G., Zivotofsky, A. Z., Das, V. E., Averbuch-Heller, L., Discenna, A. O., Poonyathalang, A., & Leigh, R. J. (1996). Comparison of Horizontal, Vertical and Diagonal Smooth Pursuit Eye Movements in Normal Human Subjects. *Vision Research*, *36*(14), 2189–2195. https://doi.org/10.1016/0042-6989(95)00302-9

Rubinstein, J. F., Singh, M., & Kowler, E. (2024). Bayesian approaches to smooth pursuit of random dot kinematograms: Effects of varying RDK noise and the predictability of RDK direction. *Journal of Neurophysiology*, *131*(2), 394–416. https://doi.org/10.1152/jn.00116.2023

Santos, E. M., & Kowler, E. (2017). Anticipatory smooth pursuit eye movements evoked by probabilistic cues. *Journal of Vision*, *17*(13), 1–16. https://doi.org/10.1167/17.13.13

Schlack, A., Krekelberg, B., & Albright, T. D. (2007). Recent History of Stimulus Speeds Affects the Speed Tuning of Neurons in Area MT. *The Journal of*

Neuroscience, *27*(41), 11009–11018.
https://doi.org/10.1523/JNEUROSCI.3165-07.2007

Schmerler, J. (1976). The Visual Perception of Accelerated Motion. *Perception, 5*(2),
167–185. https://doi.org/10.1068/p050167

Souto, D., & Kerzel, D. (2013). Like a rolling stone: Naturalistic visual kinematics
facilitate tracking eye movements. *Journal of Vision, 13*(2), 9–9.
https://doi.org/10.1167/13.2.9

Souto, D., Montagnini, A., & Masson, G. S. (2008). Scaling of anticipatory smooth
pursuit eye movements with target speed probability. *Journal of Vision, 8*(6),
665–665. https://doi.org/10.1167/8.6.665

Takeichi, N., Fukushima, J., Kurkin, S., Yamanobe, T., Shinmei, Y., & Fukushima, K.
(2003). Directional asymmetry in smooth ocular tracking in the presence of
visual background in young and adult primates. *Experimental Brain Research,*
149(3), 380–390. https://doi.org/10.1007/s00221-002-1367-3

Tychsen, L., & Lisberger, S. G. (1986). Visual motion processing for the initiation of
smooth-pursuit eye movements in humans. *Journal of Neurophysiology, 56*(4),
953–968. https://doi.org/10.1152/jn.1986.56.4.953

Voeten, C. (2020). *buildmer: Stepwise elimination and term reordering for mixed-effects*
regression [R].

Watamaniuk, S. N. J., & Heinen, S. J. (2003). Perceptual and oculomotor evidence
of limitations on processing accelerating motion. *Journal of Vision, 3*(11), 5.
https://doi.org/10.1167/3.11.5

Wu, X., Rothwell, A. C., Spering, M., & Montagnini, A. (2021). Expectations about
motion direction affect perception and anticipatory smooth pursuit
differently. *Journal of Neurophysiology, 125*(3), 977–991.
https://doi.org/10.1152/jn.00630.2020

Zago, M., Iosa, M., Maffei, V., & Lacquaniti, F. (2010). Extrapolation of vertical
target motion through a brief visual occlusion. *Experimental Brain Research,*
201(3), 365–384. https://doi.org/10.1007/s00221-009-2041-9

Supplementary material

Final models for the LMM analysis

Exp 1A:

$aSPv \sim 1 + P(HS) + (1 + P(HS) \mid participant)$

$aSPv \sim 1 + P(HS)*Tv_{N-1} + (1 + P(HS) + Tv_{N-1} \mid participant)$

Exp 1B, constant speed probability-mixtures:

$aSPv \sim 1 + P(v33) + axis + (1 + P(v33) + axis \mid participant)$

$aSPv \sim 1 + P(v33)*Tv_{N-1} + (1 + P(v33) + Tv_{N-1} \mid participant)$

Exp 2A-B, accelerating target probability-mixtures:

$aSPv \sim 1 + prob + axis + exp + axis:exp + (1 + prob + axis \mid participant)$

Exp 3, comparison between fully predictable blocks:

$aSPv \sim 1 + v0*accel + (1 + v0 \mid participant)$

Exp3, categorical model for pairwise comparisons:

$aSpv \sim 1 + condition + (1 \mid participant)$

1080

LMM analysis - result tables

1082

Exp1A: Anticipatory Parameters

	Dependent variable:
	aSPv
P(HS)	3.488^{***} (3.138, 3.838)
	t = 19.541
	p = 0.000
Constant	2.759^{***} (2.036, 3.482)
	t = 7.480
	p = 0.000
Random Effects	
Groups	3
sd(Constant)	0.77
sd(P(HS))	0.33
Note:	$^{*}p<0.01$; $^{**}p<0.001$; $^{***}p<$ 1e-04

1083

Exp 1A: Anticipatory Parameters– Sequential Effects

	Dependent variable:
	aSPv
$Tv_{N-1}[LS]$	-0.501*** (-0.663, -0.340)
	t = -6.073
	p = 0.000
P(HS)	4.357*** (3.983, 4.732)
	t = 22.827
	p = 0.000
$P(HS):Tv_{N-1}[LS]$	-2.744*** (-3.102, -2.386)
	t = -15.030
	p = 0.000
Constant	4.427*** (3.801, 5.053)
	t = 13.853
	p = 0.000
Random Effects	
Groups	3
sd(Constant)	0.70
sd(P(HS))	0.10
sd($Tv_{N-1}[LS]$)	0.29
Note:	*p$<$0.01; **p$<$0.001; ***p$<$1e-04

Exp 1B: Anticipatory Parameters – Const Speed

	Dependent variable:
	aSPv
P(v33)	2.743[***] (1.824, 3.663)
	t = 5.847
	p = 0.000
Axis[vert.]	-0.740 (-1.464, -0.016)
	t = -2.002
	p = 0.046
P(v33):Axis[vert.]	-0.674[***] (-0.985, -0.363)
	t = -4.252
	p = 0.00003
Constant	2.707[***] (1.740, 3.673)
	t = 5.490
	p = 0.00000
Random Effects	
Groups	13
sd(Constant)	1.82
sd(P(v33))	1.68
sd(Axis[vert.])	1.32
Note:	[*]p<0.01; [**]p<0.001; [***]p<1e-04

Exp 1B: Anticipatory Parameters – Const Speed - Sequential Effects

	Dependent variable:
	aSPv
$Tv_{N-1}[v11]$	-0.613*** (-0.882, -0.345)
	t = -4.482
	p = 0.00001
P(v33)	1.380* (0.437, 2.323)
	t = 2.869
	p = 0.005
$Tv_{N-1}[v11]$:P(v33)	0.823** (0.383, 1.263)
	t = 3.663
	p = 0.0003
Constant	3.948*** (3.114, 4.781)
	t = 9.279
	p = 0.000
Random Effects	
Groups	13
sd(Constant)	1.58
sd(P(v33))	1.67
sd($Tv_{N-1}[v11]$)	0.43
Note:	*p<0.01; **p<0.001; ***p<1e-04

Exp 2A-B: Anticipatory Parameters – Accelerating Target

	Dependent variable:
	aSPv
P(vdec)	1.884*** (1.111, 2.657)
	t = 4.779
	p = 0.00001
Axis[vert.]	-0.460 (-1.264, 0.343)
	t = -1.123
	p = 0.262
Exp.[const.Time]	1.683*** (1.410, 1.956)
	t = 12.086
	p = 0.000
Exp.[const.Time]:Axis[vert.]	-1.651*** (-2.027, -1.275)
	t = -8.611
	p = 0.000
Constant	3.041*** (2.099, 3.983)
	t = 6.326
	p = 0.000
Random Effects	
Groups	16
sd(Constant)	1.96
sd(P(V3))	1.58
sd(Axis[vert.])	1.66
Note:	*p<0.01; **p<0.001; ***p<1e-04

Exp 3: Anticipatory Parameters – Parametric Variables

	Dependent variable:
	aSPv
V0	0.246^{***} (0.149, 0.343)
	t = 4.963
	p = 0.00000
Accel	0.017^{*} (0.004, 0.030)
	t = 2.640
	p = 0.009
V0:Accel	0.001^{*} (0.0004, 0.002)
	t = 2.905
	p = 0.004
Constant	0.364 (-0.324, 1.052)
	t = 1.037
	p = 0.300
Random Effects	
Groups	7
sd(Constant)	0.95
sd(v0)	0.14
Groups	7
Note:	*p < 0.01; **p < 0.001; ***p < 1e-04

Exp 3: Pairwise comparisons

contrast	estimate	SE	df	t.ratio	p.value	
V1a - V1c	0.640	0.140	8749	4.559	<.0001	
V1a - V1d	0.934	0.143	8749	6.545	<.0001	
V1a - V2a	-2.236	0.136	8749	-16.396	<.0001	
V1a - V2c	-1.037	0.140	8749	-7.430	<.0001	
V1a - V2d	-1.503	0.139	8749	-10.779	<.0001	
V1a - V3a	-4.114	0.130	8749	-31.598	<.0001	
V1a - V3c	-3.320	0.135	8749	-24.626	<.0001	
V1a - V3d	-2.679	0.136	8749	-19.647	<.0001	
V1c - V1d	0.294	0.142	8749	2.065	0.0390	*n.s.
V1c - V2a	-2.876	0.136	8749	-21.149	<.0001	
V1c - V2c	-1.677	0.139	8749	-12.051	<.0001	
V1c - V2d	-2.143	0.139	8749	-15.417	<.0001	
V1c - V3a	-4.754	0.130	8749	-36.624	<.0001	
V1c - V3c	-3.960	0.134	8749	-29.472	<.0001	
V1c - V3d	-3.319	0.136	8749	-24.415	<.0001	
V1d - V2a	-3.170	0.138	8749	-22.921	<.0001	
V1d - V2c	-1.971	0.141	8749	-13.932	<.0001	
V1d - V2d	-2.437	0.141	8749	-17.251	<.0001	
V1d - V3a	-5.048	0.132	8749	-38.177	<.0001	
V1d - V3c	-4.254	0.137	8749	-31.095	<.0001	

V1d - V3d	-3.613	0.138	8749	-26.115	<.0001
V2a - V2c	1.199	0.135	8749	8.875	<.0001
V2a - V2d	0.733	0.135	8749	5.434	<.0001
V2a - V3a	-1.878	0.125	8749	-14.982	<.0001
V2a - V3c	-1.084	0.130	8749	-8.326	<.0001
V2a - V3d	-0.443	0.132	8749	-3.362	0.0008
V2c - V2d	-0.466	0.138	8749	-3.372	0.0008 *effect in the opposite direction
V2c - V3a	-3.077	0.129	8749	-23.872	<.0001
V2c - V3c	-2.283	0.134	8749	-17.100	<.0001
V2c - V3d	-1.642	0.135	8749	-12.153	<.0001
V2d - V3a	-2.611	0.129	8749	-20.288	<.0001
V2d - V3c	-1.818	0.133	8749	-13.631	<.0001
V2d - V3d	-1.176	0.135	8749	-8.719	<.0001
V3a - V3c	0.794	0.124	8749	6.413	<.0001
V3a - V3d	1.435	0.125	8749	11.442	<.0001
V3c - V3d	0.641	0.130	8749	4.927	<.0001

Degrees-of-freedom method: containment

P value adjustment: BH method for 36 tests