Atypical cortical feedback underlies failure to process contextual information in the superior colliculus of Scn2a$^{+/-}$ autism model mice

Authors: Leiron Ferrarese[1], Hiroki Asari*[1]

Affiliations:
1. Epigenetics and Neurobiology Unit, EMBL Rome, European Molecular Biology Laboratory, Monterotondo, 00015, Italy

***Corresponding author:** Hiroki Asari, E-mail: asari@embl.it

Acknowledgments: This work was supported by research grants from EMBL (H.A.) and SFARI (#875447; H.A.); and EMBL Interdisciplinary, International and Intersectorial Postdoctoral Fellowship (EI3POD; L.F.). The EMBL Gene Editing and Embryology Facility is acknowledged for the mouse line rederivation; EMBL Genetic and Viral Engineering Facility for virus production; EMBL IT Support for provision of computer and data storage servers; and the LAR facility for taking care of animals. We thank all the Asari lab members as well as Cornelius Gross, Eleonora Satta and Michael Lombardo for many useful discussions.

Author contributions: L.F. and H.A. designed the study; L.F performed experiments and analyzed the results; L.F. and H.A. wrote the manuscript.

Competing interests: The authors declare no competing financial interests.

Key words: autism spectrum disorder; superior colliculus; implicit visual learning; Bayesian inference; hierarchical Gaussian filter.

Acronyms: AAV, adeno-associated virus; ANOVA, analysis of variance; AP, anterior-posterior; ASD, autism spectrum disorder; CNMF, constrained non-negative matrix factorization; CNO, clozapine N-oxide; DAPI, 4',6-diamidino-2-phenylindole; DREADD, designer receptors exclusively activated by designer drugs; DV, dorsal-ventral; ES, early stable; FDR, false discovery rate; GECI, genetically-encoded calcium indicator; hM4Di, modified human M4 muscarinic acetylcholine receptor for inhibitory DREADD; HGF, hierarchical Gaussian filter; IR, infrared; ISI,: interstimulus interval; L5, layer 5; LS, late stable; ML, medial-lateral; PFA, paraformaldehyde; PBS, phosphate buffer saline; RGC, retinal ganglion cell; SC, superior colliculus; Scn2a, sodium voltage-gated channel alpha subunit 2; V1, primary visual cortex; VL, volatile; WT, wild-type.

1 Abstract

Atypical sensory integration and contextual learning are common symptoms in autism, but how

sensory circuits are affected remains elusive. Here we performed *in vivo* two-photon calcium

imaging and pupillometry of mice engaged in an implicit learning task in stable and volatile visual

contexts. Wild-type and Scn2a-haploinsufficient autism-model mice exhibited distinct patterns of

context-dependent modulation in pupil and neuronal dynamics in the superior colliculus, while

retinal outputs showed no contextual modulation. A hierarchical Bayesian learning model showed

that the superior colliculus largely represented expectations on the task volatility in wild-type mice,

but rather on the outcome itself in $Scn2a^{+/-}$ mice, much as suggested in autism. Chemogenetic

perturbation revealed that the cortical feedback signals mediated the observed contextual

modulation, both at the neurophysiological and behavioral levels. These results suggest that the

corticotectal input is critical for contextual sensory integration in the superior colliculus, and its

anomaly underlies abnormal sensory learning in autism.

14 Introduction

Bayesian inference provides a theoretical framework for understanding how the brain combines

prior knowledge with incoming sensory information to shape perceptual experiences and guide

behavior (Friston, 2009, 2012). In particular, a hierarchical Bayesian learning model has been

successfully applied to explain atypical behaviors observed in individuals with autism spectrum

disorder (ASD; Pellicano and Burr, 2012; Chrysaitis and Seriès, 2023): e.g., how overlearning and

overestimation of the environmental volatility result in a reduced response to unexpected events

in ASD (Lawson et al., 2017). However, the current experimental evidence supporting this

hypothesis is limited in both sample sizes and computational models (Chrysaitis and Seriès,

2023).

How do neural circuit functions relate to Bayesian inference, and how is it altered in ASD?

The superior colliculus (SC) is a promising target to address these questions. It is a midbrain hub

for the visual system (May, 2006; Cang et al., 2018), and expected to be central for visual

inference as part of a hierarchical recursive network in the brain (**Fig. 1A**; Friston, 2009, 2012).

Neurons in the superficial layers of SC receive direct visual input from the retina, and play a role

in processing visual motion and orientation (Lomber et al., 2001; Feinberg and Meister, 2015;

Inayat et al., 2015; Li et al., 2020). From the Bayesian perspective, these retinal inputs can be

considered as "*likelihood*" of the sensory evidence. Superficial SC neurons also receive a

prominent feedback signal from the primary visual cortex (V1; Wang and Burkhalter, 2013; Zingg

et al., 2017; Benavidez et al., 2021). These top-down inputs from V1 provide signals for contextual

gain modulation: e.g., enhancing the responses of SC to looming stimuli, while negatively

controlling the response gain to the saliency of complex visual scenes (Zhao et al., 2014;

Ahmadlou et al., 2017). Furthermore, nonlinearity of the corticotectal synapses serves for

regulating sensory-guided innate behavior, attentional modulation, and visual learning (Zhao et

al., 2014; Ruediger and Scanziani, 2020; Tang and Higley, 2020). Therefore, although some

controversies still remain on the modulatory role of the corticotectal inputs (Hoffmann and

Straschill, 1971; Ogasawara et al., 1984; Wang et al., 2010), we expect that they contribute to the

formation of "*prior"* beliefs in the Bayesian framework. Given the convergence of retinal inputs and

cortical feedback onto SC, one can then further posit that local SC neurons are involved in the

computation of "*posterior*" perception.

A growing body of evidence indicates the involvement of subcortical structures in ASD,

including SC (Marco et al., 2011; Johnson et al., 2016; Jure, 2022). For instance, gaze

management and attentional focus are two major functions of SC that are often altered in ASD

(Lomber et al., 2002; Kato et al., 2011). SC also plays a role in prioritizing sensory attention

(Sridharan et al., 2017) and mediating innate avoidance responses to a looming stimulus in mice

(Dean et al., 1989; Yilmaz and Meister, 2013; Wei et al., 2015; De Franceschi et al, 2016). Such

visual responses are less robust in ASD model mice (Burnett et al., 2022), highlighting anomalous

processing in the SC.

As an increasing number of ASD-associated genes have been identified (Sanders et al.,

2012; Satterstrom et al., 2020), animal models of ASD have been developed to investigate the

underlying neurobiological mechanisms (Patel et al., 2018). One of the well-established ASD

mouse models is the haploinsufficiency for *sodium voltage-gated channel alpha subunit 2*

(Scn2a$^{+/-}$ mice; Ben-Shalom et al., 2017). The *Scn2a* gene encodes the Na$_V$1.2 sodium channels

that help regulate dendritic excitability of mature cortical pyramidal neurons (Planells-Cases et al.,

2000), including those forming the corticotectal pathway (Zingg et al., 2017; Spratt et al., 2019).

Haploinsufficiency of Na$_V$1.2 reduces the backpropagation of action potentials into dendrites,

impairing synaptic plasticity and strength in layer 5 (L5) corticofugal pyramidal neurons (Spratt et

al., 2019). Therefore, Scn2a$^{+/-}$ mice are expected to have atypical corticotectal projections, leading

to abnormal neuronal modulations within the SC circuitry.

Here we developed an implicit visual learning task, similar to the ones used in human

studies (Mathys et al., 2014; Lawson et al., 2017), but for mice to form probabilistic associations

of cue-outcome stimulus sequences in stable and volatile environments. We then combined *in*

vivo two-photon calcium imaging, chemogenetic perturbation, pupillometry, and computational

modelling to characterize both neurophysiological and behavioral phenotypes of mice engaged in

this task. Wild-type (WT) mice exhibited a robust context-dependent modulation of the pupil and

neuronal dynamics in SC, but not in the retinal output signals. A hierarchical Bayesian learning

model indicated that these SC neurons represented posterior expectations on the environmental

volatility rather than those on the outcome (cue-outcome contingency) *per se*. In contrast, Scn2a$^{+/-}$

mice demonstrated no or anomalous contextual modulations in the pupil or neuronal dynamics,

respectively. Specifically, these SC neurons showed no adaptation to an increased outcome

predictability in the stable environment, and largely represented the outcome expectations in a

way consistent with the observations in ASD subjects from the hierarchical Bayesian learning

model perspective (Lawson et al., 2017). We further found that chemogenetically inhibiting the

visual cortical feedback to SC fully abolished contextual modulations in both WT and Scn2a$^{+/-}$

mice, both at the physiological and behavioral levels. These results suggest that the corticotectal

input plays a critical role in integrating contextual information in SC, and that anomaly in this

process underlies atypical sensory learning in ASD.

Results

Implicit visual learning paradigm to test behavioral and physiological anomalies in autism model mice

Individuals with autism spectrum disorder (ASD) often struggle to learn about probabilistically

aberrant events (Lawson et al., 2017). Understanding the neuronal processes underlying implicit

learning and the integration of contextual information will thus provide valuable insights into the

neurobiological mechanisms of ASD (Chrysaitis and Seriès, 2023). In this study, we focused on

the mouse visual system to study a) how neuronal circuit computation contributes to implicit

learning about visual environmental contexts; and b) how it is altered in a mouse model of ASD

(Scn2a$^{+/-}$). We thus began by designing an implicit visual learning paradigm based on Bayesian

principles (Mathys, 2011): i.e., probabilistic associative learning of cue-outcome stimulus

sequences, much as those experimental paradigms used in human studies (Lawson et al., 2017).

Figure 1: Bayesian implicit learning task to compare the visual responses of the superior colliculus between wild-type and autism model mice.
A: Rationale of experimental design. The superficial superior colliculus (SC) receives direct sensory input from reginal ganglion cells (RGCs) and feedback from the primary visual cortex (V1). In a Bayesian inference framework, one can assume that the retinal input represents the likelihood (dark red), while V1 provides information about the animal's prior belief (blue). Neurons in SC can then be considered to compute the posterior contextual perception (light green). **B**: Schematic of the implicit visual learning task. In each trial, a neutral cue (vertical or horizontal gratings; 1 s) is followed by an aversive outcome (looming and/or sweeping dark spot, randomly chosen from six different types; 2 s) or non-aversive one (omitted; 2 s) with an inter-stimulus interval (ISI) of 0.5 s. In total, the task consisted of 216 trials, including 108 aversive outcome trials (color-coded) and 108 non-aversive ones (dark gray). **C**: The conditional probability of cue-outcome contingencies (expected, gray; unexpected, light brown) during stable and volatile environments. The stable environment is twice as long as the volatile one (orange), and divided into early- and late-stable epochs (72 trials each; light gray and light blue, respectively). **D**: The task aims to let animals form expectations on cue-outcome probabilities during the stable environment, leading to increased prior and posterior beliefs in the late-stable epoch. In the volatile epoch, however, the precision of the posterior decreases as the likelihood reduces. **E:** In Scn2a[+/-] mice, sensory hypersensitivity (top, model 1) and/or imprecise priors (bottom, model 2) lead to the posterior perception dominated by sensory input (likelihood).

Our implicit learning task was carefully tailored with visual stimuli to elicit robust responses

in the mouse superior colliculus (SC; Cang et al., 2018), such as static gratings with different

orientations (Basso and May, 2017; De Franceschi and Solomon, 2018) and looming and

sweeping stimuli that trigger SC-mediated arousal and innate defensive behavior (Dean et al.,

1989; Yilmaz and Meister, 2013; Wei et al., 2015; De Franceschi et al., 2016). Specifically, we

used vertical and horizontal gratings as neutral cues; and a combination of sweeping and looming

dark spots as aversive outcomes while a gray screen (omitted stimulus) as a non-aversive

outcome (**Fig. 1B**). Since the task did not involve any reward or punishment, we exploited visual

threats in the outcome stimulus to motivate animals to recognize stimulus sequences. To increase

unpredictability, we introduced uncertainty by randomizing the type and direction of the visual

threat at each presentation (**Fig. 1B**; see Methods for details).

The task followed a Bayesian update scheme modeled by a hierarchical Gaussian filter

(HGF; Mathys, 2011; Lawson et al., 2017), consisting of a stable environment followed by a

volatile one (**Fig. 1C**). During the stable environment (144 trials in total), we presented aversive

outcome stimuli more frequently after one neutral cue stimulus than after another. Mice were then

expected to form probabilistic associations for these cue-outcome stimulus sequences. At a small

probability, trials with expected outcomes were interspersed with unexpected ones. This allowed

animals to determine whether an unexpected outcome indicates a deviation from a stable stimulus

pattern or a change in the pattern itself. In the stable environment, the stability and precision of

the sensory inputs (likelihood) remained high. Thus, the prior expectation should increase in the

early stable epoch (ES; initial 72 trials) and remain high in the late-stable epoch (LS; last 72 trials);

and so does the posterior belief about the environment, or the predictability, that can be given by

the product of the likelihood and the prior (**Fig. 1D**). In the following volatile epoch (VL, 72 trials in

total), we then introduced variations in the probability of aversive stimuli following each neutral

stimulus. This challenged the animal's expectations on the stimulus sequences. The environment

became less precise (i.e., likelihood decreased) while the prior belief remained relatively precise

because updating the prior is slow in general. Thus, the posterior precision should decrease and

shift towards the prior in wild-type (WT) animals.

In Scn2a$^{+/-}$ mice (**Fig. 1E**), we instead expect that their posterior is rather dominated by

the likelihood due to sensory hypersensitivity and/or imprecise prior belief formation as implicated

in human studies (Pellicano and Burr, 2012; Haker et al., 2016; Lawson et al., 2017). On the one

hand, hypersensitivity to sensory inputs will lead to an overly high likelihood regardless of the

stability of the environment. The posterior probability will then follow the changes in the prior,

hence its precision will increase in the LS epoch, but stay unchanged in the VL epoch (model 1).

On the other hand, if the prior expectations are incoherent and not updated during the task, the

posterior predictability will follow the changes in the likelihood. Thus, its precision will remain in

the LS epoch but decrease in the VL epoch (model 2). In either case, the prior and posterior

predictions will not become close enough in ASD, leading to a larger perception error than in WT.

Distinct contextual modulation patterns in SC between WT and Scn2a$^{+/-}$ mice

If neurons in SC compute contextual-dependent predictions about the environment, they should

reflect the posterior predictability distribution, while inputs from retinal ganglion cells (RGCs) and

the primary visual cortex (V1) represent the likelihood and the animal's prior belief, respectively.

To test this hypothesis, we conducted experiments on awake, head-fixed mice expressing

genetically-encoded calcium indicators (GECIs) in SC somata delivered via adeno-associated

viral vectors (AAVs; see Methods for details; **Fig. 2A,B**). Specifically, we exposed the mice to the

implicit visual learning task as described above, and simultaneously performed *in vivo* two-photon

calcium imaging of superficial SC through a cranial window, together with pupillometry for

behavioral monitoring (Dombeck et al., 2007; Feinberg and Meister, 2015; Molotkov et al., 2023;

Fig. 2A,B; Supplementary Fig. 1A).

On average we recorded 123±37 cells per animal (mean ± standard deviation; 27 mice)

and estimated their spiking activity using CaImAn (Pnevmatikakis et al., 2019; $Z(t)$ in *z*-score, see

Methods for details). As expected (Cang et al., 2018; De Franceschi and Solomon, 2018),

superficial SC neurons generally responded well to the onset and/or offset of the static gratings

(**Supplementary Fig. 1**). About a half of them were responsive to the looming and/or sweeping

stimuli (60±19 cells per recording), whereas in most cells we found virtually no response to the

omitted stimulus (gray screen; e.g., **Supplementary Fig. 1B**). Here we considered cells

responsive if the trial average of their visual response dynamics exceeded a threshold adjusted

for each recording (e.g., $\max[z(t)] > 1$, where $z(t) = \langle Z(t) \rangle_{\text{trials}}$; see Methods for details).

To examine the SC dynamics during the implicit learning task, we compared the aversive

stimulus responses at the population level across the three analysis epochs in our paradigm (ES,

LS, and VL; **Fig. 1C**). For each responsive cell, we calculated the average response during the

stimulus presentation period in each epoch ($\bar{z} = \langle z(t) \rangle_t$; e.g., **Fig. 2C**), and subtracted the mean

across the epochs for normalization ($\Delta z = \bar{z} - \langle \bar{z} \rangle_{\text{epochs}}$; e.g., **Fig. 2C-E**). Using this normalized

response, Δz, we found that SC neurons in WT animals responded significantly weaker to the

aversive stimuli during the LS epoch than the ES or VL epoch (**Fig. 2E**, 1,037 cells in total from

17 mice; p(ES-LS)=0.007, p(ES-VL)=0.8, p(LS-VL)=0.04, Tukey-Kramer test). This suggests that

the responses of SC neurons in WT mice are modulated by the visual contexts in two ways. First,

when the stimulus pattern is reliably predictable during the stable environment, the responses to

aversive stimuli become progressively attenuated (adaptation). Second, when the stimulus

pattern deviates from an animal's expectation during the volatile environment, the response of SC

neurons is reset to the baseline and becomes as high as in the initial ES epoch before adaptation.

Figure 2: Neurons in the superior colliculus show contextual adaptation to a stable environment in wild-type mice but not in Scn2a[+/-] mice.

A: Experimental setup. *Left*: The schematic illustrates a head-fixed mouse pre-injected with adeno-associated viruses (AAVs) carrying genetically encoded calcium indicators (e.g., jGCaMP7f). Visual stimulus was presented to the left eye. *Right:* The pupil dynamics of the left eye were recorded with an infrared (IR) camera (top; pupil outline, red). Two-photon calcium imaging of the right superior colliculus (SC) was performed through a glass window implanted with a silicone plug to expose SC (bottom). **B**: *Left:* An example frame of motion-corrected two-photon calcium imaging in the mouse SC. *Right*: A coronal slice image taken two months after AAV injections, demonstrating a high expression of jGCaMP8m (green) in the right SC, along with 4',6-diamidino-2-phenylindole stain (DAPI, blue). **C**: Deconvoluted spatial components extracted using CaImAn in a sample wild-type (WT) mouse displaying the topographic position of SC neurons responsive to aversive stimuli (top). Representative aversive stimulus responses of an SC neuron (circled in red on top), averaged over 36 aversive outcome trials for each analysis epoch (bottom: $z(t) = \langle Z(t) \rangle_{\text{trials}}$; shaded area, s.e.m.). The average over the stimulus period ($\bar{z} = \langle z(t) \rangle_t$) and s.e.m. are shown on the right for each epoch (from left to right: early stable (ES), gray; late stable (LS), light blue; volatile (VL), orange). **D**: Heatmap ($z(t)$) and the population average ($\langle z(t) \rangle_{\text{cells}}$) of the aversive stimulus responses in each epoch for the responsive SC neurons (sorted with $\max[z(t)] > 1$), recorded from a representative WT mouse (90 cells from C; shaded area, s.e.m.). **E**: Distributions of the normalized SC responses across epochs ($\Delta z = \bar{z} - \langle \bar{z} \rangle_{\text{epochs}}$) from all WT animals in the dataset (1,037 cells from 17 mice): violin plots, distribution of all cells; gray lines with open circles, average for each mouse; vertical bars, interquartile range; thick line graph with error bars, mean ± s.e.m.; ***, p<0.001; **, p<0.01; *, p<0.05 with Tukey-Kramer test for pairwise comparisons. See Methods for details. **F-H**: Corresponding data for Scn2a[+/-] mice (F, simultaneously recorded SC cell location in the image plane and a representative cell's responses; G, representative data from all simultaneously recorded responsive cells (86 cells); H, population data for Scn2a[+/-] mice, 630 cells in total from 10 mice).

Supplementary Figure 1: Neuronal activity of the mouse superior colliculus and pupil dynamics in response to cue-outcome stimuli in the implicit visual learning task.
A: Trial-by-trial response example during the early stable (ES) epoch, with 83% expected and 17% unexpected outcomes. *From top to bottom*: The spiking activity of the superior colliculus (SC) neurons ($Z(t)$ in z-score) estimated from recorded calcium dynamics; corresponding population average response ($\langle Z(t) \rangle_{\text{cells}}$; gray shade, ±1 standard deviation); and concurrent pupil area dynamics ($A(t)$). **B**: Responses of SC neurons to each of the four stimulus types presented in the ES epoch (from left to right: neutral cue stimulus associated with non-aversive outcome, non-aversive omitted stimulus, neutral cue stimulus associated with aversive outcome, and aversive stimuli). *From top to bottom*: the trial average responses of individual SC cells ($z(t) = \langle Z(t) \rangle_{\text{trials}}$), sorted by the peak aversive stimulus response values ($\max[z(t)]$, rightmost histogram; red rectangle, responsive cells with $\max[z(t)]>1$); the average population response ($\langle z(t) \rangle_{\text{cells}}$); and the average pupil dynamics ($\langle A(t) \rangle_{\text{trials}}$). Gray shade, ±1 standard deviation.

In contrast, Scn2a$^{+/-}$ mice displayed different context-dependent modulations in SC during

the implicit visual learning task: no adaptation in the stable environment, but increased neuronal

responses during the volatile environment (**Fig. 2F-H**, 630 cells in total from 10 mice; p(ES-

LS)=0.7, p(ES-VL)=5e-4, p(LS-VL)=0.009, Tukey-Kramer test). These response patterns of SC

neurons were significantly different between WT and Scn2a$^{+/-}$ mice (p(epoch x group) = 0.002, 3-

way nested analysis of variance (ANOVA); see Methods for details). This suggests that, unlike

WT animals, SC neurons in Scn2a$^{+/-}$ mice fail to attenuate their responses even when the stimulus

pattern is kept stable and predictable. They are, however, still sensitive to increased volatility, and

thus their response during the VL epoch exceeded the baseline activity in the ES epoch. This is

consistent with the observations on ASD subjects who tend to overestimate environmental

volatility during stable environments, and overreact to increased unpredictability during volatile

environments (Lawson et al., 2017).

Context-dependent modulation is absent in retinal output responses

How do SC neurons achieve such context-dependent modulations? One possibility is that the

contextual modulation in SC is inherited from RGCs. To investigate the contribution of bottom-up

RGC input to SC, we performed *in vivo* two-photon calcium imaging of RGC axon terminals in SC

(Molotkov et al., 2023) while the animals were engaged in the implicit visual learning task (**Fig.**

3A). On average we recorded 240±34 putative axonal boutons per animal (mean ± standard

deviation; 5 mice), of which 131±29 were responsive to the looming and/or sweeping stimuli

($\max[z(t)] > 1$). These RGC axon terminals in SC showed consistent responses to the aversive

stimuli regardless of the visual context (**Fig. 3B**). Across populations, we found no significant

contextual modulation across ES, LS, and VL epochs (**Fig.3C**; 655 putative RGC boutons in total

from 5 mice; p(ES-LS)=0.8, p(ES-VL)=0.6, p(LS-VL)=0.9, Tukey-Kramer test). This excludes the

possibility that the contextual modulation in SC arises from the retina.

Figure 3: Retinal ganglion cells projected to the superior colliculus do not show any context-dependent modulation during the implicit learning task.
A: Experimental setup. Animals were intraocularly injected with adeno-associated viruses (AAVs) carrying GCaMP6s and underwent cranial window surgery over the right superior colliculus (SC) to monitor the activity of retinal ganglion cell (RGC) axons *in vivo* (top). An example frame of RGC axonal imaging in SC (middle) and deconvoluted spatial components by CaImAn (bottom). **B**: Aversive stimulus responses of simultaneously recorded RGC putative boutons (top: heatmap of $z(t) = \langle Z(t) \rangle_{\text{trials}}$, sorted by $\max[z(t)]$) and the population average (bottom, $\langle z(t) \rangle_{\text{boutons}}$; 166 boutons in total from A; shaded area, s.e.m.) in each analysis epoch (shown in the same format as in **Fig. 2D,G**). **C**: Distributions of the normalized RGCs responses across epochs ($\Delta z = \bar{z} - \langle \bar{z} \rangle_{\text{epochs}}$, where $\bar{z} = \langle z(t) \rangle_t$; 655 boutons from 5 WT mice), shown in the same format as in **Fig. 2E,H**. Violin plots, distribution of all boutons; gray lines with open circles, average for each mouse; vertical bars, interquartile range; thick line graph with error bars, mean ± s.e.m. No significant difference was found by pairwise comparisons (Tukey-Kramer test).

Chemogenetically inhibiting corticotectal projection abolishes context-dependent modulation in the superior colliculus

Another possible source for the contextual modulation in SC is the corticotectal projection (**Fig. 1A**). Besides sensory input from RGCs, the superficial layer of SC receives substantial feedback signals from layer 5 (L5) pyramidal neurons in V1 (Zingg et al., 2017). These V1 L5 neurons integrate feedforward visual information through their basal dendrites and feedback contextual information via the apical dendrites (Larkum, 2013), and provide output to SC for controlling the response gain (Zhao et al., 2014; Ahmadlou et al., 2017; Tang and Higley, 2020). Cortical pyramidal neurons in Scn2a$^{+/-}$ mice have impaired synaptic functions due to a reduced expression

of the Na$_V$1.2 sodium channel (Spratt et al., 2019). This may result in a failure to incorporate visual

and contextual information in V1, which in turn may lead to an anomalous modulation of SC as

observed in our paradigm (**Fig. 2**).

To examine the role of V1 feedback in the contextual modulation of SC, we employed a

chemogenetic approach to selectively inhibit the V1 L5 corticotectal neurons using a designer

receptor exclusively activated by designer drug (DREADD; Armbruster et al., 2007). Specifically,

we achieved selective expression of modified human M4 muscarinic acetylcholine receptors for

inhibitory DREADD (hM4Di) in the target neuronal population by injecting Cre-dependent AAV8

carrying the hM4Di receptor into V1 and AAV2retro carrying Cre-recombinase into SC (**Fig. 4A,B**).

Systemic administration of the DREADD actuator, clozapine N-oxide (CNO), then activated

hM4Di, hence effectively inhibited the V1 neurons projecting to SC. As before, AAVs carrying

GECIs were also injected into SC for monitoring the visual responses of SC neurons with *in vivo*

two-photon microscopy.

Figure 4: Inhibition of the corticotectal projection abolishes context-dependent modulation in the
superior colliculus.
A: The schematic of the viral vector strategy for chemogenetic inhibition of the projection from the primary
visual cortex (V1) to the superior colliculus (SC). AAV8::hSyn-DIO-hM4Di-mCherry was injected into V1,
while AAV2retro::hSyn-Cre and AAV9::hSyn-jGCaMP8m were co-injected in SC. Systemic administration
of clozapine N-oxide (CNO) activated the hM4Di receptor, hence inhibiting the V1-to-SC projection. **B**:
Histological verification of the hM4Di receptor expression (with mCherry, red) in the layer 5 (L5) cells in V1
projecting to SC. Blue, 4',6-diamidino-2-phenylindole (DAPI); green, SC neurons expressing jGCaMP8m.
C,D: Distributions of the normalized SC responses (Δz) across epochs in WT (C, 355 neurons in total from
7 mice) or Scn2a$^{+/-}$ mice (D, 433 neurons in total from 7 mice) after chemogenetically blocking V1-to-SC
projection. Data are shown in the same format as in **Fig. 2E,H**. No significant contextual-modulation was
found in either case (Tukey-Kramer test).

Supplementary Figure 2: Context-dependent modulation in the superior colliculus is abolished by chemogenetic inhibition of the corticotectal projection.
A: Representative aversive stimulus responses ($z(t) = \langle Z(t) \rangle_{\text{trial}}$; gray shades, s.e.m.) of three SC neurons in a representative WT mouse expressing hM4Di in SC-projecting V1 neurons with intraperitoneal injection of saline (top; WT) or the DREADD actuator, CNO (bottom; WT V1-block). The average over the stimulus presentation period ($\bar{z} = \langle z(t) \rangle_t$) is shown on the right for each cell for each epoch (early stable, ES; late stable, LS; volatile, VL). Spatial arrangement of the cells, extracted from imaging data by CaImAn, is shown in the middle with red circles indicating the representative cells (#1, #4, and #20). **B,C:** Heatmap ($z(t)$; 73 cells) and the population average ($\langle z(t) \rangle_{\text{cells}}$) of the aversive stimulus responses in each epoch from the representative WT mouse after saline (B) or CNO (C) injections. **D,E**: Distributions of the normalized SC responses across epochs ($\Delta z = \bar{z} - \langle \bar{z} \rangle_{\text{epochs}}$) with saline (D, 355 cells in total from 7 mice; WT dataset) and CNO injections (E, 348 cells in total from 7 mice; WT V1-block dataset, reproduced from **Fig.4C**). Data are shown in the same format as **Fig. 2D,E**. *, p<0.05 with Tukey-Kramer test for pairwise comparisons.

Chemogenetic inhibition of the projection from V1 to SC abolished the context-dependent modulation of SC in both WT and Scn2a[+/-] mice (**Fig. 4C,D; Supplementary Fig. 2**). After the intraperitoneal injection of CNO, SC neurons did not show adaptation during the LS epoch or enhanced responses during the VL epoch (e.g., **Supplementary Fig. 2**). As a result, the overall SC response level did not significantly change throughout the implicit visual learning task (**Fig.**

4C; 355 cells in total from 7 WT mice: p(ES-LS)=0.8, p(ES-VL)=0.7, p(LS-VL)=1, Tukey-Kramer

test; **Fig. 4D**; 433 neurons from 7 Scn2a$^{+/-}$ mice: p(ES-LS)=0.7, p(ES-VL)=1, p(LS-VL)=0.9). Such

response patterns across epochs in the chemogenetically-manipulated animals were significantly

different from those in intact animals (p(epoch x group) = 0.004, 3-way nested ANOVA; see

Methods for details). These results support that feedback input from V1 plays a key role in

modulating the aversive stimulus response of SC neurons in a context-dependent manner. In

particular, we suggest that V1 projections provide SC with contextual information necessary for

achieving adaptation to expected outcomes during the stable environment.

Pupil dynamics reflect context-dependent stimulus pattern recognition in WT but

not in Scn2a$^{+/-}$ mice

Besides neuronal activities, we collected pupillometric data (**Fig. 5**) as a proxy to monitor cognitive

processes of the animals (Loewenfeld, 1958). Here we in particular interpreted pupil dynamics as

a reflection of an animal's stimulus pattern recognition (Suzuki et al., 2018; Ganea et al., 2020;

Dobbins, 2021). Indeed, all mice exhibited stereotypical pupil dynamics during the implicit visual

learning task: i.e., pupil contraction in response to the cue stimuli, and pupil dilation during the

outcomes, especially when the aversive stimuli were presented (**Supplementary Fig. 1B**,

bottom). Here we focused our analysis on the pupil dynamics during the aversive outcome trials

across the three analysis epochs (ES, LS, and VL; **Fig. 1C**) as we did for the neuronal dynamics

(**Figs. 2-4**).

In WT animals, we found a greater pupil dilation during the LS epoch than the ES or VL

epoch (**Fig. 5D,H;** q(ES-LS) = 0.04, q(ES-VL) = 0.1, q(LS-VL) = 0.02, Storey's false discovery

rate (FDR) from Wilcoxon signed-rank test; 17 mice). This supports that pupil dilation is associated

with an animal's recognition of the stimulus pattern (Suzuki et al., 2018; Ganea et al., 2020;

Dobbins, 2021). This context-dependent modulation of the pupil dynamics was abolished when

the feedback signal from V1 to SC was chemogenetically blocked (**Fig. 5F,J;** q(ES-LS)=0.1, q(ES-

VL)=0.1, q(LS-VL)=0.1; 5 mice). In contrast, Scn2a$^{+/-}$ animals generally showed a smaller pupil

size than WT animals throughout the task (10 Scn2a$^{+/-}$ mice, 0.33 ± 0.05 mm^2, mean ± s.e.m of

the median pupil area; 17 WT mice, 0.49 ± 0.04 mm^2; p = 0.037, Mann-Whitney U-test; **Fig. 5A-**

C), and displayed no context-dependent modulation (**Fig. 5E,I;** q(ES-LS)=0.2, q(ES-VL)=0.1,

q(LS-VL)=0.1; 10 mice). Chemogenetic inhibition of the corticotectal projection did not affect the

pupil dynamics in Scn2a$^{+/-}$ mice (**Fig. 5G,K;** q(ES-LS)=0.1, q(ES-VL)=0.1, q(LS-VL)=0.1; 5 mice).

The modulation pattern of the pupil dynamics across epochs was significantly different between

WT and the other three experimental groups (p(epoch x group) = 0.04; 2-way ANOVA; see

Methods for details). These findings indicate that the influence of the corticotectal signals

extended beyond the neuronal activity in SC, likely playing a role in the contextualization of the

visual scene even at the perceptual level; and that Scn2a$^{+/-}$ mice have altered V1-to-SC signaling,

resulting in a failure to properly recognize the stimulus patterns during the implicit learning task.

Taken together, we found that the modulation of the pupil dynamics (**Fig. 5**) was well linked

to that of the neuronal dynamics in SC (**Figs. 2-4**): i.e., the greater the pupil dilation was, the

weaker the SC responses were in our experimental paradigm, much as was demonstrated in

previous studies (Wang et al., 2012; Joshi et al., 2016). Most importantly, Scn2a$^{+/-}$ mice showed

impaired neurophysiological and behavioral phenotypes, both due to anomalous signaling from

V1 to SC. This highlights a crucial role of the top-down corticotectal pathway in forming perceptual

associations to recognize and properly react to visual contexts (Tang and Higley, 2020), and a

possible relationship between ASD and dysfunctions of V1 L5 projection neurons (Marco et al.,

2011; Johnson et al., 2016; Jure, 2022).

Figure 5: Volatility-dependent pupillometric response patterns in wild-type but not in Scn2a[+/-] mice.
A,B: Representative image frame of the left eye (top; red, pupil outline) of WT (A) and Scn2a[+/-] mice (B) and the pupil area dynamics $A(t)$ during visual stimulation (bottom; dashed line, median). **C**: median pupil size of 17 WT and 10 Scn2a[+/-] mice during the implicit visual task (error bar, mean ± s.e.m): *, $p < 0.05$, Mann-Whitney *U*-test. **D-G**: Pupil size dynamics of mice (average proportional change in pupil area, $\langle a(t) \rangle_{\text{mice}}$; shaded area, s.e.m.) in response to the aversive stimulus in each analysis epoch (from left to right: early stable, ES; late stable, LS; volatile, VL): D, 17 WT mice; E, 10 Scn2a[+/-] mice; F, 5 WT mice with chemogenetic inhibition of V1-to-SC projection (WT V1-block); and G, 6 Scn2a[+/-] mice with chemogenetic inhibition of V1-to-SC projection (Scn2a[+/-] V1-block). **H-K**: Normalized change in pupil size upon aversive stimulus presentation in each analysis epoch ($\Delta a = \bar{a} - \langle \bar{a} \rangle_{\text{epochs}}$): thin line, individual animal; error bars, population average and s.e.m.; H-K, from corresponding groups in D-K, respectively. *, $q < 0.05$, Storey's FDR from Wilcoxon signed-rank test.

SC neurons represent expectations on environmental volatility in WT mice, while those on outcomes in Scn2a$^{+/-}$ mice

Thus far we have analyzed the overall activity level of SC populations during the implicit visual learning task, and demonstrated that their dynamics were largely consistent with the behavior of the posterior predictability from the Bayesian inference viewpoint (**Figs. 2-4**). What do then individual neurons in SC represent? To address this question, we simulated how an ideal Bayesian observer should respond to the sequence of the cue-outcome contingencies in our experimental paradigm (Mathys, 2011; Friston, 2009, 2012), and compared the dynamics of the underlying model parameters with those of the observed neuronal responses (**Fig. 6**).

As an ideal observer model, here we employed an HGF model similar to the one used to explain the differences between subjects with and without ASD in their behavioral responses to environmental uncertainty (Lawson, et al., 2017). Using variational Bayes inversion processes, the model infers hidden states that represent beliefs about the sensory input at three hierarchical levels (**Fig. 6A**; see Methods for details). The lowest level concerns directly on the outcome (aversive versus non-aversive outcomes; X_1); the next level on the outcome probability (cue-outcome contingencies; X_2); and the highest level on the likelihood of environmental changes (stability versus volatility; X_3). Inference at each level i depends on that at a higher level, and involves the following five parameters: prior predictions on the outcome ($\hat{\mu}_i$); variance of prediction belief ($\hat{\sigma}_i$); learning rates (α_i); precision-weighted prediction errors (ε_i); and uncertainties (unc_i). In total, the HGF model has 13 parameters that are dynamically updated on a trial-to-trial basis (as the two are redundant; **Fig.6B**; see Methods for details). Each of these parameters provides an insight into distinct aspects of the Bayesian inference process at different levels of uncertainty associated with the sensory input.

Figure 6: Superior colliculus neurons in wild-type mice compute expectations about the volatility of the visual environment while those in Scn2a$^{+/-}$ mice focus instead on outcome expectations.
397
398
A: Schematic representation of the generative HGF model as a Bayesian network. The perceptual model
399
tracks a mouse's learning of the task structure across three levels. Level 1 (X_1) represents trial-wise stimulus
400
transitions from one stimulus to the next; X_2 the probability of the transitions of contingencies; and X_3 the
401
environment volatility, where k is the current trial number. Mice hold and update beliefs about the true
402
quantities at each level, with a mean μ and a variance σ; ϑ and ω are mouse-specific parameters that couple
403
the levels and determine the respective speed of belief updating about phasic volatility and transition
404
contingencies. **B**: HGF model parameter trajectories, generated with the cue-outcome contingencies of the
405
implicit visual learning task. Non-aversive outcome trials were excluded to match neuronal trajectories (in
406
C). Learning rates (Lr) at X_1-X_3; Prior predictions (Pp) at X_1-X_3; Believe variances (Bv) at X_2 and X_3; Outcome
407
uncertainty (Un) at X_1; Information uncertainty at X_2; Volatility uncertainty at X_3; Precision-weighted
408
prediction errors (Pe) at X_2 and X_3. See Methods for details. **C**: Population average of the normalized
409
neuronal trajectories for each experimental group ($\langle \Delta z_k^{res} \rangle_{cells}$). From left to right: WT, 1150 neurons from 19
410
mice; Scn2a$^{+/-}$, 523 neurons from 9 mice; WT V1-block, 289 neurons from 6 mice; Scn2a$^{+/-}$ V1-block, 386
411
neurons from 6 mice. **D**: *Left*, Neural representation of HGF parameters for each experimental group (as in
412
C), given as the percentage of SC neurons whose response trajectory was significantly correlated with each
413
HGF model parameter trajectory (from B): circle area, percentage of cells with $q<0.05$ from Storey's FDR;
414
red, positive correlation; blue, negative correlation. A bias of the correlation polarity for each HDR parameter
415
was examined with a shuffling test with FDR correction: *, $q<0.05$; **, $q<0.01$. *Right*, absolute correlation
416
between the population average neuronal trajectories ($\langle \Delta z_k^{res} \rangle_{cells}$ from C) and HGF trajectories (from B),
417
averaged for each level, $\langle |\rho| \rangle_{params|level}$. **E,F:** Population average trajectory of RGC axonal responses (E as
418
in C) and the percentage of RGC axonal boutons whose trajectories were significantly correlated with each
419
HGF parameter trajectory (F as in D). **G**: Pairwise cosine similarity index of neural representations of HGF
420
parameters between different experimental groups. See Methods for details.
421

To make a fair comparison between the model and neuronal outputs, we focused only on

the trials with aversive outcomes because SC neurons were generally not responsive to the non-

aversive stimulus in our paradigm (i.e., the omitted stimulus; see, e.g., **Supplementary Fig. 1B**).

Thus, after running the simulation with the entire trial sequence, we concatenated the output of

the aversive outcome trials alone to generate the trajectories of the HGF model parameters (**Fig.**

6B). To generate corresponding trajectories of the neuronal signals (Δz_k^{res}; **Fig. 6C**), we normalized

the observed responses of individual neurons for each of the six different types of aversive stimuli,

and further filtered out slow non-stationary components (see Methods for details). We then

conducted pairwise Pearson's correlation tests (with FDR correction) between the neuronal and

model-parameter trajectories to characterize neuronal representations of the HGF model in the

SC of WT and Scn2a$^{+/-}$ mice.

We obtained two conclusions from this analysis. First, most of the HGF parameters were

represented by some subsets of SC neurons, evidenced by statistically significant correlations

between the two trajectories (**Fig. 6D**, *left*). This was the case with both WT and Scn2a$^{+/-}$ animals,

regardless of the chemogenetic inhibition of the corticotectal projection. In contrast, hardly any

HGF parameter was well represented by the incoming signals from the retina (**Fig. 6E,F**).

Second, SC neurons in WT mice had a distinct representation of the HGF parameters,

compared to those in Scn2a$^{+/-}$ animals or when corticotectal projections were chemogenetically

blocked (**Fig. 6D,G**). Specifically, WT mice showed a significant bias in the representation of two

HGF parameters at level 3 (learning rate α_3, 3.4% of positively and 0.1% of negatively correlated

cells, q = 0.002, shuffling test; uncertainty unc_3, 0.1% of positively and 3.2% of negatively

correlated cells, q = 0.01). In contrast, Scn2a$^{+/-}$ mice showed a tendency towards a positive

correlation bias at lower levels (level 1 learning rate α_1, 2.8% of positively and 0.0% of negatively

correlated cells; level 2 uncertainty unc_2, 2.7% of positively and 0.2% of negatively correlated

cells). Blocking V1 inputs to SC generally attenuated the bias of the parameter representations in

both WT and Scn2a$^{+/-}$ animals (**Fig. 6D**). This trend was also clear when we computed Pearson's

correlation, ρ, between the population average of the neuronal trajectories ($\langle \Delta z_k^{\mathrm{res}} \rangle_{\mathrm{cells}}$; **Fig. 6C**)

and each of the HGF parameter trajectories (**Fig. 6B**). The average of the absolute correlation

values over the model parameters at each level ($\langle |\rho| \rangle_{\mathrm{params|level}}$; **Fig. 6D**, *right*) showed that SC

population trajectories in WT were on average more correlated with X_3 parameters, while those in

Scn2a$^{+/-}$ or after blocking V1-to-SC projections were with X_1 parameters. Furthermore, we used a

cosine similarity to measure how close the HGF parameter representations in SC were between

different experimental groups (see Methods for details), and found that the representations in WT

and Scn2a$^{+/-}$ animals were dissimilar between each other, but became rather similar when V1

projections to SC were chemogenetically inhibited in WT mice (**Fig. 6G**).

Taken together, these results provide additional evidence that SC neurons in WT animals

are involved in the computation of the overall contextual uncertainty in the visual environment by

incorporating information from V1. In contrast, SC neurons in Scn2a$^{+/-}$ mice focus primarily on

"lower levels" of uncertainty from recent sensory experiences due to anomalous V1 feedback

signaling. This is consistent with an oversensitivity to immediate outcomes observed in ASD

(Bertone et al., 2005; Baron-Cohen et al. 2009; Lawson et al., 2017), highlighting the importance

of feedback signaling from V1 in properly processing visual environment context.

Discussion

Many species have evolved innate behaviors such as flight or freeze responses as essential

survival strategies to evade predators (Tinbergen, 1951; LeDoux, 2012; Branco & Redgrave,

2020). Mice, for example, respond rapidly and robustly to overhead shadows as they often imply

a threat, such as an aerial predator (Yilmaz and Meister, 2013; De Franceschi et al., 2016). This

instinctive response is controlled by the superior colliculus (SC; Sahibzada et al., 1986; Evans et

al., 2018; Shang et al., 2018), a midbrain structure that directly transforms visual signals from the

retina into motor commands, hence bypassing higher cognitive processing in the neocortex (Cang

et al., 2018). This helps animals react fast to potential threats, prioritizing survival over conscious

vision. This can, however, do more harm than good to animals if their response flexibility is too

limited, much as in autism spectrum disorder (ASD; Marco et al., 2011; Johnson et al., 2016; Jure,

2022). In autumn, for instance, looming shadows arise frequently from falling leaves; however,

they are harmless and thus animals do not need to respond. Adaptive learning allows animals to

tailor their behavior according to the specific context, determining properly when to react to or

ignore potential threats, such as looming shadows, to ensure survival without unnecessary

reactions.

In this study, we dissected the neurophysiological basis of such contextual learning in wild-

type (WT) and Scn2a-haploinsufficient ASD-model mice. Following the Bayesian inference

principle (Pellicano and Burr, 2012; Friston, 2009, 2012), we developed an implicit visual learning

task and examined the visual responses of SC neurons in animals engaged in the task, with the

assumption that SC represents the posterior belief on the stimulus while inputs from retinal

ganglion cells (RGCs) and the primary visual cortex (V1) reflect the likelihood and the prior belief,

respectively (**Fig. 1**). Consistent with this assumption, WT mice showed contextual modulations

in both SC (**Fig. 2**) and pupil dynamics (**Fig. 5**): the smaller the SC responses, the larger the pupil

area, and the more precise the posterior prediction (i.e., the narrower the posterior distribution;

Fig. 1D). Contrary to our initial hypothesis, however, we did not observe any contextual

modulation in RGC input to SC (**Fig. 3**). This indicates that signals from the retina do not directly

carry information on the visual environmental context or the likelihood. We instead suggest that

the feedback signals from V1 should carry both the likelihood and the prior. Indeed,

chemogenetically blocking V1 feedback fully eliminated the contextual modulations in SC (**Fig. 4**)

as well as in pupil dynamics (**Fig. 5**). This underscores a central role of the cortical feedback in

the adaptive responses at both neurophysiological and behavioral levels.

As expected from previous studies on ASD subjects (Lawson et al., 2017), Scn2a$^{+/-}$

animals exhibited anomalous responses to the implicit learning task at both neurophysiological

and behavioral levels. In particular, SC neurons demonstrated sustained responses (no

adaptation) in a stable environment, and an over-enhanced activity in a volatile environment (**Fig.**

2). This is consistent with our weak prior model of ASD, where we expect no change in the

posterior precision during the stable epochs but a reduction during the volatile epoch (**Fig. 1E**,

model 2). This implies that posterior computation in Scn2a$^{+/-}$ is primarily driven by changes in the

likelihood (sensory input variability). Notably, such an atypical modulation pattern in Scn2a$^{+/-}$ was

abolished by blocking V1 input (**Fig. 4**). Taken together with the results from WT animals, we then

suggest that the feedback signals from V1 have two counteracting roles in modulating SC

responses (Tang and Higley, 2020) – suppression linked with stability and facilitation with volatility

– and their imbalance underlies the anomaly in Scn2a$^{+/-}$. In WT mice, on the one hand, V1

suppresses SC responses to predictable aversive stimuli in a stable environment. A milder

behavioral response will then follow so that animals can better evaluate the situation. When the

environment becomes more volatile and unpredictable, V1 then facilitates SC to regain its

response. This will help trigger a stronger behavioral response, prioritizing the animal's immediate

safety. In Scn2a$^{+/-}$ animals, on the other hand, the suppressive function of V1 feedback seems

primarily impaired, leading to an imbalance towards facilitation (**Fig. 2**). It has been shown that

reduced expression of Na$_V$1.2 channels in Scn2a$^{+/-}$ affects signal integration properties of these

layer 5 (L5) corticofugal pyramidal cells (Spratt et al., 2019). They may then fail to properly

integrate top-down cortico-cortical signals at the apical dendrites that carry contextual information,

especially about environmental stability.

The above-described scenario was further supported by our hierarchical Gaussian filter

(HGF) model analysis (**Fig. 6**). In both WT and Scn2a$^{+/-}$ animals, HGF parameters across all levels

were widely represented by subpopulations of SC neurons (**Fig. 6D**). We found, however, a

representation bias at different hierarchical levels, in particular for the learning rate α: level 3 for

WT, while level 1 for Scn2a$^{+/-}$ animals. This may explain why ASD individuals tend to focus more

on immediate sensory inputs and associated outcomes (Lawson et al., 2017; Baron-Cohen et al.

2009; Bertone et al., 2005), as they both belong to lower-level HGF inference processes (Mathys,

2011). If this is indeed the case, resulting sensory overload and/or oversensitivity to

unpredictability in ASD (Gomot et al., 2011) may be alleviated by adjusting cortical feedback to

bias neural computation towards higher-level HGF inferences, hence redirecting the focus from

immediate details to broader contexts.

Our results largely align with the Bayesian brain theory (Friston, 2009, 2012). It describes

the brain as recursive circuits that receive feedforward and feedback inputs within a hierarchical

structure, continuously updating beliefs and computing the posterior. The output is then sent

upstream as the likelihood for higher-order brain structures, and downstream as a prior belief for

lower-order regions. In our case, the RGC-SC-V1(L5) loop can be considered as a lower-level

circuit in this recursive system, with a higher-level circuit consisting of the visual thalamus-V1(L5)-

associative cortex loop (**Fig. 7**). We suggest that anomalies in Scn2a$^{+/-}$ reside in V1 L5 neurons

that compute an incomplete posterior and send it to SC as a weak prior (Tang and Higley, 2020);

and such dysfunction in V1 likely arises from further upstream, such as the anterior cingulate

cortex whose activity has been suggested to be sensitive to the environmental volatility (Behrens

et al., 2007). While we focus on a specific brain region and a specific computation involved, our

findings may extend as a more generic computational pattern of ASD within complex, recursive

feedforward-feedback interactions of sensory and associative systems. In future studies, it would

be interesting to investigate other brain areas in a similar Bayesian framework, and also test

different ASD mouse models for better understanding neurophysiological underpinnings of ASD.

Figure 7: Circuit model for hierarchical interactions in the sensory system.
A lower-level subcortical circuit of the RGC-SC-V1(L5) loop (right) interacts with a sensory cortical circuit of the lateral geniculate nucleus (LGN)-V1-associative cortex loop (middle), which in turn interacts with a higher-level cortical circuit (left). Arrows represent axonal projections, and color indicates information from the Bayesian perspective (red, likelihood; blue, prior; green, posterior).

Methods

No statistical method was used to predetermine the sample size. We performed all animal experiments under license 233/2017-PR from the Italian Ministry of Health, following the protocol approved by the Institutional Animal Care and Use Committee at European Molecular Biology Laboratory. We conducted data analyses using Python and Matlab (Mathworks), and reported all summary statistics as mean ± standard error of the mean (s.e.m.) unless specified otherwise. We considered p- and q-values significant at α < 0.05 *, 0.01 **, 0.001 ***, unless stated otherwise.

Animals

We used C57BL/6J mice (*Mus musculus*, both sex), aged approximately 5-10 weeks at the time of the initial surgery. We housed the mice under a 12-hour light/12-hour dark cycle and gave them *ad libitum* access to water and food. After the surgical procedures, we grouped the animals based on their operation day. We conducted the imaging experiments when the mice were between 8-16 weeks old.

563   We acquired SCN2A haploinsufficient mice as frozen sperm (Scn2a⁻) from Riken

564   Bioresource Research Center (RBRC10243), and revitalized the line in a C57BL/6J background.

565   The homozygous mutants (Scn2a$^{-/-}$) were lethal. We used the heterozygous mice (Scn2a$^{+/-}$) as a

566   model of autism spectrum disorder (ASD; Sandlers 2012; Ben-Shalom et al., 2017), and

567   considered their homozygous littermates (Scn2a$^{+/+}$) as wild-type (WT).

Intracranial viral injections

We locally injected pseudotyped adeno-associated viruses (AAVs) into the mouse superior

colliculus (SC). These AAVs consisted of AAV2 *rep* and AAV9, AAV5, or AAV2-retro *cap* genes,

or a hybrid of AAV1 and AAV2 *cap* genes; and were used to express genetically-encoded calcium

indicators (GECIs; jGCaMP7f, jGCaMP8m, or jRGECO1a; **Fig. 2A**) or Cre-recombinase under

the pan-neuronal human synapsin (hSyn) promoter. We also injected pseudotyped AAVs,

composed of AAV2 *rep* and AAV8 *cap*, into the primary visual cortex (V1) to express of the

inhibitory designer receptor exclusively activated by designer drugs (DREADD), hM4Di-mCherry,

under the hSyn promoter (Dana et al., 2019; **Fig. 4A**).

We performed the intracranial viral injection simultaneously with the cranial implantation,

as described below. We created a craniotomy over the right SC or V1 and inserted an injection

pipette (with a tip diameter of approximately 30 μm; WPI 1B120F-3 borosilicate glass capillary

pulled with Zeitz DMZ puller) filled with a virus solution (ranging from approximately 5x10^{12} to

4x10^{14} vg/mL in phosphate-buffered saline, PBS) through the dura. For SC, we positioned the

pipette using coordinates relative to Bregma, approximately [-4]-[-4.5] mm anterior-posterior (AP),

0.5-0.7 mm medial-lateral (ML), and slowly advanced it to a depth of about 1.25 mm. For V1, we

used the following coordinates: -3.5 mm AP, 1.8-2 mm ML, and 0.6 mm dorsal-ventral (DV). We

injected the virus solution at a rate of 1-2 nL/s for SC (0.4-0.6 μL in total) and 0.5-1 nL/s for V1

(0.25-0.4 μL in total) using a microinjection pump (either Neurostar NanoW or WPI NanoLiter

2010). After completing the injection, we waited at least 10 minutes before slowly withdrawing the

pipette, and then proceeded with the cranial window implantation procedure.

Intravitreal viral injections

We performed Intravitreal injections of AAV2, pseudotyped with a hybrid of AAV1 and AAV2

capsids, to deliver hSyn-axon-GCaMP6s expression cassette to the mouse retinal ganglion cells

(RGCs; Molotkov, et al., 2023; **Fig. 3A**). We exposed the scleral surface on the left eye and made

a small piercing with a sterile 28-30G needle in between the sclera and the cornea. We then

inserted an injection pipette (~50 μm tip diameter with 30-40° bevel) prefilled with a virus solution

(~1.5×10^{14} vg/mL in phosphate-buffered saline with 0.001% Pluronic F68 and 0.001% FastGreen)

into the vitreous chamber, approximately 1 mm deep. After a good sealing of the pipette was

formed, we injected 1.2 μL of the virus solution at a rate of 10 nL/s, using a microinjection pump.

We slowly withdrew the pipette at least 5 minutes after the completion of the injection, and covered

the treated eye with the eye ointment. We carried on the rest of the procedure and the head

implant as described in the previous paragraph on intracranial injections.

Cranial implantations

We performed the cranial window implantation over the mouse SC as described previously

(Molotkov et al., 2023). Before the surgery, we prepared a cranial window assembly by activating

the surface of a circular glass coverslip (5 mm diameter, 0.13-0.15 mm thickness; Assistent Karl

Hecht) with a laboratory corona treater (BD-20ACV Electro-Technic Products). We then fused the

activated glass coverslip to a cylindrical silicone plug (1.5 mm diameter, 0.75-1.00 mm height;

Kwik-Sil, WPI) by baking it at 70-80 °C for 24 hours.

During the implantation, we anesthetized the mice (induction: 4% isoflurane in oxygen;

maintenance: 1.5-2.0%) and positioned them in a stereotaxic apparatus (Stoelting 51625). We

used a heated plate (Supertech Physiological Temperature Controller) to maintain body

temperature and prevent hypothermia, and applied eye ointment (VitA-POS, Ursapharm) to

protect the eyes. We disinfected and removed the scalp (Betadine 10%, Meda Pharma), then

scratched and cleaned the skull surface to ensure the cement adhered well.

Using a high-speed surgical drill (OmniDrill35, WPI) with a 0.4 mm ball-tip carbide bur

(Meisinger), we made a craniotomy of approximately 3.0 mm AP by 2.5 mm ML over the right SC.

To prevent bleeding, we applied haemostatic sponges (Cutanplast, Mascia Brunelli) soaked in

sterile cortex buffer (NaCl 125 mM, KCl 5 mM, Glucose 10 mM, HEPES 10 mM, CaCl$_2$ 2 mM,

MgSO$_4$ 2 mM, pH 7.4) to the craniotomy site. We carefully positioned the cranial window

assembly, pushing the transversal sinus and posterior cortex about 0.5 mm forward, with the

silicone plug covering the medial-caudal region of the right SC (see, e.g., **Fig. 2A**). We fixed and

sealed the implant with tissue adhesive (Vetbond, 3M). We then cemented a custom-made

titanium headplate (0.8 mm thick) to the skull using a blend of acrylic cement powder (Paladur,

Kulzer) and cyanoacrylate adhesive (Loctite 401, Henkel).

After the surgery, the animal was allowed to recover from anesthesia in a warmed chamber

and then returned to its home cage. For postoperative care, we administered intraperitoneal

injections of 5 mg/kg Rimadyl (Zoetis) and 5 mg/kg Baytril (Bayer) to the mice daily for 3-5 days.

We then waited an extra 20-30 days to ensure the cranial window fully recovered before starting

in vivo two-photon imaging sessions.

Habituation

One week before the experiment began, we habituated the mice to both the experimenter and the

head-fixation procedure in a dark setting. We positioned the animals on a stable platform (non-

rotating disc; about 20 cm diameter) and let them stay still. We mounted a plexiglass tunnel over

the rear part of their bodies to increase comfort. We conducted these habituation sessions twice

a day, each lasting for 2 hours, over five to ten days. This process accustomed the mice to the

experimental setup and aimed to reduce any stress during the actual data collection.

Experimental setup

Two to seven days after training, we placed the animal on a stable platform with its head fixed

under a two-photon microscope, in front of a spherical projection screen. We used a non-rotating

disc (about 20 cm diameter) as the platform to minimize the effect of motor activities on neuronal

firing during the recordings (Schröder et al., 2020). The platform, however, allowed for vertical

excursions, which was compensated by a spring and a counter-weight adjustable for each animal

and monitored by a Hall sensor. This helped minimize head movements by preventing the mouse

from exerting excessive force on the head-holder with its paws. We excluded from the analysis

any trials if the average vertical excursion of the platform during the trial exceeded 0.25 mm. We

used an infrared (IR) camera (Imaging Source DMK23UX174) with IR illumination to monitor the

movement of the animal's left eye during recordings (**Fig. 2A**; 30 frames/s; 640-by-480 pixels).

Chemogenetics and histology

We had a subset of mice with hM4Di expressed in V1 (7 WT and 7 Scn2a$^{+/-}$; **Fig. 4;**

Supplementary Fig. 2). These animals underwent the experiment twice, with at least a two-week

interval. For both experiment repetitions, mice received intraperitoneal injections of either a saline

solution (0.9% NaCl) containing 0.2 mg/ml clozapine N-oxide (CNO), leading to a final dose of 3

mg/kg (administered in a volume of 0.1-0.3 ml, adjusted for mouse weight), or the saline alone.

We alternated the solution allocation: mice receiving saline alone in the first iteration were given

the CNO solution in the second, and vice versa. Starting 45-70 minutes post-injection, the mice

underwent the experiment for approximately one hour.

Once all the recording sessions were done, we histologically examined the expression of

hM4Di-mCherry. We anesthetized the mice terminally with a ketamine/xylazine overdose, and

perfused them intracardially with 4% paraformaldehyde (PFA) in PBS. We then harvested the

brains, incubated them for a day in 4% PFA, and for at least another day in PBS. We sliced the

fixed brains into coronal sections (100 µm thickness) with a cryostat (Leica CM3050S). We then

mounted the slices on a glass coverslip using 4',6-diamidino-2-phenylindole (DAPI)-mount. We

performed fluorescent imaging with a slide scanner (VS200 automated slide scanner, Olympus)

with a 10X dry objective (UPLAPO 10x, NA: 0.4, X-line) across three channels: DAPI, GCaMP,

and mCherry (**Fig. 2B**). Only those animals that displayed clear mCherry expression (hence

hM4Di expression) in V1 layer 5 (L5) somata (e.g., **Fig. 4B**) were included in the analysis (**Fig.**

4C,D; Supplementary Fig. 2; "V1-block" groups).

Visual stimulation

We presented visual stimuli to the mice as described previously (Boissonnet et al., 2023). We

used a custom gamma-corrected digital light processing device to project images onto a spherical

screen placed approximately 20 cm from the left eye of the animal. The projected images covered

the left monocular visual field within ±22° in elevation and ±36.5° in azimuth (**Fig. 2A**). The images

were displayed at a resolution of 1280-by-720 pixels and a frame rate of 60 Hz.

One day before the experiment, mice underwent a training process. We first exposed to

them a gray screen for at least 10 minutes to adapt to the average luminance in a head-fixed

condition under the microscope. Subsequently, we presented the following visual stimuli:

1. Still gratings: A random sequence of black and white square wave gratings in four

orientations (0°, 45°, 90°, 135°) with a spatial frequency of 0.1 cycles/°. Each stimulus

lasted for 1 second, followed by a 0.5-second interstimulus interval (ISI). This sequence

was repeated 46 times in 5 minutes.

2. Cue-outcome stable 100%: Horizontal and vertical bars (abovementioned still gratings

with 0° and 90° orientations, respectively) served as cue stimuli. Horizontal bars were

682   followed by an omitted stimulus (prolonged ISI with a gray screen for 2 seconds), while

683   vertical bars were followed by one out of four possible aversive stimuli (2 s): Sweep-left (a

684   black disk 8° in diameter, appearing at 10° from the top edge and moving horizontally from

685   the left edge to the right one at 42 °/s), Sweep-right (moving from right to left), Sweep-left-

686   loom (a Sweep-left for the first second to bring the disk at the upper center of the screen,

687   followed by the disk increasing in size for the next second at a speed of 57°/s while moving

688   its center downward until reaching the horizontal center and a final size of 64° in diameter),

689   and Sweep-right-loom (Sweep-right for the first half, followed by the loom for the second

690   half). To avoid pupillary light reflex, the average luminance of the screen was kept constant

691   at any time. For the sweep and loom stimuli, the background luminance was adjusted for

692   the relative area of the black disk at every frame. We presented three blocks of 72 cue-

693   outcome trials (approximately 5 min), where the stimulus sequence of each block was

694   pseudo-randomized with different seeds.

695 3. Cue-outcome stable 83%: The cue-outcome probabilities were changed so that horizontal-

696   omitted and vertical-aversive trials occurred with a probability of 83.4%, while horizontal-

697   aversive and vertical-omitted trials occurred with a probability of 16.6%. We presented

698   three blocks of 72 cue-outcome trials (approximately 5 min), where the stimulus sequence

699   of each block was pseudo-randomized with different seeds.

700 On the day of the experiment, mice were again habituated to the average luminance for at least

701 10 minutes. We then presented the following stimuli:

702 1. Still gratings.

703 2. Cue-outcome stable 83%, but with six different types of aversive stimuli. Besides the four

704   aversive stimuli described above, we included: Slow-loom (a black 3° disk appeared in the

705   vertical center at the top edge and increased its size at a speed of 28.5°/s while moving

706   its center downward until reaching the horizontal center and a final size of 64° in diameter)

and Fast-loom (a black 3° disk appeared in the vertical center at the top edge and stayed

still for 0.5 seconds; it then increased its size at a speed of 57°/s while moving its center

downward until reaching the horizontal center and a final size of 64° in diameter; it then

stayed still again for 0.5 seconds). In total, 216 trials were presented in three blocks (72

trials each; **Fig. 1B and Supplementary Fig. 1A**). Animals typically moved a lot at the

beginning and stayed still after a while. Thus, the first block was excluded from the analysis

to avoid any movement artifact. The second and third ones were used for the analysis as

early-stable (ES) and late-stable (LS) epochs, respectively (**Fig. 1C**).

3. Cue-outcome volatile: The cue-outcome stimulus sequences were presented with three

different contingencies: a given cue-outcome (horizontal-omitted and vertical-aversive)

probability of 16.6% for the first 24 trials; 50% for the next 24 trials; and 83.4% for the last

24 trials (**Fig. 1C**). These data (72 trials in total) were referred to as the volatile (VL) epoch

in the analysis. Collectively, the cue-outcome stimuli #2 and #3 constituted an implicit

visual learning task.

4. Random water-wave stimuli (3 minutes): These stimuli were used to generate binary

masks for signal source extraction in calcium image analysis (see below "Calcium image

analysis" section).

5. Randomly flickering black-and-white checkerboard stimuli (10 minutes): These stimuli

were used for receptive field mapping. The rectangular fields had a width of 3.7° and a

height of 2.9°. Each field was independently modulated by white noise at a frequency of 4

Hz.

In the cue-outcome stimuli (#2 and #3), the trials were evenly balanced within each analysis epoch

(72 trials each), with 36 horizontal and 36 vertical cues presented before 36 omitted and 36

aversive outcomes. Each of the six different aversive stimuli was presented six times within the

36 aversive trials. The sequences of stimuli in each epoch were pseudorandomized with different

seeds, but we used the same set of seeds for all mice. This ensures that each mouse was

presented with a new random sequence of stimuli in each epoch, but all mice were presented with

the same sets of sequences and cue-outcome contingencies (**Fig. 1B**).

For the subset of mice subject to the chemogenetics experiments (see "Chemogenetics

and histology" section above), we repeated the stimulus paradigm on the same experimental day

for CNO or saline treatment.

In vivo two-photon imaging

During the experiment, we kept the animal with its head fixed inside the rig for a maximum duration

of 2 hours (typically 2 to 5 imaging sessions per animal). We performed two-photon calcium

imaging using a galvo-resonant scanner (Scientifica HyperScope) and SciScan image acquisition

software. The microscope setup included a mode-locked Ti:sapphire tunable laser (InSight DS+,

Spectra-Physics) and a plan fluorite objective (CFI75 LWD 16X W, Nikon).

In each imaging session, we conducted single-plane time-lapse recordings with a field of

view approximately ranging from 0.65-by-0.65 mm to 0.43-by-0.43 mm at a depth of 115-130 μm

from the surface of SC. We used the excitation wavelength of 920 nm for GCaMPs, and 1,040 nm

for jRGECO1a. The average laser power under the objective ranged between 40-80 mW. We

bandpass-filtered the emitted fluorescence signal (BP 527/70 or BP 650/100 after beam-splitter

FF580-FDi01, Semrock) and detected it using a non-descanned gallium arsenide phosphide

photomultiplier tube (Hamamatsu GaAsP PMT). We acquired each frame with either 512-by-512

pixels (16-bit depth) at a rate of 30.9 Hz, or 1,024-by-1,024 pixels at a rate of 15.5 Hz (**Figs. 2B**

and 3A).

Calcium image analysis

To simplify manual alignment across different recordings, we first cropped the recorded images

from 512-by-512 pixels to 480-by-480 pixels (1.3-0.85 μm/pixel), or from 1,024-by-1,024 to 960-

by-960 (0.65-0.43 μm/pixel) using ImageJ software. We then applied a series of rigid and non-

rigid motion corrections in CaImAn (version 8 or 9; Pnevmatikakis et al., 2019).

To generate masks for cellular detection in CaImAn, we used either a segment of the

recording containing 3,000 frames from the random water-wave stimulus presentation period or

the entire recording frames. We divided the images into 6-by-6 patches for the constrained non-

negative matrix factorization (CNMF) deconvolution, resulting in a total of 36 patches. We set the

expected number of neurons per patch (*params.K*) to 5 for SC cell bodies, and set the size of the

neurons (*params.gSig*) to 5-by-5 pixels (*half size*) for 512-by-512 acquisitions, or 10-by-10 pixels

for 1,024-by-1,024 acquisitions. For RGC boutons, we set 8 for *params.K* and 2.5-by-2.5 pixels

for *params.gSig*. We manually selected putative cells that had a uniformly-filled round shape of

approximately 10-20 μm in size, and putative boutons to be below 2-3 μm. We then converted

these selected components into binary spatial masks and performed two iterations of masked

CNMF in CaImAn on the entire time-lapse recordings. We used the resulting spatial components

(*estimates.A*; e.g., **Figs. 2C,F and 3A; Supplementary Fig. 2A**) and deconvolved neural

activities, represented as spike estimations based on the df/f calcium transients (*estimates.S*; e.g.,

Supplementary Fig. 1), for subsequent analyses.

Analysis of neuronal activity contextual modulation

First, the estimated spiking activity of each putative neuronal or axonal component in each

analysis epoch was extracted from the entire time lapse and converted into z-scores (Matlab:

zscore): $Z(t)$. To compare the aversive stimulus response across analysis epochs, we then

calculated the average response across trials, $z(t) = \langle Z(t) \rangle_{\text{trials}}$, for each cell for each epoch (e.g.,

Fig. 2C,F), excluding those trials with excessive vertical platform excursion (>0.25 mm on

average). We then selected responsive cells for subsequent analyses that had $z(t)$ exceeding a

threshold in at least one analysis epoch. We varied this threshold value (8 recordings with 1, e.g.,

SFig. 1B; 22 with 1.5; and 11 with 2) to maintain a balanced representation of responsive cells

across recordings (ranging from 22 to 99 cells; 41 recordings in total).

For each responsive neuron, we took an average of $z(t)$ over time during the aversive

stimulus presentation period (2 s) for each analysis epoch (e.g., **Fig. 2D,G**): $\bar{z} = \langle z(t) \rangle_t$. We then

subtracted the average of $\bar{z}$ across the three epochs for normalization: $\Delta z = \bar{z} - \langle \bar{z} \rangle_{\text{epochs}}$, and used

this quantity Δz for the population analysis (e.g., **Fig. 2E,H**). Using this normalized activity of the

responsive cells, Δz, we then performed a 3-way analysis of variance (ANOVA; Matlab: anovan)

with a full interaction model for comparisons across animals. The three variable categories are:

Mouse (41 mice), Experimental Group (WT, Scn2a$^{+/-}$, WT V1-block, Scn2a$^{+/-}$ V1-block), and

Epochs (ES, LS, VL) with the "Mouse" nested within the "Experimental Group." For pairwise

comparisons across epochs within each experimental group, we used the Tukey-Kramer tests

(Matlab: multcompare; e.g., **Fig. 2E,H**).

Pupillometry

Pupil detection and eye-tracking were performed with a region-based convolutional neural

network (R-CNN; He et al. 2017). The network was trained by a set of 260 mouse eye video

frames with manually segmented pupil outlines (Boissonnet et al, 2023). The detected pupil was

parameterized by a 2-dimensional Gaussian curve fit to the pixel map. From these parameters,

we calculated the area of the pupil, $A(t)$, for each frame (6×10^3 pixel^2 = 1 mm^2; e.g., **Fig. 5A,B**).

Mann-Whitney *U*-test was used to compare the median pupil area during the implicit visual task

between WT and Scn2a$^{+/-}$ mice (**Fig. 5C**).

To analyze pupil size dynamics across analysis epochs, we first calculated the proportional

change in pupil size upon each trial of the aversive stimulus presentation, and then took the

average over trials for each epoch: $a(t) = \langle A(t)/A_0 \rangle_{\text{trials}}$, where A_0 is the average pupil size over

the 10 frames before the stimulus onset for each trial (**Fig. 5D-G**). We then took the average over

the stimulation presentation period (2 s) for each analysis epoch: $\bar{a} = \langle a(t) \rangle_t$, and subtracted the

mean across epochs for normalization: $\Delta a = \bar{a} - \langle \bar{a} \rangle_{\text{epochs}}$ (**Fig. 5H-K**). For pairwise comparisons

of Δa across epochs at the population level, Wilcoxon signed-rank tests (Matlab: signrank) were

used with false discovery rate (FDR) correction for multiple comparisons (Matlab: mafdr). We also

performed 2-way ANOVA (Matlab: anovan) to make a comparison across "Experimental Group"

= {WT, Scn2a$^{+/-}$, WT V1-block, Scn2a$^{+/-}$ V1-block} and "Epoch" = {ES, LS, VL} (**Fig. 5H-K**).

Hierarchical Gaussian filter model

To simulate how an ideal Bayesian observer would behave in response to the implicit visual

learning task, we implemented the hierarchical Gaussian filter (HGF) model (Mathys, 2011; **Fig.**

6A) and calculated its parameter trajectories based on the 216 cue-outcome contingencies (**Fig.**

6B). The stimuli presented during the experiments followed a pseudo-random sequence that was

identical for all animals as described in the "visual stimuli" section (**Fig. 1B,C**). Thus, the sequence

of contingencies remained the same across the subjects. For each trial k, we assigned a binary

value to the contingency: 0 for an expected outcome (vertical-aversive or horizontal-omitted) and

1 for an unexpected outcome (vertical-omitted or horizontal-aversive). These values were

represented as a binary vector u that served as the sensory input for the task. From u, the

posterior expectations at three levels (μ_i with $i = 1,2,3$) were calculated as follows.

- At level 1, the posterior is equal to the sensory input: $\mu_1 = u$.

- At level 2, the posterior expectation about $u = 1$ from the previous trial $(k-1)$ is:

$\mu_2 = \mu_2^{(k-1)} + \sigma_2^k \delta_1^k$, where $\delta_1^k = \mu_1^k - s\left(\mu_2^{(k-1)}\right)$ is the prediction error for level 1, with $s(\cdot)$

being a sigmoid function; and $\sigma_2^k = \frac{1}{1/\hat{\sigma}_2^k + \hat{\sigma}_1^k}$ is the standard deviation or variance at level 2.

- At level 3, the posterior expectation regarding environmental volatility is: $\mu_3 = \mu_3^{(k-1)} +$

$\sigma_3^{\frac{k}{2}} w_2^k \delta_2^k$, where the weight is given as $w_2^k = \dfrac{\exp\left(\kappa\mu_3^{(k-1)}+\omega_2\right)}{\sigma_2^{(k-1)}\exp\left(\kappa\mu_3^{(k-1)}+\omega_2\right)}$ and the prediction error

as $\delta_2^k = \dfrac{\sigma_2^k+\left(\mu_2^k-\mu_2^{(k-1)}\right)^2}{\sigma_2^{(k-1)}+\exp\left(k\mu_3^{(k-1)}+\omega_2\right)} - 1.$

From these posterior expectations μ_i, we derived the prior expectations at each level $\hat{\mu}_i$ that

represent the prior prediction about the probability of having $u = 1$ before the sensory input is

presented.

- At level 1, the prior prediction is the sigmoid of the posterior expectation at level 2: $\hat{\mu}_1 =$

$s(\mu_2) = \dfrac{1}{1+\exp(-\mu_2)}.$

- At level 2, the prior prediction about the probability that $\mu_1 = 1$ is: $\hat{\mu}_2 = \mu_2^{(k-1)}.$

- At level 3, the prior prediction about environment volatility, or probability in change of μ_2,

is: $\hat{\mu}_3 = \mu_3^{(k-1)}.$

From here, the variance of the belief at every level $\hat{\sigma}_i$, or the inverse of the predicted precision $\dfrac{1}{\hat{\pi}_i}$,

is given as follows:

- At level 1, the belief variance or imprecision of prediction that $u = 1$ is: $\hat{\sigma}_1^k = \hat{\mu}_1(1 - \hat{\mu}_1).$

- At level 2, the belief variance about the probability that $\mu_1 = 1$ is: $\hat{\sigma}_2^k = \sigma_2^{(k-1)} +$

$\exp\left(\kappa_2\mu_3^{(k-1)} + \omega_2\right).$

- At level 3, the belief variance about environment volatility, or probability in change of μ_2,

is: $\hat{\sigma}_3^k = \sigma_3^{(k-1)} + \exp(\omega_3).$

To calculate the precision-weighted prediction errors ϵ_i, we first calculated prediction errors at

each level:

-   level 1: $\delta_1^k = \mu_1^k - \hat{\mu}_1^k$,

-   level 2: $\delta_2^k = \left(\left(\sigma_2^k + \left(\mu_2^k - \hat{\mu}_2^k \right)^2 \right) \hat{\pi}_2^k \right) - 1$, and

-   level 3: $\delta_3^k = \left(\left(\sigma_3^k + \left(\mu_3^k - \hat{\mu}_3^k \right)^2 \right) \hat{\pi}_3^k \right) - 1$, where $\pi_i = 1/\sigma_i$.

Because precision weights on prediction errors are $\psi_2 = \sigma_2$ at level 2; and $\psi_3 = \pi_3/\hat{\pi}_2$ at level 3,

we have: $\epsilon_2 = \psi_2 \delta_1$ and $\epsilon_3 = \psi_3 \delta_2$.

The uncertainties unc_i are calculated as outcome uncertainty.

-   At level 1, this corresponds to the predicted variance: $unc_1^k = \hat{\sigma}_1^k$.

-   At level 2, the informational uncertainty is: $unc_2^k = s(\mu_2^k)\left(1 - s(\mu_2^k)\right)\sigma_2^k$.

-   At level 3, the environmental uncertainty is: $unc_3^k = \exp\left(\kappa_2 \mu_3^{(k-1)} + \omega_2\right)$.

Finally, the learning rates α_i are calculated as follows:

-   At level 1, $\alpha_1 = \left(s(\mu_2^k) - s\left(\mu_2^{(k-1)}\right)\right)/\left(u - s(\mu_2)\right)$.

-   At level 2, learning rate is equivalent to the precision weight, or variance at level 2: $\alpha_2 =$

857            $\psi_2 = \sigma_2$.

-   At level 3, $\alpha_3 = \dfrac{\exp\left(\kappa_2 \mu_3^{(k-1)} + \omega_2\right)}{2\,\psi_3}$.

In sum, the HGF model consisted of a set of these 13 parameters in five classes across three

hierarchical levels **(Fig. 6B)**:

1. Learning rates: $\alpha_1, \alpha_2, \alpha_3$

2. Prior predictions: $\hat{\mu}_1, \hat{\mu}_2, \hat{\mu}_3$

3. Variances of Belief: $\hat{\sigma}_2, \hat{\sigma}_3$

4. Uncertainties: $unc_1 (= \hat{\sigma}_1), unc_2, unc_3$

5. Precision-weighted prediction errors: ϵ_2, ϵ_3

After generating the model parameter trajectories from the cue-outcome contingency vector u, we

eliminated those trials with non-aversive outcomes to make a fair comparison with the neuronal

response trajectories that included the responses to aversive outcomes only (see below; **Fig. 6C**).

Neuronal trajectory estimation

Because we used six different types of aversive stimuli, and because the cue-outcome stimulus

patterns were presented in blocks, the observed neuronal responses had a certain systematic

bias that needed to be corrected before comparing them to the HGF model parameter trajectories.

To this end, we normalized the neuronal responses in two steps: 1) stimulus-specific normalization

and 2) non-stationarity elimination.

First, for each neuron, we calculated the average response in z-score for each aversive

stimulus trial: $\overline{z}_k = \langle Z(t)\rangle_t$, and subtracted the average per stimulus type across three epochs (i.e.,

18 trials in total per type): $\Delta z_{k|\text{type}} = \overline{z}_{k|\text{type}} - \langle \overline{z}_{k|\text{type}}\rangle_{\text{trials}|\text{type}}$. The resulting concatenated response

Δz_k, termed as "neuronal trajectory" here, represented trial-to-trial differences of the neuronal

activity, independent of the stimulus type.

These neuronal trajectories typically showed a slow decay over trials in each analysis

epoch, likely due to photobleaching. To eliminate such non-stationary components, we next fitted

a power function, $f(k) = a \cdot k^b + c$ (Matlab: fit "power2"), to the population average of all the

neuronal trajectories in the dataset, averaged across epochs. We then subtracted a scaled decay

curve, $d \cdot f(k)$, from the neuronal trajectory of each neuron. The same scalar, d, was used across

epochs to avoid altering the context-dependent neuronal activity. The residual trajectory

$(\Delta z_k^{\text{res}} = \Delta z_k - d \cdot f(k)$; **Fig.6C**) had a Pearson's correlation with the decay within ±0.05, and was

used for the subsequent correlation analyses with the HGF model parameter trajectories.

Correlation analysis between neuronal trajectories and HGF trajectories

We used Pearson's correlation (Matlab: corr) to examine the relationship between the trajectories

of neurons and the HGF model parameters. Specifically, we correlated each of the 13 HGF

parameters (**Fig. 6B**) with every individual neuronal trajectory. The associated *p*-values were then

converted to *q*-values with Storey's FDR methods (Matlab: mafdr) to determine the statistical

significance. We then counted the fraction of SC neurons significantly correlated with each HGF

parameter for each of the four experimental groups (WT, Scn2a$^{+/-}$, WT V1-block, Scn2a$^{+/-}$ V1-

block; **Fig. 6D**, left panels), which we considered as the neuronal representation of the HGF

model. A bias of the correlation polarity was tested with a shuffling test (10,000 repetitions, with

FDR correction). The same procedure was applied to RGC axonal responses (**Fig. 6E,F**).

To compare the neural representations of HGF between the experimental groups, we used

a cosine similarity index: $cos(\theta) = R_i \cdot R_j^{\mathrm{T}} / \|R_i\| \cdot \|R_j\|$, where R_i is a 1-by-13 vector for

experimental group i with each element given by the difference between the fractions of the

positively and negatively correlated cells with each HGF parameter (**Fig. 6G**).

At the population level, we also calculated Pearson's correlation, ρ, between the

population average of the neuronal trajectories $\langle \Delta z_k^{\mathrm{res}} \rangle_{\mathrm{cells}}$ (**Fig. 6C**) and each of the HGF parameter

trajectories (**Fig. 6B**). We then took the average of the absolute correlation values over the

parameters at each level, $\langle |\rho| \rangle_{\mathrm{params|level}}$, to compare how each experimental group represents the

HGF parameters across different levels (**Fig. 6D**, right panels).

References

Ahmadlou, M., Tafreshiha, A., Heimel, J.A. (2017). Visual Cortex Limits Pop-Out in the Superior
Colliculus of Awake Mice. *Cereb. Cortex* **27**, 5772–5783.

Armbruster, B.N., Li, X., Pausch, M.H., Herlitze, S., Roth, B.L. (2007). Evolving the lock to fit the
key to create a family of G protein-coupled receptors potently activated by an inert ligand. *Proc.*
Natl. Acad. Sci. U.S.A. **104**, 5163–5168.

Baron-Cohen, S., Ashwin, E., Ashwin, C., Tavassoli, T., Chakrabarti, B. (2009). Talent in autism:
hyper-systemizing, hyper-attention to detail and sensory hypersensitivity. *Phil. Trans. R. Soc. B*
364, 1377–1383.

Basso, M.A., May, P.J. (2017). Circuits for Action and Cognition: A View from the Superior
Colliculus. *Annu. Rev. Vis. Sci.* **3**, 197–226.

Benavidez, N.L., Bienkowski, M.S., Zhu, M., Garcia, L.H., Fayzullina, M., Gao, L., Bowman, I.,
Gou, L., Khanjani, N., Cotter, K.R., Korobkova, L., Becerra, M., Cao, C., Song, M.Y., Zhang, B.,
Yamashita, S., Tugangui, A.J., Zingg, B., Rose, K., Lo, D., Foster, N.N., Boesen, T., Mun, H.-S.,
Aquino, S., Wickersham, I.R., Ascoli, G.A., Hintiryan, H., Dong, H.-W. (2021). Organization of the
inputs and outputs of the mouse superior colliculus. *Nat. Commun.* **12**, 4004.

Ben-Shalom, R., Keeshen, C.M., Berrios, K.N., An, J.Y., Sanders, S.J., Bender, K.J. (2017).
Opposing Effects on Na V 1.2 Function Underlie Differences Between SCN2A Variants Observed
in Individuals With Autism Spectrum Disorder or Infantile Seizures. *Biol. Psychiatry* **82**, 224–232.

Behrens, T.E.J., Woolrich, M.W., Walton, M.E., Rushworth, M.F.S. (2007). Learning the value of
information in an uncertain world. *Nat. Neurosci.* **10**, 1214–1221.

Bertone, A., Mottron, L., Jelenic, P., Faubert, J. (2005). Enhanced and diminished visuo-spatial
information processing in autism depends on stimulus complexity. *Brain* **128**, 2430–2441.

Boissonnet, T., Tripodi, M., Asari, H. (2023). Awake responses suggest inefficient dense coding
in the mouse retina. *eLife* **12**, e78005.

Branco, T., Redgrave, P. (2020). The Neural Basis of Escape Behavior in Vertebrates. *Annu. Rev.*
Neurosci. **43**, 417–439.

Burnett, L.E., Koppensteiner, P., Symonova, O., Masson, T., Vega-Zuniga, T., Contreras, X.,
Rülicke, T., Shigemoto, R., Novarino, G., Jösch, M. (2022). Subcortical circuit dysfunctions delay
perceptual decision-making in autism models. *bioRxiv* 2022.10.11.511691.

Cang, J., Savier, E., Barchini, J., Liu, X. (2018). Visual Function, Organization, and Development
of the Mouse Superior Colliculus. *Annu. Rev. Vis. Sci.* **4**, 239–262.

Chrysaitis, A.N., Seriès, P. (2023). 10 years of Bayesian theories of autism: A comprehensive
review. *Neurosci. Biobehav. Rev.* **145**, 105022.

Dana, H., Sun, Y., Mohar, B., Hulse, B.K., Kerlin, A.M., Hasseman, J.P., Tsegaye, G., Tsang, A.,
Wong, A., Patel, R., Macklin, J.J., Chen, Y., Konnerth, A., Jayaraman, V., Looger, L.L., Schreiter,
E.R., Svoboda, K., Kim, D.S. (2019). High-performance calcium sensors for imaging activity in
neuronal populations and microcompartments. *Nat. Methods* **16**, 649–657.

Dean, P., Redgrave, P., Westby, G.W.M. (1989). Event or emergency? Two response systems in
the mammalian superior colliculus. *Trends Neurosci.* **12**, 137–147.

De Franceschi, G., Vivattanasarn, T., Saleem, A.B., Solomon, S.G. (2016). Vision Guides
Selection of Freeze or Flight Defense Strategies in Mice. *Curr. Biol.* **26**, 2150–2154.

De Franceschi, Solomon, S.G. (2018) Visual response properties of neurons in the superficial
layers of the superior colliculus of awake mouse. *J. Physiol.* **596**, 6307–6332.

Dobbins, I.G. (2021). Pupil dilation signals recognition salience. *Psychon. Bull. Rev.* **28**, 565–573.

Dombeck, D.A., Khabbaz, A.N., Collman, F., Adelman, T.L., Tank, D.W. (2007). Imaging large-
scale neural activity with cellular resolution in awake, mobile mice. *Neuron* **56**, 43–57.

Evans, D.A., Stempel, A.V., Vale, R., Ruehle, S., Lefler, Y., Branco, T. (2018). A synaptic
threshold mechanism for computing escape decisions. *Nature* **558**, 590–594.

Feinberg, E.H., Meister, M. (2015). Orientation columns in the mouse superior colliculus. *Nature*
519, 229–232.

Friston, K. (2009). The free-energy principle: a rough guide to the brain? *Trends Cogn. Sci.* **13**,
293–301.

Friston, K. (2012). Prediction, perception and agency. *Int. J. Psychophysiol.* **83**, 248–252.

Ganea, D.A., Bexter, A., Günther, M., Gardères, P.-M., Kampa, B.M., Haiss, F. (2020). Pupillary

Dilations of Mice Performing a Vibrotactile Discrimination Task Reflect Task Engagement and

Response Confidence. *Front. Behav. Neurosci.* **14**, 159.

Gomot, M., Blanc, R., Clery, H., Roux, S., Barthelemy, C., Bruneau, N. (2011). Candidate

Electrophysiological Endophenotypes of Hyper-Reactivity to Change in Autism. *J. Autism Dev.*

Disord. **41**, 705–714.

Haker, H., Schneebeli, M., Stephan, K.E. (2016). Can Bayesian Theories of Autism Spectrum

Disorder Help Improve Clinical Practice? *Front Psychiatry* **7**, 107.

He, K., Gkioxari, G., Dollár, P., Girshick, R. (2018). Mask R-CNN. *arXiv*, 1703.06870.

Hoffmann, K.P., Straschill, M. (1971). Influences of cortico-tectal and intertectal connections on

visual responses in the cat's superior colliculus. *Exp. Brain. Res.* **12**, 120–131.

Inayat, S., Barchini, J., Chen, H., Feng, L., Liu, X., Cang, J. (2015). Neurons in the Most Superficial

Lamina of the Mouse Superior Colliculus Are Highly Selective for Stimulus Direction. *J. Neurosci.*

35, 7992–8003.

Johnson, B.P., Lum, J.A.G., Rinehart, N.J., Fielding, J. (2016). Ocular motor disturbances in

autism spectrum disorders: Systematic review and comprehensive meta-analysis. *Neurosci.*

Biobehav. Rev. **69**, 260–279.

Joshi, S., Li, Y., Kalwani, R.M., Gold, J.I. (2016). Relationships between Pupil Diameter and

Neuronal Activity in the Locus Coeruleus, Colliculi, and Cingulate Cortex. *Neuron* **89**, 221–234.

Jure, R. (2022). The "Primitive Brain Dysfunction" Theory of Autism: The Superior Colliculus Role.

Front. Integr. Neurosci. **16**, 797391.

Kato, R., Takaura, K., Ikeda, T., Yoshida, M., Isa, T. (2011). Contribution of the retino-tectal

pathway to visually guided saccades after lesion of the primary visual cortex in monkeys. *Eur. J.*

Neurosci. **33**, 1952–1960.

Larkum, M. (2013). A cellular mechanism for cortical associations: an organizing principle for the

cerebral cortex. *Trends Neurosci.* **36**, 141–151.

Lawson, R.P., Mathys, C., Rees, G. (2017). Adults with autism overestimate the volatility of the
sensory environment. *Nat Neurosci.* **20**, 1293–1299.

LeDoux, J.E. (2012). Evolution of human emotion: a view through fear. *Prog. Brain. Res.* **195**,
431–442.

Li, Y.-T., Turan, Z., Meister, M. (2020). Functional Architecture of Motion Direction in the Mouse
Superior Colliculus. *Curr. Biol.* **30**, 3304–3315.e4.

Loewenfeld, I.E. (1958). Mechanisms of reflex dilatation of the pupil; historical review and
experimental analysis. *Doc. Ophthalmol.***12**, 185–448.

Lomber, S.G. (2002). Learning to see the trees before the forest: Reversible deactivation of the
superior colliculus during learning of local and global visual features. *Proc. Natl. Acad. Sci. U.S.A.*
99, 4049–4054.

Lomber, S.G., Payne, B.R., Cornwell, P. (2001). Role of the superior colliculus in analyses of
space: superficial and intermediate layer contributions to visual orienting, auditory orienting, and
visuospatial discriminations during unilateral and bilateral deactivations. *J. Comp. Neurol.* **441**,
44–57.

Marco, E.J., Hinkley, L.B.N., Hill, S.S., Nagarajan, S.S. (2011). Sensory Processing in Autism: A
Review of Neurophysiologic Finding. *Pediatr. Res.* **69**, 48R-54R.

Mathys, C. (2011). A Bayesian foundation for individual learning under uncertainty. *Front. Hum.*
Neurosci. **5**, 39.

Mathys, C.D., Lomakina, E.I., Daunizeau, J., Iglesias, S., Brodersen, K.H., Friston, K.J., Stephan,
K.E. (2014). Uncertainty in perception and the Hierarchical Gaussian Filter. *Front. Hum. Neurosci.*
8, 825.

May, P.J. (2006). The mammalian superior colliculus: laminar structure and connections. *Prog.*
Brain Res. **151**, 321–378.

Molotkov, D., Ferrarese, L., Boissonnet, T., Asari, H. (2023). Topographic axonal projection at
single-cell precision supports local retinotopy in the mouse superior colliculus. *bioRxiv*
2022.03.25.485790.

Ogasawara, K., McHaffie, J.G., Stein, B.E. (1984). Two visual corticotectal systems in cat. *J.*
Neurophysiol. **52**, 1226–1245.

Patel, J., Lukkes, J.L., Shekhar, A. (2018). Overview of genetic models of autism spectrum
disorders. *Prog. Brain Res.* **241**, 1–36.

Pellicano, E., Burr, D. (2012). When the world becomes 'too real': a Bayesian explanation of
autistic perception. *Trends Cogn. Sci.* **16**, 504–510.

Planells-Cases, R., Caprini, M., Zhang, J., Rockenstein, E.M., Rivera, R.R., Murre, C., Masliah,
E., Montal, M. (2000). Neuronal death and perinatal lethality in voltage-gated sodium channel
alpha(II)-deficient mice. *Biophys. J.* **78**, 2878–2891.

Pnevmatikakis, E.A., Soudry, D., Gao, Y., Machado, T.A., Merel, J., Pfau, D., Reardon, T., Mu,
Y., Lacefield, C., Yang, W., Ahrens, M., Bruno, R., Jessell, T.M., Peterka, D.S., Yuste, R.,
Paninski, L. (2016). Simultaneous Denoising, Deconvolution, and Demixing of Calcium Imaging
Data. *Neuron* **89**, 285–299.

Ruediger, S., Scanziani, M., (2020). Learning speed and detection sensitivity controlled by distinct
cortico-fugal neurons in visual cortex *eLife* **9**, e59247.

Sahibzada, N., Dean, P., Redgrave, P. (1986). Movements resembling orientation or avoidance
elicited by electrical stimulation of the superior colliculus in rats. *J. Neurosci.* **6**, 723–733.

Sanders, S.J., Murtha, M.T., Gupta, A.R., Murdoch, J.D., Raubeson, M.J., Willsey, A.J., Ercan-
Sencicek, A.G., DiLullo, N.M., Parikshak, N.N., Stein, J.L., Walker, M.F., Ober, G.T., Teran, N.A.,
Song, Y., El-Fishawy, P., Murtha, R.C., Choi, M., Overton, J.D., Bjornson, R.D., Carriero, N.J.,
Meyer, K.A., Bilguvar, K., Mane, S.M., Šestan, N., Lifton, R.P., Günel, M., Roeder, K., Geschwind,
D.H., Devlin, B., State, M.W. (2012). De novo mutations revealed by whole-exome sequencing
are strongly associated with autism. *Nature* **485**, 237–241.

Satterstrom, F.K., Kosmicki, J.A., Wang, J., Breen, M.S., De Rubeis, S., An, J.-Y., Peng, M.,
Collins, R., Grove, J., Klei, L., Stevens, C., Reichert, J., Mulhern, M.S., Artomov, M., Gerges, S.,
Sheppard, B., Xu, X., Bhaduri, A., Norman, U., Brand, H., Schwartz, G., Nguyen, R., Guerrero,
E.E., Dias, C., Autism Sequencing Consortium, iPSYCH-Broad Consortium, Betancur, C., Cook,
E.H., Gallagher, L., Gill, M., Sutcliffe, J.S., Thurm, A., Zwick, M.E., Børglum, A.D., State, M.W.,
Cicek, A.E., Talkowski, M.E., Cutler, D.J., Devlin, B., Sanders, S.J., Roeder, K., Daly, M.J.,

Buxbaum, J.D. (2020). Large-scale exome sequencing study implicates both developmental and
functional changes in the neurobiology of autism. *Cell* **180**, 568–584.e23.

Schröder, S., Steinmetz, N.A., Krumin, M., Pachitariu, M., Rizzi, M., Lagnado, L., Harris, K.D.,
Carandini, M. (2020). Arousal Modulates Retinal Output. *Neuron* **107**, 487-495.e9.

Shang, C., Chen, Z., Liu, A., Li, Y., Zhang, J., Qu, B., Yan, F., Zhang, Y., Liu, W., Liu, Z., Guo, X.,
Li, D., Wang, Y., Cao, P. (2018). Divergent midbrain circuits orchestrate escape and freezing
responses to looming stimuli in mice. *Nat. Commun.* **9**, 1232.

Spratt, P.W.E., Ben-Shalom, R., Keeshen, C.M., Burke, K.J., Clarkson, R.L., Sanders, S.J.,
Bender, K.J. (2019). The Autism-Associated Gene Scn2a Contributes to Dendritic Excitability and
Synaptic Function in the Prefrontal Cortex. *Neuron* **103**, 673–685.e5.

Sridharan, D., Steinmetz, N.A., Moore, T., Knudsen, E.I. (2017). Does the Superior Colliculus
Control Perceptual Sensitivity or Choice Bias during Attention? Evidence from a Multialternative
Decision Framework. *J. Neurosci.* **37**, 480–511.

Suzuki, Y., Minami, T., Nakauchi, S. (2018). Association between pupil dilation and implicit
processing prior to object recognition via insight. *Sci. Rep.* **8**, 6874.

Tang, L., Higley, M.J. (2020). Layer 5 Circuits in V1 Differentially Control Visuomotor Behavior.
Neuron **105**, 346-354.e5.

Tinbergen, N. (1951). The study of instinct, The study of instinct. Clarendon Press/Oxford
University Press, New York, NY, US.

Wang, L., Sarnaik, R., Rangarajan, K., Liu, X., Cang, J. (2010). Visual Receptive Field Properties
of Neurons in the Superficial Superior Colliculus of the Mouse. *J. Neurosci.* **30**, 16573–16584.

Wang, C.-A., Boehnke, S.E., White, B.J., Munoz, D.P. (2012). Microstimulation of the Monkey
Superior Colliculus Induces Pupil Dilation Without Evoking Saccades. *J. Neurosci.* **32**, 3629–
3636.

Wang, Q., Burkhalter, A. (2013). Stream-Related Preferences of Inputs to the Superior Colliculus
from Areas of Dorsal and Ventral Streams of Mouse Visual Cortex. *J. Neurosci.* **33**, 1696–1705.

Wei, P., Liu, N., Zhang, Z., Liu, X., Tang, Y., He, X., Wu, B., Zhou, Z., Liu, Y., Li, J., Zhang, Y.,

Zhou, X., Xu, L., Chen, L., Bi, G., Hu, X., Xu, F., Wang, L. (2015). Processing of visually evoked

innate fear by a non-canonical thalamic pathway. *Nat. Commun.* **6**, 6756.

Yilmaz, M., Meister, M. (2013). Rapid Innate Defensive Responses of Mice to Looming Visual

Stimuli. *Curr. Biol.* **23**, 2011–2015.

Zhao, X., Liu, M., Cang, J. (2014). Visual Cortex Modulates the Magnitude but Not the Selectivity

of Looming-Evoked Responses in the Superior Colliculus of Awake Mice. *Neuron* **84**, 202–213.

Zingg, B., Chou, X., Zhang, Z., Mesik, L., Liang, F., Tao, H.W., Zhang, L.I. (2017). AAV-Mediated

Anterograde Transsynaptic Tagging: Mapping Corticocollicular Input-Defined Neural Pathways

for Defense Behaviors. *Neuron* **93**, 33–47.