The scramblases VMP1 and TMEM41b are required for primitive

endoderm specification by targeting WNT signaling

Markus Holzner1, Tea Sonicki1, Hugo Hunn1, Federico Uliana2,3, Weijun Jiang1,

Vamshidhar R. Gade2, Karsten Weis2, Anton Wutz1#, and Giulio Di Minin1#

1 Institute of Molecular Health Sciences, Department of Biology, ETH Zurich, Zurich,

Switzerland

2 Institute of Biochemistry, Department of Biology, ETH Zurich, Zurich, Switzerland

3 Current address: Johannes Gutenberg University of Mainz, 55128 Mainz, Germany

corresponding authors: awutz@ethaz.ch (AW); giulio.diminin@biol.ethz.ch (GDM)

ABSTRACT

The ER resident proteins VMP1 and TMEM41b share a conserved DedA domain,

which confers lipid scramblase activity. Loss of either gene results in embryonic

lethality in mice and defects in autophagy and lipid droplet metabolism. We set out to

investigate their role in pluripotency and specification. For this purpose, we generated

Vmp1 and Tmem41b mutations in mouse embryonic stem cells (ESCs). We observed

that ESCs carrying mutations in Vmp1 and Tmem41b show robust self-renewal and

an unperturbed pluripotent expression profile but accumulate LC3-positive

autophagosomes and lipid droplets consistent with defects in autophagy and lipid

metabolism. ESCs carrying combined mutations in Vmp1 and Tmem41b can

differentiate into a wide range of embryonic cell types. However, differentiation into

primitive endoderm-like cells in culture is impaired, and the establishment of extra-

embryonic endoderm stem (XEN) cells is delayed. Mechanistically, we show the

deregulation of genes that are associated with WNT signaling. This is further confirmed

by cell surface proteome profiling, which identified a significant reduction of the WNT

receptor FZD2 at the plasma membrane in Vmp1 and Tmem41b double mutant ESCs.

Importantly, we show that transgenic expression of Fzd2 rescues XEN differentiation.

Our findings identify the role of the lipid scramblases VMP1 and TMEM41b in WNT

signaling during extra-embryonic endoderm development and characterize their distinct and overlapping functions.

INTRODUCTION

Vesicle trafficking and membrane synthesis are intricately interconnected within cellular signaling networks. Lipid scramblases, with their property to shape lipid membrane composition, play a crucial role in facilitating inter-organelle communication. Beyond their integral involvement in maintaining cell functionality, recent studies have unveiled their implications in various pathological conditions, such as viral infection [1-5], cancer development [6-11], and inflammatory diseases [12-16]. The vacuole membrane protein 1 (VMP1) and transmembrane protein 41B (TMEM41B) are lipid scramblases with a highly conserved DedA domain [17, 18]. They localize at the ER membrane and are distinguished from other ER-associated flippases or scramblases by the property to facilitate bi-directional lipid diffusion in an ATP- and Ca2+-independent manner. VMP1 and TMEM41B are necessary for autophagosome formation [19, 20], lipid droplet metabolism [17, 21], and lipoprotein formation [22]. Additional reports proposed a more global role in ER protein homeostasis [23] and vesicle transport [24]. Despite multifaceted functions, their absence does not influence cell survival significantly. Compensatory mechanisms shared by VMP1 and TMEM41B or other ER scramblases likely maintain cellular homeostasis. While VMP1 depletion leads to a pronounced defect in autophagosome assembly [19, 20], TMEM41B appears to play a pivotal role in sustaining lipid droplet homeostasis and serves as a crucial factor in RNA virus replication [3, 25]. However, a direct comparison between the activities of VMP1 and TMEM41B has not yet been performed.

VMP1 and TMEM41B are both essential during embryonic development. Vmp1 [21] and Tmem41b [26] deficient mice die around E8.5 with a severe developmental delay. Previously, defects in the visceral endoderm function have been proposed. In particular, loss of Vmp1 has been suggested to block the synthesis and release of lipoproteins from the visceral endoderm. However, mice with defects in lipoprotein biogenesis experience mortality at a later stage, E10.5 [27, 28]. This observation

suggests a distinct and earlier involvement of VMP1 and TMEM41B in embryo development.

During implantation, the mouse embryo is composed of three lineages: the trophectoderm, the primitive endoderm (PE), and the epiblast (EPI), which is primed to develop into the embryo [29]. The EPI and PE are specified within the inner cell mass (ICM) of the blastocyst. Defects in the formation of these lineages are challenging to characterize *in vivo* as they appear pre-implantation and due to the small embryo size. To investigate the role of Vmp1 and Tmem41b in early embryonic lineage specification, we generated mutations in Vmp1 and Tmem41b in ESCs. We show that a combined Vmp1 and Tmem41b mutation in ESCs results in a specific defect in PE formation and a delayed conversion to extra-embryonic endoderm cells (XEN). This delay is caused by reduced WNT signaling and can be rescued through WNT activation.

RESULTS

Vmp1 and Tmem41b mutant ESCs accumulate LC3+ autophagosomes and lipid droplets

To assess the influence of a Vmp1 or Tmem41b depletion on mouse ESCs and in early embryo development, we engineered genomic deletions using CRISPR/Cas9 targeted mutagenesis. Vmp1 was mutated by introducing a frameshift in exon 2 (Fig.1A). Loss of Vmp1 protein was confirmed in independent mutant ESC clones (Vmp1^{KO}) (Fig.S1A). To abrogate Tmem41b expression, we deleted the genomic region between exons 2 and 3 (Fig.1A), which encodes the first transmembrane domain of the protein (Tmem41b^{KO}). This mutation is predicted to compromise the protein topology. Additionally, the splicing of exon 1 to exon 4 produces a frameshift, suggesting a null allele. The deletion was confirmed on the genomic DNA (Fig.S1B). Tmem41b mRNA levels were strongly reduced (Fig.S1C-D).

Vmp1 and Tmem41b mutant ESCs showed similar morphology (Fig.1B), proliferation (Fig.S1E), and expression of pluripotency markers (Fig.S1F-G) compared to WT ESCs. We confirmed that the pluripotency network was unchanged by a transcriptomic analysis performed in individual Vmp1 and Tmem41b mutant cells. This

analysis also showed that the transcription profile of cells was marginally affected,
consistent with earlier results of ESC characterization (Fig.S1I-J and Table S2).

Considering the role of Vmp1 and Tmem41b in autophagy, we assessed the
accumulation of LC3+ autophagosomes to determine the impact of their mutations.
We found an increase of LC3+ puncta in both mutants compared to WT ESCs (Fig.1C
and Fig.S2A-B). Interestingly, the increased frequency of LC3+ puncta was similar in
Vmp1 (4X) and Tmem41b (3,5X) mutant cells. Next, mutant cells were analyzed for
the retention of lipid droplets (LD). To amplify LDs, we treated all samples with oleic
acid (OA) and visualized LDs by BODIPY staining. We observed that Vmp1 and
Tme41b mutations led to a 4-fold increase of LD numbers with respect to WT cells
(Fig.1D and Fig. S2D-E). These results suggest that both Vmp1 and Tmem41b
contribute to autophagosome and lipid droplet metabolism in ESCs, recapitulating
earlier studies performed in other cellular systems [17, 20]. The direct comparison of
both mutations revealed a similar contribution of VMP1 and TMEM41B to both
processes.

Vmp1 and Tmem41b activity is not required for ESC self-renewal
We generated double mutant cell lines to characterize potential redundant functions
between VMP1 and TMEM41B. A Tmem41b mutation was engineered in Vmp1 mutant
ESCs (Double KO) (Fig.S1A-D). Several independent mutant clones were obtained.
Notably, Double KO cells showed a regular ESC morphology (Fig.S1H) and
proliferation rate (Fig.S1E). The absence of Vmp1 and Tmem41b did not affect
secretory compartments, as ER, Golgi apparatus, and late endosomes appeared
normal in mutant cells (Fig.S3A). Transcriptomics analysis of Double KO cells showed
a limited perturbation in gene expression (Fig.1E). The combined mutations did not
result in a greater number of differentially regulated genes compared to Vmp1[KO] cells
(Fig.S3C). Minimal overlap of differentially expressed genes was detected among
Vmp1, Tmem41b, and Double KO mutant cell lines (Fig.S3B-C). Additionally, we did
not observe enrichment for genes associated with defined cellular processes (Fig.1F,
Fig.S3D-E). The pluripotency network, as also shown by IF staining (Fig.S1F-G), was
not perturbed in Double KO cells. We conclude that Vmp1 and Tmem41b are not
essential for ESC self-renewal.

Vmp1 and Tmem41b have redundant functions in autophagosome and lipid droplet formations

LC3 staining showed that Double KO cell lines present an exacerbated phenotype in autophagosome accumulation compared to single mutants. Although the total number of LC3 positive autophagosomes per cell remained unchanged (Fig.1G and Fig.S2A-B), we observed an increased size. 14% of LC3 clusters in the Double KO had a larger size (>3µm), compared to the 8% of autophagosomes across the single mutations and 3% in the WT condition (Fig.1H and Fig.S2C).

The simultaneous depletion of Vmp1 and Tmem41b also had a greater impact on LD formation. Double KO cells showed a drastic increase of LD number in the absence of oleic acid treatment, whereas in WT and Vmp1 and Tmem41b mutant ESCs, LDs became detectable after OA treatment (Fig.1I-J and Fig.S2D-E). As BODIPY could also stain other lipid-containing compartments, including autophagosomes [30], we confirmed the enriched lipid droplets in Double KO cells by refractive index tomography, which is a labelling-independent technique (Fig.1K and Fig.S2F-G). Together, these results indicate that Vmp1 and Tmem41b have overlapping functions in ESCs, as the combined mutation of both genes exacerbates autophagosome and lipid droplet accumulation.

Loss of Vmp1 and Tmem41b affects ESC potential to differentiate into primitive endoderm

To evaluate the effect of Vmp1 and Tmem41b on the differentiation potential of ESCs, we used embryoid body (EB) differentiation (Fig.2A). After 6 days of differentiation, EBs derived from WT ESCs showed increased expression of ectoderm (Sox1 and Fgf5), mesoderm (Actc1 and Brachyury) and endoderm (Gata6 and Dab2) genes (Fig.2B and Fig.S4A), while the expression of pluripotency markers (Nanog and Pou5f1) was lost (Fig.2B and Fig.S4A). We evaluated variations in the differentiation potential of mutant cells by comparing the expression of these markers by RT-qPCR. For each mutant, the differentiation potential of two independent clones was tested. Vmp1, Tmem41b, and double mutant ESCs formed well-defined EBs that resembled those of WT cells. All cell lines efficiently differentiated into the ectoderm and mesoderm lineages (Fig.2C and Fig.S4B-C). Notably, Double KO EBs showed a decreased expression of the endoderm markers Dab2 (100%) and Gata6 (57% of experiments) (Fig.2C and Fig.S4C). Neither single mutant showed a similar failure in

Dab2 expression, even if some variability was detected between the analyzed clones.
In the single mutants, we noticed a decreased endoderm specification, as shown by
Gata6 expression levels in experiments performed with Vmp1^{KO} (56% of experiments)
and Tmem41b^{KO} (58%) cells (Fig.2C and Fig.S4C). Our results indicate that combined
mutation of Vmp1 and Tmem41b leads to reduced Dab2 expression and predisposes
to variability in the expression of other endodermal genes.

We confirmed the differentiation potential of Double KO cells into ectodermal
lineages by generating SOX1-positive spinal cord organoids (Fig.S4D). Mesodermal
differentiation was inferred from the ability of double mutant EBs to form contractile
cardiomyocytes after plating (Video S1-2). We subsequently sectioned day 6 EBs to
evaluate protein expression and distribution of endoderm markers by IF. Expression
of GATA4, GATA6, and SOX17 was detected in all conditions but noticeably decreased
in Double KO compared to WT EBs (Fig.2D-E). Interestingly, decreased GATA4 and
GATA6 staining was pronounced in EB border regions, which resemble the primitive
endoderm layer [31]. In line with these findings, we confirmed the absence of DAB2 in
Double KO EBs (Fig.2F). DAB2 is specifically expressed in the primitive endoderm,
while GATA4 and GATA6 are known to be also expressed in definitive endoderm.

The significant reduction in both mRNA and protein levels of Dab2, along with
the weakened expression of endoderm markers at the edges of embryoid bodies,
provides evidence that the absence of Vmp1 and Tmem41b compromises the
formation of primitive endoderm.

Combined Vmp1 and Tmem41b mutations delay ESC differentiation into extra-
embryonic endoderm stem (XEN) cells
We tested the ability of Double KO ESCs to differentiate into extra-embryonic
endoderm stem (XEN) cells to characterize endodermal differentiation further. XEN
cells share characteristics with PE cells including cell signaling, gene expression, and
differentiation potential [32]. Double KO and WT cells were differentiated into XEN
cells with Activin A and Retinoic Acid [33]. After 6 days, WT ESCs efficiently
differentiated into XEN clusters (Fig.3A), exhibiting a stellate and refractile morphology
[33]. In Double KO cells, the formation of XEN clusters was substantially reduced. We
characterized XEN identity by analyzing the expression of XEN markers by RT-qPCR
(Fig.S5A) and immunostaining (Fig.3B and Fig.S5B). It is worth noting that the small
number of derived XEN cells from Double KO ESCs can be expanded for multiple

passages, as shown in Fig.S5C. This result suggests that the loss of Vmp1 and Tmem41b does not entirely prevent XEN specification, but rather causes a delay in the differentiation process.

WNT signaling is decreased in Vmp1 and Tmem41b mutant cells during XEN specification

To characterize the effect of Vmp1 and Tmem41b double mutation on XEN specification, we analyzed the transcriptomic profile of Double KO and WT cells during the differentiation process. On day two of differentiation, we detected the induction of XEN-associated genes [34] in WT cells. However, in Double KO cells these genes were weakly expressed (Fig.3C-D and Fig.S5E). Intriguingly, we did not observe any perturbation in the transcription of genes involved in lipid metabolism, autophagy, or ER maintenance (Fig.S5D).

Instead, Double KO cells show consistent deregulation of genes associated to the WNT pathway (Fig.3E-F). A close analysis of these genes allowed to identify several well-established WNT-target genes, including Dkk, Frzb, Sox17, Wnt3, and Ccdnd1 [35-38]. Their down-regulation strongly suggested a decreased activity of the WNT pathway in Double KO cells and might explain the observed defect in XEN specification [39]. Interestingly, Ddit3 was strongly upregulated in Double KO cells. DDIT3 acts as an inhibitor of the WNT signaling by repressing the transcriptional activity of the TCF7/TCF4 complex [40]. We confirmed increased levels of the DDIT3 protein in Double KO cells by Western analysis (Fig.3G) and quantified a fourfold increase over WT cells (Fig.3H). Considering that Ddit3 is not reported to be a WNT-targeted gene, but instead, its expression is controlled by VMP1 levels [41], we verified if DDIT3 increase might explain WNT-target gene repression in mutant cells. For this purpose, we depleted Ddit3 in WT and mutant cells by RNA interference. After two days of differentiation, the transcription of XEN-specific genes was evaluated by qPCR. Despite efficient siRNA-mediated reduction of Didt3 mRNA levels (Fig. 3I), we did not observe an increase in the expression of XEN marker genes in Double KO cells.

These results indicate that the increased level of DDIT3 in mutant cells is not the only cause of the XEN specification defect caused by Vmp1 and Tmem41b mutations.

Vmp1 and Tmem41b mutations reduce FZD2 abundance at the plasma membrane

To identify potentially affected factors of cell signaling, we investigated the influence of the Vmp1 mutation on proteins at the plasma membrane (PM) in ESCs. For this purpose, we enriched surface proteins in mutant and WT cells using aminooxy-biotinylation and characterized their abundance by TMT mass spectrometry (Fig.S6A-B). Our analysis revealed that a limited subset of proteins showed differential abundance at the plasma membrane in mutant cells, with 26 proteins significantly down-regulated and 12 up-regulated (out of 1417 detected proteins) (Fig. 4A and Table S4). This finding aligns with our RNA expression profiles, indicating that Vmp1 and Tmem41b mutations has a limited effect on the protein levels of ESCs. It also supports the conclusion that, despite defects in autophagy and lipid droplet formation, the secretory pathway remains largely intact in the absence of Vmp1. GO-term analysis highlighted that proteins more abundantly present at the PM in Vmp1 mutant cells are characterized by an ER-Golgi localization, suggesting partial defects in retrograde trafficking (Fig. 4B). Moreover, we noticed that the abundance of four GPCRs was reduced at the plasma membrane of mutant cells (Fig. 4B). Among them, we found the WNT-receptor FZD2 (Fig. 4A and Table S4). Other Frizzled receptors—FZD5, FZD7, and FZD10— were detected at the PM but remained un-affected (Fig. 4C and S6C). The decrease of FZD2 protein abundance at the PM was independent of Fzd2 transcription, as no difference in Fzd2 mRNA was detected between WT and mutant cells (Fig. S6D).

To confirm these findings and assess whether FZD2 reduction contributes to the XEN specification defects in Double KO cells, we generated ESCs with constitutive expression of HA-Fzd2 across Vmp1KO, Double KO, and WT genotypes. We evaluated HA-FZD2 translocation to the plasma membrane using a cell-impermeable biotinylation reagent followed by purification of biotinylated proteins (Fig. 4D). We confirmed that FZD2 is less abundant at the plasma membrane in Vmp1 mutant cells, validating previous MS results (Fig. 4D-E). Notably, Double KO cells exhibited an even greater reduction in FZD2 levels at the PM, suggesting an additive effect of the Vmp1 and Tmem41b mutations. Similar results were observed when analyzing HA-FZD2 enrichment at the plasma membrane in Double KO and WT cells during XEN specification (Fig. 4F). In this experiment, we noticed a marked reduction of the upper

form of the FZD2 doublets in mutant cells. This suggests that FZD2 is subjected to
different post-translational modifications in WT and mutant cells.

FZD2 over-expression rescues XEN differentiation defects in Vmp1 and
Tmem41b mutant cells
Our observation of deregulated WNT target genes and a reduced abundance of FZD2
at the PM indicated that WNT signaling defects might impair XEN specification. To
investigate if indeed the WNT signaling cascade is affected, we quantified β-CATENIN,
down-stream protein, and major signal transducer of the WNT pathway. We observed
a marked decrease of β-CATENIN protein in Double KO compared to WT cells (Fig.
5A). This difference was even stronger, specifically analyzing the cytoplasmic fraction
of β-CATENIN that is not associated with cadherins. To explore if the defect in XEN
differentiation of double mutant cells can be rescued by chemically activating WNT
signaling, we employed the GSK3 kinase inhibitor (CHIR) to prevent β-CATENIN
degradation. GSK3 inhibition had no effect on XEN specification in WT ESCs.
However, GSK3 inhibition significantly enhanced XEN cell cluster formation in double
mutant ESCs (Fig. 5B and S7A-B). Finally, to investigate whether also overexpression
of FZD2 would rescue XEN differentiation defects in Double KO cells, we generated
subclones with high FZD2 expression for both Double KO and WT ESCs (Fig. S7C).
The increased amount of FZD2 alone did not further stimulate the WNT pathway or
XEN specification in WT cells (Fig. 5C-D and S7D-F). However, FZD2 overexpression
in Double KO cells was sufficient to restore their potential for differentiation into XEN
cells. Collectively, our data implicate Vmp1 and Tmem41b in WNT signaling during
primitive endoderm specification.

DISCUSSION
Vmp1 and Tmem41b have recently been identified in a number of genetic screens
designed to study various cellular processes, including autophagy, lipid metabolism,
virus maturation, and ER stress [22]. In this study, we have unveiled the role of Vmp1
and Tmem41b in regulating protein secretion. Our findings indicate that their presence
is essential for the trafficking of the WNT receptor FZD2 to the plasma membrane. A
drastic reduction of FZD2 in mutant cells impairs WNT signals and causes an XEN cell
differentiation defect. Our data, therefore, implicate a function of Vmp1/Tmem41b in

WNT signaling. Given that Vmp1 expression and activity are influenced by various
stimuli [42, 43], the interaction between Vmp1 and Tmem41b within the secretory
pathway is critical for physiological and pathological processes.

The pleiotropic function and peculiar structure of the DedA domain found in both
scramblases have prompted multiple studies addressing its mode of action and role in
cellular homeostasis. An important point under evaluation is whether Vmp1 and
Tmem41b work cooperatively or show specialization for different cellular processes.
To address this question, we targeted Vmp1 and Tmem41b genes in ESCs and
introduced mutations abrogating their expression. This allowed us to directly compare
the specificity of Vmp1 and Tmem41b in multiple processes. We report that their
absence similarly affects autophagy and lipid droplet accumulation. Moreover, we
show that the simultaneous loss of both genes synergizes and exacerbates defects of
single mutants, particularly in LD formation. Notably, loss of both genes did not affect
self-renewal in ESCs and had only subtle effects on gene expression. Our
characterization shows that Vmp1 and Tmem41b perform distinct and overlapping
functions in ESCs. It will be interesting to understand if variating expression levels of
Vmp1 and Tmem41b can account for their redundancy or dominance in different
cellular systems.

The embryonic lethality caused by Vmp1 and Tmem41b mutations [21, 26]
during development may be explained by the defective primitive endoderm
specification that we detected *in vitro*. Embroid body experiments and the direct
differentiation of ESCs to extra-embryonic endoderm stem (XEN) cells support a
specific delay in PE formation in double mutant cells. Interestingly, it has been reported
that other mutations leading to PE delay cause retardation and embryo lethality at E8.5
[44], mirroring phenotypes of Vmp1 or Tmem41b mutant embryos. While both
scramblases are needed in mice for regular development, the depletion of both genes
is required to cause a differentiation delay in ESCs. More permissive *in vitro* growth
conditions may explain the requirement for the depletion of both genes to manifest this
phenotype in cell culture. It is worth noting that single Vmp1^{KO} or Tmem41b^{KO} cells
also showed subtle defects in endoderm specification in embryoid bodies, albeit less
consistently than Double KO cells (Fig.2C). The finding of WNT signaling deregulation
in Vmp1 and Tmem41b XEN cells explains the specification delay detected in mutant
cells. This is consistent with earlier reports that the WNT/TCF7L1 transcriptional
response is necessary *in vitro* and *in vivo* for PE expansion [39]. Retardation – not

impairment – of PE specification induced by Vmp1 or Tmem41b mutations may permit
embryo implantation but the accumulation of defects likely blocks development during
gastrulation.

Overexpression of the FZD2 receptor rescues the defects of Vmp1 and
Tmem41b double mutant cells during XEN specification. We propose that the
decreased availability of FZD2 at the plasma membrane in mutant cells causes a delay
in WNT signaling activation and PE defects. The observation that Fzd2 mutant mice
exhibit developmental defects at later stages [45] suggests the possibility of
compensatory mechanisms among different Frizzled receptors *in vivo*. Future studies
will focus on the deep characterization of other potential GPCRs subjected to
Vmp1/Teme41b regulation. While our results indicate that the increased expression of
the WNT repressor DDIT3 alone does not fully account for the delay in XEN
specification, it may still contribute to this delay during embryonic development. The
elevated levels of DDIT3 in mutant cells might exert an additive effect, further reducing
WNT pathway activity.

Vmp1 and Tmem41b emerge as key factors in the secretary machinery of cells.
Beyond their established roles in lipid and membrane metabolism, we demonstrate
their involvement in regulating protein trafficking at the plasma membrane. Notably,
this effect is transcriptional independent, as evidenced by expression analyses and
experiments in cells with constitutive FZD2 expression. Furthermore, our findings
indicate that Vmp1/Tmem41b depletion does not cause a global disruption of protein
secretion but shows a specificity for some sub-classes of secreted proteins, such as
GPCRs. In denaturing conditions, the upper forms of the FZD2 protein are less
abundant in mutant cells. This finding suggests potential post-translational
modification differences between the two conditions. GPCRs are tightly regulated at
the ER-Golgi interface, where proper folding is essential for their route in the secretory
pathway. In the Golgi, GPCRs undergo extensive glycosylation, which alters their
molecular weight and activity. These results suggest a role of Vmp1/Tmem41b in
promoting the maturation and stability of a specific class of transmembrane proteins.

The cellular systems generated in our study will help clarify the mechanisms
shared by these scramblases in controlling protein secretion. Furthermore, the role of
Vmp1 and Tmem41b in modulating the WNT cascade may extend to other signaling
networks, revealing other cross-connections between cellular pathways and metabolic
processes.

MATERIALS AND METHODS

Cell Culture

Mouse ES cells were maintained as previously described in chemically defined 2i medium with added leukemia inhibitory factor (LIF) as described in Di Minin et al. [46]. Base media composition: DMEM (Gibco, #41965039), 15% FBS (Biowest, #S1810-500) 1% sodium pyruvate (Gibco, 11360-070), 1% non-essential amino acids (Gibco, #11140-035). Lif (2000 U/mL, homemade), PD0325901 (1 µM, AxonMedchem, #391210-10-9) and Chiron (3 µM, AxonMedchem, #252917-06-9) are added fresh before use.

EB differentiation

To obtain embryonic bodies (EBs), 200k ESCs were plated on Day1 in ESCs base media without 2i and LIF (see cell culture) into a well of a Sphericalplate5D (Kugelmeiers). On Day4, EBs have formed and are transferred to a pHEMA coated 6cm culture dish and cultured until Day6. EBs are then collected and subjected to RNA extraction and RT-qPCR analysis or prepared for sectioning and immunofluorescence. In brief, EBs are fixed in 4% PFA for 30 min, cryoprotected in 10%, 20% and 30% sucrose and then embedded and frozen in Tissue Freezing Medium (Leica, #14020108926). 15 µm sections were prepared on a cryotome, mounted onto Superfrost Plus Adhesion slides (VWR international GmBH, #631-0108) and processed for immunofluorescence staining.

Cardiomyocyte differentiation

EBs were formed as previously described. On Day4, EBs were collected and seeded onto a non-treated 10cm dish, maintaining ESC base media. Cardiomyocytes randomly specified and were observed as early as on Day6 of differentiation.

Spinal Cord Organoids (SCOs) formation

For derivation of SCOs, ESCs were plated in EBs media (DMEM- F12 (Gibco) and Neurobasal Medium (Gibco) (1:1 ratio) supplemented with 200 mM L-glutamine and 10% of Knockout serum replacement) on Sphericalplate 5D dishes (Kugelmeiers AG)

403   {Holzner, 2024 #139}. On day 2, SCOs were transferred in 10 cm tissue culture dishes

404   and treated with RA (10 nM). SCOs were collected and processed for analysis at day

405   5.

Xen Differentiation

XEN cells were obtained following a protocol by Niakan et al. with minor changes [33].

In brief, 230k cells were plated in XEN medium on a gelatin-coated 3cm dish. The next

day, the media was replaced with Xen medium containing 0.1 µM retinoic acid

(Thermo Fischer, #17110052) and 10 ng/mL Activin A (Peprotech, #120-14). On Day4,

the medium was again replaced by XEN medium, no further supplements were added,

and on Day6 samples were either analyzed by IF or qPCR.

Rescue of double KO Xen differentiation was performed by adding 1 µM Chiron

(AxonMedchem, #252917-06-9) at Day2, together with the retinoic acid and Activin A.

Media was changed on day 4 to Xen medium. XEN medium composition: RPMI 1640

(Gibco, #61870010), GlutaMAX^{TM} (Gibco, #61870036), 15% FBS (Biowest, #S1810-

500), 0,1 mM ß-mercaptoethanol (Sigma, #BCCB9882).

CRISPR-Cas9 editing to generate mutant ESC lines

Vmp1: the gRNA (gagacgcatagcaatgagta) was cloned into a px330 vector (Addgene,

#158973). Per 3cm well, 300k ESCs were plated and transfected with 2.5 µg of gRNA

containing plasmid, as well as 0.25 µg of a GFP expressing plasmid (pCMV-GFP).

GFP-positive cells were enriched by FACS and plated as single cells. Colonies were

picked. Analysis by WB was used to identify identified VMP1 KO lines. Tmem41b: the

gRNAs (gtatgtttgacctgggcgaa and gagtgacatgtggaaatcag) were cloned into px330

vectors, and KO clones were derived as described above. PCR analysis of the edited

locus (FW: atcttggacaggggggagttc RV: gaccaggggttcattgtcatt and FW:

cagcacacacctttaatccagc RV: gccacatagcaagcttaagagc) identified KO clones. qPCR of

mRNA from Exon 2 (edited locus, N-terminal end) and Exon 7 (C-terminal end) further

confirmed the absence of mRNA.

Transfection and stable cell line generation

For derivation of *Fzd2*-HA ESCs the PB-HA-Fzd2-IRES-Neo plasmid [46] was used.

Stable expression of the construct was achieved in WT, Vmp1^{KO}, and DoubleKO ESCs

by co-transfection of the Piggybac and the PBase plasmids. Cells were plated by

limiting dilution and integration events selected with G418. Multiple clones
characterized by different HA-FZD2 expressions were expanded. For the
dowunregulation of Ddit3 the following siRNA were used. Cells were transfected with
Lipofectamine™ RNAiMAX Transfection Reagent.
441

442 **Proliferation assay**
443 100k ESCs were plated in a well of a MW12. 48h later, cells were collected and
444 counted. Proliferation rates were calculated as relative factor of cells collected over
445 cells seeded.
446

LC3 staining and size characterization
100k cells were plated on cover-slips, pre-treated with Matrigel (Corning, # 354277),
in N2B27 media. N2B27 composition: 50 mL Neurobasal media (Gibco, # 2348949),
50 mL Advanced DMEM F12 (Gibco, # 12634010), 4% BSA (Gibco, # 15260), 1% L-
Glutamine (Gibco, # 25030024), 1% B27 supplement (Gibco, # 2336837), 0,5% N2
supplement (Gibco, # 2328253), and 0,1 mM ß-mercapotethanol (Sigma,
#BCCB9882). The next day, samples were fixed in 4% PFA for 15 min, washed 3 times
with PBS and then treated with ice-cold MeOH for 5 min on RT. The staining was then
performed as described below. Quantification was performed with the Fiji plugin
TrackMate [47, 48] to automatically detect puncti with a diameter of 1, 2, and 3 microns
at a threshold of 600, 200 and 150, respectively. Nuclei were detected at an average
size of 15 microns with a threshold of 10. Unique particles were annotated for 5
different frames for each condition and shown as particles per cell. The experiment
was repeated 3 times.
461

LD staining and RF microscopy
Samples were plated as for the LC3 staining. For samples treated with oleic acid,
200 μM (Sigma-Aldrich, # O1008-1G) were added at plating, over-night. The next day,
samples were incubated in 5 μM BODIPY 493/503 (Invitrogen, #D3922) in PBS and
subsequently washed 3 times with PBS and fixed in 4 % PFA for 15 min. Samples
were then washed, stained with Dapi for 15 min at RT and mounted with Mowiol.
Quantification was performed using the Fiji plugin TrackMate [47, 48] detecting

droplets of 1, 2, and 3 microns in diameter at a threshold of 900. Nuclei were detected

at 10 micron and a threshold of 1.

For RF microscopy, 250k cells were plated per well of a 6 well glass bottom plate

(Cellvis, # P06-1.5H-N) in N2B27 media. After 36 h samples were treated with 200 µM

oleic acid o/n and then with BODIPY for 3 h. Samples were washed 3 times with PBS

and maintained in N2B27 for RF microscopy. The imaging was performed on

Tomocube HT-X1 instrument at 37C and 5 % CO2. At least 5-8 images were acquired

for each condition for each well. The z-stacks were acquired with 1.06 µm slices. Max

Intensity projections (MIP) are generated on the Tomostudio 3.3.9. The fluorescence

max intensity projections were resized to Refractive index tomograms for comparison.

The ROIs from each image were isolated for further analysis. Comdet plugin was used

for identifying high refractive index regions. The approximate particle size was 2 pixels

and Intensity threshold values were chosen individually for each image to avoid any

false positives or negatives.

Immunofluorescence

Samples were washed 3 times with PBS, fixed in 4% PFA for 10 min and again washed

3 times with PBS. Samples stained for LC3 were then treated for 5 min with 100%

MeOH at RT and washed again 3 times with PBS. Samples were then blocked and

permeabilized with 10% donkey serum (Jackson Immunoresearch, #017-000-121)

and 0,1% Triton X (Thermo Fischer, # HFH10) in PBS for 20 min and incubated with

primary antibodies at 4 °C over-night. Oct4 (1:100, Satna Cruz, # sc-5279), Nanog

(1:100, Abcam, # ab80892), Sox2 (1:50, R&D Systems #AF2018), Lc3 (1:200, Cell

Signalling, # 2775S), Pdi (1:200, Cell Signalling, # 3501S), Rcas1 (1:200, Cell

Signalling, # 12290S), Rab7 (1:200, Cell Signaling, # 9367S) Sox1 (1:50, R&D

Systems, # AF3369), Sox17 (1:200 R&D Systems, # AF1924-SP), Gata4 (1:100 R&D

Systems, # AF1700-SP), Gata6 (1:100 R&D Systems, # AF1700-SP), Dab2 (1:100

BD, # 610464). The next day, samples were washed 3 times with 0.1% Triton X in PBS

and then incubated with secondary antibodies and Dapi.

Western Blot Analysis

Cells were lysed in membrane buffer (150 mM NaCl, 50 mM Tris-HCl pH 7,5, 1 mM 2-

mercaptoethanol (Sigma, #M3148) and 1% CHAPS (Sigma, #C3023) in water).

Lysates were run on SDS-Page and transferred to nitrocellulose membranes. Bands

were detected with the Amersham ECL Prime Western Blotting Detection Reagents (#RPN2232) and captured on Fujifilm Super RX-N films (#47410). Antibodies for Vmp1 (Cell Signalling, #12929S) ,DDIT3 (Cell Signalling, #2895T), β-CATENIN (Santa Cruz, sc-7963), β-ACTIN (BioLegend, W16197A), E-CADHERIN (BioLegend, 67A4), H3 (Abcam, #1791), HA-epitope (Roche, 11867423001), HSP90 (Santa Cruz, #sc-101494) were used. The Quantification of western blot bands was performed with ImageJ.

RNA extraction and RT qPCR

RNA was extracted with the RNeasy Kit (Quiagen, #74104) following the manufacturer's instructions, including the on-column DNA digestion. EB and SCO samples were pre-treated with the QIAshredder (Quiagen, #79656). cDNAs were prepared using the Bio-Rad iScript Reverse Transcription Supermix for RT-qPCR (#1708841). RT qPCR samples were prepared using the KAPPA SYBR R fast (Roche, KK4611). Primer Sequences:

Tmem41b Exon2 FW: TGCAGAAGCTGGATCAGCAA, RV: GAGCCTTGGCGTCATCCATA

Tmem41b Exon7 FW: GCAGGTAGAGCGTCACAGAG, RV: TGCAGAGTTGTTCCTGCCTT

Sdha FW: CGTTATGTGAGGGGTGTGCCTT, RV: CCTTACCAAACCTTGTGTCTGGA

Sox1 FW: ACGTAGCCAACGTAGACAC, RV: ATGAGCGTCGTCCCGTGG

Fgf5 FW: GCCTGTCCTTGCTCTTCCTCAT, RV: GGAGAAGCTGCGACTGGTGA

Actc1 FW: TGTAGGACCGTGTCGAAACC, RV: CGAAACCACACACTGTTACCG

Brachyury FW: TCTCTGGTCTGTGAGCAATGGT, RV: TGCGTCAGTGGTGTGTAATGTG

Gata6 FW: GAAGCGCGTGCCTTCATC, RV: GTAGTGGTTGTGGTGTGACAGTTG

Dab2 FW: TTGATGATGTGCCTGATGCT, RV: TTTGCTTGTGTTGTCCCTGA

Sox17 FW: CCCAACACTCCTCCCAAAGTATC, RV: TTCCCTGTCTTGGTTGATTTCT

Ddit3 FW: GGCTCAAGCAGGAAATCGAG, RV: AAGTGAGAGGCTGTTGACAC

Quantification and statistical analysis

Statistical analysis was performed using Python and GraphPad Prism. Data are
presented as mean-centered and the standard deviation. All experiments were
repeated with at least 3 independent biological replicates, unless stated otherwise.
The statistical test used to evaluate significance is the Welch's t test. Statistical
significance in the figures is denoted as follows: ns: $p > 0.05$, *: $p < 0.05$, **: $p < 0.01$,
***: $p < 0.001$.
543

RNA seq
RNA samples were submitted to the Functional Genomics Center in Zurich (FGCZ) for
RNA sequencing. The quality of the RNA was determined with a Fragment Analyzer
standard sensitivity RNA measurement (SS RNA kit (15 nt), Agilent, Waldbronn,
Germany). The measured concentrations (> 50 ng/µl) and RIN (>9.9) values qualified
for a Poly-A enrichment strategy in order to generate the sequencing libraries applying
the TruSeq mRNA Stranded Library Prep Kit (Illumina, Inc, California, USA). After Poly-
A selection using Oligo-dT beads the mRNA was reverse-transcribed into cDNA. The
cDNA was fragmented, end-repaired and polyadenylated before ligation of TruSeq UD
Indices (IDT, Coralville, Iowa, USA). The quality and quantity of the amplified
sequencing libraries were validated using a Fragment Analyzer SS NGS Fragment Kit
(1–6000 bp) (Agilent, Waldbronn, Germany). The equimolar pool of samples was
spiked into a NovaSeq6000 run targeting 20M reads per sample on a S1 FlowCell
(Novaseq S1 Reagent Kit, 100 cycles, Illumina, Inc, California, USA). Reads from
RNA-seq were first preprocessed by Trimmomatic (v0.40) [49] to remove low-quality
bases and adapters. Then, reads were aligned to mouse genome GRCm38 (release
81 version from Ensembl) using HISAT2 (v2.2.1) [50]. HTSeq Count (v2.0.2) [51] was
used to count reads for each gene (Ensembl GRCm38.81), ignoring reads on
overlapped regions. Differentially expressed genes were defined by DESeq2 (v1.40.2)
[52]. MA plot was generated using custom-made scripts. Genes significantly enriched
show a fold change > 1 and FDR (adjusted p-value) < 0.01. The read counts were
transformed in RPKM for the heatmap representations of specific genes and
pathways. The genes corresponding to germ-layer differentiation markers (including
those for embryonic and extra-embryonic endoderm), biological processes, and
signaling pathways were obtained through literature mining and the KEGG database
(Table S1). The processed data are available as excel files (Table S2 and Table S3)

and raw data have been deposited on NCBI SRA database under the accession:

PRJNA1031567

Surface Proteome Profiling

Sample preparation and biotin labelling was performed following the instructions from

{Wojdyla, 2020 #140}{Weekes, 2012 #141}. In brief, 10M cells were collected per

sample using Accutase (Biolegend, #423201). Cell pellets were washed one with PBS

and then incubated, resuspended in 1 mL biotinylation buffer (97 µM aminooxy-biotin,

Biotium #90113, 1 mM sodium periodate, Thermo-scientific #20504, 0,093% aniline,

Sigma-Aldrich, #242284) for 30 min at 4 °C. Next, cells were pelleted, the reaction

quenched with 1 mM glycerol (30min, 4 °C) and washed once with PBS and lysed in

1 mL of lysis buffer (see western blot) by sonication (Dr. Hielscher sonicator, 30 sec at

80% amplitude and 80% cycle time). Lysates were pelleted, the supernatant

transferred into a fresh tube and incubated with 30 µL (dry resin) of streptavidin

agarose resin (Thermo-Scientific, #20347) for 1.5h at 4 °C. beads collected,

transferred onto a column (BioRad, #7326008) and washed with 8 mL of lysis buffer

and 4 mL of 0,5% SDS in 50 mM TEAB. Beads were resuspended in 200 µL 50 mM

Ambic with 10 mM TCEP (30 min, 37 °C), then 20 mM iodoacetamide was added (30

min. 37 °C). Beads were further washed with 4 mL of SDS buffer (see above), 8 mL

Urea buffer (6M Urea in 50 mM TEAB, pH=8,5), and with 2 mL 50 mM TEAB. Beads

were taken up in 200 µL TEAB buffer and digested with 3 µL Trypsin (Promega) over

night, 37 C. Tryptic digestion was quenched by the addition of acetonitrile (ACN).

Tandem mass tag (TMT) isobaric reagents (TMTpro 6plex Thermo Fisher Scientific)

were dissolved in anhydrous ACN to a final concentration of 20 mg/ml, of which a

unique TMT label was added (100µg). Peptides were incubated at room temperature

for one hour, TMT labeling reactions were halted by 0.3% hydroxylamine and dried.

Peptides were fractionated by high pH fraction in reverse phase (microspin column,

Nest group) following the procedure based on the high pH reversed-phase peptide

fractionation kit (Pierce).

LC-MS/MS was performed on an Orbitrap Exploris 480 mass spectrometer (Thermo

Fisher) coupled to an Vanquish Neo liquid chromatography system (Thermo Fisher).

Peptides were separated using a reverse phase column (75 µm ID x 400 mm New

Objective, in-house packed with ReproSil Gold 120 C18, 1.9 µm, Dr. Maisch GmbH)

across 180 min linear gradient from 7 to 35% (buffer A: 0.1% [v/v] formic acid; buffer
B: 0.1% [v/v] formic acid, 80% [v/v] acetonitrile). Samples were acquired in DDA mode
(Data Dependent Acquision) with MS1 scan (scan range = 350-1500, R=120K, max
injection time auto and AGC target = 100), followed by dependent MS2 scans (first
mass (m/z) 110, R = 45K, max injection time auto and AGC target = 200).Cycle time
was set to 2 sec. Peptides with charge between 2-5 were isolated (m/z = 0.7) and
fragmented (NCE 36%).

Acquired spectra were searched using the MaxQuant software version 2.1.0.0 against
mus musculus proteome reference dataset (http://www.uniprot.org/, downloaded on
03.04.2020) extended with reverse decoy sequences. The search parameters were
set to tryptic peptides, maximum two missed cleavage, carbamidomethyl as static
peptide modification, oxidation (M) and deamidation (N-terminal) and TMT6 as static
peptide modification. The MS and MS/MS mass tolerance was set to 10 ppm. False
discovery rate of < 1% was used at PSM and protein level. Abundance of 1417 proteins
was determined from the intensity of top two peptides. Intensities values were median
normalized and missing values were imputed using random sampling from a normal
distribution generated from 1% less intense values. Statistical analysis was performed
using unpaired two sided t-test and pvalues were corrected using Benjamin-Hochberg
correction. The entire dataset, including raw data, generated tables and scripts used
for the data analysis are available in the PRIDE repository: PXD053039. Matrices with
protein intensities are reported in Table S4.

Competing interest
The authors declare no competing interests.

Acknowledgments

We acknowledge the Functional Genomics Center Zurich (FGCZ) for technical support
and Sungsik Lee from ScopeM facility for help on the RF microscopy. We thank G.J.
Pereira, C. Fimiani, and U. Sutter for their help with reagents. We thank C. Ciaudo, A.
Grison, J. Corn, and K. Basler for their helpful discussions. GDM was supported by
the ETH Zurich Postdoctoral Fellowship Program as well as the Marie Curie Actions

for People COFUND Program. This work was supported by grants from the Swiss
National Science Foundation (SNF grants 31003A_152814/1 and 31003A_175643/1).
639

Author Contributions

Conceptualization: MH, AW and GDM; Data curation: MH, FU, VG and GDM; Formal
analysis: MH, TS, HH, FU, VG, and GDM; Funding acquisition: KW, AW and GDM;
Investigation: MH, AW and GDM; Methodology: MH, TS, HH, FU, VG, WJ, and GDM;
Project administration: AW, GDM; Resources: MH, KW, AW and GDM; Supervision:
KW, AW and GDM; Validation: MH, GDM; Visualization: MH, FU, VG and GDM; Writing
– original draft: MH, AW and GDM; Writing – review & editing: MH, TS, HH, FU, VG,
WJ, KW, AW and GDM.
648
649
650
651

REFERENCES

1. Trimarco, J.D., et al., *TMEM41B is a host factor required for the replication of diverse*
coronaviruses including SARS-CoV-2. PLoS Pathog, 2021. **17**(5): p. e1009599.
2. Sun, L., et al., *Genome-scale CRISPR screen identifies TMEM41B as a multi-function*
host factor required for coronavirus replication. PLoS Pathog, 2021. **17**(12): p.
e1010113.
3. Hoffmann, H.H., et al., *TMEM41B Is a Pan-flavivirus Host Factor.* Cell, 2021. **184**(1): p.
133-148 e20.
4. Kratzel, A., et al., *A genome-wide CRISPR screen identifies interactors of the autophagy*
pathway as conserved coronavirus targets. PLoS Biol, 2021. **19**(12): p. e3001490.
5. Baggen, J., et al., *Cellular host factors for SARS-CoV-2 infection.* Nat Microbiol, 2021.
6(10): p. 1219-1232.
6. Lin, W., et al., *VMP1, a novel prognostic biomarker, contributes to glioma development*
by regulating autophagy. J Neuroinflammation, 2021. **18**(1): p. 165.
7. Ropolo, A., et al., *A Novel E2F1-EP300-VMP1 Pathway Mediates Gemcitabine-Induced*
Autophagy in Pancreatic Cancer Cells Carrying Oncogenic KRAS. Front Endocrinol
(Lausanne), 2020. **11**: p. 411.
8. Folkerts, H., et al., *Elevated VMP1 expression in acute myeloid leukemia amplifies*
autophagy and is protective against venetoclax-induced apoptosis. Cell Death Dis,
2019. **10**(6): p. 421.
9. Rodriguez, M.E., et al., *A novel HIF-1alpha/VMP1-autophagic pathway induces*
resistance to photodynamic therapy in colon cancer cells. Photochem Photobiol Sci,
2017. **16**(11): p. 1631-1642.
10. Gilabert, M., et al., *Novel role of VMP1 as modifier of the pancreatic tumor cell*
response to chemotherapeutic drugs. J Cell Physiol, 2013. **228**(9): p. 1834-43.

11. Guo, L., et al., *Novel roles of Vmp1: inhibition metastasis and proliferation of*
hepatocellular carcinoma. Cancer Sci, 2012. **103**(12): p. 2110-9.
12. Zack, S.R., et al., *Vacuole Membrane Protein 1 (VMP1) Restricts NLRP3 Inflammasome*
Activation by Modulating SERCA Activity and Autophagy. Res Sq, 2023.
13. Yao, Y., et al., *WHSC1L1-mediated epigenetic downregulation of VMP1 participates in*
herpes simplex virus 1 infection-induced mitophagy impairment and
neuroinflammation. Mol Immunol, 2023. **163**: p. 63-74.
14. Li, X.Q., et al., *TMEM41B Is an Interferon-Stimulated Gene That Promotes Pseudorabies*
Virus Replication. J Virol, 2023. **97**(6): p. e0041223.
15. Loncle, C., et al., *The pancreatitis-associated protein VMP1, a key regulator of inducible*
autophagy, promotes Kras(G12D)-mediated pancreatic cancer initiation. Cell Death
Dis, 2016. **7**(7): p. e2295.
16. Jiang, X., et al., *Lack of VMP1 impairs hepatic lipoprotein secretion and promotes non-*
alcoholic steatohepatitis. J Hepatol, 2022. **77**(3): p. 619-631.
17. Li, Y.E., et al., *TMEM41B and VMP1 are scramblases and regulate the distribution of*
cholesterol and phosphatidylserine. J Cell Biol, 2021. **220**(6).
18. Ghanbarpour, A., et al., *A model for a partnership of lipid transfer proteins and*
scramblases in membrane expansion and organelle biogenesis. Proc Natl Acad Sci U S
A, 2021. **118**(16).
19. Molejon, M.I., et al., *The VMP1-Beclin 1 interaction regulates autophagy induction.* Sci
Rep, 2013. **3**: p. 1055.
20. Moretti, F., et al., *TMEM41B is a novel regulator of autophagy and lipid mobilization.*
EMBO Rep, 2018. **19**(9).
21. Morishita, H., et al., *A critical role of VMP1 in lipoprotein secretion.* Elife, 2019. **8**.
22. Hama, Y., H. Morishita, and N. Mizushima, *Regulation of ER-derived membrane*
dynamics by the DedA domain-containing proteins VMP1 and TMEM41B. EMBO Rep,
2022. **23**(2): p. e53894.
23. Tornero-Ecija, A., et al., *A Dictyostelium model for BPAN disease reveals a functional*
relationship between the WDR45/WIPI4 homolog Wdr45l and Vmp1 in the regulation
of autophagy-associated PtdIns3P and ER stress. Autophagy, 2022. **18**(3): p. 661-677.
24. Bard, F., et al., *Functional genomics reveals genes involved in protein secretion and*
Golgi organization. Nature, 2006. **439**(7076): p. 604-7.
25. Huang, D., et al., *TMEM41B acts as an ER scramblase required for lipoprotein*
biogenesis and lipid homeostasis. Cell Metab, 2021. **33**(8): p. 1655-1670 e8.
26. Van Alstyne, M., et al., *Stasimon/Tmem41b localizes to mitochondria-associated ER*
membranes and is essential for mouse embryonic development. Biochem Biophys Res
Commun, 2018. **506**(3): p. 463-470.
27. Farese, R.V., Jr., et al., *Knockout of the mouse apolipoprotein B gene results in*
embryonic lethality in homozygotes and protection against diet-induced
hypercholesterolemia in heterozygotes. Proc Natl Acad Sci U S A, 1995. **92**(5): p. 1774-
8.
28. Raabe, M., et al., *Knockout of the abetalipoproteinemia gene in mice: reduced*
lipoprotein secretion in heterozygotes and embryonic lethality in homozygotes. Proc
Natl Acad Sci U S A, 1998. **95**(15): p. 8686-91.
29. Arnold, S.J. and E.J. Robertson, *Making a commitment: cell lineage allocation and axis*
patterning in the early mouse embryo. Nat Rev Mol Cell Biol, 2009. **10**(2): p. 91-103.

30. O'Rourke, E.J., et al., *C. elegans major fats are stored in vesicles distinct from lysosome-*
related organelles. Cell Metab, 2009. **10**(5): p. 430-5.

31. Zeevaert, K., et al., *Cell Mechanics in Embryoid Bodies.* Cells, 2020. **9**(10).

32. Kruithof-de Julio, M., et al., *Regulation of extra-embryonic endoderm stem cell*
differentiation by Nodal and Cripto signaling. Development, 2011. **138**(18): p. 3885-
95.

33. Niakan, K.K., et al., *Derivation of extraembryonic endoderm stem (XEN) cells from*
mouse embryos and embryonic stem cells. Nat Protoc, 2013. **8**(6): p. 1028-41.

34. Ngondo, R.P., et al., *Argonaute 2 Is Required for Extra-embryonic Endoderm*
Differentiation of Mouse Embryonic Stem Cells. Stem Cell Reports, 2018. **10**(2): p. 461-
476.

35. Tetsu, O. and F. McCormick, *Beta-catenin regulates expression of cyclin D1 in colon*
carcinoma cells. Nature, 1999. **398**(6726): p. 422-6.

36. Du, Y.C., et al., *Induction and down-regulation of Sox17 and its possible roles during*
the course of gastrointestinal tumorigenesis. Gastroenterology, 2009. **137**(4): p. 1346-
57.

37. Zhang, Z., et al., *Secreted frizzled related protein 2 protects cells from apoptosis by*
blocking the effect of canonical Wnt3a. J Mol Cell Cardiol, 2009. **46**(3): p. 370-7.

38. Aguilera, O., et al., *The Wnt antagonist DICKKOPF-1 gene is induced by 1alpha,25-*
dihydroxyvitamin D3 associated to the differentiation of human colon cancer cells.
Carcinogenesis, 2007. **28**(9): p. 1877-84.

39. Athanasouli, P., et al., *The Wnt/TCF7L1 transcriptional repressor axis drives primitive*
endoderm formation by antagonizing naive and formative pluripotency. Nat Commun,
2023. **14**(1): p. 1210.

40. Chiu, C.S., et al., *Exploiting Honokiol-induced ER stress CHOP activation inhibits the*
growth and metastasis of melanoma by suppressing the MITF and beta-catenin
pathways. Cancer Lett, 2019. **442**: p. 113-125.

41. Wang, S., et al., *Loss of acinar cell VMP1 triggers spontaneous pancreatitis in mice.*
Autophagy, 2022. **18**(7): p. 1572-1582.

42. Wang, C., et al., *An autoregulatory feedback loop of miR-21/VMP1 is responsible for*
the abnormal expression of miR-21 in colorectal cancer cells. Cell Death Dis, 2020.
11(12): p. 1067.

43. Lo Re, A.E., et al., *Novel AKT1-GLI3-VMP1 pathway mediates KRAS oncogene-induced*
autophagy in cancer cells. J Biol Chem, 2012. **287**(30): p. 25325-34.

44. Futterer, A., et al., *Ablation of Dido3 compromises lineage commitment of stem cells in*
vitro and during early embryonic development. Cell Death Differ, 2012. **19**(1): p. 132-
43.

45. Yu, H., et al., *Frizzled 1 and frizzled 2 genes function in palate, ventricular septum and*
neural tube closure: general implications for tissue fusion processes. Development,
2010. **137**(21): p. 3707-17.

46. Di Minin, G., et al., *TMED2 binding restricts SMO to the ER and Golgi compartments.*
PLoS Biol, 2022. **20**(3): p. e3001596.

47. Ershov, D., et al., *TrackMate 7: integrating state-of-the-art segmentation algorithms*
into tracking pipelines. Nat Methods, 2022. **19**(7): p. 829-832.

48. Tinevez, J.Y., et al., *TrackMate: An open and extensible platform for single-particle*
tracking. Methods, 2017. **115**: p. 80-90.

49.   Bolger, A.M., M. Lohse, and B. Usadel, *Trimmomatic: a flexible trimmer for Illumina*
770   *sequence data.* Bioinformatics, 2014. **30**(15): p. 2114-20.
771   50.   Kim, D., et al., *Graph-based genome alignment and genotyping with HISAT2 and HISAT-*
772   *genotype.* Nat Biotechnol, 2019. **37**(8): p. 907-915.
773   51.   Putri, G.H., et al., *Analysing high-throughput sequencing data in Python with HTSeq*
774   *2.0.* Bioinformatics, 2022. **38**(10): p. 2943-2945.
775   52.   Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and*
776   *dispersion for RNA-seq data with DESeq2.* Genome Biol, 2014. **15**(12): p. 550.
777
778

FIGURE LEGENDS

Figure 1. Vmp1 and Tmem41b double KO ESCs display increased autophagosome and lipid droplet accumulation

(A) Schematic representation of the CRISPR/Cas9-mediated KO strategy for Vmp1 and Tmem41b genes. For depleting Vmp1 (top) a single gRNA inducing an indel mutation has been used. In the case of the Tmem41b gene (bottom), two gRNAs were used to delete exons 2 and 3. The sequences coding for transmembrane domains (TM) are indicated in blue, the conserved DedA domain is highlighted in yellow, red lines indicate cut sites of gRNAs. **(B)** Colony morphology of VMP1^{KO} and Tmem41^{KO} ESCs remains unchanged compared to WT ESCs. Scale bar: 100 µm. **(C)** VMP1 and Tmem41b depletion induce accumulation of autophagosomes in ESCs. The graph shows the quantification of LC3 puncta per cell of a representative experiment (see Fig.S2A). Data were collected from more than 200 cells. For each KO condition, two independent clones have been analyzed. Dots correspond to the frames analyzed per sample. Data are represented as mean ± SD. The P-value is relative to the WT sample. **(D)** Vmp1 and Tmem41b depletion leads to the accumulation of lipid droplets (LD) in ESCs. Graph quantifies LD per cell in WT and KO ESCs (see Fig.S2D). ± oleic acid treatment. At least 500 cells were analyzed for every condition; dots represent independent biological replicates. Data are represented as mean ± SD. **(E-G)** Transcriptional analysis of Double KO ESCs did not identify many deregulated genes. **(E)** Shown are differentially regulated genes in Double KO ESCS in blue as MA plot. Genes highlighted in red are pluripotency markers. **(F)** Differentially regulated genes do not cluster in essential processes. The graph shows differentially expressed genes of single and double KO ESCs by cellular processes. Dot size correlates to the number of deregulated genes within a cellular process. **(G-H)** Double KO ESCs accumulate LC3+ autophagosomes of increased size. **(G)** Shown are representative IF images of WT and double KO ESCs stained for LC3. Nuclei are stained with Dapi. Scale bar: 20 µm. **(H)** Plotted is the quantification of LC3 puncta of 3 µm in diameter. For each condition, two independent clones have been analyzed. Dots correspond to the frames analyzed per sample. Data were collected from more than 200 cells. Data are represented as mean ± SD. The P-value is relative to the Double KO sample with the highest SD. **(I-J)** Double KO ESCs accumulate LDs without oleic acid treatment. **(I)**

Live cell staining of WT, Vmp1, Tmem41b, and double KO ESCs with BODIPY. Scale

bar: 20 µm **(J)** The number of LDs per cell is plotted for all KO clones. Data are

represented as mean ± SD. **(K)** Double KO ESCs present an increased amount of LD

visualized through refractive index microscopy (top) and co-stained with BODIPY.

Figure 2. Double KO ESCs specify all germ layers, but not the primitive

endoderm

(A) Schematic representation of an Embryonic Body (EB) which consists of 3 different

lineages. The ectoderm expresses Fgf5 and Sox1 (grey), the mesoderm Actc1 and

Brachyury (blue), and the endoderm by Gata6. The surrounding region of the EB

corresponds to the primitive endoderm (green) and is marked by Gata6 and Dab2

expression. **(B)** ESCs differentiated to EBs express lineage markers for ectoderm

(Fgf5), mesoderm (Brachyury), endoderm (Gata6), and primitive endoderm (Dab2).

Graphs show mRNA levels of WT EBs normalized to the Sdha gene. Dots represent

biological independent experiments. Data are represented as mean ± SD. **(C)** Double

KO ESCs specifically show defects in expressing the primitive endoderm marker,

Dab2. qPCR analysis for Sox1, Actc1, and Dab2 mRNA expression in mutant EBs

relative to WT EBs. Two independent clones were analyzed per cell line. Each dot

represents an independent experiment. Data are represented as mean ± SD**.** The line

at 0,7 indicates an arbitrary threshold after which successful expression of the marker

is considered. Statistical significance (Welch's t test) is denoted as follows: ns: $p > 0.05$,

*: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$. **(D-F)** Double mutant EBs show decreased

expression of endoderm markers. Immunofluorescence images of EB sections stained

for the endoderm markers GATA4 and SOX17 **(D)**, GATA6 **(E)**, DAB2 **(F)**. WT in the

top panel, two independent clones for Double KO condition in the center and bottom

panels. Nuclei are stained with Dapi. Scale bar: 50 µm.

Figure 3. XEN differentiation in Double KO cells is delayed(A-B)

Double KO ESCs show a delay in XEN differentiation. **(A)** Brightfield images of XEN

cells at day 6 of differentiation in WT and Double KO cells. XEN islands are circled in

white in the Double KO condition. Scale bar: 100 µm. **(B)** Immunofluorescence images

of day 6 XEN cells stained for SOX17 and GATA4. Nuclei are stained with Dapi, Scale

bar: 150 µm. **(C-F)** Transcriptional analysis of XEN cells. **(C)** Differentially expressed

genes between Double KO and WT cells at day 2 during XEN specification. XEN

markers are labeled in red; differentially expressed genes in blue. **(D-E)** GO-Term

analysis of differentially expressed genes between Double KO and WT conditions in

ESCs and XEN cells in the context of developmental processes **(D)** and signaling

pathways **(E)**. Dot size corresponds to the number of differentially regulated genes.

(F) WNT pathway-related genes differentially expressed in Double KO cells. Heatmap

shows min to max expression of the average of three biological replicates for WT and

Double KO XEN cells. **(G-H)** VMP1 and Tmem41b depletion promotes the increase of

DDIT3 protein levels. **(G)** Western Blot analysis of DDIT3 expression in WT and

Double KO cells differentiated to XEN cells for 2 days. HSP90 is shown as loading

control. **(H)** Quantification of increased DDIT3 protein levels in three biological

replicates. Expression relative to WT XEN cell and normalized to respective HSP90

expression. Data are represented as mean ± SD. **(I)** DDIT3 silencing does not restore

XEN specification in Double KO cells. qPCR analysis at day 3 of differentiation for

Ddit3, Gata4, and Dab2 mRNA expression in mutant cells relative to WT ESCs

transfected without and with Ddit3 siRNA. Data are represented as mean ± SD (n=3).

Figure 4. Vmp1 and Tmem41b mutant ESCs show reduced plasma membrane

abundance of the WNT receptor FZD2

(A) Differences in the abundance of plasma membrane proteins in WT and VMP1[KO]

cells. Volcano plot displaying differentially expressed proteins at the plasma

membrane in WT and VMP1[KO] ESCs. Proteins with the most significant divergence

are highlighted in red. Gray lines denote the selection criteria: an absolute log2 fold-

change > 1 and a false discovery rate < 0.05. The position of FZD2 is annotated. **(B)**

GO term analysis of proteins differentially abundant at the plasma membrane of WT

and Vmp1[KO] cells. Shown are GO terms related to protein localization and molecular

function. **(C)** Abundance of Frizzled receptors identified by MS analysis in WT and

mutant cells. The average and standard error is reported for three independent

biological replicates **(D)** Western blot analysis of HA-FZD2 abundance at the PM in

WT, VMP1[KO], and Double KO ESCs. E-CADHERIN and ACTIN serve as controls for

PM purification and loading, respectively. **(E)** Quantification of HA-FZD2 protein levels

at the PM across three biological replicates relative to WT cells and normalized to the

total HA-FZD2 (Input). Data are represented as mean ± SD. **(F)** Western blot analysis

of HA-FZD2 abundance at the PM in WT and Double KO cells differentiated into XEN

cells for 2 days.

Figure 5. Activation of WNT signaling rescues the delay of Double KO cells in

XEN differentiation

(A) Cytoplasmic β-CATENIN levels are reduced in Double KO cells. Western blot

analysis was performed on whole-cell lysates (1/50) and purified cytoplasmic fractions

(1/50) from WT and Double KO cells. E-CADHERIN, Histone H3, and ACTIN were

used as controls for purification and loading. Whole-cell and purified cytoplasmic

fractions are shown with the same exposure conditions. **(B)** Chiron-dependent WNT

activation rescues the delay of Double KO XEN differentiation. IF staining for SOX17

and GATA4; nuclei are stained with Dapi. Scale Bar: 150 µm. **(C-D)** FZD2

overexpression rescues the differentiation delay in Double KO XEN cells. **(C)**

Immunofluorescence staining for SOX17 and GATA4 in WT and Double KO cells.

Nuclei are counterstained with DAPI. Scale bar: 100 µm. **(D)** Quantification of GATA4

and SOX17 positive cells during XEN differentiation, expressed as a percentage of

total cells (DAPI-stained).

SOX17 GATA4 DAPI