The scramblases VMP1 and TMEM41b are required for primitive endoderm specification by targeting WNT signaling

Markus Holzner1, Tea Sonicki1, Hugo Hunn1, Vamshidhar R. Gade2, Karsten Weis2, Anton Wutz1#, and Giulio Di Minin1#

1 Institute of Molecular Health Sciences, Department of Biology, ETH Zurich, Zurich, Switzerland

2 Institute of Biochemistry, Department of Biology, ETH Zurich, Zurich, Switzerland

corresponding authors: awutz@ethaz.ch (AW); giulio.diminin@biol.ethz.ch (GDM)

ABSTRACT

The VMP1 and TMEM41b ER resident proteins share a DedA domain, which defines their lipid scramblase activity. Loss of either gene results in embryonic lethality in mice, and defects in autophagy and lipid droplet metabolism. To understand their role in pluripotency and specification, we generated Vmp1 and Tmem41b mutant mouse Embryonic Stem Cells (ESCs). We observe that ESCs carrying mutations in Vmp1 and Tmem41b are viable, proliferate normally, and conserve a largely intact pluripotency-associated transcriptional profile. Despite clear defects in the accumulation of LC3-positive autophagosomes and lipid droplets, ESCs carrying combined mutations in Vmp1 and Tmem41b can differentiate into a wide range of embryonic cell types. However, we find that simultaneous loss of Vmp1 and Tmem41b impairs the specification of primitive endoderm-like cells. We observe a delayed differentiation of mutant ESCs into extra-embryonic stem (XEN) cells. Mechanistically, we discover that mutant cells upregulate DDIT3, a WNT inhibitor. Chemical stimulation of the WNT cascade can rescue the differentiation delay. Our findings reveal a redundant function of the lipid scramblases VMP1 and TMEM41b and identify a specific role in cell signaling during extraembryonic endoderm development.

INTRODUCTION

The role of vesicle trafficking and membrane lipid composition becomes increasingly important to understand cell signaling. Recently, the role of lipid scramblases has been implicated in viral infection, such as in the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) (Baggen et al., 2021; Hoffmann et al., 2021; Kratzel et al., 2021; Sun et al., 2021; Trimarco et al., 2021), cancer development (Folkerts et al., 2019; Gilabert et al., 2013; Guo et al., 2012; Lin et al., 2021; Rodriguez et al., 2017; Ropolo et al., 2020), and inflammatory diseases (Jiang et al., 2022; Li et al., 2023b; Loncle et al., 2016; Yao et al., 2023; Zack et al., 2023). The vacuole membrane protein 1 (VMP1) and transmembrane protein 41B (TMEM41B) are two lipid scramblases that share a highly conserved DedA domain (Ghanbarpour et al., 2021; Li et al., 2021). They localize at the ER membrane and are distinguished from other ER-associated flippases or scramblases by the property to facilitate bi-directional lipid diffusion in an ATP- and Ca2+-independent manner. Vmp1 and Tmem41b activities are necessary for autophagosome formation (Molejon et al., 2013; Moretti et al., 2018), lipid droplet metabolism (Li et al., 2021; Morishita et al., 2019), and lipoprotein formation (Hama et al., 2022). Additional reports proposed a more global role in vesicle transport (Bard et al., 2006) and ER protein homeostasis (Tornero-Ecija et al., 2022). Despite their multifaceted functions, the absence of these factors does not appear to impact cell survival significantly. Compensatory mechanisms shared by VMP1 and TMEM41b or between other ER scramblases might be involved in maintaining cellular homeostasis. A systematic comparison between the activities of Vmp1 and Tmem41b has not yet been performed. VMP1 depletion leads to pronounced defects in autophagosome assembly (Molejon et al., 2013; Moretti et al., 2018), while TMEM41B appears to play a pivotal role in maintaining lipid droplet homeostasis and serves as a crucial factor in RNA virus replication (Hoffmann et al., 2021; Huang et al., 2021).

VMP1 and TMEM41B are both essential in embryonic development. Vmp1 (Morishita et al., 2019) and Tmem41b (Van Alstyne et al., 2018) deficient mice die around E8.5 because of severe developmental delay and impaired implantation. Prior studies formulated a hypothesis where Vmp1 restricts the transfer of vital nutrients to the embryo by blocking the synthesis and release of lipoproteins from the visceral endoderm. However, the observation that mice with a deficiency in crucial genes for

lipoprotein biogenesis, such as ApoB and Mttp, experience mortality at later stages
(specifically, E10.5) (Farese et al., 1995; Raabe et al., 1998), implies a distinct and
earlier involvement of VMP1 and TMEM41B in embryo development.

During implantation, the mouse embryo is composed by three distinct lineages:
the trophectoderm, responsible for placental development; the primitive endoderm
(PE), contributing to yolk sac formation; and the epiblast (EPI), primed to develop into
the embryo itself (Arnold and Robertson, 2009). The EPI and PE are specified within
the inner cell mass (ICM) at the blastocyst stage. Defects in the formation of these
lineages are challenging to characterize *in vivo* due to their occurrence in a pre-
implantation stage and the small size of the embryo. To investigate the role of VMP1
and TMEM41B in early-embryo linage specification, we studied the ability of mutant
cells to differentiate *in vitro* towards embryonic germ layers. We found that a combined
mutation of Vmp1 and Tmem41b in ESCs results in a specific defect in PE formation
and a delayed conversion to Extra-embryonic endoderm cells (XEN). We find that this
delay can be rescued by activation of WNT signaling.

RESULTS

Vmp1 and Tmem41b knockout ESCs show accumulation of LC3+
autophagosome and lipid droplets

To assess the influence of Vmp1 and Tmem41b depletion on mouse ESCs and in early
embryo development, we generated independent knockout clones by introducing
genomic deletions via the CRISPR/Cas9 system. A mutation of the Vmp1 gene was
achieved by introducing an indel in Exon 2 (Fig.1A). This strategy aims to preserve the
expression of miR-21, which overlaps with the 3' end of Vmp1 coding sequence (Wang
et al., 2020). Edited clones were validated by WB (Fig.S1A). To abrogate Tmem41b
expression, we targeted intron 1 and 3, for removing the second and third exons
(Fig.1A), which encode the first transmembrane domain of the protein, compromising
its correct topology. Additionally, splicing of exon 1 to exon 4 results in a frameshift,
suggesting a strong or null allele. The deletion was confirmed on the genomic DNA
(Fig.S1B) and mRNA levels by RT-qPCR (Fig.S1C-D). Vmp1 and Tmem41b depleted
ESCs show similar morphology (Fig.1B), proliferation (Fig.S1E), and expression of
pluripotency markers (Fig.S1F-G) compared to WT ESCs. We confirmed that the
pluripotent network was unchanged by a transcriptomic analysis performed in Vmp1

and Tmem41b mutant cells. Interestingly, this analysis also show that the transcription profile of cells was marginally affected by the depletion of the two factors (Fig.S1I-J and Table S2).

To verify the effect of the mutations on autophagy, we assessed the accumulation of LC3+ autophagosomes. We found an increase of LC3+ puncta in both mutant conditions compared to WT ESCs (Fig.1C and Fig.S2A-B). Interestingly, the increase in the frequency of LC3+ puncta was similar in Vmp1 (4X) and Tmem41b (3,5X) depleted cells. Next, mutant cells were analyzed for the retention of lipid droplets (LD). To amplify LDs, we treated all samples with oleic acid (OA) and visualized LDs by BODIPY staining. Compared to WT ESCs, we observed increased LDs in both mutant conditions (Fig.1D and Fig.S2D). This result was more evident after OA treatment, where Vmp1 and Tme41b mutations led to an increase of 4 times of LD frequency with respect to WT cells. The results suggest that both Vmp1 and Tmem41b are equally necessary for autophagosome and lipid droplet metabolism in ESCs.

Vmp1 and Tmem41b are not required for ESC survival and self-renewal

To characterize potential compensatory mechanisms between Vmp1 and Tmem41b, we generated cell lines in which both proteins are depleted. For this purpose, we started with Vmp1 KO ESCs and deleted Tmem41b following the strategy detailed above (Double KO) (Fig.S1A-D). As previously shown in the context of the single mutants, Double KO cells show no differences in ESC morphology (Fig.S1H) and proliferation (Fig.S1E). To further characterize mutant ESCs, we analyzed secretory compartments. The morphology of ER, Golgi apparatus, and Late Endosomes (Fig.S3A) is unaffected in all mutant cells. In line with these findings, the transcription profile of Double KO cells showed a limited perturbation in gene expression (Fig.1E). Simultaneous depletion of both Vmp1 and Tmem41b does not result in a greater number of differentially regulated genes compared to Vmp1 KO cells (Fig.1F). Minimal overlap of differentially expressed genes was detected among mutant cell lines (Fig.1F and Fig.S3B), and we did not observe enrichment for genes that are involved in defined cellular processes (Fig.1G, Fig.S3C-D). The pluripotency network, as also shown by IF staining (Fig.S1F-G), was not perturbed by the simultaneous absence of Vmp1 and Tmem41b. We concluded that Vmp1 and Tmem41b are not necessary for maintaining ESC homeostasis.

Vmp1 and Tmem41b have redundant functions

We found that Double KO cell lines show an exacerbated phenotype in autophagosome and LD accumulation in respect to single mutants. In the Double KO, we did not detect an additional increase in the number of autophagosomes per cell (Fig.1H and Fig.S2A-B). However, comparing their size distribution, we found about 14% of LC3 clusters of the Double KO to be of larger size (>3μm), compared to the 8% of autophagosomes across the single mutations and 3% in the WT condition (Fig.1I and Fig.S2C). The simultaneous depletion of Vmp1 and Tmem41b had a greater impact on LD formation. Double KO cells show a high enrichment of LDs in the absence of oleic acid treatment, which was instead necessary to visualize LDs in WT and single mutant cells (Fig.1J and Fig.S2D-E). Considering that the used stain could also dye lipid-containing compartments like autophagosomes (O'Rourke et al., 2009), we confirmed enriched lipid droplets in Double KO cells with refractive index tomography, a labelling-independent technique (Fig.1K and Fig.S2F-G). Together, these results indicate that Vmp1 and Tmem41b share redundant functions, as the depletion of both genes exacerbates autophagosome and lipid droplet accumulation.

Loss of VMP1 and TMEM41b affects ESC potential to differentiate into primitive endoderm

To assess the effect of Vmp1 and Tmem41b on the differentiation potential of ESCs, we used embryoid body (EB) differentiation (Fig.2A). Vmp1, Tmem41b, and double mutant ESCs formed well-defined EBs that resembled those of WT cells. We evaluated variations in the differentiation potential of mutant cells by comparing gene expression by RT-qPCR. For each mutant, the differentiation potential of two independent clones was tested. After 6 days, genes expressed in ectoderm (Sox1 and Fgf5), mesoderm (Actc1 and Brachyury) and endoderm (Gata6 and Dab2) are upregulated during the process of differentiation (Fig.2B and Fig.S4A), while the expression of pluripotent markers (Nanog and Pou5f1) was lost (Fig.2B and Fig.S4A). All cell lines analyzed efficiently differentiated into ectoderm and mesoderm lineages (Fig.2C and Fig.S4B-C). Notably, Double KO EBs showed a decreased expression of the endoderm markers Dab2 (100%) and Gata6 (42% of experiments) (Fig.2C and Fig.S4C). Neither single mutant showed Dab2 downregulation even if a decreased endoderm specification was detected, as shown by Gata6 expression levels in

experiments performed with Vmp1^{KO} (45%) and Tmem41b^{KO} (40%) cells (Fig.2C and

Fig.S4C). Our results indicate that combined mutation of Vmp1 and Tmem41b leads

to abrogated Dab2 expression and predisposes variability in the expression of other

endodermal genes.

We confirmed the ability of Double KO cells to differentiate into ectodermal

lineages by generating SOX1-positive spinal cord organoids (Fig.S4D). Mesodermal

differentiation was inferred by the ability of double mutant EBs to form contractile

cardiomyocytes after plating on gelatin-coated dishes (Video S1-2). We subsequently

performed immunostaining on sections of day 6 EBs to evaluate the protein expression

and distribution of several endoderm markers. Expression of GATA4, GATA6, and

SOX17 was detected but noticeably decreased in Double KO compared to WT EBs

(Fig.2D-E). Loss of GATA4 and GATA6 staining was pronounced at the border of the

EBs, which resembles the primitive endoderm layer (Zeevaert et al., 2020). In line with

these findings, we confirmed the absence of DAB2 in Double KO EBs (Fig.2F). Dab2

is mainly expressed In the primitive endoderm, while the other markers are known to

be expressed in both primitive and definitive endoderm.

The significant reduction in both mRNA and protein levels of Dab2, along with

the weakened expression of endoderm markers at the edges of embryoid bodies,

provides evidence that the absence of Vmp1 and Tmem41b results in a particular

deficiency in the formation of primitive endoderm.

Vmp1 and Tmem41b mutations delay ESC differentiation into Extra Embryonic

Stem Cells (XEN)

To further characterize endodermal differentiation, we examined the impact of

depleting Vmp1 and Tmem41b on extra-embryonic stem (XEN) cell specification. XEN

cells resemble the primitive endoderm (PE) of the blastocyst and allow to recapitulate

in vitro the signaling and transcriptional program activated during PE specification

(Kruithof-de Julio et al., 2011). Double KO and WT cells were differentiated into XEN

cells upon treatment with Activin A and Retinoic Acid (Niakan et al., 2013). After 6 days,

we observed that the majority (>80%) of ESCs differentiated to XEN clusters

presenting stellate and refractile morphology (Niakan et al., 2013) in the WT sample

(Fig.3A). In Double KO cells, the formation of these clusters was strongly impaired.

We confirmed XEN identity and the Double KO defect by analyzing the expression of

XEN markers by RT-qPCR (Fig.S5A) and immunostaining (Fig.3B and Fig.S5B). It is

worth noting that the small number of derived XEN cells from Double KO ESCs can
be expanded for multiple passages, as shown in Fig. S5C. This result suggests that
the absence of Vmp1 and Tmem41b does not completely prevent XEN specification
but causes a delayed entry into XEN differentiation.
206

Lack of WNT signal in Vmp1 and Tmem41b mutant cells impairs XEN specification

To chracterize the effect of Vmp1 and Tmem41b depletion on XEN specification, we
analyzed the transcription profile of Double KO and WT cells during the differentiation
process. We performed RNA sequencing on cells on the second day of differentiation
to identify differences in gene expression. Already at this time point, the expression of
XEN-dependent genes (Ngondo et al., 2018) was strongly down-regulated in Double
KO cells (Fig.3C-D and Fig.5E). At the transcriptional level, depletion of Vmp1 and
Tmem41b did not significantly impact processes involving lipid metabolism,
autophagy, or ER maintenance (Fig.S5D).

During the XEN differentiation process, we observed specific deregulation of
genes related to the WNT pathway in Double KO cells (Fig.3E-F). A close analysis of
WNT pathway-deregulated genes allowed to identify several well-established WNT-
target genes, such as Frzb, Sox17, Wnt3, and Ccdnd1 (Du et al., 2009; Tetsu and
McCormick, 1999; Zhang et al., 2009). Their down-regulation strongly suggests a
decrease in activity of the WNT pathway in mutant cells which is a plausible
explanation for the XEN specification defects (Athanasouli et al., 2023). Interestingly,
we detected Ddit3 among the upregulated genes in Double KO cells. DDIT3 is an
inhibitor of the WNT signaling by repressing the transcriptional activity of the
TCF7/TCF4 complex (Chiu et al., 2019). We confirmed higher level of DDIT3 protein
in Double KO cells (Fig.4A), and quantified a fourfold increase over WT cells (Fig.4B).

To investigate if XEN differentiation of double mutant cells can be rescued by
WNT signaling, we chemically activated WNT signaling. We used a GSK kinase
inhibitor (CHIR) as a WNT agonist. GSK inhibition did not affect XEN specification in
wild-type ESCs. However, WNT activation significantly enhanced the formation of XEN
cells in double mutant ESCs (as shown in Fig. 3D-F and Fig. S5B). Collectively, our
data indicate how Vmp1 and Tmem41b affect WNT signaling, and explain the delay
detected during primitive endoderm specification.
235

DISCUSSION

Vmp1 and Tmem41b have been recently identified in genetic screens designed to study a variety of cellular processes, such as autophagy, lipid metabolism, virus maturation, pathway modulation, and ER stress. The pleiotropic function and peculiar structure of the DedA domain have prompted multiple studies addressing its mode of action and role in cellular homeostasis. An important point under evaluation is whether Vmp1 and Tmem41b work cooperatively or show specialization for different cellular processes. By deriving ESCs lacking the expression of Vmp1 and Tmem41b, we reported that their absence has a similar effect on autophagy and lipid droplet accumulation. Moreover, we show that simultaneous depletion of both genes exacerbate single mutant defects, particularly in the context of LD formation. Our findings support the existence of a partial redundancy between Vmp1 and Tmem41b activity. It would be tempting to speculate that variable expression levels could account for the contrasting cell dependencies observed across different cellular systems. While Tmem41b's transcriptional regulation is still largely unknown, Vmp1 expression is known to be influenced by external stimuli (Lo Re et al., 2012; Wang et al., 2020).

We find that ESCs can self-renew and maintain a substantial differentiation potential independent of TMEM41B and VMP1. Double knockout cells differentiate efficiently into lineages of the three germ layers despite severe defects in autophagy and lipid droplet metabolism (Mizushima and Levine, 2010). It would be interesting to further investigate how the lipid composition equilibrium between the inner and outer leaflets of ER membranes is maintained in the absence of Vmp1 and Tmem41b. Other ER-resident scramblases are Ca2+ dependent, proposing that compensation of Vmp1 and Tmem41b absence might require an increased energy status and/or could trigger supplementary signaling events.

We verified that ESCs lacking Vmp1 and Tmem41b show a strong delay in the differentiation of extraembryonic (XEN) stem cells *in vitro*. Embryoid body experiments further support defective primitive endoderm specification, which may account for the embryonic lethality observed in Vmp1 and Tmem41b KO embryos (Morishita et al., 2019; Van Alstyne et al., 2018). More permissive growth conditions may explain the requirement for simultaneous depletion of both genes *in vitro*, while the absence of one factor is sufficient to cause the primitive endoderm delay in embryos. Defects in endoderm specification were also observed in Embryoid bodies derived from single Vmp1[KO] or Tmem41b[KO] cells but with lower consistency than those derived from

Double KO cells. The finding of WNT signaling deregulation in Vmp1 and Tmem41b XEN cells, explains the specification delay detected in mutant cells: it has been shown that WNT/TCF7L1 transcriptional response is necessary *in vitro* and *in vivo* to promote PE expansion (Athanasouli et al., 2023). Retardation – not impairment – of PE specification induced by Vmp1 or Tmem41b mutations might still lead to embryo implantation but also to the accumulation of defects, which are incompatible with the gastrulation stage.

The direct dependence of the embryo's phenotype on defects on autophagy or lipid/lipoprotein distribution is unlikely. Both KO embryos die at day E8.5 with severe signs of developmental delay, which is already evident at E7.5. Mutant mice with defects in autophagy or LD metabolism either reach the pup stage in the former case (Kuma et al., 2017) or at least die at E10.5 after proper gastrulation and during organogenesis (Farese et al., 1995; Raabe et al., 1998). We propose that PE defects in Vmp1/Tmem41b depleted cells are caused by the increased activity of the WNT repressor DDIT3 which we show to be stabilized at the transcriptional and protein level in KO cells (Fig.4F). DDIT3 is activated via the PERK/ATF4 pathway by ER stress (Rozpedek et al., 2016) or Calcium-dependent mechanisms upon Vmp1 depletion in the absence of ER stress (Li et al., 2023a). Notably, downstream activation of the WNT cascade through chemicals fully rescues the defects of Vmp1 and Tmem41b mutant cells during XEN specification. Thereby, this study presents novel evidence illustrating the roles of Vmp1 and Tmem41b in acting at the interface of cell homeostasis and signaling modulation, adding to their scramblase activity the potential to modulate developmental processes.

MATERIALS AND METHODS

Cell Culture

Mouse ES cells were maintained as previously described in chemically defined 2i medium with added leukemia inhibitory factor (LIF) as described in Di Minin et al. (Di Minin et al., 2022). Base media composition: DMEM (Gibco, #41965039), 15% FBS (Biowest, #S1810-500) 1% sodium pyruvate (Gibco, 11360-070), 1% non-essential amino acids (Gibco, #11140-035). Lif (2000 U/mL, homemade), PD0325901 (1 µM, AxonMedchem, #391210-10-9) and Chiron (3 µM, AxonMedchem, #252917-06-9) are added fresh before use.

EB differentiation

To obtain embryonic bodies (EBs), 200k ESCs were plated on Day1 in ESCs base media without 2i and LIF (see cell culture) into a well of a Sphericalplate5D (Kugelmeiers). On Day4, EBs have formed and are transferred to a pHEMA coated 6cm culture dish and cultured until Day6. EBs are then collected and subjected to RNA extraction and RT-qPCR analysis or prepared for sectioning and immunofluorescence. In brief, EBs are fixed in 4% PFA for 30 min, cryoprotected in 10%, 20% and 30% sucrose and then embedded and frozen in Tissue Freezing Medium (Leica, #14020108926). 15 µm sections were prepared on a cryotome, mounted onto Superfrost Plus Adhesion slides (VWR international GmBH, #631-0108) and processed for immunofluorescence staining.

Cardiomyocyte differentiation

EBs were formed as previously described. On Day4, EBs were collected and seeded onto a non-treated 10cm dish, maintaining ESC base media. Cardiomyocytes randomly specified and were observed as early as on Day6 of differentiation.

Spinal Cord Organoids (SCOs) formation

For derivation of SCOs, ESCs were plated in EBs media (DMEM- F12 (Gibco) and Neurobasal Medium (Gibco) (1:1 ratio) supplemented with 200 mM L-glutamine and 10% of Knockout serum replacement) on Sphericalplate 5D dishes (Kugelmeiers AG). On day 2, SCOs were transferred in 10 cm tissue culture dishes and treated with RA (10 nM). SCOs were collected and processed for analysis at day 5.

Xen Differentiation

XEN cells were obtained following a protocol by Niakan et al. with minor changes (Niakan et al., 2013). In brief, 230k cells were plated in XEN medium on a gelatin-coated 3cm dish. The next day, the media was replaced with Xen medium containing 0.1 µM retinoic acid (Thermo Fischer, #17110052) and 10 ng/mL Activin A (Peprotech, #120-14). On Day4, the medium was again replaced by XEN medium, no further supplements were added, and on Day6 samples were either analyzed by IF or qPCR. Rescue of double KO Xen differentiation was performed by adding 1 µM Chiron (AxonMedchem, #252917-06-9) at Day2, together with the retinoic acid and Activin A.

Media was changed on day 4 to Xen medium. XEN medium composition: RPMI 1640 (Gibco, #61870010), GlutaMAX™ (Gibco, #61870036), 15% FBS (Biowest, #S1810-500), 0,1 mM ß-mercaptoethanol (Sigma, #BCCB9882).

CRISPR-Cas9 editing to generate mutant ESC lines

Vmp1: the gRNA (gagacgcatagcaatgagta) was cloned into a px330 vector (Addgene, #158973). Per 3cm well, 300k ESCs were plated and transfected with 2.5 µg of gRNA containing plasmid, as well as 0.25 µg of a GFP expressing plasmid (pCMV-GFP). GFP-positive cells were enriched by FACS and plated as single cells. Colonies were picked. Analysis by WB was used to identify identified VMP1 KO lines. Tmem41b: the gRNAs (gtatgtttgacctgggcgaa and gagtgacatgtggaaatcag) were cloned into px330 vectors, and KO clones were derived as described above. PCR analysis of the edited locus (FW: atcttggacaggggggagttc RV: gaccaggggttcattgtcatt and FW: cagcacacacctttaatccagc RV: gccacatagcaagcttaagagc) identified KO clones. qPCR of mRNA from Exon 2 (edited locus, N-terminal end) and Exon 7 (C-terminal end) further confirmed the absence of mRNA.

Proliferation assay

100k ESCs were plated in a well of a MW12. 48h later, cells were collected and counted. Proliferation rates were calculated as relative factor of cells collected over cells seeded.

LC3 staining and size characterization

100k cells were plated on cover-slips, pre-treated with Matrigel (Corning, # 354277), in N2B27 media. N2B27 composition: 50 mL Neurobasal media (Gibco, # 2348949), 50 mL Advanced DMEM F12 (Gibco, # 12634010), 4% BSA (Gibco, # 15260), 1% L-Glutamine (Gibco, # 25030024), 1% B27 supplement (Gibco, # 2336837), 0,5% N2 supplement (Gibco, # 2328253), and 0,1 mM ß-mercapotethanol (Sigma, #BCCB9882). The next day, samples were fixed in 4% PFA for 15 min, washed 3 times with PBS and then treated with ice-cold MeOH for 5 min on RT. The staining was then performed as described below. Quantification was performed with the Fiji plugin TrackMate (Ershov et al., 2022; Tinevez et al., 2017) to automatically detect puncti with a diameter of 1, 2, and 3 microns at a threshold of 600, 200 and 150, respectively. Nuclei were detected at an average size of 15 microns with a threshold of 10. Unique

particles were annotated for 5 different frames for each condition and shown as particles per cell. The experiment was repeated 3 times.

LD staining and RF microscopy

Samples were plated as for the LC3 staining. For samples treated with oleic acid, 200 µM (Sigma-Aldrich, # O1008-1G) were added at plating, over-night. The next day, samples were incubated in 5 µM BODIPY 493/503 (Invitrogen, #D3922) in PBS and subsequently washed 3 times with PBS and fixed in 4 % PFA for 15 min. Samples were then washed, stained with Dapi for 15 min at RT and mounted with Mowiol. Quantification was performed using the Fiji plugin TrackMate (Ershov et al., 2022; Tinevez et al., 2017) detecting droplets of 1, 2, and 3 microns in diameter at a threshold of 900. Nuclei were detected at 10 micron and a threshold of 1.

For RF microscopy, 250k cells were plated per well of a 6 well glass bottom plate (Cellvis, # P06-1.5H-N) in N2B27 media. After 36 h samples were treated with 200 µM oleic acid o/n and then with BODIPY for 3 h. Samples were washed 3 times with PBS and maintained in N2B27 for RF microscopy. The imaging was performed on Tomocube HT-X1 instrument at 37C and 5 % CO_2. At least 5-8 images were acquired for each condition for each well. The z-stacks were acquired with 1.06 µm slices. Max Intensity projections (MIP) are generated on the Tomostudio 3.3.9. The fluorescence max intensity projections were resized to Refractive index tomograms for comparison. The ROIs from each image were isolated for further analysis. Comdet plugin was used for identifying high refractive index regions. The approximate particle size was 2 pixels and Intensity threshold values were chosen individually for each image to avoid any false positives or negatives.

Immunofluorescence

Samples were washed 3 times with PBS, fixed in 4% PFA for 10 min and again washed 3 times with PBS. Samples stained for LC3 were then treated for 5 min with 100% MeOH at RT and washed again 3 times with PBS. Samples were then blocked and permeabilized with 10% donkey serum (Jackson Immunoresearch, #017-000-121) and 0,1% Triton X (Thermo Fischer, # HFH10) in PBS for 20 min and incubated with primary antibodies at 4 °C over-night. Oct4 (1:100, Satna Cruz, # sc-5279), Nanog (1:100, Abcam, # ab80892), Sox2 (1:50, R&D Systems #AF2018), Lc3 (1:200, Cell Signalling, # 2775S), Pdi (1:200, Cell Signalling, # 3501S), Rcas1 (1:200, Cell

Signalling, # 12290S), Rab7 (1:200, Cell Signaling, # 9367S) Sox1 (1:50, R&D Systems, # AF3369), Sox17 (1:200 R&D Systems, # AF1924-SP), Gata4 (1:100 R&D Systems, # AF1700-SP), Gata6 (1:100 R&D Systems, # AF1700-SP), Dab2 (1:100 BD, # 610464). The next day, samples were washed 3 times with 0.1% Triton X in PBS and then incubated with secondary antibodies and Dapi.

Western Blot Analysis

Cells were lysed in membrane buffer (150 mM NaCl, 50 mM Tris-HCl pH 7,5, 1 mM 2-mercaptoethanol (Sigma, #M3148) and 1% CHAPS (Sigma, #C3023) in water). Lysates were run on SDS-Page and transferred to nitrocellulose membranes. Bands were detected with the Amersham ECL Prime Western Blotting Detection Reagents (#RPN2232) and captured on Fujifilm Super RX-N films (#47410). Antibodies for Vmp1 (Cell Signalling, #12929S) were used at 1:1000 and DDIT3 1:1000 (Cell Signalling, #2895T). Quantification of western blot bands was performed with ImageJ, normalized to HSP90 (Santa Cruz, #sc-101494).

RNA extraction and RT qPCR

RNA was extracted with the RNeasy Kit (Quiagen, #74104) following the manufacturer's instructions, including the on-column DNA digestion. EB and SCO samples were pre-treated with the QIAshredder (Quiagen, #79656). cDNAs were prepared using the Bio-Rad iScript Reverse Transcription Supermix for RT-qPCR (#1708841). RT qPCR samples were prepared using the KAPPA SYBR R fast (Roche, KK4611). Primer Sequences:
Tmem41b Exon2 FW: TGCAGAAGCTGGATCAGCAA, RV: GAGCCTTGGCGTCATCCATA
Tmem41b Exon7 FW: GCAGGTAGAGCGTCACAGAG, RV: TGCAGAGTTGTTCCTGCCTT
Sdha FW: CGTTATGTGAGGGGTGTGCCTT, RV: CCTTACCAAACCTTGTGTCTGGA
Sox1 FW: ACGTAGCCAACGTAGACAC, RV: ATGAGCGTCGTCCCGTGG
Fgf5 FW: GCCTGTCCTTGCTCTTCCTCAT, RV: GGAGAAGCTGCGACTGGTGA
Actc1 FW: TGTAGGACCGTGTCGAAACC, RV: CGAAACCACACACTGTTACCG

Brachyury FW: TCTCTGGTCTGTGAGCAATGGT, RV:

TGCGTCAGTGGTGTGTAATGTG

Gata6 FW: GAAGCGCGTGCCTTCATC, RV: GTAGTGGTTGTGGTGTGACAGTTG

Dab2 FW: TTGATGATGTGCCTGATGCT, RV: TTTGCTTGTGTTGTCCCTGA

Sox17 FW: CCCAACACTCCTCCCAAAGTATC, RV:

TTCCCTGTCTTGGTTGATTTCT

RNA seq

RNA samples were submitted to the Functional Genomics Center in Zurich (FGCZ) for

RNA sequencing. The quality of the RNA was determined with a Fragment Analyzer

standard sensitivity RNA measurement (SS RNA kit (15 nt), Agilent, Waldbronn,

Germany). The measured concentrations (> 50 ng/µl) and RIN (>9.9) values qualified

for a Poly-A enrichment strategy in order to generate the sequencing libraries applying

the TruSeq mRNA Stranded Library Prep Kit (Illumina, Inc, California, USA). After Poly-

A selection using Oligo-dT beads the mRNA was reverse-transcribed into cDNA. The

cDNA was fragmented, end-repaired and polyadenylated before ligation of TruSeq UD

Indices (IDT, Coralville, Iowa, USA). The quality and quantity of the amplified

sequencing libraries were validated using a Fragment Analyzer SS NGS Fragment Kit

(1–6000 bp) (Agilent, Waldbronn, Germany). The equimolar pool of samples was

spiked into a NovaSeq6000 run targeting 20M reads per sample on a S1 FlowCell

(Novaseq S1 Reagent Kit, 100 cycles, Illumina, Inc, California, USA). Reads from

RNA-seq were first preprocessed by Trimmomatic (v0.40) (Bolger et al., 2014) to

remove low-quality bases and adapters. Then, reads were aligned to mouse genome

GRCm38 (release 81 version from Ensembl) using HISAT2 (v2.2.1) (Kim et al., 2019).

HTSeq Count (v2.0.2) (Putri et al., 2022) was used to count reads for each gene

(Ensembl GRCm38.81), ignoring reads on overlapped regions. Differentially

expressed genes were defined by DESeq2 (v1.40.2) (Love et al., 2014). MA plot was

generated using custom-made scripts. Genes significantly enriched show a fold

change > 1 and FDR (adjusted p-value) < 0.01. The read counts were transformed in

RPKM for the heatmap representations of specific genes and pathways. The genes

corresponding to germ-layer differentiation markers (including those for embryonic and

extra-embryonic endoderm), biological processes, and signaling pathways were

obtained through literature mining and the KEGG database (Table S1). The processed

data are available as excel files (Table S2 and Table S3) and raw data have been

deposited on NCBI SRA database under the accession: PRJNA1031567

Competing interest

476 The authors declare no competing interests.

Acknowledgments

We acknowledge the Functional Genomics Center Zurich (FGCZ) for technical support

and Sungsik Lee from ScopeM facility for help on the RF microscopy. We thank G.J.

Pereira, C. Fimiani, and U. Sutter for their help with reagents. We thank C. Ciaudo, A.

Grison, J. Corn, and K. Basler for their helpful discussions. GDM was supported by

the ETH Zurich Postdoctoral Fellowship Program as well as the Marie Curie Actions

for People COFUND Program. This work was supported by grants from the Swiss

National Science Foundation (SNF grants 31003A_152814/1 and 31003A_175643/1).

Author Contributions

Conceptualization: MH, AW and GDM; Data curation: MH, VG and GDM; Formal

analysis: MH, TS, HH, VG, and GDM; Funding acquisition: KW, AW and GDM;

Investigation: MH, AW and GDM; Methodology: MH, TS, HH, VG, and GDM; Project

administration: AW, GDM; Resources: MH, KW, AW and GDM; Supervision: KW, AW

and GDM; Validation: MH, GDM; Visualization: MH, VG and GDM; Writing – original

draft: MH, AW and GDM; Writing – review & editing: MH, TS, HH, VG, KW, AW and

GDM.

REFERENCES

Arnold, S.J., and Robertson, E.J. (2009). Making a commitment: cell lineage allocation and axis patterning in the
early mouse embryo. Nat Rev Mol Cell Biol *10*, 91-103.
Athanasouli, P., Balli, M., De Jaime-Soguero, A., Boel, A., Papanikolaou, S., van der Veer, B.K., Janiszewski, A.,
Vanhessche, T., Francis, A., El Laithy, Y.*, et al.* (2023). The Wnt/TCF7L1 transcriptional repressor axis drives
primitive endoderm formation by antagonizing naive and formative pluripotency. Nat Commun *14*, 1210.
Baggen, J., Vanstreels, E., Jansen, S., and Daelemans, D. (2021). Cellular host factors for SARS-CoV-2 infection.
Nat Microbiol *6*, 1219-1232.
Bard, F., Casano, L., Mallabiabarrena, A., Wallace, E., Saito, K., Kitayama, H., Guizzunti, G., Hu, Y., Wendler, F.,
Dasgupta, R.*, et al.* (2006). Functional genomics reveals genes involved in protein secretion and Golgi
organization. Nature *439*, 604-607.
Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics *30*, 2114-2120.

Chiu, C.S., Tsai, C.H., Hsieh, M.S., Tsai, S.C., Jan, Y.J., Lin, W.Y., Lai, D.W., Wu, S.M., Hsing, H.Y., Arbiser, J.L., *et al.*
(2019). Exploiting Honokiol-induced ER stress CHOP activation inhibits the growth and metastasis of melanoma
by suppressing the MITF and beta-catenin pathways. Cancer Lett *442*, 113-125.

Di Minin, G., Holzner, M., Grison, A., Dumeau, C.E., Chan, W., Monfort, A., Jerome-Majewska, L.A., Roelink, H.,
and Wutz, A. (2022). TMED2 binding restricts SMO to the ER and Golgi compartments. PLoS Biol *20*, e3001596.

Du, Y.C., Oshima, H., Oguma, K., Kitamura, T., Itadani, H., Fujimura, T., Piao, Y.S., Yoshimoto, T., Minamoto, T.,
Kotani, H., *et al.* (2009). Induction and down-regulation of Sox17 and its possible roles during the course of
gastrointestinal tumorigenesis. Gastroenterology *137*, 1346-1357.

Ershov, D., Phan, M.S., Pylvanainen, J.W., Rigaud, S.U., Le Blanc, L., Charles-Orszag, A., Conway, J.R.W., Laine, R.F.,
Roy, N.H., Bonazzi, D., *et al.* (2022). TrackMate 7: integrating state-of-the-art segmentation algorithms into
tracking pipelines. Nat Methods *19*, 829-832.

Farese, R.V., Jr., Ruland, S.L., Flynn, L.M., Stokowski, R.P., and Young, S.G. (1995). Knockout of the mouse
apolipoprotein B gene results in embryonic lethality in homozygotes and protection against diet-induced
hypercholesterolemia in heterozygotes. Proc Natl Acad Sci U S A *92*, 1774-1778.

Folkerts, H., Wierenga, A.T., van den Heuvel, F.A., Woldhuis, R.R., Kluit, D.S., Jaques, J., Schuringa, J.J., and
Vellenga, E. (2019). Elevated VMP1 expression in acute myeloid leukemia amplifies autophagy and is protective
against venetoclax-induced apoptosis. Cell Death Dis *10*, 421.

Ghanbarpour, A., Valverde, D.P., Melia, T.J., and Reinisch, K.M. (2021). A model for a partnership of lipid transfer
proteins and scramblases in membrane expansion and organelle biogenesis. Proc Natl Acad Sci U S A *118*.

Gilabert, M., Vaccaro, M.I., Fernandez-Zapico, M.E., Calvo, E.L., Turrini, O., Secq, V., Garcia, S., Moutardier, V.,
Lomberk, G., Dusetti, N., *et al.* (2013). Novel role of VMP1 as modifier of the pancreatic tumor cell response to
chemotherapeutic drugs. J Cell Physiol *228*, 1834-1843.

Guo, L., Yang, L.Y., Fan, C., Chen, G.D., and Wu, F. (2012). Novel roles of Vmp1: inhibition metastasis and
proliferation of hepatocellular carcinoma. Cancer Sci *103*, 2110-2119.

Hama, Y., Morishita, H., and Mizushima, N. (2022). Regulation of ER-derived membrane dynamics by the DedA
domain-containing proteins VMP1 and TMEM41B. EMBO Rep *23*, e53894.

Hoffmann, H.H., Schneider, W.M., Rozen-Gagnon, K., Miles, L.A., Schuster, F., Razooky, B., Jacobson, E., Wu, X.,
Yi, S., Rudin, C.M., *et al.* (2021). TMEM41B Is a Pan-flavivirus Host Factor. Cell *184*, 133-148 e120.

Huang, D., Xu, B., Liu, L., Wu, L., Zhu, Y., Ghanbarpour, A., Wang, Y., Chen, F.J., Lyu, J., Hu, Y., *et al.* (2021).
TMEM41B acts as an ER scramblase required for lipoprotein biogenesis and lipid homeostasis. Cell Metab *33*,
1655-1670 e1658.

Jiang, X., Fulte, S., Deng, F., Chen, S., Xie, Y., Chao, X., He, X.C., Zhang, Y., Li, T., Li, F., *et al.* (2022). Lack of VMP1
impairs hepatic lipoprotein secretion and promotes non-alcoholic steatohepatitis. J Hepatol *77*, 619-631.

Kim, D., Paggi, J.M., Park, C., Bennett, C., and Salzberg, S.L. (2019). Graph-based genome alignment and
genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol *37*, 907-915.

Kratzel, A., Kelly, J.N., V'Kovski, P., Portmann, J., Bruggemann, Y., Todt, D., Ebert, N., Shrestha, N., Plattet, P., Staab-
Weijnitz, C.A., *et al.* (2021). A genome-wide CRISPR screen identifies interactors of the autophagy pathway as
conserved coronavirus targets. PLoS Biol *19*, e3001490.

Kruithof-de Julio, M., Alvarez, M.J., Galli, A., Chu, J., Price, S.M., Califano, A., and Shen, M.M. (2011). Regulation
of extra-embryonic endoderm stem cell differentiation by Nodal and Cripto signaling. Development *138*, 3885-
3895.

Kuma, A., Komatsu, M., and Mizushima, N. (2017). Autophagy-monitoring and autophagy-deficient mice.
Autophagy *13*, 1619-1628.

Li, T., Zhao, H., Guo, G., Xia, S., and Wang, L. (2023a). VMP1 affects endoplasmic reticulum stress sensitivity via
differential modulation of the three unfolded protein response arms. Cell Rep *42*, 112209.

Li, X.Q., Zeng, L., Liang, D.G., Qi, Y.L., Yang, G.Y., Zhong, K., Chu, B.B., and Wang, J. (2023b). TMEM41B Is an
Interferon-Stimulated Gene That Promotes Pseudorabies Virus Replication. J Virol *97*, e0041223.

Li, Y.E., Wang, Y., Du, X., Zhang, T., Mak, H.Y., Hancock, S.E., McEwen, H., Pandzic, E., Whan, R.M., Aw, Y.C., *et al.*
(2021). TMEM41B and VMP1 are scramblases and regulate the distribution of cholesterol and
phosphatidylserine. J Cell Biol *220*.

Lin, W., Sun, Y., Qiu, X., Huang, Q., Kong, L., and Lu, J.J. (2021). VMP1, a novel prognostic biomarker, contributes
to glioma development by regulating autophagy. J Neuroinflammation *18*, 165.

Lo Re, A.E., Fernandez-Barrena, M.G., Almada, L.L., Mills, L.D., Elsawa, S.F., Lund, G., Ropolo, A., Molejon, M.I.,
Vaccaro, M.I., and Fernandez-Zapico, M.E. (2012). Novel AKT1-GLI3-VMP1 pathway mediates KRAS oncogene-
induced autophagy in cancer cells. J Biol Chem *287*, 25325-25334.

Loncle, C., Molejon, M.I., Lac, S., Tellechea, J.I., Lomberk, G., Gramatica, L., Fernandez Zapico, M.F., Dusetti, N., Urrutia, R., and Iovanna, J.L. (2016). The pancreatitis-associated protein VMP1, a key regulator of inducible autophagy, promotes Kras(G12D)-mediated pancreatic cancer initiation. Cell Death Dis *7*, e2295.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550.

Mizushima, N., and Levine, B. (2010). Autophagy in mammalian development and differentiation. Nat Cell Biol *12*, 823-830.

Molejon, M.I., Ropolo, A., Re, A.L., Boggio, V., and Vaccaro, M.I. (2013). The VMP1-Beclin 1 interaction regulates autophagy induction. Sci Rep *3*, 1055.

Moretti, F., Bergman, P., Dodgson, S., Marcellin, D., Claerr, I., Goodwin, J.M., DeJesus, R., Kang, Z., Antczak, C., Begue, D.*, et al.* (2018). TMEM41B is a novel regulator of autophagy and lipid mobilization. EMBO Rep *19*.

Morishita, H., Zhao, Y.G., Tamura, N., Nishimura, T., Kanda, Y., Sakamaki, Y., Okazaki, M., Li, D., and Mizushima, N. (2019). A critical role of VMP1 in lipoprotein secretion. Elife *8*.

Ngondo, R.P., Cirera-Salinas, D., Yu, J., Wischnewski, H., Bodak, M., Vandormael-Pournin, S., Geiselmann, A., Wettstein, R., Luitz, J., Cohen-Tannoudji, M.*, et al.* (2018). Argonaute 2 Is Required for Extra-embryonic Endoderm Differentiation of Mouse Embryonic Stem Cells. Stem Cell Reports *10*, 461-476.

Niakan, K.K., Schrode, N., Cho, L.T., and Hadjantonakis, A.K. (2013). Derivation of extraembryonic endoderm stem (XEN) cells from mouse embryos and embryonic stem cells. Nat Protoc *8*, 1028-1041.

O'Rourke, E.J., Soukas, A.A., Carr, C.E., and Ruvkun, G. (2009). C. elegans major fats are stored in vesicles distinct from lysosome-related organelles. Cell Metab *10*, 430-435.

Putri, G.H., Anders, S., Pyl, P.T., Pimanda, J.E., and Zanini, F. (2022). Analysing high-throughput sequencing data in Python with HTSeq 2.0. Bioinformatics *38*, 2943-2945.

Raabe, M., Flynn, L.M., Zlot, C.H., Wong, J.S., Veniant, M.M., Hamilton, R.L., and Young, S.G. (1998). Knockout of the abetalipoproteinemia gene in mice: reduced lipoprotein secretion in heterozygotes and embryonic lethality in homozygotes. Proc Natl Acad Sci U S A *95*, 8686-8691.

Rodriguez, M.E., Catrinacio, C., Ropolo, A., Rivarola, V.A., and Vaccaro, M.I. (2017). A novel HIF-1alpha/VMP1-autophagic pathway induces resistance to photodynamic therapy in colon cancer cells. Photochem Photobiol Sci *16*, 1631-1642.

Ropolo, A., Catrinacio, C., Renna, F.J., Boggio, V., Orquera, T., Gonzalez, C.D., and Vaccaro, M.I. (2020). A Novel E2F1-EP300-VMP1 Pathway Mediates Gemcitabine-Induced Autophagy in Pancreatic Cancer Cells Carrying Oncogenic KRAS. Front Endocrinol (Lausanne) *11*, 411.

Rozpedek, W., Pytel, D., Mucha, B., Leszczynska, H., Diehl, J.A., and Majsterek, I. (2016). The Role of the PERK/eIF2alpha/ATF4/CHOP Signaling Pathway in Tumor Progression During Endoplasmic Reticulum Stress. Curr Mol Med *16*, 533-544.

Sun, L., Zhao, C., Fu, Z., Fu, Y., Su, Z., Li, Y., Zhou, Y., Tan, Y., Li, J., Xiang, Y.*, et al.* (2021). Genome-scale CRISPR screen identifies TMEM41B as a multi-function host factor required for coronavirus replication. PLoS Pathog *17*, e1010113.

Tetsu, O., and McCormick, F. (1999). Beta-catenin regulates expression of cyclin D1 in colon carcinoma cells. Nature *398*, 422-426.

Tinevez, J.Y., Perry, N., Schindelin, J., Hoopes, G.M., Reynolds, G.D., Laplantine, E., Bednarek, S.Y., Shorte, S.L., and Eliceiri, K.W. (2017). TrackMate: An open and extensible platform for single-particle tracking. Methods *115*, 80-90.

Tornero-Ecija, A., Tabara, L.C., Bueno-Arribas, M., Anton-Esteban, L., Navarro-Gomez, C., Sanchez, I., Vincent, O., and Escalante, R. (2022). A Dictyostelium model for BPAN disease reveals a functional relationship between the WDR45/WIPI4 homolog Wdr45l and Vmp1 in the regulation of autophagy-associated PtdIns3P and ER stress. Autophagy *18*, 661-677.

Trimarco, J.D., Heaton, B.E., Chaparian, R.R., Burke, K.N., Binder, R.A., Gray, G.C., Smith, C.M., Menachery, V.D., and Heaton, N.S. (2021). TMEM41B is a host factor required for the replication of diverse coronaviruses including SARS-CoV-2. PLoS Pathog *17*, e1009599.

Van Alstyne, M., Lotti, F., Dal Mas, A., Area-Gomez, E., and Pellizzoni, L. (2018). Stasimon/Tmem41b localizes to mitochondria-associated ER membranes and is essential for mouse embryonic development. Biochem Biophys Res Commun *506*, 463-470.

Wang, C., Peng, R., Zeng, M., Zhang, Z., Liu, S., Jiang, D., Lu, Y., and Zou, F. (2020). An autoregulatory feedback loop of miR-21/VMP1 is responsible for the abnormal expression of miR-21 in colorectal cancer cells. Cell Death Dis *11*, 1067.

Yao, Y., Gu, J., Li, M., Li, G., Ai, J., and Zhao, L. (2023). WHSC1L1-mediated epigenetic downregulation of VMP1
participates in herpes simplex virus 1 infection-induced mitophagy impairment and neuroinflammation. Mol
Immunol *163*, 63-74.
Zack, S.R., Nikolaienko, R., Cook, B., Melki, R., Zima, A.V., and Campbell, E.M. (2023). Vacuole Membrane Protein
1 (VMP1) Restricts NLRP3 Inflammasome Activation by Modulating SERCA Activity and Autophagy. Res Sq.
Zeevaert, K., Elsafi Mabrouk, M.H., Wagner, W., and Goetzke, R. (2020). Cell Mechanics in Embryoid Bodies. Cells
9.
Zhang, Z., Deb, A., Zhang, Z., Pachori, A., He, W., Guo, J., Pratt, R., and Dzau, V.J. (2009). Secreted frizzled related
protein 2 protects cells from apoptosis by blocking the effect of canonical Wnt3a. J Mol Cell Cardiol *46*, 370-377.
628
629

 FIGURE LEGENDS

Figure 1. Vmp1 and Tmem41b double KO ESCs display increased
autophagosome and lipid droplet accumulation

(A) Schematic representation of the CRISPR/Cas9-mediated KO strategy for Vmp1
and Tmem41b genes. For depleting Vmp1 (top) a single gRNA inducing an indel
mutation has been used. In the case of the Tmem41b gene (bottom), two gRNAs were
used to delete exons 2 and 3. The sequences coding for transmembrane domains
(TM) are indicated in blue, the conserved DedA domain is highlighted in yellow, red
lines indicate cut sites of gRNAs. **(B)** Colony morphology of VMP1^{KO} and Tmem41^{KO}
ESCs remains unchanged compared to WT ESCs. Scale bar: 100 μm. **(C)** VMP1 and
Tmem41b depletion induce accumulation of autophagosomes in ESCs. The graph
shows the quantification of LC3 puncta per cell of a representative experiment (see
Fig.S2A). Data were collected from more than 200 cells. For each KO condition, two
independent clones have been analyzed. Dots correspond to the frames analyzed per
sample. Data are represented as mean ± SD **(D)** Vmp1 and Tmem41b depletion leads
to the accumulation of lipid droplets (LD) in ESCs. Graph quantifies LD per cell in WT
and KO ESCs (see Fig.S2D). ± oleic acid treatment. At least 500 cells were analyzed
for every condition; dots represent independent biological replicates. Data are
represented as mean ± SD. **(E-G)** Transcriptional analysis of Double KO ESCs did not
identify many deregulated genes. **(E)** Shown are differentially regulated genes in
Double KO ESCS in blue as MA plot. Genes highlighted in red are pluripotency
markers. **(F)** Diagram showing uniquely and collectively differentially expressed genes
in Vmp1, Tmem41b, and Double KO ESCs. **(G)** Differentially regulated genes do not
cluster in essential processes. The graph shows differentially expressed genes of
single and double KO ESCs by cellular processes. Dot size correlates to the number
of deregulated genes within a cellular process. **(H-I)** Double KO ESCs accumulate
LC3+ autophagosomes of increased size. **(H)** Shown are representative IF images of
WT and double KO ESCs stained for LC3. Nuclei are stained with Dapi. Scale bar:
20 μm. **(I)** Plotted is the quantification of LC3 puncta of 3 μm in diameter. For each
condition, two independent clones have been analyzed. Dots correspond to the frames
analyzed per sample. Data were collected from more than 200 cells. Data are
represented as mean ± SD. **(J)** Double KO ESCs accumulate LDs without oleic acid

treatment. The number of LDs per cell is plotted for all KO clones. Data are
represented as mean ± SD. **(K)** Double KO ESCs present an increased amount of LD
visualized through refractive index microscopy (top) and co-stained with BODIPY.
667
668 **Figure 2. Double KO ESCs specify all germ layers, but not the primitive**
669 **endoderm**
670
(A) Schematic representation of an Embryonic Body (EB) which consists of 3 different
lineages. The ectoderm expresses Fgf5 and Sox1 (grey), the mesoderm Actc1 and
Brachyury (blue), and the endoderm by Gata6 and Dab2 (yellow). **(B)** ESCs
differentiated to EBs express lineage markers for ectoderm (Fgf5), mesoderm
(Brachyury), and endoderm (Gata6). Graphs show mRNA levels of WT EBs
normalized to the Sdha gene. Dots represent biological independent experiments.
Data are represented as mean ± SD. **(C)** Double KO ESCs specifically show defects
in expressing the primitive endoderm marker, Dab2. qPCR analysis for Sox1, Actc1
and Dab2 mRNA expression in mutant EBs relative to WT EBs. Two independent
clones were analyzed per cell line. Each dot represents an independent experiment.
Data are represented as mean ± SD**.** The line at 0,7 indicates an arbitrary threshold
after which successful expression of the marker is considered. **(D-F)** Double mutant
EBs show decreased expression of endoderm markers. Immunofluorescence images
of EB sections stained for the endoderm markers GATA4 and SOX17 **(D)**, GATA6 **(E)**,
DAB2 **(F)**. WT in the top panel, two independent clones for Double KO condition in the
center and bottom panel. Nuclei are stained with Dapi. Scale bar: 50 μm.
687
688 **Figure 3. XEN differentiation in Double KO cells is delayed**
689
(A-B) Double KO ESCs show a delay in XEN differentiation. **(A)** Brightfield images of
XEN cells at day 6 of differentiation in WT and in Double KO cells. XEN islands are
circled in white in the Double KO condition. Scale bar: 100 μm. **(B)**
Immunofluorescence images of day 6 XEN cells stained for SOX17 and GATA4. Nuclei
are stained with Dapi, Scale bar: 150 μm. **(C-F)** Transcriptional analysis of XEN cells.
(C) Differentially expressed genes between Double KO and WT cells at day 2 during
XEN specification. XEN markers are labeled in red, differentially expressed genes in
blue. **(D-E)** GO-Term analysis of differential expressed genes between Double KO and

 WT conditions in ESCs and XEN cells in the context of developmental processes **(D)**

 and signaling pathways **(E)**. Dot size corresponds to differentially regulated genes. **(F)**

 WNT pathway related genes differentially expressed in Double KO cells. Heatmap

 shows min to max expression of three biological replicates for WT and Double KO

 XEN cells.

 Figure 4. Activation of WNT signaling fully rescues the delay of Double KO cells

 in XEN differentiation

 (A-B) VMP1 and Tmem41b depletion promotes increase of DDIT3 protein levels. **(A)**

 Western Blot analysis of DDIT3 expression in WT and Double KO cells differentiated

 to XEN cells for 2 days. HSP90 is shown as loading control. **(B)** Quantification of

 increased DDIT3 protein levels in three biological replicates. Expression relative to WT

 XEN cell and normalized to respective HSP90 expression. Data are represented as

 mean ± SD. **(C-E)** Chiron-dependent WNT activation rescues the delay of Double KO

 XEN differentiation. **(C)** Brightfield images show XEN clusters of double KO XEN cells

 (highlighted by white lines) with and w/o Chiron treatment. Scale bar: 100 μm. IF

 staining for SOX17 and GATA4 **(D)** or DAB2 and GATA6 **(E)**; nuclei are stained with

 Dapi. Scale Bar: 150 μm. **(F)** Schematic overview highlighting the correlation between

 loss of Vmp1 and Tmem41b and decreased WNT signaling via DDIT3.

Figure1

Figure 3

A

DDIT3

HSP90

WT   Double KO

B

DDIT3

FC protein level

WT   Double KO

C

NT   +CHIR

D

WT   Double KO

NT

+CHIR

SOX17 GATA4 DAPI

E

Double KO

DAB2 GATA6 DAPI

F

TMEM41B

VMP1

LD ↑
Autophagy ↓

Ddit3

WNT
signaling

XEN
spec.