The scramblases VMP1 and TMEM41b are required for primitive

endoderm specification by targeting WNT signaling

Markus Holzner1, Tea Sonicki1, Hugo Hunn1, Vamshidhar R. Gade2, Karsten Weis2,

Anton Wutz1#, and Giulio Di Minin1#

1 Institute of Molecular Health Sciences, Department of Biology, ETH Zurich, Zurich,

Switzerland

2 Institute of Biochemistry, Department of Biology, ETH Zurich, Zurich, Switzerland

corresponding authors: awutz@ethaz.ch (AW); giulio.diminin@biol.ethz.ch (GDM)

12
13

14 Keywords: Vmp1, Tmem41b, WNT pathway, ESCs, Primitive endoderm, Autophagy

15 and Lipid droplets

17 Characters (with space): 33500

18 Abstract: 158 words

19 4 figures

20
21

SUMMARY

The ER resident proteins VMP1 and TMEM41b share a DedA domain, which confers

lipid scramblase activity. Loss of either gene results in embryonic lethality in mice and

defects in autophagy and lipid droplet metabolism. To understand their role in

pluripotency and specification, we generated Vmp1 and Tmem41b mutant mouse

embryonic stem cells (ESCs). We observe that ESCs carrying mutations in Vmp1 and

Tmem41b are viable, proliferate normally, and maintain a largely intact pluripotency-

associated transcriptional profile. Despite clear defects in the accumulation of LC3-

positive autophagosomes and lipid droplets, ESCs carrying combined mutations in

Vmp1 and Tmem41b can differentiate into a wide range of embryonic cell types.

However, the combined loss of Vmp1 and Tmem41b impairs the specification of

primitive endoderm-like cells. We observe a delayed differentiation of mutant ESCs

into extra-embryonic endoderm stem (XEN) cells. Mechanistically, we discover that

mutant cells upregulate DDIT3, a WNT inhibitor. Chemical stimulation of the WNT

cascade can rescue the differentiation delay. Our findings reveal a redundant function

of the lipid scramblases VMP1 and TMEM41b and identify a specific role in signaling

during extra-embryonic endoderm development.

INTRODUCTION

Vesicle trafficking and membrane synthesis are intricately interconnected within cellular signaling networks. Lipid scramblases, with their property to shape lipid membrane composition, play a crucial role in facilitating inter-organelle communication. Beyond their integral involvement in maintaining cell functionality, recent studies have unveiled their implications in various pathological conditions, such as viral infection, cancer development, and inflammatory diseases (Hama et al., 2022). The vacuole membrane protein 1 (VMP1) and transmembrane protein 41B (TMEM41B) are lipid scramblases with a highly conserved DedA domain (Ghanbarpour et al., 2021; Li et al., 2021). They localize at the ER membrane and are distinguished from other ER-associated flippases or scramblases by the property to facilitate bi-directional lipid diffusion in an ATP- and Ca^{2+}-independent manner. VMP1 and TMEM41B are necessary for autophagosome formation (Molejon et al., 2013; Moretti et al., 2018), lipid droplet metabolism (Li et al., 2021; Morishita et al., 2019), and lipoprotein formation (Hama et al., 2022). Additional reports proposed a more global role in ER protein homeostasis (Tornero-Ecija et al., 2022) and vesicle transport (Bard et al., 2006). Despite multifaceted functions, their absence does not influence cell survival significantly. Compensatory mechanisms shared by VMP1 and TMEM41B or other ER scramblases likely maintain cellular homeostasis. While VMP1 depletion leads to a pronounced defect in autophagosome assembly (Molejon et al., 2013; Moretti et al., 2018), TMEM41B appears to play a pivotal role in sustaining lipid droplet homeostasis and serves as a crucial factor in RNA virus replication (Hoffmann et al., 2021; Huang et al., 2021). However, a systematic comparison between the activities of VMP1 and TMEM41B has not yet been performed.

VMP1 and TMEM41B are both essential during embryonic development. Vmp1 (Morishita et al., 2019) and Tmem41b (Van Alstyne et al., 2018) deficient mice die around E8.5 following severe developmental delay. Prior studies formulated a hypothesis where VMP1 restricts the transfer of vital nutrients to the embryo by blocking the synthesis and release of lipoproteins from the visceral endoderm. Nonetheless, mice with defects in lipoprotein biogenesis experience mortality at later stages (Farese et al., 1995; Raabe et al., 1998), specifically at E10.5. This observation implies a distinct and earlier involvement of VMP1 and TMEM41B in embryo development.

During implantation, the mouse embryo is composed of three lineages: the
trophectoderm, which gives rise to the placenta; the primitive endoderm (PE),
contributing to the yolk sac; and the epiblast (EPI), which is primed to develop into the
embryo (Arnold and Robertson, 2009). The EPI and PE are specified within the inner
cell mass (ICM) of the blastocyst. Defects in the formation of these lineages are
challenging to characterize *in vivo* as they appear pre-implantation and due to the
small embryo size. To investigate the role of Vmp1 and Tmem41b in early embryo
lineage specification, we studied the ability of mutant cells to differentiate *in vitro*
towards embryonic germ layers. We found that a combined Vmp1 and Tmem41b
mutation in ESCs results in a specific defect in PE formation and a delayed conversion
to extra-embryonic endoderm cells (XEN). This delay is caused by imbalanced WNT
signaling and can be rescued through artificial WNT activation.

RESULTS

Vmp1 and Tmem41b mutant ESCs accumulate LC3+ autophagosomes and lipid
droplets
To assess the influence of a Vmp1 or Tmem41b depletion on mouse ESCs and in early
embryo development, we engineered genomic deletions using CRISPR/Cas9 targeted
mutagenesis. Vmp1 was mutated by introducing a frameshift in exon 2 (Fig.1A). Loss
of Vmp1 protein was confirmed in independent mutant ESC clones (Vmp1^{KO})
(Fig.S1A). To abrogate Tmem41b expression, we deleted the genomic region between
exons 2 and 3 (Fig.1A), which encodes the first transmembrane domain of the protein
(Tmem41b^{KO}). This mutation is predicted to compromise the protein topology.
Additionally, the splicing of exon 1 to exon 4 produces a frameshift, suggesting a null
allele. The deletion was confirmed on the genomic DNA (Fig.S1B). Tmem41b mRNA
levels were strongly reduced (Fig.S1C-D).
Vmp1 and Tmem41b mutant ESCs showed similar morphology (Fig.1B),
proliferation (Fig.S1E), and expression of pluripotency markers (Fig.S1F-G) compared
to WT ESCs. We confirmed that the pluripotency network was unchanged by a
transcriptomic analysis performed in individual Vmp1 and Tmem41b mutant cells. This
analysis also showed that the transcription profile of cells was marginally affected,
consistent with earlier results of ESC characterization (Fig.S1I-J and Table S2).

To test the effect of Vmp1 and Tmem41b mutations on autophagy, we
measured the accumulation of LC3+ autophagosomes. We found an increase of LC3+
puncta in both mutants compared to WT ESCs (Fig.1C and Fig.S2A-B). Interestingly,
the increased frequency of LC3+ puncta was similar in Vmp1 (4X) and Tmem41b
(3,5X) mutant cells. Next, mutant cells were analyzed for the retention of lipid droplets
(LD). To amplify LDs, we treated all samples with oleic acid (OA) and visualized LDs
by BODIPY staining. We observed that Vmp1 and Tme41b mutations led to an
increase of 4 times LD frequency with respect to WT cells. These results suggest that
both Vmp1 and Tmem41b contribute to autophagosome and lipid droplet metabolism
in ESCs, recapitulating earlier studies performed in other cellular systems (Li et al.,
2021; Moretti et al., 2018). The direct comparison of mutation effects revealed a similar
contribution of VMP1 and TMEM41B in both processes.

Vmp1 and Tmem41b activity is not required for ESC self-renewal
We generated double mutant cell lines to characterize potential redundant functions
between VMP1 and TMEM41B. A Tmem41b mutation was engineered in Vmp1 mutant
ESCs (Double KO) (Fig.S1A-D). Several independent mutant clones were obtained.
Notably, Double KO cells showed a regular ESC morphology (Fig.S1H) and
proliferation rate (Fig.S1E). The absence of Vmp1 and Tmem41b did not affect
secretory compartments, as ER, Golgi apparatus, and Late Endosomes appeared
normal in mutant cells (Fig.S3A). Double KO cells showed a limited perturbation in
gene expression (Fig.1E). The combined mutations did not result in a greater number
of differentially regulated genes compared to Vmp1^{KO} cells (Fig.S3C). Minimal overlap
of differentially expressed genes was detected among Vmp1, Tmem41b and Double
KO mutant cell lines (Fig.S3B-C). Additionally, we did not observe enrichment for
genes that are involved in defined cellular processes (Fig.1F, Fig.S3D-E). The
pluripotency network, as also shown by IF staining (Fig.S1F-G), was not perturbed in
Double KO cells. We conclude that Vmp1 and Tmem41b are not essential for
maintaining ESC pluripotency.

Vmp1 and Tmem41b have redundant functions
LC3 staining showed that Double KO cell lines present an exacerbated phenotype in
autophagosome accumulation compared to single mutants. In the Double KO, we did
not detect an additional increase in the number of autophagosomes per cell (Fig.1G

and Fig.S2A-B). However, 14% of LC3 clusters in the Double KO had a larger size (>3µm), compared to the 8% of autophagosomes across the single mutations and 3% in the WT condition (Fig.1H and Fig.S2C).

The simultaneous depletion of Vmp1 and Tmem41b had a greater impact on LD formation. Double KO cells showed a drastic increase of LD number in the absence of oleic acid treatment, whereas in WT and Vmp1 and Tmem41b mutant conditions, LDs were clearly observed just after OA treatment (Fig.1I-J and Fig.S2D-E). As BODIPY could also stain lipid-containing compartments like autophagosomes (O'Rourke et al., 2009), we confirmed the enriched lipid droplets in Double KO cells with refractive index tomography, a labelling-independent technique (Fig.1K and Fig.S2F-G). Together, these results indicate that Vmp1 and Tmem41b share redundant functions, as the combined mutation of both genes exacerbates autophagosome and lipid droplet accumulation.

Loss of Vmp1 and Tmem41b affects ESC potential to differentiate into primitive endoderm

To assess the effect of Vmp1 and Tmem41b on the differentiation potential of ESCs, we used embryoid body (EB) differentiation (Fig.2A). After 6 days of differentiation, EBs derived from WT ESCs showed increased expression of ectoderm (Sox1 and Fgf5), mesoderm (Actc1 and Brachyury) and endoderm (Gata6 and Dab2) genes (Fig.2B and Fig.S4A), while the expression of pluripotency markers (Nanog and Pou5f1) was lost (Fig.2B and Fig.S4A). We evaluated variations in the differentiation potential of mutant cells by comparing the expression of these markers by RT-qPCR. For each mutant, the differentiation potential of two independent clones was tested. Vmp1, Tmem41b, and double mutant ESCs formed well-defined EBs that resembled those of WT cells. All cell lines efficiently differentiated into the ectoderm and mesoderm lineages (Fig.2C and Fig.S4B-C). Notably, Double KO EBs showed a decreased expression of the endoderm markers Dab2 (100%) and Gata6 (57% of experiments) (Fig.2C and Fig.S4C). Neither single mutant showed a similar failure in Dab2 expression, even if some variability was detected between the analyzed clones. In the single mutants, we noticed a decreased endoderm specification, as shown by Gata6 expression levels in experiments performed with Vmp1[KO] (56% of experiments) and Tmem41b[KO] (58%) cells (Fig.2C and Fig.S4C). Our results indicate that combined

mutation of Vmp1 and Tmem41b leads to reduced Dab2 expression and predisposes
to variability in the expression of other endodermal genes.

We confirmed the differentiation potential of Double KO cells into ectodermal
lineages by generating SOX1-positive spinal cord organoids (Fig.S4D). Mesodermal
differentiation was inferred from the ability of double mutant EBs to form contractile
cardiomyocytes after plating (Video S1-2). We subsequently sectioned day 6 EBs to
evaluate protein expression and distribution of endoderm markers by IF. Expression
of GATA4, GATA6, and SOX17 was detected but noticeably decreased in Double KO
compared to WT EBs (Fig.2D-E). Decreased GATA4 and GATA6 staining was
pronounced in EB border regions, which resemble the primitive endoderm layer
(Zeevaert et al., 2020). In line with these findings, we confirmed the absence of DAB2
in Double KO EBs (Fig.2F). Dab2 is mainly expressed in the primitive endoderm, while
the other markers are known to be expressed in both primitive and definitive
endoderm.

The significant reduction in both mRNA and protein levels of Dab2, along with
the weakened expression of endoderm markers at the edges of embryoid bodies,
provides evidence that the absence of Vmp1 and Tmem41b compromises the
formation of primitive endoderm.

Combined Vmp1 and Tmem41b mutations delay ESC differentiation into extra-
embryonic endoderm stem (XEN) cells
We tested the ability of Double KO ESCs to differentiate into extra-embryonic
endoderm stem (XEN) cells to characterize endodermal differentiation further. XEN
cells are a useful model of PE cells, with which they share signaling programs, gene
expression, and differentiation potential (Kruithof-de Julio et al., 2011). Double KO and
WT cells were differentiated into XEN cells with Activin A and Retinoic Acid (Niakan et
al., 2013). After 6 days, WT ESCs efficiently differentiated into XEN clusters (Fig.3A),
exhibiting a stellate and refractile morphology (Niakan et al., 2013). In Double KO cells,
the formation of XEN clusters was substantially reduced. We characterized XEN
identity by analyzing the expression of XEN markers by RT-qPCR (Fig.S5A) and
immunostaining (Fig.3B and Fig.S5B). It is worth noting that the small number of
derived XEN cells from Double KO ESCs can be expanded for multiple passages, as
shown in Fig.S5C. This result suggests that the loss of Vmp1 and Tmem41b does not

entirely prevent XEN specification, but rather causes a delay in the differentiation process.

WNT signaling defect in Vmp1 and Tmem41b mutant cells impair XEN specification

To characterize the effect of Vmp1 and Tmem41b double mutation on XEN specification, we analyzed the transcriptomic profile of Double KO and WT cells during the differentiation process. On day two of differentiation, we detected the induction of XEN-associated genes (Ngondo et al., 2018) in WT cells. These genes, however, were only poorly expressed in Double KO cells (Fig.3C-D and Fig.5E). Intriguingly, we did not observe any further perturbation in the transcription of genes involved in lipid metabolism, autophagy, or ER maintenance (Fig.S5D).

Instead, Double KO cells show specific deregulation of genes related to the WNT pathway (Fig.3E-F). A close analysis of these genes allowed to identify several well-established WNT-target genes, including Frzb, Sox17, Wnt3, and Ccdnd1 (Du et al., 2009; Tetsu and McCormick, 1999; Zhang et al., 2009). Their down-regulation strongly suggested a decreased activity of the WNT pathway in Double KO cells and might explain the observed defect in XEN specification (Athanasouli et al., 2023). Interestingly, Ddit3 was strongly upregulated in Double KO cells. DDIT3 acts as an inhibitor of the WNT signaling by repressing the transcriptional activity of the TCF7/TCF4 complex (Chiu et al., 2019). We confirmed increased levels of the DDIT3 protein in Double KO cells (Fig.4A) and quantified a fourfold increase over WT cells (Fig.4B).

Finally, we attempted to rescue the defect in XEN differentiation of double mutant cells by chemically activating WNT signaling. We employed the GSK3β kinase inhibitor (CHIR), as a WNT agonist. GSK3β inhibition did not affect XEN specification in WT ESCs. However, WNT activation significantly enhanced XEN cell cluster formation in double mutant ESCs (Fig. 3D-F and Fig. S5B). Collectively, our data indicate how Vmp1 and Tmem41b affect WNT signaling and explain the delay in primitive endoderm specification.

DISCUSSION

Vmp1 and Tmem41b have been recently identified in genetic screens designed to study various cellular processes, like autophagy, lipid metabolism, virus maturation, pathway modulation, and ER stress. The pleiotropic function and peculiar structure of the DedA domain have prompted multiple studies addressing its mode of action and role in cellular homeostasis. An important point under evaluation is whether Vmp1 and Tmem41b work cooperatively or show specialization for different cellular processes. To address this question, we targeted Vmp1 and Tmem41b genes in ESCs and introduced mutations abrogating their expression. This allowed us to directly compare the specificity of Vmp1 and Tmem41b in multiple processes. We report that their absence similarly affects autophagy and lipid droplet accumulation. Moreover, we show that the simultaneous loss of both genes exacerbates defects of single mutants, particularly in LD formation. Our findings unveil a partial redundancy between Vmp1 and Tmem41b activity. It will be interesting to understand if variating expression levels of Vmp1 and Tmem41b can account for their redundancy or dominance in different cellular systems. This is especially relevant since Vmp1 expression is known to be influenced by cell signaling (Lo Re et al., 2012; Wang et al., 2020).

The embryonic lethality caused by Vmp1 and Tmem41b mutations (Morishita et al., 2019; Van Alstyne et al., 2018) during development may be explained by the defective primitive endoderm specification that we detected *in vitro*. Embryoid body experiments and the direct differentiation of ESCs to extra-embryonic endoderm stem (XEN) cells support a specific delay in PE formation in double mutant cells. Interestingly, it has been reported that other mutations leading to PE delay cause retardation and embryo lethality at E8.5 (Futterer et al., 2012), mirroring phenotypes of Vmp1 or Tmem41b mutant embryos. While both scramblases are needed in mice for regular development, the depletion of both genes is required to cause a differentiation delay in ESCs. More permissive *in vitro* growth conditions may explain the requirement for the depletion of both genes to manifest this phenotype in cell culture. It is worth noting that single Vmp1[KO] or Tmem41b[KO] cells also showed subtle defects in endoderm specification in embryoid bodies, albeit less consistently than Double KO cells. (Fig.2C). The finding of WNT signaling deregulation in Vmp1 and Tmem41b XEN cells explains the specification delay detected in mutant cells: WNT/TCF7L1 transcriptional response is necessary *in vitro* and *in vivo* to promote PE expansion (Athanasouli et al., 2023). Retardation – not impairment – of PE

specification induced by Vmp1 or Tmem41b mutations may permit embryo implantation but the accumulation of defects likely blocks development during gastrulation.

It is unlikely that issues in autophagy or lipid/lipoprotein metabolism alone cause embryo arrest. Both KO embryos die earlier than day E8.5 with severe developmental delay, which is already evident at E7.5. Mutant mice with autophagy or LD metabolism defects reach the pup stage (Kuma et al., 2017) or die at E10.5 after gastrulation and during organogenesis (Farese et al., 1995; Raabe et al., 1998), respectively. We propose that PE defects in Vmp1/Tmem41b depleted cells are caused by the increased expression of the WNT repressor DDIT3 which we show at the transcriptional and protein level in KO cells (Fig.4F). DDIT3 can be activated via the PERK/ATF4 pathway by ER stress (Rozpedek et al., 2016) or calcium-dependent mechanisms upon Vmp1 depletion in the absence of ER stress (Li et al., 2023). Notably, downstream activation of the WNT cascade fully rescues the defects of Vmp1 and Tmem41b double mutant cells during XEN specification.

This work presents novel evidence illustrating the roles of Vmp1 and Tmem41b in acting at the interface of cell homeostasis and signaling. Applying the generated cellular systems, future studies will help clarify redundant mechanisms shared by these scramblases. Furthermore, the role of Vmp1 and Tmem41b in modulating the WNT cascade may extend to other signaling networks, revealing other cross-connections between cellular pathways and metabolic processes.

MATERIALS AND METHODS

Cell Culture

Mouse ES cells were maintained as previously described in chemically defined 2i medium with added leukemia inhibitory factor (LIF) as described in Di Minin et al. (Di Minin et al., 2022). Base media composition: DMEM (Gibco, #41965039), 15% FBS (Biowest, #S1810-500) 1% sodium pyruvate (Gibco, 11360-070), 1% non-essential amino acids (Gibco, #11140-035). Lif (2000 U/mL, homemade), PD0325901 (1 µM, AxonMedchem, #391210-10-9) and Chiron (3 µM, AxonMedchem, #252917-06-9) are added fresh before use.

EB differentiation

To obtain embryonic bodies (EBs), 200k ESCs were plated on Day1 in ESCs base media without 2i and LIF (see cell culture) into a well of a Sphericalplate5D (Kugelmeiers). On Day4, EBs have formed and are transferred to a pHEMA coated 6cm culture dish and cultured until Day6. EBs are then collected and subjected to RNA extraction and RT-qPCR analysis or prepared for sectioning and immunofluorescence. In brief, EBs are fixed in 4% PFA for 30 min, cryoprotected in 10%, 20% and 30% sucrose and then embedded and frozen in Tissue Freezing Medium (Leica, #14020108926). 15 µm sections were prepared on a cryotome, mounted onto Superfrost Plus Adhesion slides (VWR international GmBH, #631-0108) and processed for immunofluorescence staining.

Cardiomyocyte differentiation

EBs were formed as previously described. On Day4, EBs were collected and seeded onto a non-treated 10cm dish, maintaining ESC base media. Cardiomyocytes randomly specified and were observed as early as on Day6 of differentiation.

Spinal Cord Organoids (SCOs) formation

For derivation of SCOs, ESCs were plated in EBs media (DMEM- F12 (Gibco) and Neurobasal Medium (Gibco) (1:1 ratio) supplemented with 200 mM L-glutamine and 10% of Knockout serum replacement) on Sphericalplate 5D dishes (Kugelmeiers AG). On day 2, SCOs were transferred in 10 cm tissue culture dishes and treated with RA (10 nM). SCOs were collected and processed for analysis at day 5.

Xen Differentiation

XEN cells were obtained following a protocol by Niakan et al. with minor changes (Niakan et al., 2013). In brief, 230k cells were plated in XEN medium on a gelatin-coated 3cm dish. The next day, the media was replaced with Xen medium containing 0.1 µM retinoic acid (Thermo Fischer, #17110052) and 10 ng/mL Activin A (Peprotech, #120-14). On Day4, the medium was again replaced by XEN medium, no further supplements were added, and on Day6 samples were either analyzed by IF or qPCR. Rescue of double KO Xen differentiation was performed by adding 1 µM Chiron (AxonMedchem, #252917-06-9) at Day2, together with the retinoic acid and Activin A. Media was changed on day 4 to Xen medium. XEN medium composition: RPMI 1640

(Gibco, #61870010), GlutaMAX^{TM} (Gibco, #61870036), 15% FBS (Biowest, #S1810-

500), 0,1 mM ß-mercaptoethanol (Sigma, #BCCB9882).

CRISPR-Cas9 editing to generate mutant ESC lines

Vmp1: the gRNA (gagacgcatagcaatgagta) was cloned into a px330 vector (Addgene,

#158973). Per 3cm well, 300k ESCs were plated and transfected with 2.5 µg of gRNA

containing plasmid, as well as 0.25 µg of a GFP expressing plasmid (pCMV-GFP).

GFP-positive cells were enriched by FACS and plated as single cells. Colonies were

picked. Analysis by WB was used to identify identified VMP1 KO lines. Tmem41b: the

gRNAs (gtatgtttgacctgggcgaa and gagtgacatgtggaaatcag) were cloned into px330

vectors, and KO clones were derived as described above. PCR analysis of the edited

locus (FW: atcttggacaggggggagttc RV: gaccaggggttcattgtcatt and FW:

cagcacacacctttaatccagc RV: gccacatagcaagcttaagagc) identified KO clones. qPCR of

mRNA from Exon 2 (edited locus, N-terminal end) and Exon 7 (C-terminal end) further

confirmed the absence of mRNA.

Proliferation assay

100k ESCs were plated in a well of a MW12. 48h later, cells were collected and

counted. Proliferation rates were calculated as relative factor of cells collected over

cells seeded.

LC3 staining and size characterization

100k cells were plated on cover-slips, pre-treated with Matrigel (Corning, # 354277),

in N2B27 media. N2B27 composition: 50 mL Neurobasal media (Gibco, # 2348949),

50 mL Advanced DMEM F12 (Gibco, # 12634010), 4% BSA (Gibco, # 15260), 1% L-

Glutamine (Gibco, # 25030024), 1% B27 supplement (Gibco, # 2336837), 0,5% N2

supplement (Gibco, # 2328253), and 0,1 mM ß-mercapotethanol (Sigma,

#BCCB9882). The next day, samples were fixed in 4% PFA for 15 min, washed 3 times

with PBS and then treated with ice-cold MeOH for 5 min on RT. The staining was then

performed as described below. Quantification was performed with the Fiji plugin

TrackMate (Ershov et al., 2022; Tinevez et al., 2017) to automatically detect puncti

with a diameter of 1, 2, and 3 microns at a threshold of 600, 200 and 150, respectively.

Nuclei were detected at an average size of 15 microns with a threshold of 10. Unique

particles were annotated for 5 different frames for each condition and shown as particles per cell. The experiment was repeated 3 times.

LD staining and RF microscopy

Samples were plated as for the LC3 staining. For samples treated with oleic acid, 200 µM (Sigma-Aldrich, # O1008-1G) were added at plating, over-night. The next day, samples were incubated in 5 µM BODIPY 493/503 (Invitrogen, #D3922) in PBS and subsequently washed 3 times with PBS and fixed in 4 % PFA for 15 min. Samples were then washed, stained with Dapi for 15 min at RT and mounted with Mowiol. Quantification was performed using the Fiji plugin TrackMate (Ershov et al., 2022; Tinevez et al., 2017) detecting droplets of 1, 2, and 3 microns in diameter at a threshold of 900. Nuclei were detected at 10 micron and a threshold of 1.

For RF microscopy, 250k cells were plated per well of a 6 well glass bottom plate (Cellvis, # P06-1.5H-N) in N2B27 media. After 36 h samples were treated with 200 µM oleic acid o/n and then with BODIPY for 3 h. Samples were washed 3 times with PBS and maintained in N2B27 for RF microscopy. The imaging was performed on Tomocube HT-X1 instrument at 37C and 5 % CO_2. At least 5-8 images were acquired for each condition for each well. The z-stacks were acquired with 1.06 µm slices. Max Intensity projections (MIP) are generated on the Tomostudio 3.3.9. The fluorescence max intensity projections were resized to Refractive index tomograms for comparison. The ROIs from each image were isolated for further analysis. Comdet plugin was used for identifying high refractive index regions. The approximate particle size was 2 pixels and Intensity threshold values were chosen individually for each image to avoid any false positives or negatives.

Immunofluorescence

Samples were washed 3 times with PBS, fixed in 4% PFA for 10 min and again washed 3 times with PBS. Samples stained for LC3 were then treated for 5 min with 100% MeOH at RT and washed again 3 times with PBS. Samples were then blocked and permeabilized with 10% donkey serum (Jackson Immunoresearch, #017-000-121) and 0,1% Triton X (Thermo Fischer, # HFH10) in PBS for 20 min and incubated with primary antibodies at 4 °C over-night. Oct4 (1:100, Satna Cruz, # sc-5279), Nanog (1:100, Abcam, # ab80892), Sox2 (1:50, R&D Systems #AF2018), Lc3 (1:200, Cell Signalling, # 2775S), Pdi (1:200, Cell Signalling, # 3501S), Rcas1 (1:200, Cell

Signalling, # 12290S), Rab7 (1:200, Cell Signaling, # 9367S) Sox1 (1:50, R&D Systems, # AF3369), Sox17 (1:200 R&D Systems, # AF1924-SP), Gata4 (1:100 R&D Systems, # AF1700-SP), Gata6 (1:100 R&D Systems, # AF1700-SP), Dab2 (1:100 BD, # 610464). The next day, samples were washed 3 times with 0.1% Triton X in PBS and then incubated with secondary antibodies and Dapi.

Western Blot Analysis

Cells were lysed in membrane buffer (150 mM NaCl, 50 mM Tris-HCl pH 7,5, 1 mM 2-mercaptoethanol (Sigma, #M3148) and 1% CHAPS (Sigma, #C3023) in water). Lysates were run on SDS-Page and transferred to nitrocellulose membranes. Bands were detected with the Amersham ECL Prime Western Blotting Detection Reagents (#RPN2232) and captured on Fujifilm Super RX-N films (#47410). Antibodies for Vmp1 (Cell Signalling, #12929S) were used at 1:1000 and DDIT3 1:1000 (Cell Signalling, #2895T). Quantification of western blot bands was performed with ImageJ, normalized to HSP90 (Santa Cruz, #sc-101494).

RNA extraction and RT qPCR

RNA was extracted with the RNeasy Kit (Quiagen, #74104) following the manufacturer's instructions, including the on-column DNA digestion. EB and SCO samples were pre-treated with the QIAshredder (Quiagen, #79656). cDNAs were prepared using the Bio-Rad iScript Reverse Transcription Supermix for RT-qPCR (#1708841). RT qPCR samples were prepared using the KAPPA SYBR R fast (Roche, KK4611). Primer Sequences:

Tmem41b Exon2 FW: TGCAGAAGCTGGATCAGCAA, RV:
GAGCCTTGGCGTCATCCATA

Tmem41b Exon7 FW: GCAGGTAGAGCGTCACAGAG, RV:
TGCAGAGTTGTTCCTGCCTT

Sdha FW: CGTTATGTGAGGGGTGTGCCTT, RV:
CCTTACCAAACCTTGTGTCTGGA

Sox1 FW: ACGTAGCCAACGTAGACAC, RV: ATGAGCGTCGTCCCGTGG

Fgf5 FW: GCCTGTCCTTGCTCTTCCTCAT, RV: GGAGAAGCTGCGACTGGTGA

Actc1 FW: TGTAGGACCGTGTCGAAACC, RV: CGAAACCACACACTGTTACCG

Brachyury FW: TCTCTGGTCTGTGAGCAATGGT, RV:
TGCGTCAGTGGTGTGTAATGTG

Gata6 FW: GAAGCGCGTGCCTTCATC, RV: GTAGTGGTTGTGGTGTGACAGTTG

Dab2 FW: TTGATGATGTGCCTGATGCT, RV: TTTGCTTGTGTTGTCCCTGA

Sox17 FW: CCCAACACTCCTCCCAAAGTATC, RV:

TTCCCTGTCTTGGTTGATTTCT

Quantification and statistical analysis

Statistical analysis was performed using Python and GraphPad Prism. Data are

presented as mean-centered and the standard deviation. All experiments were

repeated with at least 3 independent biological replicates, unless stated otherwise.

The statistical test used to evaluate significance is the Welch's *t* test. Statistical

significance in the figures is denoted as follows: ns: *p*> 0.05, *: *p* < 0.05, **: *p* < 0.01,

***: *p* < 0.001.

RNA seq

RNA samples were submitted to the Functional Genomics Center in Zurich (FGCZ) for

RNA sequencing. The quality of the RNA was determined with a Fragment Analyzer

standard sensitivity RNA measurement (SS RNA kit (15 nt), Agilent, Waldbronn,

Germany). The measured concentrations (> 50 ng/µl) and RIN (>9.9) values qualified

for a Poly-A enrichment strategy in order to generate the sequencing libraries applying

the TruSeq mRNA Stranded Library Prep Kit (Illumina, Inc, California, USA). After Poly-

A selection using Oligo-dT beads the mRNA was reverse-transcribed into cDNA. The

cDNA was fragmented, end-repaired and polyadenylated before ligation of TruSeq UD

Indices (IDT, Coralville, Iowa, USA). The quality and quantity of the amplified

sequencing libraries were validated using a Fragment Analyzer SS NGS Fragment Kit

(1–6000 bp) (Agilent, Waldbronn, Germany). The equimolar pool of samples was

spiked into a NovaSeq6000 run targeting 20M reads per sample on a S1 FlowCell

(Novaseq S1 Reagent Kit, 100 cycles, Illumina, Inc, California, USA). Reads from

RNA-seq were first preprocessed by Trimmomatic (v0.40) (Bolger et al., 2014) to

remove low-quality bases and adapters. Then, reads were aligned to mouse genome

GRCm38 (release 81 version from Ensembl) using HISAT2 (v2.2.1) (Kim et al., 2019).

HTSeq Count (v2.0.2) (Putri et al., 2022) was used to count reads for each gene

(Ensembl GRCm38.81), ignoring reads on overlapped regions. Differentially

expressed genes were defined by DESeq2 (v1.40.2) (Love et al., 2014). MA plot was

generated using custom-made scripts. Genes significantly enriched show a fold

change > 1 and FDR (adjusted p-value) < 0.01. The read counts were transformed in
RPKM for the heatmap representations of specific genes and pathways. The genes
corresponding to germ-layer differentiation markers (including those for embryonic and
extra-embryonic endoderm), biological processes, and signaling pathways were
obtained through literature mining and the KEGG database (Table S1). The processed
data are available as excel files (Table S2 and Table S3) and raw data have been
deposited on NCBI SRA database under the accession: PRJNA1031567
485

486 **Competing interest**
487 The authors declare no competing interests.
488

Acknowledgments
We acknowledge the Functional Genomics Center Zurich (FGCZ) for technical support
and Sungsik Lee from ScopeM facility for help on the RF microscopy. We thank G.J.
Pereira, C. Fimiani, and U. Sutter for their help with reagents. We thank C. Ciaudo, A.
Grison, J. Corn, and K. Basler for their helpful discussions. GDM was supported by
the ETH Zurich Postdoctoral Fellowship Program as well as the Marie Curie Actions
for People COFUND Program. This work was supported by grants from the Swiss
National Science Foundation (SNF grants 31003A_152814/1 and 31003A_175643/1).
497

Author Contributions
Conceptualization: MH, AW and GDM; Data curation: MH, VG and GDM; Formal
analysis: MH, TS, HH, VG, and GDM; Funding acquisition: KW, AW and GDM;
Investigation: MH, AW and GDM; Methodology: MH, TS, HH, VG, and GDM; Project
administration: AW, GDM; Resources: MH, KW, AW and GDM; Supervision: KW, AW
and GDM; Validation: MH, GDM; Visualization: MH, VG and GDM; Writing – original
draft: MH, AW and GDM; Writing – review & editing: MH, TS, HH, VG, KW, AW and
GDM.

REFERENCES

Arnold, S.J., and Robertson, E.J. (2009). Making a commitment: cell lineage allocation and axis patterning in the early mouse embryo. Nat Rev Mol Cell Biol *10*, 91-103.

Athanasouli, P., Balli, M., De Jaime-Soguero, A., Boel, A., Papanikolaou, S., van der Veer, B.K., Janiszewski, A., Vanhessche, T., Francis, A., El Laithy, Y.*, et al.* (2023). The Wnt/TCF7L1 transcriptional repressor axis drives primitive endoderm formation by antagonizing naive and formative pluripotency. Nat Commun *14*, 1210.

Bard, F., Casano, L., Mallabiabarrena, A., Wallace, E., Saito, K., Kitayama, H., Guizzunti, G., Hu, Y., Wendler, F., Dasgupta, R.*, et al.* (2006). Functional genomics reveals genes involved in protein secretion and Golgi organization. Nature *439*, 604-607.

Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics *30*, 2114-2120.

Chiu, C.S., Tsai, C.H., Hsieh, M.S., Tsai, S.C., Jan, Y.J., Lin, W.Y., Lai, D.W., Wu, S.M., Hsing, H.Y., Arbiser, J.L.*, et al.* (2019). Exploiting Honokiol-induced ER stress CHOP activation inhibits the growth and metastasis of melanoma by suppressing the MITF and beta-catenin pathways. Cancer Lett *442*, 113-125.

Di Minin, G., Holzner, M., Grison, A., Dumeau, C.E., Chan, W., Monfort, A., Jerome-Majewska, L.A., Roelink, H., and Wutz, A. (2022). TMED2 binding restricts SMO to the ER and Golgi compartments. PLoS Biol *20*, e3001596.

Du, Y.C., Oshima, H., Oguma, K., Kitamura, T., Itadani, H., Fujimura, T., Piao, Y.S., Yoshimoto, T., Minamoto, T., Kotani, H.*, et al.* (2009). Induction and down-regulation of Sox17 and its possible roles during the course of gastrointestinal tumorigenesis. Gastroenterology *137*, 1346-1357.

Ershov, D., Phan, M.S., Pylvanainen, J.W., Rigaud, S.U., Le Blanc, L., Charles-Orszag, A., Conway, J.R.W., Laine, R.F., Roy, N.H., Bonazzi, D.*, et al.* (2022). TrackMate 7: integrating state-of-the-art segmentation algorithms into tracking pipelines. Nat Methods *19*, 829-832.

Farese, R.V., Jr., Ruland, S.L., Flynn, L.M., Stokowski, R.P., and Young, S.G. (1995). Knockout of the mouse apolipoprotein B gene results in embryonic lethality in homozygotes and protection against diet-induced hypercholesterolemia in heterozygotes. Proc Natl Acad Sci U S A *92*, 1774-1778.

Futterer, A., Raya, A., Llorente, M., Izpisua-Belmonte, J.C., de la Pompa, J.L., Klatt, P., and Martinez, A.C. (2012). Ablation of Dido3 compromises lineage commitment of stem cells in vitro and during early embryonic development. Cell Death Differ *19*, 132-143.

Ghanbarpour, A., Valverde, D.P., Melia, T.J., and Reinisch, K.M. (2021). A model for a partnership of lipid transfer proteins and scramblases in membrane expansion and organelle biogenesis. Proc Natl Acad Sci U S A *118*.

Hama, Y., Morishita, H., and Mizushima, N. (2022). Regulation of ER-derived membrane dynamics by the DedA domain-containing proteins VMP1 and TMEM41B. EMBO Rep *23*, e53894.

Hoffmann, H.H., Schneider, W.M., Rozen-Gagnon, K., Miles, L.A., Schuster, F., Razooky, B., Jacobson, E., Wu, X., Yi, S., Rudin, C.M.*, et al.* (2021). TMEM41B Is a Pan-flavivirus Host Factor. Cell *184*, 133-148 e120.

Huang, D., Xu, B., Liu, L., Wu, L., Zhu, Y., Ghanbarpour, A., Wang, Y., Chen, F.J., Lyu, J., Hu, Y.*, et al.* (2021). TMEM41B acts as an ER scramblase required for lipoprotein biogenesis and lipid homeostasis. Cell Metab *33*, 1655-1670 e1658.

Kim, D., Paggi, J.M., Park, C., Bennett, C., and Salzberg, S.L. (2019). Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol *37*, 907-915.

Kruithof-de Julio, M., Alvarez, M.J., Galli, A., Chu, J., Price, S.M., Califano, A., and Shen, M.M.
(2011). Regulation of extra-embryonic endoderm stem cell differentiation by Nodal and Cripto
signaling. Development *138*, 3885-3895.

Kuma, A., Komatsu, M., and Mizushima, N. (2017). Autophagy-monitoring and autophagy-
deficient mice. Autophagy *13*, 1619-1628.

Li, T., Zhao, H., Guo, G., Xia, S., and Wang, L. (2023). VMP1 affects endoplasmic reticulum stress
sensitivity via differential modulation of the three unfolded protein response arms. Cell Rep
42, 112209.

Li, Y.E., Wang, Y., Du, X., Zhang, T., Mak, H.Y., Hancock, S.E., McEwen, H., Pandzic, E., Whan,
R.M., Aw, Y.C.*, et al.* (2021). TMEM41B and VMP1 are scramblases and regulate the
distribution of cholesterol and phosphatidylserine. J Cell Biol *220*.

Lo Re, A.E., Fernandez-Barrena, M.G., Almada, L.L., Mills, L.D., Elsawa, S.F., Lund, G., Ropolo,
A., Molejon, M.I., Vaccaro, M.I., and Fernandez-Zapico, M.E. (2012). Novel AKT1-GLI3-VMP1
pathway mediates KRAS oncogene-induced autophagy in cancer cells. J Biol Chem *287*, 25325-
25334.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550.

Molejon, M.I., Ropolo, A., Re, A.L., Boggio, V., and Vaccaro, M.I. (2013). The VMP1-Beclin 1
interaction regulates autophagy induction. Sci Rep *3*, 1055.

Moretti, F., Bergman, P., Dodgson, S., Marcellin, D., Claerr, I., Goodwin, J.M., DeJesus, R., Kang,
Z., Antczak, C., Begue, D.*, et al.* (2018). TMEM41B is a novel regulator of autophagy and lipid
mobilization. EMBO Rep *19*.

Morishita, H., Zhao, Y.G., Tamura, N., Nishimura, T., Kanda, Y., Sakamaki, Y., Okazaki, M., Li, D.,
and Mizushima, N. (2019). A critical role of VMP1 in lipoprotein secretion. Elife *8*.

Ngondo, R.P., Cirera-Salinas, D., Yu, J., Wischnewski, H., Bodak, M., Vandormael-Pournin, S.,
Geiselmann, A., Wettstein, R., Luitz, J., Cohen-Tannoudji, M.*, et al.* (2018). Argonaute 2 Is
Required for Extra-embryonic Endoderm Differentiation of Mouse Embryonic Stem Cells. Stem
Cell Reports *10*, 461-476.

Niakan, K.K., Schrode, N., Cho, L.T., and Hadjantonakis, A.K. (2013). Derivation of
extraembryonic endoderm stem (XEN) cells from mouse embryos and embryonic stem cells.
Nat Protoc *8*, 1028-1041.

O'Rourke, E.J., Soukas, A.A., Carr, C.E., and Ruvkun, G. (2009). C. elegans major fats are stored
in vesicles distinct from lysosome-related organelles. Cell Metab *10*, 430-435.

Putri, G.H., Anders, S., Pyl, P.T., Pimanda, J.E., and Zanini, F. (2022). Analysing high-throughput
sequencing data in Python with HTSeq 2.0. Bioinformatics *38*, 2943-2945.

Raabe, M., Flynn, L.M., Zlot, C.H., Wong, J.S., Veniant, M.M., Hamilton, R.L., and Young, S.G.
(1998). Knockout of the abetalipoproteinemia gene in mice: reduced lipoprotein secretion in
heterozygotes and embryonic lethality in homozygotes. Proc Natl Acad Sci U S A *95*, 8686-
8691.

Rozpedek, W., Pytel, D., Mucha, B., Leszczynska, H., Diehl, J.A., and Majsterek, I. (2016). The
Role of the PERK/eIF2alpha/ATF4/CHOP Signaling Pathway in Tumor Progression During
Endoplasmic Reticulum Stress. Curr Mol Med *16*, 533-544.

Tetsu, O., and McCormick, F. (1999). Beta-catenin regulates expression of cyclin D1 in colon
carcinoma cells. Nature *398*, 422-426.

Tinevez, J.Y., Perry, N., Schindelin, J., Hoopes, G.M., Reynolds, G.D., Laplantine, E., Bednarek,
S.Y., Shorte, S.L., and Eliceiri, K.W. (2017). TrackMate: An open and extensible platform for
single-particle tracking. Methods *115*, 80-90.

Tornero-Ecija, A., Tabara, L.C., Bueno-Arribas, M., Anton-Esteban, L., Navarro-Gomez, C.,
Sanchez, I., Vincent, O., and Escalante, R. (2022). A Dictyostelium model for BPAN disease
reveals a functional relationship between the WDR45/WIPI4 homolog Wdr45l and Vmp1 in
the regulation of autophagy-associated PtdIns3P and ER stress. Autophagy *18*, 661-677.

Van Alstyne, M., Lotti, F., Dal Mas, A., Area-Gomez, E., and Pellizzoni, L. (2018).
Stasimon/Tmem41b localizes to mitochondria-associated ER membranes and is essential for
mouse embryonic development. Biochem Biophys Res Commun *506*, 463-470.

Wang, C., Peng, R., Zeng, M., Zhang, Z., Liu, S., Jiang, D., Lu, Y., and Zou, F. (2020). An
autoregulatory feedback loop of miR-21/VMP1 is responsible for the abnormal expression of
miR-21 in colorectal cancer cells. Cell Death Dis *11*, 1067.

Zeevaert, K., Elsafi Mabrouk, M.H., Wagner, W., and Goetzke, R. (2020). Cell Mechanics in
Embryoid Bodies. Cells *9*.

Zhang, Z., Deb, A., Zhang, Z., Pachori, A., He, W., Guo, J., Pratt, R., and Dzau, V.J. (2009).
Secreted frizzled related protein 2 protects cells from apoptosis by blocking the effect of
canonical Wnt3a. J Mol Cell Cardiol *46*, 370-377.

    FIGURE LEGENDS

    Figure 1. Vmp1 and Tmem41b double KO ESCs display increased
    autophagosome and lipid droplet accumulation

    (A) Schematic representation of the CRISPR/Cas9-mediated KO strategy for Vmp1
    and Tmem41b genes. For depleting Vmp1 (top) a single gRNA inducing an indel
    mutation has been used. In the case of the Tmem41b gene (bottom), two gRNAs were
    used to delete exons 2 and 3. The sequences coding for transmembrane domains
    (TM) are indicated in blue, the conserved DedA domain is highlighted in yellow, red
    lines indicate cut sites of gRNAs. **(B)** Colony morphology of VMP1^{KO} and Tmem41^{KO}
    ESCs remains unchanged compared to WT ESCs. Scale bar: 100 µm. **(C)** VMP1 and
    Tmem41b depletion induce accumulation of autophagosomes in ESCs. The graph
    shows the quantification of LC3 puncta per cell of a representative experiment (see
    Fig.S2A). Data were collected from more than 200 cells. For each KO condition, two
    independent clones have been analyzed. Dots correspond to the frames analyzed per
    sample. Data are represented as mean ± SD. The P-value is relative to the WT sample.
    (D) Vmp1 and Tmem41b depletion leads to the accumulation of lipid droplets (LD) in
    ESCs. Graph quantifies LD per cell in WT and KO ESCs (see Fig.S2D). ± oleic acid
    treatment. At least 500 cells were analyzed for every condition; dots represent
    independent biological replicates. Data are represented as mean ± SD. **(E-G)**
    Transcriptional analysis of Double KO ESCs did not identify many deregulated genes.
    (E) Shown are differentially regulated genes in Double KO ESCS in blue as MA plot.
    Genes highlighted in red are pluripotency markers. **(F)** Differentially regulated genes
    do not cluster in essential processes. The graph shows differentially expressed genes
    of single and double KO ESCs by cellular processes. Dot size correlates to the number
    of deregulated genes within a cellular process. **(G-H)** Double KO ESCs accumulate
    LC3+ autophagosomes of increased size. **(G)** Shown are representative IF images of
    WT and double KO ESCs stained for LC3. Nuclei are stained with Dapi. Scale bar:
    20 µm. **(H)** Plotted is the quantification of LC3 puncta of 3 µm in diameter. For each
    condition, two independent clones have been analyzed. Dots correspond to the frames
    analyzed per sample. Data were collected from more than 200 cells. Data are
    represented as mean ± SD. The P-value is relative to the Double KO sample with the
    highest SD. **(I-J)** Double KO ESCs accumulate LDs without oleic acid treatment. **(I)**

Live cell staining of WT, Vmp1, Tmem41b, and double KO ESCs with BODIPY. Scale bar: 20 μm **(J)** The number of LDs per cell is plotted for all KO clones. Data are represented as mean ± SD. **(K)** Double KO ESCs present an increased amount of LD visualized through refractive index microscopy (top) and co-stained with BODIPY.

Figure 2. Double KO ESCs specify all germ layers, but not the primitive endoderm

(A) Schematic representation of an Embryonic Body (EB) which consists of 3 different lineages. The ectoderm expresses Fgf5 and Sox1 (grey), the mesoderm Actc1 and Brachyury (blue), and the endoderm by Gata6. The surrounding region of the EB corresponds to the primitive endoderm (green) and is marked by Gata6 and Dab2 expression. **(B)** ESCs differentiated to EBs express lineage markers for ectoderm (Fgf5), mesoderm (Brachyury), endoderm (Gata6), and primitive endoderm (Dab2). Graphs show mRNA levels of WT EBs normalized to the Sdha gene. Dots represent biological independent experiments. Data are represented as mean ± SD. **(C)** Double KO ESCs specifically show defects in expressing the primitive endoderm marker, Dab2. qPCR analysis for Sox1, Actc1, and Dab2 mRNA expression in mutant EBs relative to WT EBs. Two independent clones were analyzed per cell line. Each dot represents an independent experiment. Data are represented as mean ± SD**.** The line at 0,7 indicates an arbitrary threshold after which successful expression of the marker is considered. **(D-F)** Double mutant EBs show decreased expression of endoderm markers. Immunofluorescence images of EB sections stained for the endoderm markers GATA4 and SOX17 **(D)**, GATA6 **(E)**, DAB2 **(F)**. WT in the top panel, two independent clones for Double KO condition in the center and bottom panels. Nuclei are stained with Dapi. Scale bar: 50 μm.

Figure 3. XEN differentiation in Double KO cells is delayed

(A-B) Double KO ESCs show a delay in XEN differentiation. **(A)** Brightfield images of XEN cells at day 6 of differentiation in WT and Double KO cells. XEN islands are circled in white in the Double KO condition. Scale bar: 100 μm. **(B)** Immunofluorescence images of day 6 XEN cells stained for SOX17 and GATA4. Nuclei are stained with Dapi, Scale bar: 150 μm. **(C-F)** Transcriptional analysis of XEN cells. **(C)** Differentially

expressed genes between Double KO and WT cells at day 2 during XEN specification.

XEN markers are labeled in red; differentially expressed genes in blue. **(D-E)** GO-Term

analysis of differentially expressed genes between Double KO and WT conditions in

ESCs and XEN cells in the context of developmental processes **(D)** and signaling

pathways **(E)**. Dot size corresponds to the number of differentially regulated genes.

(F) WNT pathway-related genes differentially expressed in Double KO cells. Heatmap

shows min to max expression of three biological replicates for WT and Double KO

XEN cells.

Figure 4. Activation of WNT signaling fully rescues the delay of Double KO cells

in XEN differentiation

(A-B) VMP1 and Tmem41b depletion promotes the increase of DDIT3 protein levels.

(A) Western Blot analysis of DDIT3 expression in WT and Double KO cells

differentiated to XEN cells for 2 days. HSP90 is shown as loading control. **(B)**

Quantification of increased DDIT3 protein levels in three biological replicates.

Expression relative to WT XEN cell and normalized to respective HSP90 expression.

Data are represented as mean ± SD. **(C-E)** Chiron-dependent WNT activation rescues

the delay of Double KO XEN differentiation. **(C)** Brightfield images show XEN clusters

of Double KO XEN cells (highlighted by white lines) with and w/o Chiron treatment.

Scale bar: 100 µm. IF staining for SOX17 and GATA4 **(D)** or DAB2 and GATA6 **(E)**;

nuclei are stained with Dapi. Scale Bar: 150 µm. **(F)** Schematic overview highlighting

the correlation between loss of Vmp1 and Tmem41b and decreased WNT signaling

via DDIT3.

Figure 4

A

DDIT3
HSP90

WT Double KO

B

DDIT3 Fold

***

WT Double KO

C

NT   +CHIR

D

WT   Double KO

NT

+CHIR

SOX17 GATA4 DAPI

E

Double KO

DAB2 GATA6 DAPI

F

TMEM41B
VMP1

LD ↑
Autophagy ↓

Ddit3

WNT
signaling

XEN
spec.