Automated assessments can support regional Red Lists but need to be

applied with caution: A case study from central Germany

Alexander Zizka[1]*, Indra Starke-Ottich[2], David Eichenberg[3], Dirk Bönsel[2], Georg Zizka[2,4]

[1]Biodiversity of Plants Lab, Department of Biology, Philipps-University Marburg, Karl-von-

Frisch-Straße 8, Marburg, Germany

[2]Botany and Molecular Evolution, Senckenberg Research Institute and Natural History

Museum, Frankfurt am Main, Germany

[3]Independent Researcher, Leipzig, Germany

[4]Institute for Ecology, Evolution and Diversity, Department of Biology, Goethe University

Frankfurt am Main, Germany

* Corresponding author: alexander.zizka@biologie.uni-marburg.de

Abstract

Aim. Expert-based regional Red Lists (RL) carry conservation legislation globally. Yet, they are often difficult to reproduce and their regular compilation by a dwindling number of expert assessors burdens many regional conservation authorities. Here we batch-estimate RL indicators and extinction risk for >1,100 plant species and test the potential of this automated approach to support the expert-based RL process.

Location. State of Hesse, Central Germany

Methods. First, we estimated current population status, short-term population trend, long-term population trend, and extinction risk by binning existing occurrence probabilities modelled at three time slices with cut-off values derived from RL methodology. Subsequently we compared the results with the latest version of the Hessian expert-based RL using summary statistics and selected example species.

Results. We find the assessments of extinction risk to agree in c. 60% of the cases, mostly for species not threatened with extinction. Existing mismatch was by one category in most cases, but up to 6 categories in some cases (mean: 1.6 categories). Furthermore, agreement was highest for extreme categories and very abundant species.

Main conclusions. Automated assessments were simplistic for many rare and taxonomically challenging species, but we considered them more accurate than the expert assessments for species with intermediate population size and for species of anthropogenic habitats. Furthermore, the automated assessments are particularly informative for the estimation of long-term and short-term population trends, for which experts are often left to guesstimate based on little data.

Key words

Introduction

Regional Red Lists (RL) quantify species' risk of local extinction and drive conservation action on the state and national level in many countries (Azam et al., 2016; Bennun et al., 2018; Betts et al., 2020; Jiang et al., 2020; Rodrigues et al., 2006). For instance, in Germany, the National Strategy for Biodiversity, a primary policy to counteract an accelerating national biodiversity loss (Habel et al., 2016; Jandt et al., 2022; Kamp et al., 2021), is based partly on RL.

RL follow a simple procedure: Specialist(s) with expertise in the biota of a region assess the extinction risk of species following fixed criteria, and, as a result, deliver a documented extinction risk category for each species. The criteria differ among RL, but are virtually always based on an assessment of habitat availability, range size, population size, and population trends through time or combinations thereof (IUCN Standards and Petitions Subcommittee, 2017; Schippmann, 2020). For instance, the RL of Germany classifies species into one of nine extinction risk categories based on three indicators: current populations status, long-term population trend (percent change in the last 50-150 years) and short-term population trend (10-25 years; Ludwig et al., 2006; Starke-Ottich et al., 2019).

The laborious process to estimate indicators and derive extinction risk assessment based on expert-opinion are the root for multiple challenges for RL (IUCN Standards and Petitions Subcommittee, 2017): First, taxonomic bias. Available expertise for animal and plant groups is proportional to their appeal to scientists and the general public. Therefore, charismatic mammals, amphibians, birds, and, to a lesser extent, flowering plants are better assessed than more elusive groups, such as soil biota and most invertebrates (Cowie et al., 2022). Likewise, even within individual plant and animal groups, assessments are more elaborate for well-studied genera and species. Second, low reproducibility. While expert's opinion can provide a high-quality interpretation of all available information, individual experts may judge population trends and extinction risk differently and therefore deliver different assessments based on identical data. Third, uncertainty for species with intermediate population size. Estimating population dynamics through time requires detailed data and understanding of a species' biology. Experts usually have a good understanding for extremely rare and very rare species, because those species are of particular interest to the specialist community and therefore most scrutinized. Likewise, common and very common species can be assumed to not to be threatened with extinction. In contrast, species with intermediate population size challenge expert assessments: on the one hand they are too common to be

of particular interest to specialists, but on the other hand they are rare enough, so that
unobserved population declines may put them under risk of extinction.

The challenges to RL are exacerbated by the need for regular re-assessment. RL
assessments need to be repeated every 10 years to account for changes in species status in
a changing world: new threats and the effectiveness of conservation measures need to be
quantified to ensure effective species conservation (Rondinini et al., 2014). Since RL are
limited by expert availability and often rely on volunteer contributions, many RL are (at least
partly) outdated (Cowie et al., 2022; Rondinini et al., 2014). For instance, the last available
RL for plants in several German states date to the late 1990ies and early 2000s. This effect
is likely to intensify in the future, because the number of available experts is dwindling
(Frobel & Schlumprecht, 2013; Starke-Ottich et al., 2019).

A possible remedy to the lack of person-power and data on species' population status for RL
assessments can be to use species distribution models (Cazalis et al., 2022). SDM can
estimate species geographic ranges from few geo-referenced occurrence records and
environmental variables in the absence of complete sampling for many species at once
(Franklin, 2010). SDM-based range estimates can then be used to approximate current
population status, and its change over time, which is encouraged by some RL guidelines
(Starke-Ottich et al., 2019). SDM encompass a multitude of algorithms from simple climate
envelopes to ensemble approaches combining multiple machine learning techniques. In a
simple form (Area of Habitat) SDM based analyses are widely used in global RL
assessments (IUCN Standards and Petitions Subcommittee, 2017). Despite their flexibility
many SDM may be unsuitable to estimate temporal trends in species' distribution or
population for RL due to the nature of the underlying data: If the effort to observe species
differs among time slices, as it most often does, inferred trends would be distorted. The
FRESCALO (an SDM in the broad sense) estimates the probability of a species occurrence
based on environmental similarity and species co-occurrence across neighbouring locations
(Hill, 2012), and efficiently corrects for this observation bias by harnessing common species
as benchmark to estimate sampling effort and correct for its distortion (Isaac et al., 2014); an
approach that has proven suitable for to estimate trends in plant species distribution across
Germany (Eichenberg et al., 2021).

The application of modelled species occurrence probabilities by the FRESCALO algorithm or
comparable approaches (short as SDM hereafter) for RL is particularly promising in
combination with a second class of automated data analysis techniques: automated
assessments (AA). AA standardize the interpretation of species occurrence dynamics to

batch estimate the extinction risk of many species. Similar to SDM, AA exist in varying complexity (Cazalis et al., 2022; Henry et al., 2023), from simple approaches to estimate range size indicators from species occurrence records (Bachman et al., 2011, 2020; Dauby et al., 2017) to artificial intelligence using a broad range of data (Walker et al., 2022; Zizka et al., 2021, 2022). SDM and AA may be applied individually or in combination to assist Red Listing, and, as data-driven baseline, have the potential to guide assessment effort (Cazalis et al., 2023), increase reproducibility (Cazalis et al., 2022) and reduce the time needed for assessments by an order of magnitude (Silva et al., 2022).

Despite their potential, SDM and AA have only been sporadically used for RL. Likely, the reason is that both methods add complexity to the assessment process and pose methodological hurdles for assessors (which are mostly taxonomic specialists). Furthermore, AA are under active development, so that no standard for their application exists, and their performance for RL is untested. For instance, AA have so far predominantly been developed for the global IUCN Red List (with few academic exceptions, e.g. Ralimanana et al., 2022; Schmidt et al., 2017) and their reliability on the regional scale remains questionable.

Here, we ask the question *"How well can the combination of SDM and AA reproduce expert-based estimates of current population status, long-term and short-term population trends and extinction risk on the regional scale?"*. We mimic the expert-based Red Listing process by batch-applying RL thresholds to approximations of species current population status, long-term and short-term population trend derived from modelled distributions ranges of >1,100 plant species from the state of Hesse in central Germany.

Methods

Study area

Hesse is a state in central Germany covering c. 21,100 km²; mid-elevation mountains, and the floodplains of the Rhine and Main rivers are the major habitat types. We chose Hesse as model region for three reasons:

1. A long-lasting tradition of scientific floristic exploration (since the 1600s) and of formal Red Listing of flowering plants (five version of the state's RL have been published since 1976: Buttler et al., 1996; Hemm et al., 2008; Hessische Landesanstalt für Umwelt, 1976; Kalheber et al., 1980; Starke-Ottich et al., 2019) and therefore many data are available on species populations trends and threat status.
2. The RL of Hesse is compiled for four subregions separately, increasing data availability.

148   3. Up-to-date distribution models of the Hessian flora exist in high resolution (25 km^2)
149   and at three time slices (1960-1987, 1988-1996, 1997-2017; Eichenberg et al.,
150   2021).

Automated regional Red List (ARL)

We generated an automated Red List for the Flora of Hessen (ARL) by estimating current
population situation, short-term population trend and long-term population trend ("RL
indicators" hereafter) and extinction risk for the four subregions and the entire state by
thresholding modelled species distributions.

First, we obtained modelled occurrence probabilities of individual plant species in a 5x5 km^2
grid across Germany at three time slices: 1960-1987, 1988-1996, 1997-2017 from
Eichenberg et al. (2021). The modelled occurrence probabilities were based on 29 million
occurrence records (from the whole of Germany) from a variety of sources and equally
distributed among the three time-slices. Species occurrence probabilities were then
modelled based on ecological similarity from 76 environmental variables and species co-
occurrences across grid-cells using the FRESCALO algorithm, accounting for sampling bias.
The underlying method may be applied to any geographic area, taxon and time slice, given
enough data are available (See Eichenberg et al., 2021 for details of the methodology). For
our study, we extracted the state-wide, gridded occurrence probabilities of all species that
occupied any Hessian grid cell (Fig. 1A).

Next, we estimated species' ranges in Hesse at each time slice based on the modelled
occurrence probabilities. To do so, we sampled 200 replicates from the occurrence
probability of each species in each grid cell at each time-slice with an independent binomial
draw and then summarized the mean number of grid cells occupied by in each time-slice per
species.

Then, we used the modelled species ranges per time slice to approximate RL indicators: We
estimated species current population status by the proportion of grid cells occupied by the
species in 1997-2017; the short-term population trend by the percentage of grid-cells lost or
gained from 1988-1996 to 1997-2017 (15 years); and the long-term population trend by the
percentage of grid-cells lost or gained from 1960-1987 to 1997-2017 (34 years). We then
used the cut-off values proposed by the German Federal Ministry for Nature Conservation
(Ludwig et al., 2006) and the methodology of the 5[th] version of the Hessian Red List for
vascular plants ("HRL"; Starke-Ottich et al., 2019) to convert the numeric values into RL
categories for each indicator (Table 1). For instance, a current population status of 12% grid

cells occupied translated into the current population status category "rare" (S), and a
decrease in cells occupied of 30% from 1988-1996 to 1997-2017 translated into the short-
term population change category "strong decrease" (↓↓). Notably, the approximation of
population trends from modelled species distributions is a simplification: It may lead to
erroneous estimates, if species occur over a relatively large range but in low densities, or in
limited areas with large populations. Similarly, a constant geographic range through time
may conceal a decline of number of individuals within this range. However, the official
Hessian Red List (HRL) explicitly encourages the use modelled species distributions in this
way (Starke-Ottich et al., 2019). Additionally, ARL has the advantage of being explicitly
defined, documented and hence reproducible, in contrast to expert-opinion based estimates
of population trends which often originate from a similar, "range-size-for-population-size"
proxy.
197
Finally, we combined the categorical estimates of RL indicators into the seven most relevant
extinction risk categories used for the 5[th] version of expert-based Hessian Red List (0, 1, 2,
3, R, V, *) following the same methodology used in HRL (Starke-Ottich et al., 2019). Each
combination of RL indicators translated into a specific extinction risk category. For instance,
the combination of current population status "very rare" (ss), long-term population trend
"stagnant" (=), and short-term population trend "moderate decrease" (↓), ss=(↓), resulted in
the extinction risk category "threatened" (3). Similarly, the combination of "very common"
(sh), "moderate decrease" (<) and "distinct increase" (↑), sh<↑, resulted in the extinction risk
category "not threatened" (*).
207
Expert-based regional Red List (HRL)
HRL is published by Hesse's Agency for Nature Conservation, Environment and Geology
every 10 years and includes assessments of all native species in the state (but neither
neophytes nor established archeophytes). In contrast to earlier versions, the 5[th] version of
HRL is based on methodology provided by the German Federal Ministry for Nature
Conservation for Red Listing (Ludwig et al., 2006) to increase comparability with other
German states. The next paragraphs give a short overview on the methodology of HRL, see
Ludwig (2006) and Starke-Ottich et al. (2019) for details.
216
HRL treated four subregions within Hesse separately, including Northwest (NW), Northeast
(NO), Southwest (SW), Southeast (SO) (Fig. 1; Buttler et al., 1996). The regions represented
dominant landscape types in the state (Klausing, 1988), and therefore differed in size: NO
covered 52% of Hesse, SO covered only 9%. While regions NW, NO, and SO principally

comprised mid-elevation mountain habitats, region SW mostly comprised the lowlands of the
Rhine and Main river floodplains.

For each subregion and species, HRL provided estimates of current population status
(extinct to very common), short-term population trend (very strong decrease to strong
increase), long-term population trend (very strong decrease to strong increase) (Table 1)
based on the evaluation of a total of 12 experts. The relevant time-span for the long-term
population trend was 50-150 years. Assessments for this period were based on information
from herbarium collections and literature. For the short-term population trend the relevant
time span was 10-25 years, and assessment was based on changes compared to the 4[th]
version of the HRL (Hemm et al., 2008). HRL only provided RL indicators for the subregions,
but not for the entire state.

Based on the three RL indicators HRL classified each species in each subregion into one of
nine extinction risk categories (Ludwig et al., 2006): extinct or lost (0), threatened with
extinction (1), highly threatened (2), threatened (3), threat of unknown extent (G), extremely
rare (R), near threatened (V), not threatened (*) and data deficient (D). The same extinction
risk categories are provided for the entire state based on the most positive evaluation in a
region. While the methodology (Ludwig et al., 2006) in principle allowed the inclusion of
specific risk factors, these were not used for HRL.

Comparing HRL and ARL
To compare ARL and HRL, we first harmonized the taxonomic reference between both lists
since the data underlying the ARL used a different taxonomic backbone than HRL (HRL:
Jäger, 2017; ARL: GermanSL; Jansen & Dengler, 2008 version 1.4,
https://germansl.infinitenature.org/). Next, we removed species which we found
incomparable between HRL and ARL including: (1) subspecies and varieties; (2) hybrids; (3)
species which names we could not harmonize between HRL and ARL because one name in
one list corresponded to multiple accepted names in the other list; (4) taxonomically difficult
groups, for instance apomictic species in *Rubus* and *Taraxacum*; (5) species in HRL with no
modelled distribution available from Eichenberg et al. (2021); (6) species predicted as
present in Hesse by ARL, but that were never considered present in Hesse by the HRL
(species formerly present and now considered extinct in HRL are included in the analyses);
(7) species that were considered present in Hesse by HRL but never by ARL.

To test the agreement between ARL und HRL, we lumped cases from all subregions and
compared the accuracy and magnitude of error for RL indicators and extinction risk. The

magnitude of error was the mismatch in number of ordinal classes, e. g. HRL: * and ARL: V

amounted to an error of 1, HRL: * and ARL: 1 amounted to an error of 4 (Table 1). To

calculate accuracy, we excluded any instances (an instance is the estimate of an RL

indicator or extinction risk in a specific region) classified as "?" in HRL or ARL. Likewise, for

calculating the accuracy of the current population status we excluded species classified as

"common" (h) in ARL, since this category was not used by HRL (which only uses

"moderately common", mh). Lumping the "h" category with the categories "sh" or "mh" was

not advisable, because records in "h" could either be close to "sh" or "mh". Additionally, to

calculate the magnitude of error, we excluded instances classified in category "R", because

this category was not easily set in ordinal relation with the other categories. Finally, we

compared the modelled percentage of grid-cells covered for current population status, and

the percentage change in coverage for short-term population trend and long-term population

trend with the categorized estimates from HRL. We also compared the match of the

extinction risk assessment between ARL and HRL for species on the state level (RL

indicators are not provided on the state level by HRL).

To identify systematic reasons for ARL-HRL mismatch, we first compared the accuracy of

estimation and magnitude of error among the current population status classes from HRL

(expected higher uncertainty for species with intermediate population size). Furthermore, we

identified instances for which ARL and HRL disagreed most to discuss possible

improvements of ARL and the integration of ARL into the next version of HRL. We defined

instances as "disagreeing most", if: they were considered as "not threatened" (*) in one list

and as one of "extinct" (0), "threatened with extinction" (1), or "extremely rare" (R) in the

other list (160 instances); or their summed magnitude of error across populations indicators

was more than 5 categories (82 instances); or ARL predicted a positive short-term or long-

term trend, since these are rare in HRL and only one expert used them at all (395

instances).

We consider identifying mismatches as the major contribution of automated assessments to

RL in the future (besides increased reproducibility), since it will allow a more efficient use of

effort by pointing experts to species which need increased scrutiny. Unfortunately, identifying

mismatches is also unsatisfying as it leaves the central question "Which one is right: HRL or

ARL?", unanswered. To give a rough hint on the answer to this question, we mimic what we

anticipate as future standard procedure, and use our expertise of the local flora (including

one contributing expert and the lead editor of HRL) to loosely score either HRL or ARL as

"likely more accurate" for cases in which both approaches disagreed most. To do so, we

explicitly consider new data which has emerged since 2019 (including on
www.inaturalist.org) when the last version of HRL was published.

Software used

We did all analyses in R (R Core Team, 2022) and used the caret (Kuhn, 2022), cowplot
(Wilke, 2020), plotly (Sievert, 2020), tidyverse v1.3.0 (Wickham et al., 2019), readxl v1.3.1
(Wickham & Bryan, 2022), viridis v0.5.1 (Garnier et al., 2021), and writexl (Ooms, 2021)
packages.

Results

We obtained modelled occurrence probabilities for 1,255 species in Hesse (Fig. 1A) and
information for 1,933 taxa from HRL (Fig. 1B). During matching we removed a total of 1,040
taxa with 4,111 instances. Of these, 147 taxa (730 instances) were subspecies or varieties;
39 species (195 instances) were hybrids; 38 species (190 instances) couldn't be harmonized
between HRL and ARL; 236 species (1,180 instances) belonged to taxonomically difficult
groups*;* 259 species (1,295 instances) had no modelled distribution; 348 species (519
instances) were not considered present in Hesse by HRL. In the end we retained 1,139
species with 5,023 instances (Fig. 1C) for the comparison of HRL and ARL, corresponding to
40.8% of the 2,790 taxa included in HRL (see Supplementary material S1 and S2 for a full
list of matched species and Supplementary material S3 for a list of removed taxa).

For the current population status our analyses included 3,848 instances of which ARL and
HRL agreed for 1,420 instances (36.9%, Fig. 2A). When excluding the 729 instances with
assessment "h" in ARL (See Methods), a total of 3,119 instances remained, with 1,420
instances of identical assessment (44,3%). In a striking 1,539 instances ARL suggested a
higher current population status than HRL (49,3%; magnitude of error of 1 category in 1,226
instances/39,3%, 2 categories: 285/9,1%, 3 categories: 14/0,4%, 4: 4/0,1%, Fig. 2A). In
contrast, ARL suggested a lower current populations status in 170 instances (5,5%;
magnitude of error of 1 category in 150 instances/4,8%, 2 categories: 16/0,5% and 3
categories: 4/0,1%).

For the short-term population trend our analyses included 3,463 instances, of which HRL
and ARL agreed in 1,723 instances (nearly 50%, Fig. 2B). In 471 instances ARL suggested
a more positive short-term population trend than HRL (magnitude of error of 1 category in
404 instances/11,7%, 2 categories: 57/1,65%, 3 categories: 9/0,3%). In contrast, in 1,253
instances ARL suggested a more negative trend than HRL (magnitude of error of 1 category

in 865 instances/25,1%, 2 categories: 286/8,3%, 3 categories: 100/2,9%, 4 categories:

2/0,05%, Fig. 3). The mismatch was particularly severe in the HRL categories (V) and =, for

which modelled assessment span from ↑ to ↓↓↓ (Fig. 2B).

For the long-term population trend our analyses included 3,463 instances, of which HRL and

ARL agreed in 2,237 instances (65%), 1,902 of them in category "=" (Fig. 2C). ARL

suggested a more positive long-term population trend than HRL in 550 instances (17,7%;

magnitude of error of 1 category in 462 instances/13,4%, 2: 75/2,2%, 3: 13/0,4%). In

contrast, ARL suggested a more negative trend in 655 instances (19%; magnitude of error of

1 category in 609 instances/17,7%, 2 categories: 43/1,2%, 3 categories: 3/0,1%).

For extinction risk assessments our analysis included 3,452 instances, of which ARL and

HRL agreed in 2,741 instances on the binary threatened/not threatened level (79%, Fig. 3A)

and in 2,147 instances (62%) on the detailed level of extinction risk categories, 1,816 of

them in the category not threatened (*). ARL suggested a lower extinction risk than HRL in

567 instances (16.5%). In contrast, ARL suggested a higher extinction risk than HRL in 555

instances (16%); 6% of instances were considered in extinction risk category (R). In cases of

mismatch between ARL and HRL, the mean magnitude of error was 1.65 categories. The

agreement in extinction risk assessments between ARL and HRL was similar in all regions,

with 60% in NO and NW, 63% in SW and 66% in SO.

For the extinction risk across the entire state, our analysis included 1,135 species, of which

ARL and HRL agreed for 847 species on the binary threatened/non-threatened level (229

threatened and 618 not threatened; accuracy of ARL: 0.84, Fig. 3B). HRL considered 288 of

species as threatened (3: 103, 2:114, 1: 69) and 737 not threatened (*: 644, V: 93), 33

species as rare (R), 2 species as data deficient (?), and 79 species as extinct (ex). In

contrast, ARL considered 415 species threatened (3: 136, 2: 116, 1: 163), 676 species not

threatened (*: 585, V: 91), 44 species as rare, and 4 species as extinct for Hesse.

Concerning the detailed extinction risk categories, the ARL and HRL agreed on 643 species

(56,7%). For 266 species (23,4%) ARL suggested a higher extinction risk than HRL

(magnitude of error of 1 category: 146 species/12,9%, 2 categories: 77/6,8%, 3 categories:

27/2,4%, 4 categories: 11/0,9%, and 5 categories: 5/0,4%). In contrast, for 226 species

(20%), ARL suggested a lower extinction risk than HRL (magnitude of error of 1 category:

120 species/10,6%, 2 categories: 63/5,6%, 3 categories: 26/2,3%, 4 categories: 10/0,9%, 5

categories: 6/0,5%, and 6 categories: 1/0,09%). Notably, ARL considered more than double

the number of species in the highest threat category than HRL. Per class sensitivity varied

between 0.26 and 0.79 and was much higher for the extreme classes (* = 0.79, 1 = 0.71) as compared to the intermediate classes (V: 0.24, 3: 0.32, 2: 0.26).

For the relation between current population status and ARL-HRL agreement of extinction risk estimates, we found the overall agreement across regions to be the lower, the rarer a species was (Fig. 4). The agreement between ARL and HRL was high for very common species (775 out of 778 instances, accuracy: 0.99) and common species (550 out of 653 instances, accuracy: 0.84) but decreased to 91 out of 332 species for extremely rare species (accuracy: 0.34). Similarly, the mean magnitude of error was the higher, the rarer a species was, and increased from 1.3 for very common species to 1.8 for extremely rare species (Fig. 4).

For the species which disagreed most between we judged ARL to estimate the current population situation as more accurate in 246 cases and HRL to be more accurate in 536 cases (Supplementary material S4).

Discussion

Here we compared estimates of RL indicators and extinction risk from an Automated Red List (ARL) based on species distribution models and automated assessments with a traditional expert-based Red List (HRL) in the state of Hesse, a well-documented study area in central Europe. We found both approaches to agree for extinction risk in between 57%-62% of cases and the estimates for individual RL indicators consistent in around half of the cases (current population status: 44,3%, short-term trend: 50%, long-term trend: 65%, Fig. 2). In many cases the mismatch was small (difference of 1 category in 23,5% of the instances) but could be considerable in some cases (up to 6 categories, Fig 3). We found the discrepancies between ARL and HRL to increase with species rarity (Fig. 4).

In the following paragraphs, we use our results to discuss possible improvements of ARL and the potential to integrate ARL in the next version of HRL (due 2029) as well as its potential for expert-based RL in general, along six themes.

Very Rare species ("es" and "ss")
Examples from this category often occured in forests or dry lawns, such as *Vicia sylvatica* (HRL: *; ARL: 2), *Iris sibirica* (HRL: 2; ARL: *), *Moenchia erecta* (HRL: 2; ARL: *), and *Hepatica nobilis* (HRL: *, ARL: 2). In addition to a potential mis-approximation of population trends by range size (see Methods) a main reason for mismatch of this category is that

documented occurrence records are often scarce for rare species (which by definition are found less frequently or may be particularly protected and hence not be documented). The performance of SDM, including the FRESCALO algorithm, depends to the amount of data available for model training (Wisz et al., 2008). Hence, a lower performance of ARL is expected for rare species. Additionally, the modelled occurrence probabilities used for ARL were based on distribution data from across Germany, so that edge effects may cause the model to predict species that only occur in neighbouring states as present in Hessen.

In the future, the limitation from data availability will be remedied by rapidly increasing data from ongoing efforts to compile existing data (e.g. https://www.gbif.org, https://land.gbif.de/) and to harness verified data from citizen science platforms (e.g., https://www.inaturalist.org/, https://floraincognita.com/) which will continuously improve ARL performance for rare species by increasing the data basis for SDM. Additionally, ARL performance can increase by using SDMs tailored to the study area. Such models may be implemented as part of an integrated pipeline for automated regional red listing (comparable to what sRedList does for the IUCN Red List, https://sredlist.eu), and additionally be combined with predictive automated assessment methods which can directly process incomplete information on population size and trends together with species functional traits and evolutionary relationship (Cazalis et al., 2022).

Species with intermediate population size (s, mh, h)

Examples for this category for which ARL and HRL differed in the estimation of current population status are *Salvia pratensis, Populus alba* (HRL: ss, ARL: h), *Arum maculatum, Dipsacus fullonum, Echium vulgare, Primula veris* (HRL: s, ARL: h), *Digitaria ischaemum, Saponaria officinalis, Sedum album* (HRL: ss, ARL: mh). Likely, a reason for the mismatch is limited data availability to experts and a relatively strong performance of ARL. Experts can usually assign very rare and very common species confidently to extinction risk categories and approximate trends in their population status. In contrast, recognizing trends in rare to common species is more challenging, since less information is available to experts, because they usually receive less attention (Starke-Ottich et al., 2019) but often are particularly threatened (Jansen et al., 2020). Hence, these species may increase or decrease substantially before changes are picked up by experts, losing valuable time for conservation. In the future, ARL estimates of RL indicators have the potential to serve as data-driven, reproducible baseline to inform experts during the assessment process in particular for long-term and short-term population trend.

Species of azonal (small-scale) habitats

Examples in this category are species from wet meadows, bogs, flooding meadows, dry lawns (following www.floraweb.de) which ARL often considered less threatened than HRL, such as *Equisetum pratense* (HRL: 3, ARL: *), *Carex flava, Iris sibirica, Moenchia erecta* (HRL: 2, ARL: *), *Crepis pulchra* (HRL: R, ARL: *). There may be multiple non-mutually exclusive explanations for this mismatch. First, these habitats have received considerable conservation and research attention by specialists in the last years making HRL particularly precise and accurate. Second, these habitats are often protected and, in some cases, hard to access, so that particularly few collection and observation records (but see Sperle & Bruelheide, 2021) as basis for the modelled distributions exist. Third, modelling the ranges for species of azonal, small-scale habitats is challenging. Our SDM approach is comprehensive, as it includes 76 variables to measure ecological similarity and includes species co-occurrences to model occurrence probabilities among grid cells (Eichenberg et al., 2021), but due to the 25 km^2 resolution it may fail to capture microhabitat conditions and therefore systematic underestimate the occurrence probability of species of azonal habitats. In the future higher resolution SDM may improve the match between ARL and HRL, as could the relaxation of the neighbourhood criterion in the modelling algorithm, yet ARL may remain of limited use for these species.

Species in taxonomically difficult groups and aggregate species

Examples in this category are *Alchemilla vulgaris* (NW: ARL: sh, HRL: s)*, Bistorta officinalis* (SW: ARL: h, HRL: ss)*, Carex flava* (SO: ARL: h, HRL: ss; SW: ARL: mh, HRL: es)*,* and *Galium mollugo* (SW: ARL: sh, HRL: ss). The reasons are likely an unsatisfactory taxonomic matching between HRL and ARL and the high identification uncertainty for these species. Although we have taken great care to resolve the taxonomies between the two datasets and have excluded particularly complex groups from the analyses, the correct assessment remains a challenge. Another interesting example is *Nymphaea alba*, where a high percentage of recorded occurrences was of hybrid origin or even misidentified, which is correctly reflected in the HRL but not ARL (Starke-Ottich et al., 2019). As a route forward for ARL, further effort for taxonomic harmonisation before analyses may remedy this problem, yet it will likely persist, since a correct identification of species in taxonomic difficult groups cannot be guaranteed when using data from multiple sources including for instance citizen science.

Species related to cultivation or anthropogenic habitats

Examples in this category for which ARL overestimates extinction risk (as compared to HRL and our post-hoc expert judgement) are *Juglans regia*, *Setaria viridis*, *Setaria pumila*, and

Solanum nigrum. One reason for this systematic error was the low number of occurrence
records included as basis for the modelling in ARL for these species, likely caused by the
disregard of data collectors due to an ascribed low value or relevance for RL. In contrast,
examples of species occurring naturally in agricultural fields or ruderal or urban habitats for
which HRL underestimated current distribution range, were *Anchusa officinalis, Barbarea*
stricta, Bryonia dioica, Filago germanica, Bupleurum rotundifolium, Galeopsis ladanum,
Geranium molle, Odontites vernus. An explanation could be, that especially urban habitats
and their flora receive less attention from experts in the course of Red list assessments.
These results suggest, that currently ARL estimates can supplement expert-based
assessments for species native in anthropogenic habitats, but are of limited value for
cultivated species. In the future, information on cultivation or habitat could be used to
exclude species a priori from ARL or more sophisticated ARL algorithms could incorporated
this information in the prediction.

Quantifying uncertainty and methodological considerations
All Red Listing approaches rely on arbitrarily chosen thresholds, either to convert continuous
modelled distributions (in the case of ARL) or abstract expert knowledge (in the case of
HRL) into the distinct categories required for Red Lists. While we have carefully chosen the
thresholds for ARL following the HRL methodology as much as possible, the guidelines are
necessarily vague (since they ought to apply to many taxa) and different thresholds could
change the results of ARL. Compared to HRL, ARL has the advantage, that the thresholds
are clearly documented and therefore the assessments are reproducible. Furthermore, it is
conceivable to statistically quantify this uncertainty in the future, for instance by integrating
assessments over multiple thresholds.

The definition of the reference time points for estimating short term and long-term population
trends add uncertainty. There are several reasons for this, which affect HRL and ARL
equally. First, less data is available further back in time. The recommended baseline for the
long-term trend is 50-150 years (Ludwig et al., 2006). This is conceptually challenging for
ARL, where theoretically some, but considerably less data exist from this time as basis for
modelling, and environmental variables used for modelling may have changed, but likely
even more severe for HRL, where experts may have some experience on the upper bound
of this interval and general information from specimen or (often grey) literature may be
accessible (e.g., Händler, 2019). In practice, for ARL we had to rely the first time slice with
modelled species distributions available, which was from 1960-1987 as baseline for the long-
term trend (on average 50 years ago at the time of the analyses).

Conclusion

While we consider expert assessments more accurate in many cases, particularly for very rare species and taxonomic difficult groups, our results indicate that the combination of SDM and automated assessments can provide valuable, reproducible input to the expert-assessment process, particularly for estimating short-term and long-term population trend and for species with intermediate abundance or from anthropogenic habitats. The value of automated assessment will potentially increase in the future, as more distribution data become available and if quantification of uncertainty and predictive approaches were to be included in an analyses pipeline specifically tailored for regional Red Lists. In Supplementary material S5-S11 we provide interactive graphs to identify differences between the automated assessment and expert-based regional RL to facilitate the integration of automated assessments into the regional Red List.

References

Azam, C., Gigot, G., Witte, I., & Schatz, B. (2016). National and subnational Red Lists in European and Mediterranean countries: Current state and use for conservation. *Endangered Species Research*, *30*, 255–266. https://doi.org/10.3354/esr00740

Bachman, S., Moat, J., Hill, A., Torre, J. de la, & Scott, B. (2011). Supporting Red List threat assessments with GeoCAT: Geospatial conservation assessment tool. *ZooKeys*, *150*, 117–126. https://doi.org/10.3897/zookeys.150.2109

Bachman, S., Walker, B., Barrios, S., Copeland, A., & Moat, J. (2020). Rapid Least Concern: Towards automating Red List assessments. *Biodiversity Data Journal*, *8*, e47018. https://doi.org/10.3897/BDJ.8.e47018

Bennun, L., Regan, E. C., Bird, J., van Bochove, J.-W., Katariya, V., Livingstone, S., Mitchell, R., Savy, C., Starkey, M., Temple, H., & Pilgrim, J. D. (2018). The Value of the IUCN Red List for Business Decision-Making. *Conservation Letters*, *11*(1), e12353. https://doi.org/10.1111/conl.12353

Betts, J., Young, R. P., Hilton-Taylor, C., Hoffmann, M., Rodríguez, J. P., Stuart, S. N., & Milner-Gulland, E. j. (2020). A framework for evaluating the impact of the IUCN Red

List of threatened species. *Conservation Biology*, *34*(3), 632–643.
https://doi.org/10.1111/cobi.13454

Buttler, K., Frede, A., Kubosch, R., Gregor, T., Hand, R., Cezanne, R., & Hodvina, S. (1996).
Rote Liste der Farn- und Samenpflanzen Hessens, 3. Fassung. Hessisches
Ministerium des Innern und für Landwirtschaft, Forsten und Naturschutz.

Cazalis, V., Di Marco, M., Butchart, S. H. M., Akçakaya, H. R., González-Suárez, M., Meyer,
C., Clausnitzer, V., Böhm, M., Zizka, A., Cardoso, P., Schipper, A. M., Bachman, S.
P., Young, B. E., Hoffmann, M., Benítez-López, A., Lucas, P. M., Pettorelli, N.,
Patoine, G., Pacifici, M., … Santini, L. (2022). Bridging the research-implementation
gap in IUCN Red List assessments. *Trends in Ecology & Evolution*, *37*(4), 359–370.
https://doi.org/10.1016/j.tree.2021.12.002

Cazalis, V., Santini, L., Lucas, P. M., González-Suárez, M., Hoffmann, M., Benítez-López,
A., Pacifici, M., Schipper, A. M., Böhm, M., Zizka, A., Clausnitzer, V., Meyer, C.,
Jung, M., Butchart, S. H. M., Cardoso, P., Mancini, G., Akçakaya, H. R., Young, B.
E., Patoine, G., & Di Marco, M. (2023). Prioritizing the reassessment of data-deficient
species on the IUCN Red List. *Conservation Biology*, *37*(6), e14139.
https://doi.org/10.1111/cobi.14139

Cowie, R., Bouchet, P., & Fontaine, B. (2022). The Sixth Mass Extinction: Fact, fiction or
speculation? *Biological reviews of the Cambridge Philosophical Society*, *97*, 0–000.
https://doi.org/10.1111/brv.12816

Dauby, G., Stévart, T., Droissart, V., Cosiaux, A., Deblauwe, V., Simo-Droissart, M., Sosef,
M. S. M., Lowry, P. P., Schatz, G. E., Gereau, R. E., & Couvreur, T. L. P. (2017).
ConR: An R package to assist large-scale multispecies preliminary conservation
assessments using distribution data. *Ecology and Evolution*, *7*(24), 11292–11303.
https://doi.org/10.1002/ece3.3704

Eichenberg, D., Bowler, D. E., Bonn, A., Bruelheide, H., Grescho, V., Harter, D., Jandt, U.,
May, R., Winter, M., & Jansen, F. (2021). Widespread decline in Central European

568   plant diversity across six decades. *Global Change Biology, 27*(5), 1097–1110.

569   https://doi.org/10.1111/gcb.15447

570   Franklin, J. (2010). *Mapping Species Distributions: Spatial Inference and Prediction.*

571   Cambridge University Press. https://doi.org/10.1017/CBO9780511810602

572   Frobel, K., & Schlumprecht, H. (2013). Erosion der Artenkenner—Ergebnise einer Befragung

573   und notwendige Reaktionen. *Naturschutz und Landschaftsplanung, 48*(4), 105–113.

574   Garnier, Simon, Ross, Noam, Rudis, Robert, Camargo, Pedro, A., Sciaini, Marco, Scherer, &

575   Cédric. (2021). *viridis—Colorblind-Friendly Color Maps for R.*

576   https://doi.org/10.5281/zenodo.4679424

577   Habel, J. C., Segerer, A., Ulrich, W., Torchyk, O., Weisser, W. W., & Schmitt, T. (2016).

578   Butterfly community shifts over two centuries. *Conservation Biology, 30*(4), 754–762.

579   https://doi.org/10.1111/cobi.12656

580   Händler, A. (2019). *Die aktuelle Verbreitung und Bestandssituation von Pflanzen der Roten*

581   *Liste Hessens 2008, im Vergleich mit den Angaben von Dillenius (1719), Walther*

582   *(1802) sowie Heyer & Rossmann (1860-1863) in Gießen und Umgebung* [Master

583   Thesis]. GEB-IDN/14834. http://geb.uni-giessen.de/geb/volltexte/2019/14834/

584   Hemm, K., Barth, U., Buttler, K., Frede, A., Kubasch, T., Gregor, T., Hand, R., Cezanne, R.,

585   Hodvina, S., Mahn, D., Nawrath, S., Huck, S., & Uebeler, M. (2008). *Rote Liste der*

586   *Farn- und Samenpflanzen Hessens, 4. Fassung.* Hessisches Ministerium für Umwelt,

587   ländlichen Raum und Verbraucherschutz.

588   Henry, E. G., Santini, L., Butchart, S., Gonzalez-Suarez, M., Lucas, P. M., Benitez-Lopez,

589   A., Mancini, G., Jung, M., Cardoso, P., Zizka, A., Meyer, C., Akcakaya, H. R.,

590   Berryman, A., Cazalis, V., & Di Marco, M. (2023). *Modelling the probability of*

591   *meeting IUCN Red List criteria to support reassessments* [Preprint]. Ecology.

592   https://doi.org/10.1101/2023.06.08.544254

593   Hessische Landesanstalt für Umwelt. (1976). *Rote Liste der bestandsgefährdeten Farn- und*

594   *Blütenpflanzen Hessens (Stand: März 1976).* Hessische Landesanstalt für Umwelt.

Hill, M. O. (2012). Local frequency as a key to interpreting species occurrence data when

recording effort is not known. *Methods in Ecology and Evolution*, *3*(1), 195–205.

https://doi.org/10.1111/j.2041-210X.2011.00146.x

Isaac, N. J. B., van Strien, A. J., August, T. A., de Zeeuw, M. P., & Roy, D. B. (2014).

Statistics for citizen science: Extracting signals of change from noisy ecological data.

Methods in Ecology and Evolution, *5*(10), 1052–1060. https://doi.org/10.1111/2041-

210X.12254

IUCN Standards and Petitions Subcommittee. (2017). *Guidelines for Using the IUCN Red

List—Categories and Criteria. Version 13. Prepared by the Standards and Petitions

Subcommittee. Http://www.iucnredlist.org/documents/RedListGuidelines.pdf* (S. 1–

60).

Jäger, E. (Hrsg). (2017). *Rothmaler – Exkursionsflora von Deutschland, Gefäßpflanzen:

Grundband, 21. Aufl.* Spektrum.

Jandt, U., Bruelheide, H., Jansen, F., Bonn, A., Grescho, V., Klenke, R. A., Sabatini, F. M.,

Bernhardt-Römermann, M., Blüml, V., Dengler, J., Diekmann, M., Doerfler, I., Döring,

U., Dullinger, S., Haider, S., Heinken, T., Horchler, P., Kuhn, G., Lindner, M., … Wulf,

M. (2022). More losses than gains during one century of plant biodiversity change in

Germany. *Nature*, *611*(7936), Article 7936. https://doi.org/10.1038/s41586-022-

05320-w

Jansen, F., Bonn, A., Bowler, D. E., Bruelheide, H., & Eichenberg, D. (2020). Moderately

common plants show highest relative losses. *Conservation Letters*, *13*(1), e12674.

https://doi.org/10.1111/conl.12674

Jansen, F., & Dengler, J. (2008). GermanSL – Eine universelle taxonomische Referenzliste

für Vegetationsdatenbanken. *Tuexenia*, 239–253.

Jiang, Z., Jiang, J., Wang, Y., Zhang, E., Zhang, Y., & Cai, B. (2020). Significance of country

red lists of endangered species for biodiversity conservation. *Biodiversity Science*,

28(5), 558. https://doi.org/10.17520/biods.2020149

Kalheber, H., Korneck, D., Müller, R., Nieschalk, A., Nieschalk, C., Sauer, H., & Seibig, A.

(1980). *Rote Liste der in Hessen ausgestorbenen, verschollenen und gefährdeten*

Farn- und Blütenpflanzen. 2. Fassung, Stand 31. 12. 1979. Hess. Landesanstalt für

Umwelt.

Kamp, J., Frank, C., Trautmann, S., Busch, M., Dröschmeister, R., Flade, M., Gerlach, B.,

Karthäuser, J., Kunz, F., Mitschke, A., Schwarz, J., & Sudfeldt, C. (2021). Population

trends of common breeding birds in Germany 1990–2018. *Journal of Ornithology*,

162(1), 1–15. https://doi.org/10.1007/s10336-020-01830-4

Klausing, O. (1988). Die Naturräume Hessens mit einer Karte der naturräumlichen

Gliederung 1:200 000. [2. Aufl.]. *Umweltplanung, Arbeits- Umweltschutz*, *67*, 1–43.

Kuhn, M. (2022). *caret: Classification and Regression Training*. https://CRAN.R-

project.org/package=caret

Ludwig, G., Haupt, H., Gruttke, H., & Binot-Hafke, M. (2006). Methodische Anleitung zur

Erstellung Roter Listen gefährdeter Tiere, Pflanzen und Pilze. *BfN Skripten*, *191*, 1–

97.

Ooms, J. (2021). *writexl: Export Data Frames to Excel „xlsx" Format*. https://CRAN.R-

project.org/package=writexl

R Core Team. (2022). *R: A Language and Environment for Statistical Computing*. R

Foundation for Statistical Computing. https://www.R-project.org/

Ralimanana, H., Perrigo, A. L., Smith, R. J., Borrell, J. S., Faurby, S., Rajaonah, M. T.,

Randriamboavonjy, T., Vorontsova, M. S., Cooke, R. S. C., Phelps, L. N., Sayol, F.,

Andela, N., Andermann, T., Andriamanohera, A. M., Andriambololonera, S.,

Bachman, S. P., Bacon, C. D., Baker, W. J., Belluardo, F., … Antonelli, A. (2022).

Madagascar's extraordinary biodiversity: Threats and opportunities. *Science*,

378(6623), eadf1466. https://doi.org/10.1126/science.adf1466

Rodrigues, A. S. L., Pilgrim, J. D., Lamoreux, J. F., Hoffmann, M., & Brooks, T. M. (2006).

The value of the IUCN Red List for conservation. *Trends in Ecology & Evolution*,

21(2), 71–76. https://doi.org/10.1016/j.tree.2005.10.010

Rondinini, C., Marco, M. D., Visconti, P., Butchart, S. H. M., & Boitani, L. (2014). Update or

Outdate: Long-Term Viability of the IUCN Red List. *Conservation Letters*, *7*(2), 126–

130. https://doi.org/10.1111/conl.12040

Schippmann, U. (2020). *Bibliography of National Red Lists for Vascular Plants.* (547. Aufl.).

Bundesamt für Naturschutz. https://doi.org/10.19217/skr547

Schmidt, M., Zizka, A., Traoré, S., Ataholo, M., Chatelain, C., Daget, P., Dressler, S., Hahn,

K., Kirchmair, I., Krohmer, J., Mbayngone, E., Müller, J. V., Nacoulma, B.,

Ouédraogo, A., Ouédraogo, O., Sambaré, O., Schuman, K., Wieringa, J. J., Zizka,

G., & Thiombiano, A. (2017). Diversity, distribution and preliminary conservation

status of the flora of Burkina Faso. *Phytotaxa Monographs*, *304*(1), 1–215.

Sievert, C. (2020). *Interactive Web-Based Data Visualization with R, plotly, and shiny*.

Chapman and Hall/CRC. https://plotly-r.com

Silva, S. V., Andermann, T., Zizka, A., Kozlowski, G., & Silvestro, D. (2022). Global

Estimation and Mapping of the Conservation Status of Tree Species Using Artificial

Intelligence. *Frontiers in Plant Science*, *13*.

https://www.frontiersin.org/articles/10.3389/fpls.2022.839792

Sperle, T., & Bruelheide, H. (2021). Climate change aggravates bog species extinctions in

the Black Forest (Germany). *Diversity and Distributions*, *27*(2), 282–295.

https://doi.org/10.1111/ddi.13184

Starke-Ottich, I., Gregor, T., Barth, U., Blatt, H., Böger, K., Bönsel, D., Cezanne, R., Frede,

A., Gottschlich, G., Hemm, K., Hodvina, S., Jansen, W., Kubosch, R., Mahn, D., &

Uebeler, M. (2019). *Rote Liste der Farn- und Samenpflanzen Hessens—5. Fassung*.

Hessisches Landesamt für Naturschutz, Umwelt und Gerologie.

https://www.hlnug.de/fileadmin/dokumente/naturschutz/Rote_Listen/RL_HE_Farn-

_und_Samenpflanzen_Fassung5_2019.pdf

Walker, B. E., Leão, T. C. C., Bachman, S. P., Lucas, E., & Nic Lughadha, E. (2022).

Evidence-based guidelines for automated conservation assessments of plant

species. *Conservation Biology*, *37*(1), e13992. https://doi.org/10.1111/cobi.13992

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D., François, R., Grolemund,

G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T. L., Miller, E., Bache, S.

M., Müller, K., Ooms, J., Robinson, D., Seidel, D. P., Spinu, V., … Yutani, H. (2019).

Welcome to the tidyverse. *Journal of Open Source Software*, *4*(43), 1686.

https://doi.org/10.21105/joss.01686

Wickham, H., & Bryan, J. (2022). *readxl: Read Excel Files*. https://CRAN.R-

project.org/package=readxl

Wilke, C. O. (2020). *cowplot: Streamlined Plot Theme and Plot Annotations for „ggplot2".*

https://CRAN.R-project.org/package=cowplot

Wisz, M. S., Hijmans, R. J., Li, J., Peterson, A. T., Graham, C. H., Guisan, A., & Group, N. P.

S. D. W. (2008). Effects of sample size on the performance of species distribution

models. *Diversity and Distributions*, *14*(5), 763–773. https://doi.org/10.1111/j.1472-

4642.2008.00482.x

Zizka, A., Andermann, T., & Silvestro, D. (2022). IUCNN – Deep learning approaches to

approximate species' extinction risk. *Diversity and Distributions*, *28*(2), 227–241.

https://doi.org/10.1111/ddi.13450

Zizka, A., Silvestro, D., Vitt, P., & Knight, T. M. (2021). Automated conservation assessment

of the orchid family with deep learning. *Conservation Biology*, *35*(3), 897–908.

https://doi.org/10.1111/cobi.13616

Data accessibility statement

The data are available in Supplementary material S1 and S2. All modelled occurrences and

analysis scripts will be available from a zenodo repository upon publication.

Tables

Table. 1. Categories of current population status, long-term trend and short-term trend used in the 5[th] version of the Hessian Red List for vascular plants (HRL) in the automated assessment in this study (ARL), and the cut-off values used to categorize modelled RL indicators. Risk factors were not used.

Current Population status		
Symbol	Category name	Percent grid-cells occupied
ex	Extinct	0
es	Extremely rare	0-1%
ss	Very rare	1-5%
s	Rare	5-15%
mh	Moderately common	15-35%
h	Common	35-70%
sh	Very common	75-100%
?	Unknown	-
Short-term population trend		
Symbol	Category name	Change in cells occupied between 1988-1996 and 1997-2017
↓↓↓	Very strong decrease	-100 - -35%
↓↓	Strong decrease	-35 - -25%
(↓)	Moderate decrease, or of unknown magnitude	-25 - -10%
=	Stagnant	-10 - +10%
↑	Distinct increase	> +10%
?	Data deficient	-
Long-term population trend		
Symbol	Category name	Change in cells occupied between 1960-1987 and 1997-2017
<<<	Very strong decrease	-100 - -80%
<<	Strong decrease	-80 - -55%
<	Moderate decrease	-55 - -30%
(<)	Decrease of unknown mangitude	-
=	Stagnant	-30 - +30%
>	Distinct increase	> +30%
?	Data deficient	-
Extinction risk		
Symbol	Category name	Corresponding IUCN category
0	Extinct or lost	RE
1	Threatened with extinction	CR
2	Strongly threatened	EN
3	Threatened	VU
G	Threat of unkown magnitude	[I]
R	Extremely rare	[R]
V	Near threatened	NT
D	Data deficient	DD
*	Not threatened	LC

Figures

Figure 1. The study area, the state of Hesse in central Germany. **A)** Modelled richness of all
species with distribution models available, as the cumulative occurrence probability in 25
km^2 grid cells. **B)** The number species in each subregion assessed in the 5th version of the
Hessian Red List of vascular plants (HRL). **C)** The number of species included in the
comparison between automated assessment and HRL. The inlay in A shows the position of
the study area in Europe.

715
Figure 2. Comparison of expert-based (HRL) and modelled (ARL) RL indicators. **A)** Current
population status, **B)** Short-term population trend, C) Long-term population trend. Each
datapoint represents an instance (species per subregion). Supplementary material S6-S8
are interactive versions of the subfigures indicating the species names and modelled
distribution for each datapoint.

Figure 3. Error matrix comparing the extinction risk estimates from our automated
assessment (ARL) with the 5[th] version of the Hessian Red List of vascular plants (HRL). A)
Extinction risk assessments for species per subregion (instance). B) State-wide extinction
risk for 1,135 species. Numbers and colours indicate the number of species in each
category. See Supplementary material S11 for an interactive versions of subfigure B
indicating the species names.

729
730 **Figure 4.** The accuracy of ARL compared to HRL depending on the species current
731 population status. Inlets show the error matrices for each current population status category
732 (HRL on the x-axis and ARL on the y-axis) and the distribution of the magnitude of error (by
733 how many categories the assessments disagree in how many cases). Data from all
734 subregions are shown together.