Distinct Activation Mechanisms of CXCR4 and ACKR3 Revealed by Single-Molecule Analysis of their Conformational Landscapes

Christopher T. Schafer[a,1], Raymond F. Pauszek III[b,2], Martin Gustavsson[a,3], Tracy M. Handel[a]*, and David P. Millar[b]*

[a] Skaggs School of Pharmacy and Pharmaceutical Sciences, Department of Pharmacology, University of California San Diego, La Jolla, CA 92037.
[b] Department of Integrative Structural and Computational Biology, The Scripps Research Institute, La Jolla, CA 92037.
[1] Present address: Amsterdam Institute for Molecular and Life Sciences, Department of Medicinal Chemistry, Faculty of Science, Vrije Universiteit Amsterdam, 1081 HV Amsterdam, Netherlands
[2] Present address: LUMICKS, 1105 AG Amsterdam, Netherlands
[3] Present address: Department of Biomedical Sciences, Faculty of Health and Medical Sciences, University of Copenhagen, 2200 Copenhagen, Denmark

*Correspondence and requests for materials should be addressed to D.P.M. (millar@scripps.edu) or T.M.H. (thandel@health.ucsd.edu)

Keywords: Chemokine receptor, single-molecule FRET, GPCR, ACKR3, conformational dynamics

This PDF file includes:
  Main Text
  Figures 1 to 6

Abstract

The canonical chemokine receptor CXCR4 and atypical receptor ACKR3 both respond to CXCL12 but induce different effector responses to regulate cell migration. While CXCR4 couples to G proteins and directly promotes cell migration, ACKR3 is G protein-independent and scavenges CXCL12 to regulate extracellular chemokine levels and maintain CXCR4 responsiveness, thereby indirectly influencing migration. The receptors also have distinct activation requirements. CXCR4 only responds to wild-type CXCL12 and is sensitive to mutation of the chemokine. By contrast, ACKR3 recruits GPCR kinases (GRKs) and β-arrestins and promiscuously responds to CXCL12, CXCL12 variants, other peptides and proteins, and is relatively insensitive to mutation. To investigate the role of conformational dynamics in the distinct pharmacological behaviors of CXCR4 and ACKR3, we employed single-molecule FRET to track discrete conformational states of the receptors in real-time. The data revealed that apo-CXCR4 preferentially populates a high-FRET inactive state, while apo-ACKR3 shows little conformational preference and high transition probabilities among multiple inactive, intermediate and active conformations, consistent with its propensity for activation. Multiple active-like ACKR3 conformations are populated in response to agonists, compared to the single CXCR4 active-state. This and the markedly different conformational landscapes of the receptors suggest that activation of ACKR3 may be achieved by a broader distribution of conformational states than CXCR4. Much of the conformational heterogeneity of ACKR3 is linked to a single residue that differs between ACKR3 and CXCR4. The dynamic properties of ACKR3 may underly its inability to form productive interactions with G proteins that would drive canonical GPCR signaling.

Main Text

Introduction

CXC chemokine receptor 4 (CXCR4) is one of the most intensively studied chemokine receptors due to its central role in driving cell migration during development and immune responses, and in cancer where it promotes tumor growth and metastasis (1-4). As a class A G protein-coupled receptor (GPCR), CXCR4 activates inhibitory Gαi protein signaling pathways to directly control cell movement in response to the chemokine CXCL12 (5). The activated receptor is also phosphorylated by GPCR kinases (GRKs), which promotes arrestin recruitment and cessation of the G protein signal. CXCR4 often works together with atypical chemokine receptor 3 (ACKR3, formerly CXCR7) which indirectly influences migration by scavenging CXCL12 to regulate available extracellular levels of the agonist and in turn the responsiveness of CXCR4 (6-8). In the absence of ACKR3 scavenging, excessive CXCL12 stimulation of CXCR4 leads to downregulation, resulting in profound effects on neuronal cell migration and development (9-11).

In contrast to CXCR4 and with some exceptions noted (12, 13), ACKR3 lacks G protein activity and instead is considered to be arrestin-biased (14). However, we and others (9) have shown that arrestins are dispensable for chemokine scavenging while GRK phosphorylation is critical for this function (15), suggesting that ACKR3 might be better described as GRK-biased. The molecular basis for its inability to couple to G proteins in most cell types remains an unanswered question. Our recently determined structures of ACKR3-ligand complexes showed the expected hallmarks of GPCR activation including "microswitch residues" in active state configurations, displacement of transmembrane helix 6 (TM6) away from the helical bundle and an open intracellular pocket, consistent with a receptor that should be able to activate G proteins (16). Accordingly, we replaced the intracellular loops (ICLs) of ACKR3 with those of CXCR2, a canonical G protein-coupled chemokine receptor; however, these changes did not lead to G protein activation (16), suggesting that the lack of coupling is not due to the absence of specific residue interactions. CXCL12 also adopts a distinct pose when bound to ACKR3 compared to CXCR4 and all other chemokines in chemokine-receptor complexes, but since small molecules induce similar biased effector responses, the chemokine pose cannot explain the effector-coupling bias (16). Having excluded other mechanisms we therefore surmised that the inability of ACKR3 to activate G proteins may be due to differences in receptor dynamics. Consistent with this hypothesis, the ICLs observed in ACKR3-agonist complexes are disordered, which may preclude productive effector coupling. The dynamic nature of ACKR3 is also suggested by its considerable constitutive activity in recruiting β-arrestins and its high level of constitutive internalization (16-18).

In addition to their distinct effector interactions, ACKR3 and CXCR4 have dramatically different susceptibilities to activation by different ligands. CXCR4 is activated by a single chemokine agonist, CXCL12. Moreover, modifications of the CXCL12 N-terminal signaling domain (e.g. single point mutations as in $CXCL12_{P2G}$ or multiple mutations as in $CXCL12_{LRHQ}$) transform the chemokine into an antagonist of CXCR4 (19, 20). Mutational analysis, modeling, and new structures of CXCR4 suggest that ligand activation involves a precise network of interacting residues that stabilize the active receptor conformation (21-25). By contrast, ACKR3 is activated by CXCL12 as well as its variants (including $CXCL12_{P2G}$ and $CXCL12_{LRHQ}$ (19, 20)), other chemokines (CXCL11) (26), other proteins (adrenomedullin, BAM22) (27, 28), and opioid peptides (29). In fact, most ligands for ACKR3 act as agonists, which is best explained by a non-specific "distortion" mechanism

of activation whereby any ligand that breaches the binding pocket causes helical movements that are permissive to GRK phosphorylation and arrestin recruitment. A distortion mechanism is consistent with the different binding poses observed for a small molecule agonist compared to the CXCL12 N-terminus in the receptor orthosteric pocket (16), and ligand bulk rather than specific interactions between agonist and ACKR3 being required for activation. We hypothesize that this distortion mechanism would also be facilitated by a receptor that is conformationally dynamic, by enabling nonspecific ligand interactions and more than a single conformation to promote activation.

To investigate the role of conformational dynamics in the distinct pharmacological behaviors of ACKR3 and CXCR4, we developed a single-molecule Förster resonance energy transfer (smFRET) approach. Many ensemble methods such as EPR, NMR and fluorescence-based methods have provided considerable insights into GPCR dynamics, conformational heterogeneity, and exchange between distinct structural states (30-36). However, these methods are limited in their ability to resolve the sequence of state-to-state transitions except in rare cases where transitions can be temporally coordinated with high precision (37). By contrast smFRET enables detection of sparsely populated states, reveals the sequence of state-to-state transitions and provides kinetic information through analysis of state dwell times. For example, smFRET studies of the β_2 adrenergic receptor (β_2AR) revealed a dynamic equilibrium between inactive and active conformations that was responsive to agonist and G protein binding (38). More recent smFRET studies of the glucagon receptor (39) and the A_{2A} receptor ($A_{2A}R$) (40) have documented the existence of distinct intermediate conformations in addition to inactive and active receptor conformations.

Here we present the first smFRET study of the chemokine receptors CXCR4 and ACKR3. Our experimental system allows real-time observation of the conformational fluctuations of individual receptor molecules in a native-like lipid environment and assessment of the differences in the conformational dynamics of the two receptors in their apo states and in response to ligands. Our results indicate that ACKR3 is more dynamic and conformationally heterogeneous than CXCR4 or other class A GPCRs previously studied, which may explain its activation prone nature and lack of G protein coupling. In contrast, CXCR4 appears less flexible, consistent with a more restricted, structurally-defined activation mechanism. Together these data characterize the molecular differences between CXCR4 and ACKR3 and suggest that enhanced conformational dynamics plays an important role in the atypical function of ACKR3.

Results
Development of smFRET experimental systems for CXCR4 and ACKR3.
To visualize conformational fluctuations of CXCR4 and ACKR3 by smFRET, cysteine residues were introduced into CXCR4 at positions $150^{4.40}$ in TM4 and $233^{6.29}$ in TM6, and at $159^{4.40}$ and $245^{6.28}$ in ACKR3 (Figs. S1A & B) (numbers in superscript refer to the Ballesteros-Weinstein numbering scheme for GPCRs) for covalent labeling with FRET donor (D, Alexa Fluor 555, A555) and acceptor (A, Cyanine5, Cy5) fluorophores. Single receptor molecules of labeled CXCR4 or ACKR3 were reconstituted into phospholipid nanodiscs to mimic the native membrane bilayer environment. Nanodiscs have been used in a wide variety of structural and biophysical studies of integral membrane proteins, including ACKR3, and reconstitute protein-lipid interactions lost in detergent systems (41-43). The nanodisc-receptor complexes were then tethered to a quartz slide through

150   biotinylation of the nanodisc membrane scaffolding protein (MSP) (Fig. 1A). To promote
151   monomeric receptor incorporation in each nanodisc, we utilized MSP1E3D1, which forms
152   nanodiscs with a diameter of approximately 13 nm (44, 45).
153
Crystal and cryo-electron microscopy (cryo-EM) structures of homologous class A
GPCRs, such as β_2AR, in inactive and active conformations, reveal that TM6 moves
outwards from the TM helical bundle during activation, whereas the position of TM4
remains relatively fixed (46). Accordingly, we anticipated that labeling at the positions
indicated above would be sensitive to transitions between inactive and active receptor
conformations and would give rise to different donor-acceptor distances and apparent
FRET efficiencies that could be resolved by smFRET measurements (shown
schematically in Fig. 1B). These positions are similar to those used effectively for β_2AR
(38), $A_{2A}R$, (40), and the μ-opioid receptor (47) for monitoring their conformational
dynamics by smFRET. Importantly, neither wild-type (WT) CXCR4 nor ACKR3 exhibited
significant labeling, obviating the need to remove any of the native cysteine residues in
either receptor. Moreover, the double cysteine receptor mutants retained CXCL12-
promoted β-arrestin2 recruitment (Figs. S1C & D). Only the E_{max} for arrestin recruitment
to CXCL12-stimulated ACKR3 was significantly altered by the mutations, while all other
pharmacological parameters were the same as for WT receptors.
169
<u>ACKR3 exhibits greater conformational dynamics than CXCR4.</u>
Receptor-nanodisc complexes were imaged on the slide surface using smFRET
microscopy by exciting the A555 donor with a green (532 nm) laser and monitoring the
resulting emission from both A555 and the Cy5 acceptor over time on separate segments
of a CCD camera. Several hundred individual receptor-nanodisc complexes were typically
observed in the field of view. Individual receptor molecules labeled with a single A555
donor and a single Cy5 acceptor were identified by single-step photobleaching transitions
(abrupt loss of acceptor fluorescence signal or simultaneous loss of fluorescence signal
in both channels), as shown for two representative ACKR3-nanodisc complexes in Fig.
1C. Additionally, anti-correlated changes in donor and acceptor emission prior to
photobleaching confirmed that FRET occurred between A555 and Cy5. Additional
examples of single-molecule traces for ACKR3 and CXCR4 are shown in Fig. S2. Dwell
times spent in one FRET level prior to switching to another level are generally in the
seconds range (Figs. 1C & S2).
184
The apparent FRET efficiencies from many individual receptor-nanodisc complexes,
recorded in the presence or absence of different ligands, were globally analyzed using
Hidden Markov Models (HMMs) assuming the presence of two, three, four, or five FRET
states (Materials and Methods). To evaluate the appropriate level of model complexity,
each resulting apparent FRET efficiency distribution was fit with a Gaussian Mixture Model
(GMM) and the corresponding Bayesian Information Criterion (BIC) was calculated. The
BIC is a statistical measure of the likelihood that the model describes the data, while also
penalizing the addition of parameters that could lead to overfitting of noise. Theoretically,
this value will be at a minimum for the model with the appropriate number of states. As an
additional criterion, we carefully compared the FRET distributions following the initial HMM
analysis for the different models and confirmed that peak positions were consistent across
the different experimental conditions.
197

These analyses indicated that three FRET states were sufficient to fit the smFRET data
for CXCR4 under all conditions (Figs. S3 & S4). The resulting apparent FRET efficiency
distributions are presented in Figs. 2A & B. The predominant high-FRET state (E_{app} = 0.85)
observed in the apo-state (Fig. 2A) suggests that TM4 and 6 are in close physical
proximity, which is consistent with the conformation of inactive GPCRs (48, 49).
Accordingly, we interpret this FRET state as the inactive conformation of CXCR4 and
designate it as R. The minor low-FRET state (E_{app} = 0.19) reflects outward movement of
TM6 away from TM4, as expected for an active receptor conformation (designated R*)
(46, 50) and consistent with the limited basal activity of CXCR4. Moreover, there was a
major population shift from the high-FRET state to the low-FRET state upon addition of
CXCL12$_{WT}$ (Fig. 2B), consistent with the expected stabilization of active GPCR
conformations by agonists (51, 52). To gain insight into the nature of the mid-FRET state
(E_{app} = 0.59), we examined the connectivity between all three FRET states. Two-
dimensional transition density probability (TDP) plots revealed that the three FRET states
were connected in a sequential fashion (Figs. 2A & B), indicating that the transitions
occurred within the same molecules. Notably, these observations exclude the possibility
that the mid-FRET state arises from different local fluorophore environments (hence FRET
efficiencies) for the two possible labeling orientations of the introduced cysteines:
assuming two receptor conformations, this model would produce four distinct FRET states,
but only two cross peaks in the TDP plot. Another significant observation is that direct
transitions between R and R* states were rarely observed (Figs. 2A & B). Taken together,
these results suggest that the mid-FRET state represents an intermediate receptor
conformation (designated R') that lies on the pathway between inactive and active
conformations. In the apo-state, transitions were mostly observed between states R and
R' (Fig. 2A), while in the presence of CXCL12$_{WT}$ the most frequent transitions were
observed between the R' and R* states (Fig. 2B).
224
In contrast, the apparent FRET efficiency histogram for ACKR3 in the apo state (Fig. 2C)
was much broader than the corresponding histogram for CXCR4 (Fig. 2A), indicating
greater conformational heterogeneity. Moreover, the smFRET data for ACKR3 could not
be described by three FRET states and the inclusion of a fourth state was necessary (Figs.
S5 & S6). The FRET distributions recovered by four-state global analysis are presented
in Figs. 2C & D. The positions of the high-FRET (E_{app} = 0.85) and low-FRET (E_{app} = 0.11)
peaks are similar to the corresponding peaks observed in CXCR4 and are likewise
assigned to inactive (R) and active (R*) conformations, respectively. Consistent with these
assignments, the R* state increased in population at the expense of the R state in the
presence of the chemokine agonists CXCL12$_{WT}$ (Fig. 2D) or CXCL11 (Fig. S7B). TDP
plots indicated that the four FRET states in ACKR3 were connected in a sequential fashion
and reside along the receptor activation pathway (Figs. 2C & D). Similar to the arguments
presented above for CXCR4, the intermediate FRET states are most likely discrete
receptor conformations and not arising from mixed labeling of the two introduced
cysteines. Accordingly, the mid-FRET peaks (E_{app} = 0.39 and E_{app} = 0.66) are assigned to
two intermediate receptor conformations, designated R*' and R', respectively. Notably, the
R*' state was not observed in CXCR4 (Figs. 2A & B). The R' state in ACKR3 decreased
in population in the presence of chemokine agonists (Fig. 2D, Fig. S7B), suggesting that
this state represents an inactive intermediate receptor conformation, consistent with its
position on the conformational pathway (closest to R). In striking contrast to apo-CXCR4,
apo-ACKR3 populated the four conformational states more or less equally, and all
possible sequential conformational transitions were observed (Fig. 2C). Thus, ACKR3 is

intrinsically more conformationally heterogenous and dynamic than CXCR4. In the
presence of CXCL12$_{WT}$, ACKR3 showed more frequent R' ↔ R*' and R*' ↔ R* transitions
compared with the apo-receptor, accounting for the population shift towards the R* state
(Figs. 2C & D).
251
<u>Effect of small-molecule ligands on conformational states of CXCR4 and ACKR3.</u>
The small-molecule ligand IT1t is reported to act as an inverse agonist of CXCR4 (53-55).
However, the conformational distribution of CXCR4 showed little change to the overall
apparent FRET profile, although R' ↔ R* transitions appeared in the TDP plot (Figs. 3A
& B, Fig. S8). This suggests that the small molecule does not suppress CXCR4 basal
signaling by changing the conformational equilibrium. Instead IT1t appears to increase
transition probabilities which may impair G protein coupling by CXCR4.
259
Treatment of ACKR3 with the small-molecule agonist VUF15485 (56) shifted the
conformational distribution of the receptor towards the active R* state (Figs. S7C & S8),
as expected for an agonist and supporting the assignment of the R* FRET state. In
contrast, treatment of ACKR3 with the small-molecule inverse agonist VUF16840 (52)
shifted the conformational distribution to the inactive R conformation, with a concomitant
decrease of R* (Fig. 3D, Fig. S8), consistent with the expected effect of an inverse agonist
and with suppression of the basal activity of ACKR3 by this ligand (52). The intermediate
R*' population also decreased in the presence of the inverse agonist (Fig. 3D, Fig. S8),
suggesting that this state represents an active-like receptor conformation, consistent with
its placement on the conformational pathway (closest to R*). The suppression of active
receptor conformations was also evident in the TDP, which revealed fewer transitions
between R*' and R* conformations relative to the apo receptor (Figs. 3C & D).
272
<u>CXCL12 N-terminal mutants promote active receptor conformations despite their</u>
<u>contrasting pharmacological effects on CXCR4 and ACKR3.</u>
CXCR4 is sensitive to N-terminal mutations of CXCL12 while ACKR3 is relatively
insensitive. For example, the variant CXCL12$_{P2G}$, containing a proline to glycine mutation
in the second position, and CXCL12$_{LRHQ}$, where the first three residues of CXCL12$_{WT}$ are
replaced with the four-residue motif LRHQ starting with L0, are antagonists of CXCR4 but
agonists of ACKR3 (19, 20). To gain further insight into the ligand-dependent responses
of CXCR4 and ACKR3, we examined how these mutant chemokines influence the
conformational states and dynamics of both receptors.
282
Surprisingly, CXCL12$_{P2G}$ promoted a shift to the active R* conformation of CXCR4
compared to the apo-receptor (R* increased by 16%, Figs. 4A & B, Fig. S8), although the
shift was less pronounced than observed for CXCL12$_{WT}$ (28% increase in R*, Fig. 2B, Fig.
S8). The shift is also evident in the TDPs where the state-to-state transitions involving the
R* active state were more probable for the CXCL12$_{P2G}$ complex compared with apo-
CXCR4 (Figs. 4A & B). The presence of CXCL12$_{LRHQ}$ had a more subtle effect on CXCR4:
the active R* conformation increased by 9% (Figs. 4A & C, Fig. S8). Despite the ability of
CXCL12$_{P2G}$ and CXCL12$_{LRHQ}$ to stabilize the active R* conformation of CXCR4, both
variants are known to act as antagonists (20). This suggests that the CXCL12 mutants
inhibit CXCR4 coupling to G proteins not by suppressing the active receptor population
but rather by increasing the dynamics of the receptor state-to-state transitions. Our results
suggest that the helical movements considered classic signatures of the active state may
not be sufficient for CXCR4 to engage productively with G proteins..

296
In the case of ACKR3, $CXCL12_{P2G}$ induced a modest increase in the population of the
active R* conformation relative to the apo-receptor (R* increased by 5%, Figs. 4D & E,
Fig. S8), consistent with the ability of this CXCL12 variant to act as an agonist of ACKR3
(20). Additionally, $CXCL12_{P2G}$ also promoted formation of the active-like intermediate R*'
conformation (R*' increased by 6%, Figs. 4D & E, Fig. S8), suggesting that activation of
ACKR3 can be achieved by populating the R*' state, not just the R* state, which is
consistent with a flexible, distortion activation mechanism. Together, $CXCL12_{P2G}$
increased active (R*) and active-like (R*') conformations by 11%, somewhat less than
observed for $CXCL12_{WT}$ (24%, Fig. 2C, Fig. S8). $CXCL12_{LRHQ}$ also promoted the active
(R*) and active-like (R*') conformations of ACKR3 (R* + R*' increased by 11%, Fig. 4F,
Fig. S8). Additionally, the probabilities of R ↔ R' transitions were suppressed relative to
apo receptor in the presence of $CXCL12_{P2G}$, while the probabilities for R'* ↔ R* transitions
were enhanced in the presence of the $CXCL12_{LRHQ}$ (Figs. 4E & F). These differences may
be a consequence of the longer residence time of the LRHQ mutant on the receptor (57).
The agonism observed for both chemokine variants (19, 20) suggests that both the active
R* and active-like R*' conformations of ACKR3 are sufficient for GRK phosphorylation and
arrestin recruitment.
314
<u>ACKR3 constitutive activity is linked to receptor conformational heterogeneity.</u>
As noted above, apo-ACKR3 displays little conformational selectivity, with similar
occupancies observed for all four FRET states (Fig. 2C). We hypothesized that this might
be related to the constitutive activity of the receptor (13, 16) and tied to the presence of
Tyr at position $257^{6.40}$ (Fig. 5A), which is a hydrophobic residue (V, I, or L) in all other
chemokine receptors. In many other class A GPCRs, mutating the residue at 6.40 results
in constitutive activity (32, 58-60) and previous analysis of rhodopsin suggests that this is
a consequence of lowering the energy barrier for transitions between different receptor
conformations (61). In our previous work, we showed that mutation of $Y257^{6.40}$ to leucine,
the corresponding amino acid in CXCR4, reduces constitutive arrestin recruitment to
ACKR3, while preserving the ability of the receptor to be activated by CXCL12 (Fig. S9)
(16). Adding this single-point mutation to our ACKR3 smFRET construct had a significant
impact, converting the broad conformational distribution of the WT apo-receptor to a
narrower distribution concentrated in the high-FRET region (Figs. 5B & C). Overall, the
FRET histogram is similar to what we observe for apo-CXCR4 (Fig. 2A), although the
active conformation is still split between R* and R*' states (the latter unique to ACKR3)
(Fig. 5C). State-to-state transitions were also suppressed relative to WT ACKR3 (Figs. 5B
& C). This result indicates that $Y257^{6.40}$ is a major determinant of the broad conformational
heterogeneity of ACKR3.
334
$CXCL12_{WT}$ promoted active and active-like conformations of $Y257^{6.40}L$ ACKR3 (R* + R*'
increased by 16%, Figs. 5C & E), although the effect was somewhat reduced in
comparison to WT ACKR3 (R* + R*' increased by 24%, Figs. 5B & D). Consistent with
this, the mutant receptor recruits β-arrestin in response to $CXCL12_{WT}$ with an E_{max} value
that is only slightly reduced relative to the WT receptor (Fig. S9, ~80% of WT) (16),
suggesting again that the population of the R*' intermediate conformational state may
contribute to receptor activation.
342
Discussion

The molecular basis for the atypical pharmacological behavior of ACKR3 is not well
understood. Since structures of ACKR3 show intracellular loop disorder, and progressive
structural substitutions within the loops fail to promote G protein coupling (16), we recently
proposed that the atypical nature of ACKR3 may be related to receptor conformational
dynamics (16, 62). Consistent with this hypothesis, in the present study we found that the
conformational dynamics of ACKR3 and the canonical GPCR CXCR4 are indeed
markedly different. Our smFRET results revealed four distinct conformations of apo-
ACKR3 with approximately equal populations: inactive R, active R*, R' (inactive-like) and
R*' (active-like) (Fig. 2C). The state-to-state TDP plots (Fig. 2C) further reinforced the
notion that ACKR3 is a flexible receptor that readily exchanges between different
conformational states, consistent with our previous structural studies where a flexible
intracellular interface in the absence of interaction partners was observed (16). In contrast,
apo-CXCR4 primarily populates the inactive R state, with only a single intermediate state
(R'), and relatively few transitions between states (Fig. 2A). These observations imply that
ACKR3 has a relatively flat energy landscape, with similar energy minima for the different
conformations, whereas the energy landscape of CXCR4 is more rugged (Fig. 6). For both
receptors, the energy barriers between states are sufficiently high that transitions occur
relatively slowly with seconds long dwell times (Figs. 1C and S2).
362
The relatively flat energy landscape of ACKR3 may account for the propensity of ACKR3
to be activated by a diverse range of ligands. In principle, ligands could promote activation
by providing more energetically favorable binding interactions with the receptor in the
active R* or R*' conformations relative to the inactive R or R' conformations. Alternatively,
ligands could destabilize the inactive R or R' conformations of the receptor, which would
also shift the receptor population to the active conformation(s). Our results do not
distinguish between these two possibilities. Regardless, the similar free energies of
different receptor conformations in ACKR3 (Fig. 6) implies that relatively little energy is
required to switch the receptor from one state to another. Moreover, activation can be
achieved by populating either the R* or R*' state. Thus, independent of specific binding
poses and receptor interactions, a variety of different ligands may be able to tip the
balance between different receptor conformations.
375
The TDP plots demonstrate that the R' state in CXCR4 and the R' and R*' states in ACKR3
are intermediate species that lie on the pathway between inactive R and active R* receptor
conformations. SmFRET studies of the glucagon receptor (39) and the $A2_{AR}$ (40) (both
labeled in TM4 and TM6, as here) also revealed a discrete intermediate receptor
conformation. In principle, the intermediates in CXCR4 and ACKR3 could represent partial
movements of TM6 from the inactive to active conformation or more subtle conformational
changes altering the photophysical characteristics of the probes without drastically altering
the donor-acceptor distance. Either possibility leads to detectable changes in apparent
FRET efficiency and reflect discrete conformational steps on the activation pathway;
however, it is not possible to resolve specific structural changes from the data.
Interestingly, the R*' intermediate appears to be unique to ACKR3. Moreover, this state
responds to agonist and inverse agonist ligands in the same manner as the active R*
state: agonists increase the population of the R*' state, while an inverse agonist decreases
the population. In contrast, the single intermediate state in CXCR4 (R') is not responsive
or only weakly responsive to ligands. The presence of both active (R*) and active-like (R*')
conformations in ACKR3 may be responsible for its activation prone nature and ligand
promiscuity.

Structural features that make ACKR3 conformationally flexible include Y257$^{6.40}$, which we
previously showed contributes to the constitutive activity of ACKR3 (16). Similar to the
constitutively active M257$^{6.40}$Y mutant of rhodopsin, Y$^{6.40}$ in ACKR3 stacks against Y$^{5.58}$
and Y$^{7.53}$ and stabilizes the active-like conformation (63). Additionally, the broad
conformational distribution and high probability of state-to-state transitions of ACKR3
parallel the dramatically lower energy barrier of the M257$^{6.40}$Y rhodopsin mutant relative
to WT rhodopsin (61). By contrast, mutation of Y257$^{6.40}$ to leucine, the corresponding
amino acid in CXCR4, promoted the dominance of an inactive high-FRET receptor
conformation similar to CXCR4 (Figs. 2A & 5C); it also reduced the transition probabilities
(Figs. 5B & C) and the ability of ACKR3 to constitutively recruit arrestin (16).

In addition to the contribution of Y257$^{6.40}$ to the dynamic behavior of ACKR3, a disulfide
bond between C34 in the receptor N-terminus and C287 in extracellular loop 3 (ECL3),
observed in all other reported chemokine receptor-chemokine structures (64-66) is
conspicuously missing in cryo-EM structures of ACKR3 with chemokine and small
molecule agonists (16). Furthermore, ACKR3 remains functional in the absence of these
cysteines (67) unlike other chemokine receptors (68, 69). The disulfide may constrain the
relative positions of TM1 and TM6/7 and the opening of the orthosteric pocket; therefore,
its absence may confer ACKR3 with greater conformational flexibility than other
chemokine receptors, consistent with our observations. It may also allow ACKR3 to be
activated by diverse ligands.

Our results provide insights into the linkage between receptor conformation and ligand
pharmacology in CXCR4. The chemokine variants CXCL12$_{P2G}$ and CXCL12$_{LRHQ}$ are
reported to act as antagonists of CXCR4 (19, 20), and the small molecule IT1t acts as an
inverse agonist (54-56). Surprisingly, none of these ligands inhibit formation of the active
R* conformation of CXCR4. In fact, the chemokine variants both stabilize and increase
this state to some degree, although less effectively than CXCL12$_{WT}$. Thus, the antagonism
and inverse agonism of these ligands does not appear to be linked exclusively to receptor
conformation, suggesting that the ligands inhibit coupling of G proteins to CXCR4 or
disrupt the ligand-receptor-G protein interaction network required for signaling (Fig. S10)
(21, 23). Interestingly, these ligands also increase the probabilities of state-to-state
transitions (Figs. 3B & 4B), suggesting that enhanced conformational exchange prevents
the receptor from productively engaging G proteins. Similarly, ACKR3 is naturally dynamic
and lacks G protein coupling, suggesting a common mechanism of G protein antagonism.
Future smFRET studies performed in the presence of G proteins should be informative in
this regard.

An unusual aspect of ACKR3 behavior is the failure to activate G proteins. Instead, ACKR3
recruits arrestins in response to phosphorylation of its C-terminal tail. Why ACKR3 does
not couple to G proteins, at least in most cells, is unclear and not readily explained by
differences in primary sequence, since insertion of a DRY box motif and substitution of all
ICLs from a canonical GPCR failed to confer G protein activity (16). It is possible that the
active receptor conformation clashes sterically with the G protein as suggested by docking
of G proteins to structures of ACKR3 (16). Alternatively, the receptor dynamics and
conformational transitions revealed here may prevent formation of productive contacts
between ACKR3 and G protein that are required for coupling, even though G proteins
appear to constitutively associate with the receptor (13, 16, 70). An important caveat is

that our study does not report on the dynamics of the other TM helices and H8, some of
which are known to participate in arrestin interactions (31, 32). Lack of a well-organized
intracellular pocket due to frequent conformational transitions may also explain why the
fingerloop of arrestin is not observed to interact with the pocket, in contrast with other
GPCRs (71, 72), but instead inserts into membranes/micelles adjacent to the receptor
(62). Nevertheless, arrestins are still recruited to CXCL12-stimulated ACKR3 due to GRK
phosphorylation of the receptor C-terminal tail (15, 73). Since GRKs also interact with the
cytoplasmic pocket to facilitate phosphorylation, it remains to be determined how
dynamics might decouple G protein activation and arrestin binding to the receptor
cytoplasmic pocket (62) while supporting pocket-mediated GRK activity. However, given
the fleeting interaction between GRKs and GPCRs (74), rapid state sampling by ACKR3
may not necessarily be detrimental to GRK engagement and phosphorylation.
Furthermore, conformational intermediates in addition to the fully active receptor have
been shown to be targets for GRK phosphorylation, such as the early photoactivated
rhodopsin metarhodopsin I (75). A more constrained system may be necessary to promote
productive interactions between ACKR3 and G proteins. Along these lines, a local
increase of membrane pressure in certain cell environments could explain the apparent
ability of ACKR3 to activate G proteins in astrocytes and glioma cells (12, 13). The atypical
activation of ACKR3 does not appear to be dependent on any singular receptor feature
and is likely a combination of several factors.
462
The pharmacological behavior of ACKR3 resembles the human cytomegalovirus
chemokine receptor US28, which also recognizes diverse chemokines, constitutively
internalizes, and displays multiple functional conformations (76). Similar to ACKR3, US28
appears to be activated by distortion of the orthosteric binding pocket rather than by
specific side chain contacts between receptor and ligand, which is supported by multiple
active conformations observed for both apo-US28 and US28 with different agonists (76-
79). Whether US28 also has a relatively flat energy landscape like ACKR3 remains to be
seen. The conformational dynamics and activation mechanisms revealed here for ACKR3
may also be operative in other chemokine receptors that respond to multiple ligands and
have considerable constitutive activity, such as CCR1, CCR2, and CCR3 (64, 80). Finally,
the ability of ACKR3 to be activated by populating more than one conformational state
may explain why antagonizing the receptor by targeting the orthosteric binding pocket has
proven to be challenging; in contrast, the specific requirements for CXCR4 agonism has
permitted the development of many orthosteric antagonists but few agonists (81). Drug
discovery efforts aimed at inhibiting ACKR3 may therefore require allosteric strategies.
478
479 **Materials and Methods**
480 Unless otherwise stated all chemicals and reagents were purchased from SigmaAldrich
481 or Fisher Scientific. Methoxy e-Coelenterazine (Prolume Purple) was purchased from
482 Nanolight Technologies (Prolume LTD).
483
Cloning
Human ACKR3 (residues 2-362) preceded by an N-terminal HA signal sequence and
followed by C-terminal 10His and FLAG tags was cloned into the pFasBac vector for
purification from *Sf9* cells. Human CXCR4 (residues 2-352) with an N-terminal FLAG tag
and C-terminal 10His was inserted into pFasBac for purification. For cell-based assays,
ACKR3 (residues 2-362) or CXCR4 (residues 2-352) were inserted into pcDNA3.1
expression vector with an N-terminal FLAG tag and followed with a C-terminal *Renilla*

luciferase II (ACKR3_rlucII and CXCR4_rlucII). Site-directed mutagenesis was performed
by overlap extension and confirmed by Sanger sequencing. No native cysteines were
substituted for either CXCR4 or ACKR3.
494

<u>Arrestin recruitment by BRET</u>
Arrestin recruitment to ACKR3 and CXCR4 was detected using a BRET2 assay as
previously described (15, 57). Briefly, HEK293T cells (ATCC) were plated at 750k/well in
a 6-well dish in Dulbecco's modified eagle media (DMEM) with 10% fetal bovine serum
(FBS) and transfected 24 hrs later with 50 ng receptor_rlucII DNA, 1 µg GFP10_β-
arrestin2 (a kind gift from N. Heveker, Université de Montréal, Canada), and 1.4 µg empty
pcDNA3.1 vector using TransIT-LT1 transfection system (MirusBio) and expressed for 40
hrs. The cells were then washed with PBS (137 mM NaCl, 2.7 mM KCl, 10 mM Na_2HPO_4,
1.8 mM KH_2PO_4, pH 7.4) and mechanically lifted in Tyrode's buffer (25 mM HEPES, 140
mM NaCl, 1 mM $CaCl_2$, 2.7 mM KCl, 12 mM $NaHCO_3$, 5.6 mM Glucose, 0.5 mM $MgCl_2$,
0.37 mM NaH_2PO_4, pH 7.5). 100k cells were plated per 96-well white BRET plate (BD
Fisher) and reattached for 45 min at 37 °C. GFP expression was checked using a
SpectraMax M5 plate fluorometer (Molecular Devices) with 485 nm excitation, 538 nm
emission, and 530 nm cutoff. 5 µM Prolume Purple substrate was subsequently added
and total luminescence detected using a TECAN Spark Luminometer (TECAN Life
Sciences) at 37 °C. CXCL12 was then added to each well at the indicated final
concentrations and BRET was read using default BRET2 settings (blue emission 360-440
nm, red emission 505-575 nm) and an integration time of 0.5 sec. Experiments were
baseline matched and normalized to the E_{max} of WT receptor. The reported data is the
average of three independent experiments performed in duplicate. Points were fit to a
sigmoidal dose-response model using SigmaPlot 11.0 (Systat Software, Inc).
516

<u>Receptor purification, labeling, and nanodisc reconstitution</u>
M159$^{4.40}$C/Q245$^{6.28}$C ACKR3 (WT and Y257$^{6.40}$L) and L150$^{4.40}$C/Q233$^{6.29}$C CXCR4 were
purified from *Sf9* cells (Expression Systems) as previously described (16). Briefly, *Sf9* cells
were infected with baculovirus (prepared using Bac-to-Bac Baculovirus Expression
System, Invitrogen) containing either the mutant ACKR3 or CXCR4. Cells were harvested
after 48 hrs and membranes dounce homogenized in hypotonic buffer (10 mM HEPES pH
7.5, 10 mM $MgCl_2$, 20 mM KCl) followed three more times with hypotonic buffer with 1 M
NaCl. The membranes were spun down at 50k x g for 30 min and resuspended between
each round of douncing. After the final round, membranes were incubated with 100 µM
CCX662 (Chemocentryx Inc.) for ACKR3 or 100 µM IT1t for CXCR4 and solubilized in 50
mM HEPES pH 7.5, 400 mM NaCl, 0.75/0.15% dodecyl maltoside/cholesteryl
hemisuccinate (DDM/CHS) with a protease inhibitor tablet (Roche) for 4 hrs. Insoluble
material was then removed by centrifugation at 50k x g for 30 min and Talon resin
(Clontech) with 20 mM imidazole overnight binding at 4 °C. The resin was then transferred
to a column and washed with WB1 (50 mM HEPES pH 7.5, 400 mM NaCl, 0.1/0.02%
DDM/CHS, 10% glycerol, 20 mM imidazole) followed by WB2 (WB1 with 0.025/0.005%
DDM/CHS) and finally eluted with WB2 with 250 mM imidazole. The imidazole was
removed by desalting column (PD MiniTrap G-25, GE Healthcare). Final protein
concentration was determined by A_{280} using an extinction coefficient of 75000 $M^{-1}cm^{-1}$
(ACKR3) and 58850 $M^{-1}cm^{-1}$ (CXCR4). Samples were snap frozen in liquid nitrogen, and
stored at -80 °C until use.
538

When ready to prepare samples for smFRET measurements, two nanomoles of receptor
was thawed and incubated with fourteen nanomoles of Alexa Fluor 555 C2 maleimide
(A555) and Cy5 maleimide overnight at 4 °C with rotation. The next morning, free label
was removed by dilution using a 100k Da cut-off spin concentrator (Amicon) and the
sample concentrated to ~100 µl. Label incorporation was evaluated by measuring the
absorbance at A_{280} (ε_{280} = 75000 $M^{-1}cm^{-1}$ for ACKR3 and 58850 $M^{-1}cm^{-1}$ for CXCR4), A_{555}
(ε_{555} = 150000 $M^{-1}cm^{-1}$), and A_{645} (ε_{645} = 250000 $M^{-1}cm^{-1}$) to detect the concentrations of
the labeled receptor, A555, and Cy5, respectively. The contribution of the fluorophores to
A_{280} was removed before determining labeled receptor concentration. The entire sample
was used for nanodisc reconstitution.
549
Labeled receptors were reconstituted into biotinylated MSP1E3D1 nanodiscs as
previously described (16). Briefly, 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phosphocholine
(POPC, Avanti) and 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phospho-(1'-rac-glycerol) (POPG,
Avanti) were prepared in a 3:2 POPC:POPG ratio and solubilized in ND buffer (25 mM
HEPES pH 7.5, 150 mM NaCl, 180 mM cholate). MSP1E3D1 was expressed and purified
as previously described (82) and biotinylated with EZ-Link NHS-polyethylene glycol 4
(PEG4)-Biotin (Thermo Fisher) per manufacturer instructions. The receptors, MSP1E3D1,
and lipids were combined at a molar ratio of 0.1:1:110 for ACKR3:MSP:lipids respectively.
Additional ND buffer was added to keep the final cholate concentration >20 mM. After 30
min at 4 °C, 200 mg of Biobeads (Bio-Rad) were added and incubated for 3-6 hrs. The
sample was then loaded on a Superdex 200 10/300 GL column equilibrated with 25 mM
HEPES pH 7.5, 150 mM NaCl and fractions containing nanodisc complexes were
combined. 200 µl Talon resin was added with 20 mM imidazole (final concentration) and
the samples were incubated for 16 hrs at 4 °C. The resin was then transferred to a Micro
Bio-Spin Column (Bio-Rad) and washed with 25 mM HEPES pH 7.5, 150 mM NaCl, 20
mM imidazole and eluted with 25 mM HEPES pH 7.5, 150 mM NaCl, 250 mM imidazole.
Imidazole was removed by buffer-exchange using 100k Da spin concentrators and 25 mM
HEPES pH 7.5, 150 mM NaCl. Sample were concentrated to ~1 µM and stored at 4 °C
until use.
569
<u>CXCL12 purification from *E. Coli*</u>
CXCL12 was expressed and purified as previously described (15, 16). Briefly, the
chemokines were expressed by IPTG induction in BL21(DE3)pLysS cells. The cells were
collected by centrifugation, resuspended in 50 mM tris pH 7.5, 150 mM NaCl and lysed by
sonication. Inclusion bodies were then collected by centrifugation, resuspended in
equilibration buffer (50 mM tris, 6 M guanidine-HCl pH 8.0), sonicated to release the
chemokines and the samples centrifuged again to pellet insoluble material. The
supernatant was then passed over a Ni-nitrilotriacetic acid (NTA) column equilibrated with
equilibration buffer to bind the His-tagged chemokines. The column was washed with
wash buffer (50 mM MES pH 6.0, 6 M guanidine-HCl) and eluted with 50 mM acetate pH
4.0, 6 M guanidine-HCl. The chemokine-containing elutions were pooled and dithiothreitol
(DTT) added to a final concentration of 4 mM. After incubating 10 min, the solution was
added dropwise into refolding buffer (50 mM tris pH 7.5, 500 mM arginine-HCl, 1 mM
EDTA, 1 mM oxidized glutathione) and incubated at room temperature for 4 hrs before
dialyzing against 20 mM tris pH 8.0, 50 mM NaCl. To remove the N-terminal purification
tag, enterokinase was added and the sample incubated at 37 °C for 5 days. Uncleaved
chemokine and free tags were removed by reverse Ni-NTA and eluted with wash buffer.
Finally, the sample was purified on a reverse-phase C18 column equilibrated with 75%

buffer A (0.1% trifluoroacetic acid (TFA)) and 25% buffer B (0.1% TFA, 90% acetonitrile)
and eluted by a linear gradient of buffer B. The pure protein was lyophilized and stored at
-80 °C until use.
591
smFRET microscopy
smFRET experiments were performed on a custom built prism-based TIRF microscope
as previously described (83). Briefly, a flow cell was assembled on a quartz slide
passivated with polyethylene glycol (PEG) and a small fraction of biotinylated PEG, after
which neutravidin was introduced (84). Labeled ACKR3 or CXCR4 in biotinylated
nanodiscs were diluted into trolox buffer (25 mM HEPES pH 7.5, 150 mM NaCl, 1 mM
propyl gallate, 5 mM trolox), flowed into the sample chamber, and incubated for 5 min at
room temperature. The samples were then washed twice with imaging buffer (trolox buffer
with 2 mM protocatechuic acid, 50 nM protocatechuate-3,4-dioxygenase). Movies were
collected with 100 ms integration time using a custom single-molecule data acquisition
program to control the CCD camera (Andor). Single-molecule donor and acceptor
emission traces were extracted from the recordings using custom IDL (Interactive Data
Language) scripts. The software packages used to control the CCD camera and extract
time trajectories were provided by Dr. Taekjip Ha. In all cases, five initial apo-receptor
movies were recorded at different locations on the slide and then ligand was flowed into
the cell by two washes with chemokine or small molecule in imaging buffer at final
concentrations of 500 nM for chemokines (CXCL12$_{WT}$, CXCL12$_{P2G}$, CXCL12$_{LRHQ}$,
CXCL11) and 1 µM for the small molecules (IT1t, VUF16840, VUF15485). Ten more
movies were collected for each condition at different locations on the slide. The data
presented are a composite of at least three individual slides and treatments.
612
FRET trajectories were generated and analyzed using custom software written in-house
(https://github.com/rpauszek/smtirf). Donor and acceptor traces for each molecule were
corrected for donor bleed through and background signal and apparent FRET efficiencies
were calculated as $E_{app} = I_A/(I_A+I_D)$, where E_{app} is the apparent FRET efficiency at each
time point and I_D and I_A are the corresponding donor and acceptor fluorophore intensities,
respectively. Traces were screened manually for single donor and acceptor bleach steps
and anti-correlated behavior between fluorophores to confirm the presence of single
receptors within the identified particles and single donor and acceptor labeling. All traces
for a particular protein/ligand combination were analyzed globally by a single Hidden
Markov Model assuming two, three, four, or five states and shared variance as previously
described (83). Briefly, each model was trained on all selected trajectories for a given
sample simultaneously using an expectation-maximization method (85). Once a model
was trained, the Viterbi algorithm (85) was used to determine the most likely hidden path
for each trajectory. This labeled state path was then used to aggregate all data points
belonging to a particular state in order to compile composite histograms of apparent FRET
efficiency, using a Kernel Density estimation algorithm (Python package Scikit-learn,
version 1.2.2) with a Gaussian kernel and a bandwidth of 0.04. The relative populations
of distinct FRET states were directly obtained during compilation of the corresponding
histograms. The resulting apparent FRET efficiency histograms were fit with a Gaussian
Mixture Model (Scikit-learn) and the corresponding Bayesian Information Criterion (BIC)
was calculated as described (86). Transition density probability plots revealing the
connectivity among individual FRET states were calculated as described (87).
635
636   Data availability

637 All relevant data supporting this study are included in the Article or Supplementary files
638 and are available from the authors upon request.
639

Acknowledgments
640
We gratefully acknowledge the initial work on this project undertaken by Chunxia Zhao
and Rajan Lamichhane. Additionally, we thank Handel lab members Cheyanne Shinn,
Catherina Salanga, and Nicholas Chimileski for providing the chemokines CXCL11 and
$CXCL12_{P2G}$, respectively, R. Leurs (Vrije Universiteit Amsterdam) for the small molecules
VUF16840 and VUF15485, and N. Heveker (Université de Montréal) for the GFP10_β-
arrestin2 plasmid. This work was supported by National Institutes of Health grants R01
GM133157 (D.P.M./T.M.H), R01 CA254402 (T.M.H.), R01 AI161880 (T.M.H.), F32
GM137505 (C.T.S.), F32 GM115017 (R.F.P.) and T32 AI007354 (R.F.P.). Additional
support was from the Robertson Foundation/Cancer Research Institute Irvington
Postdoctoral Fellowship (M.G.).

Author Contributions:
Conception and design of the research: C.T.S., T.M.H., and D.P.M.
Methodology: C.T.S., R.F.P., and M.G.
Software: R.F.P.
Validation: C.T.S. and R.F.P.
Formal Analysis: C.T.S. and R.F.P.
Investigation: C.T.S.
Writing-Original Draft Preparation: C.T.S., T.M.H., and D.P.M.
Writing-Reviewing & Editing: C.T.S., R.F.P., M.G., T.M.H., and D.P.M.
Visualization: C.T.S. and R.F.P.
Supervision: T.M.H. and D.P.M.
Funding Acquisition: T.M.H. and D.P.M.
664

665 **Competing Interest Statement:** T.M.H. is a cofounder of Lassogen Inc. and serves on
666 the Scientific Advisory Boards of Abilita Bio, and Abalone Bio. The terms of these
667 arrangements have been reviewed and approved by the University of California, San
668 Diego in accordance with its conflict of interest policies. The other authors declare that
669 they have no competing interests.
670

References
671
1.   S. Chatterjee, B. Behnam Azad, S. Nimmagadda, The intricate role of CXCR4 in
cancer. *Adv Cancer Res* **124**, 31-82 (2014).
2.   F. Balkwill, The significance of cancer cell expression of the chemokine receptor
CXCR4. *Semin Cancer Biol* **14**, 171-179 (2004).
3.   U. M. Domanska *et al.*, A review on CXCR4/CXCL12 axis in oncology: no place
to hide. *Eur J Cancer* **49**, 219-230 (2013).
4.   N. Kawaguchi, T. T. Zhang, T. Nakanishi, Involvement of CXCR4 in Normal and
Abnormal Development. *Cells* **8**, 185 (2019).
5.   I. Kufareva, M. Gustavsson, Y. Zheng, B. S. Stephens, T. M. Handel, What Do
Structures Tell Us About Chemokine Receptor Function and Antagonism? *Annu
Rev Biophys* **46**, 175-198 (2017).
6.   J. W. Griffith, C. L. Sokol, A. D. Luster, Chemokines and chemokine receptors:
positioning cells for host defense and immunity. *Annu Rev Immunol* **32**, 659-702
(2014).

7.   R. J. Nibbs, G. J. Graham, Immune regulation by atypical chemokine receptors.
Nat Rev Immunol **13**, 815-829 (2013).

8.   A. Vacchini, M. Locati, E. M. Borroni, Overview and potential unifying themes of
the atypical chemokine receptor family. *J Leukoc Biol* **99**, 883-892 (2016).

9.   F. Saaber *et al.*, ACKR3 Regulation of Neuronal Migration Requires ACKR3
Phosphorylation, but Not beta-Arrestin. *Cell Rep* **26**, 1473-1488.e1479 (2019).

10.   S. Lau *et al.*, A negative-feedback loop maintains optimal chemokine
concentrations for directional cell migration. *Nat Cell Biol* **22**, 266-273 (2020).

11.   M. Wong *et al.*, Dynamic Buffering of Extracellular Chemokine by a Dedicated
Scavenger Pathway Enables Robust Adaptation during Directed Tissue
Migration. *Dev Cell* **52**, 492-508 e410 (2020).

12.   V. Odemis *et al.*, The presumed atypical chemokine receptor CXCR7 signals
through G(i/o) proteins in primary rodent astrocytes and human glioma cells. *Glia*
60, 372-381 (2012).

13.   A. Fumagalli *et al.*, The atypical chemokine receptor 3 interacts with Connexin 43
inhibiting astrocytic gap junctional intercellular communication. *Nat Commun* **11**,
4855 (2020).

14.   S. Rajagopal *et al.*, Beta-arrestin- but not G protein-mediated signaling by the
"decoy" receptor CXCR7. *Proc Natl Acad Sci U S A* **107**, 628-632 (2010).

15.   C. T. Schafer, Q. Chen, J. J. G. Tesmer, T. M. Handel, Atypical Chemokine
Receptor 3 'Senses' CXC Chemokine Receptor 4 Activation Through GPCR
Kinase Phosphorylation. *Mol Pharmacol* **104**, 174-186 (2023).

16.   Y. C. Yen *et al.*, Structures of atypical chemokine receptor 3 reveal the basis for
its promiscuity and signaling bias. *Sci Adv* **8**, eabn8063 (2022).

17.   B. E. Hopkins *et al.*, Effects of Small Molecule Ligands on ACKR3 Receptors.
Mol Pharmacol **102**, 128-138 (2022).

18.   U. Naumann *et al.*, CXCR7 functions as a scavenger for CXCL12 and CXCL11.
PLoS One **5**, e9175 (2010).

19.   M. S. Hanes *et al.*, Dual targeting of the chemokine receptors CXCR4 and
ACKR3 with novel engineered chemokines. *J Biol Chem* **290**, 22385-22397
(2015).

20.   A. Jaracz-Ros *et al.*, Differential activity and selectivity of N-terminal modified
CXCL12 chemokines at the CXCR4 and ACKR3 receptors. *J Leukoc Biol* **107**,
1123-1135 (2020).

21.   T. Ngo *et al.*, Crosslinking-guided geometry of a complete CXC receptor-
chemokine complex and the basis of chemokine subfamily selectivity. *PLoS Biol*
18, e3000656 (2020).

22.   M. P. Wescott *et al.*, Signal transmission through the CXC chemokine receptor 4
(CXCR4) transmembrane helices. *Proc Natl Acad Sci U S A* **113**, 9928-9933
(2016).

23.   B. S. Stephens, T. Ngo, I. Kufareva, T. M. Handel, Functional anatomy of the full-
length CXCR4-CXCL12 complex systematically dissected by quantitative model-
guided mutagenesis. *Sci Signal* **13**, eaay5024 (2020).

24.   Y. Liu *et al.*, Cryo-EM structure of monomeric CXCL12-bound CXCR4 in the
active state. *Cell Rep* **43**, 114578 (2024).

25.   K. Saotome *et al.*, Structural insights into CXCR4 modulation and
oligomerization. *Nat Struct Mol Biol* 10.1038/s41594-024-01397-1 (2024).

26. J. M. Burns *et al.*, A novel chemokine receptor for SDF-1 and I-TAC involved in
cell survival, cell adhesion, and tumor development. *J Exp Med* **203**, 2201-2213
(2006).
27. K. R. Klein *et al.*, Decoy receptor CXCR7 modulates adrenomedullin-mediated
cardiac and lymphatic vascular development. *Dev Cell* **30**, 528-540 (2014).
28. Y. Ikeda, H. Kumagai, A. Skach, M. Sato, M. Yanagisawa, Modulation of
circadian glucocorticoid oscillation via adrenal opioid-CXCR7 signaling alters
emotional behavior. *Cell* **155**, 1323-1336 (2013).
29. M. Meyrath *et al.*, The atypical chemokine receptor ACKR3/CXCR7 is a broad-
spectrum scavenger for opioid peptides. *Nat Commun* **11**, 3033 (2020).
30. J. J. Liu, R. Horst, V. Katritch, R. C. Stevens, K. Wuthrich, Biased signaling
pathways in beta2-adrenergic receptor characterized by 19F-NMR. *Science* **335**,
1106-1110 (2012).
31. L. M. Wingler *et al.*, Angiotensin Analogs with Divergent Bias Stabilize Distinct
Receptor Conformations. *Cell* **176**, 468-478 e411 (2019).
32. J. F. Fay, D. L. Farrens, Structural dynamics and energetics underlying allosteric
inactivation of the cannabinoid receptor CB1. *Proc Natl Acad Sci U S A* **112**,
8469-8474 (2015).
33. X. Yao *et al.*, Coupling ligand structure to specific conformational switches in the
beta2-adrenoceptor. *Nat Chem Biol* **2**, 417-422 (2006).
34. M. Elgeti, W. L. Hubbell, DEER Analysis of GPCR Conformational Heterogeneity.
Biomolecules **11**, 778 (2021).
35. L. M. Wingler *et al.*, Angiotensin and biased analogs induce structurally distinct
active conformations within a GPCR. *Science* **367**, 888-892 (2020).
36. A. P. Ray, N. Thakur, N. G. Pour, M. T. Eddy, Dual mechanisms of cholesterol-
GPCR interactions that depend on membrane phospholipid composition.
Structure **31**, 836-847 e836 (2023).
37. C. T. Schafer, J. F. Fay, J. M. Janz, D. L. Farrens, Decay of an active GPCR:
Conformational dynamics govern agonist rebinding and persistence of an active,
yet empty, receptor state. *Proc Natl Acad Sci U S A* **113**, 11961-11966 (2016).
38. G. G. Gregorio *et al.*, Single-molecule analysis of ligand efficacy in beta(2)AR-G-
protein activation. *Nature* **547**, 68-73 (2017).
39. K. Krishna Kumar *et al.*, Negative allosteric modulation of the glucagon receptor
by RAMP2. *Cell* **186**, 1465-1477 e1418 (2023).
40. D. D. Fernandes *et al.*, Ligand modulation of the conformational dynamics of the
A(2A) adenosine receptor revealed by single-molecule fluorescence. *Sci Rep* **11**,
5910 (2021).
41. I. G. Denisov, S. G. Sligar, Nanodiscs for structural and functional studies of
membrane proteins. *Nat Struct Mol Biol* **23**, 481-486 (2016).
42. I. G. Denisov, S. G. Sligar, Nanodiscs in Membrane Biochemistry and
Biophysics. *Chem Rev* **117**, 4669-4713 (2017).
43. S. A. Eberle, M. Gustavsson, Bilayer lipids modulate ligand binding to atypical
chemokine receptor 3. *Structure* **32**, 1174-1183 e1175 (2024).
44. H. Tsukamoto, A. Sinha, M. DeWitt, D. L. Farrens, Monomeric rhodopsin is the
minimal functional unit required for arrestin binding. *J Mol Biol* **399**, 501-511
(2010).
45. S. A. Eberle, M. Gustavsson, A Scintillation Proximity Assay for Real-Time
Kinetic Analysis of Chemokine-Chemokine Receptor Interactions. *Cells* **11**, 1317
(2022).

46. S. G. Rasmussen *et al.*, Crystal structure of the beta2 adrenergic receptor-Gs
protein complex. *Nature* **477**, 549-555 (2011).
47. J. Zhao *et al.*, Ligand efficacy modulates conformational dynamics of the micro-
opioid receptor. *Nature* **629**, 474-480 (2024).
48. K. Palczewski *et al.*, Crystal structure of rhodopsin: A G protein-coupled receptor.
Science **289**, 739-745 (2000).
49. S. G. Rasmussen *et al.*, Crystal structure of the human beta2 adrenergic G-
protein-coupled receptor. *Nature* **450**, 383-387 (2007).
50. D. L. Farrens, C. Altenbach, K. Yang, W. L. Hubbell, H. G. Khorana,
Requirement of rigid-body motion of transmembrane helices for light activation of
rhodopsin. *Science* **274**, 768-770 (1996).
51. A. B. Kleist *et al.*, Conformational selection guides beta-arrestin recruitment at a
biased G protein-coupled receptor. *Science* **377**, 222-228 (2022).
52. O. Otun *et al.*, Conformational dynamics underlying atypical chemokine receptor
3 activation. *Proc Natl Acad Sci U S A* **121**, e2404000121 (2024).
53. C. Perpina-Viciano *et al.*, Kinetic Analysis of the Early Signaling Steps of the
Human Chemokine Receptor CXCR4. *Mol Pharmacol* **98**, 72-87 (2020).
54. C. E. Mona *et al.*, Design, synthesis, and biological evaluation of CXCR4 ligands.
Org Biomol Chem **14**, 10298-10311 (2016).
55. E. M. Rosenberg, Jr. *et al.*, Characterization, Dynamics, and Mechanism of
CXCR4 Antagonists on a Constitutively Active Mutant. *Cell Chem Biol* **26**, 662-
673 e667 (2019).
56. A. Zarca *et al.*, Pharmacological characterization and radiolabeling of VUF15485,
a high-affinity small-molecule agonist for the atypical chemokine receptor
ACKR3. *Mol Pharmacol* **105**, 301-312 (2024).
57. M. Gustavsson, D. P. Dyer, C. Zhao, T. M. Handel, Kinetics of CXCL12 binding
to atypical chemokine receptor 3 reveal a role for the receptor N terminus in
chemokine binding. *Sci Signal* **12**, eaaw3657 (2019).
58. M. Han, S. O. Smith, T. P. Sakmar, Constitutive activation of opsin by mutation of
methionine 257 on transmembrane helix 6. *Biochemistry* **37**, 8253-8261 (1998).
59. X. Han, S. D. Tachado, H. Koziel, W. A. Boisvert, Leu128(3.43) (I128) and
Val247(6.40) (V247) of CXCR1 are critical amino acid residues for g protein
coupling and receptor activation. *PLoS One* **7**, e42765 (2012).
60. M. Cui *et al.*, Crystal structure of a constitutive active mutant of adenosine A(2A)
receptor. *IUCrJ* **9**, 333-341 (2022).
61. H. Tsukamoto, D. L. Farrens, A constitutively activating mutation alters the
dynamics and energetics of a key conformational change in a ligand-free G
protein-coupled receptor. *J Biol Chem* **288**, 28207-28216 (2013).
62. Q. Chen *et al.*, ACKR3-arrestin2/3 complexes reveal molecular consequences of
GRK-dependent barcoding. *bioRxiv* 10.1101/2023.07.18.549504 (2023).
63. X. Deupi *et al.*, Stabilized G protein binding site in the structure of constitutively
active metarhodopsin-II. *Proc Natl Acad Sci U S A* **109**, 119-124 (2012).
64. Z. Shao *et al.*, Molecular insights into ligand recognition and activation of
chemokine receptors CCR2 and CCR3. *Cell Discov* **8**, 44 (2022).
65. L. Qin *et al.*, Structural biology. Crystal structure of the chemokine receptor
CXCR4 in complex with a viral chemokine. *Science* **347**, 1117-1122 (2015).
66. K. Liu *et al.*, Structural basis of CXC chemokine receptor 2 activation and
signalling. *Nature* **585**, 135-140 (2020).

67.   M. Szpakowska *et al.*, Mutational analysis of the extracellular disulphide bridges
of the atypical chemokine receptor ACKR3/CXCR7 uncovers multiple binding
and activation modes for its chemokine and endogenous non-chemokine
agonists. *Biochem Pharmacol* **153**, 299-309 (2018).
68.   L. S. Ai, F. Liao, Mutating the four extracellular cysteines in the chemokine
receptor CCR6 reveals their differing roles in receptor trafficking, ligand binding,
and signaling. *Biochemistry* **41**, 8332-8341 (2002).
69.   C. Limatola *et al.*, Cysteine residues are critical for chemokine receptor CXCR2
functional properties. *Exp Cell Res* **307**, 65-75 (2005).
70.   A. Levoye, K. Balabanian, F. Baleux, F. Bachelerie, B. Lagane, CXCR7
heterodimerizes with CXCR4 and regulates CXCL12-mediated G protein
signaling. *Blood* **113**, 6085-6093 (2009).
71.   D. P. Staus *et al.*, Structure of the M2 muscarinic receptor-beta-arrestin complex
in a lipid nanodisc. *Nature* **579**, 297-302 (2020).
72.   W. Huang *et al.*, Structure of the neurotensin receptor 1 in complex with beta-
arrestin 1. *Nature* **579**, 303-308 (2020).
73.   P. Sarma *et al.*, Molecular insights into intrinsic transducer-coupling bias in the
CXCR4-CXCR7 system. *Nat Commun* **14**, 4808 (2023).
74.   A. Pulvermuller, K. Palczewski, K. P. Hofmann, Interaction between
photoactivated rhodopsin and its kinase: stability and kinetics of complex
formation. *Biochemistry* **32**, 14082-14088 (1993).
75.   R. Paulsen, J. Bentrop, Activation of rhodopsin phosphorylation is triggered by
the lumirhodopsin-metarhodopsin I transition. *Nature* **302**, 417-419 (1983).
76.   T. W. M. De Groof *et al.*, Selective targeting of ligand-dependent and -
independent signaling by GPCR conformation-specific anti-US28 intrabodies. *Nat*
Commun **12**, 4357 (2021).
77.   J. S. Burg *et al.*, Structural biology. Structural basis for chemokine recognition
and activation of a viral G protein-coupled receptor. *Science* **347**, 1113-1117
(2015).
78.   T. F. Miles *et al.*, Viral GPCR US28 can signal in response to chemokine
agonists of nearly unlimited structural degeneracy. *Elife* **7** (2018).
79.   N. Tsutsumi *et al.*, Atypical structural snapshots of human cytomegalovirus
GPCR interactions with host G proteins. *Sci Adv* **8**, eabl5442 (2022).
80.   C. T. Gilliland, C. L. Salanga, T. Kawamura, J. Trejo, T. M. Handel, The
chemokine receptor CCR1 is constitutively active, which leads to G protein-
independent, beta-arrestin-mediated internalization. *J Biol Chem* **288**, 32194-
32210 (2013).
81.   M. Lefrancois *et al.*, Agonists for the Chemokine Receptor CXCR4. *ACS Med*
Chem Lett **2**, 597-602 (2011).
82.   T. K. Ritchie *et al.*, Chapter 11 - Reconstitution of membrane proteins in
phospholipid bilayer nanodiscs. *Methods Enzymol* **464**, 211-231 (2009).
83.   R. F. Pauszek, 3rd, R. Lamichhane, A. Rajkarnikar Singh, D. P. Millar, Single-
molecule view of coordination in a multi-functional DNA polymerase. *Elife* **10**
(2021).
84.   R. Lamichhane, A. Solem, W. Black, D. Rueda, Single-molecule FRET of protein-
nucleic acid and protein-protein complexes: surface passivation and
immobilization. *Methods* **52**, 192-200 (2010).
85.   L. R. Rabiner, A Tutorial on Hidden Markov-Models and Selected Applications in
Speech Recognition. *P Ieee* **77**, 257-286 (1989).

86.   G. Schwarz, Estimating the Dimension of a Model. *Ann. Statist.* **6**, 461-464
(1978).
87.   S. A. McKinney, C. Joo, T. Ha, Analysis of single-molecule FRET trajectories
using hidden Markov modeling. *Biophys J* **91**, 1941-1951 (2006).

883
884

885 **Figures and Tables**
886
887

888
Figure 1. Experimental design of the smFRET system. A) A single receptor molecule
(blue) was labeled with donor (D) and acceptor (A) fluorophores, inserted into a
phospholipid (yellow) nanodisc (green), and immobilized on a quartz slide via biotin (tan
circle)-neutravidin (grey rectangle) attachment. A prism facilitates total internal reflection
of the excitation laser to excite only donor fluorophores close to the surface. B) Cartoon
depicting inactive (left) and active (right) receptor conformations. C) Two representative
single-molecule time traces for apo-ACKR3. In both examples, the donor (green) and
acceptor (red) intensities are shown in the top panel and the corresponding apparent
FRET efficiency (black) is shown in the bottom panel.
898
899
900

Figure 2. ACKR3 exhibits greater conformational flexibility compared to CXCR4. A) Apparent FRET efficiency histogram of apo-CXCR4 (left, black trace) resolved into three distinct conformational states: a high-FRET state corresponding to the inactive receptor conformation (R, blue), a low-FRET active receptor conformation (R*, red) and an intermediate conformation (R', gray). The fractional populations of each state obtained from global analysis are indicated. The receptor is mostly in the inactive conformation. A transition density probability (TDP) plot (right) displays the relative probabilities of transitions from an initial FRET state (x-axis) to a final FRET state (y-axis). For apo-CXCR4, transitions between R and R' states are observed most frequently. B) Addition of CXCL12$_{WT}$ to CXCR4 shifted the conformational distribution to the low-FRET R* state and resulted in more transitions between all three FRET states. C) The broad apparent FRET efficiency histogram of apo-ACKR3 (left, black trace) is resolved into four distinct conformational states: inactive R (blue), active R* (red), inactive-like R' (light blue), and active-like R*' (pink). Little conformational preference is observed among these states. Moreover, all possible sequential state-to-state transitions are observed (right). D) Addition of CXCL12$_{WT}$ to ACKR3 shifted the conformational distribution to the low-FRET R* state, which was also reflected in the transition probabilities. In all cases, data sets represent the analysis of at least three independent experiments.

Figure 3. A small molecule inhibitor shifts the ACKR3 conformational population to the inactive FRET state, while CXCR4 is largely unaffected. A) FRET distributions and TDP of apo-CXCR4 repeated from Fig. 2A for comparison. B) Treatment of CXCR4 with the inhibitor IT1t had little impact on the FRET distribution, but increased transition probabilities compared to the apo-receptor. C) FRET distributions and TDP of apo-ACKR3 repeated from Fig. 2C for comparison. D) Treatment of ACKR3 with VUF16480, an inverse agonist, shifted the conformational distribution and TDP to the high-FRET inactive R conformation. Data sets represent the analysis of at least three independent experiments. The overall apparent FRET efficiency envelopes for the samples are represented by the black traces.

939
Figure 4. CXCL12 variants containing mutations to the N-terminus promoted active receptor conformations in both CXCR4 and ACKR3. A) FRET distributions and TDP for apo-CXCR4 repeated from Fig. 2A for reference. B) Addition of CXCL12$_{P2G}$ to CXCR4 promoted a shift to the low-FRET active (R*) conformation and an increase in state-to-state transition probabilities. C) CXCL12$_{LRHQ}$ led to a more subtle shift to the R* conformation of CXCR4 without affecting the transition probabilities. D) FRET distributions and TDP for apo-ACKR3 repeated from Fig. 2C. E) Treatment of ACKR3 with CXCL12$_{P2G}$ displayed a shift to low-FRET, R*' and R* states, while reducing the transition probabilities for R' ↔ R*' and R*' ↔ R* transitions relative to the apo-receptor. F) CXCL12$_{LRHQ}$ treatment of ACKR3 shifted the FRET distribution to the low-FRET R* active state and promoted R*' ↔ R* transitions relative to the apo-receptor. In all cases, the data sets represent the analysis of at least three independent experiments. The overall apparent FRET efficiency envelopes for the samples are represented by the black traces.

Figure 5. Replacement of Y257⁶·⁴⁰ with the corresponding residue in CXCR4 (leucine)
reduces conformational heterogeneity of ACKR3. A) Structure of ACKR3 bound with
CXCL12$_{WT}$ (PDBID: 7SK3) highlighting the location of Y257⁶·⁴⁰ (purple) (16). B) Apparent
FRET efficiency distributions and TDP of WT ACKR3 in the apo-state, repeated from Fig.
2C. C) The mutation Y257⁶·⁴⁰L shifted the conformational landscape of the apo-receptor
to the high-FRET inactive R conformation at the expense of active R* and active-like R*'
conformations, and also reduced the probability of state-to-state transitions. D) Apparent
FRET efficiency distributions and TDP of WT ACKR3 treated with CXCL12$_{WT}$, repeated
from Fig. 2D. E) Treatment of Y257⁶·⁴⁰L ACKR3 with CXCL12$_{WT}$ promoted more low-FRET
active R* and active-like R*' states. Data sets represent the analysis of at least three
independent experiments. The overall apparent FRET efficiency envelopes for the
samples are represented by the black traces.

Figure 6. Schematic illustration of the conformational energy landscapes for CXCR4 and ACKR3, highlighting the differences in the responsiveness of the two receptors to ligands. CXCR4 populates three distinct conformations, shown here as wells on the energy landscape. Apo-CXCR4 is predominantly in the inactive R state. The receptor is converted incompletely to R* with CXCL12$_{WT}$ treatment, while the small molecule inhibitor IT1t has little impact on the conformational distribution. Though CXCL12$_{P2G}$ is an antagonist of CXCR4, the ligand promoted a detectable shift to the active R* state, suggesting TM6 movement is not sufficient for CXCR4 activation. In contrast, ACKR3 populates four distinct conformations and shows little preference among them in the apo-form. The inverse agonist, VUF16840, shifts the population to the inactive R conformation, while the agonists CXCL12$_{WT}$ and CXCL12$_{P2G}$ promote the R* and R*' populations of ACKR3. Despite stabilizing different levels of the active R* state and active-like intermediate R*' state, both CXCL12$_{WT}$ and CXCL12$_{P2G}$ are agonists of ACKR3. The flexibility of ACKR3 may contribute to the ligand-promiscuity of this atypical receptor. Figure created with biorender.com