Functional characterization of the disease-associated *CCL2* rs1024611G-rs13900T

haplotype: The role of the RNA-binding protein HuR

Feroz Akhtar[1], Joselin Hernandez Ruiz[2][¶], Ya-Guang Liu[3], Roy G. Resendez[1], Denis Feliers[4][‡],

Liza D. Morales[5], Alvaro Diaz-Badillo[1], Donna M. Lehman[1], Rector Arya[1], Juan Carlos Lopez-

Alvarenga[6], John Blangero[5], Ravindranath Duggirala[1], and Srinivas Mummidi[1]*

[1]Department of Health and Behavioral Sciences, Texas A&M University- San Antonio, Texas,

USA, [2] Utah Center for Genetic Discovery, Department of Human Genetics, University of Utah,

Salt Lake City, Utah, USA, [3]Department of Pathology, School of Medicine, University of Texas

Health San Antonio, San Antonio, Texas, USA, [4]Department of Medicine, School of Medicine,

University of Texas Health San Antonio, San Antonio, Texas, USA, [5] South Texas Diabetes

and Obesity Institute, Department of Genetics, School of Medicine, University of Texas Rio

Grane Valley, Brownsville, USA, [6]Department of Population Health and Biostatistics, School

of Medicine, University of Texas Rio Grande Valley, Harlingen, Texas, USA,

[‡]Current Address: Novartis Institute for Biomedical Research San Diego, San Diego, CA.

* **CORRESPONDING AUTHOR:**

Srinivas Mummidi, D.V.M., Ph.D.

Email: srinivas.mummidi@tamusa.edu

Abstract

CC-chemokine ligand 2 (CCL2) is involved in the pathogenesis of several diseases associated

with monocyte/macrophage recruitment, such as HIV-associated neurocognitive disorder

(HAND), tuberculosis, and atherosclerosis. The rs1024611 (alleles:A>G; G is the risk allele)

polymorphism in the *CCL2 cis*-regulatory region is associated with increased CCL2

expression in vitro and ex vivo, leukocyte mobilization in vivo, and deleterious disease

outcomes. However, the molecular basis for the rs1024611-associated differential CCL2

expression remains poorly characterized. It is conceivable that genetic variant(s) in linkage

disequilibrium (LD) with rs1024611 could mediate such effects. Previously, we used rs13900

(alleles: C>T) in the *CCL2* 3' untranslated region (3' UTR) that is in perfect LD with rs1024611

to demonstrate allelic expression imbalance (AEI) of *CCL2* in heterozygous individuals. Here

we tested the hypothesis that the rs13900 could modulate *CCL2* expression by altering mRNA

turnover and/or translatability. The rs13900 T allele conferred greater stability to the *CCL2*

transcript when compared to the rs13900 C allele. The rs13900 T allele also had increased

binding to Human Antigen R (HuR), an RNA-binding protein, in vitro and ex vivo. The rs13900

alleles imparted differential activity to reporter vectors and influenced the translatability of the

reporter transcript. We further demonstrated a role for HuR in mediating allele-specific effects

on CCL2 expression in overexpression and silencing studies. The presence of the

rs1024611G-rs13900T conferred a distinct transcriptomic signature related to inflammation

and immunity. Our studies suggest that the differential interactions of HuR with rs13900 could

modulate CCL2 expression and explain the interindividual differences in CCL2-mediated

disease susceptibility.

Introduction

Identification of functional and/or causal genetic variants continues to pose a significant challenge in the post-genome-wide association studies (post-GWAS) era(MacArthur et al., 2017; Tam et al., 2019; Visscher et al., 2017). While emphasis has been placed on polymorphisms that map to *cis*-elements in enhancers and promoters, genetic variants that disrupt cis-elements in RNA binding protein (RBP) motifs in 3' UTR have received much less attention. For many genes, the interactions of their 3' UTRs with specific stabilizing and destabilizing RBPs play a critical role in modulating post-transcriptional events such as mRNA turnover and translatability (Mayr, 2019; Schwerk and Savan, 2015). The lack of a mechanistic understanding of how SNPs localizing to RBP motifs could impact gene expression impedes a greater understanding of the variability in disease susceptibility and outcomes.

Post-transcriptional mechanisms are thought to play a crucial role in the initiation and resolution of the inflammatory response(Anderson, 2010; Yoshinaga and Takeuchi, 2019). Such regulation can profoundly affect gene expression levels; modest changes in mRNA stability can lead to significant effects on mRNA and protein abundance(Buccitelli and Selbach, 2020; Ross, 1995). Notably, in a recent genome-scale study using mouse dendritic cells, post-transcriptional mRNA degradation was a salient feature of inflammatory and immune signaling genes, as well as targets of NF-kappaB signaling following lipopolysaccharide (LPS) stimulation(Rabani et al., 2011). In addition, polymorphisms in non-coding regions such as the 3' UTR can have a significant impact on mRNA stability and translatability. For example, a genome-wide study of variation in gene-specific mRNA decays in lymphoblastoid cell lines across individuals found about 195 genetic variants that are specifically associated with variation in mRNA decay rates, called "rdQTLs" (RNA Decay Quantitative Trait Loci)(Pai et al., 2012). The authors estimated that 35% of the most significant eQTL (Expression Quantitative Trait Loci) single nucleotide polymorphisms (SNPs) are associated with decay rates(Pai et al., 2012). In another study, Duan *et al.* reported that ~37% of gene expression differences among individuals may be attributed to RNA half-life

differences(Duan et al., 2013). Farh et al., reported that the 3' UTRs are highly enriched for

eQTL candidate causal SNPs (>1,500) relative to other transcribed SNPs(Farh et al., 2015).

However, the molecular mechanisms by which these polymorphisms alter mRNA stability or

translatability are poorly understood.

CCL2 is a potent monocyte chemoattractant produced by various cell types, either

constitutively or following activation. CCL2 expression can be regulated by inflammatory

molecules (e.g., IL-1, TNFα, LPS, and IFNγ) and growth factors (e.g., PDGF). While the cells

of monocyte-macrophage lineage are a major source of CCL2(Yoshimura et al., 1989), other

cell types, such as fibroblasts, astrocytes, epithelial, and endothelial cells, also are an

important source(Deshmane et al., 2009). CCL2 mediates recruitment of monocytes, memory

T-cells, and dendritic cells to the site of inflammation(Gschwandtner, Derler and Midwood,

2019; Melgarejo et al., 2009) and there is substantial evidence implicating CCL2 as a key

mediator in macrophage-mediated diseases. Rovin et al. described a SNP in the 5'-regulatory

region of *CCL2* annotated as rs1024611 (dbSNP database; originally designated as –

2518A>G(Rovin, Lu and Saxena, 1999) or –2578A>G(Gonzalez et al., 2002)) that was

associated with increased plasma CCL2 expression(Rovin, Lu and Saxena, 1999). This SNP

is associated with increased serum CCL2 levels, enhanced macrophage recruitment to

tissues, and progression to HIV-associated dementia(Gonzalez et al., 2002). Other studies

showed that the rs1024611 G allele is associated with increased CCL2 levels in the plasma,

urine, and cerebrospinal fluid in health and disease, and in tissues such as liver and skin(Cho

et al., 2004; Fenoglio et al., 2004; Joven et al., 2006; Letendre et al., 2004; McDermott et al.,

2005). The rs1024611 polymorphism has been associated with several diseases, including

myocardial infarction(McDermott et al., 2005), carotid atherosclerosis(Alonso-Villaverde et al.,

2004), pulmonary tuberculosis(Flores-Villanueva et al., 2005), severe acute

pancreatitis(Cavestro et al., 2010), lupus nephritis(Tucci et al., 2004), asthma susceptibility

and severity(Szalai et al., 2001), Crohn's disease (CD)(Palmieri et al., 2010), and Alzheimer's

disease(Fenoglio et al., 2004), and infections by Japanese Encephalitis virus(Chowdhury and

Khan, 2017) and SARS-CoV-1(Tu et al., 2015). Notably, rs3091315, a GWAS risk variant for

CD(Franke et al., 2010) and inflammatory bowel disease (IBD)(Liu et al., 2015) is in a high

linkage disequilibrium (LD) with both rs1024611 (D′=1.0, r^2=0.98) and rs13900 (D′=1.0,

r^2=0.98) in the CEU population. Given the importance of disease associations with rs1024611,

significant efforts have been made by other groups and us to understand the molecular basis

of this differential CCL2 expression associated with this polymorphism(Gonzalez et al., 2002;

Mummidi, Bonello and Ahuja, 2009; Page et al., 2011; Pham et al., 2012; Wright et al., 2008).

However, these studies did not provide a mechanistic link between the rs1024611

polymorphism and CCL2 expression, giving rise to the possibility that SNPs in strong LD with

rs1024611 could be mediating these effects.

To identify potential functional SNPs that could explain the variability in CCL2 expression, we

developed an extensive LD map of the *CCL2* genomic locus and reported that a SNP

designated as the rs13900 (NM_002982.4:c.*65=) in the *CCL2* 3' UTR is in perfect LD with

rs1024611 and can serve as its proxy(Pham et al., 2012). We further showed that *CCL2* is

subjected to allelic expression imbalance (AEI), with the expression of the rs13900 T allele

relatively higher when compared to the rs13900 C allele(Pham et al., 2012). In this study, we

show that the differential binding of Human Antigen R (HuR), an RBP previously implicated in

CCL2 expression, leads to altered stability and translatability of *CCL2* transcripts and MDM

with rs13900 T allele have a distinct transcriptomic signature providing a mechanistic

explanation for increased CCL2 expression in individuals with the rs1024611G-rs13900T

haplotype and inter-individual differences in disease susceptibility associated with this

haplotype.

Results

Individuals heterozygous for rs13900 show AEI of *CCL2*

We and others have previously reported a perfect linkage disequilibrium between rs1024611 in the *CCL2* cis-regulatory region and rs13900 in its 3′ UTR and that rs13900 can serve as a proxy for the disease-associated rs1024611. We further showed that *CCL2* exhibits allelic expression imbalance (AEI) in heterozygous individuals, with the rs13900 T allele (alternative allele) having a higher expression than the rs13900 C allele (reference allele) (Figure 1A). For this study, we recruited 47 healthy unrelated individuals (18–35-year-old) who were screened for rs13900 (Figure S1A and B). Supplementary Table 1 shows the genotype and allele frequencies for rs13900 polymorphism. We found that the rs13900 C allele was at a higher frequency than the rs13900 T allele and the genotype frequencies were in line with Hardy-Weinberg equilibrium ($P \geq 0.05$). Subsequent experiments were conducted on 15 homozygous and heterozygous individuals. We reconfirmed that *CCL2* exhibits AEI using data from heterozygous individuals. For this we used total RNA obtained from LPS treated PBMC as described previously(Pham et al., 2012). LPS induced *CCL2* expression in PBMCs as confirmed by qRT-PCR, with about 4.3-fold increase at 1 h, 6.09-fold increase at 3 h, and 1.94-fold increase at 6 h (Fig. S2). Given the high expression of *CCL2* transcript after a 3 h treatment with LPS, we used this time point for all subsequent experiments. AEI was measured by quantifying the relative amount of the two alleles i.e., alternative allele (T) to reference allele (C), measured from the chromatogram after normalization of peak intensity using PeakPeaker v.2.0 (Figure 1B and 1C). This strategy ensures a direct comparison between the amount of *CCL2* mRNA that is transcribed from each allele or haplotype and that each allele is equally subjected the effects of any external factors. gDNA was utilized as a control. The boxplot illustrates a notable difference in the detected levels of the C and T allele in the gDNA and cDNA with a higher expression of T allele relative to C allele ($P < 0.005$) (Figure 1D).

CCL2 mRNA transcripts bearing rs13900 C and T alleles have different stability

Previous studies have shown that _CCL2_ mRNA is subjected to post-transcriptional regulation

through modulation of mRNA stability(Hao and Baltimore, 2009). Here, we determined

whether the rs13900 modulates mRNA stability which may in part explain AEI. We used

purified monocytes from four heterozygous individuals that were left untreated or stimulated

with LPS for 3 h. Cells were either harvested after 3 h of LPS stimulation (considered as t=0)

or cultured in the presence or absence of the transcriptional inhibitor actinomycin D for an

additional 1, 2, or 4 h. Total RNA was isolated at each time point to assess _CCL2_ transcript,

calculated as fold induction over unstimulated cells. _CCL2_ mRNA levels showed a strong

upregulation following LPS stimulation ($P < 0.05$) (Figure 2A). Actinomycin D treatment

revealed the kinetics of _CCL2_ transcript degradation by determining the mRNA half-life, which

represents the time (expressed in hours) at which mRNA expression is 50% of the initial level

(Figure 2B; $t_{1/2}$=Ln (0.5)/slope). For _CCL2_, $t_{1/2}$=1.763 h and is in line with previously published

studies that _CCL2_ mRNA stability is modulated following inflammatory and cytokine

stimuli(Hao and Baltimore, 2009; Zhai et al., 2008).

To rule out the confounding effects of preexisting mRNA, the relative stability of rs13900 C-

and T-allele bearing transcripts in heterozygous individuals were evaluated using nascent

RNA. Using nascent RNA allows accurate determination of mRNA decay by eliminating the

effects of preexisting mRNA. Briefly, nascent RNA was obtained from LPS stimulated

monocytes cultured in the presence of actinomycin D and ethylene uridine (EU). The EU RNA

was then subjected to a click reaction. The click reaction adds a biotin handle to nascent RNA

which is then captured by streptavidin beads. cDNA was synthesized from the captured

nascent RNA, PCR-amplified and expression of the individual alleles was assessed as

described above. Data from three individuals showed that the transcript bearing rs13900 T

allele was enriched relative to rs13900 C allele in the nascent transcripts isolated from ActD-

treated cells ($P < 0.05$) after 4 h (Figure 2C).

Bioinformatic analyses of rs13900

While previous experimental studies showed that *CCL2* 3′ UTR binds HuR, it is not known whether rs13900 disrupts or alters HuR binding(Lebedeva et al., 2011). Therefore, we analyzed the rs13900 flanking region using various bioinformatic software to mine existing whole genome datasets (e.g., PAR-CLIP datasets) and to predict any mRNA structural changes and altered RBP motifs. We used AURA to examine the colocalization between rs13900 and HuR (Figure 3A). As genetic variants can also alter RBP binding in mRNA, we used ViennaRNA package to predict changes in the *CCL2* secondary structure due to rs13900. As shown in Figure 3B, the presence of rs13900 T allele is predicted to alter the *CCL2* transcript secondary structure (red arrow). The POSTAR3 suite also incorporates HOMER(Heinz et al., 2010) for motif analysis and RNAcontext(Kazan et al., 2010) that identifies not only known but also predicts relative binding and structural preferences of RBPs. HOMER motif analysis(Heinz et al., 2010) identified a HuR binding motif in the region flanking the rs13900 (Fig. 3C). Figure 3D shows the relative structural preference of HuR to different structural contexts identified by RNAcontext, where the alphabets P, L, U, M indicate that the nucleotide is paired (P), in a hairpin loop (L), in an unstructured (or external) region (U), or Miscellaneous (M). The M category includes various unpaired contexts such as nucleotide localizing to a bulge, internal loop or multiloop. The rs13900 C allele to rs13900 T allele transition is predicted to form a stem (Figure 3B) which is predicted to increase HuR binding. Analysis using RBP-Var2 predicted that the rs13900 is likely to affect RBP binding, RNA secondary structure and expression and assigned it a score of 1e. Taken together, our bioinformatic analyses suggested that the rs13900 allele could potentially alter the binding of HuR to *CCL2* 3′ UTR.

Differential binding of HuR to rs13900 C and T alleles *in vitro*

To experimentally verify our bioinformatic findings on rs13900, we utilized REMSA to determine whether the region of *CCL2* 3′ UTR that flanks rs13900 binds *in vitro* to HuR and if

there are allelic differences in binding. Purified labeled single-stranded oligoribonucleotides

bearing either rs13900 C or rs13900 T alleles were incubated with 10 μg of whole cell extracts.

We tested whether the bound complexes contained HuR by performing antibody mediated

supershift assays. As shown in Figure 4A and 4B, a predominant shift was observed in the

presence of rs13900 T allele (lane 8). Notably, there was an approximately 7-fold difference

($P < 0.005$) in oligoribonucleotide/HuR/antibody complexes generated with T allele and those

generated with rs13900 C allele (lane 4). The specificity of the complex formation was

confirmed by using a non-specific antibody (Figure 4A, lanes 3 & 7). To rule out the possibility

that additional RBPs are involved in this complex formation, we used purified HuR protein in

mobility shift assays. These studies with purified protein further confirmed that

oligoribonucleotides with rs13900 C or T alleles differentially bound HuR (Figure 4C). Taken

together, these studies provided experimental validation that the rs13900 may influence

differences in binding affinity to HuR (Figure 4D).

Differential binding of HuR to rs13900 C and T alleles *ex vivo*

We next tested the hypothesis that the rs13900 is associated with altered binding affinity to

HuR *ex vivo*. For this, we performed RNA immunoprecipitation in monocyte/macrophages

derived from four heterozygous individuals. Immunoprecipitations were performed using

cytoplasmic lysates of macrophages treated with LPS using an affinity purified HuR antibody

or IgG. Figure 5A shows the relative enrichment of HuR in the immunoprecipitated fraction

compared to the IgG control. The HuR immunoprecipitated fraction showed significant

enrichment ($P < 0.05$) of the region encompassing rs13900 when compared to the IgG control

(Figure 5B-C). *CCL2* transcript was enriched ~10-fold in the HuR immunoprecipitated fraction

in comparison to the IgG bound fraction (Figure 5C). Notably, transcripts containing rs13900

T allele were enriched relative to transcripts containing rs13900 C allele ($P < 0.05$) in the

anti-HuR RIP complexes (Figure 5D).

rs13900 T allele confers increased mRNA stability in reporter assays

As HuR is implicated in mRNA stability of many mRNA transcripts including *CCL2*, we tested the hypothesis that *CCL2* 3′ UTR influences its stability and that rs13900 modulates this effect. We constructed reporter plasmids that harbor the *CCL2* 3′ UTR containing either rs13900 C or rs13900 T allele (Figure 6A) and nucleofected them into HEK293 cells as described in Materials and Methods. Luciferase activity was measured after 24 h post-transfection (Figure 6B). Our results indicated that the presence of *CCL2* 3′ UTR significantly ($P < 0.05$) reduced the luciferase activity. However, cells transfected with plasmids bearing the rs13900 T allele showed higher luciferase activity when compared with cells transfected with plasmids bearing the rs13900 C allele suggesting that the presence of rs13900 T allele conferred increased stability to the transcript. We next analyzed the influence of HuR overexpression in reporter vectors containing *CCL2* 3′ UTR with either rs13900 C or T alleles. Overexpression of HuR led to a significant increase in luciferase activity of the reporter vector bearing rs13900 T allele ($P < 0.05$), However, HuR overexpression had no significant effect on the luciferase activity of the reporter vector bearing the rs13900 C allele (Figure 6C). Conversely, we examined the effect of HuR on the expression of the reporter assay by co-transfection of HuR siRNA and luciferase reporter constructs. While HuR knockdown had no effect on luciferase activity of the reporter construct bearing rs13900 C allele, it caused a significant reduction in luciferase activity of construct bearing the rs13900 T allele ($P < 0.05$) (Figure 6D). The overexpression and knockdown of the HuR in the nucleofected cells was confirmed using Western blots (Figure S3).

Role of HuR-rs13900 interactions in *CCL2* mRNA translatability

The differential interactions with HuR by rs13900 C and rs13900 T alleles could potentially alter *CCL2* mRNA translatability as HuR could increase mRNA translation(Schultz et al., 2020). Therefore, we assessed the relative allelic enrichment in the monosomal and polysomal fractions obtained from MDMs from two individuals exhibiting AEI. Translationally active and inactive pools of RNA were fractionated by isolating the monosomal and polysomal fractions

from MDMs on sucrose gradients after the cells were treated with cycloheximide to block

translation. The distribution of *CCL2* mRNA in macrophages cultured in presence or absence

of LPS for 3 h is depicted in Supplementary Figure S4 panels A and B. As previously reported

by others, LPS treatment resulted in a distinct shift of the mRNAs from monosomal fraction to

the polysomal fraction (Schott et al., 2014). Consistent with this prior report, 60.4% of the

CCL2 mRNA was associated with polysomal fraction following stimulation of cells with LPS.

We assessed the differences in allelic enrichment in cytosolic, monosomal, and polysomic

fractions and found that the polysomal fractions showed the enrichment of the rs13900 T allele

(Supplementary Table 2). However, this differential loading of rs13900 T allele was noted for

cytosolic and monosomal fractions.

To further address this question, we used a reporter-based system to assess the effect of

rs13900 C and T alleles on the translatability. To measure translatability, luciferase mRNA

and protein were measured simultaneously, and translatability was calculated as luciferase

activity normalized by the luciferase mRNA levels after adjusting for protein and 18S rRNA

(Figure 7A-C). Our results suggest that the rs13900 may alter the mRNA translatability in

addition to the transcript stability.

Differential effect of HuR over-expression on the *CCL2* rs13900 T allele

Our *in vitro* and *ex vivo* data indicate that HuR positively regulates *CCL2* transcript stability by

increased binding to the T allele. To determine a direct functional relationship between HuR

and *CCL2* AEI, we used a lentiviral overexpression system. We either transduced HuR specific

(pCMV6-HuR) or non-specific control shRNAs (CMV-null) into the monocytes obtained from

donors who were either homozygous for rs13900 C or T allele. Ready to use GFP-tagged

pCMV6-HuR or CMV-null lentiviral particles were transduced into macrophages in presence

of polybrene at an MOI of 1. Cells were processed 72 h following virus addition for the analysis

of HuR and *CCL2* mRNA expression. Using this lentiviral system, both high transduction

efficiency (90%) and high expression levels were achieved in primary human macrophages

(Figure S5 A-B). Compared with CMV-Null particles, substantial overexpression of HuR was

noted with pCMV6-HuR lentiviral particles (Figure 8A). HuR overexpression was associated

with a higher expression of *CCL2* mRNA in persons homozygous for T allele relative to those

who were homozygous for C allele (Figure 8B).

Discussion

Genome-wide association studies (GWASs) have led to the discovery of numerous disease

susceptibility genetic variants/genes and biological pathways involved in specific diseases,

including many with immunological etiologies(Buniello et al., 2019; Manolio, 2010; Visscher et

al., 2017). Most of the disease-associated genetic variants identified are present in the non-

coding regions of the genome(Maurano et al., 2012; Zhang and Lupski, 2015; Zou et al.,

2019). However, characterizing the functional consequence of these genetic variants in the

regulatory regions remains a significant challenge to date. About 3.7% of the non-coding

variants localize to the untranslated regions(Steri et al., 2018), suggesting post-transcriptional

mechanisms such as mRNA stability and translatability could determine disease

susceptibility(Maurano et al., 2012).

The *CCL2* locus exemplifies the difficulty in dissecting the functional consequences of cis-

regulatory and non-coding genetic variants. Our analysis of regulatory elements in an ~20 kb

region upstream of the human *CCL2* coding region identified highly conserved enhancers and

other regulatory elements in the *CCL2* locus(Bonello et al., 2011). We identified SNPs with

strong LD in this "super-enhancer" and determined their impact on transcriptional activity,

which was minimal(Pham et al., 2012). For example, we tested the role of rs7210316 and

rs9889296, which had an $r^2 > 0.9$ with rs1024611 in CEU population and were located ~6.3

kb and ~9.2 kb upstream of it(Pham et al., 2012). The genetic regions that encompass and

flank these two SNPs either had no or minimal transcriptional activity and lacked the epigenetic

markers that have been traditionally associated with gene activation. Additionally, conflicting

context-dependent results have been obtained for the transcriptional activities associated with

reporter vectors containing rs1024611 A and G alleles, which may have been due to the use

of different lengths of the *CCL2* cis-regulatory regions employed in the transcriptional assays.

To avoid such context-dependent confounding, we used chromatin annotation, a powerful

approach to identify functional SNPs, to generate reporter constructs that span a 6 kb cis-

regulatory region(Pham et al., 2012). This approach allowed us to directly compare the roles

of four correlated SNPs (rs1860190, rs2857654, rs1024611, and rs2857656) on *CCL2*

transcriptional activity. We found no significant differences in transcription strength between

these two constructs when transfected into primary human astrocytes (Pham et al., 2012).

Furthermore, to understand the basis for increased CCL2 expression associated with the

rs1024611 G allele, we and others have demonstrated that several transcription factors bind

differentially to *CCL2* polymorphisms, including IRF-1, PARP-1, STAT-1, Prep1/Pbx

complexes(Gonzalez et al., 2002; Mummidi, Bonello and Ahuja, 2009; Page et al., 2011;

Wright et al., 2008). However, these previous studies could not unequivocally demonstrate

the mechanistic link between differential binding of transcription factor and transcriptional

activity and the physiological relevance of implicated transcription factor in CCL2 expression.

Our comprehensive studies outlined here demonstrate that rs13900 T allele differentially binds

to the RBP HuR, which could alter CCL2 mRNA stability and gene expression. Furthermore,

we provide multiple lines of functional evidence for rs13900/HuR role in *CCL2* AEI, including

reporter vector assays and HuR overexpression and depletion experiments.

Our previous finding that rs13900 is in perfect LD with rs1024611 has allowed us to exploit

the powerful AEI technique and mechanistically link the *CCL2* rs1024611G-rs13900T

haplotype to increased expression(Pham et al., 2012). These findings, together with previous

knowledge that *CCL2* is subjected to post-transcriptional regulation(Hao and Baltimore, 2009)

and regulated by RBP(Fan et al., 2011), raised the possibility that rs13900 could be functional

and impact mRNA stability and translatability. In this study, we determined the half-life of *CCL2*

to be 1.76 h, reinforcing the fact that post-transcriptional regulation is a key feature of *CCL2*

expression (Figure 2B). Since transcription and RNA degradation are tightly linked, and

transcriptional inhibition may lead to mRNA decay, we assessed the allelic differences in

expression in nascent RNA obtained from heterozygous individuals. We found that the *CCL2*

transcripts with rs13900 T allele have increased stability relative to those with rs13900 C allele

(Figure 2C). In addition, both rs1024611 and rs13900 have suggestive associations with *CCL2*

expression in *cis*-expression quantitative trait loci (eQTL) analyses. For example, rs1024611

and rs13900 are associated with CCL2 expression in MDM infected with *Listeria*

monocytogenes(Nedelec et al., 2016) ($-\log10(p) = 3.19$; Effect size = 0.29±0.084)(Kwong et

al., 2021). Remarkably, rs13900 is also identified as a cis-splicing quantitative trait locus (cis-

sQTL) for *CCL2* in both untreated and treated monocytes(Alasoo et al., 2018; Kwong et al.,

2021; Quach et al., 2016). Recent evidence suggests that genetic variation influencing RNA

splicing could play an important role in determining complex phenotypic traits(Li et al., 2016).

While rs13900 serves as an sQTL for several *CCL2* transcripts variants, we provide an

illustrative example here. rs13900 showed significant associations with the canonical *CCL2*

transcript, ENST00000225831 in LPS treated monocytes ($-\log10(p) = 9.88$; Effect size =

0.57±0.084) and with a CCL2 transcript ENST00000580907 ($-\log10(p) = 9.88$; Effect size =

-0.57±0.084) that encodes a truncated CCL2 protein(Kwong et al., 2021). While the

importance of these associations needs more in-depth studies, our observation that *CCL2*

transcripts with rs13900 T allele have a slower degradation could explain the increased CCL2

expression in individuals with rs1024611G-rs13900 haplotype as modest changes in mRNA

stability can lead to significant effects on mRNA and protein abundance.

Post-transcriptional gene expression is regulated through interactions between the *cis*-

elements in mRNAs and their cognate RBPs(Wu and Brewer, 2012). Studies indicate

presence of post-transcriptional operons or regulons – unique subsets of RNAs that associate

with RBPs, which coordinate their localization, translation, and degradation. Our

bioinformatics analysis and experimental data confirm HuR binds to the region spanning the

rs13900. iClip data from HeLa cells(Wang et al., 2010), suggested that the RBPs TIAL1 (T cell

intracellular antigen-1 like protein) and U2AF65 (U2 snRNP auxiliary factor large subunit)

could also bind to this region of the *CCL2* transcript(Mao et al., 2016). TIAL1 has been

proposed to act as a cellular sensor and has been associated with a transcriptome associated

with control of inflammation, cell-cell signaling, immune-suppression, angiogenesis,

metabolism and cell proliferation(Reyes, Alcalde and Izquierdo, 2009). The role of TIAL1 in

CCL2 expression is not known and additional studies are needed to determine if its binding

contributes to *CCL2* AEI. U2AF65 is a widely expressed splicing factor that associates with

RNA polymerase II to bind upstream 3′ splice sites to facilitate splice site pairing in higher

eukaryotes(Hollander et al., 2016). In addition, a change in RNA structure due to a SNP could

indirectly alter accessibility of additional regions to RBPs and further studies are required to

determine any such changes(Shatoff and Bundschuh, 2020).

The Hu family contains 4 members, of which HuR is ubiquitously distributed. HuR is reported

to bind variably sized hairpin loops rich in uracil(Uren et al., 2011). HuR shuttles between the

nucleus and cytoplasm and is likely involved in the transport and stabilization of mRNA. HuR

action is antagonized by RNA destabilizing proteins such as TTP (tristetraprolin) and AUF

(ARE/poly(U)-binding/degradation factor 1)(Wu and Brewer, 2012). The role of HuR in

chemokine gene regulation has been extensively studied in airway epithelium. Fan *et al*.

identified that *CCL2* is one of the top targets for HuR(Fan et al., 2011). They demonstrated

that HuR associated with *CCL2* 3′ UTR *in vitro* and *CCL2* expression could be modulated by

changes in HuR levels. Also, HuR levels influenced *CCL2* mRNA decay. In a mouse model of

alcoholic liver disease, HuR was found to play a key role in NOX4 mediated increase in *CCL2*

mRNA stability(Sasaki et al., 2017). Notably, HuR has also been implicated in mRNA stability

of CX3CL1(Matsumiya et al., 2010) and IL-8(Choi et al., 2009) among others. A recent study

showed that CCL2 itself can induce nuclear to cytoplasmic translocation of HuR and stabilize

vascular endothelial growth factor-A (*VEGFA*) mRNA in CD14[+]CD16[low] inflammatory

monocytes, thus raising the possibility that CCL2 may stabilize its own message in an

autocrine fashion(Morrison et al., 2014). These studies further bolster a role for HuR in *CCL2*

expression and support our finding that differential binding to HuR alters its expression level.

While there are several examples of genetic variants that influence mRNA stability(Akdeli et

al., 2014; Duan et al., 2013; Vilmundarson et al., 2021; Wang, Pitarque and Ingelman-

Sundberg, 2006), very few studies have examined the role of RBPs such as HuR in

differentially influencing gene expression levels in humans(Steri et al., 2018). A notable

exception is the report showing that RBP AUF1 regulates the allele-specific stability of

thymidylate synthase(Pullmann et al., 2006). Another study showed that a type 2 diabetes

associated polymorphism in the 3′ UTR of *PPP1R3* alters distance between two ARE motifs

and results in differential binding of protein complexes and may be associated with altered

mRNA stability(Xia, Bogardus and Prochazka, 1999). Vilmundarson et al. demonstrated that

HuR differentially modulates *IRF2BP2* translation through a 3′ UTR polymorphism that is

associated with increased coronary artery calcification(Vilmundarson et al., 2021). Our

exploratory studies did not resolve whether the rs13900 allelic variation leads to differences

in polysomal loading and additional studies are required to address this issue. Another

important mechanism by which disease associated genetic variants in the UTRs could

influence mRNA stability and translatability is by disrupting or creating microRNA binding

sites(Huang et al., 2019). We examined the region flanking the rs13900 and found that there

are several predicted binding sites for miRNAs. Among these, several miRNAs have minimum

free energy ≤ –25 kcal/mol suggesting that they can potentially play a role in *CCL2* expression

and will be investigated in future studies. Of note, a recent study showed that RBPs may play

an important role in miRNA mediated gene regulation(Kim et al., 2021).

RBPs regulate the protein expression from a given mRNA by modulation of its half-life,

subcellular localization, and ribosomal recruitment(Glisovic et al., 2008). However, mRNA

abundance and stability are not always predictive of protein synthesis: relative mRNA

abundance/stability and translation levels of a given gene can vary significantly and

determined by an assortment of post-transcriptional events(Liu, Beyer and Aebersold, 2016).

Our previous findings and results from the present study suggest that there is a close

association between *CCL2* rs1024611-rs13900 haplotype, mRNA, and protein expression.

Nevertheless, we cannot completely rule out the contribution of transcription-based

mechanisms to the allelic expression imbalance at the *CCL2* locus. Previous studies have

suggested that there is a high level of conservation for interactions between RBPs and their

target molecules and that RNA-mediated gene regulation is less evolvable than transcriptional

regulation(Payne, Khalid and Wagner, 2018). Thus, it is remarkable that the rs13900 alters

HuR binding and impacts both transcript stability and translatability. Our bioinformatic analysis

shows that rs13900 could potentially lead to dramatic changes in the secondary structure of

CCL2 mRNA (Fig. 3B). A recent study using bioinformatic analyses has shown that SNPs may

affect RNA-protein interactions from outside binding motifs through altered RNA secondary

structure(Shatoff and Bundschuh, 2020). We cannot rule out the possibility that HuR may

differentially bind to other regions of *CCL2* transcript due to changes in secondary structure

and therefore will be examined in future studies.

Both rs1024611 and rs13900 are in high LD with rs3091315, which is detected in the GWA

studies of CD and IBD(de Lange et al., 2017; Franke et al., 2010; Kenny et al., 2012; Liu et

al., 2015; Palmieri et al., 2010). rs3091315 is classified as an upstream risk variant and is

located in the intergenic region between *CCL2* and *CCL7*. The GWA risk allele for CD is

rs3091315-A which is correlated with rs1024611(A) and rs13900(C) alleles. While the

biological role of *CCL2* in CD and IBD are well recognized(Darkoh et al., 2014; Maharshak et

al., 2010; Martin et al., 2019), the results from genetic association studies are not consistent

and the disease outcomes could potentially differ by population studied(Chen et al., 2016).

In conclusion, our study shows that the disease associations mediated by the *CCL2*

rs1024611-rs13900 haplotype may be due to altered mRNA stability mediated through

differential binding of an RBP. Given the importance of mRNA stability in immune homeostasis

such mechanisms could play a critical role in inter-individual differences in disease

pathogenesis.

Materials and Methods

Recruitment of study participants, primary cell culture, and genotyping

All research involving human subjects were approved by the Institutional Review Boards

(IRBs) of the University of Texas Health San Antonio, San Antonio, Texas and University of

Texas Rio Grande Valley, Edinburg, Texas. Written, informed consent from each individual for

participation in our study was obtained following the approval by the IRBs. A total of 47

unrelated individuals were recruited into the study. Data and samples from the study

participants were obtained at the First Outpatient Research Unit (FORU), UT Health San

Antonio, San Antonio, Texas. The rs13900 polymorphism was detected by TaqMan

predesigned SNP genotyping assay (Applied Biosystems, CA, USA, Cat. No. 4351379).

Briefly, genomic DNA samples were obtained from peripheral blood of healthy study

participants using QIAamp DNA Blood Mini Kits (QIAGEN, CA) in accordance with the

manufacturer's protocol, checked for quality and concentration, and stored at −80ºC until

used. Genotyping of rs13900 was performed on 10 ng of genomic DNA using TaqMan

genotyping Master mix in a 10 µL reaction volume. PCR was performed on the Quantstudio

12K Flex (Applied Biosystems) in 384-well plates. The temperature cycling conditions

consisted of an initial enzyme heat-activation step of 10 min at 95ºC and 40 cycles of a 3-step

amplification profile of 20 sec at 95ºC for denaturation, 1 min at 60ºC for annealing, and 30

sec for 72ºC for extension. QS12K software (Applied Biosystems, CA, USA). was used to

score the alleles. Whole blood samples were collected from study participants who were either

homozygous (either C/C or T/T) or heterozygous (C/T) for rs13900 at the initial screening for

the genotype status, followed by recalling selected individuals for detailed studies. Peripheral

blood mononuclear cells (PBMC) were isolated by a Ficoll density gradient centrifugation.

Total genomic DNA and RNA were isolated from 0.5×10^6 PBMC following stimulation with LPS

for 3 h as described previously(Pham et al., 2012; Sharif et al., 2007). CD14+ monocytes were

purified using EasySep Human Monocytes Isolation Kit (negative selection kit; Stem cell

Technologies) according to manufacturer's instructions. Monocytes were treated with M-CSF

for 72 h to induce macrophage differentiation, and flow cytometric measurement of surface

markers CD64, CD206, and CD44 was used to confirm the differentiation.

Real-time quantitative PCR (RT-qPCR) assays

Total RNA was isolated using the RNeasy plus mini kit (Qiagen) according to manufacturer's

instructions. 10 μL of total RNA sample (500 ng) was converted to cDNA using MultiScribe

Reverse Transcriptase (Applied Biosystems) with random primers. Real-time quantitative

PCR was performed using TaqMan gene expression assays. Relative expression levels were

calculated by applying the $2^{-\Delta\Delta Ct}$ method using 18S rRNA as a reference.

Capture of nascent RNA

Newly synthesized RNA was captured by isolating Nascent RNA using Click-It-Nascent RNA

kit (Invitrogen) according to the manufacturer's recommended protocol. Briefly, PBMC or

monocyte derived macrophages (MDMs) from heterozygous individuals were stimulated with

lipopolysaccharide (LPS) for 3 h and incubated with actinomycin D (ActD) and ethylene uridine

(EU) which is an alkyne–modified nucleoside and a uridine analog. The EU RNA was

subjected to a click reaction that adds a biotin handle which was then captured by streptavidin

beads. The captured RNA was used for cDNA synthesis (Superscript Vilo kit, Invitrogen), PCR

amplification, and allelic quantification.

AEI assessment of *CCL2*

Total RNA and genomic DNA were isolated from primary immune cells (PBMC and

macrophages) after they were treated with LPS for 3 h. Total RNA was reverse transcribed as

described above. The region encompassing rs13900 marker was amplified by PCR in a 50 μL

reaction mixture containing 2 μL of cDNA or genomic DNA, 25 μL AmpliTaq Gold 360 Master

Mix and 10 μM each of the forward and reverse primers. The nucleotide sequence of the

forward primer and reverse primers were 5′-ACCTGGACAAGCAAACCCAA-3′ and 5′-

ACCCTCAAAACATCCCAGGG-3′. The following temperature conditions were used: initial

denaturation at 95°C for 10 min followed by 40 cycles of denaturation at 95°C for 30 s,

annealing at 53.6°C for 30 s, extension at 72°C for 30 s, and final extension at 72°C for 7 min.

The amplicons were purified with The GeneJET PCR Purification Kit (ThermoFisher Scientific)

and were subjected to Sanger sequencing (Macrogen, USA) using the sequencing primer 5′-

GCAAACCCAAACTCCGAAGAC-3′. The degree of AEI for CCL2 was expressed as a ratio of

reference allele (REF; major) to alternative allele (ALT; minor). PeakPicker,v.2.0 was used

with default settings to quantify the relative amount of the two alleles from the chromatogram

after peak intensity normalization(Ge et al., 2005).

Bioinformatic analyses

The ex vivo binding of RBPs to region flanking rs13900 and CCL2 3′ UTR was examined using

Atlas of UTR Regulatory Activity (AURA)(Dassi et al., 2014). The resources included in

POSTAR3 were used to interrogate the RNA binding motifs that overlap the rs13900(Zhao et

al., 2022). RBP-var2 was used to explore the potential functional consequences(Mao et al.,

2016), and the ViennaRNA package(Lorenz et al., 2011) was used to predict changes in *CCL2*

secondary structure due to rs13900.

RNA electrophoretic mobility shift assay (REMSA)

Oligoribonucleotides spanning the rs13900 C or rs13900 T alleles (IDT) labeled with infrared

dyes were used to determine the differential binding of HuR. The sequences of the

oligoribonucleotide   with   rs13900   C   allele   is

rCrUrUrUrCrCrCrCrArGrArCrArCrCrCrUrGrUrUrUrUrArUrUrU and oligoribonucleotide with

rs13900 T allele is rCrUrUrUrCrCrCrCrArGrArCrArCrCrUrUrGrUrUrUrUrArUrUrU. The

oligoribonucleotides were incubated with either 10 μg HuR overexpressing HeLa cell whole

cell extracts or 25-200 μM ELAVL1 human recombinant protein (Origene, Rockville, MD) in a

20 μL reaction mixture with 20 units of RNasin and 1X RNA binding buffer containing 20 mM

HEPES (pH 7.6), 3 mM $MgCl_2$, 40 mM KCl, 5% glycerol and 2 mM dithiothreitol, 20 μg yeast

tRNA. The reaction was incubated on ice for 10 min and super-shift assay was performed by

adding 2 μL of anti-HuR antibody (3A2, mouse monoclonal IgG antibody, Santa Cruz

Biotechnology). After antibody addition, the complexes were incubated on ice for 15 min and

were resolved by electrophoresis under non-denaturing conditions in 5% polyacrylamide gels

with 0.5 × TBE running buffer. Gels were then analyzed using Li-Cor Odyssey CLX.

Cell culture and transfection

HEK-293 cell line was maintained in EMEM (ATCC) containing 10% heat-inactivated FBS,

penicillin (100 U/mL) and streptomycin (100mg, mL) (GIBCO). Cells were cultured at 37⁰C in

humidified air containing 5% CO_2. For HuR overexpression and silencing studies, cells were

seeded in six-well plates in serum-free EMEM at a density of ~ 250,000 cells per well. The

cells were transfected when they reached ~ 70 to 80% confluency, using lipofectamine 3000

(Thermo Fisher Scientific) according to the manufacturer's recommended protocol. Briefly,

lipofectamine 3000 was diluted in OptiMEM and allowed to complex with either 0.5 μg of

pCMV6-HuR or pCMV-Entry plasmid or 50 μM of HuR siRNA or control siRNA for 15 min at

room temperature. The plasmid/siRNA-lipofectamine 3000 mixture was then added to the cells

in a final volume of 2 mL of complete medium. After incubation for 72 h, cells were harvested

and used for further processing.

Western blotting

Cell lysates were prepared in chilled RIPA buffer (25 mM Tris-HCL pH 7.6, 150 mM NaCl, 1%

NP-40, 1% sodium deoxycholate, 0.1% SDS; Thermo Scientific, Rockford, IL) containing

protease inhibitor (complete Mini, EDTA-free Protease inhibitor cocktail tablets, Roche

Diagnostics, Indianapolis, IN). Cells were lysed on ice for 10-15 min followed by centrifugation

at 12,000 × g for 15 min, and the cleared supernatant was collected and stored at −20ºC. Cell

lysates with equal amounts of total protein (15-20 µg) were loaded and separated on a SDS

polyacrylamide gel (10–15%) and were electrophoretically transferred onto nitrocellulose

membranes (Thermo Scientific). The membranes were blocked with non-fat dry milk in TBST

(50 mM Tris-HCl, pH 7.4, 150 mM NaCl, 0.2% Tween 20) and were incubated overnight with

anti-HuR primary antibodies (diluted 1:1000). Following incubation, the membranes were

washed and then further incubated with anti-mouse ß-actin for one h at room temperature,

followed by HRP labeled goat anti-mouse or goat anti-rabbit antibodies (1:1000; Santa Cruz).

Protein bands were detected using a chemiluminescence (ECL kit) method (Pierce) and

visualized on X-ray film (Kodak).

Reporter assays

Site-specific mutagenesis was done on CCL2 LightSwitch 3′ UTR reporter (Active Motif) to

generate constructs containing rs13900 C or rs13900 T alleles. The nucleotide sequence of

the complete constructs was verified by Sanger sequencing (Macrogen, USA). 0.5 µg of these

constructs were nucleofected into HEK293 cells using 4D Nucleofector (Lonza) and luciferase

activity was measured 24 h post transfection using Spectramax plate reader (Molecular

Devices). The efficiency of the nucleofection was verified by confocal microscopy, and it was

greater than 90%. Given the high efficiency of the nucleofection and to avoid cross-talk with

co-transfected control plasmids(Farr and Roman, 1992), luciferase data across samples was

normalized using the protein content of the lysates using BCA protein assay. mRNA

translatability was assessed using methods as described previously(Zhang et al., 2017).

Polysome profiling

Monocytes differentiated primary macrophages were obtained from heterozygous individuals

and were treated with LPS or PBS for 3 h to induce CCL2 expression. Cells were treated with

0.1 mM cycloheximide, a translation inhibitor, for 3 min, harvested and washed with ice-cold

PBS containing 0.1 mM cycloheximide, and pelleted by centrifugation for 5 min at 500 x g.

The pellets were either stored at −80ºC or immediately used for cytoplasmic lysate

preparation. The cell pellet was resuspended in 500 µL lysis buffer containing 150 mM NaCl,

50 mM Tris-HCl pH 7.5, 10 mM KCl, 10 mM $MgCl_2$, 0.2% NP-40, 2 mM dithiothreitol, 2 mM

sodium orthovanadate, 1 mM phenylmethylsulfonyl fluoride, and 80 units/mL RNaseOUT.

After 10 min incubation on ice with occasional inverting every two minutes, samples were

centrifuged at 12,000 x g for 10 min at 4°C to pellet the nuclei and the post-nuclear

supernatants. The optical density at 254 nm was measured, and volumes corresponding to

the same OD254 were used. Post-nuclear supernatants were laid on top of a 10-50% sucrose

gradient. The gradient was centrifuged for 90 min at 200,000 x g. After the ultracentrifugation,

ten fractions were collected from top to bottom. RNA was extracted from each fraction and

RT-qPCR was performed using *CCL2* mRNA specific primers. For analysis, fraction one was

considered as cytoplasmic lysate, fractions 2-4 were considered monosomal fraction, and

fractions 6-10 as polysomal fraction. The percent (%) distribution for the *CCL2* mRNAs across

the gradients was calculated using the differences in the cycle threshold (ΔCt) values using

the following formula(Nayak et al., 2013):

% of mRNA A in each fraction = $2^{\Delta CT\ fraction\ X}$ x 100/Sum, where ΔCT fraction X = CT (fraction

1) − CT (fraction X)

Lentiviral transduction of primary macrophages

626   Ready to use GFP- tagged pCMV6-HuR or CMV-null lentiviral particles (Amsbio, Cambridge,

627   MA) were transduced into 0.5×10^6 macrophages in the presence of polybrene (60µg/ml) at an

628   MOI of 1. The cells were processed for HuR and CCL2 expression 72 h after transduction.

Statistical Analyses

631   All values in figures are presented as the mean ± SEM. Results were analyzed with ANOVA

632   and posthoc contrasts with Fisher (Least Significant Difference (LSD) test or Student's t test.

633   Statistical analyses were carried out in the Sigma Plot 12.0 software.

634 **Acknowledgments**

635 We thank the participants of the study for their time and support.

  Figure Captions

  Figure.1. rs13900 heterozygous individuals exhibit AEI in *CCL2*. (A) Schematic depicting

  distal, proximal regulatory elements extending 3 kb on either side of CCL2 gene and the LD

  between regulatory polymorphism rs1024611 and the transcribed polymorphism rs13900.

  rs1024611 is located 2578 base pairs upstream of the *CCL2* translation start site and rs13900

  located in the *CCL2* 3′ UTR. **(B)** Allelic expression imbalance (AEI) in heterozygous donors is

  measured as a ratio of alternative allele (ALT) to reference allele (REF) in a transcribed

  polymorphism. **(C)** Representative chromatograms obtained following Sanger sequencing of

  PCR products obtained from genomic DNA (gDNA) and reverse transcription-PCR of mRNA

  (cDNA) from three individuals heterozygous for rs13900. gDNA and mRNA were obtained

  from PBMC treated with LPS for 3 h as previously described.   The allelic ratios shown were

  determined by PeakPicker analysis. Peakpicker calculates allelic ratios by dividing the peak

  height of the alternate allele (rs13900 T allele) by that of reference allele (rs13900 C allele).

  The gDNA peaks were used for normalization. **(D)** Allelic ratio for cDNA and gDNA in six

  individuals heterozygous for rs13900 after treatment with LPS for 3 h. Statistical significance

  for the difference in the level of expression between the alleles was determined using

  Student's *t*-test ($P < 0.003$).

  Figure 2. rs13900T confers greater stability to *CCL2* mRNA. (A) *CCL2* mRNA expression

  in peripheral monocytes of heterozygous individuals (n = 4) after treatment with LPS for 3 h

  and then incubated with 5 µg Act D for indicated times. mRNA was detected by RT-PCR.

  Results, normalized to 18S rRNA levels, are expressed as fold increase over unstimulated

  cells (CNT). Levels shown in bar graph represent mean ± SEM of result at time 0. (*$P = 0.019$)

  versus unstimulated cells (N = 4). **(B)** *CCL2* mRNA half-life, calculated for each condition as

  the time (in hours) required for the transcript to decrease to 50% of its initial abundance [t 1/2

  = Ln (0.5)/slope]. **(C)** Nascent RNA was isolated from treated monocytes from three individuals

  in the presence and absence of ActD. Allelic ratio was determined after 4 h of incubation with

or without ActD. Expression of rs13900 T allele was much higher in ActD-treated samples.

The difference between the groups were assessed by ANOVA with Fisher LSD method (*$P <$

0.05, ** $P < 0.005$).

Figure 3. Bioinformatic analysis of the rs13900. (A) Validation RBP binding sites and

polymorphism located on the 3′ UTR of *CCL2* transcript from the Atlas of UTR Regulatory

Activity and analysis of ENCODE genome-wide data sets detected specific enrichment of HuR

(ELAVL1) at the region that contains the rs13900. **(B)** Predicted changes in the secondary

structure using Vienna RNA package 2.0; changes in secondary structure are indicated by

arrows. **(C)** Sequence logo of HuR binding site as determined by HOMER. **(D)** Relative

structural preference of HuR to different structural contexts, the letters P, L, U, M indicate that

the nucleotide is paired (P), in a hairpin loop (L), in an unstructured (or external) region (U),

and Miscellaneous (M) region.

Figure 4. The rs13900 T allele shows increased in vitro binding of HuR. (A) REMSA with

labelled oligoribonucleotide containing either rs13900 C or T allele and whole cell extracts

from 293T cells. § denotes free probe; black arrow, bound probe; red arrow, supershift. **(B)**

Representative quantitative densitometric analysis of the antibody shifted complexes

suggested increased HuR binding to the oligoribonucleotide bearing rs13900 T allele. The

signals in the bound fraction(s) were normalized using the free probe (N = 4). The top panel

represents the data from four independent experiments (mean ± SEM). Statistical analyses

were performed using Student's t test (*$P < 0.001$). The bottom panel shows the relative fold

enrichment of the bound protein complexes to the oligoribonucleotide containing the rs13900

T allele relative to that containing the rs13900 C allele . Statistical significance was calculated

using Student t test (*$P < 0.001$) **(C)** REMSA with labelled oligoribonucleotides containing

either rs13900 T or C allele and purified HuR protein. **(D)** Plot showing the fraction of bound

rs13900 C or rs13900 T oligoribonucleotides with increasing HuR concentration.

Figure 5. rs13900 C and T alleles are associated with differential binding to HuR ex vivo.

(A) HuR enrichment in immunoprecipitated material from macrophages stimulated with LPS.

(B) *CCL2* 3′ UTR was detected at significant levels in the samples precipitated by α-HuR

antibody when compared to the control IgG. **(C)** *CCL2* mRNA expression in anti-HuR antibody

enriched immunoprecipitated material analyzed by RT-qPCR (N = 4). Statistical significance

was calculated using Student t-test (*$P < 0.005$). The error bars represent SEM. **(D)** Relative

expression levels for rs13900 C and T alleles in macrophages stimulated with LPS (N = 4).

Statistical significance was calculated using a Student t-test (*$P < 0.005$).

Figure 6. Differential effects of rs13900 alleles in reporter assays and role of HuR. (A)

Schematic representation of the luciferase reporter vectors containing *CCL2* 3' UTR with either

rs13900 C or T allele. **(B)** HEK-293 cells were transfected with the equal quantities of *CCL2*

3′ UTR reporter vectors and luciferase activity was measured 48 hrs later. The relative

luciferase activities of the 3′ UTR reporter plasmids were expressed as percent reduction in

the luminescence when compared to the control vector that was set to 100% after normalizing

for the protein content of the lysates. The error bars indicate the standard error of mean and

statistical significance was calculated using two-tailed Student's *t* test (*$P < 0.05$). **(C)** HEK-

293T cells were transfected with either pCMV6-HuR (0.5 µg) or pCMV-Entry (0.5 µg) and after

72 hours they were co-transfected with the two plasmid constructs (0.5 µg). Twenty-four hours

after transfection the relative change in luciferase activity was determined. **(D)** Cells were co-

transfected with 125 pmol HuR siRNA or control siRNA and with the two plasmid constructs

(0.5 µg). Twenty-four hours after transfection the relative change in luciferase activity was

determined (normalized to total protein concentration, data from four independent

experiments (mean ± SEM; N = 4). Statistical analyses were performed using Fisher LSD

method (*$P < 0.05$).

Figure 7. Influence of rs13900 C and T alleles on *CCL2* translatability. (A) HEK-293 cells

were transfected by nucleofection with control and *CCL2* 3′ UTR reporter constructs with

rs13900 C and T alleles. The nucleofected cells were plated separately and harvested for total

RNA isolation or lysed for mRNA level or protein level expression of luciferase respectively

after 24 h. The reporter mRNA levels from the transfected 293 T cells were quantified by qRT-

PCR, and 18S rRNA was used for normalization. **(B)** The relative luciferase activities of the 3′

UTR reporter plasmids were expressed as percent reduction in the luminescence when

compared to the control vector that was set to 100% after normalizing for the protein content

of the lysates. **(C)** mRNA translatability was calculated as luciferase activity normalized by the

reporter luciferase mRNA level. The error bars indicate the standard error of mean from six

independent experiments (N = 6) and statistical significance was calculated using ANOVA and

post hoc contrast with Fisher LSD method. (*$P < 0.05$, **$P < 0.005$).

Figure 8. HuR differentially regulates *CCL2* haplotypes. (A) HuR expression in primary

human macrophages following lentiviral transduction. Macrophages obtained from four

individuals who are homozygous for either rs13900 C or rs13900 T allele were transduced

with either CMV-null or HuR expressing lentiviral particles (pCMV6-HuR) for 72 hours followed

by LPS stimulation for 3 h. **(B)** *CCL2* expression determined by RT-qPCR. Error bars

represent SEM. Statistical analyses were performed using Student's t test   (* $P < 0.05$).

Description of the supplemental data

The Supplemental data includes Tables S1-S2 and Figures S1-S5.

Funding

This study was supported by Grant from National Institute of Health (5R01AI119131)

Declaration of interests

The authors declare no competing interests.

Web resources

AURA, http://aura.science.unitn.it/

POSTAR3, http://postar.ncrnalab.org

RBP-var2, http://159.226.67.237/sun/RBP-Var/index.php/Rbp/index

ViennaRNA package, https://www.tbi.univie.ac.at/RNA/

References

Akdeli, N., Riemann, K., Westphal, J., Hess, J., Siffert, W., Bachmann, H.S., 2014. A 3'UTR
polymorphism modulates mRNA stability of the oncogene and drug target Polo-like Kinase 1. Mol
Cancer 13, 87.
Alasoo, K., Rodrigues, J., Mukhopadhyay, S., Knights, A.J., Mann, A.L., Kundu, K., Consortium, H.,
Hale, C., Dougan, G., Gaffney, D.J., 2018. Shared genetic effects on chromatin and gene expression
indicate a role for enhancer priming in immune response. Nat Genet 50, 424-431.
Alonso-Villaverde, C., Coll, B., Parra, S., Montero, M., Calvo, N., Tous, M., Joven, J., Masana, L., 2004.
Atherosclerosis in patients infected with HIV is influenced by a mutant monocyte chemoattractant
protein-1 allele. Circulation 110, 2204-2209.
Anderson, P., 2010. Post-transcriptional regulons coordinate the initiation and resolution of
inflammation. Nat Rev Immunol 10, 24-35.
Bonello, G.B., Pham, M.H., Begum, K., Sigala, J., Sataranatarajan, K., Mummidi, S., 2011. An
evolutionarily conserved TNF-alpha-responsive enhancer in the far upstream region of human CCL2
locus influences its gene expression. J Immunol 186, 7025-7038.
Buccitelli, C., Selbach, M., 2020. mRNAs, proteins and the emerging principles of gene expression
control. Nat Rev Genet 21, 630-644.
Buniello, A., MacArthur, J.A.L., Cerezo, M., Harris, L.W., Hayhurst, J., Malangone, C., McMahon, A.,
Morales, J., Mountjoy, E., Sollis, E., Suveges, D., Vrousgou, O., Whetzel, P.L., Amode, R., Guillen, J.A.,
Riat, H.S., Trevanion, S.J., Hall, P., Junkins, H., Flicek, P., Burdett, T., Hindorff, L.A., Cunningham, F.,
Parkinson, H., 2019. The NHGRI-EBI GWAS Catalog of published genome-wide association studies,
targeted arrays and summary statistics 2019. Nucleic Acids Res 47, D1005-D1012.
Cavestro, G.M., Zuppardo, R.A., Bertolini, S., Sereni, G., Frulloni, L., Okolicsanyi, S., Calzolari, C.,
Singh, S.K., Sianesi, M., Del Rio, P., Leandro, G., Franze, A., Di Mario, F., 2010. Connections between
genetics and clinical data: Role of MCP-1, CFTR, and SPINK-1 in the setting of acute, acute recurrent,
and chronic pancreatitis. Am J Gastroenterol 105, 199-206.
Chen, S., Deng, C., Hu, C., Li, J., Wen, X., Wu, Z., Li, Y., Zhang, F., Li, Y., 2016. Association of MCP-1-
2518A/G polymorphism with susceptibility to autoimmune diseases: a meta-analysis. Clin Rheumatol
35, 1169-1179.
Cho, M.L., Kim, J.Y., Ko, H.J., Kim, Y.H., Kim, W.U., Cho, C.S., Kim, H.Y., Hwang, S.Y., 2004. The MCP-1
promoter -2518 polymorphism in Behcet's disease: correlation between allele types, MCP-1
production and clinical symptoms among Korean patients. Autoimmunity 37, 77-80.
Choi, H.J., Yang, H., Park, S.H., Moon, Y., 2009. HuR/ELAVL1 RNA binding protein modulates
interleukin-8 induction by muco-active ribotoxin deoxynivalenol. Toxicol Appl Pharmacol 240, 46-54.
Chowdhury, P., Khan, S.A., 2017. Significance of CCL2, CCL5 and CCR2 polymorphisms for adverse
prognosis of Japanese encephalitis from an endemic population of India. Sci Rep 7, 13716.
Darkoh, C., Comer, L., Zewdie, G., Harold, S., Snyder, N., Dupont, H.L., 2014. Chemotactic
chemokines are important in the pathogenesis of irritable bowel syndrome. PLoS One 9, e93144.
Dassi, E., Re, A., Leo, S., Tebaldi, T., Pasini, L., Peroni, D., Quattrone, A., 2014. AURA 2: Empowering
discovery of post-transcriptional networks. Translation (Austin) 2, e27738.
de Lange, K.M., Moutsianas, L., Lee, J.C., Lamb, C.A., Luo, Y., Kennedy, N.A., Jostins, L., Rice, D.L.,
Gutierrez-Achury, J., Ji, S.G., Heap, G., Nimmo, E.R., Edwards, C., Henderson, P., Mowat, C.,
Sanderson, J., Satsangi, J., Simmons, A., Wilson, D.C., Tremelling, M., Hart, A., Mathew, C.G.,
Newman, W.G., Parkes, M., Lees, C.W., Uhlig, H., Hawkey, C., Prescott, N.J., Ahmad, T., Mansfield,
J.C., Anderson, C.A., Barrett, J.C., 2017. Genome-wide association study implicates immune
activation of multiple integrin genes in inflammatory bowel disease. Nat Genet 49, 256-261.
Deshmane, S.L., Kremlev, S., Amini, S., Sawaya, B.E., 2009. Monocyte chemoattractant protein-1
(MCP-1): an overview. J Interferon Cytokine Res 29, 313-326.
Duan, J., Shi, J., Ge, X., Dolken, L., Moy, W., He, D., Shi, S., Sanders, A.R., Ross, J., Gejman, P.V., 2013.
Genome-wide survey of interindividual differences of RNA stability in human lymphoblastoid cell
lines. Sci Rep 3, 1318.

Fan, J., Ishmael, F.T., Fang, X., Myers, A., Cheadle, C., Huang, S.K., Atasoy, U., Gorospe, M., Stellato,
C., 2011. Chemokine transcripts as targets of the RNA-binding protein HuR in human airway
epithelium. J Immunol 186, 2482-2494.
Farh, K.K., Marson, A., Zhu, J., Kleinewietfeld, M., Housley, W.J., Beik, S., Shoresh, N., Whitton, H.,
Ryan, R.J., Shishkin, A.A., Hatan, M., Carrasco-Alfonso, M.J., Mayer, D., Luckey, C.J., Patsopoulos,
N.A., De Jager, P.L., Kuchroo, V.K., Epstein, C.B., Daly, M.J., Hafler, D.A., Bernstein, B.E., 2015.
Genetic and epigenetic fine mapping of causal autoimmune disease variants. Nature 518, 337-343.
Farr, A., Roman, A., 1992. A pitfall of using a second plasmid to determine transfection efficiency.
Nucleic Acids Res 20, 920.
Fenoglio, C., Galimberti, D., Lovati, C., Guidi, I., Gatti, A., Fogliarino, S., Tiriticco, M., Mariani, C.,
Forloni, G., Pettenati, C., Baron, P., Conti, G., Bresolin, N., Scarpini, E., 2004. MCP-1 in Alzheimer's
disease patients: A-2518G polymorphism and serum levels. Neurobiol Aging 25, 1169-1173.
Flores-Villanueva, P.O., Ruiz-Morales, J.A., Song, C.H., Flores, L.M., Jo, E.K., Montano, M., Barnes,
P.F., Selman, M., Granados, J., 2005. A functional promoter polymorphism in monocyte
chemoattractant protein-1 is associated with increased susceptibility to pulmonary tuberculosis. J
Exp Med 202, 1649-1658.
Franke, A., McGovern, D.P., Barrett, J.C., Wang, K., Radford-Smith, G.L., Ahmad, T., Lees, C.W.,
Balschun, T., Lee, J., Roberts, R., Anderson, C.A., Bis, J.C., Bumpstead, S., Ellinghaus, D., Festen, E.M.,
Georges, M., Green, T., Haritunians, T., Jostins, L., Latiano, A., Mathew, C.G., Montgomery, G.W.,
Prescott, N.J., Raychaudhuri, S., Rotter, J.I., Schumm, P., Sharma, Y., Simms, L.A., Taylor, K.D.,
Whiteman, D., Wijmenga, C., Baldassano, R.N., Barclay, M., Bayless, T.M., Brand, S., Buning, C.,
Cohen, A., Colombel, J.F., Cottone, M., Stronati, L., Denson, T., De Vos, M., D'Inca, R., Dubinsky, M.,
Edwards, C., Florin, T., Franchimont, D., Gearry, R., Glas, J., Van Gossum, A., Guthery, S.L.,
Halfvarson, J., Verspaget, H.W., Hugot, J.P., Karban, A., Laukens, D., Lawrance, I., Lemann, M.,
Levine, A., Libioulle, C., Louis, E., Mowat, C., Newman, W., Panes, J., Phillips, A., Proctor, D.D.,
Regueiro, M., Russell, R., Rutgeerts, P., Sanderson, J., Sans, M., Seibold, F., Steinhart, A.H., Stokkers,
P.C., Torkvist, L., Kullak-Ublick, G., Wilson, D., Walters, T., Targan, S.R., Brant, S.R., Rioux, J.D.,
D'Amato, M., Weersma, R.K., Kugathasan, S., Griffiths, A.M., Mansfield, J.C., Vermeire, S., Duerr,
R.H., Silverberg, M.S., Satsangi, J., Schreiber, S., Cho, J.H., Annese, V., Hakonarson, H., Daly, M.J.,
Parkes, M., 2010. Genome-wide meta-analysis increases to 71 the number of confirmed Crohn's
disease susceptibility loci. Nat Genet 42, 1118-1125.
Ge, B., Gurd, S., Gaudin, T., Dore, C., Lepage, P., Harmsen, E., Hudson, T.J., Pastinen, T., 2005. Survey
of allelic expression using EST mining. Genome Res 15, 1584-1591.
Glisovic, T., Bachorik, J.L., Yong, J., Dreyfuss, G., 2008. RNA-binding proteins and post-transcriptional
gene regulation. FEBS Lett 582, 1977-1986.
Gonzalez, E., Rovin, B.H., Sen, L., Cooke, G., Dhanda, R., Mummidi, S., Kulkarni, H., Bamshad, M.J.,
Telles, V., Anderson, S.A., Walter, E.A., Stephan, K.T., Deucher, M., Mangano, A., Bologna, R., Ahuja,
S.S., Dolan, M.J., Ahuja, S.K., 2002. HIV-1 infection and AIDS dementia are influenced by a mutant
MCP-1 allele linked to increased monocyte infiltration of tissues and MCP-1 levels. Proc Natl Acad Sci
U S A 99, 13795-13800.
Gschwandtner, M., Derler, R., Midwood, K.S., 2019. More Than Just Attractive: How CCL2 Influences
Myeloid Cell Behavior Beyond Chemotaxis. Front Immunol 10, 2759.
Hao, S., Baltimore, D., 2009. The stability of mRNA influences the temporal order of the induction of
genes encoding inflammatory molecules. Nat Immunol 10, 281-288.
Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre, C., Singh, H.,
Glass, C.K., 2010. Simple combinations of lineage-determining transcription factors prime cis-
regulatory elements required for macrophage and B cell identities. Mol Cell 38, 576-589.
Hollander, D., Naftelberg, S., Lev-Maor, G., Kornblihtt, A.R., Ast, G., 2016. How Are Short Exons
Flanked by Long Introns Defined and Committed to Splicing? Trends Genet 32, 596-606.

Huang, Z., Shi, J., Gao, Y., Cui, C., Zhang, S., Li, J., Zhou, Y., Cui, Q., 2019. HMDD v3.0: a database for experimentally supported human microRNA-disease associations. Nucleic Acids Res 47, D1013-D1017.

Joven, J., Coll, B., Tous, M., Ferre, N., Alonso-Villaverde, C., Parra, S., Camps, J., 2006. The influence of HIV infection on the correlation between plasma concentrations of monocyte chemoattractant protein-1 and carotid atherosclerosis. Clin Chim Acta 368, 114-119.

Kazan, H., Ray, D., Chan, E.T., Hughes, T.R., Morris, Q., 2010. RNAcontext: a new method for learning the sequence and structure binding preferences of RNA-binding proteins. PLoS Comput Biol 6, e1000832.

Kenny, E.E., Pe'er, I., Karban, A., Ozelius, L., Mitchell, A.A., Ng, S.M., Erazo, M., Ostrer, H., Abraham, C., Abreu, M.T., Atzmon, G., Barzilai, N., Brant, S.R., Bressman, S., Burns, E.R., Chowers, Y., Clark, L.N., Darvasi, A., Doheny, D., Duerr, R.H., Eliakim, R., Giladi, N., Gregersen, P.K., Hakonarson, H., Jones, M.R., Marder, K., McGovern, D.P., Mulle, J., Orr-Urtreger, A., Proctor, D.D., Pulver, A., Rotter, J.I., Silverberg, M.S., Ullman, T., Warren, S.T., Waterman, M., Zhang, W., Bergman, A., Mayer, L., Katz, S., Desnick, R.J., Cho, J.H., Peter, I., 2012. A genome-wide scan of Ashkenazi Jewish Crohn's disease suggests novel susceptibility loci. PLoS Genet 8, e1002559.

Kim, S., Kim, S., Chang, H.R., Kim, D., Park, J., Son, N., Park, J., Yoon, M., Chae, G., Kim, Y.K., Kim, V.N., Kim, Y.K., Nam, J.W., Shin, C., Baek, D., 2021. The regulatory impact of RNA-binding proteins on microRNA targeting. Nat Commun 12, 5057.

Kwong, A., Boughton, A.P., Wang, M., VandeHaar, P., Boehnke, M., Abecasis, G., Kang, H.M., 2021. FIVEx: an interactive eQTL browser across public datasets. Bioinformatics.

Lebedeva, S., Jens, M., Theil, K., Schwanhausser, B., Selbach, M., Landthaler, M., Rajewsky, N., 2011. Transcriptome-wide analysis of regulatory interactions of the RNA-binding protein HuR. Mol Cell 43, 340-352.

Letendre, S., Marquie-Beck, J., Singh, K.K., de Almeida, S., Zimmerman, J., Spector, S.A., Grant, I., Ellis, R., Group, H., 2004. The monocyte chemotactic protein-1 -2578G allele is associated with elevated MCP-1 concentrations in cerebrospinal fluid. J Neuroimmunol 157, 193-196.

Li, Y.I., van de Geijn, B., Raj, A., Knowles, D.A., Petti, A.A., Golan, D., Gilad, Y., Pritchard, J.K., 2016. RNA splicing is a primary link between genetic variation and disease. Science 352, 600-604.

Liu, J.Z., van Sommeren, S., Huang, H., Ng, S.C., Alberts, R., Takahashi, A., Ripke, S., Lee, J.C., Jostins, L., Shah, T., Abedian, S., Cheon, J.H., Cho, J., Dayani, N.E., Franke, L., Fuyuno, Y., Hart, A., Juyal, R.C., Juyal, G., Kim, W.H., Morris, A.P., Poustchi, H., Newman, W.G., Midha, V., Orchard, T.R., Vahedi, H., Sood, A., Sung, J.Y., Malekzadeh, R., Westra, H.J., Yamazaki, K., Yang, S.K., International Multiple Sclerosis Genetics, C., International, I.B.D.G.C., Barrett, J.C., Alizadeh, B.Z., Parkes, M., Bk, T., Daly, M.J., Kubo, M., Anderson, C.A., Weersma, R.K., 2015. Association analyses identify 38 susceptibility loci for inflammatory bowel disease and highlight shared genetic risk across populations. Nat Genet 47, 979-986.

Liu, Y., Beyer, A., Aebersold, R., 2016. On the Dependency of Cellular Protein Levels on mRNA Abundance. Cell 165, 535-550.

Lorenz, R., Bernhart, S.H., Honer Zu Siederdissen, C., Tafer, H., Flamm, C., Stadler, P.F., Hofacker, I.L., 2011. ViennaRNA Package 2.0. Algorithms Mol Biol 6, 26.

MacArthur, J., Bowler, E., Cerezo, M., Gil, L., Hall, P., Hastings, E., Junkins, H., McMahon, A., Milano, A., Morales, J., Pendlington, Z.M., Welter, D., Burdett, T., Hindorff, L., Flicek, P., Cunningham, F., Parkinson, H., 2017. The new NHGRI-EBI Catalog of published genome-wide association studies (GWAS Catalog). Nucleic Acids Res 45, D896-D901.

Maharshak, N., Hart, G., Ron, E., Zelman, E., Sagiv, A., Arber, N., Brazowski, E., Margalit, R., Elinav, E., Shachar, I., 2010. CCL2 (pM levels) as a therapeutic agent in Inflammatory Bowel Disease models in mice. Inflamm Bowel Dis 16, 1496-1504.

Manolio, T.A., 2010. Genomewide association studies and assessment of the risk of disease. N Engl J Med 363, 166-176.

Mao, F., Xiao, L., Li, X., Liang, J., Teng, H., Cai, W., Sun, Z.S., 2016. RBP-Var: a database of functional
variants involved in regulation mediated by RNA-binding proteins. Nucleic Acids Res 44, D154-163.
Martin, J.C., Chang, C., Boschetti, G., Ungaro, R., Giri, M., Grout, J.A., Gettler, K., Chuang, L.S., Nayar,
S., Greenstein, A.J., Dubinsky, M., Walker, L., Leader, A., Fine, J.S., Whitehurst, C.E., Mbow, M.L.,
Kugathasan, S., Denson, L.A., Hyams, J.S., Friedman, J.R., Desai, P.T., Ko, H.M., Laface, I., Akturk, G.,
Schadt, E.E., Salmon, H., Gnjatic, S., Rahman, A.H., Merad, M., Cho, J.H., Kenigsberg, E., 2019. Single-
Cell Analysis of Crohn's Disease Lesions Identifies a Pathogenic Cellular Module Associated with
Resistance to Anti-TNF Therapy. Cell 178, 1493-1508 e1420.
Matsumiya, T., Ota, K., Imaizumi, T., Yoshida, H., Kimura, H., Satoh, K., 2010. Characterization of
synergistic induction of CX3CL1/fractalkine by TNF-alpha and IFN-gamma in vascular endothelial
cells: an essential role for TNF-alpha in post-transcriptional regulation of CX3CL1. J Immunol 184,
4205-4214.
Maurano, M.T., Humbert, R., Rynes, E., Thurman, R.E., Haugen, E., Wang, H., Reynolds, A.P.,
Sandstrom, R., Qu, H., Brody, J., Shafer, A., Neri, F., Lee, K., Kutyavin, T., Stehling-Sun, S., Johnson,
A.K., Canfield, T.K., Giste, E., Diegel, M., Bates, D., Hansen, R.S., Neph, S., Sabo, P.J., Heimfeld, S.,
Raubitschek, A., Ziegler, S., Cotsapas, C., Sotoodehnia, N., Glass, I., Sunyaev, S.R., Kaul, R.,
Stamatoyannopoulos, J.A., 2012. Systematic localization of common disease-associated variation in
regulatory DNA. Science 337, 1190-1195.
Mayr, C., 2019. What Are 3' UTRs Doing? Cold Spring Harb Perspect Biol 11.
McDermott, D.H., Yang, Q., Kathiresan, S., Cupples, L.A., Massaro, J.M., Keaney, J.F., Jr., Larson,
M.G., Vasan, R.S., Hirschhorn, J.N., O'Donnell, C.J., Murphy, P.M., Benjamin, E.J., 2005. CCL2
polymorphisms are associated with serum monocyte chemoattractant protein-1 levels and
myocardial infarction in the Framingham Heart Study. Circulation 112, 1113-1120.
Melgarejo, E., Medina, M.A., Sanchez-Jimenez, F., Urdiales, J.L., 2009. Monocyte chemoattractant
protein-1: a key mediator in inflammatory processes. Int J Biochem Cell Biol 41, 998-1001.
Morrison, A.R., Yarovinsky, T.O., Young, B.D., Moraes, F., Ross, T.D., Ceneri, N., Zhang, J., Zhuang,
Z.W., Sinusas, A.J., Pardi, R., Schwartz, M.A., Simons, M., Bender, J.R., 2014. Chemokine-coupled
beta2 integrin-induced macrophage Rac2-Myosin IIA interaction regulates VEGF-A mRNA stability
and arteriogenesis. J Exp Med 211, 1957-1968.
Mummidi, S., Bonello, G.B., Ahuja, S.K., 2009. Confirmation of differential binding of Interferon
Regulatory Factor-1 (IRF-1) to the functional and HIV disease-influencing -2578 A/G polymorphism in
CCL2. Genes Immun 10, 197-198; author reply 199.
Nayak, B.K., Feliers, D., Sudarshan, S., Friedrichs, W.E., Day, R.T., New, D.D., Fitzgerald, J.P., Eid, A.,
Denapoli, T., Parekh, D.J., Gorin, Y., Block, K., 2013. Stabilization of HIF-2alpha through redox
regulation of mTORC2 activation and initiation of mRNA translation. Oncogene 32, 3147-3155.
Nedelec, Y., Sanz, J., Baharian, G., Szpiech, Z.A., Pacis, A., Dumaine, A., Grenier, J.C., Freiman, A.,
Sams, A.J., Hebert, S., Page Sabourin, A., Luca, F., Blekhman, R., Hernandez, R.D., Pique-Regi, R.,
Tung, J., Yotova, V., Barreiro, L.B., 2016. Genetic Ancestry and Natural Selection Drive Population
Differences in Immune Responses to Pathogens. Cell 167, 657-669 e621.
Page, S.H., Wright, E.K., Jr., Gama, L., Clements, J.E., 2011. Regulation of CCL2 expression by an
upstream TALE homeodomain protein-binding site that synergizes with the site created by the A-
2578G SNP. PLoS One 6, e22052.
Pai, A.A., Cain, C.E., Mizrahi-Man, O., De Leon, S., Lewellen, N., Veyrieras, J.B., Degner, J.F., Gaffney,
D.J., Pickrell, J.K., Stephens, M., Pritchard, J.K., Gilad, Y., 2012. The contribution of RNA decay
quantitative trait loci to inter-individual variation in steady-state gene expression levels. PLoS Genet
8, e1003000.
Palmieri, O., Latiano, A., Salvatori, E., Valvano, M.R., Bossa, F., Latiano, T., Corritore, G., di Mauro, L.,
Andriulli, A., Annesec, V., 2010. The -A2518G polymorphism of monocyte chemoattractant protein-1
is associated with Crohn's disease. Am J Gastroenterol 105, 1586-1594.
Payne, J.L., Khalid, F., Wagner, A., 2018. RNA-mediated gene regulation is less evolvable than
transcriptional regulation. Proc Natl Acad Sci U S A 115, E3481-E3490.

Pham, M.H., Bonello, G.B., Castiblanco, J., Le, T., Sigala, J., He, W., Mummidi, S., 2012. The rs1024611
regulatory region polymorphism is associated with CCL2 allelic expression imbalance. PLoS One 7,
e49498.
Pullmann, R., Jr., Abdelmohsen, K., Lal, A., Martindale, J.L., Ladner, R.D., Gorospe, M., 2006.
Differential stability of thymidylate synthase 3'-untranslated region polymorphic variants regulated
by AUF1. J Biol Chem 281, 23456-23463.
Quach, H., Rotival, M., Pothlichet, J., Loh, Y.E., Dannemann, M., Zidane, N., Laval, G., Patin, E.,
Harmant, C., Lopez, M., Deschamps, M., Naffakh, N., Duffy, D., Coen, A., Leroux-Roels, G., Clement,
F., Boland, A., Deleuze, J.F., Kelso, J., Albert, M.L., Quintana-Murci, L., 2016. Genetic Adaptation and
Neandertal Admixture Shaped the Immune System of Human Populations. Cell 167, 643-656 e617.
Rabani, M., Levin, J.Z., Fan, L., Adiconis, X., Raychowdhury, R., Garber, M., Gnirke, A., Nusbaum, C.,
Hacohen, N., Friedman, N., Amit, I., Regev, A., 2011. Metabolic labeling of RNA uncovers principles of
RNA production and degradation dynamics in mammalian cells. Nat Biotechnol 29, 436-442.
Reyes, R., Alcalde, J., Izquierdo, J.M., 2009. Depletion of T-cell intracellular antigen proteins
promotes cell proliferation. Genome Biol 10, R87.
Ross, J., 1995. mRNA stability in mammalian cells. Microbiol Rev 59, 423-450.
Rovin, B.H., Lu, L., Saxena, R., 1999. A novel polymorphism in the MCP-1 gene regulatory region that
influences MCP-1 expression. Biochem Biophys Res Commun 259, 344-348.
Sasaki, Y., Dehnad, A., Fish, S., Sato, A., Jiang, J., Tian, J., Schroder, K., Brandes, R., Torok, N.J., 2017.
NOX4 Regulates CCR2 and CCL2 mRNA Stability in Alcoholic Liver Disease. Sci Rep 7, 46144.
Schott, J., Reitter, S., Philipp, J., Haneke, K., Schafer, H., Stoecklin, G., 2014. Translational regulation
of specific mRNAs controls feedback inhibition and survival during macrophage activation. PLoS
Genet 10, e1004368.
Schultz, C.W., Preet, R., Dhir, T., Dixon, D.A., Brody, J.R., 2020. Understanding and targeting the
disease-related RNA binding protein human antigen R (HuR). Wiley Interdiscip Rev RNA 11, e1581.
Schwerk, J., Savan, R., 2015. Translating the Untranslated Region. J Immunol 195, 2963-2971.
Sharif, O., Bolshakov, V.N., Raines, S., Newham, P., Perkins, N.D., 2007. Transcriptional profiling of
the LPS induced NF-kappaB response in macrophages. BMC Immunol 8, 1.
Shatoff, E., Bundschuh, R., 2020. Single nucleotide polymorphisms affect RNA-protein interactions at
a distance through modulation of RNA secondary structures. PLoS Comput Biol 16, e1007852.
Steri, M., Idda, M.L., Whalen, M.B., Orru, V., 2018. Genetic variants in mRNA untranslated regions.
Wiley Interdiscip Rev RNA 9, e1474.
Szalai, C., Kozma, G.T., Nagy, A., Bojszko, A., Krikovszky, D., Szabo, T., Falus, A., 2001. Polymorphism
in the gene regulatory region of MCP-1 is associated with asthma susceptibility and severity. J Allergy
Clin Immunol 108, 375-381.
Tam, V., Patel, N., Turcotte, M., Bosse, Y., Pare, G., Meyre, D., 2019. Benefits and limitations of
genome-wide association studies. Nat Rev Genet 20, 467-484.
Tu, X., Chong, W.P., Zhai, Y., Zhang, H., Zhang, F., Wang, S., Liu, W., Wei, M., Siu, N.H., Yang, H., Yang,
W., Cao, W., Lau, Y.L., He, F., Zhou, G., 2015. Functional polymorphisms of the CCL2 and MBL genes
cumulatively increase susceptibility to severe acute respiratory syndrome coronavirus infection. J
Infect 71, 101-109.
Tucci, M., Barnes, E.V., Sobel, E.S., Croker, B.P., Segal, M.S., Reeves, W.H., Richards, H.B., 2004.
Strong association of a functional polymorphism in the monocyte chemoattractant protein 1
promoter gene with lupus nephritis. Arthritis Rheum 50, 1842-1849.
Uren, P.J., Burns, S.C., Ruan, J., Singh, K.K., Smith, A.D., Penalva, L.O., 2011. Genomic analyses of the
RNA-binding protein Hu antigen R (HuR) identify a complex network of target genes and novel
characteristics of its binding sites. J Biol Chem 286, 37063-37066.
Vilmundarson, R.O., Duong, A., Soheili, F., Chen, H.H., Stewart, A.F.R., 2021. IRF2BP2 3'UTR
Polymorphism Increases Coronary Artery Calcification in Men. Front Cardiovasc Med 8, 687645.
Visscher, P.M., Wray, N.R., Zhang, Q., Sklar, P., McCarthy, M.I., Brown, M.A., Yang, J., 2017. 10 Years
of GWAS Discovery: Biology, Function, and Translation. Am J Hum Genet 101, 5-22.

Wang, J., Pitarque, M., Ingelman-Sundberg, M., 2006. 3'-UTR polymorphism in the human CYP2A6
gene affects mRNA stability and enzyme expression. Biochem Biophys Res Commun 340, 491-497.
Wang, Z., Kayikci, M., Briese, M., Zarnack, K., Luscombe, N.M., Rot, G., Zupan, B., Curk, T., Ule, J.,
2010. iCLIP predicts the dual splicing effects of TIA-RNA interactions. PLoS Biol 8, e1000530.
Wright, E.K., Jr., Page, S.H., Barber, S.A., Clements, J.E., 2008. Prep1/Pbx2 complexes regulate CCL2
expression through the -2578 guanine polymorphism. Genes Immun 9, 419-430.
Wu, X., Brewer, G., 2012. The regulation of mRNA stability in mammalian cells: 2.0. Gene 500, 10-21.
Xia, J., Bogardus, C., Prochazka, M., 1999. A type 2 diabetes-associated polymorphic ARE motif
affecting expression of PPP1R3 is involved in RNA-protein interactions. Mol Genet Metab 68, 48-55.
Yoshimura, T., Yuhki, N., Moore, S.K., Appella, E., Lerman, M.I., Leonard, E.J., 1989. Human monocyte
chemoattractant protein-1 (MCP-1). Full-length cDNA cloning, expression in mitogen-stimulated
blood mononuclear leukocytes, and sequence similarity to mouse competence gene JE. FEBS Lett
244, 487-493.
Yoshinaga, M., Takeuchi, O., 2019. Post-transcriptional control of immune responses and its
potential application. Clin Transl Immunology 8, e1063.
Zhai, Y., Zhong, Z., Chen, C.Y., Xia, Z., Song, L., Blackburn, M.R., Shyu, A.B., 2008. Coordinated
changes in mRNA turnover, translation, and RNA processing bodies in bronchial epithelial cells
following inflammatory stimulation. Mol Cell Biol 28, 7414-7426.
Zhang, F., Lupski, J.R., 2015. Non-coding genetic variants in human disease. Hum Mol Genet 24,
R102-110.
Zhang, X., Chen, X., Liu, Q., Zhang, S., Hu, W., 2017. Translation repression via modulation of the
cytoplasmic poly(A)-binding protein in the inflammatory response. Elife 6.
Zhao, W., Zhang, S., Zhu, Y., Xi, X., Bao, P., Ma, Z., Kapral, T.H., Chen, S., Zagrovic, B., Yang, Y.T., Lu,
Z.J., 2022. POSTAR3: an updated platform for exploring post-transcriptional regulation coordinated
by RNA-binding proteins. Nucleic Acids Res 50, D287-D294.
Zou, J., Hormozdiari, F., Jew, B., Castel, S.E., Lappalainen, T., Ernst, J., Sul, J.H., Eskin, E., 2019.
Leveraging allelic imbalance to refine fine-mapping for eQTL studies. PLoS Genet 15, e1008481.

1056

Figure 1

Figure 2

Figure 3

A

```
0  20  40  60  80  100 120 140 160 180 200 220 240 260 280 300 320 340 360
```

CTGCAGCTAACTTATTTTCCCCTAGCTTTCCCCAGACACCCTGTTTTATTTTATTATAATGAATTTTGTT

SNP
RBP

rs181021073   rs13900

ELAVL1
TIA1
TIAL1
MSI1

B

0 ▮▮▮▮ 3.3

C

bits

D

Relative preference

L   M   P   U

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

	CCL2 (rs13900 C/T) Control (42)
Wild allele [CC]	18 (42.8%)
Heterozygous [CT]	16 (38.09%)
Rare allele [TT]	8 (19.04%)
MAF	0.38
PIC	0.36
Heterozygosity	0.47
No of alleles	84
Hardy Weinberg P	0.459

Supplementary Table 1. Allele carriages and allele frequencies of rs13900 in healthy volunteers

	gDNA	Cytosol	Monosome	Polysome
D-1	0.975806	0.872946	0.873425	1.778429
D-2	1.034091	1.557778	1.441919	1.562622

Supplementary Table 2. Differential loading of rs13900 C>T onto polysome isolated from macrophages of heterozygous donors. C>T ratio < 1 indicates increased levels of the minor T allele relative to C allele. Result shows enrichment of rs13900T allele on polysomes.

Figure S1

Figure S2

Figure S3

A

Figure S5

A

B