Deployment of a *Vibrio cholerae* ordered transposon mutant library in a quorum-
competent genetic background

Nkrumah A. Grant[1,2,3*], Gracious Yoofi Donkor[1], Jordan T. Sontz,[4] William Soto[5], Christopher
M. Waters[2,3,4*]

[1]Department of Microbiology, University of Illinois Urbana-Champaign, Urbana, IL
[2]Department of Microbiology, Genetics, and Immunology, Michigan State University, East
Lansing, MI
[3]BEACON Center for the Study of Evolution in Action, Michigan State University, East
Lansing, MI
[4]MSU College of Osteopathic Medicine, Michigan State University, East Lansing, MI
[5]Department of Biology, College of William and Mary, Williamsburg, VA

[*]Correspondence to: grantnkr@illinois.edu or watersc3@msu.edu

Abstract

Vibrio cholerae, the causative agent of cholera, has sparked seven pandemics in recent centuries,
with the current one being the most prolonged. *V. cholerae's* pathogenesis hinges on its ability to
switch between low and high cell density gene regulatory states, enabling transmission between
host and the environment. Previously, a transposon mutant library for *V. cholerae* was created to
support investigations aimed toward uncovering the genetic determinants of its pathogenesis.
However, subsequent sequencing uncovered a mutation in the gene *luxO* of the parent strain,
rendering mutants unable to exhibit high cell density behaviors. In this study, we used chitin-
independent natural transformation to move transposon insertions from these low cell density
mutants into a wildtype genomic background. Library transfer was aided by a novel gDNA
extraction we developed using thymol, which also showed high lysis-specificity for *Vibrio*. The
resulting Grant Library comprises 3,102 unique transposon mutants, covering 79.8% of *V.
cholerae's* open reading frames. Whole genome sequencing of randomly selected mutants
demonstrates 100% precision in transposon transfer to cognate genomic positions of the recipient
strain. Notably, in no instance did the *luxO* mutation transfer into the wildtype background. Our
research uncovered density-dependent epistasis in growth on inosine, an immunomodulatory
metabolite secreted by gut bacteria that is implicated in enhancing gut barrier functions.
Additionally, Grant Library mutants retain the plasmid that enables rapid, scarless genomic
editing. In summary, the Grant Library reintroduces organismal relevant genetic contexts absent

in the low cell density locked library equivalent.

Significance Statement

Ordered transposon mutant libraries are essential tools for catalyzing research by providing

access to null mutants of all non-essential genes. Such a library was previously generated for

Vibrio cholerae, but whole genome sequencing revealed that this library was made using a

parent strain that is unable to exhibit cell-cell communication known as quorum sensing. Here,

we utilize natural competence combined with a novel, high-throughput genomic DNA extraction

method to regenerate the signaling incompetent *V. cholerae* ordered transposon mutant library in

quorum sensing competent strain. Our library provides researchers with a powerful tool to

understand *V. cholerae* biology within a genetic context that influences how it transitions from

an environmentally benign organism to a disease-causing pathogen.

Introduction

Vibrio cholerae is a human pathogen and the causative agent of the acute diarrheal disease

cholera. While cholera is a global threat, the disease is endemic in developing countries, where it

is estimated that there are between 1.3 to 4.0 million *V. cholerae* annual infections, resulting in

21,000 – 143,000 deaths (1, 2). Implementation of disease intervention strategies, including

increasing access to clean drinking water, provision of adequate healthcare infrastructure, and

educating citizens in communities at risk of disease spread, remains a top priority in the fight

toward eradicating cholera. As such, the Global Task Force on Cholera Control was founded in

2017 to support efforts aimed at reducing cholera burden by 90% before the year 2030 (3, 4).

To date there have been seven cholera pandemics. The seventh pandemic began in 1961

and is ongoing, marking it as the longest running cholera pandemic (5). *V. cholerae* strains are

grouped by surface antigens, including O polysaccharide, such that there are >200 serogroups

(6). Of these serogroups, only two, O1 and O139, host pandemic *V. cholerae* strains (7, 8).

Furthermore, pandemic strains from the O1 serogroup are further stratified into two biotypes

with the fifth and sixth pandemics caused by the Classical and the seventh caused by El Tor (9).

Like most vibrios, *V. cholerae* predominantly possesses two chromosomes (10, 11).

Whole genome sequencing of strains before and after the emergence of seventh pandemic *V.*

cholerae strains has proved to be an invaluable resource toward understanding its origin and

evolution. These analyses show *V. cholerae* El Tor originated from a nonpathogenic strain (12),

becoming a pathogen after acquiring two genomic islands (VSP-I and VSP-II), a prophage that
expresses cholera toxin (CTXφ), and 12 additional mutations that enhanced its transmissibility
(9, 13, 14). Recent work has also shown that these genomic changes contribute to seventh
pandemic *V. cholerae's* long-term persistence in the estuarian environments they occupy (15).
75

Key to El Tor's transition to a pathogen is that it is naturally competent, granting it the
ability to take up DNA from the environment which can be integrated into the genome by
homologous recombination (16). Natural competence is a highly regulated process, and four
environmental factors are key determinants for driving this process. These include nutrient
limitation, extracellular signaling molecules, high cell density, and the presence of chitin (17–
19), which together coordinate the expression of several regulators governing competence (20–
23). Recent work has shown that expressing the master regulator of competence TfoX can
circumvent the chitin requirement for natural competence in *V. cholerae* (19, 24). This discovery
has allowed exploitation of chitin-independent natural competence as a genetic tool for editing
Vibrio spp. genomes (25–27), facilitating our understanding of the functional relationships
between genes and furthering our understanding of how *V. cholerae* interfaces with its
environment.
88

Our understanding of *V cholerae's* ecology and evolution has also benefited from
transposon mutagenesis. In contrast to conventionally laborious genetic engineering approaches
for disrupting single genes to study their phenotypic effects, transposon mutagenesis allows the
simultaneous construction and screening of tens of thousands of mutants. Researchers have taken
advantage of transposon mutagenesis to construct nonredundant (ordered) mutant libraries,
where each nonessential gene has been inactivated and the individual mutants are arrayed across
the wells of 96-well microtiter plates (28–31). An ordered mutant *V. cholerae* library was
constructed in 2008 (32). The ordered library consists of >3,100 mutants, each with a Tn5-based
transposon insertion that confers kanamycin resistance. The mutant library has provided a public
resource enhancing *V. cholerae* genotypic and phenotypic investigations, supporting identifying
novel therapeutic targets for vaccine development (33), and increasing our understanding of the
V. cholerae virulome (34, 35), amongst other discoveries (36, 37).
101

Although significant research advances have been made in our understanding of *V.*
cholerae using mutants from this ordered mutant transposon library, the library was constructed
in a laboratory acquired quorum sensing mutant of *V. cholerae*. Quorum sensing, the process of
cell-to-cell communication in bacteria, allows cells to undergo global changes in gene expression

as the bacteria transition from low to high cell density. This transition can influence behaviors

like host immunity escape (38), virulence factor production (39–41), biofilm formation (42),

infectivity and environmental dissemination (43), and protection against phage predation (44).

This transition is mediated by extracellular autoinducers (45, 46) which impact the activity of

cytoplasmic regulators, including LuxO and HapR.

In the low-cell-density state, LuxO is phosphorylated leading to inhibition of *hapR*

mRNA translation via regulatory sRNAs, whereas at high-cell-density, LuxO is

dephosphorylated and inactive, resulting in HapR protein translation and the switch to high-cell-

density (41, 47). Recently, a presumably laboratory acquired mutation in *luxO, luxOG319S* was

discovered in some laboratory isolates of the widely used *V. cholerae* strain C6706 that locked

this strain in the low-cell-density state (48). After sequence validating mutants from the 2008

library, we observed the *luxO*G319S mutation, indicating it was present in the parent strain from

which the library was constructed. Thus, the 2008 ordered mutant library (32) is only partially

representative of how *V. cholerae* senses and responds to its environment and a new library in a

quorum sensing competent strain would greatly enhance our understanding of *V. cholerae*

ecology and evolution.

To overcome this limitation of the 2008 library, we use chitin-independent natural

competence to regenerate the defective ordered *V. cholerae* library in a quorum competent

C6706 genomic background. To streamline the transfer process, we developed a cost effective

gDNA extraction method using thymol (49). Compared to other methods, our approach is both

economical ($0.64 per reaction vs $2.85 per reaction) and efficient, enabling up to 200 gDNA

extractions in as little as 2.5 hours, in contrast to 25 reactions requiring >6 hours using

conventional methods. Using our methods, we successfully regenerated the library with a 99%

success rate, as indicated by growth on antibiotic selection growth media. Sequence validation of

randomly selected mutants from the regenerated library show that the transposon insertions

recombine into our recipient strain with 100% precision and transfer of the *luxOG319S* mutation

never occurred. Of significance, assays performed using paired transposon mutants from each

library provide evidence for density-dependent epistasis in several flagellar genes, and when

grown on inosine – a metabolite secreted by gut bacteria – illustrating the utility of this new

library. Lastly, we show that strains from the library generated in this work – hereafter the Grant

Library – can undergo additional edits using chitin-independent natural competence. Taken

together, the Grant Library represents an added contribution in our efforts toward understanding

V. cholerae behaviors under transcriptional regulatory states that are representative of how this
pathogen evolves and interfaces with its environment.
142
143   **Materials and Methods**
144
Bacterial matings
To generate the library recipient strain used in this study, NG001, we mated wildtype *V. cholerae*
C6706 with an *E. coli* S17-λpir donor strain harboring plasmid pMMB-*tfoX-qstR* (generously
provided by Ankur Dalia, Indiana University). In short, we mixed equal volumes of donor and
recipient strains in the center of a Luria-Bertani agar plate which we incubated at 37°C for three
hours. Afterwards, we added 1 mL of LB media containing chloramphenicol (for selection of
plasmid in recipient strain) and polymyxin B (for counterselection of the donor *E. coli* strain) to
the plate, washed, and collected the mating spot into a borosilicate glass culture tube, which we
incubated overnight at 37°C with orbital shaking at 210 RPM.
154
155   **Media and buffers**
156
157   *Thymol cell lysis agent*
158   We prepared thymol at a stock concentration of 350 mM by dissolving thymol crystals in
159   dimethyl sulfoxide (DMSO). Thymol stocks were made in 15 mL which we stored at room
160   temperature on the lab bench until they were exhausted. We observed no differences in the lytic
161   activity of thymol during storage.
162
gDNA extraction buffer
We prepared solutions of a gDNA extraction buffer in two 500 mL volume parts. We prepared
solution one by mixing 80 mL of 1M Tris (pH 8.0), 56 mL of 0.5M EDTA (pH 8.0), 46.7 g
NaCl, and 3 g sodium metabisulfite, which we stirred until dissolved. We then adjusted this
solution to the final volume with dH_2O and autoclaved for 25 minutes. We prepared solution
two, sodium acetate (2.5M, pH 5.2), by dissolving 102.5 g of the salt in dH_2O, which we
sterilized using filtration. Prior to each extraction, we prepared a master mixture of solution one
and solution two by mixing 21 mL and 31.5 mL for every 100 samples processed, respectively.
171
Minimal media with inosine
We prepared inosine at a stock concentration of 133 mM by mixing the compound in sterile
dH_2O. We then applied gentle heating and stirred the mixture until the solution was homogenous,

which we then filter sterilized. We prepared 1x M9 minimal media amended with 0.1 mM $CaCl_2$,

and 2 mM $MgSO_4$, to which we added inosine to a 20 mM final concentration.

Strain revival

We revived strains from the nonredundant library ((32); henceforth referred to as donor library)

in microplate format to maintain parity between it and that produced in this work. The donor

library was generously provided by the laboratories of Vic DiRita (plates: 2, 4-6, 8-10, 12-17,

19-33), Michigan State University, and Bonnie Bassler (plates: 1, 3, 7, 11, 18, 34), Princeton

University. Briefly, microtiter plates from the donor library were slightly thawed and 10 µl from

each well was inoculated into a deep-well microplate containing 550 µl of selective media

(Luria-Bertani broth + Kanamycin (100 µg/ml) in each of the 96 wells. We revived the recipient

library strain, NG001, in a 250 ml baffled flask containing 25 ml of selective media (Luria-

Bertani broth + chloramphenicol (10 µg/ml)) to maintain plasmid *pmmB-tfoX-qstR* and 100

µg/ml isopropyl β-D-1-thiogalactopyranoside (IPTG) to induce natural competence. Unless

noted otherwise, we grew all cultures overnight (18 – 24h) at 37°C with orbital shaking at 210

RPM.

Whole gDNA preparation

The non-redundant donor library is comprised of 35 96-well plates of *Vibrio cholerae* mutants.

To extract genomic DNA from the donor library in a cost efficient and high-throughput fashion,

we developed an extraction method using a modified recipe of the gDNA extraction buffer

developed by Mantel and Sweigart, 2019 (accessed September, 2022, at

https://www.protocols.io/view/quick-amp-dirty-dna-extraction-4r3l287zql1y/v1). In brief, we

pelleted cells from overnight cultures of the donor library by centrifuging the deep-well plates at

4°C for 20 minutes. We then removed ~500 µl of supernatant from each well, resuspended the

cell pellet in 10 µl thymol (~35 mM final concentration), and incubated the plates at 64°C for 15

minutes. Thereafter, we added 500 µl of gDNA buffer to each well and incubated the plates at

64°C for an additional 45 minutes.

To clear culture supernatants of cellular debris, we transferred cell lysates from each well

into an E-Z 96™ Lysate Clearance Plate (https://www.omegabiotek.com/product/e-z-96-lysate-

clearance-plate/). We then used a vacuum manifold to filter the supernatant through the plate,

which we collected in a second deep-well plate containing 200 µl of 100% cold isopropanol per

well. We then incubated the plates for 15 minutes at room temperature to precipitate genomic

DNA. Afterwards, we again used a vacuum manifold to filter the cell lysate containing our

gDNA through an E-Z 96™ DNA plate (https://www.omegabiotek.com/product/e-z-96-dna-
plates/?cn-reloaded=1). According to the manufacturer, "E-Z 96™ plates are silica glass fiber
plates that can bind up to 50 µg of genomic DNA or 20 µg of plasmid DNA per prep." After this
initial filtration, we washed our bound DNA, in sequence, using 500 µl of 70% and 95% ice cold
ethanol. We then added 100 µl of 1x TE buffer to each well and eluted genomic DNA from the
DNA plates into PCR plates by centrifugation (max speed, one minute).
216

Chitin-independent natural competence for gDNA transfer
After 18-24 h of overnight growth, we diluted the library recipient strain (NG001) 1:4 in 0.5x
instant ocean to which we added chloramphenicol (5 µg/ml) and IPTG (100 µg/ml). We then
added 200 µl of these competent cells to each well of a deep-well 96-well plate and added the
gDNA extracted from the donor library to a matched recipient well (final dilution 1:3). We then
incubated the deep-well plates overnight in a standing incubator at 30°C. Following the
overnight incubation, we added 400 µl of selective media (Luria-Bertani broth +
chloramphenicol (5 µg/mL) + kanamycin (100 µg/mL)) to each well and incubated the plates at
37°C in a standing incubator for 48 h (Fig. 1). We determined transformation success by spot
plating 3 µl of each well onto selective LB agar plates (Luria-Bertani agar + chloramphenicol (5
µg/mL) + kanamycin (100 µg/mL)), which we grew overnight at 37°C (Fig. S1).
228

Grant Library transformation efficiency analysis
We created a comprehensive plate indices file (Dataset S1) covering all possible wells within the
donor library (P1A1 to P34D6; 3,210 wells). Using this file, we developed a script to compare it
with a modified version of supplementary dataset 3 from Cameron et al., 2008 (Dataset S2),
which allowed us to identify empty wells (86) and those containing mutants (3,124). Our
comparative analysis also flagged duplicate entries in supplementary dataset 3 of the donor
library (Dataset S3).
236

While constructing the Grant Library, we meticulously documented the outcomes of
growth on selective media following chitin-independent natural transformation of the donor
library: True Positives (3,102 wells), False Negatives (22 wells expected to grow but didn't),
True Negatives (71 blank wells that didn't transfer), and False Positives (15 blank wells that
transferred unexpectedly). These values served as inputs for a binary confusion matrix (50),
providing a comprehensive assessment.
243

244   **TnFGL3 transposon insertion site identification**

Genome Sequencing

We randomly selected 23 mutants from the Grant Library to identify the transposon insertion
site. Strains were incubated overnight in 3 ml of LB + Kanamycin (100 µg/ml) and DNA was
extracted from 1 ml of culture using a Wizard ® Genomic DNA Purification Kit following the
manufacturer's instructions. DNA was rehydrated overnight at 4°C in 50 µl of TE buffer. Library
preparation was performed by SeqCoast Genomics using an Illumina DNA Prep tagmentation kit
and unique dual indexes. Samples were multiplexed and sequenced on the Illumina NextSeq2000
platform using a 300-cycle flow cell kit to produce 2x150 bp paired reads. Read demultiplexing,
read trimming, and run analytics were performed using DRAGEN v3.10.11, an on-board analysis
software on the NextSeq2000. Raw sequencing data generated during this study are available in
the following public Github repository (https://github.com/NkrumahG/Grant-Library-
Construction)

Reference genomes

The reference genomes for *V. cholerae* strains used in this study were downloaded from NCBI
GenBank. The chromosome accession numbers for chromosomes I and II are, NZ_CP028827.1
(2,975,504 bp) and NZ_CP028828.1 (1,072,331 bp) for strain N16961, and NZ_CP046844
(3,019,938 bp) and NZ_CP046845 (1,070,359 bp) for strain C6706, respectively. The sequence
map for plasmid pSC189 was downloaded from addgene
(https://www.addgene.org/browse/sequence/36279/), and the sequence for pMMB-*tfoX-qstR* was
provided by Ankur Dalia (Indiana University).

Mutation calling

We used consensus mode in *breseq* ((51), version 0.37) to align the paired-end reads for each of
the sequenced mutants to the reference genomes. The computational resources used to perform
breseq were provided by the Institute of Cyber-Enabled Research at Michigan State University.

Ortholog identification

We used Mauve ((52), snapshot _2015-02-25) to identify orthologs in whole genome sequences
of C6706 and N16961. Each pair of chromosomes were aligned using the progressive Mauve
method with default settings. In the Supplemental Information, we have included lists of all
orthologous gene pairs in *V. cholerae* N16961 and C6706 (Dataset 4) and the identity of genes
unique to N16961 (Dataset S5) and C6706 (Dataset S6). Additionally, we used Geneious to

279 extract, align, and visually display (Supplementary Information, Fig. S2 and S3) orthologous
280 gene pairs for the transposon mutants sequenced in this study.
281
282 **LuxO gain of function mutation identification**
283 Using supplementary dataset 3 provided for the donor library (32), we identified all potential
284 transposon mutants located within a 50 kb range, both upstream and downstream of *luxO*.
285 Following this selection, we cultured these 92 mutants in selective media (LB + Kanamycin, 100
286 μg/mL) for 24 hours under standard conditions.
287
After the incubation, we extracted gDNA using the thymol-based gDNA extraction
method we developed to construct the Grant Library. The extracted genomic DNA was then
diluted 1:2, which we used as template for PCR amplification of the *luxO* gene. The *luxO* gene
was amplified with Q5 polymerase using the following primers: forward primer 5`
GGCTATGCAACATAATCAATCTTG-3` and reverse primer 5`-
GCTTTGGTTGATCCATTCTCTCAT-3`. PCR conditions included an initial denaturation at
98℃ for 2 minutes, followed by 29 cycles of denaturing at 98°C for 30 seconds, annealing at
63℃ for 30 seconds, and extension at 72°C for 1 minute. A final extension step was
performed at 72°C for 5 minutes.
297
Following the PCR amplification, we validated amplification of *luxO* using gel
electrophoresis. We then used the Qiaquick PCR purification system to purify our *luxO* PCR
product. Sanger sequencing of the purified *luxO* allele was performed using the reverse primer
5`-TGCGATAGATGGTTGACGGG-3` by the Roy J. Carver Biotechnology Center at the
University of Illinois at Urbana-Champaign. We imported the Sanger Sequence read data into
Geneious (53) and aligned it to *luxO* (*V. cholerae* chromosome I, C6706), allowing
comprehensive analysis and visualization of our sequenced mutants, with special attention to
amino acid position 319, the site of the *luxO* gain of function mutation *luxOG319S* (48).
306
Motility Assay
We screened mutants with presumptive TnFLG3 transposon insertions in known motility genes.
Strains were grown in a deep-welled 96-well plate overnight at 37°C in selective media (LB +
Kanamycin) and on the following day, we used toothpicks to manually stab each strain into
motility plates containing LB and 0.35% agar. Plates were incubated at 37°C for 24 h, and the
motility zones were recorded with a gel imager. The area of the motility zones was measured

computationally using the Hough Circle Transform package from the UCB vision plugin in Fiji

(54).

Inosine Growth Assay

We randomly selected 5-paired 96-well plates from each of the Grant and Cameron et al.

transposon mutant libraries and incubated them overnight in LB plus kanamycin (100 µg/mL).

We also revived the parent strains used to construct each library in the same fashion. On the

following day, we recorded mutants that didn't grow overnight and then back diluted each well

from the plates 1:100 into M9 minimal media supplemented with 20 mM inosine. We incubated

all plates with linear shaking at 37°C. We monitored growth of the parent strains at an OD_{600} for

10 hours, with measurements taken at 5-minute intervals. For the paired library mutants, we

measured the 10 h endpoint optical density.

Plasmid Curing Assay

We inoculated three cultures of NG001 from frozen glycerol stocks into LB media supplemented

with chloramphenicol (10 µg/ml) and incubated them overnight. On the following day, we back

diluted each culture 1000-fold into either LB, LB media containing chloramphenicol (10 µg/ml),

or LB containing IPTG (100 µM). For the duration of the experiment, we passaged each culture

in the same media. In parallel, we serially diluted stationary phase cultures in 0.5x Instant Ocean

and spot plated 3.0 µl from each dilution onto LB or LB agar containing chloramphenicol. The

percent of population with the *pMMB-tfoX-qstR* plasmid (plasmid carriage) was calculated as a

ratio of the dilutions where colonies grew on LB with chloramphenicol relative to growth on LB

alone.

Data accessibility

All data generated, and analysis scripts used in this work are available in the GitHub repository

at https://github.com/NkrumahG/Grant-Library-Construction.

Results and Discussion

Chitin-independent transfer of transposons from donor library strains into a quorum-

competent *V. cholerae* genetic background is remarkably efficient. We hypothesized that the

transposon insertions within the genomes of the quorum-incompetent *V. cholerae* strain

generated in Cameron et. al., 2008 could be transferred to a quorum-competent wildtype *V.*

cholerae strain using natural competence. To test this, we extracted gDNA from each mutant in

donor library using an in-house method involving thymol lysis and alcohol precipitation.

Subsequently, we co-incubated this gDNA with a wildtype *V. cholerae* strain (C6706) carrying a

plasmid that expresses the natural competence master regulator, TfoX, upon induction with IPTG

(Fig. 1, Materials and Methods).

The donor library consists of 34 plates arrayed from P1A1 to P34D6, for a total 3,210

possible wells containing transposon mutants. Upon examination of supplementary data table 3

for the donor library, we discovered that 86 wells were not listed, and some of the mutants were

recorded multiple times. With these considerations, we expected growth for 3,124 unique

transposon mutants after conducting our natural transformation protocol.

To verify the presence of transposon insertions in the Grant library mutants, we spot-

plated each mutant on LB medium supplemented with kanamycin. Of the 3,124 unique

transposon mutants we expected to grow on the plates, 3,102 mutants (99.2%) exhibited growth

(Fig. S1). None of the 22 missing mutants, which were expected to grow, showed growth in

selective media during overnight revival of donor library freezer stock. This suggests that these

mutants might no longer viable perhaps due to low culture density upon freezing or death

resulting from repeated freeze-thaw cycling during long-term storage of the donor library. In the

cases where we anticipated no growth, 71/86 (82.0%) did not grow on selective media.

To evaluate the performance of our library transfer approach, we utilized a confusion

matrix (Fig. 2). The results of this analysis indicate that our approach was highly accurate and

precise, with a high recall (F1 score = 0.9941) and a low False Discovery Rate (0.0070). These

results unequivocally demonstrate the remarkable efficiency of using chitin-independent natural

competence to transfer the transposons into a quorum-competent strain. The transference process

was made possible through our in-house gDNA extraction method, involving thymol cell lysis

and alcohol precipitation, affirming the suitability of this gDNA preparation for downstream

applications. Indeed, this suitability was confirmed when we employed thymol-extracted DNA

as a template in PCR, with no observed inhibitions.

Transposons from the donor library homologously recombine into the recipient genome

with 100% precision. The frequency of genome edits by homologous recombination in *Vibrio*

spp. is highest when there are at least 2 kB arms of homology flanking the genomic sequence

being altered (25). Given this requirement, the observation that the recipient strains grew after

incubation on selective media (Fig. S1) was suggestive that transposons correctly integrated into
the homologous site of the recipient's genome. To validate our supposition, we randomly
selected 23 mutants, isolated gDNA, and sequenced whole genomes. Subsequently, we aligned
the sequenced reads to four reference sequences using breseq (51). The reference sequences
included *V. cholerae* chromosomes I and II, and the plasmid conferring natural competence. A
fourth reference sequence (pSC189) containing sequences associated with the transposon was
also included as a "genomic lure" to capture reads that mapped to the transposon. When
examining the breseq output, we specifically looked for instances where new junctions were
created between chromosomes I or II (depending on which chromosome the gene targeted for
homologous recombination was located) and pSC189. Such a junction would indicate a
transposon integration event. Fig. 3 is an example breseq output from one such alignment, which
reports the transposon insertion site in chromosome II, including the basepair position it is
located.
396
The parent strain of both the donor and recipient libraries is *V. cholerae* El Tor C6706.
However, when the donor library was generated in 2008, transposon insertion sites were mapped
to reference genomes of *V. cholerae* El Tor strain N16961 because it was the only *V. cholerae*
that had been sequenced for *Vibrio spp.* at that time (55). In this work, we mapped transposon
insertion sites to reference genomes of strain C6706. To identify whether transposons insertions
were in the correct site of our recipient strains, we performed whole genomic comparative
analysis of *V. cholerae* N16961 and C6706. When we performed this analysis, we found
chromosomes I and II differ in size between strains and are not syntenic (Fig. 4). Out of the
2,649 genes annotated in our reference genomes for chromosome I, 2,558 (96.6%) of genes are
shared between N16961 and C6706. Additionally, 21 genes (0.79%) are unique to N16961 and
70 (2.64%) to C6706. On chromosome II, 1,052 out of 1,064 annotated genes (98.9%) are shared
between strains, where 10 (0.94%) and 2 (0.19%) of genes are unique to N16961 and C6706,
respectively. Importantly, this analysis provided the genomic position for each ortholog of
N16961 genes present in C6706 (Dataset S1).
411
We used the orthologous sites to analyze each transposon mutant we sequenced (Dataset
S1, Fig. S2 and S3). Using the breseq annotation data, we calculated the position of the
transposon within each gene as a percent of basepairs into the ORF of C6706. These position
values were previously reported for each strain in the donor library strain N16961. To assess the
relationship, we performed a linear regression of our sequenced C6706 insertions with the
previously described N16961 from the donor library, expecting a perfect linear correlation.

Although 19/23 mutants had an exact match, some of our values diverged from the line of best fit

(Fig. 5), although there was a strong statistical correlation between the variables (Kendall's

coefficient $\tau = 0.752$, $n = 23$, and $P < 0.0001$). Upon further investigation of this anomaly, we

discovered two categories of false negatives. In category one (Fig. S3, Panels A and B), we

observed two cases where genes were annotated in N16961 but not in C6706. Consequently,

these data points lacked a calculated value for the transposon insertion site in C6706 and thus fell

on the y-axis. In the second category, two false negatives were attributed to a difference in the

size of the gene orthologs between N16961 and C6706 (Fig. S3, Panels C and D). This

discrepancy resulted in a shift of the relative position reported for the transposon. After

accounting for these deviations and correcting the data accordingly, we confirmed that the

transposon inserted into the cognate position of the Grant Library mutants we sequenced with

100% precision.

Co-transformation of the *luxOG319S* gain of function mutation does not occur. Since we

employed natural competence to transfer the transposon mutations, there is a possibility that the

luxOG319S mutation from the donor library might have been co-transformed along with the

transposon into the Grant library. However, unlike the strong selection pressure for the

transposon mutation, selection for the *luxO* mutation would be indirect and weaker, depending

on whether the *luxO* mutation provides a fitness advantage to cells allowing them to outcompete

wildtype cells during outgrowth in selective media.

In the 23 genome sequences analyzed above, none had the *luxOG319S* mutation.

However, we did consider that the frequency of transfer of *luxO* mutation could be elevated in

genes for which the transposon is within close genomic proximity to *luxO*. To test this, we

isolated gDNA from all 94 mutants within a 50kb window up- and downstream of *luxO* (Dataset

S7). We then PCR amplified *luxO* and performed Sanger sequencing. In no instance did the

luxOG319S gain of function mutation transfer from the donor to the recipient library parent

strain (Dataset S8). This suggests that proximity of the transposon mutations to *luxO* does not

influence transfer of the gain-of-function mutation, however we do recommend sequencing *luxO*

as a best practice when drawing comparisons between donor and recipient library strains.

Natural competence in Grant library strains is IPTG inducible. When generating the Grant

library, we supplemented the outgrowth media with chloramphenicol and kanamycin to select for

maintenance of the plasmid conferring IPTG-induced natural competence, pMMB-*tfoX-qstR*, and

the TnFLG3 transposon, respectively. We reasoned that a library established with strains

harboring this plasmid would be advantageous to the research community as it would facilitate

additional genome edits using chitin-independent natural competence and homologous

recombination upon induction with IPTG. In proof of concept for this idea, we randomly selected

10 mutants from the Grant library and performed a knock-in experiment of a trimethoprim (tm)

resistance gene. In short, we used thymol to isolate whole gDNA from a tm resistant strain and

added 16.2 µg of whole gDNA to our competent cells, which we prepared following our protocol

described in Materials and Methods.

After incubating our competent cells with gDNA overnight, we used spot plates to assess

antibiotic phenotypes. All the randomly selected library mutants grew on LB plates

supplemented with trimethoprim between dilution factors of $10^{-3} – 10^{-8}$ (Fig. 6). When also

plated on growth media supplemented with kanamycin, there was parity in the growth

phenotypes observed on tm plates for all but one mutant (strain 8C5). When we examined the

images of the kanamycin supplemented spot plates when the library was constructed (Fig. S1),

we found mutant 8C5 didn't grow there either. We posit that the spurious growth of strain 8C5

likely reflects a frozen subpopulation of our library recipient strain that was able to withstand

kanamycin during outgrowth or contamination introduced during the spot-plate assay. Taken

together, all the library mutants we tested could undergo additional genome edits with overnight

induction of natural competence.

Plasmid conferring natural competence is readily cured from Grant library strains with

IPTG induction. Selecting for Grant library transposon insertion mutants retaining the pMMB-

tfoX-qstR plasmid allows for additional genomic edits to be constructed in each strain using

chitin-independent natural competence. However, plasmid carriage might impose fitness costs

that can vary between mutants due to genetic conflicts with each disrupted gene or complicate

efforts to introduce alternative plasmids. Taking this into account, we sought to establish whether

we could cure Grant library strains of the plasmid. To this end, we serially passaged the recipient

strain (NG001) carrying the plasmid in 1) LB media, 2) LB supplemented with chloramphenicol

and not IPTG, or 3) LB without chloramphenicol but supplemented with IPTG. We hypothesized

that amplifying plasmid carriage cost by inducing its expression in the absence of antibiotic

selection (treatment three) would lead to rapid curing.

In agreement with our hypothesis, the number of cells carrying the plasmid decreased by

≈ 94% after 120 h of serial passage in LB media supplemented with IPTG alone (Fig. 7). In

contrast, plasmid carriage in cells passaged in LB alone or in LB containing chloramphenicol

decreased by ≈ 39% and ≈ 11%, respectively. These data show that the pMMB-*tfox-qstR* plasmid
can be readily cured from Grant library strains.

Differential motility responses of recipient library mutants reveal unappreciated epistasis
exists in donor library. *V. cholerae* exhibits high motility by virtue of a single polar flagellum,
a characteristic that contributes significantly to its pathogenesis. *V. cholerae* isolated from active
infections display increased motility compared to lab-grown strains (56) and demonstrated a
heightened ability to colonize hosts (57). Moreover, motility allows *V. cholerae* to survive in its
aquatic environment, enabling it to swim freely as planktonic cells or form biofilms in response
to environmental stress. Furthermore, previous work has demonstrated the involvement of at
least 40 genes in flagellum biosynthesis and motility, classified into temporally distinct classes (I
through IV) (58–60). Notably, some of these genes are known to trigger an immune response
(56).

Motility is an easily screened phenotype, and the functional consequences of null
flagellar gene mutants are well established. Therefore, Cameron et al., 2008 (32), employed
motility assays as one of the validation metrics for their constructed library. In this context, we
also measured motility of the Grant library null flagellar gene mutants. By performing the assay
using paired mutants from the donor and recipient libraries, we reasoned that we would foremost
independently confirm the motility phenotypes reported by Cameron et. al., 2008. Furthermore,
previous studies have indicated that genes in classes III and IV of the flagellum biosynthesis
hierarchy exhibit cell-density dependent phenotypes (61). Thus, the motility assay would serve to
provide supporting evidence for or against our hypothesis that quorum sensing epistasis is a
missing feature of the donor library.

Out of the 33 genes examined, we successfully replicated the motility phenotype reported
for the donor library's nonmotile mutants in all but one strain (*fliR*) (Fig. 8). Additionally, we
observed that both libraries exhibited nonmotile behavior for *rpoN* (class I) and *flaA,* (class III)
that are essential for expression of the flagellar biosynthesis genes (Fig. 8). Furthermore, motility
was also abolished for the σ-54 dependent transcriptional regulator transposon mutants of *flrA*
(class I) in both libraries, consistent with previous observations (62). When comparing motility
within classes, paired null mutants from the Cameron and Grant libraries displayed similar
movement on the plates for all classes, except for class III, where a statistically significant
difference was observed (Table 1, two-tailed *t*-test: $P = 0.012$). Additionally, two mutant pairs
(*flgM* and *flgN*) within class IV biosynthesis genes exhibited opposite motility phenotypes.

While these opposing effects were observed within the same class, the net effect of class IV
genes canceled each other out in the statistical analysis, so this class comparison was
insignificant. Taken together, our results suggest quorum sensing epistasis with motility genes
results in the different motility outcomes as the donor library is locked in a low-cell density
genomic background.
528
Growth on inosine is quorum dependent. Inosine is an essential nucleoside that plays an
important role in purine biosynthesis and gene translation (63). Recent studies have
demonstrated that various species of gut bacteria, including *Lactobacillus* (64) and
Bifidobacterium, secrete inosine in significant quantities during growth (65). This production not
only contributes to systemic immunomodulatory and protective functions (66) but also enhances
mucosal barrier functions within the gut (65). Additionally, inosine has been found to inhibit the
expansion of pathogenic *Enterobacteriaceae* through PPARγ activation, further highlighting its
importance in preventing gut dysbiosis (67).
537
In an unrelated screen analyzing for quorum-dependent growth effects using Phenotype
Microarrays for Microorganisms (Biolog), we observed wild-type C6706 and a locked high-cell
density (Δ*luxO*) mutant exhibited enhanced growth when grown on inosine as the sole carbon
source compared to a locked low-cell density mutant (Δ*hapR*). As that the *luxOG319S* gain-of-
function mutant locks the Cameron library into the low-cell density state, we reasoned screening
of mutants in the Grant library for growth on inosine could yield previously unappreciated
epistatic effects between quorum sensing and central metabolism. Given *V. cholerae's*
manifestation of disease as an enteric pathogen – where inosine production is high – we sought
to investigate whether these differences extended to other mutants from the recipient and donor
transposon libraries. This exploration may shed light on the pathogenesis of *V. cholerae* and its
interaction with inosine metabolism. Moreover, any significant differences between the
transposon mutants would underscore the importance of using a library that doesn't mask
phenotypic states relevant to *V.* cholerae's ecology and evolution.
551
We began our investigation by Sanger sequencing the *lux*O locus of the parent strains for
the donor and recipient libraries. The sequence chromatograms (Fig. 8*A*) validated the glycine-
to-serine gain-of-function mutation at amino acid position 319 in the donor library parent. We
then grew replicate cultures of the parent strains in M9 minimal media supplemented with 20
mM inosine (Dataset S9). There was no significant difference in the growth rates of the parent
strains. However, the recipient library parent strain bearing the ancestral *luxO* allele grew to an

endpoint optical density that was on average ≈ 23% higher than the donor library parent strain

(Fig. 8*B*) and the relationship was statistically significant (Two-Sample unpaired $t = 29.34$, $df =$

16, and $P < 0.0001$).

We then randomly selected and revived five-paired plates from the donor and recipient

libraries and grew them in inosine for 24 h (Dataset S10). In agreement with the experiments

performed with the parent strains, the Grant library transposon mutants grew to an endpoint

optical density that was 35% higher than the Cameron et al. library (Fig. 8*C*) and the relationship

was statistically significant (Two-Sample unpaired $t = -20.34$, $df = 904$, and $P < 0.0001$).

Reaction norms describe the phenotypic expression of a single genotype across an

environmental gradient (68). Given our experimental design and the isogenicity of the donor and

recipient strains, we adopted this method to visualize deviations between paired library mutants

grown on inosine (Fig. 9*C*, grey lines). We focused our attention on mutants whose growth on

inosine were above and below two standard deviations (top and bottom 2.5% of the data,

respectively) away from the mean for each library (Fig. 10).

The data shows strong evidence of quorum dependent epistasis. When we looked at the

bottom 2.5% of all the data, seven of the thirteen transposon mutants were shared between

libraries and each failed to grow on inosine (Tables S1 – S2). The functional roles for the

associated genes included amino acid biosynthesis, energy metabolism, and regulatory functions.

That these mutants failed to grow independent of the *luxO* genetic background likely reflects the

essentiality of these genes when inosine is used as the sole carbon source. For the remaining six

genes, five grew better on inosine in the Grant Library genomic background and one marginally

less. Interestingly, when examining the top 2.5% of the data for Cameron or Grant Library paired

mutants, there was strong anticorrelation. That is, where growth was high for a mutant in one

library, its growth was lower in the opposite. In stark contrast to the bottom 2.5% of genes, only

one transposon mutant (gene disruption of *fusA-2*), was shared within the top 2.5% of each

dataset, and there was more variation in the functional roles of all called mutants (Tables S3 –

S4). We hypothesize that the non-linearity in the top 2.5% is due to differences in the

transcription of genes at low and high cell densities.

Collectively, our inosine findings show that ecologically relevant phenotypes can be

masked for *V. cholerae* in quorum dependent contexts and are supported by several other

research groups who as well have observed nuanced phenotypes in the *luxOG319S* genetic

background (48).

Conclusion

Bacteria employ quorum sensing (QS) to regulate gene expression based on population density.

In the case of *V. cholerae*, the etiological agent of cholera, QS controls various phenotypes as the

population transitions between low and high cell density. Researchers previously constructed an

ordered mutant library, systematically disrupting every non-essential gene in *V. cholerae* with a

transposon insertion. However, unbeknownst to them, this library was created in a strain with a

mutation that rendered the mutants incapable of transitioning between low and high cell density.

In this study, we successfully transferred transposon insertions from these *V. cholerae*

non-redundant ordered mutants into a wildtype genetic background using chitin-independent

natural transformation. The resulting Grant Library comprises 3,102 mutants, covering

approximately 79.8% of the ORFs annotated in *V. cholerae*.

In addition to encoding a functional quorum sensing system, another notable advantage

of the Grant Library is that we selected for the transposon insertion and the plasmid that confers

IPTG-inducible natural competence during outgrowth in selective media. Accordingly, upon

induction with IPTG, mutants from the Grant Library can undergo additional genome edits when

co-incubated with gDNA. Furthermore, growth in the absence of selection leads to rapid curing

of the competence plasmid if it is not needed. These features make the Grant Library an

invaluable resource for studying pleiotropic and epistatic genetic interactions in *V. cholerae*.

Indeed, the non-linearity between paired mutants when grown on inosine demonstrates this

utility. Ongoing work in the Grant lab involves utilizing the two transposon libraries described in

this work, and a third high cell density locked library we are generating, to deconstruct gene *x*

transcriptional state *x* environmental interactions, including synthetically lethal genetic

combinations. Such information will be invaluable toward generating new therapeutic options to

treat *V. cholerae* infections.

Construction of the Grant Library was aided by a novel, in-house gDNA extraction

method we developed based on thymol cell lysis. Our use of thymol allowed extraction of gDNA

in a highly efficient and economical way. Indeed, N.A.G. has performed as many as 576 manual

gDNA extractions in a single workday. Additionally, at the concentrations used in our studies,

gDNA extracted with thymol does not negatively impact its use in other downstream

applications, including PCR amplification, whole genome and Sanger sequencing. Given
thymol's cell killing efficacy in *Vibrio*, we tested the treatment on several other biological
specimens including bacteria in gram -negative and -positive staining classes and yeast. Among
our test strains, gDNA yield was highest for *V. cholerae* (~ 22 ng/µl), although most strains
yielded some gDNA (2 – 8 ng/µl) (data not shown). That *Vibrio* is highly susceptible to thymol
treatment is an interesting avenue for further research as its understanding can provide additional
insight into its antibacterial activity, potentially contributing to novel disease control strategies.
In summary, the Grant Library complements the non-redundant library made in the quorum-
incompetent strain, and the methods used to construct it, a valuable approach toward
understanding *V. cholerae* evolutionary genetics.
637

Acknowledgements
We thank Ewen Cameron and the John Mekalanos lab for constructing the 2008 *V. cholerae* non-
redundant library, Vic DiRita for granting access to a copy of the collection, and Bonnie Bassler
and Julie Valastyan for providing plates that were missing from the DiRita collection and the
gain of function *luxOG319S* parent strain. We also thank Ankur Dalia for providing the plasmid
conferring natural competence. This work was supported by National Institutes of Health Grants
GM139537 and AI158433 to C.M.W and a NIGMS Diversity Supplement to N.A.G. Financial
support was also received from The BEACON Center for the Study of Evolution in Action at
Michigan State University (to N.A.G).

References
1. M. Ali, A. R. Nelson, A. L. Lopez, D. A. Sack, Updated Global Burden of Cholera in
Endemic Countries. *PLOS Neglected Tropical Diseases* **9**, e0003832 (2015).

2. I. Ilic, M. Ilic, Global Patterns of Trends in Cholera Mortality. *Trop Med Infect Dis* **8**, 169
(2023).

3. D. Legros, Global Cholera Epidemiology: Opportunities to Reduce the Burden of Cholera
by 2030. *J Infect Dis* **218**, S137–S140 (2018).

4. Roadmap 2030 – Global Task Force on Cholera Control. Available at:
https://www.gtfcc.org/about-cholera/roadmap-2030/ [Accessed 22 October 2023].

5. D. Lippi, E. Gotuzzo, S. Caini, Cholera. *Microbiology Spectrum* **4**,
10.1128/microbiolspec.poh-0012–2015 (2016).

6. S. Yamai, T. Okitsu, T. Shimada, Y. Katsube, Distribution of serogroups of *Vibrio cholerae*
non-O1 non-O139 with specific reference to their ability to produce cholera toxin, and
addition of novel serogroups. *Kansenshogaku Zasshi* **71**, 1037–1045 (1997).

7. A. Safa, G. B. Nair, R. Y. C. Kong, Evolution of new variants of *Vibrio cholerae* O1.
Trends in Microbiology **18**, 46–54 (2010).

8. J. B. Kaper, J. G. Morris, M. M. Levine, Cholera. *Clin Microbiol Rev* **8**, 48–86 (1995).

9. D. Hu, *et al.*, Origins of the current seventh cholera pandemic. *Proceedings of the National*
Academy of Sciences **113**, E7730–E7739 (2016).

10. K. Okada, T. Iida, K. Kita-Tsukamoto, T. Honda, *Vibrios* Commonly Possess Two
Chromosomes. *J Bacteriol* **187**, 752–757 (2005).

11. S. Sozhamannan, T. Waldminghaus, Exception to the exception rule: synthetic and
naturally occurring single chromosome *Vibrio cholerae*. *Environmental Microbiology* **22**,
4123–4132 (2020).

12. D. K. Karaolis, R. Lan, P. R. Reeves, The sixth and seventh cholera pandemics are due to
independent clones separately derived from environmental, nontoxigenic, non-O1 *Vibrio*
cholerae. *J Bacteriol* **177**, 3191–3198 (1995).

13. S. M. Faruque, *et al.*, Emergence and evolution of *Vibrio cholerae* O139. *Proceedings of*
the National Academy of Sciences **100**, 1304–1309 (2003).

14. T. Ramamurthy, *et al.*, Revisiting the Global Epidemiology of Cholera in Conjunction With
the Genomics of *Vibrio cholerae*. *Front Public Health* **7**, 203 (2019).

15. S. N. Sakib, G. Reddi, S. Almagro-Moreno, Environmental Role of Pathogenic Traits in
Vibrio cholerae. *J Bacteriol* **200**, e00795-17 (2018).

16. Y. Sun, E. E. Bernardy, B. K. Hammer, T. Miyashiro, Competence and Natural
Transformation in *Vibrios*. *Mol Microbiol* **89**, 10.1111/mmi.12307 (2013).

17. R. L. Marvig, M. Blokesch, Natural transformation of *Vibrio cholerae* as a tool -
Optimizing the procedure. *BMC Microbiology* **10**, 155 (2010).

18.   S. Stutzmann, M. Blokesch, Comparison of chitin-induced natural transformation in
686   pandemic *Vibrio cholerae* O1 El Tor strains. *Environ Microbiol* **22**, 4149–4166 (2020).

687   19.   K. L. Meibom, M. Blokesch, N. A. Dolganov, C.-Y. Wu, G. K. Schoolnik, Chitin Induces
688   Natural Competence in *Vibrio cholerae*. *Science* **310**, 1824–1827 (2005).

689   20.   S. S. Watve, J. Thomas, B. K. Hammer, CytR Is a Global Positive Regulator of
690   Competence, Type VI Secretion, and Chitinases in *Vibrio cholerae*. *PLoS One* **10**,
691   e0138834 (2015).

692   21.   M. Lo Scrudato, S. Borgeaud, M. Blokesch, Regulatory elements involved in the expression
693   of competence genes in naturally transformable *Vibrio cholerae*. *BMC Microbiology* **14**,
694   327 (2014).

695   22.   M. L. Scrudato, M. Blokesch, The Regulatory Network of Natural Competence and
696   Transformation of *Vibrio cholerae*. *PLOS Genetics* **8**, e1002778 (2012).

697   23.   E. S. Antonova, B. K. Hammer, Genetics of Natural Competence in *Vibrio cholerae* and
698   other Vibrios. *Microbiol Spectr* **3**, 3.3.20 (2015).

699   24.   A. B. Dalia, D. W. Lazinski, A. Camilli, Identification of a Membrane-Bound
700   Transcriptional Regulator That Links Chitin and Natural Competence in *Vibrio cholerae*.
701   *mBio* **5**, 10.1128/mbio.01028-13 (2014).

702   25.   T. N. Dalia, *et al.*, Multiplex genome editing by natural transformation (MuGENT) for
703   synthetic biology in *Vibrio natriegens*. *ACS Synth Biol* **6**, 1650–1655 (2017).

704   26.   A. B. Dalia, "Natural Cotransformation and Multiplex Genome Editing by Natural
705   Transformation (MuGENT) of *Vibrio cholerae*" in *Vibrio Cholerae: Methods and*
706   *Protocols*, Methods in Molecular Biology., A. E. Sikora, Ed. (Springer, 2018), pp. 53–64.

707   27.   A. B. Dalia, E. McDonough, A. Camilli, Multiplex genome editing by natural
708   transformation. *Proc Natl Acad Sci U S A* **111**, 8937–8942 (2014).

709   28.   T. Baba, *et al.*, Construction of *Escherichia coli* K-12 in-frame, single-gene knockout
710   mutants: the Keio collection. *Mol Syst Biol* **2**, 2006.0008 (2006).

29. M. M. Pearson, S. Pahil, V. S. Forsyth, A. E. Shea, H. L. T. Mobley, Construction of an
Ordered Transposon Library for Uropathogenic *Proteus mirabilis* HI4320. *Microbiol Spectr*
10, e03142-22.

30. B. Ramage, *et al.*, Comprehensive Arrayed Transposon Mutant Library of *Klebsiella*
pneumoniae Outbreak Strain KPNIH1. *Journal of Bacteriology* **199**, 10.1128/jb.00352-17
(2017).

31. P. D. Fey, *et al.*, A Genetic Resource for Rapid and Comprehensive Phenotype Screening of
Nonessential *Staphylococcus aureus* Genes. *mBio* **4**, 10.1128/mbio.00537-12 (2013).

32. D. E. Cameron, J. M. Urbach, J. J. Mekalanos, A defined transposon mutant library and its
use in identifying motility genes in *Vibrio cholerae*. *Proceedings of the National Academy*
of Sciences **105**, 8736–8741 (2008).

33. H. Rui, *et al.*, Reactogenicity of live-attenuated *Vibrio cholerae* vaccines is dependent on
flagellins. *Proceedings of the National Academy of Sciences* **107**, 4359–4364 (2010).

34. R. W. Bogard, B. W. Davies, J. J. Mekalanos, MetR-Regulated *Vibrio cholerae* Metabolism
Is Required for Virulence. *mBio* **3**, 10.1128/mbio.00236-12 (2012).

35. T. G. Dong, B. T. Ho, D. R. Yoder-Himes, J. J. Mekalanos, Identification of T6SS-
dependent effector and immunity proteins by Tn-seq in *Vibrio cholerae*. *Proceedings of the*
National Academy of Sciences **110**, 2623–2628 (2013).

36. A. J. Van Alst, V. J. DiRita, Aerobic Metabolism in *Vibrio cholerae* Is Required for
Population Expansion during Infection. *mBio* **11**, 10.1128/mbio.01989-20 (2020).

37. S. T. Miyata, D. Unterweger, S. P. Rudko, S. Pukatzki, Dual Expression Profile of Type VI
Secretion System Immunity Genes Protects Pandemic *Vibrio cholerae*. *PLOS Pathogens* **9**,
e1003752 (2013).

38. A. Hsiao, Z. Liu, A. Joelsson, J. Zhu, *Vibrio cholerae* virulence regulator-coordinated
evasion of host immunity. *Proceedings of the National Academy of Sciences* **103**, 14542–
14547 (2006).

39. F. P. Rothenbacher, J. Zhu, Efficient responses to host and bacterial signals during *Vibrio*
cholerae colonization. *Gut Microbes* **5**, 120–128 (2014).

40. S. Saha, S. Aggarwal, D. V. Singh, Attenuation of quorum sensing system and virulence in
Vibrio cholerae by phytomolecules. *Frontiers in Microbiology* **14** (2023).

41. J. Zhu, *et al.*, Quorum-sensing regulators control virulence gene expression in *Vibrio
cholerae*. *Proceedings of the National Academy of Sciences* **99**, 3129–3134 (2002).

42. J. Zhu, J. J. Mekalanos, Quorum Sensing-Dependent Biofilms Enhance Colonization in
Vibrio cholerae. *Developmental Cell* **5**, 647–656 (2003).

43. M. Jemielita, N. S. Wingreen, B. L. Bassler, Quorum sensing controls *Vibrio cholerae*
multicellular aggregate formation. *eLife* **7**, e42057 (2018).

44. M. M. Hoque, *et al.*, Quorum Regulated Resistance of *Vibrio cholerae* against
Environmental Bacteriophages. *Sci Rep* **6**, 37956 (2016).

45. H. Boyaci, *et al.*, Structure, Regulation, and Inhibition of the Quorum-Sensing Signal
Integrator LuxO. *PLoS Biol* **14**, e1002464 (2016).

46. R. Herzog, N. Peschek, K. S. Fröhlich, K. Schumacher, K. Papenfort, Three autoinducer
molecules act in concert to control virulence gene expression in *Vibrio cholerae*. *Nucleic
Acids Res* **47**, 3171–3183 (2019).

47. S. A. Jung, C. A. Chapman, W.-L. Ng, Quadruple Quorum-Sensing Inputs Control *Vibrio
cholerae* Virulence and Maintain System Robustness. *PLOS Pathogens* **11**, e1004837
(2015).

48. S. Stutzmann, M. Blokesch, Circulation of a Quorum-Sensing-Impaired Variant of *Vibrio
cholerae* Strain C6706 Masks Important Phenotypes. *mSphere* **1**, 10.1128/msphere.00098-
16 (2016).

49. A. Escobar, M. Pérez, G. Romanelli, G. Blustein, Thymol bioactivity: A review focusing on
practical applications. *Arabian Journal of Chemistry* **13**, 9243–9269 (2020).

50. J. B. Brown, Classifiers and their Metrics Quantified. *Mol Inform* **37**, 1700127 (2018).

51. D. E. Deatherage, J. E. Barrick, Identification of mutations in laboratory evolved microbes
from next-generation sequencing data using breseq. *Methods Mol Biol* **1151**, 165–188
(2014).

52.   A. C. E. Darling, B. Mau, F. R. Blattner, N. T. Perna, Mauve: Multiple Alignment of
767   Conserved Genomic Sequence With Rearrangements. *Genome Res* **14**, 1394–1403 (2004).

768   53.   M. Kearse, *et al.*, Geneious Basic: An integrated and extendable desktop software platform
769   for the organization and analysis of sequence data. *Bioinformatics* **28**, 1647–1649 (2012).

770   54.   J. Schindelin, *et al.*, Fiji: an open-source platform for biological-image analysis. *Nat
771   Methods* **9**, 676–682 (2012).

772   55.   J. F. Heidelberg, *et al.*, DNA sequence of both chromosomes of the cholera pathogen *Vibrio
773   cholerae*. *Nature* **406**, 477–483 (2000).

774   56.   M. Grognot, A. Mittal, M. Mah'moud, K. M. Taute, *Vibrio cholerae* Motility in Aquatic
775   and Mucus-Mimicking Environments. *Applied and Environmental Microbiology* **87**,
776   e01293-21 (2021).

777   57.   F. Khan, N. Tabassum, R. Anand, Y.-M. Kim, Motility of *Vibrio* spp.: regulation and
778   controlling strategies. *Appl Microbiol Biotechnol* **104**, 8187–8208 (2020).

779   58.   M. A. Echazarreta, K. E. Klose, *Vibrio* Flagellar Synthesis. *Frontiers in Cellular and
780   Infection Microbiology* **9** (2019).

781   59.   K. A. Syed, *et al.*, The *Vibrio cholerae* Flagellar Regulatory Hierarchy Controls Expression
782   of Virulence Factors. *J Bacteriol* **191**, 6555–6570 (2009).

783   60.   S. Zhu, S. Kojima, M. Homma, Structure, gene regulation and environmental response of
784   flagella in *Vibrio*. *Frontiers in Microbiology* **4** (2013).

785   61.   S. Beyhan, A. D. Tischler, A. Camilli, F. H. Yildiz, Transcriptome and Phenotypic
786   Responses of *Vibrio cholerae* to Increased Cyclic di-GMP Level. *J Bacteriol* **188**, 3600–
787   3613 (2006).

788   62.   D. S. Millikan, E. G. Ruby, FlrA, a σ54-Dependent Transcriptional Activator in *Vibrio
789   fischeri*, Is Required for Motility and Symbiotic Light-Organ Colonization. *J Bacteriol* **185**,
790   3547–3557 (2003).

791   63.   S. Srinivasan, A. G. Torres, L. Ribas de Pouplana, Inosine in Biology and Disease. *Genes
792   (Basel)* **12**, 600 (2021).

64.   Y. Liu, *et al.*, *Limosilactobacillus reuteri* and *Lacticaseibacillus rhamnosus* GG
differentially affect gut microbes and metabolites in mice with Treg deficiency. *Am J*
Physiol Gastrointest Liver Physiol **320**, G969–G981 (2021).

65.   L. F. Mager, *et al.*, Microbiome-derived inosine modulates response to checkpoint inhibitor
immunotherapy. *Science* **369**, 1481–1489 (2020).

66.   G. Haskó, M. V. Sitkovsky, C. Szabó, Immunomodulatory and neuroprotective effects of
inosine. *Trends Pharmacol Sci* **25**, 152–157 (2004).

67.   D. Li, *et al.*, Gut microbiota-derived inosine from dietary barley leaf supplementation
attenuates colitis through PPARγ signaling activation. *Microbiome* **9**, 83 (2021).

68.   M. Pigliucci, C. J. Murren, C. D. Schlichting, Phenotypic plasticity and evolution by
genetic assimilation. *Journal of Experimental Biology* **209**, 2362–2367 (2006).

Figure Legends

Figure 1. Thymol-Assisted gDNA extraction and chitin-independent natural transformation.
Overnight cultures of the donor library were grown in 550 µl of selective media (LB +
kanamycin). After overnight growth, the samples were centrifuged, the supernatant was
aspirated, and the cell pellets were incubated for one hour at 64°C in a novel gDNA extraction
buffer containing thymol. gDNA was then isolated from the clarified supernatant using an
alcohol precipitation method, followed by resuspension of the gDNA in 1x TE buffer. In parallel,
a quorum-competent wildtype *V. cholerae* strain carrying the plasmid conferring natural
competence (pMMB-tfoX-qstR) was grown in selective media (LB + chloramphenicol)
supplemented with IPTG. The next day, competent cells were diluted 1:4 in 0.5x Instant Ocean,
to which chloramphenicol and IPTG were added. Subsequently, 200 µl of cells were dispensed
into the wells of a deep-well 96-well plate and co-incubated with 100 µl of our gDNA extraction
at 30°C for 18 – 24 h. For a detailed description, refer to the Materials and Methods section.

Figure 2. Analysis of Library Transfer. The outcomes of growth on selective media for each
mutant were recorded after chitin-independent natural transformation of the donor library. These
values were utilized as inputs to construct a 2x2 confusion matrix, categorizing the results as
follows: True Positives (3,102 wells), False Negatives (22 wells), True Negatives (71 wells), and

False Positives (15 wells). The accompanying table presents the calculated measures, the values
of which were computed using standard derivations.

Figure 3. Representative sample of breseq output showing new junction evidence supporting
transposon insertion. The figure shows a portion of the summary.html file generated for a breseq
analysis run. In this example, a new junction has been called between chromosome II
(NZ_CP046845) of *V. cholerae* strain C6706 and the plasmid (pSC189) containing
representative sequences of the transposon. Insertions have left and right boundaries, where the
chromosome and sequence meet. The boundaries are marked with an asterisk. The annotation
column shows the position of the transposon within the gene (GPY04_RS15290), which we
converted to a value called "percent into open reading frame." In this example, that value is
41.43% ((225/543) * 100)). This value was calculated for all strains sequenced in this study.

Figure 4. Genomic comparative analysis of *V. cholerae* N16961 and C6706 for chromosome I
(A) and chromosome II (B). Genbank files for whole-genome sequences of chromosomes I and
II of both strains were downloaded from NCBI. The analysis involved a comparative
examination of genetic variations and conservation patterns between the two strains. *V. cholerae*
strain N16961 is shown with a gray background and C6706 a white one. Genome segments are
shown as colored colinear blocks centered around the "positive" and "negative" strands of each
chromosome. Lines connecting the segments show their orthologous alignments between the
strains, including inversions. Panel C is a Venn diagram summarizing the relationship of genes
for each strain and chromosome. The data underlying the illustration in panel C, including the
genomic position of orthologous gene pairs, is provided in the Supplementary Information
(Datasets S2–S4).

Figure 5. Correlation between transposon insertion sites in *V. cholerae* N16961 and C6706.
Each point represents the transposon insertion site, calculated as percent into open reading frame.
The values for strain N16961 were reported in supplementary dataset 3 of Cameron et al. (2008),
and in this work, they were calculated using the data provided in the annotation column of the
breseq output (Fig. 3). The correlation analysis yields a strong positive correlation with
Kendall's coefficient $\tau = 0.752$ ($n = 23$, and $P \ll 0.0001$), indicating a significant relationship
between the insertion sites of the two strains.

Figure 6. Chromosomal integration of a trimethoprim resistant gene in select mutants from the
Grant library using IPTG-induced natural competence. We induced natural competence in

wildtype *V. cholerae* strain C6706 carrying plasmid pMMB-*tfoX-qstR* (NG001) and 10 randomly

selected mutants from the Grant library according to procedure described in Materials and

Methods. These strains were then co-incubated with gDNA extracted from a strain carrying a

trimethoprim resistant cassette and on the following day spotted on antibiotic selection media.

All the strains carrying the plasmid grew on plates supplemented with trimethoprim (Panel A).

On the plate with kanamycin (Panel B) or kanamycin + trimethoprim (Panel C), all strains grew

except strain 8C5, indicating that this strain does not have a transposon insertion.

Figure 7. Effects of media composition on plasmid loss in *V. cholerae*. We serially propagated

cultures of the Grant Library parent strain in LB media with or without antibiotics or IPTG

induction as indicated in the figure. The percent of population with the plasmid conferring

natural competence was calculated as a ratio of the dilutions where colonies grew on selective

media (LB + chloramphenicol (Cam) relative to non-selective media (LB). Error bars are the 95

% confidence intervals on a sample size of $n = 3$ for each treatment.

Figure 8. Motility behavior of flagellar mutants from the Grant and Cameron et al. (2008)

libraries. Paired mutants with transposons in flagellar genes from both libraries were screened for

motility in nutrient-rich plates containing 0.35% agar. The plates were imaged, and the area

moved by each mutant from an inoculum site was calculated using the procedure described in the

Materials and Methods. These areas were then normalized to a value relative to the wild-type

strains for each library from 3 replicate assays in all but four cases (*flgM, flhB, fliR*, and *flgF*),

which had two replicates each. The dashed red line is the normalization for the wildtype strains.

Colored boxes highlight the class each gene belongs to in the flagellar biosynthesis gene cluster

hierarchy, and red stars highlight the expected non-motile phenotype for transposon insertions

that lie within genes encoding sigma factors required to transcribe downstream flagellar

biosynthesis genes.

Figure 9. Genotypic and phenotypic characterization of Grant and Cameron library *V. cholerae*

strains on inosine. (A) *luxO* sequence chromatogram for *V. cholerae* C6706 parent strains of

Grant (recipient) and Cameron (donor) transposon libraries. Red arrow shows location of gain-

of-function mutation resulting in a Glycine-to-Serine transition at amino acid position 319 in the

Cameron library. (B) Growth of parent strains on inosine. Quorum-competent Grant library

strain grows to a higher OD (Two-Sample unpaired $t = 29.34$, $df = 16$, and $P << 0.0001$). (C)

Boxplots of end-point optical density values for paired mutants from Grant and Cameron

libraries grown on inosine. Each point represents a single mutant and the connecting lines the

reaction norm for each mutant with respect to the library sampled from. Like the parent strains

(B), Grant Library mutants grows to a higher OD than the Cameron et al. Library strains (Two-

Sample unpaired t = -20.34, df = 904, and P << 0.0001).

Figure 10. Reaction norm for paired transposon mutants grown on inosine. Each point on the

graph represents the 10-hour endpoint optical density for mutants from the Cameron and Grant

libraries grown on inosine, with lines connecting paired mutants. We identified outliers by

focusing on the top and bottom 2.5% of the data, calculated using the mean and standard

deviation for each library. Relationships are plotted for points that deviated two standard

deviations from each mean.

Donor library gDNA extraction

| Overnight culture of donor library (550 µl) | Thymol cell lysis and DNA neutralization (1h) | Lysate clearance and alcohol precipitation | DNA binding and ethanol wash | DNA elution (centrifuge, 1 min) |

Chitin-independent natural competence

| Chitin-independent competent cells are prepared by diluting an overnight culture of donor recipient library strain grown under IPTG induction 1:4 in Instant Ocean buffer containing chloramphenicol (5 µg/mL) and IPTG | Dispense 200 µl of competent cells into deep-well plates and add gDNA from donor library (100 µl) | Incubate in a standing incubator overnight (18-24 h) at 37°C |

	Has transposon	Does not have transposon
Identified as having the transposon	3102	15
Identified as not having the transposon	22	71

Measure	Value
Sensitivity	0.9930
Specificity	0.8256
Precision	0.9952
Negative Predictive Value	0.7634
False Positive Rate	0.1744
False Discovery Rate	0.0048
False Negative Rate	0.0070
Accuracy	0.9885
F1 Score	0.9941
Matthews Correlation Coefficient	0.7880

	seq id	position	reads (cov)	reads (cov)	score	skew	freq	annotation	gene	product
								Unassigned new junction evidence		
?	NZ_CP046845	151866 =	2 (0.050)	37 (0.780)	15/290	0.0	97.4%	coding (225/543 nt)	GPY04_RS15290	N-acetyltransferase
* ?	pSC189	= 4058	0 (0.000)					intergenic (+121/+620)	aph(3')-II (or nptII)/traJ	aminoglycoside phosphotransferase from Tn5/oriT-recognizing protein
?	NZ_CP046845	= 151868	2 (0.050)	25 (0.520)	14/292	0.0	96.2%	coding (227/543 nt)	GPY04_RS15290	N-acetyltransferase
* ?	pSC189	1802 =	0 (0.000)					intergenic (–/-41)	–/3xFLAG	–/three tandem FLAG(R) epitope tags, followed by an enterokinase cleavage site

WT -gDNA
WT +gDNA
Strain 5B2
Strain 6F4
Strain 8C5
Strain 15F6
Strain 17G10
Strain 19F5
Strain 19F12
Strain 27B10
Strain 28C4
Strain 28D5