Deployment of a *Vibrio cholerae* ordered transposon mutant library in a quorum-
competent genetic background

Nkrumah A. Grant[1,2,3*], Jordan T. Sontz,[4] Christopher M. Waters[2,3,4*]

[1]Department of Microbiology, University of Illinois Urbana-Champaign, Urbana, IL
[2]Department of Microbiology and Molecular Genetics, Michigan State University, East Lansing,
MI
[3]BEACON Center for the Study of Evolution in Action, Michigan State University, East
Lansing, MI
[4]MSU College of Osteopathic Medicine, Michigan State University, East Lansing, MI

[*]Correspondence to: grantnkr@illinois.edu or watersc3@msu.edu

Abstract

Cholera is a severe diarrheal disease that claims approximately 30,500 lives annually, mainly in
regions with limited access to clean water. *Vibrio cholerae*, the causative agent, has sparked
seven pandemics in recent centuries, with the current one being the most prolonged. *V. cholerae*
pathogenesis hinges on its ability to switch between low and high cell density gene regulatory
states, enabling transmission between host and the environment.
Previous researchers created a transposon mutant library for *V. cholerae* to support investigations
aimed toward uncovering the genetic determinants of its pathogenesis. However, subsequent
sequencing uncovered a mutation in the gene *luxO* of the parent strain, rendering mutants unable
to exhibit high cell density behaviors. In this study, we used chitin-independent natural
transformation to move transposon insertions from these low cell density mutants into a wildtype
genomic background. Library transfer was aided by a novel gDNA extraction we developed
using thymol, which also showed marked vibrio-specific activity. The resulting Grant Library
comprises 3,102 unique transposon mutants, covering 79.8% of *V. cholerae's* open reading
frames. Whole genome sequencing of randomly selected mutants demonstrates 100% precision
in transposon transfer to cognate genomic positions of the recipient strain. Notably, *luxO*
mutations transferred at a low frequency only with insertions near *luxO*. Our research uncovered
density-dependent epistasis in motility genes, relevant to *V. cholerae* pathogenesis. Additionally,
Grant Library mutants retain a plasmid conferring natural competence, enabling rapid, scarless
genomic editing. In summary, the Grant Library reintroduces organismal relevant genetic
contexts absent in the low cell density locked library equivalent.

Introduction

Vibrio cholerae is a human pathogen and the causative agent of the acute diarrheal disease cholera. While cholera is a global threat, the disease is endemic in developing countries, where it is estimated that there are between 1.3 to 4.0 million *V. cholerae* annual infections, resulting in 21,000 – 143,000 deaths (1, 2). Implementation of disease intervention strategies, including increasing access to clean drinking water, provision of adequate healthcare infrastructure, and educating citizens in communities at risk of disease spread, remains a top priority in the fight toward eradicating cholera. As such, the Global Task Force on Cholera Control was founded in 2017 to support efforts aimed at reducing cholera burden by 90% before the year 2030 (3, 4).

To date there have been seven cholera pandemics. The seventh pandemic began in 1961 and is ongoing, marking it as the longest running cholera pandemic (5). *V. cholerae* strains are grouped by surface antigens, including O polysaccharide, such that there are >200 serogroups (6). Of these serogroups, only two, O1 and O139, host pandemic *V. cholerae* strains (7, 8). Furthermore, pandemic strains from the O1 serogroup are further stratified into two biotypes with the fifth and sixth pandemics caused by the Classical and the seventh caused by El Tor (9).

Like most vibrios, *V. cholerae* predominantly possesses two chromosomes (10, 11). Whole genome sequencing of strains before and after the emergence of seventh pandemic *V. cholerae* strains has proved to be an invaluable resource toward understanding its origin and evolution. These analyses show *V. cholerae* El Tor originated from a nonpathogenic strain (12), becoming a pathogen after acquiring two genomic islands (VSP-I and VSP-II), a prophage that expresses cholera toxin (CTXφ), and 12 additional mutations that enhanced its transmissibility (9, 13, 14). Recent work has also shown that these genomic changes contribute to seventh pandemic *V. cholerae's* long-term persistence in the estuarian environments they occupy (15).

Key to El Tor's transition to a pathogen is that it is naturally competent, granting it the ability to take up DNA from the environment which can be integrated into the genome by homologous recombination (16). Natural competence is a highly regulated process, and four environmental factors are key determinants for driving this process. These include nutrient limitation, extracellular signaling molecules, high cell density, and the presence of chitin (17–19), which together coordinate the expression of the four regulators (HapR, TfoX, CRP, and CytR) governing competence (20–23). Recent work has shown that expressing the master regulator TfoX can circumvent the chitin requirement for natural competence in *V. cholerae* (19, 24). This discovery has allowed exploitation of chitin-independent natural competence as a

genetic tool for editing *Vibrio spp.* genomes (25–27), facilitating our understanding of the

functional relationships between genes and furthering our understanding of how *V. cholerae*

interfaces with its environment.

Our understanding of *V cholerae's* ecology and evolution has also benefited from

transposon mutagenesis. In contrast to conventionally laborious genetic engineering approaches

for disrupting single genes to study their phenotypic effects, transposon mutagenesis allows the

simultaneous construction and screening of tens of thousands of mutants. Researchers have taken

advantage of transposon mutagenesis to construct nonredundant (ordered) mutant libraries,

where each nonessential gene has been inactivated and the individual mutants are arrayed across

the wells of 96-well microtiter plates (28–31). An ordered mutant *V. cholerae* library was

constructed in 2008 (32). The ordered library consists of >3,100 mutants, each with a Tn5-based

transposon insertion that confers kanamycin resistance. The mutant library has provided a public

resource enhancing *V. cholerae* genotypic and phenotypic investigations, supporting identifying

novel therapeutic targets for vaccine development (33), and increasing our understanding of the

V. cholerae virulome (34, 35), amongst other discoveries (36, 37).

Although significant research advances have been made in our understanding of *V.*

cholerae using mutants from this ordered mutant transposon library, the library was constructed

in a laboratory acquired quorum sensing mutant of *V. cholerae*. Quorum sensing, the process of

cell-to-cell communication in bacteria, allows cells to undergo global changes in gene expression

as the bacteria transition from low to high cell density. This transition can influence behaviors

like host immunity escape (38), virulence factor production (39–41), biofilm formation (42),

infectivity and environmental dissemination (43), and protection against phage predation (44).

This transition is mediated by extracellular autoinducers (45, 46) which impact the activity of

cytoplasmic regulators, including LuxO and HapR .

In the low-cell-density state, LuxO is phosphorylated leading to inhibition of *hapR*

mRNA translation via regulatory sRNAs, whereas at high-cell-density, LuxO is

dephosphorylated and inactive, resulting in HapR protein translation and the switch to high-cell-

density (41, 47). Recently, a presumably laboratory acquired mutation in *luxO, luxOG319S* was

discovered in some laboratory isolates of the widely used *V. cholerae* strain C6706 that locked

this strain in the low-cell-density state (48). After sequence validating mutants from the 2008

library, we observed the *luxO*G319S mutation, indicating it was present in the parent strain from

which the library was constructed. Thus, the 2008 ordered mutant library (32) is only partially

representative of how *V. cholerae* senses and responds to its environment and a new library in a
quorum sensing competent strain would greatly enhance our understanding of *V. cholerae*
ecology and evolution.
109
To overcome this limitation of the 2008 library, we use chitin-independent natural
competence to regenerate the defective ordered *V. cholerae* library in a quorum competent
C6706 genomic background. To streamline the transfer process, we developed a cost effective
gDNA extraction method using thymol (49). Compared to other methods, our approach is both
economical ($0.64 per reaction vs $2.85 per reaction) and efficient, enabling up to 200 gDNA
extractions in as little as 2.5 hours, in contrast to 25 reactions requiring >6 hours using
conventional methods. Using our methods, we successfully regenerated the library with a 99%
success rate, as indicated by growth on antibiotic selection growth media. Sequence validation of
randomly selected mutants from the regenerated library show that the transposon insertions
recombine into our recipient strain with 100% precision. Transfer of the *luxOG319S* mutation
was minimal and only occurred in genes located close to the transposon insertion site. Of
significance, we provide evidence for density-dependent epistasis in several genes within class
III of the flagellum biosynthesis gene clusters, an observation that undoubtedly extends to other
phenotypes, illustrating the utility of this new library. Lastly, we show that strains from the
library generated in this work – hereafter the Grant Library – can undergo additional edits using
chitin-independent natural competence. Taken together, the Grant Library represents an added
contribution in our efforts toward understanding *V. cholerae* behaviors under transcriptional
regulatory states that are representative of how this pathogen evolves and interfaces with its
environment.
129
130 **Materials and Methods**
131
Bacterial matings
To generate the library recipient strain used in this study, NG001, we mated wildtype *V. cholerae*
C6706 with an *E. coli* S17-λpir donor strain harboring plasmid pMMB-*tfoX-qstR* (generously
provided by Ankur Dalia, Indiana University). In short, we mixed equal volumes of donor and
recipient strains in the center of a Luria-Bertani agar plate which we incubated at 37°C for three
hours. Afterwards, we added 1 mL of LB media containing chloramphenicol (10 μg/mL, for
selection of plasmid in recipient strain) and polymyxin B (55.5 μg/mL, for counterselection of
the donor *E. coli* strain) to the plate, washed, and collected the mating spot into a borosilicate
glass culture tube, which we incubated overnight at 37°C with orbital shaking at 210 RPM.

Media and buffers

Thymol cell lysis agent
We prepared thymol at a stock concentration of 350 mM by dissolving thymol crystals in dimethyl sulfoxide (DMSO). Thymol stocks were made in 15 mL which we stored at room temperature on the lab bench until they were exhausted. We observed no differences in the lytic activity of thymol during storage.

gDNA extraction buffer
We prepared solutions of a gDNA extraction buffer in two 500 mL volume parts. We prepared solution one by mixing 80 mL of 1M Tris (pH 8.0), 56 mL of 0.5M EDTA (pH 8.0), 46.7 g NaCl, and 3 g sodium metabisulfite, which we stirred until dissolved. We then adjusted this solution to the final volume with dH_2O and autoclaved for 25 minutes. We prepared solution two, sodium acetate (2.5M, pH 5.2), by dissolving 102.5 g of the salt in dH_2O, which we sterilized using filtration. Prior to each extraction, we prepared a master mixture of solution one and solution two by mixing 21 mL and 31.5 mL for every 100 samples processed, respectively.

Strain revival
We revived strains from the nonredundant library ((32); henceforth referred to as donor library) in microplate format to maintain parity between it and that produced in this work. The donor library was generously provided by the laboratories of Vic DiRita (plates: 2, 4-6, 8-10, 12-17, 19-33), Michigan State University, and Bonnie Bassler (plates: 1, 3, 7, 11, 18, 34), Princeton University. Briefly, microtiter plates from the donor library were slightly thawed and 10 µl from each well was inoculated into a deep-well microplate containing 550 µl of selective media (Luria-Bertani broth + Kanamycin (100 µg/ml) in each of the 96 wells. We revived the recipient library strain, NG001, in a 250 ml baffled flask containing 25 ml of selective media (Luria-Bertani broth + chloramphenicol (10 µg/ml)) to maintain plasmid *pmmB-tfoX-qstR* and 100 µg/ml isopropyl β-D-1-thiogalactopyranoside (IPTG) to induce natural competence. Unless noted otherwise, we grew all cultures overnight (18 – 24h) at 37°C with orbital shaking at 210 RPM.

Whole gDNA preparation
The non-redundant donor library is comprised of 34 96-well plates of *Vibrio cholerae* mutants. To extract genomic DNA from the donor library in a cost efficient and high-throughput fashion, we developed an extraction method using a modified recipe of the gDNA extraction buffer

developed by Mantel and Sweigart, 2019 (accessed September, 2022, at

https://www.protocols.io/view/quick-amp-dirty-dna-extraction-4r3l287zql1y/v1). In brief, we

pelleted cells from overnight cultures of the donor library by centrifuging the deep-well plates at

4°C for 20 minutes. We then removed ~500 µl of supernatant from each well, resuspended the

cell pellet in 10 µl thymol (~35 mM final concentration), and incubated the plates at 64°C for 15

minutes. Thereafter, we added 500 µl of gDNA buffer to each well and incubated the plates at

64°C for an additional 45 minutes.

To clear culture supernatants of cellular debris, we transferred cell lysates from each well

into an E-Z 96™ Lysate Clearance Plate (https://www.omegabiotek.com/product/e-z-96-lysate-

clearance-plate/). We then used a vacuum manifold to filter the supernatant through the plate,

which we collected in a second deep-well plate containing 200 µl of 100% cold isopropanol per

well. We then incubated the plates for 15 minutes at room temperature to precipitate genomic

DNA. Afterwards, we again used a vacuum manifold to filter the cell lysate containing our

gDNA through an E-Z 96™ DNA plate (https://www.omegabiotek.com/product/e-z-96-dna-

plates/?cn-reloaded=1). According to the manufacturer, "E-Z 96™ plates are silica glass fiber

plates that can bind up to 50 µg of genomic DNA or 20 µg of plasmid DNA per prep." After this

initial filtration, we washed our bound DNA, in sequence, using 500 µl of 70% and 95% ice cold

ethanol. We then added 100 µl of 1x TE buffer to each well and eluted genomic DNA from the

DNA plates into PCR plates by centrifugation (max speed, one minute).

Chitin-independent natural competence for gDNA transfer

After 18-24 h of overnight growth, we diluted the library recipient strain (NG001) 1:4 in 0.5x

instant ocean to which we added chloramphenicol (5 µg/ml) and IPTG (100 µg/ml). We then

added 200 µl of these competent cells to each well of a deep-well 96-well plate and added the

gDNA extracted from the donor library to a matched recipient well (final dilution 1:3). We then

incubated the deep-well plates overnight in a standing incubator at 30°C. Following the

overnight incubation, we added 400 µl of selective media (Luria-Bertani broth +

chloramphenicol (5 µg/mL) + kanamycin (100 µg/mL)) to each well and incubated the plates at

37°C in a standing incubator for 48 h (Fig. 1). We determined transformation success by spot

plating 3 µl of each well onto selective LB agar plates (Luria-Bertani agar + chloramphenicol (5

µg/mL) + kanamycin (100 µg/mL)), which we grew overnight at 37°C (Fig. S1).

Grant Library transformation efficiency analysis

We created a comprehensive plate indices file (Dataset S1) covering all possible wells within the

donor library (P1A1 to P34D6; 3,210 wells). Using this file, we developed a script to compare it

with a modified version of supplementary dataset 3 from Cameron et al., 2008 (Dataset S2),

which allowed us to identify empty wells (86) and those containing mutants (3,124). Our

comparative analysis also flagged duplicate entries in supplementary dataset 3 of the donor

library (Dataset S3).

While constructing the Grant Library, we meticulously documented the outcomes of

growth on selective media following chitin-independent natural transformation of the donor

library: True Positives (3,102 wells), False Negatives (22 wells expected to grow but didn't),

True Negatives (71 blank wells that didn't transfer), and False Positives (15 blank wells that

transferred unexpectedly). These values served as inputs for a binary confusion matrix (50),

providing a comprehensive assessment.

TnFGL3 transposon insertion site identification

Genome Sequencing

We randomly selected 23 mutants from the Grant library to identify the transposon insertion site.

Strains were incubated overnight in 3 ml of LB + Kanamycin (100 µg/ml) and DNA was

extracted from 1 ml of culture using a Wizard ® Genomic DNA Purification Kit following the

manufacturer's instructions. DNA was rehydrated overnight at 4°C in 50 µl of TE buffer. Library

preparation was performed by SeqCoast Genomics using an Illumina DNA Prep tagmentation kit

and unique dual indexes. Samples were multiplexed and sequenced on the Illumina NextSeq2000

platform using a 300-cycle flow cell kit to produce 2x150 bp paired reads. Read demultiplexing,

read trimming, and run analytics were performed using DRAGEN v3.10.11, an on-board analysis

software on the NextSeq2000. Raw sequencing data generated during this study are available in

the following public Github repository (https://github.com/NkrumahG/Grant-Library-

Construction)

Reference genomes

The reference genomes for *V. cholerae* strains used in this study were downloaded from NCBI

GenBank. The chromosome accession numbers for chromosomes I and II are, NZ_CP028827.1

(2,975,504 bp) and NZ_CP028828.1 (1,072,331 bp) for strain N16961, and NZ_CP046844

(3,019,938 bp) and NZ_CP046845 (1,070,359 bp) for strain C6706, respectively. The sequence

map for plasmid pSC189 was downloaded from addgene

(https://www.addgene.org/browse/sequence/36279/), and the sequence for pMMB-*tfoX-qstR* was

provided by Ankur Dalia (Indiana University).

Mutation calling

We used consensus mode in *breseq* ((51), version 0.37) to align the paired-end reads for each of

the sequenced mutants to the reference genomes. The computational resources used to perform

breseq were provided by the Institute of Cyber-Enabled Research at Michigan State University.

Ortholog identification

We used Mauve ((52), snapshot _2015-02-25) to identify orthologs in whole genome sequences

of C6706 and N16961. Each pair of chromosomes were aligned using the progressive Mauve

method with default settings. In the Supplemental Information, we have included lists of all

orthologous gene pairs in *V. cholerae* N16961 and C6706 (Dataset S4) and the identity of genes

unique to N16961 (Dataset S5) and C6706 (Dataset S6). Additionally, we used Geneious to

extract, align, and visually display (Supplementary Information, Fig. S2 and S3) orthologous

gene pairs for the transposon mutants sequenced in this study.

LuxO gain of function mutation identification

Using supplementary dataset 3 provided for the donor library (32), we identified all potential

transposon mutants located within a 50 kb range, both upstream and downstream of *luxO*.

Subsequently, we developed a script that randomly picked 10 mutants from each side of *luxO*,

using 5 kb windows. Following this selection, we cultured these mutants in selective media (LB

+ Kanamycin, 100 µg/mL) for 24 hours under standard conditions.

After the incubation, we extracted gDNA using the Wizard Genomic DNA Purification

Kit, following the manufacturer's instructions. The extracted genomic DNA was then diluted 1:2,

which we used as template for PCR amplification of the *luxO* gene. The *luxO* gene was

amplified with Q5 polymerase using the following primers: forward primer 5`

GGCTATGCAACATAATCAATCTTG-3` and reverse primer 5`-

GCTTTGGTTGATCCATTCTCTCAT-3`. PCR conditions included an initial denaturation at

98℃ for 2 minutes, followed by 29 cycles of denaturing at 98℃ for 30 seconds, annealing at

63℃ for 30 seconds, and extension at 72℃ for 1 minute. A final extension step was

performed at 72℃ for 5 minutes.

Following the PCR amplification, we validated amplification of *luxO* using gel
electrophoresis. We then used the Wizard SV Gel and PCR Clean-Up System to purify our *luxO*
PCR product. Sanger sequencing of the purified *luxO* allele was performed using the reverse
primer 5`-TGCGATAGATGGTTGACGGG-3` by the Research Technology Support Facility
(RTSF) at Michigan State University. We imported the Sanger Sequence read data into Geneious
(53) and aligned it to *luxO* (*V. cholerae* chromosome I, C6706), allowing comprehensive
analysis and visualization of our sequenced mutants, with special attention to amino acid position
319, the site of the *luxO* gain of function mutation *luxOG319S* (48).
287
Motility Assay
We screened mutants with presumptive TnFLG3 transposon insertions in known motility genes.
Strains were grown in a deep-welled 96-well plate overnight at 37°C in selective media (LB +
Kanamycin) and on the following day, we used toothpicks to manually stab each strain into
motility plates containing LB and 0.35% agar. Plates were incubated at 37°C for 24 h, and the
motility zones were recorded with a gel imager. The area of the motility zones was measured
computationally using the Hough Circle Transform package from the UCB vision plugin in Fiji
(54).
296
297 **Data accessibility**
298 All of the data generated including Datasets S1-S6, and analysis scripts used in this work are
299 available in the GitHub repository at https://github.com/NkrumahG/Grant-Library-Construction.
300
301
302 **Results and Discussion**
303
Chitin-independent transfer of transposons from donor library strains into a quorum-
competent *V. cholerae* genetic background is remarkably efficient. We hypothesized that the
transposon insertions within the genomes of the quorum-incompetent *V. cholerae* strain
generated in Cameron et. al., 2008 could be transferred to a quorum-competent wildtype *V.*
cholerae strain using natural competence. To test this, we extracted gDNA from each mutant in
donor library using an in-house method involving thymol lysis and alcohol precipitation.
Subsequently, we co-incubated this gDNA with a wildtype *V. cholerae* strain (C6706) carrying a
plasmid that expresses the natural competence master regulator, TfoX, upon induction with IPTG
(Materials and Methods).
313

314   The donor library consists of 34 plates arrayed from P1A1 to P34D6, for a total 3,210
315   possible wells containing transposon mutants. Upon examination of supplementary data table 3
316   for the donor library, we discovered that 86 wells were not listed, and some of the mutants were
317   recorded multiple times. With these considerations, we expected growth for 3,124 unique
318   transposon mutants after conducting our natural transformation protocol.
319
To verify the presence of transposon insertions in the Grant library mutants, we spot-
plated each mutant on LB medium supplemented with kanamycin. Of the 3,124 unique
transposon mutants we expected to grow on the plates, 3,102 mutants (99.2%) exhibited growth
(Fig. S1). None of the 22 missing mutants, which were expected to grow, showed growth in
selective media during overnight revival of donor library freezer stock. This suggests that these
mutants might no longer viable perhaps due to low culture density upon freezing or death
resulting from repeated freeze-thaw cycling during long-term storage of the donor library. In the
cases where we anticipated no growth, 71/86 (82.0%) did not grow on selective media.
328
To evaluate the performance of our library transfer approach, we utilized a confusion
matrix (Fig. 2). The results of this analysis indicate that our approach was highly accurate and
precise, with a high recall (F1 score = 0.9941) and a low False Discovery Rate (0.0070). These
results unequivocally demonstrate the remarkable efficiency of using chitin-independent natural
competence to transfer the transposons into a quorum-competent strain. The transference process
was made possible through our in-house gDNA extraction method, involving thymol cell lysis
and alcohol precipitation, affirming the suitability of this gDNA preparation for downstream
applications. Indeed, this suitability was confirmed when we employed thymol-extracted DNA
as a template in PCR, with no observed inhibitions.
338
Transposons from the donor library homologously recombine into the recipient genome
with 100% precision. The frequency of genome edits by homologous recombination in *Vibrio*
spp. is highest when there are at least 2 kb arms of homology flanking the genomic sequence
being altered (25). Given this requirement, the observation that the recipient strains grew after
incubation on selective media (Fig. S1) was suggestive that transposons correctly integrated into
the homologous site of the recipient's genome. To validate our supposition, we randomly
selected 23 mutants, isolated gDNA, and sequenced whole genomes. Subsequently, we aligned
the sequenced reads to four reference sequences using breseq (51). The reference sequences
included *V. cholerae* chromosomes I and II, and the plasmid conferring natural competence. A
fourth reference sequence (pSC189) containing sequences associated with the transposon was

also included as a "genomic lure" to capture reads that mapped to the transposon. When

examining the breseq output, we specifically looked for instances where new junctions were

created between chromosomes I or II (depending on which chromosome the gene targeted for

homologous recombination was located) and pSC189. Such a junction would indicate a

transposon integration event. Fig. 3 is an example breseq output from one such alignment, which

reports the transposon insertion site in chromosome II, including the basepair position it is

located.

The parent strain of both the donor and recipient libraries is *V. cholerae* El Tor C6706.

However, when the donor library was generated in 2008, transposon insertion sites were mapped

to reference genomes of *V. cholerae* El Tor strain N16961 because it was the only *V. cholerae*

that had been sequenced for *Vibrio spp.* at that time (55). In this work, we mapped transposon

insertion sites to reference genomes of strain C6706. To identify whether transposons insertions

were in the correct site of our recipient strains, we performed whole genomic comparative

analysis of *V. cholerae* N16961 and C6706. When we performed this analysis, we found

chromosomes I and II differ in size between strains and are not syntenic (Fig. 4). Out of the

2,649 genes annotated in our reference genomes for chromosome I, 2,558 (96.6%) of genes are

shared between N16961 and C6706. Additionally, 21 genes (0.79%) are unique to N16961 and

70 (2.64%) to C6706. On chromosome II, 1,052 out of 1,064 annotated genes (98.9%) are shared

between strains, where 10 (0.94%) and 2 (0.19%) of genes are unique to N16961 and C6706,

respectively. Importantly, this analysis provided the genomic position for each ortholog of

N16961 genes present in C6706 (Dataset S1).

We used the orthologous sites to analyze each transposon mutant we sequenced (Dataset

S1, Fig. S2 and S3). Using the breseq annotation data, we calculated the position of the

transposon within each gene as a percent of basepairs into the ORF of C6706. These position

values were previously reported for each strain in the donor library strain N16961. To assess the

relationship, we performed a linear regression of our sequenced C6706 insertions with the

previously described N16961 from the donor library, expecting a perfect linear correlation.

Although 19/23 mutants had an exact match, some of our values diverged from the line of best fit

(Fig. 5), although there was a strong statistical correlation between the variables (Kendall's

coefficient $\tau = 0.752$, $n = 23$, and $P < 0.0001$). Upon further investigation of this anomaly, we

discovered two categories of false negatives. In category one (Fig. S3, Panels A and B), we

observed two cases where genes were annotated in N16961 but not in C6706. Consequently,

these data points lacked a calculated value for the transposon insertion site in C6706 and thus fell

on the y-axis. In the second category, two false negatives were attributed to a difference in the
size of the gene orthologs between N16961 and C6706 (Fig. S3, Panels C and D). This
discrepancy resulted in a shift of the relative position reported for the transposon. After
accounting for these deviations and correcting the data accordingly, we confirmed that the
transposon inserted into the cognate position of the Grant Library mutants we sequenced with
100% precision.
390
Co-transformation of the *luxOG319S* gain of function mutation is an infrequent event.
Since we employed natural competence to transfer the transposon mutations, there is a possibility
that the *luxOG319S* mutation from the donor library might have been co-transformed along with
the transposon into the Grant library. However, unlike the strong selection pressure for the
transposon mutation, selection for the lux mutation would be indirect and weaker, depending on
whether the lux mutation provides a fitness advantage to cells allowing them to outcompete
wildtype cells during outgrowth in selective media.
398
In the 23 genome sequences analyzed above, none had the *luxOG319S* mutation.
However, we did consider that the frequency of transfer of *luxO* mutation could be elevated in
genes for which the transposon is within close genomic proximity to *luxO*. To test this, we
isolated gDNA from 20 randomly sampled transposon mutants in 5kb windows up to 50 kb up-
and downstream of *luxO* (Table S1). We then PCR amplified *luxO* and performed Sanger
sequencing. In only one instance did the *luxOG319S* mutation transfer into a Grant library strain
(Fig. 6). The absolute distance of this transposon mutant is 12,363 away from *luxO*, and three
mutants closer to *luxO* in this analysis did not transfer the mutation. This suggests that proximity
of the transposon mutations to *luxO* increases the chance of transfer of the *luxO* mutation, but its
transfer remains partially stochastic.
409
In summary, our investigation into the potential co-transfer of the *luxOG319S* mutation
with the transposon has shown that while it is possible, it is an infrequent event. Among the 23
genomes we analyzed and the additional Sanger sequencing data we collected, the *luxOG319S*
mutation was observed in only 1 out of 45 (0.02%) genomes, emphasizing the relative rarity of
this co-transfer event. However, when examining transposon insertions near *luxO* from the Grant
library, we recommend the sequence of *luxO* be verified.
416
Natural competence in Grant library strains is IPTG inducible. When generating the Grant
library, we supplemented the outgrowth media with chloramphenicol and kanamycin to select for

maintenance of the plasmid conferring IPTG-induced natural competence, pMMB-*tfoX-qstR*, and
the TnFLG3 transposon, respectively. We reasoned that a library established with strains
harboring this plasmid would be advantageous to the research community as it would facilitate
additional genome edits using chitin-independent natural competence and homologous
recombination upon induction with IPTG. In proof of concept for this idea, we randomly selected
10 mutants from the Grant library and performed a knock-in experiment of a trimethoprim (tm)
resistance gene. In short, we used thymol to isolate whole gDNA from a tm resistant strain and
added 16.2 µg of whole gDNA to our competent cells, which we prepared following our protocol
described in Materials and Methods.
428
After incubating our competent cells with gDNA overnight, we used spot plates to assess
antibiotic phenotypes. All the randomly selected library mutants grew on LB plates
supplemented with trimethoprim between dilution factors of $10^{-3} - 10^{-8}$ (Fig. 7). When also
plated on growth media supplemented with kanamycin, there was parity in the growth
phenotypes observed on tm plates for all but one mutant (strain 8C5). When we examined the
images of the kanamycin supplemented spot plates when the library was constructed (Fig. S1),
we found mutant 8C5 didn't grow there either. We posit that the spurious growth of strain 8C5
likely reflects a frozen subpopulation of our library recipient strain that was able to withstand
kanamycin during outgrowth or contaimination introduced during the spot-plate assay. Taken
together, all the library mutants we tested could undergo additional genome edits with overnight
induction of natural competence.
440
Differential motility responses of recipient library mutants reveal unappreciated epistasis
exists in donor library. *V. cholerae* exhibits high motility by virtue of a single polar flagellum,
a characteristic that contributes significantly to its pathogenesis. *V. cholerae* isolated from active
infections display increased motility compared to lab-grown strains (56) and demonstrated a
heightened ability to colonize hosts (57). Moreover, motility allows *V. cholerae* to survive in its
aquatic environment, enabling it to swim freely as planktonic cells or form biofilms in response
to environmental stress. Furthermore, previous work has demonstrated the involvement of least
40 genes in flagellum biosynthesis and motility, classified into temporally distinct classes (I
through IV) (58–60). Notably, some of these genes are known to trigger an immune response
(56).
451
Motility is an easily screened phenotype, and the functional consequences of null
flagellar gene mutants are well established. Therefore, Cameron et al., 2008 (32), employed

motility assays as one of the validation metrics for their constructed library. In this context, we

also measured motility of the Grant library null flagellar gene mutants. By performing the assay

using paired mutants from the donor and recipient libraries, we reasoned that we would foremost

independently confirm the motility phenotypes reported by Cameron et. al., 2008. Furthermore,

previous studies have indicated that genes in classes III and IV of the flagellum biosynthesis

hierarchy exhibit cell-density dependent phenotypes (61). Thus, the motility assay would serve to

provide supporting evidence for or against our hypothesis that quorum sensing epistasis is a

missing feature of the donor library.

Out of the 33 genes we examined, we successfully replicated the motility phenotype

reported for the donor library's nonmotile mutants in all but one strain (*fliR*) (Fig. 8).

Additionally, we observed that both libraries exhibited nonmotile behavior for *rpoN* (class I) and

flaA,(class III) that are essential for expression of the flagellar biosynthesis genes (Fig. 8).

Furthermore, motility was also abolished for the σ-54 dependent transcriptional regulator

transposon mutants of *flrA* (class I) in both libraries, consistent with previous observations (62).

When comparing motility within classes, paired null mutants from the Cameron and Grant

libraries displayed similar movement on the plates for all classes, except for class III, where a

statistically significant difference was observed (Table 1, two-tailed *t*-test: P = 0.012).

Additionally, two mutant pairs (*flgM* and *flgN*) within class IV biosynthesis genes exhibited

opposite motility phenotypes. While these opposing effects were observed within the same class,

the net effect of class IV genes canceled each other out in the statistical analysis, so this class

comparison was insignificant. Taken together, our results suggest quorum sensing epistasis with

motility genes results in the different motility outcomes as the donor library is locked in a low-

cell density genomic background. Our finding is supported by several other research groups who

as well have observed nuances in different phenotypes resulting from differences in genomic

background (48)

Conclusion

Bacteria employ quorum sensing (QS) to regulate gene expression based on population density.

In the case of *V. cholerae*, the etiological agent of cholera, QS controls various phenotypes as the

population transitions between low and high cell density. Researchers previously constructed an

ordered mutant library, systematically disrupting every non-essential gene in *V. cholerae* with a

transposon insertion. However, unbeknownst to them, this library was created in a strain with a

mutation that rendered the mutants incapable of transitioning between low and high cell density.

In this study, we successfully transferred transposon insertions from these *V. cholerae*
non-redundant ordered mutants into a wildtype genetic background using chitin-independent
natural transformation. The resulting Grant Library comprises 3,102 mutants, covering
approximately 79.8% of the ORFs annotated in *V. cholerae*.
493
In addition to encoding a functional quorum sensing system, another notable advantage
of the Grant Library is that we selected for the transposon insertion and the plasmid that confers
IPTG-inducible natural competence during outgrowth in selective media. Accordingly, upon
induction with IPTG, mutants from the Grant Library can undergo additional genome edits when
co-incubated with gDNA. Furthermore, growth in the absence of selection leads to rapid curing
of the competence plasmid if it is not needed. These features make the Grant Library an
invaluable resource for studying pleiotropic and epistatic genetic interactions in *V. cholerae*,
including combinations that are synthetically lethal. Such information will be invaluable toward
generating new therapeutic options to treat *V. cholerae* infections.
503
Construction of the Grant Library was aided by a novel, in-house gDNA extraction
method we developed based on thymol cell lysis. Our use of thymol allowed extraction of gDNA
in a highly efficient and economical way, producing DNA that was usable in several downstream
applications. Indeed, as many as 576 manual gDNA extractions were performed in a single
workday. We have tested the efficiency of thymol treatment on several other biological
specimens including bacteria in gram -negative and -positive staining classes and yeast. Among
our test strains, gDNA yield was highest for *V. cholerae* (~ 22 ng/µl), although most strains
yielded some gDNA (2 – 8 ng/µl) (data not shown). The apparent vibrio-specificity of thymol
treatment is an interesting avenue for further research as its understanding can provide additional
insight into its antibacterial activity, potentially contributing to novel disease control strategies.
In summary, the Grant Library complements the non-redundant library made in the quorum-
incompetent strain, and the methods used to construct it, a valuable approach toward
understanding *V. cholerae* evolutionary genetics.
517

Acknowledgements
We thank Ewen Cameron and the John Mekalanos lab for constructing the 2008 *V. cholerae* non-
redundant library, Vic DiRita for granting access to a copy of the collection, and Bonnie Bassler
and Julie Valastyan for providing plates that were missing from the DiRita collection. We also
thank Ankur Dalia for providing the plasmid conferring natural competence. This work was
supported by National Institutes of Health Grants GM139537 and AI158433 to C.M.W and a

NIGMS Diversity Supplement to N.A.G. Financial support was also received from The

BEACON Center for the Study of Evolution in Action at Michigan State University (to N.A.G).

Figures and Tables

529
Figure 1. Thymol-Assisted gDNA extraction and chitin-independent natural transformation.

Overnight cultures of the donor library were grown in 550 µl of selective media (LB +

kanamycin). After overnight growth, the samples were centrifuged, the supernatant was

aspirated, and the cell pellets were incubated for one hour at 64°C in a novel gDNA extraction

buffer containing thymol. gDNA was then isolated from the clarified supernatant using an

alcohol precipitation method, followed by resuspension of the gDNA in 1x TE buffer. In parallel,

a quorum-competent wildtype *V. cholerae* strain carrying the plasmid conferring natural

competence (pMMB-*tfoX-qstR*) was grown in selective media (LB + chloramphenicol)

supplemented with IPTG. The next day, competent cells were diluted 1:4 in 0.5x Instant Ocean,

to which chloramphenicol and IPTG were added. Subsequently, 200 µl of cells were dispensed

into the wells of a deep-well 96-well plate and co-incubated with 100 µl of our gDNA extraction

buffer at 30°C for 18 – 24 h. For a detailed description, refer to the Materials and Methods
section.

	Has transposon	Does not have transposon
Identified as having the transposon	3102	15
Identified as not having the transposon	22	71

Measure	Value
Sensitivity	0.9930
Specificity	0.8256
Precision	0.9952
Negative Predictive Value	0.7634
False Positive Rate	0.1744
False Discovery Rate	0.0048
False Negative Rate	0.0070
Accuracy	0.9885
F1 Score	0.9941
Matthews Correlation Coefficient	0.7880

Figure 2. Analysis of Library Transfer. The outcomes of growth on selective media for each
mutant were recorded after chitin-independent natural transformation of the donor library. These
values were utilized as inputs to construct a 2x2 confusion matrix, categorizing the results as
follows: True Positives (3,102 wells), False Negatives (22 wells), True Negatives (71 wells), and
False Positives (15 wells). The accompanying table presents the calculated measures, the values
of which were computed using standard derivations.

Unassigned new junction evidence									
	seq id	position	reads (cov)	reads (cov)	score	skew	freq	annotation	gene	product
?	NZ_CP046845	151866 =	2 (0.050)					coding (225/543 nt)	GPY04_RS15290	N-acetyltransferase
?	pSC189	= 4058	0 (0.000)	37 (0.780)	15/290	0.0	97.4%	intergenic (+121/+620)	aph(3')-II (or nptII)/traJ	aminoglycoside phosphotransferase from Tn5/oriT-recognizing protein
?	NZ_CP046845	= 151868	2 (0.050)					coding (227/543 nt)	GPY04_RS15290	N-acetyltransferase
?	pSC189	1802 =	0 (0.000)	25 (0.520)	14/292	0.0	96.2%	intergenic (–/-41)	–/3xFLAG	–/three tandem FLAG(R) epitope tags, followed by an enterokinase cleavage site

Figure 3. Representative sample of breseq output showing new junction evidence supporting transposon insertion. The figure shows a portion of the summary.html file generated for a breseq analysis run. In this example, a new junction has been called between chromosome II (NZ_CP046845) of *V. cholerae* strain C6706 and the plasmid (pSC189) containing representative sequences of the transposon. Insertions have left and right boundaries, where the chromosome and sequence meet. The boundaries are marked with an asterisk. The annotation column shows the position of the transposon within the gene (GPY04_RS15290), which we converted to a value called "percent into open reading frame." In this example, that value is 41.43% ((225/543) * 100)). This value was calculated for all strains sequenced in this study.

Figure 4. Genomic comparative analysis of *V. cholerae* N16961 and C6706 for chromosome I
(A) and chromosome II (B). Genbank files for whole-genome sequences of chromosomes I and
II of both strains were downloaded from NCBI. The analysis involved a comparative
examination of genetic variations and conservation patterns between the two strains. *V. cholerae*
strain N16961 is shown with a gray background and C6706 a white one. Genome segments are
shown as colored colinear blocks centered around the "positive" and "negative" strands of each
chromosome. Lines connecting the segments show their orthologous alignments between the
strains, including inversions. Panel C is a Venn diagram summarizing the relationship of genes
for each strain and chromosome. The data underlying the illustration in panel C, including the
genomic position of orthologous gene pairs, is provided in the Supplementary Information
(Datasets S2–S4).

Figure 5. Correlation between transposon insertion sites in *V. cholerae* N16961 and C6706. Each point represents the transposon insertion site, calculated as percent into open reading frame. The values for strain N16961 were reported in supplementary dataset 3 of Cameron et al. (2008), and in this work, they were calculated using the data provided in the annotation column of the breseq output (Fig. 3). The correlation analysis yields a strong positive correlation with Kendall's coefficient $\tau = 0.752$ ($n = 23$, and $P \ll 0.0001$), indicating a significant relationship between the insertion sites of the two strains.

Figure 6. Sanger sequence genome alignments. We randomly selected Grant library transposon mutants in 5kb windows up to 50 kp up- and downstream of *luxO*. Subsequently, we isolated gDNA and processed samples for Sanger sequencing. Each row shows a randomly selected transposon mutant (Tn) and their absolute distance downstream (Ds) or upstream (Us) of *luxO*. The gain of function mutation in *luxO* is annotated as a glycine to serine mutation at amino acid position 319. Our Sanger sequencing analysis identified this mutation in one mutant (W3_TnDs_12363), indicating the mutation transfers at low frequency and only in genes within close proximity to *luxO*.

Figure 7. Chromosomal integration of a trimethoprim resistant gene in select mutants from the
Grant library using IPTG-induced natural competence. We induced natural competence in
wildtype *V. cholerae* strain C6706 carrying plasmid pMMB-*tfoX-qstR* (NG001) and 10 randomly
selected mutants from the Grant library according to procedure described in Materials and
Methods. These strains were then co-incubated with gDNA extracted from a strain carrying a
trimethoprim resistant cassette and on the following day spotted on antibiotic selection media.
All the strains carrying the plasmid grew on plates supplemented with trimethoprim (Panel A).
On the plate with kanamycin (Panel B) or kanamycin + trimethoprim (Panel C), all strains grew
except strain 8C5, indicating that this strain does not have a transposon insertion.

Figure 8. Motility behavior of flagellar mutants from the Grant and Cameron et al. (2008)
libraries. Paired mutants with transposons in flagellar genes from both libraries were screened for
motility in nutrient-rich plates containing 0.35% agar. The plates were imaged, and the area
moved by each mutant from an inoculum site was calculated using the procedure described in the
Materials and Methods. These areas were then normalized to a value relative to the wild-type
strains for each library from 3 replicate assays in all but four cases (*flgM*, *flhB, fliR*, and *flgF*),
which had two replicates each. The dashed red line is the normalization for the wildtype strains.
Colored boxes highlight the class each gene belongs to in the flagellar biosynthesis gene cluster
hierarchy, and red stars highlight the expected non-motile phenotype for transposon insertions
that lie within genes encoding sigma factors required to transcribe downstream flagellar
biosynthesis genes.

Table 1. Two-sided *t*-test results comparing the motility of paired Grant and Cameron et. al., (2008). Flagellar mutants assayed in this study are grouped by the flagellar biosynthesis gene cluster (I-IV) they belong to.

Cluster	Sample size (n)	Mean difference	df	*t*	*P*
I	3	0.452	2	1.64	0.242
II	13	-0.045	12	-0.37	0.720
III	14	0.999	13	2.92	0.012
IV	3	0.072	2	0.09	0.935

References

1. M. Ali, A. R. Nelson, A. L. Lopez, D. A. Sack, Updated Global Burden of Cholera in Endemic Countries. *PLOS Neglected Tropical Diseases* **9**, e0003832 (2015).

2. I. Ilic, M. Ilic, Global Patterns of Trends in Cholera Mortality. *Trop Med Infect Dis* **8**, 169 (2023).

3. D. Legros, Global Cholera Epidemiology: Opportunities to Reduce the Burden of Cholera by 2030. *J Infect Dis* **218**, S137–S140 (2018).

4. , Roadmap 2030 – Global Task Force on Cholera Control (October 22, 2023).

5. D. Lippi, E. Gotuzzo, S. Caini, Cholera. *Microbiology Spectrum* **4**, 10.1128/microbiolspec.poh-0012–2015 (2016).

6. S. Yamai, T. Okitsu, T. Shimada, Y. Katsube, Distribution of serogroups of *Vibrio cholerae* non-O1 non-O139 with specific reference to their ability to produce cholera toxin, and addition of novel serogroups. *Kansenshogaku Zasshi* **71**, 1037–1045 (1997).

7. A. Safa, G. B. Nair, R. Y. C. Kong, Evolution of new variants of *Vibrio cholerae* O1. *Trends in Microbiology* **18**, 46–54 (2010).

8. J. B. Kaper, J. G. Morris, M. M. Levine, Cholera. *Clin Microbiol Rev* **8**, 48–86 (1995).

9.   D. Hu, *et al.*, Origins of the current seventh cholera pandemic. *Proceedings of the National*
Academy of Sciences **113**, E7730–E7739 (2016).

10.   K. Okada, T. Iida, K. Kita-Tsukamoto, T. Honda, Vibrios Commonly Possess Two
Chromosomes. *J Bacteriol* **187**, 752–757 (2005).

11.   S. Sozhamannan, T. Waldminghaus, Exception to the exception rule: synthetic and
naturally occurring single chromosome *Vibrio cholerae*. *Environmental Microbiology* **22**,
4123–4132 (2020).

12.   D. K. Karaolis, R. Lan, P. R. Reeves, The sixth and seventh cholera pandemics are due to
independent clones separately derived from environmental, nontoxigenic, non-O1 *Vibrio*
cholerae. *J Bacteriol* **177**, 3191–3198 (1995).

13.   S. M. Faruque, *et al.*, Emergence and evolution of *Vibrio cholerae* O139. *Proceedings of*
the National Academy of Sciences **100**, 1304–1309 (2003).

14.   T. Ramamurthy, *et al.*, Revisiting the Global Epidemiology of Cholera in Conjunction With
the Genomics of *Vibrio cholerae*. *Front Public Health* **7**, 203 (2019).

15.   S. N. Sakib, G. Reddi, S. Almagro-Moreno, Environmental Role of Pathogenic Traits in
Vibrio cholerae. *J Bacteriol* **200**, e00795-17 (2018).

16.   Y. Sun, E. E. Bernardy, B. K. Hammer, T. Miyashiro, Competence and Natural
Transformation in Vibrios. *Mol Microbiol* **89**, 10.1111/mmi.12307 (2013).

17.   R. L. Marvig, M. Blokesch, Natural transformation of *Vibrio cholerae* as a tool -
Optimizing the procedure. *BMC Microbiology* **10**, 155 (2010).

18.   S. Stutzmann, M. Blokesch, Comparison of chitin-induced natural transformation in
pandemic *Vibrio cholerae* O1 El Tor strains. *Environ Microbiol* **22**, 4149–4166 (2020).

19.   K. L. Meibom, M. Blokesch, N. A. Dolganov, C.-Y. Wu, G. K. Schoolnik, Chitin Induces
Natural Competence in *Vibrio cholerae*. *Science* **310**, 1824–1827 (2005).

20.   S. S. Watve, J. Thomas, B. K. Hammer, CytR Is a Global Positive Regulator of
Competence, Type VI Secretion, and Chitinases in *Vibrio cholerae*. *PLoS One* **10**,
e0138834 (2015).

21. M. Lo Scrudato, S. Borgeaud, M. Blokesch, Regulatory elements involved in the expression
of competence genes in naturally transformable *Vibrio cholerae*. *BMC Microbiology* **14**,
327 (2014).

22. M. L. Scrudato, M. Blokesch, The Regulatory Network of Natural Competence and
Transformation of *Vibrio cholerae*. *PLOS Genetics* **8**, e1002778 (2012).

23. E. S. Antonova, B. K. Hammer, Genetics of Natural Competence in *Vibrio cholerae* and
other Vibrios. *Microbiol Spectr* **3**, 3.3.20 (2015).

24. A. B. Dalia, D. W. Lazinski, A. Camilli, Identification of a Membrane-Bound
Transcriptional Regulator That Links Chitin and Natural Competence in *Vibrio cholerae*.
mBio **5**, 10.1128/mbio.01028-13 (2014).

25. T. N. Dalia, *et al.*, Multiplex genome editing by natural transformation (MuGENT) for
synthetic biology in *Vibrio natriegens*. *ACS Synth Biol* **6**, 1650–1655 (2017).

26. A. B. Dalia, "Natural Cotransformation and Multiplex Genome Editing by Natural
Transformation (MuGENT) of *Vibrio cholerae*" in *Vibrio Cholerae: Methods and
Protocols*, Methods in Molecular Biology., A. E. Sikora, Ed. (Springer, 2018), pp. 53–64.

27. A. B. Dalia, E. McDonough, A. Camilli, Multiplex genome editing by natural
transformation. *Proc Natl Acad Sci U S A* **111**, 8937–8942 (2014).

28. T. Baba, *et al.*, Construction of *Escherichia coli* K-12 in-frame, single-gene knockout
mutants: the Keio collection. *Mol Syst Biol* **2**, 2006.0008 (2006).

29. M. M. Pearson, S. Pahil, V. S. Forsyth, A. E. Shea, H. L. T. Mobley, Construction of an
Ordered Transposon Library for Uropathogenic *Proteus mirabilis* HI4320. *Microbiol Spectr*
10, e03142-22.

30. B. Ramage, *et al.*, Comprehensive Arrayed Transposon Mutant Library of *Klebsiella
pneumoniae* Outbreak Strain KPNIH1. *Journal of Bacteriology* **199**, 10.1128/jb.00352-17
(2017).

31. P. D. Fey, *et al.*, A Genetic Resource for Rapid and Comprehensive Phenotype Screening of
Nonessential *Staphylococcus aureus* Genes. *mBio* **4**, 10.1128/mbio.00537-12 (2013).

32.   D. E. Cameron, J. M. Urbach, J. J. Mekalanos, A defined transposon mutant library and its
use in identifying motility genes in *Vibrio cholerae*. *Proceedings of the National Academy*
of Sciences **105**, 8736–8741 (2008).

33.   H. Rui, *et al.*, Reactogenicity of live-attenuated *Vibrio cholerae* vaccines is dependent on
flagellins. *Proceedings of the National Academy of Sciences* **107**, 4359–4364 (2010).

34.   R. W. Bogard, B. W. Davies, J. J. Mekalanos, MetR-Regulated *Vibrio cholerae* Metabolism
Is Required for Virulence. *mBio* **3**, 10.1128/mbio.00236-12 (2012).

35.   T. G. Dong, B. T. Ho, D. R. Yoder-Himes, J. J. Mekalanos, Identification of T6SS-
dependent effector and immunity proteins by Tn-seq in *Vibrio cholerae*. *Proceedings of the*
National Academy of Sciences **110**, 2623–2628 (2013).

36.   A. J. Van Alst, V. J. DiRita, Aerobic Metabolism in *Vibrio cholerae* Is Required for
Population Expansion during Infection. *mBio* **11**, 10.1128/mbio.01989-20 (2020).

37.   S. T. Miyata, D. Unterweger, S. P. Rudko, S. Pukatzki, Dual Expression Profile of Type VI
Secretion System Immunity Genes Protects Pandemic *Vibrio cholerae*. *PLOS Pathogens* **9**,
e1003752 (2013).

38.   A. Hsiao, Z. Liu, A. Joelsson, J. Zhu, *Vibrio cholerae* virulence regulator-coordinated
evasion of host immunity. *Proceedings of the National Academy of Sciences* **103**, 14542–
14547 (2006).

39.   F. P. Rothenbacher, J. Zhu, Efficient responses to host and bacterial signals during *Vibrio*
cholerae colonization. *Gut Microbes* **5**, 120–128 (2014).

40.   S. Saha, S. Aggarwal, D. V. Singh, Attenuation of quorum sensing system and virulence in
Vibrio cholerae by phytomolecules. *Frontiers in Microbiology* **14** (2023).

41.   J. Zhu, *et al.*, Quorum-sensing regulators control virulence gene expression in *Vibrio*
cholerae. *Proceedings of the National Academy of Sciences* **99**, 3129–3134 (2002).

42.   J. Zhu, J. J. Mekalanos, Quorum Sensing-Dependent Biofilms Enhance Colonization in
Vibrio cholerae. *Developmental Cell* **5**, 647–656 (2003).

43.   M. Jemielita, N. S. Wingreen, B. L. Bassler, Quorum sensing controls *Vibrio cholerae*
multicellular aggregate formation. *eLife* **7**, e42057 (2018).

44.   M. M. Hoque, *et al.*, Quorum Regulated Resistance of *Vibrio cholerae* against
744   Environmental Bacteriophages. *Sci Rep* **6**, 37956 (2016).

745   45.   H. Boyaci, *et al.*, Structure, Regulation, and Inhibition of the Quorum-Sensing Signal
746   Integrator LuxO. *PLoS Biol* **14**, e1002464 (2016).

747   46.   R. Herzog, N. Peschek, K. S. Fröhlich, K. Schumacher, K. Papenfort, Three autoinducer
748   molecules act in concert to control virulence gene expression in *Vibrio cholerae*. *Nucleic*
749   *Acids Res* **47**, 3171–3183 (2019).

750   47.   S. A. Jung, C. A. Chapman, W.-L. Ng, Quadruple Quorum-Sensing Inputs Control *Vibrio*
751   *cholerae* Virulence and Maintain System Robustness. *PLOS Pathogens* **11**, e1004837
752   (2015).

753   48.   S. Stutzmann, M. Blokesch, Circulation of a Quorum-Sensing-Impaired Variant of *Vibrio*
754   *cholerae* Strain C6706 Masks Important Phenotypes. *mSphere* **1**, 10.1128/msphere.00098-
755   16 (2016).

756   49.   A. Escobar, M. Pérez, G. Romanelli, G. Blustein, Thymol bioactivity: A review focusing on
757   practical applications. *Arabian Journal of Chemistry* **13**, 9243–9269 (2020).

758   50.   J. B. Brown, Classifiers and their Metrics Quantified. *Mol Inform* **37**, 1700127 (2018).

759   51.   D. E. Deatherage, J. E. Barrick, Identification of mutations in laboratory evolved microbes
760   from next-generation sequencing data using breseq. *Methods Mol Biol* **1151**, 165–188
761   (2014).

762   52.   A. C. E. Darling, B. Mau, F. R. Blattner, N. T. Perna, Mauve: Multiple Alignment of
763   Conserved Genomic Sequence With Rearrangements. *Genome Res* **14**, 1394–1403 (2004).

764   53.   M. Kearse, *et al.*, Geneious Basic: An integrated and extendable desktop software platform
765   for the organization and analysis of sequence data. *Bioinformatics* **28**, 1647–1649 (2012).

766   54.   J. Schindelin, *et al.*, Fiji: an open-source platform for biological-image analysis. *Nat*
767   *Methods* **9**, 676–682 (2012).

768   55.   J. F. Heidelberg, *et al.*, DNA sequence of both chromosomes of the cholera pathogen *Vibrio*
769   *cholerae*. *Nature* **406**, 477–483 (2000).

56. M. Grognot, A. Mittal, M. Mah'moud, K. M. Taute, *Vibrio cholerae* Motility in Aquatic
and Mucus-Mimicking Environments. *Applied and Environmental Microbiology* **87**,
e01293-21 (2021).

57. F. Khan, N. Tabassum, R. Anand, Y.-M. Kim, Motility of *Vibrio* spp.: regulation and
controlling strategies. *Appl Microbiol Biotechnol* **104**, 8187–8208 (2020).

58. M. A. Echazarreta, K. E. Klose, *Vibrio* Flagellar Synthesis. *Frontiers in Cellular and
Infection Microbiology* **9** (2019).

59. K. A. Syed, *et al.*, The *Vibrio cholerae* Flagellar Regulatory Hierarchy Controls Expression
of Virulence Factors. *J Bacteriol* **191**, 6555–6570 (2009).

60. S. Zhu, S. Kojima, M. Homma, Structure, gene regulation and environmental response of
flagella in *Vibrio*. *Frontiers in Microbiology* **4** (2013).

61. S. Beyhan, A. D. Tischler, A. Camilli, F. H. Yildiz, Transcriptome and Phenotypic
Responses of *Vibrio cholerae* to Increased Cyclic di-GMP Level. *J Bacteriol* **188**, 3600–
3613 (2006).

62. D. S. Millikan, E. G. Ruby, FlrA, a σ54-Dependent Transcriptional Activator in *Vibrio
fischeri*, Is Required for Motility and Symbiotic Light-Organ Colonization. *J Bacteriol* **185**,
3547–3557 (2003).

787