1

6    **Spatial trajectories of coffee harvesting in large-scale plantations: ecological and**

7    **management drivers and implications**

9    Emilio Mora Van Cauwelaert[a, b, *], Denis Boyer[c], Estelí Jiménez-Soto[d] , Cecilia González González[a]

10   and Mariana Benítez[a, *]

11       a.   Laboratorio Nacional de Ciencias de la Sostenibilidad, Instituto de Ecología, Universidad Nacional Autónoma de

12            México, Mexico City, C.P. 04510, México

13       b.   Posgrado en Ciencias Biomédicas, Universidad Nacional Autónoma de México, Mexico City, C.P. 04510, México

14       c.   Instituto de Física, Universidad Nacional Autónoma de México, Mexico City, C.P. 04510, México

15       d.   Geography, Environmental Science and Policy, School of Geosciences, College of Arts and Sciences, University

16            of South Florida USF, USA

18           *mbenitez@iecologia.unam.mx, *emiliomora92@ciencias.unam.mx

**ABSTRACT**

CONTEXT: Coffee is produced under different management systems and scales of production

categorized as Syndromes of Production. The "Capitalist Syndrome" is characterized by the high use

of capital and labor inputs to increase agricultural outputs. This syndrome results in practices like high

planting densities that may promote the development and dispersal of plant pathogens like coffee leaf

rust. The spatial arrangement of coffee trees drives the spatial movement of the harvesters, who can

bear and disperse pathogens across and within plantations. In most capitalist coffee plantations,

harvesters work multiple hours to maximize the daily harvest, which might increase their dispersal

potential. However, their spatial movement has not yet been described, nor its relationship with the

scale or management of the plantation, and even less its ecological implications for pathogens

dispersal.

OBJECTIVE: We describe and analyze the daily spatial movement of coffee harvesters in two large-

scale capitalist plantations: an organic and a conventional plantation.

METHODS: Using state-space models, we recorded and analyzed the spatial movements of

harvesters. We then constructed a driver tree for harvest dynamics, which incorporated qualitative

variables related to climate, coffee plants, and management aspects reported by the harvesters.

RESULTS AND CONCLUSIONS: Our model differentiated two kinds of movements: 1) when trees

have berries, harvesters remain in the coffee rows or areas nearby (Collect state; 94-98% of the steps);

2) when not, harvesters make longer steps within the harvesting location or move to another area

(Search state; 2-6% of the steps). In the organic plantation, the Search state had a longer-tailed step-

length distribution than in the conventional plantation, resulting in a significantly higher visited area

per worker ($p<0.05$). This might be related to a) a lower fruit load or percentage of trees with ripe

fruits when we took the data or b) smaller harvesting locations ("*pantes*") per number of harvesters.

Harvesting movements that explore a wider area, either by visiting more plants or by changing

locations on the same day, could create more foci of CLR infection across the plantation.

SIGNIFICANCE: Our results highlight practices that can reduce the possible impact of human

dispersal of pathogens like shorter harvesting trajectories by working fewer hours a day or avoiding

harvesting at the end of the maturation season when few trees have berries and harvesters have to

travel longer distances. This calls for organic coffee management that could prevent diseases, increase

diversity, and guarantee just and safe conditions for workers.

**KEYWORDS**

Syndromes of production; state-space models; harvesting movement; coffee leaf rust dispersal

## 1. INTRODUCTION

In recent decades, the increased industrialization of agroecosystems has gathered significant attention due to its profound negative social and ecological consequences (Raven and Wagner, 2021). Agricultural industrialization and environmental change have led to increases in pest populations (Andow, 1983), biodiversity decline (Raven and Wagner, 2021), and are also associated with a higher likelihood of zoonotic disease outbreaks (Jones et al., 2013; Perfecto et al., 2023). Adding to these challenges are the heightened demands on human labor and precarious social conditions experienced in large-scale industrial agricultural systems (Mapes, 2010), a social phenomenon that imparts more complexity to agroecosystem management and processes.

Coffee agroecosystems exhibit a large variety of management practices that reinforce each other and that can be defined as "syndromes of production", a conceptual framework useful to evaluate social and ecological dynamics across distinct agricultural paradigms (Andow and Hidaka, 1989; Moguel y Toledo, 1999; Ong and Liao, 2020; Vandermeer and Perfecto, 2012). Coffee production syndromes can be mainly distinguished by shade management (shade or sun coffee plantations; Vandermeer and Perfecto, 2012) or by the type of land tenure and scale of production (large-scale capitalist or small-scale peasant-owned coffee plantations; Ong and Liao, 2020). In this sense, coffee agroecosystems are model systems to understand how different syndromes of production affect ecological processes such as their planned and associated biodiversities (Mas and Dietsch, 2004) but also social aspects like the worker's conditions (Jiménez-Soto, 2021).

The capitalist syndrome follows a maximization of profit rule (Ong and Liao, 2020); it is typically based upon large-scale and high-density planting farms with a high input of human labor and exploitation, although it may have specific ecological practices according to organic or conventional coffee markets (Perfecto et al., 2019). This syndrome of coffee production results in practices that may promote the development and spread of important pests (Avelino et al., 2006; Hajian-Forooshani et al., 2016). For instance, the coffee leaf rust (CLR) can travel across and within the coffee plantations through direct leaf contact, water splash, or local turbulent wind conditions (Becker and Kranz, 1977; Vandermeer et al., 2018), but its effectiveness of dispersal and impact can be modulated by management practices like the planting density (Mora Van Cauwelaert et al., 2023), the presence of wind-barrier trees in the plantation (Gagliardi et al., 2020), or the biotic communities of natural enemies (Jackson et al., 2012), all determined to some degree by the syndrome of production (Vandermeer and Perfecto, 2012; McCook and Vandermeer). In this regard, increased agricultural simplification and industrialization of coffee agroecosystems have led to massive outbreaks of CLR around the globe (Avelino et al., 2015; McCook and Vandermeer, 2015).

Some studies have proposed that harvesters can also bear CLR spores during harvesting (Becker and Kranz, 1977; Waller, 1981). Interestingly, the range of movement and duration of the daily harvest are also determined by the management, the scale, the land ownership, and by the

agricultural characteristics of the coffee plantations. In most landlord-owned large-scale coffee
plantations, harvesters are paid by piecework and work several hours to maximize the daily harvest
(Jiménez-Soto, 2021; López Echeverría, 2006). These practices affect the social and health conditions
of the workers, which are often precarious. Besides, characteristics like high planting density can also
drive the spatial movement during harvesting. We argue that the spatial movement of the harvesters
under capitalist labor-intensive conditions could also have an ecological impact, through modulating
the human dispersal of pathogens like coffee leaf rust (CLR). In fact, other studies have stressed the
relation between the landscape structure, individual movement behavior, and pathogen transmission
for predicting and understanding disease dynamics (Manlove et al., 2022; White et al., 2018).

However, the spatial movement of the harvesters in plantations within this capitalist
syndrome of production has not been described yet, nor its relationship with the management of the
plantation, and even less its ecological implications for CLR or other potential pest dispersal. Here we
present a description and qualitative analysis of the spatial movement of farmworkers during
harvesting in two large-scale landlord-owned plantations (~300 ha) with two different types of
management: an organic shaded production coffee plantation and a sun-grown production with few
scattered trees. Specifically, we asked 1) how to statistically characterize the spatial trajectories of
workers during harvesting, 2) how different features of the plantations, determined within a specific
syndrome of production, affect these trajectories, and 3) what are the socio-ecological consequences
of the harvesting movement in these large scale plantations.

We first recorded and analyzed with state-space models the daily spatial movements of the
harvesters, as well as their differences between both plantations. We then constructed a driver tree for
harvest dynamics, which incorporated qualitative variables related to climate, as well as coffee
biology and management aspects reported by the harvesters during interviews and noted through our
own observations. We also highlighted the relationship between these variables and the syndrome of
production. Finally, we put in conversation the patterns and dynamics of the spatial movement of
harvesters with the social conditions associated with the large-scale capitalist syndrome and the
possible implications in pathogen dynamics such as CLR spread.

## 2. METHODS

### 2.1. Site of study

We selected two large-scale coffee plantations located in the Soconusco region of Chiapas, Mexico. This region has a tropical wet and dry climate (Koppen classification, Aw), with a rainy season from May to October, an annual mean temperature of 27 °C, and a precipitation of 2800 mm (Villarreal-Treviño et al., 2015). The first one (organic plantation, henceforth) is an organic shaded coffee plant plantation with a high plant diversity of up to 200 species. The second one (the conventional plantation) has a lower shade cover dominated by few *Inga sp*, uses pesticides, and has a high rate of coffee tree renovation to increase the productivity of regions affected by coffee leaf rust (CLR). Both plantations are approximately 300 ha, are landlord-owned and produce different coffee varieties for exportation. Harvesting is carried out by permanent or temporary workers from Central America, primarily Guatemala and El Salvador, from October to February, depending on the previous rainy season. Workers are paid by piecework, either by weight or volume, and live in precarious conditions (Jiménez-Soto, 2021). During harvest, the foreman assigns the harvesters and their families specific areas called "*pante*" (a Nahuatl word referring to agricultural locations of variable size) to work during one or multiple days. Harvesters remain in these locations during the day or travel to another area when they finish harvesting all the trees in the *pante*.

### 2.2. Data Collection

We carried out field research for three weeks between October and November of 2021. We accompanied 12 families of harvesters during their harvesting day, six on each plantation, after their explicit consent to our research. The trajectories were labeled from C1 to C6 and from O1 to O6 for the conventional and organic plantations, respectively. We worked with them for six hours (including resting and eating pauses), completing three tasks: i) We recorded the movement of one harvester from each family by marking the location of every newly harvested tree, ii) we carried out semi-structured walking interviews (Evans and Jones, 2011) with each family and participant observation during harvesting to assess the factors that determined the different trajectories and to gather information about the whole syndrome of production, and iii) we participated in harvest, as one team member acquired the necessary harvesting skills along with experienced harvesters and donated their daily harvest to contribute to the family process.

### 2.2.1. Trajectories data

We added a waypoint for each consecutive visited tree using the satellite remote sensing Global Positioning System (GPS) Garmin 65. This system is ideal to analyze fine-scale movement patterns as its relative distance error between two consecutive recorded points is small (<0.5 m) (Breed and Severns, 2015). The recorded tracks were homogenized and curated by removing the pauses and

normalizing the UTM coordinates. We defined the step length as the distance between two harvested

trees; this distance provides a good proxy of the general movement of harvesters and is also relevant

to understand how patterns of movement could explain the dispersal of CLR or other potential pests

from one tree to another. Every pair of trees closer than one meter was considered as the same point

(Vandermeer et al., 2018).

### 2.2.2 Walking interviews

We designed a questionnaire to gather information on the organization of harvest and applied it as a

walking interview with the harvesters (Evans and Jones, 2011). We included questions about the

period of the harvesting season, the number of times each plot was harvested, and the differences in

the harvesting dynamics along the season between both plantations (organic and conventional). In

particular, we asked who decided the size of the groups, the starting point, and the daily length of

harvest. We also asked the harvesters directly about the factors that affected the motion of harvesting

within each plant and between the plants. We added questions about the CLR control in both

plantations and, last but not least, about the origins and social conditions of the harvesters. Before this

process, we explained the purpose of the study and gave an informative letter to each of the

participants to obtain their informed consent to participate in the study. The data was complemented

with personal observations on the characteristics and times of harvesting per plant, the size of the

harvesting groups, and the differences in topography and ecological management between both

plantations like the planting pattern of coffee trees.

### 2.3. Movement analysis and qualitative drivers of motion

We analyzed the harvester movement data using state-space models (Patterson et al., 2008) and

organized the factors that influence the motion in a qualitative driver tree. All data analyses and

figures were done in *Rstudio* 2023.03.1+446 using *plyr, dplyr, tidyverse, ggplot2, patchwork* libraries,

and *Inkscape 1.0 and yEd 3.23.2.* All code and data required to reproduce results in the work can be

accessed at https://github.com/tenayuco/harvestDistribution and it is further explained in

supplementary material.

### 2.3.1. Movement analysis, state-space models and differences between the plantations

We analyzed the harvesters' movement using multi-state distribution models. State-space models are

typically used in the context of movement ecology analysis to characterize the trajectories of

individual animals that are assumed to have different behaviors or states (eg. foraging, exploring,

migrating) during the recorded movement (Michelot and Blackwell, 2019; Morales et al., 2004;

Patterson et al., 2008). These behaviors are derived by characteristics like the relative angle between

two steps, or the step length. We used this approach to statistically quantify differences in the daily

movement of the harvesters (for the complete details, refer to Supplementary Material).

We first fitted the step length frequency distributions of the six recorded trajectories of each
plantation to models of one and two-states distributions (i.e. models assuming that the observed
movements are generated by one or by two behavioral states during harvest). In the two-state case, we
termed the behaviors as "Collect" and "Search". In all cases, we used two different fitting
distributions, namely, the Gamma (G(x)) and Weibull (W(x)) functions (see Equations (1-3);
Fig.S1.3; Table S1.2). For *x* and *α* >*0* , these normalized probability density functions (PDFs) are
given by the following formula:

$$W(x) = \frac{\alpha}{\lambda}\left(\frac{x}{\lambda}\right)^{(\alpha-1)} e^{-(x/\lambda)^{\alpha}}$$    (1)

$$G(x) = \frac{\frac{1}{\lambda}\left(\frac{x}{\lambda}\right)^{(\alpha-1)} e^{-(x/\lambda)}}{\Gamma(\alpha)}$$    (2)

$$\Gamma(\alpha) = \int_{0}^{\infty} y^{(\alpha-1)} e^{(-y)} dy$$    (3)

where *x* is the step length (in m), *α* is a shape parameter (no units)**,** *λ* a scale parameter (in m) and
*Γ(α)* the gamma function (Pishro-Nik, 2014). In the two-state case, the Search state should typically
be characterized by a significantly larger *λ* than the Collect state. In each fitting process, we used the
forward algorithm implemented in the *movehmm* library (Michelot et al., 2016) with 1000
combinations of prior parameters from uniform distributions that included the mean and the
maximum step lengths of the data (Table 1). For the analysis, the steps were time-irregular and we
assumed no differences between the harvesters. We compared the Gamma and Weibull models, as
well as the one-state and two-state models, using the Akaike (AIC) criteria (Michelot et al., 2016). As
the two-states distribution of the Gamma family (equation 2-3) generated a lower AIC for both
plantations, we chose it to assign to each step of the trajectories one of the two states (Search and
Collect), using Hidden Markov Models (HMM) and the Viterbi algorithm (Zucchini et al., 2016).
Hidden Markov models (HMMs) assume that the distribution that generates an observation Zt (a step-
length) depends on the state St of an underlying and unobserved Markov process. In brief, this
procedure jointly fits the two-state distribution of step-lengths and establishes a sequence of states
using a maximum likelihood procedure (Patterson et al. 2008; Zucchini et al., 2016). It first assigns to
each step the most likely state and then corrects the state using a global decoding procedure with the
whole sequence (Zucchini et al., 2016). Finally, we explored the differences in the obtained
distributions for each state (Search and Collect) between both plantations and their implications in the
number of trees and surface area visited. We divided the total area into squares with different sizes
(from 1 to 40 m wide) to account for the effect of the grain size and assumed that a square was visited
if at least one visited tree fell within it. We tested the significance of the difference between the means

for each measurement using the Wilcoxon test (w) or the Student test (st) depending on the normality
of the residuals.
240

### 2.3.2. Qualitative drivers of the spatial trajectories

To understand the qualitative drivers of harvesters' trajectories we organized the variables obtained
during the walking interviews to analyze their relation with the longitude, the direction of the
trajectories, or the presence of short and long tailed step-length distributions. We separated the
variables into two categories: environmental or coffee biology-related (e.g. rainfall or coffee
interplant ripening percentage) and those that could be directly affected by management decisions or,
in general, by the syndrome of production (e.g. coffee variety, size of the *pantes*). Finally, we
connected these variables when they influenced each other and categorized the connection into direct
and indirect drivers. Direct drivers included those whose modification affected the trajectory. Indirect
factors affected direct factors (but not the trajectory itself).
251

## 3. RESULTS

*3.1 The movement of the harvesters can be described by two-state distribution models with qualitative differences between the plantations*

Harvesters visited 94.5 trees on average (± 25.6) during the three to four hours recorded daily, excluding pauses. The average time between each visited tree was 2.3 min (± 1.8 mins). The spatial trajectories of the 12 harvesters were largely determined by the planting pattern of coffee trees (Fig.1). Within these patterns some harvesters followed more or less specific coffee tree rows (C_1 to C_3, C_5, O_6) or jumped from one row to another or other regions, depicting more randomized trajectories (C_4, C_6, O_2 to O_5) (Fig. 1). The aggregated distribution of the step lengths of harvesters in each plantation (*n*=6) is represented in Fig. 2. In both plantations the distribution is skewed to the right. In the conventional plantation the mean step length is 4.2 m and the median is 3.2 m whereas in the organic plantation these values are 6 m and 4.5 m, respectively. In some cases, harvesters could travel up to 50 m from one tree to another (conventional plantation), or even more than 100 m (organic plantation) (see inset boxplots in Fig.2).

Models with two distinct state distributions provided a better fit to the aggregated step length distribution as they exhibited lower general AICs, for both Gamma and Weibull distributions, than models with only one state (Table 1 and Fig. S1.5). In particular, the Gamma family with two states had the lowest AIC in both plantations (Table 1). Each step along the trajectories was assigned a state in Fig. 1 according to the distributions of Fig.2. In our work, the states are interpreted as a proxy for the situation during harvest. We named the first state "Collect", where the worker remains in the same vicinity of the *pante* harvesting nearby plants (Fig.1). The PDF of this state is qualitatively similar in both plantations (Table 1, Fig. 2) and encompasses steps from 1 to 8.6 m in the conventional plantation (94% of the steps) and from 1 to 18 m in the organic plantation (98% of the steps) (Fig.2). In the second state, the "Search" state, harvesters traveled to another zone of the *pante* or to another *pante* when the current one had been completely harvested (Fig. 1). This second Search state is described by two qualitatively different PDFs depending on the plantation (Table 1, Fig.2). In the conventional plantation, it includes steps from 7.4 to 44 m and represents movements within the *pante* (6% of steps, dark blue line in Fig. 1 and Fig.2). In the organic plantation, this state includes steps from 21 to 117 m long and represents mainly movements between *pantes* or relocalizations (2% of steps, dark blue line in Fig. 1 and Fig.2). Also note that in the organic plantation, the shape parameter α is close to 1 in the Search state (exponential distribution) while it is above 3 in the Collect state (distribution with a maximum at a finite step length). In the conventional plantation, the difference between the two states is less pronounced.

**Table 1-** Best models for each plantation and distribution family, for one and two-state models (State
1=Collect, State 2=Search). We added the prior parameters for parameter P1 (mean and shape for
gamma and weibull, respectively) and P2 (standard deviation and scale for gamma and weibull,
respectively) (Michelot et al., 2016). We show the minimal negative likelihood (minNegL), AIC
values and the final parameters obtained and transformed to shape and scale following the following
relations: $\alpha = \mu^2/\sigma^2$, $\lambda = \sigma^2/\mu$, where α:shape, λ:scale, μ:mean, σ:standard deviation. The parameters in
bold are the ones we used for the analysis (equations (2)-(3)).

| Farm | Number of states | Family | Priors P1 | Priors P2 | minNegL | AIC | State 1 | | State 2 | |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | | | | | Shape (α) | Scale (λ) | Shape (α) | Scale (λ) |
| Conventional | Two states | gamma | U(0.1, 10) | U(0.1, 20) | **1060** | **2134** | **4,95** | **0,70** | **2,45** | **4,10** |
| | | weibull | U(0, 2.7) | U(0.1, 15) | 1079 | 2172 | 1,46 | 10,12 | 2,42 | 3,87 |
| | One state | gamma | U(0.1, 10) | U(0.1, 20) | 1120 | 2244 | 2,85 | 1,22 | NA | NA |
| | | weibull | U(0, 2.7) | U(0.1, 15) | 1170 | 2344 | 1,47 | 4,66 | NA | NA |
| Organic | Two states | gamma | U(0.1, 10) | U(0.1, 20) | **1553** | **3119** | **3,20** | **1,56** | **1,14** | **29,00** |
| | | weibull | U(0, 2.7) | U(0.1, 15) | 1571 | 3156 | 1,96 | 5,47 | 0,97 | 21,25 |
| | One state | gamma | U(0.1, 10) | U(0.1, 20) | 1674 | 3351 | 1,92 | 3,68 | NA | NA |
| | | weibull | U(0, 2.7) | U(0.1, 15) | 1722 | 3448 | 1,16 | 6,35 | NA | NA |

**Figure 1. Harvesting trajectories from the conventional (C) and from the organic (O) plantations.** We normalized the UTM coordinates from 0 to 150 m to represent the trajectories. The colors represent the most likely state to which each step belongs for each of the farms (orange: Collect, dark blue: Search; see the text for a full explanation). Each black point represents one harvested tree.

**Figure 2. Histograms and box plots (insets) of the step lengths for the trajectories in each**
**plantation and the probability density function of the gamma distributions for each of the two**
**states**. In the conventional plantation (C), the parameters of the gamma distribution ($\alpha$, $\lambda$) are (4.95,
0.7) in the Collect state (orange line) and (2.45, 4.10) in the Search state (dark blue line). In the
organic plantation (O) these parameters become (3.2, 1.56) for the Collect state t; and (1.14, 29) for
the Search state . See Table 1 and equations (2)-(3).

*3.2. Harvested trees and visited area per day are higher in the organic plantation*

The average distance between trees within each *pante* is only slightly larger in the organic plantation
(C: 3.8 ± 2.11 m; O: 4.8 ± 2.6 m; not significant). In this sense, the marked differences in the PDFs
distributions of Search and Collect states between both plantations are not completely determined by
the planting pattern. The average number of trees visited per unit of time were higher in the organic
than in the conventional plantation but the differences were not significant (C: 85 (± 19) trees per day,
O: 104 (± 29) trees per day; C: 25 (± 7) trees per hour; O: 29 (± 5) trees per hour). The higher number
of harvested trees did not seem to correspond to differences in harvest weight per worker, which
implies that, in the organic plantation, each tree had less mature berries when the data was taken.

The average traveled area per worker (approximated as the number of visited squares of given
side length) was significantly higher in the organic plantation, for square units of 3, 5, 10 and 20 m
side (Fig. 3). When the unit area is too small (1 $m^2$) or when the unit is too large (1600 $m^2$ or 40 m
side) no significant differences are observed. The higher visited surface area might be the
consequence of a higher presence of long steps (or relocalizations) during the harvesting day (Fig. 1
and Fig. 2). In some specific trajectories harvesters traveled up to 100 meters to reach the following
tree.

**Figure 3. Surface area visited per harvester, obtained from visited squares of different sizes.** Each dot represents the number of visited squares per harvester in each of the plantations (C-gray: conventional, O-white: organic). For the square unit areas where the differences between the two plantations were significant we added a * (p<0.05) and the used test of comparison between the two means depending on the normality of the residuals (st: Student, w: Wilcoxon, n= 6).

*3.3. Differences in trajectories of the harvesters are mediated by climate, the coffee biology and plantation variables, and most of them are modulated by management decisions.*

The spatial movement of the harvesters, in particular the presence of longer steps (mainly in the state Search), as well as the daily length of the trajectory, can be directly and indirectly mediated, at least, by climate, topography, coffee biology characteristics, and coffee management variables (Fig. 4; Table 2). The relationship between these variables give us some insights on the determinants of the syndrome of production and the reasons that might explain the differences and commonalities in the trajectory between both plantations.

According to the interviews and our additional observations (solid and dotted lines in Fig. 4, respectively), the harvester's trajectory might be directly affected by the number of mature berries in each of the trees. This number is determined by the fruit load (total number of berries per tree) and the proportion of mature berries per tree. Here we use "fruit load" to refer to the number of mature berries per tree (D2). When trees have more berries, harvesters visit fewer trees and *pantes* per day, and their final trajectory is shorter (Table 2). Besides, to maximize their daily harvest, harvesters keep the same row when all the trees have berries, or jump from one row to another or even to another *pante*, when

some trees are still immature. We refer to this as the interplant ripening percentage (i.e. the percentage

of trees with ripe berries, D1). Low ripening percentage results in more randomized patterns or larger

step lengths along the day (state Search) (Fig.1). Ripening percentage and fruit load depend on the

coffee variety and age (N3, N5) and the date of harvest (N2, N4). Around the middle of the season,

each coffee variety has a peak of berry maturation and synchronization between the trees. When we

collected our data, trees were more homogeneously loaded in the conventional plantation than in the

organic plantation, explaining in part the higher number of harvested trees in the organic plantation

and the presence of larger step-lengths in the state Search (Fig. 1). This pattern could have been

reversed in another period. It is worth mentioning that coffee ripening is mostly mediated by the

rainfall regularity during the flowering season (N1) and the coffee variety (N5).

On the other hand, the spatial movement of harvesters is built upon the initial planting pattern

(in rows, in groups, etc) and the *pante* size and shape (D5, D7). Both processes depend on the type of

plantation (sun or shade-grown coffee) (N8, N11), the coffee variety (N7), and the topography of the

field (N10, N11). Plant renovation dynamics also affect the planting pattern (N8). In the organic

plantation individual plant renovations are more common, as well as the presence of other trees apart

from coffee. This also defines more irregular patterns than in the conventional plantation without

shade trees, where whole locations are removed and renovated every few years to increase short-term

productivity. This reflects on the different degrees of randomization of the trajectories of the

harvesters during harvest (Fig.1).

The number of harvesters (D3) per *pante* and the *pante* size (D7) could also determine the

general pattern as it affects how fast families can harvest one location during the day and travel to

another pante (long-steps in Fig. 1). The number of harvesters changes across the harvesting season

following the coffee maturation process (N6). Finally, the number of working hours (D6) or the

difficulty of harvesting a specific coffee variety (D4) also affect the trajectory length.

Most of these variables can be directly or indirectly affected by management decisions and

ultimately depend on the specific syndrome of production of both plantations (green boxes in Fig. 4).

For example, owners specify the starting and ending date of the harvesting season. Owners also

decide, mainly with profit-maximizing criteria, on the coffee varieties, the shade management, or the

dynamics of plant renovation. As we saw, these factors will indirectly affect the final trajectory of the

harvesters. Management can also determine the motion of harvesters by changing the working hours

or the number of people by *pante*, or the distance between *pantes*. In this sense, the recorded

trajectories of both plantations, their lengths, and patterns are the result of two different shade and

management practices within a specific syndrome of production.

**Figure 4. Qualitative drivers of the spatial trajectories of the harvesters during harvesting**. The

nodes represent the variables that interplay with the trajectory and the edges the causal relationships

between them. The color of the boxes indicates if the variable is environmental or coffee biology

related (gray), and if it can be directly modified by the management of the plantation (green). Note

that the modification of these variables affects all the variables downstream. The letter in the label of

the edges indicates whether the effect on the trajectory is direct (D) or indirect (N). The explanation of

each label is summarized in Table 2.

**Table 2**- Direct (D) and indirect (N) drivers of the trajectories of the harvesters. The relation between

the nodes is shown in Fig. 4.

| Variable | Affected by | Description | Edge | Source |
|---|---|---|---|---|
| | | **Direct effects** | | |
| **Trajectory** | Interplant ripening percentage | If the majority of plants have red berries, the harvester stays in the same row. When some of them have immature berries, he jumps from one row to another to maximize harvest | D1 | Harvester |
| | Fruit load | With a higher fruit load per tree, harvesters visit less trees and *pantes* per day | D2 | Harvester |
| | Number of harvesters | With multiple harvesters on the field, *pantes* are finished more rapidly. Individual dynamics are randomized and move away from the planting pattern. | D3 | Pers. Obs |
| | Coffee variety and age | Some varieties are harder to harvest making harvesters visit less trees and *pantes* per day | D4 | Harvester |
| | Planting pattern | The trajectory is built upon the initial planting pattern | D5 | Pers. Obs. |
| | Working hours | More working hours implies more visited trees and *pantes* per day | D6 | Pers. Obs. |
| | *Pante'*s size and shape | The shape of the *pante* defines how often a harvester "turns" or keeps straight. Smaller *pantes* are harvested more rapidly | D7 | Harvester |
| | | **Indirect effects** | | |
| **Interplant ripening percentage** | Rainfall regularity | Ripening is determined by the temporal and spatial regularity of rainfall during flowering | N1 | Pers. Obs. |
| | Date of the harvest | Ripening is variable across the harvesting season | N2 | Harvester |
| | Coffee variety and age | Some varieties present a more uniform ripening than others | N3 | Harvester |
| **Fruit load** | Date of harvest | Fruit load is variable across the harvesting season | N4 | Harvester |
| | Coffee variety and age | Some varieties present more berries than other | N5 | Harvester |
| **Number of harvesters** | Date of harvest | At the beginning or end of the harvesting season, there are less harvesters in the field | N6 | Harvester |
| **Planting** | Coffee variety and age | Some varieties are sow closer to each other modifying the general pattern | N7 | Pers. Obs. |

| | | | | |
|---|---|---|---|---|
| **pattern** | Plant renovation | The plant renovation dynamics randomizes (when individual trees are removed) or uniforms (when complete locations are removed) the general pattern along the years. | N8 | Pers. Obs. |
| | Sun or Shade coffee | Polycultures have a higher distance between trees than monocultures and tend to be more heterogeneous | N9 | Pers. Obs. |
| | Topography of the field | Higher variability in the topography within a plot creates non-uniform planting patterns | N10 | Pers. Obs. |
| *Pante* **size and shape** | Sun or Shade coffee | The presence of trees in polycultures defines less uniform *pantes* | N11 | Harvester |
| | Topography of the field | Topography conditions the boundaries of the *pante* | N12 | Pers. Obs. |
| **Plant renovation** | Coffee variety and age | The rate of plant renovation depends on the age of the plantation and the productive years of each variety | N13 | Pers. Obs. |

## 4. DISCUSSION AND CONCLUSIONS

Coffee is produced under a large variety of management practices, land tenure systems, and scales. In Mexico, one finds community-owned rustic coffee plantations as well as large-scale landlord-owned sun, and sometimes shaded, coffee plantations (Moguel and Toledo, 1999). Here we draw upon the conceptual framework of the "syndromes of production" to evaluate some of the social and ecological dynamics within a given agricultural paradigm (Andow and Hidaka, 1989; Ong and Liao, 2020; Vandermeer and Perfecto, 2012). In particular, we maintain that the two studied plantations (organic and conventional) in the Sonocusco region are part of a general capitalist syndrome of production characterized by large-scale plantations where landlords extract surplus value from the work of permanent and temporary workers and sell their product to conventional or organic markets (López Echeverría, 2006; Ong and Liao, 2020). The distinction in their markets reflects differences in their specific management and ecological practices. The organic plantation has an integrated management system that results in higher associated biodiversity, pest control, and lower plant renovation, but also in better worker's health conditions by removing pesticides (Soto-Pinto et al., 2000; Jiménez-Soto, 2021). Besides, it values quality over quantity in coffee production, which reduces the plant renovation rate and prioritizes planting high-quality arabica plants.

Our study aimed to understand how distinct agricultural managements in large-scale capitalist plantations drive differences in patterns of human movement that can potentially impact CLR spread in the environment, as well as working conditions. We found that two-state space models can describe harvester's trajectories well. This is in line with our observations in the field where we differentiate two kinds of movements: when trees have berries, harvesters remain in the same rows or nearby (Collect state: short to medium steps); when no berries are in sight or when they have finished a *pante*, harvesters move to another zone of the *pante* or to another *pante* chosen by the foreman (Search state: medium to long steps). These behaviors are similar to general foraging patterns in other agroecosystems (Reynolds et al., 2018) or to the trajectories generated by deterministic models where a walker visits the closest unvisited site among a collection of point-like sites irregularly distributed in a landscape (Santos et al., 2007). The organic shaded plantation had longer steps both in the Collect state (within the *pante*) and specially in the Search state (relocalizations between *pantes*) in the same amount of time and, according to our qualitative analysis this could be related to *a)* a lower fruit load per tree or low percentage of trees with berries when we took the data or *b)* smaller *pantes* or the number of harvesters per *pante*. This resulted in a higher number of visited areas of different sizes (Fig. 3). The heterogeneity in trajectories in this plantation seems to be related to the presence of other trees, plant renovation dynamics, or even to the topographic context, as also discussed by Hajian-Forooshani and Vandermeer, (2021) and Soto-Pinto et al., (2000).

In this sense, the spatial movements of the harvesters are the result of their interactions with
biological and climatic factors and also depend on plantation characteristics and management
decisions. This goes along the line of the unifying paradigm of movement ecology first introduced in
the context of animals (Nathan et al 2008). For example, ripening can be affected by the regularity of
the rainfall during flowering and the temperature (Kath et al., 2021), or by the distribution of shade
(Henrique Marquezine Leite et al., 2022). Nevertheless, the owners decide when to end the harvesting
season. As was reported by other studies, it is common for the owners to encourage harvesters to stay
until the end of the season by increasing the value of the harvest load, or to directly force them to
remain in the fields by delaying their payment (Jiménez-Soto, 2021). This promotes harvesting when
plants have a low percentage of trees with ripe fruits and low fruit load per tree, setting the scenario
for longer daily trajectories with a higher proportion of medium to long steps across the region.

There are several potential socio-ecological implications of the different harvesters'
trajectories on coffee rust dispersal. Here we did not report any data related to CLR or other potential
pests but we can hypothesize that visiting more trees per day, without the possibility of changing
clothing could increase the dispersal and impact of CLR (as observed by Becker and Kranz (1977)).
Besides, CLR or other pathogens are also dispersed from plant to plant through water splash or leaf-
to-leaf contact, creating localized foci of infection (Vandermeer et al., 2018). Changing *pante* several
times (Fig. 1, organic plantation) could connect these foci of infection or bear the pathogens to
different parts of the landscape and start new infections. The square units depicted in Fig. 3 could be
thought of as a proxy of these foci. In this sense, the spatial heterogeneity in coffee maturation creates
fragmented host-landscapes that interact with the movement of harvesters and could increase the CLR
or other pathogens' outbreaks (as suggested by White et al., 2018). Nonetheless, these hypotheses
need to be tested, theoretically or empirically. These implications would not mean promoting
managements that could homogenize the coffee ripening or fruit charge (e.g., homogenous sun coffee
varieties, controlled irrigations) to reduce the spatial movement of harvesters during the harvesting.
Coffee production and the incidence of specific pathogens as CLR are multifactorial processes
(Avelino et al., 2006) and decisions that could reduce human dispersal can promote another
dimension of the epidemic. For example, removing the shade trees to avoid irregular ripening
(Henrique Marquezine Leite et al., 2022) also increases the CLR wind-mediated dispersal (Gagliardi
et al., 2020). They also do not mean placing responsibility on the harvesters as they are temporary or
permanent workers paid daily by piecework, working precariously for several hours to maximize the
daily harvest in landlord-owned plantations guided by the generation of profit.

Our findings call for organic coffee management that could prevent diseases, increase
diversity, and guarantee just and safe conditions for workers. In particular, to reduce the possible
impact of human dispersal of pathogens during harvesting, we suggest, among other ideas, the need
for shorter trajectories by working fewer hours a day, avoiding harvesting at the beginning or end of
the coffee maturation season when few trees have berries and harvesters have to travel longer

distances, or even skipping infected plants. These measures can reduce short-term productivity but
sustain long-term productivity. In this sense, to be implemented, coffee farmers should overcome the
business-as-usual politics of large-scale capitalist coffee agroecosystems.
476
477 **5. ACKNOWLEDGEMENTS**
478

EMVC is a doctoral student from the Programa de Doctorado en Ciencias Biomédicas, Universidad
Nacional Autónoma de México, and has received CONACyT scholarship 686776. MB acknowledges
financial support from UNAM-DGAPA-PAPIIT (IN207819). EMVC, CGG and MB thanks Gustavo
Bautista, Gabriel Domínguez and Elisa Lotero for the discussions and precious help during the
fieldwork. EMVC especially thanks Elisa Lotero for her patience and support and the people from La
Parcela for their ideas and friendship. Finally the authors dedicate this work to all the harvester
families in both plantations and in other regions, that feed the world under social conditions that must
be overturned .

487
488

490 **6. REFERENCES**

492 Andow, D., 1983. The extent of monoculture and its effects on insect pest populations with particular
493  reference to wheat and cotton. Agriculture, Ecosystems and Environment 9, 25–35.
494  https://doi.org/10.1016/0167-8809(83)90003-8

495 Andow, D.A., Hidaka, K., 1989. Experimental natural history of sustainable agriculture: syndromes of
496  production. Agriculture, Ecosystems & Environment 27, 447–462.
497  https://doi.org/10.1016/0167-8809(89)90105-9

498 Avelino, J., Cristancho, M., Georgiou, S., Imbach, P., Aguilar, L., Bornemann, G., Läderach, P.,
499  Anzueto, F., Hruska, A.J., Morales, C., 2015. The coffee rust crises in Colombia and Central
500  America (2008–2013): impacts, plausible causes and proposed solutions. Food Security 7,
501  303–321. https://doi.org/10.1007/s12571-015-0446-9

502 Avelino, J., Zelaya, H., Merlo, A., Pineda, A., Ordoñez, M., Savary, S., 2006. The intensity of a
503  coffee rust epidemic is dependent on production situations. Ecological Modelling 197, 431–
504  447. https://doi.org/10.1016/j.ecolmodel.2006.03.013

505 Becker, S., Kranz, J., 1977. Comparative Studies on Dispersal of Hemileia vastratrix in Kenya.
506  Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz / Journal of Plant Diseases and
507  Protection 84, 526–539. https://www.jstor.org/stable/ 43213550

508  Breed, G. A., & Severns, P. M. (2015). Low relative error in consumer-grade GPS units make
509  them ideal for measuring small-scale animal movement patterns. *PeerJ*, *3*, e1205.
510  https://doi.org/10.7717/peerj.1205

511 Evans, J., Jones, P., 2011. The walking interview: Methodology, mobility and place. Applied
512  Geography 31, 849–858. https://doi.org/10.1016/j.apgeog.2010.09.005

513 Gagliardi, S., Avelino, J., Beilhe, L.B., Isaac, M.E., 2020. Contribution of shade trees to wind
514  dynamics and pathogen dispersal on the edge of coffee agroforestry systems: A functional
515  traits approach. Crop Protection 130, 105071. https://doi.org/10.1016/j.cropro.2019.105071

516 Hajian-Forooshani, Z., Rivera Salinas, I.S., Jiménez-Soto, E., Perfecto, I., Vandermeer, J., 2016.
517  Impact of Regionally Distinct Agroecosystem Communities on the Potential for Autonomous
518  Control of the Coffee Leaf Rust. Environ Entomol 45, 1521–1526.
519  https://doi.org/10.1093/ee/nvw125

520 Hajian-Forooshani, Z., Vandermeer, J., 2021. Emergent spatial structure and pathogen epidemics: the
521  influence of management and stochasticity in agroecosystems. Ecological Complexity 45,
522  100872. https://doi.org/10.1016/j.ecocom.2020.100872

523 Henrique Marquezine Leite, P., Evangelista De Oliveira, R., Durrer Bigaton, A., Vilela, G.,
524  Fontanetti, A., 2022. Shade influence of the temporary canopy of an agroforestry system on
525  coffee fruit ripening. CS 16, 1–11. https://doi.org/10.25186/.v16i.1974

Jackson, D., Skillman, J., Vandermeer, J., 2012. Indirect biological control of the coffee leaf rust,
Hemileia vastatrix, by the entomogenous fungus Lecanicillium lecanii in a complex coffee
agroecosystem.        Biological        Control        61,        89–97.
https://doi.org/10.1016/j.biocontrol.2012.01.004

Jiménez-Soto, E., 2021. The political ecology of shaded coffee plantations: conservation narratives
and the everyday-lived-experience of farmworkers. Journal of Peasant Studies 48, 1284–
1303. https://doi.org/10.1080/03066150.2020.1713109

Jones, B.A., Grace, D., Kock, R., Alonso, S., Rushton, J., Said, M.Y., McKeever, D., Mutua, F.,
Young, J., McDermott, J., Pfeiffer, D.U., 2013. Zoonosis emergence linked to agricultural
intensification and environmental change. Proc. Natl. Acad. Sci. U.S.A. 110, 8399–8404.
https://doi.org/10.1073/pnas.1208059110

Kath, J., Mittahalli Byrareddy, V., Mushtaq, S., Craparo, A., Porcel, M., 2021. Temperature and
rainfall impacts on robusta coffee bean characteristics. Climate Risk Management 32,
100281. https://doi.org/10.1016/j.crm.2021.100281

López Echeverría, M.E., 2006. Identidad, autonomía y cultura: El espíritu del capitalismo en las
Fincas      Cafetaleras      Alemanas      en      el      Soconusco      (1850-2006).
http://tesiuami.izt.uam.mx/uam/aspuam/presentatesis.php?
recno=13595&docs=UAMI13595.pdf

Manlove, K., Wilber, M., White, L., Bastille-Rousseau, G., Yang, A., Gilbertson, M. L., ... &
Pepin, K. M. (2022). Defining an epidemiological landscape that connects movement ecology
to pathogen transmission and pace-of-life. *Ecology Letters, 25*(8), 1760-1782.
https://doi.org/10.1111/ele.14032

Mapes, K., 2010. Sweet tyranny: Migrant labor, industrial agriculture, and imperial politics.
University of Illinois Press.

Mas, A.H., Dietsch, T.V., 2004. Linking Shade Coffee Certification to Biodiversity Conservation:
Butterflies and Birds in Chiapas, Mexico. Ecological Applications 14, 642–654.
https://doi.org/10.1890/02-5225

McCook, S., Vandermeer, J., 2015. The Big Rust and the Red Queen: Long-term perspectives on
coffee rust research. Phytopathology 105, 1164–1173. https://doi.org/10.1094/PHYTO-04-15-
0085-RVW

Michelot, T., Blackwell, P.G., 2019. State-switching continuous-time correlated random walks.
Methods Ecol Evol 10, 637–649. https://doi.org/10.1111/2041-210X.13154

Michelot, T., Langrock, R., Patterson, T.A., 2016. moveHMM: an R package for the statistical
modelling of animal movement data using hidden Markov models. Methods Ecol Evol 7,
1308–1315. https://doi.org/10.1111/2041-210X.12578

Moguel, P., Toledo, V.M., 1999. Biodiversity conservation in traditional coffee systems of Mexico.
Conservation Biology 13, 11–21. https://doi.org/10.1046/j.1523-1739.1999.97153.x

Mora Van Cauwelaert, E., González González, C., Boyer, D., Hajian-Forooshani, Z., Vandermeer, J.,
Benítez, M., 2023. Dispersal and plant arrangement condition the timing and magnitude of
coffee rust infection. Ecological Modelling 475, 110206.
https://doi.org/10.1016/j.ecolmodel.2022.110206

Morales, J.M., Haydon, D.T., Frair, J., Holsinger, K.E., Fryxell, J.M., 2004. Extracting More Out of
Relocation Data: Building Movement Models as Mixtures of Random Walks. Ecology 85,
2436–2445. https://doi.org/10.1890/03-0269

Nathan, R., Getz, W. M., Revilla, E., Holyoak, M., Kadmon, R., Saltz, D., & Smouse, P. E.
(2008). A movement ecology paradigm for unifying organismal movement research.
Proceedings of the National Academy of Sciences, 105(49), 19052-19059.
https://doi.org/10.1073/pnas.0800375105

Ong, T.W.Y., Liao, W., 2020. Agroecological Transitions: A Mathematical Perspective on a
Transdisciplinary Problem. Front. Sustain. Food Syst. 4, 91.
https://doi.org/10.3389/fsufs.2020.00091

Patterson, T. A., Thomas, L., Wilcox, C., Ovaskainen, O., & Matthiopoulos, J. (2008). State–
space models of individual animal movement. Trends in ecology & evolution, 23(2), 87-94.
https://doi.org/10.1016/j.tree.2007.10.009

Perfecto, I., Chaves, L.F., Fitch, G.M., Hajian-Forooshani, Z., Iuliano, B., Li, K., Medina, N., Morris,
J., Otero-Jiménez, N., Rivera-Salinas, I.S., Chenyang, S., Vandermeer, J., White, A., and
Williams-Guillén, K. 2023. Looking beyond land-use and land-cover change: Zoonoses
emerge in the agricultural matrix. One Earth, 6(9), 1131-1142.
https://doi.org/10.1016/j.oneear.2023.08.010

Perfecto, I., Jiménez-Soto, M. E., & Vandermeer, J. 2019. Coffee landscapes shaping the
Anthropocene: forced simplification on a complex agroecological landscape. Current
Anthropology, 60(S20), S236-S250. https://doi.org/10.1086/703413

Pishro-Nik, H. (2014). Introduction to probability, statistics, and random processes (p. 732). Blue
Bell, PA, USA: Kappa Research, LLC

Raven, P. H., & Wagner, D. L. (2021). Agricultural intensification and climate change are
rapidly decreasing insect biodiversity. Proceedings of the National Academy of Sciences,
118(2), e2002548117.https://doi.org/10.1073/pnas.2002548117

Reynolds, A., Ceccon, E., Baldauf, C., Medeiros, T.K., Miramontes, O., 2018. Lévy foraging patterns
of rural humans. PLoS ONE 13, 1–16. https://doi.org/10.1371/journal.pone.0199099

Santos, M.C., Boyer, D., Miramontes, O., Viswanathan, G.M., Raposo, E.P., Mateos, J.L., Da Luz,
M.G.E., 2007. Origin of power-law distributions in deterministic walks: The influence of
landscape geometry. Phys. Rev. E 75, 061114. https://doi.org/10.1103/PhysRevE.75.061114

Soto-Pinto, L., Perfecto, I., Castillo-Hernandez, J., Caballero-Nieto, J., 2000. Shade effect on coffee
production at the northern Tzeltal Zone of the state of Chiapas, Mexico. Agriculture,

Ecosystems and Environment 80, 61–69. https://doi.org/10.1016/S0167-8809(00)00134-1

Vandermeer, J., Hajian-Forooshani, Z., Perfecto, I., 2018. The dynamics of the coffee rust disease: an

epidemiological approach using network theory. European Journal of Plant Pathology 150,

1001–1010. https://doi.org/10.1007/s10658-017-1339-x

Vandermeer, J., Perfecto, I., 2012. Syndromes of production in agriculture: Prospects for social-

ecological regime change. Ecology and Society 17. https://doi.org/10.5751/ES-04813-170439

Villarreal-Treviño, C., Penilla-Navarro, R.P., Vázquez-Martínez, M.G., Moo-Llanes, D.A., Ríos-

Delgado, J.C., Fernández-Salas, I., Rodríguez, A.D., 2015. Larval habitat characterization of

Anopheles darlingi from its northernmost geographical distribution in Chiapas, Mexico.

Malar J 14, 517. https://doi.org/10.1186/s12936-015-1037-0

Waller, J.M., 1981. The recent spread of some tropical plant diseases. Tropical Pest Management 27,

360–362. https://doi.org/10.1080/09670878109413805

White, L. A., Forester, J. D., & Craft, M. E. (2018). Disease outbreak thresholds emerge from

interactions between movement behavior, landscape structure, and epidemiology.

*Proceedings of the National Academy of Sciences*, *115*(28), 7374-7379.

https://doi.org/10.1073/pnas.1801383115

Zucchini, W., MacDonald, I.L., Langrock, R., 2016. Hidden markov models for time series: an

introduction using R, Second edition. ed, Monographs on statistics and applied probability.

CRC Press, Taylor & Francis Group, Boca Raton.