Interplay between harvesting, planting density and ripening time affects

coffee leaf rust dispersal and infection

Emilio Mora Van Cauwelaert[a, b, *], Kevin Li[c], Zachary Hajian-Forooshani[d], John Vandermeer[e]

and Mariana Benítez[a, *]

a.   Laboratorio Nacional de Ciencias de la Sostenibilidad, Instituto de Ecología, Universidad Nacional

Autónoma de México, Mexico City, C.P. 04510, México

b. Posgrado en Ciencias Biomédicas, Universidad Nacional Autónoma de México, Mexico City, C.P. 04510,

México

c. School of Environment and Sustainability, University of Michigan, Ann Arbor, MI 48108, USA

d. German Centre for Integrative Biodiversity Research (iDiv), Martin Luther University Halle, Leipzig,

Germany

e.   Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, Michigan, U.S.A

*mbenitez@iecologia.unam.mx, *emiliomora92@ciencias.unam.mx

ABSTRACT

The relationship between the spatial structure of hosts and the individual movement of

vectors is central to developing a comprehensive representation of pathogen dynamics. Here

we focus on the dynamics of coffee leaf rust (*CLR; Hemileia vastatrix*), which is one of the

principal coffee diseases around the globe. This fungus disperses between plants or regions

through direct contact, water splash, or local turbulent wind conditions. Some studies have

proposed that coffee harvesters can also bear CLR during harvesting. However, it is not

clear how relevant these movements are for CLR dispersal and plot-level infection in plots

with different planting densities. Besides, there are bidirectional interactions between the

plantation characteristics (e.g. planting density and ripening synchronicity) and the

movement of harvesters that might modify their dispersal potential. Overall, there is a lack

of mechanistic models to represent these interactions. Here we present a computational

model to explore the role of coffee ripening synchronicity in reproducing the different

movement trajectories observed in the field. We then evaluate how these harvesting

trajectories modify rust dispersal and change CLR infection in plots with increasing planting

densities and a constant contact-mediated rust dispersal mechanism. For both

synchronization scenarios, the distribution of step lengths is positively skewed. Interplant

asynchronous ripening and scarcity of trees with berries generate trajectories with medium

to long steps contrary to scenarios with synchronous ripening and without scarcity. The

harvest dispersal significantly increases coffee plot infection (up to 15%) compared to

scenarios where only local mechanisms for CLR dispersal are present. This effect is maximal

for trajectories with medium to long steps from the asynchronous scenario and in

plantations with medium planting densities. In particular, medium-sized steps are more

likely to create new foci and networks of contact-mediated infected plants. The interaction

between the proportion and the size of infected networks results in a nonlinear behavior of

rust increase due to harvest. Our results aim to spur discussion on practices that could

reduce the impact of harvesting and specific trajectories, in scenarios where one can benefit

from asynchronous maturation of berries and shaded plantations. Some examples of

practices include reducing the medium to long-distance movements during the same day,

for instance, by avoiding harvesting at the end of the harvest season when trees ripening

asynchronicity and CLR levels are higher.

KEYWORDS

Coffee leaf rust dispersal; harvesting trajectories; interplant ripening; mechanistic modeling;

planting density

1. INTRODUCTION

The life cycle of pathogens in plants can be divided broadly into two stages: the infection stage where the organism invades the host and reproduces, and the dispersal stage where new replicates of the pathogen move from one host to the other (Avelino et al., 2004; Pangga et al., 2011). The rate and extent at which both stages are completed largely determine the magnitude of the epidemic and the crop loss (Avelino et al., 2004). Contrary to epidemics in non-sessile organisms, the host's density and spatial arrangement play a central role in plants, as they can modify the range and dynamics of the pathogen's dispersal (Keeling, 1999; Vandermeer et al., 2018; Papaïx et al.; 2015). In agriculture, this spatial structure depends on management factors (Beasley et al., 2022), ecological dynamics (Li et al., 2016), and also on the topography of the field, among others (Soto-Pinto et al., 2000). Additionally, the interaction of the arrangement of the hosts and the disease depends on the characteristic transmission's distance of dispersal (Park et al., 2001; Hajian-Forooshani and Vandermeer, 2021). Few studies compare the effect of different mechanisms of pathogen transmission across spatially explicit landscapes (White et al., 2018). Studying these spatial relationships is crucial for an ecological approach to pest management that does not rely on synthetic pesticides, or solely on resistant plant varieties that focus on the infection stage (Avelino et al; 2004).

Coffee leaf rust (CLR), caused by *Hemileia vastatrix* Berk. & Br is one of the principal coffee diseases and has caused devastating loss of production in different coffee sites around the globe (McCook and Vandermeer, 2015; Talhinhas et al., 2017). Its mechanisms for dispersal have been amply studied (Becker and Kranz, 1977; Kushalappa and Eskes, 1989; Li et al., 2023; Vandermeer et al., 2018) and thus provide a great model to explore the disease's spatially explicit dynamics and its relations with the host's spatial arrangement and density (Beasley et al., 2022; Gagliardi et al., 2020; Hajian-Forooshani and Vandermeer, 2021; Mora Van Cauwelaert et al., 2023a). CLR forms urediniospores that can travel to neighboring

coffee plants through direct contact, water splash, or local turbulent wind conditions (Kushalappa and Eskes, 1989). The urediniospores may also ascend to the atmosphere and travel to other plantations (Becker and Kranz, 1977; Vandermeer and Rohani, 2014). For each of these ways of dispersal, there are agroecological management proposals to reduce the spread, like reducing the density or clustering of the plantations (Beasley et al., 2022; Ehrenbergerová et al., 2018), avoiding regular or extremely dense patterns that can make the epidemic thrive through contact or splash-mediated dispersal (Hajian-Forooshani and Vandermeer, 2021; Mora Van Cauwelaert et al., 2023a; Li et al., 2023), but also planting different trees within the plots and in the borders to reduce local and regional wind dispersal (Avelino et al., 2022; Boudrot et al., 2016; Gagliardi et al., 2020)

It has been observed that workers can also disperse CLR within the plot and between regions (Becker and Kranz, 1977; Ramírez-Camejo et al., 2022; Schieber and Zentmyer, 1984). At the plot scale, this relates to the fact that some foci of infection occur close to harvesting paths and human settlements (Waller, 1982, 1972) and that the build-up of the epidemic correlates with the harvesting season (Avelino et al., 1991; Mora Van Cauwelaert et al., 2023a). However, it is unclear how relevant the dispersal by harvesters is for the whole epidemic in contrast to other modes of dispersal, and what agroecological measures could be implemented to reduce its impact.

Besides, the dynamics of harvesting in coffee plantations depend on the scale, on specific ecological and management characteristics, but also on land tenure regimes and goals of production; or on what has been called "the syndrome of production" (Andow and Hidaka, 1989; Perfecto et al., 2019). In large-scale and landlord-owned plantations, the trajectory of paid workers during harvest relies, among other factors, on the ripening synchronization of the berries between the trees in the plot (Mora Van Cauwelaert et al., 2023b). Distinct trajectories, with short or long steps between plants, could play a differential role in CLR dispersal and its impact on plantations (Mora Van Cauwelaert et al., 2023b).

Here we present a mathematical model to study the workers' spatial movement

during coffee harvesting and its contribution to CLR dispersal and infection at a plot level.

We first explore how the spatial trajectories drawn by the workers in large-scale plantations

(Mora Van Cauwelaert et al., 2023b) could be reproduced in simulated coffee plots with

different berries maturation synchronicity. We then evaluate how these trajectories modify

rust dispersal and affect the final CLR infection in plots with increasing planting densities.

Finally, we discuss the implications of our results in a broader context of different

syndromes of coffee production.

2. METHODS

The objective of implementing a computational modeling approach was twofold: i) test if the

temporal differences in ripening between the trees are sufficient to reproduce the reported

trajectories of the workers during harvest (as previously speculated by Mora Van

Cauwelaert et al., 2023b), and ii) explore the effect of these trajectories for dispersal and

average rust infection in coffee plots. We first simulated coffee plots with different planting

densities, interplant ripening scenarios and a closest-neighbor harvesting rule to study the

role of ripening on harvesting trajectories. Then, we defined a harvesting-based mechanism

of rust dispersal to evaluate the impact of harvest in the average coffee plot rust infection

alongside a contact-mediated rust dispersal mechanism. In the following lines, we describe

the specificities of the model (Fig. 1), the simulated scenarios, and the analyses made for

each objective.

2.1 Model Description

We represent one hectare coffee plots as 100 x 100 unit squares with non-periodic

boundaries, with N randomly placed points representing the plants (randomized

arrangements are observed particularly in old plantations; Hajian-Forooshani and

Vandermeer, 2021). As we fix the area, N represents the average planting density per ha.

Plants within the simulated plantation are assumed to be homogeneous, except for two

variables: the ripening and the infection status.

Coffee ripening and harvesting motion. Each plant can have coffee berries (*present*)

or not (*absent*) (Fig.1A). In order to explore the role of ripening on harvesting dynamics, we

defined two coffee interplant ripening scenarios. In the asynchronous scenario (**A**), 50%

randomly selected plants have ripe berries at the date of harvest. In the synchronous

scenario (**S**), all the plants have ripe berries at harvest (Fig. 1A). The presence of berries

defines the harvesting motion. Each worker starts harvesting in a plant with berries, collects

them (changing the status of berries of that plant to *absent*), and moves to the closest plant

with berries (Fig.1A). This cycle repeats until half of the plants in the plots are visited. In the

asynchronous scenario, this leads to a plot without any berries left, and in the synchronous

scenario half of the plants will still have berries (Fig. 1A). According to previous studies one

worker collects around 30 plants per hour, or roughly 250 plants in eight hours (i.e. one

harvesting day; Mora Van Cauwelaert et al., 2023b). In the following simulations, we adjust

the number of workers in the plots to always represent one harvesting day with 250

harvesting steps per worker (e.g. if we have 2000 plants, we will need four workers to visit

half of the plants in a day).

Rust dispersal and infection. Each plant has one of two states of infection:

susceptible or *infected*. In our model we used *infected* to refer to plants that are infected and

infective (with rust spores present). When an infected plant infects susceptible neighbors at

time step t, these become infective at $t+1$. Each unit represents roughly 30 days (i.e. one

month), this is, the time needed for rust to complete a cycle of infection (Leguizamon et al.,

1998). An infected plant can disperse and infect the neighboring plants by two mechanisms

(Fig. 1B). Firstly, they disperse to all 1.5 m apart or less susceptible neighbors through

contact-mediated dispersal. This mimics the local dispersion of rust either by contact

between two nearby plants or by splash (as proposed by Vandermeer et al., 2018). The

process creates networks of infected nearby plants where each pair of plants are 1.5 m apart

or less (gray circles in Fig. 1B). Secondly, when a worker harvests an infected plant and

reaches a susceptible plant in the following harvesting step, it will become infected (yellow

circle in Fig.1B). We call this the *worker-mediated dispersal*. Both mechanisms are assumed

to be instantaneous in relation to time step t.

Whole infection dynamics. To analyze the effect of harvesting in rust infection, we

simulate a whole season dynamic where each plot started with an initial percentage of

infected plants and got through twelve cycles of contact-mediated dispersal and infection

(control scenario, Fig. 1C). This represents the maximum rust spread during one year,

without any harvesting and without acknowledging for leaf loss or other mechanisms or

climate factors that in practice reduce the general infection (Kushalappa and Eskes, 1989). In

the harvesting scenarios, we add a punctual harvesting day event a T_H after five time steps (t

= 5). These five cycles mimic the five months from the beginning of CLR buildup to the peak

of harvesting (Mora Van Cauwelaert et al., 2023a). At the date of harvest (T_H), almost all the

plants in the same contact-mediated networks with one or more initially infected plants are

infected (gray circles in Fig. 1C). During harvest, new infections emerge (yellow circles in

Fig. 1C). After harvest, seven new cycles of contact-mediated dispersal and infection take

place in which the newly infected plants spread the rust across their contact-mediated

networks (black circles in Fig.1B).

2.2. Scenarios and measurements

We first characterized the trajectories and the distribution of steps simulated under two

scenarios of coffee ripening (A and S) and five planting densities (500, 1000, 2000, 3000, and

5000 plants per ha; Soto-Pinto et al., 2000), adding up to 10 scenarios. We replicated each

scenario 30 times to account for the random initial arrangement of plants. We compared the

trajectories drawn in plantations with 2000 plants/ha with published data of plantations

with equivalent densities and different levels of ripening synchronicity (Mora Van

Cauwelaert et al., 2023b). In particular, we took two examples from a large-scale organic

plantation with an asynchronous ripening at the date of harvest ("organic plantation",

henceforth) and two from a slightly shaded, large-scale non-organic plantation with

synchronous ripening at the date of harvest (conventional plantation). Both plantations had

a planting density between 2000 to 3000 plants/ha. Secondly, we used our model to explore

the effect of these simulated trajectories for each coffee ripening condition and planting

density, on the plot average rust infection. We ran each scenario 30 times for the whole

season (**t= 12**) starting with 20% of initially infected plants and compared them to non-

harvesting scenarios (five additional scenarios, one for each planting density; Fig.1C).

Finally, to further explore the mechanisms behind our results, we registered the networks of

nearby plants (where each pair of plants are 1.5 m apart or less) infected through the contact-

mediated dispersal mechanism before harvest (gray circles in Fig. 1B) and after the harvest

event (black circles in Fig. 1B). We calculated their proportion and their mean size in terms

of number of nodes and included the product of both magnitudes as a proxy for the number

of infected plants after the harvest event. We also recorded the newly infected plants per

step lengths during harvest (yellow circles in Fig. 1B).

The model and simulations were implemented in the *Python* 3.10.6 programming

language, using the modules *NumPy, SciPy, Pandas, and Seaborn*, and ran on the LANCIS

facilities, at the Ecology Institute of UNAM. The data analyses and figures were done in

Rstudio 2023.03.1+446 using *plyr, dplyr, tidyverse, ggplot2, patchwork* libraries, and *Inkscape 1.0.*

All code and data required to reproduce results in the work can be accessed at

https://github.com/tenayuco/toyModel_harvest.

Fig. 1. Model structure and dynamics. *A) Coffee ripening and harvesting motion.* We illustrate the two coffee ripening scenarios explained in the text. In the asynchronous scenario half of the plants in the plot have ripe berries (red circles) at the date of harvest, and in the synchronous scenario all the plants have ripe berries. The presence of berries determines the trajectory of the worker (black arrows) during harvest. *B. Dispersal and infection.* Plants can be susceptible (white circles) or can be infected before the harvest (gray circles), during the harvest (yellow circles) or after the harvest (black circles). Plants can be infected through contact-mediated dispersal when they are 1.5 m apart or less from an infected plant (gray circles), or through worker-mediated dispersal (yellow circle). *C) Whole model dynamic.* We depict the start of the simulation (T_0), the five cycles of contact dispersal and infection until the time of harvest (T_H) and the two possible scenarios afterwards: the control scenario where no harvesting takes place, and the harvest scenario where harvesters can create new infections (yellow circles) that might disperse to the neighbors after new seven cycles of contact-mediated dispersal and infection (black circles).

3. RESULTS

3.1. Interplant coffee ripening synchronization and berry scarcity produce two qualitatively different harvesting trajectories previously documented in large-scale plantations

The deterministic movement of the simulated workers across randomly distributed plants generates random trajectories (Fig. 2 and Fig.S1.1). For both synchronization scenarios, the distribution of step length is positively skewed and can be approximated with continuous lognormal and exponential distributions (Fig. 2, Fig.S1.2 and Fig. S1.3). Within each planting density, the mode is equivalent (Table S1.1). In the asynchronous ripening scenario with 2000 plants/ha, 27% of the steps are smaller than 1.5 m, 95% are below 9 m (short to medium steps), and there are some unlikely long steps (30 to 100 m) that produce a heavy-tailed distribution (Fig.2, first and second column). In the synchronous scenario, 51% of the steps are smaller than 1.5 m, 95% are below 4 m (short steps), and all steps are smaller than 30 m, creating a short-tailed distribution (Fig.2, first and second column). In the asynchronous scenario, the workers explore a larger area than in the synchronous scenario (Fig. 2, second column).

Interestingly, the presence of longer steps (>30 m) is not directly related to the initial low density of plants with berries in the asynchronous scenario. In fact they are not present in any of the equivalent and sometimes lower planting densities in the synchronous scenarios (Fig. S1.1). Long steps are concentrated near the end of the harvesting process for all planting densities, suggesting they are related to an increasing scarcity of trees with berries during the harvesting (Fig.2, third column and Fig. S1. 4).

Figure 2. Simulated harvesting trajectories under two scenarios of coffee ripening in plots with 2000 plants/ha. We present the step length distribution (first column), the XY paths (second column), and the step lengths along the percentage of trees with berries (third column). *The step length distribution* summarizes the results of 30 repetitions per scenario. The boxplot was included as an aid to visualize longer steps. The modes of the skewed distributions are 1.7 m and 1.3 m for the asynchronous and synchronous scenarios, and the means are 3.2 m (± 5.2) and 1.7 m (± 1.2), respectively. For the presented data, the number of workers is 4. *The XY path* shows the spatially explicit movement of the workers for one of the repetitions in each coffee ripening scenario. Each color represents one worker. In *the step length along the trajectory* we plot the length of each step during harvesting for each worker in relation to the increasing scarcity of trees with berries due to harvest. In the asynchronous scenario, no trees with berries remain at the end of harvest, in the synchronous, half of the trees will still have berries (Fig. 1A). The trajectories and step length distributions for other planting densities are presented in Fig. S1.1, Fig. S1.3, and Fig. S1.4).

The generated trajectories per worker at a planting density of 3000 plants/ha are

qualitatively similar to in-field trajectories drawn by workers in two large-scale coffee

plantations with equivalent planting density (3000 to 4000 plants/ha) but that differed in

their ecological management and their ripening synchronicity at the date of harvest (Fig. 3).

The modeled asynchronous trajectory is equivalent to the trajectories generated in an

organic plantation with an asynchronous interplant ripening at the date of harvest (O in Fig.

3, green line). The synchronous model is qualitatively similar to the trajectories generated in

a plot where ripening was more homogeneous (C in Fig. 3, blue line). Although other

processes and variables may affect trajectories, this result implies that the coffee ripening

differences and the scarcity of trees with berries at the end of harvesting in our model could

be *sufficient* to generate the different step lengths observed in the field. In our model, when

the ripening between plants is asynchronous, workers generate a higher number of medium

steps to reach plants with berries, and longer steps between coffee plots when a zone is

depleted (Fig. 3). Contrarily, synchronous ripening or a short harvesting process (before

reaching the scarcity of trees; Fig. 2) causes workers to stay in the same coffee rows and area

during the day (Fig.3).

Fig. 3. Comparison of the model with in-field trajectories. A. Two examples of the modeled
trajectories drawn by one worker under the asynchronous and the synchronous scenarios, with N=
3000 plants/ha. **B.** Examples of in-field registered trajectories from Mora Van Cauwelaert et al.
(2023b). The organic plantation (green line) is a large-scale shaded and organic plantation with an
asynchronous ripening at the date of harvest. The conventional plantation (blue line) is a slightly
shaded, non-organic large-scale plantation with synchronous ripening at the date of harvest. We
depict the number of visited plants for each trajectory (N) (see Mora Van Cauwelaert et al., 2023b for
more details). Both plantations are around 300 ha, with a planting density between 3000 to 4000
plants per ha.

3.2. Dispersal during harvesting increases final rust infection, especially in the asynchronous scenario and in plots with medium planting densities.

In this section we analyze the effect of the different generated simulated trajectories (with medium and long steps from the asynchronous ripening scenario, and short steps from the synchronous scenario) on the average rust in the plot.

Higher planting density increases the plot-average infection in scenarios where only contact-mediated dispersal is present (control; white line in Fig. 4A). Nonetheless, for each planting density, adding the harvesting event increases the final infection compared to the control, specially in the asynchronous ripening scenario (orange and blue lines in Fig. 4A). The difference between the average rust with harvesting and without harvesting follows a non-linear curve with increasing planting density for both kinds of harvesting scenarios (orange and blue boxes in Fig. 4B). The maximum effect of harvesting in the asynchronous scenario lies at 2000 plants/ha (~15% average rust increase; blue box in Fig. 4B), and between 1000 and 2000 plants/ha for trajectories in the synchronous scenario (~ 8% increase; orange box in Fig. 4B). At very low densities (500 plants/ha) the difference between both scenarios is not distinct from zero. At 5000 plants/ha, the impact of harvesting drops for both scenarios as almost all the plants are already infected by contact (Fig4. A and B). The difference between harvesting and non-harvesting scenarios remains with initial conditions from 5 to 30% of initially infected plants with maxima from 3000 to 1000 plants/ha (Fig. S1.5). At very low (1%) and very high (50%) initial infection scenarios, the effect of harvesting is more pronounced at higher densities (5000 plants/ha) and minimal densities (500 plants/ha), respectively (Fig. S1. 5)

A

B

Figure 4. Effect of different harvesting trajectories on the average plot rust infection. *A.* Full time
series of the average rust for three planting densities, for the control and harvesting trajectories
generated under the two coffee ripening scenarios (white: control, blue: asynchronous ripening
scenario, orange: synchronous ripening scenario). The width of each line represents the variability (±
one standard deviation around the mean) for the 30 repetitions. The time series for 1000 and 3000
plants/ha were equivalent and are not shown. **B.** Rust's increase due to the harvesting trajectories for
each of the ripening scenarios (harvesting - control), for the five different planting densities (blue:
asynchronous ripening, orange: synchronous ripening) and 30 repetitions. In all simulations, we
started with 20% of infected plants at T_0.

3.3. Medium size steps increase the proportion of infected networks of plants in the asynchronous scenario; the interaction between the proportion and size of infected networks explains the nonlinear behavior of rust increase

The proportion of infected networks of plants (where each pair of plants are 1.5 m apart or less) after harvest (# infected network AH/planting density) decreases as the planting density increases for both coffee ripening scenarios (subplot i in Fig. 5A, black circles in Fig. 5B). This makes sense since the connectivity between plants increases with density, making almost all the plants already infected before harvest due to the initial contact-mediated dispersal (gray circles in Fig. 5B). In the asynchronous ripening scenario, there are significantly more infected networks due to harvest trajectories than in the synchronous scenario (for planting densities higher than 500 plants/ha). This is due to the high proportion of medium sized steps between 1.5 and 8 m (58% of the steps) in trajectories generated under the asynchronous ripening (Fig.2), that potentially create new infected networks and contribute to the general infection (blue ribbon in Fig. 5C). In the synchronous scenario, 95% of steps in the trajectories are smaller or equal to 4 m (and 51% smaller than 1.5 m; Fig.2) and are more likely to disperse rust in the same contact-mediated networks, this is, in plants that are already infected (gray circles in Fig. 5B; orange ribbon in Fig.5C). Larger steps (>30 m) of trajectories in the asynchronous scenario, even if they connect distant zones of the plot, are not relevant for the infection as their proportion is very low (Fig. 5C).

On the other hand, the average size of new infected networks after harvest (# infected plants/infected network AH) increases with planting density as the connectivity increases (subplot ii in Fig. 5A, black circles in Fig. 5B). In this case, we found no significant differences between both harvesting scenarios. The estimated percentage of infected plants after harvest (% of plants AH/planting density; as the product of i and ii; subplot iii in Fig. 5A) follows a non-linear behavior in both scenarios equivalent to the observed increase of infection in Fig. 4. In this sense, the opposite behavior between the proportion of infected

networks after harvest and their mean size might explain the nonlinear behavior in rust

increase due to harvesting, with increasing planting densities. The higher rust increase in the

asynchronous scenario is solely related to a higher proportion of infected networks after

harvest due to the medium-sized steps.

For scenarios with very low initial infection (1%), almost no plants are infected

before harvest, so the main driver to increase the general infection will be the size of the

infected networks after harvest maximized at 5000 plants/ha (Fig. S1.4). On the contrary, at

very high initial infections (50%), almost all the plants are infected before harvest, and only

at very low densities (500 plants/ha), new plants can get infected by harvesting (Fig. S1.4).

Fig. 5. Proportion and mean size of infected networks after harvest and proportion of infective steps during harvest. **A.** We plot for each harvesting scenario (blue circle: asynchronous ripening, orange triangle: synchronous ripening) the proportion of infected networks after harvest (infected network AH/density; subplot i), their mean size (infected plants/infected network AH; subplot ii) and their product (100*infected plants AH/density; subplot iii). **B**. We show a 30 m2 zone of the plot in 500, 2000 and 5000 plants/ha densities to illustrate the non infected plants (white circles), the infected networks before harvest (gray circles) and the infected networks of plants after harvest and the cycles of contact- mediated dispersal (black circles). C. We show the proportion (mean ± standard deviation) of infective steps (steps that infected a new plant; yellow circles in Fig.1C) for each step length, three planting densities, and two harvesting scenarios.

4. DISCUSSION AND CONCLUSIONS

Many studies have stressed the relation between spatial structure of hosts, individual movement behavior, and pathogen transmission to predict and understand disease dynamics, though disease models with mechanistic representations remain rare (Manlove et al., 2022; White et al., 2018). Understanding the relationship between the spatial arrangement of the hosts and the different modes of dispersal of CLR is central to developing preventive strategies to reduce the impact of the pathogen on crop production (Hajian-Forooshani and Vandermeer, 2021).

Here we present a model to study the determinants of the movement of the workers in the field during harvesting and its relevance in CLR dispersal and infection at the plot level. Our results show that we can reproduce random walks, medium or large relocalizations by regulating the coffee ripening synchronicity and the scarcity of trees with berries at the end of harvest. These conditions could be *sufficient* to generate the different step lengths observed in the field. Relocalizations in deterministic walks on a random set of points have been documented for other systems (Brown et al., 2007) and simulated in rectangle domains (Boyer, 2008; Santos et al., 2007) or varying the size of the targets (Boyer et al., 2006). These studies present specific widths or target size uniformity ranges where the modeled workers fail to find closer unvisited points and "jump" to another zone of the area (Boyer, 2008; Boyer et al., 2006). Concerning the CLR impact, we found that increasing the planting density where only contact-mediated dispersal was present increases the final CLR at the plot level. This is in line with other studies (Hajian-Forooshani and Vandermeer, 2021; Zambolim, 2016; Mora Van Cauwelaert, et al. 2023) and emphasizes the importance of planting at lower densities to avoid big bursts of the pathogen (Ehrenbergerová et al., 2018). The harvest event magnifies the final CLR infection for all scenarios and densities. Its effect is more important for medium densities (2000 to 3000 plants/ha), which is a common

density of coffee plots owned by small-scale farmers (Soto-Pinto et al., 2000). If densities are

too high, harvesting becomes irrelevant as the plot is already infected. Other modes of

dispersal, such as wind-mediated relocalizations from neighbor plantations, were not

included (as in (Hajian-Forooshani and Vandermeer, 2021; Vandermeer and Rohani, 2014)).

This assumption is reasonable for the rainy seasons where local contact is more relevant to

build the initial epidemic (Boudrot et al., 2016). The different types of trajectories generated

under the two coffee ripening scenarios changed the plot average CLR and varied

nonlinearly with increasing planting density. Movements with medium and long steps in

the asynchronous ripening scenario promote new foci of infection (Zadoks and Van Den

Bosch, 1994) that eventually become newly infected networks with optimum size and

number at planting densities between 2000 to 3000 plants/ha. In particular, the high

proportion of medium-sized steps thrives the infection due to its high probability to infect

distant coffee networks. This movement is common in plots with asynchronous ripening

where workers jump from one row to the other to maximize harvest (Mora Van Cauwelaert

et al., 2023b). In this sense, the spatial heterogeneity in coffee maturation creates fragmented

host spatial patterns that interact with the movement of harvesters and lead to CLR

outbreaks at landscape levels (as suggested by White et al., 2018). It is important to note that

we modeled a random arrangement of the plants in the lattice. This assumption is likely

valid for old plantations and shaded coffee managements (Moguel and Toledo, 1999) but

should be revisited for aggregated or uniform patterns.

While we observe an increase in CLR in the asynchronous scenario, our results do

not imply that asynchronous ripening leads *per se* to higher rust. On the one hand, the long

steps are not generated due to the asynchronous ripening alone, but mainly due to the

scarcity of closer trees at the end of harvest. In fact, when we run the synchronous scenario

in a scarcity scenario, longer steps also appear (data not shown). This would also be the case

for large-scale landlord owned plantations where the workers are paid by daily harvest

(Jiménez-Soto, 2021); even if the following plant is hundreds of meters distant, it becomes

necessary for them to reach it within the same day (Mora Van Cauwelaert et al., 2023b). In

these plantations it is also common to force the workers to harvest at the end of the season

when only few scattered trees still have berries (Mora Van Cauwelaert et al., 2023b). This

would promote medium-sized steps, the main drivers of higher plot level infection. On the

other hand, the simulated trajectories in the two coffee ripening scenarios suppose otherwise

similar conditions on the plot. But in the field, asynchronous ripening is associated with

characteristics that change these conditions and that can reduce the general rust impact. For

example, the susceptibility of coffee plants to rust increases with fruit charge (Salgado et al.,

2008). Hence, the asynchrony in the maturation of berries due to erratic rainfalls, or simply

due to different coffee varieties (DaMatta et al., 2007), generates a mixed spatial pattern of

resistant and susceptible plants that could reduce rust dispersal. Besides, shade trees in the

plot form different patches of maturation (and the potential for long steps during harvest)

but also create barriers to wind-mediated rust dispersal (Gagliardi et al., 2020).

Our study aims to spur discussion on practices that can reduce the impact of

harvesting and specific trajectories, in scenarios where one can benefit from asynchronous

maturation of berries and shaded plantations (Ehrenbergerová et al., 2018; Liebig et al., 2019;

Soto-Pinto et al., 2000). This is crucial as almost all the plantations have or will have this

kind of ripening (DaMatta et al., 2007; Masarirambi et al., 2009), especially in the predicted

climate scenarios of increasingly erratic rainfalls (Avelino et al., 2015; Huang et al., 2012;

Rana et al., 2014). Along with planting different genetic varieties (Silva et al., 2006) or having

other species of trees inside the plot and between them (Avelino et al., 2022), our results

show the importance of reducing medium to long-distance movements during the same day

by avoiding harvesting at the end of the harvest season when trees ripening asynchronicity

and CLR levels are higher (Avelino et al., 1991; Mora Van Cauwelaert, 2023b); or simply

skipping rust-infected trees and working fewer hours a day, even if this represents a short

term decrease in harvested berries. These practices presuppose the right to decide how and

when to harvest the coffee plantations.

We hope this mechanistic model will contribute to educating our intuition on the

relationships between host spatial disposition and dispersal, and it will help us build

working hypotheses for an agroecological approach to pest management that does not rely

solely on the control of the infection stage of the pathogen, but also on preventive measures

that focus on the dispersal of the fungus.

5. ACKNOWLEDGMENTS

EMVC is a doctoral student from the Programa de Doctorado en Ciencias Biomédicas,

Universidad Nacional Autónoma de México, and has received CONACyT scholarship

686776 and support from the Programa de Apoyo a los Estudios del Posgrado (PAEP) for his

research stay at the University of Michigan (USA). MB acknowledges financial support from

UNAM-DGAPA-PAPIIT (IN207819). JV acknowledges financial support from the US

National Science Foundation, grant number DEB-1853261. EMVC thanks the people from La

Parcela and Perfectomeer lab for their great ideas and friendship. EMVC and MBK are also

thankful to Gustavo Bautista, Gabriel Domínguez, Elisa Lotero and Cecilia González for the

discussions and help during the fieldwork. The authors thank Denis Boyer and Eugenio

Azpeitia for their valuable feedback on previous versions of the manuscript and Rodrigo

García Herrera for his technical support.

6. CRediT

EMVC: Conceptualization, Methodology, Software, Validation, Formal Analysis,

Investigation, Writing-Original Draft, Visualization. **KL**: Conceptualization, Methodology,

Writing - Review & Editing. **ZHF**: Conceptualization, Methodology, Software, Writing -

Review & Editing. **JV:** Conceptualization, Methodology, Writing - Review & Editing,

Supervision, Funding acquisition. **MB:** Conceptualization, Methodology, Writing - Review

& Editing, Supervision, Project administration, Funding acquisition

7.REFERENCES

Andow, D.A., Hidaka, K., 1989. Experimental natural history of sustainable agriculture:

syndromes of production. Agriculture, Ecosystems & Environment 27, 447–462.

https://doi.org/10.1016/0167-8809(89)90105-9

Avelino, J., Cristancho, M., Georgiou, S., Imbach, P., Aguilar, L., Bornemann, G., Läderach,

P., Anzueto, F., Hruska, A.J., Morales, C., 2015. The coffee rust crises in Colombia

and Central America (2008–2013): impacts, plausible causes and proposed solutions.

Food Security 7, 303–321. https://doi.org/10.1007/s12571-015-0446-9

Avelino, J., Gagliardi, S., Perfecto, I., Isaac, M., Liebig, T., Vandermeer, J., Merle, I., Hajian-

Forooshani, Z., Tabeuguia Motisi, N., 2022. Tree effects on coffee leaf rust at field and

landscape scales. Plant Disease 107, 1–56. https://doi.org/10.1094/pdis-08-21-1804-

fe

Avelino, J., Muller, R.A., Cilas, C., Velasco Pascual, H., 1991. Développement et

comportement de la rouille orangée du caféier (Hemileia vastatrix Berk. et Br.) dans

des plantations en cours de modernisation, plantées de variétés naines, dans le sud-

est du Mexique. Café, Cacao, Thé 35, 21–37. https://agritrop.cirad.fr/416561/

Avelino, J., Willocquet, L., Savary, S., 2004. Effects of crop management patterns on coffee

rust epidemics. Plant Pathology 53, 541–547. https://doi.org/10.1111/j.1365-

3059.2004.01067.x

Beasley, E.M., Aristizábal, N., Bueno, E.M., White, E.R., 2022. Spatially explicit models

predict coffee rust spread in fragmented landscapes. Landscape Ecology 37, 2165–

2178. https://doi.org/10.1007/s10980-022-01473-1

Becker, S., Kranz, J., 1977. Comparative Studies on Dispersal of Hemileia vastratrix in Kenya.

Zeitschrift fur Planzenschultz- Journal of Plant Diseases and Protection 84, 526–539.

https://www.jstor.org/stable/ 43213550

Boudrot, A., Pico, J., Merle, I., Granados, E., Vílchez, S., Tixier, P., De Melo Virginio Filho, E.,

Casanoves, F., Tapia, A., Allinne, C., Rice, R.A., Avelino, J., 2016. Shade effects on the

dispersal of airborne Hemileia vastatrix uredospores. Phytopathology 106, 527–580.

https://doi.org/10.1094/PHYTO-02-15-0058-R

Boyer, D., 2008. Intricate dynamics of a deterministic walk confined in a strip. Europhys.

Lett. 83, 20001. https://doi.org/10.1209/0295-5075/83/20001

Boyer, D., Ramos-Fernández, G., Miramontes, O., Mateos, J.L., Cocho, G., Larralde, H.,

Ramos, H., Rojas, F., 2006. Scale-free foraging by primates emerges from their

interaction with a complex environment. Proceedings of the Royal Society B:

Biological Sciences 273, 1743–1750. https://doi.org/10.1098/rspb.2005.3462

Brown, C.T., Liebovitch, L.S., Glendon, R., 2007. Lévy Flights in Dobe Ju/'hoansi Foraging

Patterns. Hum Ecol 35, 129–138. https://doi.org/10.1007/s10745-006-9083-4

DaMatta, F.M., Ronchi, C.P., Maestri, M., Barros, R.S., 2007. Ecophysiology of coffee growth

and production. Braz. J. Plant Physiol. 19, 485–510. https://doi.org/10.1590/S1677-

04202007000400014

Ehrenbergerová, L., Kučera, A., Cienciala, E., Trochta, J., Volařík, D., 2018. Identifying key

factors affecting coffee leaf rust incidence in agroforestry plantations in Peru.

Agroforestry Systems 92, 1551–1565. https://doi.org/10.1007/s10457-017-0101-x

Gagliardi, S., Avelino, J., Beilhe, L.B., Isaac, M.E., 2020. Contribution of shade trees to wind

dynamics and pathogen dispersal on the edge of coffee agroforestry systems: A

functional traits approach. Crop Protection 130, 105071.
https://doi.org/10.1016/j.cropro.2019.105071

Hajian-Forooshani, Z., Vandermeer, J., 2021. Emergent spatial structure and pathogen
epidemics: the influence of management and stochasticity in agroecosystems.
Ecological Complexity 45, 100872. https://doi.org/10.1016/j.ecocom.2020.100872

Huang, W.-C., Chiang, Y., Wu, R.-Y., Lee, J.-L., Lin, S.-H., 2012. The Impact of Climate
Change on Rainfall Frequency in Taiwan. Terr. Atmos. Ocean. Sci. 23, 553–564.
https://doi.org/10.3319/TAO.2012.05.03.04(WMH)

Jiménez-Soto, E., 2021. The political ecology of shaded coffee plantations: conservation
narratives and the everyday-lived-experience of farmworkers. Journal of Peasant
Studies 48, 1284–1303. https://doi.org/10.1080/03066150.2020.1713109

Keeling, M.J., 1999. The effects of local spatial structure on epidemiological invasions.
Proceedings of the Royal Society B: Biological Sciences 266, 859–867.
https://doi.org/10.1098/rspb.1999.0716

Kushalappa, A.C., Eskes, A.B., 1989. Advances in Coffee Rust Research. Annual Review of
Phytopathology 27, 503–531. https://doi.org/10.1146/annurev.py.27.090189.002443

Leguizamón-Caycedo, J.E., Orozco-Gallego, L., Gómez-Gómez, L., 1998. Períodos de
incubación (pi) y de latencia (pl) de la roya del cafeto en la zona cafetera central de
Colombia. Cenicafé 49, 325–339.
https://www.cenicafe.org/es/publications/arc049(04)325-339.pdf

Li, K., Hajian-Forooshani, Z., Vandermeer, J., Perfecto, I., 2023. Coffee leaf rust (Hemileia
vastatrix) is spread by rain splash from infected leaf litter in a semi-controlled
experiment. J. Plant Pathol. 105, 667–672. https://doi.org/10.1007/s42161-023-01404-
2

Li, K., Vandermeer, J., Perfecto, I., 2016. Disentangling endogenous versus exogenous

pattern formation in spatial ecology: A case study of the ant Azteca sericeasur in

southern Mexico. R. Soc. Open Sci. 3, 160073. https://doi.org/10.1098/rsos.160073

Liebig, T., Ribeyre, F., Läderach, P., Poehling, H.M., van Asten, P., Avelino, J., 2019.

Interactive effects of altitude, microclimate and shading system on coffee leaf rust.

Journal of Plant Interactions 14, 407–415.

https://doi.org/10.1080/17429145.2019.1643934

Manlove, K., Wilber, M., White, L., Bastille-Rousseau, G., Yang, A., Gilbertson,

M.L.J., Craft, M.E., Cross, P.C., Wittemyer, G., Pepin, K.M., 2022. Defining an

epidemiological landscape that connects movement ecology to pathogen

transmission and pace-of-life. Ecol. Lett. 25, 1760–1782.

https://doi.org/10.1111/ele.14032

Masarirambi, M.T., Chingwara, V., Shongwe, V.D., 2009. The effect of irrigation on

synchronization of coffee (Coffea arabica L.) flowering and berry ripening at

Chipinge, Zimbabwe. Physics and Chemistry of the Earth, Parts A/B/C 34, 786–789.

https://doi.org/10.1016/j.pce.2009.06.013

McCook, S., Vandermeer, J., 2015. The Big Rust and the Red Queen: Long-term perspectives

on coffee rust research. Phytopathology 105, 1164–1173.

https://doi.org/10.1094/PHYTO-04-15-0085-RVW

Moguel, P., Toledo, V.M., 1999. Biodiversity conservation in traditional coffee systems of

Mexico. Conservation Biology 13, 11–21. https://doi.org/10.1046/j.1523-

1739.1999.97153.x

Mora Van Cauwelaert, E., González González, C., Boyer, D., Hajian-Forooshani, Z.,

Vandermeer, J., Benítez, M., 2023a. Dispersal and plant arrangement condition the

timing and magnitude of coffee rust infection. Ecological Modelling 475, 110206.

https://doi.org/10.1016/j.ecolmodel.2022.110206

Mora Van Cauwelaert, E., Boyer, D., Jimenez-Soto, E., González González, C.., & Benítez, M. (2023b). Spatial trajectories of coffee harvesting in large-scale plantations: ecological and management drivers and implications. *bioRxiv*, 2023-10. https://doi.org/10.1101/2023.10.31.564942

Pangga, I.B., Hanan, J., Chakraborty, S., 2011. Pathogen dynamics in a crop canopy and their evolution under changing climate: Pathogen dynamics and evolution. Plant Pathology 60, 70–81. https://doi.org/10.1111/j.1365-3059.2010.02408.x

Papaïx, J., Burdon, J.J., Zhan, J., Thrall, P.H., 2015. Crop pathogen emergence and evolution in agro-ecological landscapes. Evolutionary Applications 8, 385–402. https://doi.org/10.1111/eva.12251

Park, A.W., Gubbins, S., Gilligan, C.A., 2001. Invasion and persistence of plant parasites in a spatially structured host population. Oikos 94, 162–174. https://doi.org/10.1034/j.1600-0706.2001.10489.x

Perfecto, I., Jiménez-Soto, M. E., & Vandermeer, J. 2019. Coffee landscapes shaping the Anthropocene: forced simplification on a complex agroecological landscape. *Current Anthropology*, *60*(S20), S236-S250. https://doi.org/10.1086/703413

Ramírez-Camejo, L.A., Keith, L.M., Matsumoto, T., Sugiyama, L., Fukada, M., Brann, M., Moffitt, A., Liu, J., Aime, M.C., 2022. Coffee Leaf Rust (Hemileia vastatrix) from the Recent Invasion into Hawaii Shares a Genotypic Relationship with Latin American Populations. JoF 8, 189. https://doi.org/10.3390/jof8020189

Rana, A., Foster, K., Bosshard, T., Olsson, J., Bengtsson, L., 2014. Impact of climate change on rainfall over Mumbai using Distribution-based Scaling of Global Climate Model projections. Journal of Hydrology: Regional Studies 1, 107–128. https://doi.org/10.1016/j.ejrh.2014.06.005

Salgado, P.R., Favarin, J.L., Leandro, R.A., De Lima Filho, O.F., 2008. Total phenol concentrations in coffee tree leaves during fruit development. Scientia Agricola 65, 354–359. https://doi.org/10.1590/S0103-90162008000400005

Santos, M.C., Boyer, D., Miramontes, O., Viswanathan, G.M., Raposo, E.P., Mateos, J.L., Da Luz, M.G.E., 2007. Origin of power-law distributions in deterministic walks: The influence of landscape geometry. Phys. Rev. E 75, 061114. https://doi.org/10.1103/PhysRevE.75.061114

Schieber, E., Zentmyer, G.A., 1984. Coffee Rust in Western Hemisphere. Plant Disease 68, 89–93.

Silva, M.D.C., Várzea, V., Guerra-Guimarães, L., Azinheira, H.G., Fernandez, D., Petitot, A.S., Bertrand, B., Lashermes, P., Nicole, M., 2006. Coffee resistance to the main diseases: Leaf rust and coffee berry disease. Brazilian Journal of Plant Physiology 18, 119–147. https://doi.org/10.1016/j.ijepes.2005.11.005

Soto-Pinto, L., Perfecto, I., Castillo-Hernandez, J., Caballero-Nieto, J., 2000. Shade effect on coffee production at the northern Tzeltal Zone of the state of Chiapas, Mexico. Agriculture, Ecosystems and Environment 80, 61–69. https://doi.org/10.1016/S0167-8809(00)00134-1

Talhinhas, P., Batista, D., Diniz, I., Vieira, A., Silva, D.N., Loureiro, A., Tavares, S., Pereira, A.P., Azinheira, H.G., Guerra-Guimarães, L., Várzea, V., Silva, M. do C., 2017. The coffee leaf rust pathogen Hemileia vastatrix: one and a half centuries around the tropics. Molecular Plant Pathology 18, 1039–1051. https://doi.org/10.1111/mpp.12512

Vandermeer, J., Hajian-Forooshani, Z., Perfecto, I., 2018. The dynamics of the coffee rust disease: an epidemiological approach using network theory. European Journal of Plant Pathology 150, 1001–1010. https://doi.org/10.1007/s10658-017-1339-x

Vandermeer, J., Rohani, P., 2014. The interaction of regional and local in the dynamics of the

coffee rust disease. http://arxiv.org/abs/1407.8247

Waller, J.M., 1982. Coffee rust-epidemiology and control. Crop Protection 1, 385–404.

https://doi.org/10.1016/0261-2194(82)90022-9

Waller, J.M., 1972. Coffee rust in latin america. PANS Pest Articles and News Summaries 18,

402–408. https://doi.org/10.1080/09670877209412699

White, L. A., Forester, J. D., & Craft, M. E. (2018). Disease outbreak thresholds

emerge from interactions between movement behavior, landscape structure, and

epidemiology. Proceedings of the National Academy of Sciences, *115*(28), 7374-7379.

https://doi.org/10.1073/pnas.1801383115

Zadoks, J.C., Van Den Bosch, F., 1994. On the Spread of Plant Disease: A Theory on Foci.

Annu. Rev. Phytopathol. 32, 503–521.

https://doi.org/10.1146/annurev.py.32.090194.002443

Zambolim, L., 2016. Current status and management of coffee leaf rust in Brazil. Tropical

Plant Pathology 41, 1–8. https://doi.org/10.1007/s40858-016-0065-9

690