Early Steps of Protein Disaggregation by Hsp70 Chaperone and Class B J-Domain Proteins are Shaped by Hsp110

Wiktoria Sztangierska[1], Hubert Wyszkowski[1], Maria Pokornowska[1], Klaudia Kochanowicz[1], Michał Rychłowski[1], Krzysztof Liberek[1, 2], Agnieszka Kłosowska[1, 2]

[1]Intercollegiate Faculty of Biotechnology of University of Gdańsk and Medical University of Gdańsk, University of Gdańsk, Gdańsk 80-307, Poland

[2]Corresponding authors:
Krzysztof Liberek, Intercollegiate Faculty of Biotechnology of University of Gdańsk and Medical University of Gdańsk, University of Gdańsk, Abrahama 58, 80-307, Gdańsk, Poland, Tel: +48 58 523 63 46, Email: krzysztof.liberek@ug.edu.pl

Agnieszka Kłosowska, Intercollegiate Faculty of Biotechnology of University of Gdansk and Medical University of Gdańsk, University of Gdańsk, Abrahama 58, 80-307, Gdańsk, Poland, Tel: +48 58 523 63 59, Email: agnieszka.klosowska@ug.edu.pl

Abstract

Hsp70 is a key cellular system counteracting protein misfolding and aggregation, associated with stress, ageing and disease. Hsp70 solubilizes aggregates and aids protein refolding through substrate binding and release cycles regulated by co-chaperones: J-domain proteins (JDPs) and Nucleotide Exchange Factors (NEFs). Here, we elucidate the collaborative impact of Hsp110 NEFs and different JDP classes throughout Hsp70-dependent aggregate processing. We show that Hsp110 plays a major role at initial stages of disaggregation, determining its final efficacy. The NEF catalyses the recruitment of thick Hsp70 assemblies onto aggregate surface, which modifies aggregates into smaller species more readily processed by chaperones. Hsp70 stimulation by Hsp110 is much stronger with Class B than Class A JDPs and requires the auxiliary interaction between Class B JDP and the Hsp70 EEVD motif. Furthermore, we demonstrate for the first time that Hsp110 disrupts the JDP-Hsp70 interaction. Such destabilisation of chaperone complexes at the aggregate surface might improve disaggregation, but also lead to the inhibition above the substoichiometric Hsp110 optimum. Thus, balanced interplay between the co-chaperones and Hsp70 is critical to unlock its disaggregating potential.

Introduction

During stress, protein homeostasis is perturbed by protein misfolding and aggregation. Accumulation of protein aggregates disrupts cellular functions and contributes to ageing and is a hallmark of many neurodegenerative diseases (Hartl *et al*, 2011; Morimoto, 2008). A protein quality control system, relying on a network of cooperating chaperones, has evolved to moderate stress-induced proteotoxicity and rebalance proteostasis.

To refold proteins from aggregates into their native state, Eukaryotes use a system comprising Hsp70, an ATP-dependent chaperone, and its co-chaperones: an Hsp110 nucleotide exchange factor (NEF) and a J-domain protein (JDP/Hsp40), together with an Hsp100 disaggregase (Glover & Lindquist, 1998). The process is initiated by a JDP, which delivers Hsp70 to an aggregated substrate and whose J-domain induces ATP hydrolysis in Hsp70, resulting in conformational changes that stabilise the interaction with aggregates (Rohland *et al*, 2022). Next, Hsp100 interacts with the Hsp70-aggregate complex and the aggregate-trapped polypeptides are translocated through the Hsp100 hexamer. The disentangled and released polypeptides can fold back into their native conformation, alone or with further aid of chaperones (Lum *et al*, 2004; Schaupp *et al*, 2007; Seyffer *et al*, 2012; Weibezahn *et al*, 2004; Zietkiewicz *et al*, 2004).

Hsp110 co-chaperone boosts the Hsp70 activity by stimulating nucleotide exchange and substrate release (Dragovic *et al*, 2006; Raviol *et al*, 2006; Shaner *et al*, 2005). It belongs to the Hsp70 superfamily, with

identical domain organisation to Hsp70 but distinct size and arrangement of the nucleotide-binding domain
(NBD) and substrate-binding domain (SBD) (Easton *et al*, 2000; Liu & Hendrickson, 2007). In contrast to
Hsp70, Hsp110 is unable to refold denatured proteins, yet it may bind to and prevent aggregation of certain
misfolding substrates (Garcia *et al*, 2017; Goeckeler *et al*, 2008; Goeckeler *et al*, 2002; Oh *et al*, 1997; Oh
et al, 1999; Polier *et al*, 2010; Xu *et al*, 2012). In its well-established NEF function, SBD of Hsp110 interacts
with the ADP form of Hsp70, which mediates the release of the nucleotide. Subsequent ATP binding to
NBD of Hsp70 prompts the opening of the SBD domain and substrate dissociation, resetting Hsp70 for
another round of the cycle (Andreasson *et al*, 2008a; Laufen *et al*, 1999; Liberek *et al*, 1991; Mayer &
Bukau, 2005; Mayer & Gierasch, 2019).

In vivo studies on yeast chaperones demonstrated that the Hsp110 co-chaperone Sse1 plays a key role
in Hsp70 recruitment to aggregates and is essential for protein disaggregation (Kaimal *et al*, 2017). Hsp110
is a major cytoplasmic NEF of Hsp70 and the deletion of its both paralogs, *SSE1* and *SSE2*, is lethal and
the growth can only be supported, albeit with less efficacy, by overexpression of *FES1,* another
cytoplasmic NEF from an unrelated armadillo family (Abrams *et al*, 2014; Raviol *et al.*, 2006).

The importance of Hsp110 is also manifested *in vitro* in a low disaggregation activity of the yeast Hsp70
system without Hsp110 and the complete interdependence between their human orthologs (Glover &
Lindquist, 1998; Mattoo *et al*, 2013; Nillegoda *et al*, 2015; Rampelt *et al*, 2012). Recent studies on the
human system revealed that Hsp110 function in disaggregation of amyloid fibrils is not limited to its NEF
activity but it may also affect the architecture of chaperone complexes, inducing Hsp70 clustering, although
it is not clear whether similar effects occur during disaggregation of non-fibrillar, stress-associated
aggregates. The effect is not well understood, but it presumably increases entropic pulling of aggregated
polypeptides, ultimately leading to their disentanglement by the Hsp70 system, efficient even without the
Hsp100 disaggregase. Developing an Hsp70 system that is self-sufficient in disaggregation could have
compensated for the loss of the Hsp100 disaggregase in a common metazoan ancestor (Mattoo *et al.*,
2013; Shorter, 2011).

Another factor that potentiates the Hsp70 system is the diversity of JDP paralogs, assigned to cytoplasmic
classes A and B, which differently regulate Hsp70 (Lu & Cyr, 1998; Nillegoda *et al.*, 2015). The main
distinction between class A and B JDPs is an auxiliary interaction site between CTD1 domain of class B
JDPs and the C-terminal EEVD motif of Hsp70 (Yu *et al*, 2015). Class B JDPs are additionally regulated
by an autoinhibitory mechanism, in which Hsp70 binding by the J-domain is restricted by a neighbouring
helix. Upon binding to the C-terminal EEVD motif of Hsp70, the J-domain is released and can interact with
the NBD of Hsp70 (Faust *et al*, 2020; Wentink *et al*, 2020). We have recently shown that the yeast class A
JDP Ydj1 and class B JDP Sis1 exhibit diverse mechanisms during Hsp70 binding to aggregated
substrates. Ydj1, in accordance with the classical model of the Hsp70 ATPase cycle, binds misfolded
polypeptides and loads Hsp70 onto aggregates. Unlike that, Sis1 only weakly binds protein substrates but

due to a the more complex interaction with Hsp70, it loads more Hsp70 molecules onto aggregates, which

results in more efficient disaggregation (Wyszkowski *et al*, 2021).

Despite the vast knowledge on how Hsp110 serves as a regulator of the ATPase cycle of Hsp70

(Andreasson *et al.*, 2008a; Andreasson *et al*, 2008b; Dragovic *et al.*, 2006; Raviol *et al.*, 2006; Shaner *et*

al, 2006) little is known about the function of Hsp110 considering the mechanism of Hsp70 interaction with

different JDP classes.

Here, we demonstrate that the interplay between Hsp110 and Hsp70 in disaggregation strictly relies on

the class of a JDP and unravel the critical role of the B-class specific interaction with the EEVD motif of

Hsp70 for the Hsp110-dependent stimulation. Furthermore, we elucidate differential contribution of the

NEF across different phases of protein recovery from aggregates. We employ Bio-Layer Interferometry to

investigate the Hsp110 impact on the formation of chaperone complexes at the aggregate surface and we

assess changes in aggregates properties associated with the abundant chaperone binding. Finally, we

address a question of the competition between the NEF and JDP co-chaperones. Our findings shed new

light on the mechanisms behind the potentiation and inhibition of the disaggregation activity of Hsp70 by

its co-chaperones.

Results

Stimulation of the Hsp70 disaggregation activity by Hsp110 depends on the class of JDP

Our recent studies showed that during the recovery of aggregated proteins, the Hsp70 chaperone system

exhibits different mechanisms of action with class A and class B JDP co-chaperones (Wyszkowski *et al.*,

2021). To better understand the interplay within the Hsp70 chaperone network, we addressed how Hsp110

affects Hsp70 with members of different JDP classes. First, we tested the effect of Sse1, the most

abundant yeast NEF, on the recovery of a model protein substrate, aggregated luciferase, by Hsp70 (Ssa1)

and class A (Ydj1) or class B (Sis1) JDPs. Compared with Ssa1-Sis1, which exhibits delayed start of

disaggregation characteristic for class B JDPs, the initial luciferase recovery was significantly faster, with

higher overall output, in the presence of Sse1 (Figure 1A). In contrast, Sse1 slightly decreased the

disaggregation efficacy in the case of Ssa1-Ydj1 (Figure 1A). With another aggregated substrate, GFP,

Sse1 improved the disaggregation activity with either of the JDPs, albeit for Ssa1-Sis1, the stimulation was

almost 4 times higher than with Ydj1 (Figure 1—figure supplement 1A).

Addition of the Hsp104 disaggregase to the system increased the total disaggregation efficacy and

enhanced the positive effects of Sse1 in the case of either JDP and either aggregated substrate

(Figure 1—figure supplement 1B, C). The chaperone system comprising Hsp104-Ssa1-Sis1 with Sse1

yielded approximately 3 times higher luciferase and GFP recovery rates than without the NEF, while in the

case of Hsp104-Ssa1-Ydj1, the simulation by Sse1 was much weaker, from marginal to twofold, depending

on the protein substrate (Figure 1—figure supplement 1B, C). Thus, the positive effect of the Hsp110 nucleotide exchange factor on protein recovery from aggregates is substantially more pronounced when the Hsp70 chaperone system comprises a class B JDP.

Theoretically, the stimulation by Sse1 might occur at one or many stages of protein disaggregation – starting from the association of the Hsp70 system with an aggregate surface, through polypeptide disentanglement, to the final folding of the released substrate. Previous studies show that Ssa1-Ydj1 is much more effective than Ssa1-Sis1 in restoring the native structure of unfolded luciferase (Lu & Cyr, 1998), which could represent the final step of disaggregation – the polypeptide folding after translocation by Hsp104. To test how Sse1 contributes to the protein folding by the Hsp70 system, we measured the recovery of denatured, non-aggregated luciferase (Imamoglu *et al*, 2020). Curiously, neither of the systems: Ssa1-Sis1 nor Ssa1-Ydj1, was stimulated in luciferase folding by Sse1 (Figure 1—figure supplement 1D), suggesting that the NEF plays an important role at earlier stages of disaggregation.

We recently showed that JDP co-chaperones determine the association kinetics and the size of chaperone complexes formed at the aggregate surface, which contributes to the total effectiveness of disaggregation (Wyszkowski *et al.*, 2021). To gain insight into the influence of the Hso110 on the assembly of the protein disaggregation complex, we analysed the binding of the Hsp70 system to heat-aggregated luciferase using bio-layer interferometry (BLI). The presence of Sse1 increased the rate of Ssa1-Sis1 association with an aggregate-covered biosensor approximately 2 times and resulted in a 50% thicker protein layer (Figure 1B). In contrast, Sse1 did not substantially influence the binding level of Ssa1-Ydj1 (Figure 1B). Similar effects were observed with heat-aggregated GFP or heat-aggregated yeast lysate proteins immobilised on the sensor, indicating that they are not substrate-specific (Figure 1—figure supplement 1E, F). The binding of chaperones to the aggregates required ATP (Figure 1B), which is consistent with the ATP-dependence of Hsp70 (Figure 1B). Sse1 did not stimulate the binding of Ssa1 or JDPs alone, nor did it interact with the aggregate on its own (Figure 1—figure supplement 1G).

We also assessed the evolutionary conservation of the observed trends and applied BLI to investigate how Hsp105, a human Hsp110, affects aggregate binding by the Hsp70 system. Similarly as the yeast system, Hsc70-DNAJB4 (orthologous to Ssa1-Sis1) exhibited delayed binding and the addition of Hsp105 resulted in much faster association and a 6-fold increase in the binding level (Figure 1C). In the case of Hsc70-DNAJA2 (orthologous to Ssa1-Ydj1), the binding also increased upon Hsp105 addition, however, the stimulation was three times less pronounced than with DNAJB4 (Figure 1C). In contrast to the yeast proteins, the stimulation occurred with both JDP classes, which is in line with the stronger dependence of the human Hsp70 system on the NEF in protein disaggregation (Figure 1—figure supplement 1H).

The BLI results suggest that the Hsp110 nucleotide exchange factor greatly improves Hsp70 binding to aggregates, especially with class B JDPs. What could be the basis for this specificity? A major

discrimination factor between the two JDP classes is the stable interaction between class B JDPs and the

EEVD-motif of Hsp70 (Yu *et al.*, 2015). Perturbation of this interaction restricts the J-domain-dependent

Hsp70 activation, which can be restored through the E50A mutation in Sis1 (Yu *et al.*, 2015) or the mutation

in DNAJB4 Helix 5 (Faust *et al.*, 2020). Previous studies showed that Ssa1$^{\Delta EEVD}$-Sis1^{E50A} resembles

Ssa1-Ydj1 in aggregate binding and disassembly (Wyszkowski *et al.*, 2021), therefore we asked what

effect this disruption has on the stimulation by Sse1. We observed that the Sis1 E50A-Ssa1 ΔEEVD system

is strongly inhibited by Sse1 in protein disaggregation (Figure 1—figure supplement 1I) and the level of

aggregate binding is also negatively affected (Figure 1—figure supplement 1J). This suggests that the

Hsp70-JDP stimulation by Hsp110 is functionally linked to the Sis1-Ssa1 EEVD interaction.

We next asked whether Hsp110 contributes to the extra chaperone layer at the aggregate surface in a

way that it comprises more Hsp70 molecules. To evaluate the amount of bound Hsp70, we carried out an

analogous BLI experiment with fluorescently labelled Ssa1 (Ssa1*A488). After protein dissociation to the

basal level, the total dissociated Ssa1*A488 was quantified based on fluorescence. In the presence of

Sis1 and Sse1, the amount of Ssa1*A488 was nearly 2 times higher than in the presence of Sis1 only,

whereas Sse1 did not influence the amount Ssa1*A488 bound to the biosensor when Ydj1 was applied

(Figure 1—figure supplement 2A). The increase in the fluorescence signal corresponded to that of BLI

(Figure 1—figure supplement 2A), suggesting a major contribution of Ssa1 to the thickness of the

chaperone complex forming on the aggregate surface.

Since the presence of Sse1 substantially increased the association of Ssa1 with aggregates, we asked if

the problem limiting Hsp70 binding that is overcome by Hsp110 is the insufficient availability of Hsp70

molecules targeting the aggregate. To address that, we tested whether a similar effect to the Sse1-induced

stimulation of Ssa1 binding can be obtained by increasing Ssa1 concentration itself. Sse1 stimulated Ssa1

binding much above the level that could be achieved for the saturating concentration of Ssa1 in the

absence of Sse1 (Figure 1—figure supplement 2B). This indicates that the mechanism of stimulation by

Hsp110 is more complex than an enrichment of the pool of Hsp70 molecules capable of substrate binding

and might involve generating more Hsp70-binding sites.

To explore whether the observed effects are unique to Hsp110, we examined another cytosolic NEF that

belongs to the Armadillo type family, Fes1. It is structurally distinct from Hsp110, with a C-terminal

armadillo domain that triggers nucleotide exchange in Hsp70 through a different mechanism (Gowda *et al*,

2018). Fes1 has weaker affinity for Hsp70 and lower nucleotide exchange activity than Sse1 (Dragovic *et

al.*, 2006). Unlike Sse1, Fes1 does not directly bind protein substrates, but its N-terminal RD domain is

involved in substrate release from Hsp70 (Gowda *et al.*, 2018). When we carried out luciferase

disaggregation by Ssa1-Sis1, the level of stimulation achieved by Sse1 required 10 times more Fes1

(Figure 1—figure supplement 3A). Similarly, 1 µM Fes1 stimulated Ssa1 binding to luciferase aggregates

on the BLI sensor to the level achieved with 0.1 µM Sse1 (Figure 1—figure supplement 3B, C, 2A). This

result suggests that the substrate-binding activity specific to Hsp110 is not necessary to increase the

Sis1-Ssa1 binding to aggregates and that the effective NEF concentration is negatively correlated with its

affinity for Hsp70.

Sse1 leads to Hsp70-depenent reduction of aggregate size

Recently, we have reported that larger Hsp70-JDP assemblies at the aggregate surface, dependent on

Sis1 and its interaction with EEVD, can modify aggregates into misfolded protein species that are more

amenable to disaggregation (Wyszkowski *et al.*, 2021). To assess whether Hsp110 further stimulates such

aggregate-remodelling activity, and this way contributes to the more efficient protein recovery, we used a

variant of Hsp104 with abrogated interaction with Hsp70, D484K F508A (Hsp104^{mut}). Hsp104^{mut} does not

require Hsp70 for allosteric activation and aggregate binding and it can serve as an indicator of

Hsp70-dependent aggregate modification by facilitating final reactivation of its products

(Chamera *et al*, 2019). When heat-aggregated GFP was initially incubated with Ssa1-Sis1-Sse1, which

yielded very low protein recovery, we added Hsp104^{mut} and observed much faster and more effective GFP

reactivation than when the substrate was first incubated with Ssa1-Sis1 only (Figure 2A). In an analogous

experiment with Ydj1, we also observed stimulation by Sse1, yet not as strong as in the presence of Sis1

(Figure 2A).

Aggregate remodelling by Hsp70-JDP, improved by Hsp110, might induce changes limited to the

aggregate surface, such as partial polypeptide disentanglement that uncovers additional

chaperone-binding sites, or also lead to global rearrangements, changing aggregate size and total

exposed surface area, e. g. through partial aggregate dissolution and fragmentation. To shed light on this,

we visualised the aggregates of luciferase C-terminally fused with GFP using fluorescence microscopy.

After heat denaturation, approximately 80% of luciferase-GFP aggregates had the size of 2 µm or more

(Figure 2B). Incubation with Ssa1-Sis1 for 1 h decreased the fraction of aggregates larger than 2 µm to

25,8%, whereas the addition of Sse1 reduced it to 12,5% (Figure 2B). At the same time, the activity

remained at the level of 1-2% of the native luciferase-GFP and only the presence of Hsp104-Hsp70 led to

its substantial recovery (Figure 2—figure supplement 1A). The reduced size of luciferase-GFP aggregates

by Hsp70 depended on ATP and the aggregate size was only moderately affected by Ssa1-Ydj1,

irrespectively of the presence of Sse1 (Figure 2B).

We also measured the size of aggregates using dynamic light scattering (DLS). Heat-denatured luciferase

formed aggregates with hydrodynamic diameter of approximately 2000 nm (Figure 2—figure supplement

1B), while the DLS signal of approximately 10 nm corresponded to the native luciferase and chaperones,

consistent with their theoretical size (Figure 2—figure supplement 1B). Incubation with Ssa1-Sis1 for 1 h

slightly reduced the 2000 nm peak and produced small-size aggregates of approximately 30 nm

(Figure 2—figure supplement 1B). The presence of Sse1 increased the amount of 30 nm aggregates more

than 2-fold and significantly reduced the signal at 2000 nm (Figure 2—figure supplement 1B). The change in aggregate size was not observed unless ATP was present (Figure 2—figure supplement 1B). In agreement with the microscopy data, Ssa1-Ydj1 with or without Sse1 did not have any considerable effect on the size of the aggregates (Figure 2—figure supplement 1B).

Taken together, Sse1 boosts the aggregate-remodelling activity of Ssa1, specifically with class B JDP Sis1, causing major reduction in aggregate size.

Hormetic effects of Sse1 in Hsp70 disaggregation activity

Contrary to the strong stimulation by Hsp110 of the Hsp70 system comprising class B JDP, its impact on Hsp70 with class A JDP strongly varied with experimental setup (Figure 1A,B, Figure 1—figure supplement 1A, B, C). Knowing that the effects of Sse1 on the disaggregation by Ssa1-Ydj1 are concentration-dependent, with stimulation at substoichiometric amount of Sse1 (Dragovic *et al.*, 2006), we asked, to what degree the optimum Hsp110 level depends on the class of a JDP. When we titrated the Hsp70 system comprising either Sis1 or Ydj1, we observed an inhibition of luciferase disaggregation at increased Sse1 concentrations, yet the system with Ydj1 was much more sensitive to Sse1 (IC50 = 0.1 µM), while Ssa1-Sis1 was still stimulated at 0.3 µM Sse1 (molar ratio Ssa1:Sse1 1:0.3) (Figure 3A). When we included the Hsp104 disaggregase, both Hsp104-Ssa1-Ydj1 and Hsp104-Ssa1-Sis1 systems were stimulated at low and inhibited at high Sse1 levels, with the highest yield at 0.05 µM and 0.2 µM of Sse1, respectively (Figure 3—figure supplement 1A). A similar biphasic effect was observed for human orthologs, with Hsc70-DNAJB4 tolerating higher Hsp105 concentration than Hsc70-DNAJA2 (Figure 3B).

In contrast to protein disaggregation, the reactivation of denatured, non-aggregated luciferase by Ssa1 with either Sis1 or Ydj1 was only negatively affected across a range of Sse1 concentrations (Figure 3C), suggesting that the positive contribution of Hsp110 takes place before polypeptides get released from an aggregate. The negative effect of Sse1 on protein folding could potentially mask a positive effect of saturating Sse1 on aggregate modification, e. g. through entropic pulling. We thus analysed the size of aggregates after their incubation with Ssa1-Sis1 and 1 µM Sse1, however it was unaffected (Figure 3—figure supplement 1B, C), suggesting that higher Sse1 levels also limit the aggregate-remodelling activity of Hsp70. Consistently, when we measured Hsp70 association with luciferase aggregated on the BLI sensor across Sse1 concentrations, we observed the biphasic effect for Ssa1-Ydj1 and Ssa1-Sis1, with the maximum binding at 0.05 µM and 0.2 µM of Sse1, respectively (Figure 3D). Thus, Hsp110 promotes Hsp70 assembly at aggregates and their modification only at sub-stoichiometric concentrations, with peak performance at higher levels of Sse1 in the presence of class B than class A JDPs.

To establish whether the inhibition that occurs at higher Sse1 concentrations is directly associated with the interaction between Sse1 and Ssa1, or rather between Sse1 and a substrate, we used the previously

characterised Sse1-2 variant (N572Y E575A), with disrupted interaction with the A300 residue of Ssa1, which reduces Hsp70 binding to 20% and nucleotide exchange to 5% of that of the wild type (Polier *et al*, 2008). The variant has been reported to partially compensate for the *SSE1/SSE2* double deletion (Polier *et al.*, 2008). Consistently with the weak Ssa1 binding affinity of Sse1-2 (Figure 3—figure supplement 1D), Ssa1-Sis1 required 20-times higher concentration of the Sse1-2 variant than Sse1 WT to reach the same GFP disaggregation activity and no inhibition was observed up to 2 µM of Sse1-2 (Figure 3E). Also, the Sse1-2 variant had a much weaker impact on luciferase disaggregation (Figure 3—figure supplement 1E). Nonetheless, 1 µM Sse1-2 was sufficient to obtain a similar effect as 0.1 µM WT Sse1 in stimulating Ssa1-Sis1 binding to the aggregated luciferase in the BLI assay (Figure 3—figure supplement 1F).

This suggests that it is the high-affinity interaction between Hsp70 and the NEF that enables strong stimulation already at very low Hsp110 levels. On the other hand, such strong interaction is associated with the inhibition of the disaggregation activity when Hsp110 concentration exceeds sub-stoichiometric proportion to Hsp70.

Hsp110 limits JDP interaction with Hsp70

Since all NEFs induce polypeptide release from Hsp70, excessive dissociation from the substrate seems the most apparent explanation of Hsp70 inhibition by Hsp110. However, the strong dependence of the impact of Hsp110 on the class of JDP prompted us to search for other potential mechanisms behind the low Hsp70 chaperone activity under high Hsp110 concentration. Knowing that the sensitivity to Sse1 (Figure 3A, D) is negatively correlated with the strength of the Hsp70-JDP interaction, higher in the case class B JDP due to its binding of the EEVD motif of Hsp70 (Wyszkowski *et al.*, 2021), we explored the possibility that Hsp110 restricts the formation of the Hsp70-JDP complex.

To test this, we immobilised Sis1 on the BLI sensor and monitored Ssa1 binding across a range of Sse1 concentrations. The incubation with 0.1 µM Sse1 was almost inert, however, at 1:1 Ssa1:Sse1 ratio, the binding greatly diminished (Figure 4A). The reduced affinity of Sse1-2 for Ssa1 resulted in its milder negative effect on Ssa1 binding to Sis1 (Figure 4—figure supplement 1A), in agreement with its impact on disaggregation (Figure 3E). The degree of inhibition by the two Sse1 variants (Figure 4—figure supplement 1B) correlated with their capacity to exchange nucleotides (Dragovic *et al.*, 2006; Polier *et al.*, 2008).

In an analogous way, we analysed the interaction between the human Hsc70 and sensor-bound DNAJB4. With an increasing Hsp105 concentration, the binding signal declined to reach 85% at 1:1 Hsc70:Hsp105 ratio (Figure 4—figure supplement 1C), indicating that the negative Hsp110 impact on the Hsp70-JDP interaction exhibits a similar pattern across Fungi and Metazoa.

Since Sse1 inhibits Ssa1 binding to Sis1, we asked whether the disaggregation rate limitation imposed by Sse1 depends on the level of Sis1. To verify this, we incubated GFP aggregates with the Hsp70 system comprising various Sse1 and Sis1 concentrations (Figure 4B, C, Figure 4—figure supplement 1D). The susceptibility of the Hsp70 system to elevated Sse1 concentration significantly decreased with increasing Sis1 concentration and at 2 µM Sis1, the stimulation of the initial disaggregation rate was still observed at 2 µM Sse1 (Figure 4—figure supplement 1D). Interestingly, in the absence of Sse1, the disaggregation rate dropped with increasing Sis1 concentrations and the trend was reversed at higher Sse1 levels, with an inflection point at approximately 0.5 µM of Sse1 (Figure 4—figure supplement 1E).

Thus, Sse1 and Sis1 exhibit an apparent competition for Ssa1 binding. These results indicate that the balance between the JDPs and NEF co-chaperones is critical for the performance of the Hsp70 chaperone system in protein disaggregation.

Discussion

Our results uncover a comprehensive picture of the role of Hsp110 co-chaperones in aggregate processing by the Hsp70 system. Potentiation of the Hsp70 activity by the NEF strongly depends on the class of the J-domain protein and the preference for class B, conserved in yeast and human orthologs, relies on Hsp70 binding through the EEVD motif. Furthermore, we demonstrate that the Hsp110-dependent stimulation of disaggregation is limited to its initial stages: chaperone recruitment to aggregates and their disassembly, but not final protein folding (Figure 5). The initial Hsp110-dependent loading of more Hsp70 onto aggregates correlates with their remodelling into smaller aggregated species, which improves their recognition by the Hsp104 disaggregase. We also gained insight into the biphasic impact of Hsp110: with increasing Sse1 level, the stimulation is overshadowed by inhibition and the contribution of each trend depends on the phase of disaggregation and the composition of the disaggregating system, with a crucial role of the NEF's affinity for Hsp70. Finally, we show that the disaggregation inhibition by Hsp110 involves disruption of the Hsp70-JDP interaction, suggesting competition between JDP and Hsp110 co-chaperones.

The balance between the partners within the Hsp70 system is key for efficient disaggregation, as an individual co-chaperone with a critical function at one step may inhibit another. To investigate this differential contribution, we used biochemical assays reflecting individual stages of protein disaggregation. The results of the BLI experiments show that Hsp110 with class B JDP (JDP[B]) greatly increases Hsp70 aggregate binding (Figure 1B, C, Figure 1—figure supplement 1E, F, 2A), which correlates with the final reactivation yield (Figure 1A, Figure 1—figure supplement 1H), implying that the initiation phase determines the overall disaggregation efficiency. On the other hand, the final folding of a soluble substrate by Ssa1 relies much more on the class A JDP (JDP[A]) Ydj1 than on Sis1 (Figure 1—figure supplement 1D)

(Lu & Cyr, 1998), and not on Sse1 altogether (Figure 1—figure supplement 1D), which is in line with

previous reports that its human ortholog, Hsp105 inhibits luciferase folding (Rauch & Gestwicki, 2014).

The higher degree of stimulation with class B JDP has been previously observed, albeit without broader

insight, for Hsp110 and other NEF BAG (Rampelt et al., 2012; Rauch & Gestwicki, 2014). We show that

the distinguishing feature fundamental for this specificity is the auxiliary interaction between the CTD

domain of Sis1 and the EEVD motif of Hsp70. The Sis1 E50A variant, featuring derepressed J-domain

(Yu et al., 2015), together with Ssa1 ΔEEVD exhibited a similar aggregate binding pattern with or without

Sse1 to the system with Ydj1 (Figure 1B, Figure 1—figure supplement 1E, F, J). However, while Ydj1 can

bind misfolded proteins and prevent aggregation, Sis1 cannot (Lu & Cyr, 1998). When Sis1 is deprived of

Ssa1 binding through EEVD, which has a major contribution to the stability of their interaction

(Wyszkowski et al., 2021), the effect of Sse1 on disaggregation is detrimental (Figure 1—figure

supplement 1I). This underlines the importance of the Hsp70-JDP complex stability for the stimulation of

substrate binding and disaggregation by Sse1.

In our previous work, we reported sigmoidal aggregate binding kinetics characteristic for the Ssa1-Sis1

system, suggesting that the binding gradually generates more chaperone binding sites (Wyszkowski et al.,

2021). Now, we show that this apparent cooperativity is conserved in yeast and human, implying

mechanistic similarities between the systems (Figure 1B, C). While the s-shaped Ssa1-Sis1 binding to the

aggregate results in the association of more Ssa1 than with Ydj1 (Figure 1—figure supplement 2A),

Hsp110 shortens the initial lag phase and boosts the binding efficacy, loading a thicker layer of Hsp70

molecules onto aggregates (Figure 1B,C, Figure 1—figure supplement 2A), similarly as has been shown

for amyloid fibrils (Beton et al., 2022). This corresponds with Hsc70 clustering on amyloid in the presence

of Hsp110, reported by Wentink and co-workers (2020). Clustered, densely packed Hsp70 has been

proposed to generate entropic pulling effect that leads to fibril fragmentation (Goloubinoff & De Los Rios,

2007; Sousa & Lafer, 2019; Wentink et al., 2020). Likewise, we observed remodelling of amorphous

aggregates into smaller aggregated species, which resembles amyloid disassembly in terms of the

dependence on the NEF activity of Hsp110, the class B JDP and its interaction with the Hsp70 EEVD motif

(Figure 2B, Figure 2—figure supplement 1B) (Beton et al, 2022). We propose that the local extraction of

polypeptides by Hsp70-JDP[B]-Hsp110, which penetrates the surface and takes apart amorphous

aggregates, although not enough for full protein recovery, could successfully generate more manageable

substrates for chaperones, as we demonstrated by applying the derepressed Hsp104 variant

(Figure 2A, 5). This Hsp70 activity, albeit yielding very low protein reactivation (Figure 2—figure

supplement 1A), might be relevant for the fragmentation of a thinner, linear amyloid structure.

Such clustering-augmented aggregate remodelling and fragmentation, possibly through expanding the

total effective aggregate surface, leads to the much improved protein reactivation by Hsp104 (Figure 1—

figure supplement 1B, C, Figure 2—figure supplement 1A). This would explain the previous findings that

although Sse1 is not necessary for Hsp104 recruitment to aggregates, it is essential for Hsp104-dependent

aggregate clearance in the cell (Kaimal et al., 2017). Apparently, despite the more heterogenic and

dynamic nature of cellular aggregates than of those generated *in vitro*, their disaggregation by Hsp104

largely relies on the Hsp110-boosted processing by the Hsp70 system.

Above a certain Hsp110 level, however, the disaggregation is inhibited. The susceptibility to the NEF

depends on the phase of the process and the composition of the Hsp70 system (Figure 3A-D, Figure 3—

figure supplement 1A), which may explain the discrepancies between different studies of Hsp110's impact

on disaggregation (Dragovic *et al.*, 2006; Polier *et al.*, 2008; Raviol *et al.*, 2006; Shorter, 2011) As well

established, the most effective is substoichiometric Hsp110 proportion to Hsp70 (Gao *et al*, 2015; Polier

et al., 2008; Wentink *et al.*, 2020) which corresponds with the physiological conditions, where the ratio

between all cytoplasmic Hsp70 and Hsp110 paralogs involved in disaggregation ranges from

approximately 10 : 1.7 in yeast (Brownridge *et al*, 2013) to 10 : 2.4 in humans (Finka & Goloubinoff, 2013).

According to our data, the optimum, at which the stimulation and inhibition curves intersect, is shifted to

higher levels: a) during the initiation versus final protein folding (Figure 3A, C), (b) with class B JDP, in

comparison with class A (Figure 3A, B, D), c) with Hsp104, comparing with the system without the

disaggregase (Figure 3A, Figure 3—figure supplement 1A) and d) for the human Hsp70 system, comparing

with the yeast chaperones (Figure 3A, B). The latter trend might result from the stronger dependence of

the human Hsp70 chaperone on the nucleotide exchange activity, possibly associated with the lower basal

rate of nucleotide dissociation than that reported for Ssa1 (Dragovic et al., 2006). Possibly, due to these

problems with ADP release, the positive impact of Hsp110 on Hsc70 is manifested both in the presence

of class A and B JDPs and the inhibition occurs at much higher Hsp110 levels. Accordingly, it is tempting

to speculate that the human system, lacking Hsp104, has evolved towards better tolerance for Hsp110 to

boost the disaggregation activity of the Hsp70 system alone, e.g. to counteract amyloid toxicity.

Excessive dissociation of Hsp70 from a substrate, promoted by nucleotide exchange, may explain the

system's susceptibility to Hsp110. Here, we propose an additional mechanism behind this inhibition. The

fact that the NEF and JDP binding occurs at mutually exclusive nucleotide states of Hsp70 has raised a

question of competition between the two co-chaperones (Sousa & Lafer, 2019), which, to our knowledge,

has never been addressed before. On the other hand, class B JDP goes beyond the classical model of

the Hsp70 cycle due to the auxiliary interaction with the EEVD motif, which is not nucleotide-sensitive and

theoretically might not be exclusive with Hsp110 binding (Wyszkowski *et al.*, 2021; Yu *et al.*, 2015).

Nonetheless, our results clearly demonstrate that Hsp110 disrupts the JDP's complex with Hsp70

(Figure 4A, Figure 4—figure supplement 1C). This way, Hsp110 may cause the release of Hsp70 from an

aggregate, as well as of a JDP, unless it remains tethered via another interaction, e. g. through the other

JDP subunit in complex with an aggregate-bound Hsp70 molecule (Figure 5). The fate of the displaced

proteins is yet to be established, presumably at low Hsp110 levels, they may re-bind at another site on the

substrate or to the aggregate-bound chaperones. At high Hsp110 levels, the JDP[B]-Hsp70 interaction is very strongly affected, similarly as aggregate binding and disaggregation (Figure 3A, B, D, 4A, B), suggesting that the proteins released by Hsp110 do not re-bind or that any subsequent binding is non-productive.

The apparent competition between the co-chaperones is reflected in the fact that the inhibition by Hsp110 is moderated at increasing Sis1 level (Figure 4C). Interestingly, the class B JDP also exerts a biphasic effect on disaggregation, and optimal concentrations of the two co-chaperones are strongly interdependent (Figure 4B, C, Figure 4—figure supplement 1D, E). The highest disaggregation yield occurs at Ssa1 : Sis1 : Sse1 ratio closest to the average reported in the cytosol, 10 : 0.3 : 1.7 (Figure 4C) (Brownridge *et al.*, 2013), pointing to a possibility that the proportions between the co-chaperones have undergone evolutionary fine-tuning to develop higher tolerance to stress.

The sub-stoichiometric optimum is unique to Hsp110 (Gao *et al.*, 2015; Wentink *et al.*, 2020), while other NEF, Fes1, requires 10 times higher level to reach the same degree of stimulation, similarly as the Sse1-2 variant with reduced affinity for Hsp70. Fes1's ability to improve Hsp70 loading onto aggregate is surprising, regarding findings of Wentink et al., (2020) who demonstrated, using Hsp110 and BAG1 variants, that Hsp70 clustering on amyloid requires a bulky NEF. Although Fes1 is only 33 kDa, the stretch of its armadillo repeats and different mode of interaction with NBD could possibly lead to similar excluded volume effects as Hsp110. Nevertheless, Fes1 is present in a cell at approximately one-fifth the level of Sse1, and even its 8.4-fold overexpression from 2 plasmids, shown by Kaimal et al., was insufficient to complement the thermosensitive phenotype of *sse1-200 sse2Δ* (Kaimal *et al.*, 2017), as it was 6 times too low to reach the level effective in disaggregation *in vitro* (Figure 1—figure supplement 3A). It is worth noting that Fes1 has been reported to target substrates to degradation (Gowda *et al.*, 2018; Gowda *et al*, 2013) and apparently its role in Hsp70 recruitment to disaggregation is minor.

An abundant association of Hsp70-JDP[B] with aggregate surface, although generating strong pulling effect to disentangle polypeptides, could also mask access of chaperones to the newly emerging sites adjacent to and buried beneath the complex. We demonstrate that Hsp110 could theoretically uncover such sites not only through Hsp70 dissociation from the substrate due to the nucleotide exchange (Dragovic et al., 2006), but also by disrupting the JDP[B]-Hsp70 interaction (Figure 5). Four Hsp70 binding sites in a JDP[B] dimer allow to form an extensive network of interactions (Wyszkowski *et al.*, 2021). We speculate that Hsp110 introduces plasticity into this network, enabling the chaperone complex to infiltrate cavities emerging within the aggregate in a fluid-like manner, binding to the uncovered chaperone-binding sites and pulling up to the aggregate fragmentation (Figure 5). Such behaviour would require only a small destabilising effect of the NEF that would not dissociate the complex completely (Figure 4A), similarly as observed at the Hsp110 concentration that supports the most effective aggregate binding and disaggregation (Figure 3A, D). Further experimental verification of this scenario and future studies of the

1 dynamics within the Hsp70 chaperone system will be critical to understand and combat the stress- and

2 disease- related protein aggregation.

Materials and methods

Proteins

Sse1 (Andreasson *et al.*, 2008a), Ssa1 (Andreasson *et al.*, 2008a), Sis1 (Wyszkowski *et al.*, 2021), Ydj1

(Lipinska *et al*, 2013), Hsp104 (Lipinska *et al.*, 2013), His-tagged luciferase (Chamera *et al.*, 2019), GFP

(Zietkiewicz *et al.*, 2004) were purified using published protocols. The same protocol as for Sse1 was used

for Fes1 (Andreasson *et al.*, 2008a). DNAJB4, DNAJA2, Hsc70 and Hsp105 were purified as described in

the work by Nillegoda et al. (2015). To obtain His-tagged chaperones, a step of proteolytic cleavage of the

tag was omitted. Sse1 N572Y N575A was constructed by introduction of point mutations using PCR-

specific mutagenesis (Qiagen) and confirmed with sequencing. Fluc-EGFP and EGFP (parent vector

pCIneo-Fluc-EGFP) were cloned into pET22b plasmid. Fluc-EGFP and His-tagged EGFP were purified

using Ni-NTA agarose (Protino), followed by anion exchange chromatography using Q-Sepharose (Q

Sepharose Fast Flow, GE Healthcare). Untagged luciferase and its substrate were purchased from

Promega (E1702).

Heat-aggregated luciferase and luciferase-GFP reactivation

Luciferase or luciferase-GFP (30 µM) was denatured in the buffer A (25 mM HEPES-KOH pH 8.0, 75 mM

KCl and 15 mM $MgCl_2$) supplemented with 6 M Urea at 25 °C for 15 minutes. Next, it was transferred to

48 °C for 10 minutes and subsequently 25-fold diluted into the buffer A. After 15 minutes of incubation at

25 °C, the reactions were initiated by the addition of the mix of chaperones, which were used at the

following concentrations: 1 µM Ssa1, 1 µM Sis1, 1 µM Ydj1, 0.1 µM Sse1, 1 µM Fes1, 3 µM Hsc70, 1 µM

DNAJB4, 1 µM DNAJA2 and 0.3 µM Hsp105, if not stated otherwise. Luminescence was measured using

Sirius Luminometer (Berthold).

Unfolded luciferase reactivation

25 Method for spontaneous folding of luciferase was adapted from (Imamoglu *et al.*, 2020). Briefly, 10 µM of

26 luciferase was denatured in 5 M GuHCl and 10 mM DTT at 25 °C for 1 hour. To initiate the spontaneous

27 folding, the luciferase was 100-fold diluted into the folding buffer (25 mM HEPES-KOH pH 7.5,

28 100 mM KCl, 10 mM Mg(OAc)$_2$, 2 mM DTT, 0.05% TWEEN20). Folding was performed with chaperones

29 used at 1 µM concentration, except for 0.1 µM Sse1, unless it was stated otherwise.

Renaturation of heat-aggregated GFP

Recovery of GFP aggregates was performed as previously described (Zietkiewicz *et al.*, 2004). Briefly,

GFP (74 µM) was thermally inactivated at 85 °C for 15 min. The reactivation reaction was carried out at

25 °C in the renaturation buffer (25 mM HEPES-KOH (pH 8.0), 7% (v/v) glycerol, 60 mM potassium

glutamate, 7 mM DTT, 15 mM MgOAc, 10 mM ATP), with an ATP regeneration system comprising 1.2 µM

creatine kinase and 20 mM creatine phosphate. Disaggregation of the 100-times diluted GFP aggregates

was initiated by adding the chaperone proteins at the following concentrations: Ssa1 (1 µM), Ydj1 (1 µM),

Sis1 (1 µM), Sse1 (0.1 µM), Hsp104 WT (1 µM), Hsp104^{mut} (0.15 µM), unless indicated otherwise. GFP

fluorescence was detected in a Beckman Coulter DTX880 microplate reader.

Fluorescent transmitted light microscopy

Luciferase-GFP (14.6 µM) was incubated in the buffer A with 6-M Urea at 25 °C for 15 min. Then, it was

transferred to 48 °C for 10 minutes and 10-fold diluted into the buffer A containing 5 mM ATP and 2 mM

DTT. After 15 minutes of incubation at 25°C, the reaction was initiated upon addition of the mix of

chaperones at 1 µM concentration, except for Sse1 used at 0.1 µM. The final concentration of

luciferase-GFP aggregates in the reaction was 0.3 µM. After 1 hour of incubation with the chaperones, the

reaction was arrested upon addition of 200 mM NaCl and transferred on ice. Specimens were imaged

using a confocal laser scanning microscope Leica SP8X with a 100× oil immersion lens (Leica, Germany).

Presented data show results from three independent experiments. Each sample within the repeat was

photographed ten times. Data analysis was performed with Leica LAS X software.

Dynamic light scattering

Luciferase (23,7 µM) was incubated in the buffer A with 6 M Urea at 25 °C for 15 min, then it was transferred

to 48 °C for 10 min. Subsequently, it was 10x diluted with the buffer A containing 5 mM ATP and 1 mM

DTT and incubated for 15 min at 25 °C. Then, it was subjected to a DLS measurement to confirm the

presence of the aggregates, size of which ranged between 1000 and 3000 nm. The reaction was initiated

upon an addition of a mix of chaperones used at 1 µM concentration, except for Sse1 used at 0.1 µM. The

final concentration of luciferase was 0.6 µM. Dynamic Light Scattering was measured using the ZetaSizer

NanoS instrument (Malvern) after 1 h of incubation with chaperones at 25 °C. For each sample, three

independent measurements were performed, and particle size distributions were calculated as percent

within a range between 0.4 and 10,000 nm.

Bio-Layer Interferometry Experiments

All the BLI experiments described below were performed as previously described (Wyszkowski *et al.*,

2021) using the BLItz and Octet K2 instruments.

Binding of chaperones to luciferase aggregates

Initially, the Ni-NTA biosensor (ForteBio Dip and Read) was hydrated in the buffer A for 10 minutes and

subsequently immersed in the same buffer containing 6 M Urea and 8.2 μM His-tagged luciferase for 10

minutes, resulting in an anchoring layer of ~6 nm. Next, the biosensor was washed with the buffer A and

immersed in the buffer A with 1.6 μM of native His-tagged luciferase and incubated for 10 minutes at

44°C. Finally, the biosensor was equilibrated with the buffer A supplemented with 5 mM ATP and 2 mM

DTT for 5 minutes. The layer of luciferase aggregates reached ~16 nm and was stable. Binding and

dissociation of chaperones were performed in the buffer A supplemented with 5 mM ATP and 2 mM DTT

at 25 °C.

Binding of chaperones to the GFP aggregates

To prepare the sensor with GFP aggregates, after the initial hydration of the Ni-NTA biosensor (ForteBio

Dip and Read) in the buffer A (25 mM HEPES-KOH pH 8.0, 75 mM KCl and 15 mM $MgCl_2$) for 10 minutes,

it was immersed in the buffer A with 9 M urea containing 12.5 μM of His-tagged GFP and incubated at

85 °C for 15 minutes. After washing with the buffer A for 5 minutes, it was immersed in the buffer A

containing 4.2 μM of His-tagged GFP and incubated at 85 °C. Next, the sensor was equilibrated with the

buffer A supplemented with 5 mM ATP and 2 mM DTT for 5 minutes. The aggregate layer thickness

was ~40 nm. Association and dissociation of chaperones were performed in the same buffer at 25 °C.

Binding of chaperones to the immobilized yeast lysate

Yeast lysate was prepared from overnight culture of W303 yeasts in the YPD medium, according to

the published protocol (Wyszkowski *et al.*, 2021). The Ni-NTA biosensor (ForteBio Dip and Read) was

initially hydrated in the buffer A for 10 minutes, following incubation in the same buffer supplemented with

6 M Urea and 8.2 μM His-tagged luciferase for 10 minutes, which resulted in biolayer thickness of

~6 nm. Next, the biosensor was washed with the buffer A and immersed in the same buffer containing

soluble yeast proteins (5 mg/ml). After 15 minutes of incubation at 55 °C, the biosensor was

equilibrated with the buffer A supplemented with 5 mM ATP and 2 mM DTT, reaching biolayer

thickness of ~30 nm. Association and dissociation of chaperones were performed in the same buffer

at 25 °C.

BLI with fluorescently labelled protein

Ssa1 was incubated with 10x molar excess of Alexa Fluor 488 C5-maleimide (Invitrogen) for 2 h at 4 °C.

An excess of the label was removed with a desalting column (PD-10, GE Healthcare). The BLI experiment

was performed as with unlabelled proteins and the fluorescence of the dissociated Ssa1 (A488*) was

measured using Beckman Coulter DTX 880.

BLI of direct protein-protein interactions

The biosensor was initially hydrated in the buffer A for 10 minutes. Next, it was immersed in the buffer A containing the indicated protein (0.4 μM His-Sis1, 2.5 μM His-Sse1 or 1 μM His-Sse1-2) until binding reached saturation, which corresponded to a layer thickness of ~15 nm for His-Sis1, ~6 nm for His-See1 and ~4 nm for His-Sse1-2. Subsequently, it was washed with the buffer A supplemented with 5 mM ATP and 2 mM DTT for 5 minutes and then immersed in the same buffer with the mix of chaperones at the indicated concentrations.

Acknowledgements

We thank Bernd Bukau (Heidelberg University), Kevin A. Morano (UTHealth Houston), F. Ulrich Hartl (Max Planck Institute of Biochemistry) and Claes Andréasson (Stockholm University) for sharing plasmids. We thank Katarzyna Bury, Gabriel Petelski, Maciej Małolepszy, Katarzyna Kalinowska and Dominik Purzycki for technical suport. This work was supported by a grant of the Polish National Science Centre (2019/35/B/NZ1/01475).

References

Abrams JL, Verghese J, Gibney PA, Morano KA (2014) Hierarchical functional specificity of cytosolic heat shock protein 70 (Hsp70) nucleotide exchange factors in yeast. *J Biol Chem* 289: 13155-13167 doi: 10.1074/jbc.M113.530014

Andreasson C, Fiaux J, Rampelt H, Druffel-Augustin S, Bukau B (2008a) Insights into the structural dynamics of the Hsp110-Hsp70 interaction reveal the mechanism for nucleotide exchange activity. *Proc Natl Acad Sci U S A* 105: 16519-16524 doi: 10.1073/pnas.0804187105

Andreasson C, Fiaux J, Rampelt H, Mayer MP, Bukau B (2008b) Hsp110 is a nucleotide-activated exchange factor for Hsp70. *J Biol Chem* 283: 8877-8884 doi: 10.1074/jbc.M710063200

Beton JG, Monistrol J, Wentink A, Johnston EC, Roberts AJ, Bukau BG, Hoogenboom BW, Saibil HR (2022) Cooperative amyloid fibre binding and disassembly by the Hsp70 disaggregase. *EMBO J* 41: e110410 doi: 10.15252/embj.2021110410

Brownridge P, Lawless C, Payapilly AB, Lanthaler K, Holman SW, Harman VM, Grant CM, Beynon RJ, Hubbard SJ (2013) Quantitative analysis of chaperone network throughput in budding yeast. *Proteomics* 13: 1276-1291 doi: 10.1002/pmic.201200412

Chamera T, Klosowska A, Janta A, Wyszkowski H, Obuchowski I, Gumowski K, Liberek K (2019) Selective Hsp70-Dependent Docking of Hsp104 to Protein Aggregates Protects the Cell from the Toxicity of the Disaggregase. *Journal of Molecular Biology* 431: 2180-2196 doi: 10.1016/j.jmb.2019.04.014

Dragovic Z, Broadley SA, Shomura Y, Bracher A, Hartl FU (2006) Molecular chaperones of the Hsp110 family act as nucleotide exchange factors of Hsp70s. *EMBO J* 25: 2519-2528 doi: 10.1038/sj.emboj.7601138

Easton DP, Kaneko Y, Subjeck JR (2000) The hsp110 and Grp1 70 stress proteins: newly recognized relatives of the Hsp70s. *Cell Stress Chaperones* 5: 276-290 doi: 10.1379/1466-1268(2000)005<0276:thagsp>2.0.co;2

Faust O, Abayev-Avraham M, Wentink AS, Maurer M, Nillegoda NB, London N, Bukau B, Rosenzweig R
(2020) HSP40 proteins use class-specific regulation to drive HSP70 functional diversity. *Nature* 587: 489-
+ doi: 10.1038/s41586-020-2906-4

Finka A, Goloubinoff P (2013) Proteomic data from human cell cultures refine mechanisms of chaperone-
mediated protein homeostasis. *Cell Stress Chaperones* 18: 591-605 doi: 10.1007/s12192-013-0413-3

Gao X, Carroni M, Nussbaum-Krammer C, Mogk A, Nillegoda NB, Szlachcic A, Guilbride DL, Saibil HR,
Mayer MP, Bukau B (2015) Human Hsp70 Disaggregase Reverses Parkinson's-Linked alpha-Synuclein
Amyloid Fibrils. *Mol Cell* 59: 781-793 doi: 10.1016/j.molcel.2015.07.012

Garcia VM, Nillegoda NB, Bukau B, Morano KA (2017) Substrate binding by the yeast Hsp110 nucleotide
exchange factor and molecular chaperone Sse1 is not obligate for its biological activities. *Mol Biol Cell* 28:
2066-2075 doi: 10.1091/mbc.E17-01-0070

Glover JR, Lindquist S (1998) Hsp104, Hsp70, and Hsp40: a novel chaperone system that rescues
previously aggregated proteins. *Cell* 94: 73-82 doi:

Goeckeler JL, Petruso AP, Aguirre J, Clement CC, Chiosis G, Brodsky JL (2008) The yeast Hsp110,
Sse1p, exhibits high-affinity peptide binding. *FEBS Lett* 582: 2393-2396 doi: 10.1016/j.febslet.2008.05.047

Goeckeler JL, Stephens A, Lee P, Caplan AJ, Brodsky JL (2002) Overexpression of yeast Hsp110
homolog Sse1p suppresses ydj1-151 thermosensitivity and restores Hsp90-dependent activity. *Mol Biol
Cell* 13: 2760-2770 doi: 10.1091/mbc.02-04-0051

Goloubinoff P, De Los Rios P (2007) The mechanism of Hsp70 chaperones: (entropic) pulling the models
together. *Trends in Biochemical Sciences* 32: 372-380 doi: 10.1016/j.tibs.2007.06.008

Gowda NKC, Kaimal JM, Kityk R, Daniel C, Liebau J, Öhman M, Mayer MP, Andréasson C (2018)
Nucleotide exchange factors Fes1 and HspBP1 mimic substrate to release misfolded proteins from Hsp70.
Nature structural & molecular biology 25: 83-89 doi: 10.1038/s41594-017-0008-2

Gowda NKC, Kandasamy G, Froehlich MS, Dohmen RJ, Andréasson C (2013) Hsp70 nucleotide
exchange factor Fes1 is essential for ubiquitin-dependent degradation of misfolded cytosolic proteins.
Proceedings of the National Academy of Sciences 110: 5975-5980 doi: doi:10.1073/pnas.1216778110

Hartl FU, Bracher A, Hayer-Hartl M (2011) Molecular chaperones in protein folding and proteostasis.
Nature 475: 324-332 doi: 10.1038/nature10317

Imamoglu R, Balchin D, Hayer-Hartl M, Hartl FU (2020) Bacterial Hsp70 resolves misfolded states and
accelerates productive folding of a multi-domain protein. *Nat Commun* 11: 365 doi: 10.1038/s41467-019-
14245-4

Kaimal JM, Kandasamy G, Gasser F, Andreasson C (2017) Coordinated Hsp110 and Hsp104 Activities
Power Protein Disaggregation in Saccharomyces cerevisiae. *Molecular and Cellular Biology* 37UNSP
e00027-17
10.1128/MCB.00027-17

Laufen T, Mayer MP, Beisel C, Klostermeier D, Mogk A, Reinstein J, Bukau B (1999) Mechanism of
regulation of hsp70 chaperones by DnaJ cochaperones. *Proc Natl Acad Sci U S A* 96: 5452-5457 doi:
10.1073/pnas.96.10.5452

Liberek K, Marszalek J, Ang D, Georgopoulos C, Zylicz M (1991) Escherichia-Coli Dnaj and Grpe Heat-
Shock Proteins Jointly Stimulate Atpase Activity of Dnak. *P Natl Acad Sci USA* 88: 2874-2878 doi: DOI
10.1073/pnas.88.7.2874

Lipinska N, Zietkiewicz S, Sobczak A, Jurczyk A, Potocki W, Morawiec E, Wawrzycka A, Gumowski K, Slusarz M, Rodziewicz-Motowidlo S *et al* (2013) Disruption of ionic interactions between the nucleotide binding domain 1 (NBD1) and middle (M) domain in Hsp100 disaggregase unleashes toxic hyperactivity and partial independence from Hsp70. *J Biol Chem* 288: 2857-2869 doi: 10.1074/jbc.M112.387589

Liu Q, Hendrickson WA (2007) Insights into Hsp70 chaperone activity from a crystal structure of the yeast Hsp110 Sse1. *Cell* 131: 106-120 doi: 10.1016/j.cell.2007.08.039

Lu Z, Cyr DM (1998) Protein folding activity of Hsp70 is modified differentially by the Hsp40 co-chaperones Sis1 and Ydj1. *Journal of Biological Chemistry* 273: 27824-27830 doi: DOI 10.1074/jbc.273.43.27824

Lum R, Tkach JM, Vierling E, Glover JR (2004) Evidence for an unfolding/threading mechanism for protein disaggregation by Saccharomyces cerevisiae Hsp104. *J Biol Chem* 279: 29139-29146 doi: 10.1074/jbc.M403777200

Mattoo RUH, Sharma SK, Priya S, Finka A, Goloubinoff P (2013) Hsp110 is a bona fide chaperone using ATP to unfold stable misfolded polypeptides and reciprocally collaborate with Hsp70 to solubilize protein aggregates. *J Biol Chem* 288: 21399-21411 doi: 10.1074/jbc.M113.479253

Mayer MP, Bukau B (2005) Hsp70 chaperones: cellular functions and molecular mechanism. *Cell Mol Life Sci* 62: 670-684 doi: 10.1007/s00018-004-4464-6

Mayer MP, Gierasch LM (2019) Recent advances in the structural and mechanistic aspects of Hsp70 molecular chaperones. *Journal of Biological Chemistry* 294: 2085-2097 doi: https://doi.org/10.1074/jbc.REV118.002810

Morimoto RI (2008) Proteotoxic stress and inducible chaperone networks in neurodegenerative disease and aging. *Genes Dev* 22: 1427-1438 doi: 10.1101/gad.1657108

Nillegoda NB, Kirstein J, Szlachcic A, Berynskyy M, Stank A, Stengel F, Arnsburg K, Gao XC, Scior A, Aebersold R *et al* (2015) Crucial HSP70 co-chaperone complex unlocks metazoan protein disaggregation. *Nature* 524: 247-+ doi: 10.1038/nature14884

Oh HJ, Chen X, Subjeck JR (1997) Hsp110 protects heat-denatured proteins and confers cellular thermoresistance. *J Biol Chem* 272: 31636-31640 doi: 10.1074/jbc.272.50.31636

Oh HJ, Easton D, Murawski M, Kaneko Y, Subjeck JR (1999) The chaperoning activity of hsp110. Identification of functional domains by use of targeted deletions. *J Biol Chem* 274: 15712-15718 doi: 10.1074/jbc.274.22.15712

Polier S, Dragovic Z, Hartl FU, Bracher A (2008) Structural basis for the cooperation of Hsp70 and Hsp110 chaperones in protein folding. *Cell* 133: 1068-1079 doi: 10.1016/j.cell.2008.05.022

Polier S, Hartl FU, Bracher A (2010) Interaction of the Hsp110 molecular chaperones from S. cerevisiae with substrate protein. *J Mol Biol* 401: 696-707 doi: 10.1016/j.jmb.2010.07.004

Rampelt H, Kirstein-Miles J, Nillegoda NB, Chi K, Scholz SR, Morimoto RI, Bukau B (2012) Metazoan Hsp70 machines use Hsp110 to power protein disaggregation. *Embo Journal* 31: 4221-4235 doi: 10.1038/emboj.2012.264

Rauch JN, Gestwicki JE (2014) Binding of human nucleotide exchange factors to heat shock protein 70 (Hsp70) generates functionally distinct complexes in vitro. *J Biol Chem* 289: 1402-1414 doi: 10.1074/jbc.M113.521997

1 Raviol H, Sadlish H, Rodriguez F, Mayer MP, Bukau B (2006) Chaperone network in the yeast cytosol:
2 Hsp110 is revealed as an Hsp70 nucleotide exchange factor. *EMBO J* 25: 2510-2518 doi:
3 10.1038/sj.emboj.7601139
4
5 Rohland L, Kityk R, Smalinskaite L, Mayer MP (2022) Conformational dynamics of the Hsp70 chaperone
6 throughout key steps of its ATPase cycle. *Proc Natl Acad Sci U S A* 119: e2123238119 doi:
7 10.1073/pnas.2123238119
8
9 Schaupp A, Marcinowski M, Grimminger V, Bosl B, Walter S (2007) Processing of proteins by the
10 molecular chaperone Hsp104. *J Mol Biol* 370: 674-686 doi: 10.1016/j.jmb.2007.04.070
11
12 Seyffer F, Kummer E, Oguchi Y, Winkler J, Kumar M, Zahn R, Sourjik V, Bukau B, Mogk A (2012) Hsp70
13 proteins bind Hsp100 regulatory M domains to activate AAA+ disaggregase at aggregate surfaces. *Nature*
14 *structural & molecular biology* 19: 1347-1355 doi: 10.1038/nsmb.2442
15
16 Shaner L, Sousa R, Morano KA (2006) Characterization of Hsp70 binding and nucleotide exchange by the
17 yeast Hsp110 chaperone Sse1. *Biochemistry* 45: 15075-15084 doi: 10.1021/bi061279k
18
19 Shaner L, Wegele H, Buchner J, Morano KA (2005) The yeast Hsp110 Sse1 functionally interacts with the
20 Hsp70 chaperones Ssa and Ssb. *J Biol Chem* 280: 41262-41269 doi: 10.1074/jbc.M503614200
21
22 Shorter J (2011) The mammalian disaggregase machinery: Hsp110 synergizes with Hsp70 and Hsp40 to
23 catalyze protein disaggregation and reactivation in a cell-free system. *Plos One* 6: e26319 doi:
24 10.1371/journal.pone.0026319
25
26 Sousa R, Lafer EM (2019) The Physics of Entropic Pulling: A Novel Model for the Hsp70 Motor Mechanism.
27 *Int J Mol Sci* 2010.3390/ijms20092334
28
29 Weibezahn J, Tessarz P, Schlieker C, Zahn R, Maglica Z, Lee S, Zentgraf H, Weber-Ban EU, Dougan DA,
30 Tsai FT *et al* (2004) Thermotolerance requires refolding of aggregated proteins by substrate translocation
31 through the central pore of ClpB. *Cell* 119: 653-665 doi: 10.1016/j.cell.2004.11.027
32
33 Wentink AS, Nillegoda NB, Feufel J, Ubartaite G, Schneider CP, De Los Rios P, Hennig J, Barducci A,
34 Bukau B (2020) Molecular dissection of amyloid disaggregation by human HSP70. *Nature* 587: 483-+ doi:
35 10.1038/s41586-020-2904-6
36
37 Wyszkowski H, Janta A, Sztangierska W, Obuchowski I, Chamera T, Klosowska A, Liberek K (2021) Class-
38 specific interactions between Sis1 J-domain protein and Hsp70 chaperone potentiate disaggregation of
39 misfolded proteins. *Proc Natl Acad Sci U S A* 11810.1073/pnas.2108163118
40
41 Xu X, Sarbeng EB, Vorvis C, Kumar DP, Zhou L, Liu Q (2012) Unique peptide substrate binding properties
42 of 110-kDa heat-shock protein (Hsp110) determine its distinct chaperone activity. *J Biol Chem* 287: 5661-
43 5672 doi: 10.1074/jbc.M111.275057
44
45 Yu HY, Ziegelhoffer T, Osipiuk J, Ciesielskil SJ, Baranowski M, Zhou M, Joachimiak A, Craig EA (2015)
46 Roles of Intramolecular and Intermolecular Interactions in Functional Regulation of the Hsp70 J-protein
47 Co-Chaperone Sis1. *Journal of Molecular Biology* 427: 1632-1643 doi: 10.1016/j.jmb.2015.02.007
48
49 Zietkiewicz S, Krzewska J, Liberek K (2004) Successive and synergistic action of the Hsp70 and Hsp100
50 chaperones in protein disaggregation. *J Biol Chem* 279: 44376-44383 doi: 10.1074/jbc.M402405200
51

Figure 1. Impact of Hsp110 on protein disaggregation by Hsp70 system depends on class of JDP.
(A) Refolding of aggregated luciferase by Ssa1-Sis1 (1 µM Ssa1, 1 µM Sis1) +/- µM Sse1 (0.1 µM) (left) or Ssa1-Ydj1 (1 µM Ssa1, 1 µM Ydj1) +/- Sse1 (0.1 µM) (right). Error bars show SD from three independent repeats. Luciferase activity was measured at indicated time points and normalized to the native activity. (B) Sensor-bound luciferase aggregates incubated with Ssa1-Sis1 (1 µM Ssa1, 1 µM Sis1) +/- Sse1 (0.1 µM) or Ssa1-Ydj1 (1 µM Ssa1, 1 µM Ydj1) +/- Sse1 (0.1µM), with or without ATP, as indicated. (C) Binding of Hsc70-DNAJB4 (3 µM Hsc70, 1 µM DNAJB4) or Hsc70-DNAJA2 (3 µM Hsc70, 1 µM DNAJA2) with or without Hsp105 (0.3 µM) to the heat-aggregated luciferase immobilized on the BLI biosensor. (B)-(C) The lines represent the average of three replicates, the shades designate SD and the dashed lines indicate the start of chaperones binding and dissociation steps.

1
Figure 2. Sse1 promotes modification of aggregates by the Hsp70 system. (A) Initial incubation of
heat-aggregated GFP aggregates with the Hsp70 system, followed by the addition of the Hsp104 D484K
F508A variant (0.15 μM). Recovery was initiated by the addition of the mix of indicated chaperones:
1 μM Ssa1, 1 μM Sis1, 1 μM Ydj1 or 0.1 μM Sse1. Dashed lines indicate the beginning of the incubation
with the Hsp104 variant. Curves show average values and shades indicate SD from three replicates.
(B) Fluorescence microscopy images of luc-GFP monitored upon addition of Ssa1-Sis1 +/- Sse1 or
Ssa1-Ydj1 +/- Sse1. Chaperones were used at 1 μM concentration, except for 0.1 μM Sse1. Left panels
show controls of the luciferase-GFP aggregates alone and upon the addition of the Hsp70 system without
ATP. Quantification of the fraction of aggregates > 2 μm is from three independent replicates. Two-tailed
t test was performed: *p < 0.05, ns: not significant.
12

Figure 3. Susceptibility of Hsp70 to Hsp110 depends on JDP class and phase of disaggregation.
(A) Titration of Sse1 in the refolding of aggregated luciferase by Ssa1-Sis1 (1 µM Ssa1, 1 µM Sis1) (red)
or Ssa1-Ydj1 (1 µM Ssa1, 1 µM Ydj1) (blue). Activity of luciferase was measured after 1 h and normalized
to that of the native protein. (B) Incubation of Hsc70-DNAJB4 (3 µM Hsc70, 1 µM DNAJB4) (green) or
Hsc70-DNAJA2 (3 µM Hsc70, 1 µM DNAJA2) (orange) with luciferase aggregates at increasing
concentrations of Hsp105. Luciferase activity was measured after 4 h and normalized to the activity of the
native protein. (C) Folding of non-aggregated luciferase diluted from 5 M GuHCl (grey), spontaneous or
mediated by the Hsp70 system comprising Ssa1-Sis1 (1 µM Ssa1, 1 µM Sis1) (red) or Ydj1-Ssa1
(1 µM Ssa1, 1 µM Ydj1) (blue) with increasing concentrations of Sse1. Activity of luciferase was measured
after 2 h and normalized to the native protein. (D) Binding of Ssa1-Sis1 (1 µM Ssa1, 1 µM Sis1) or
Ssa1-Ydj1 Ssa1 (1 µM Ssa1, 1 µM Ydj1) in the presence of Sse1 at the indicated concentrations to the
sensor covered with luciferase aggregates. Right panel shows a plot of the binding signal prior to the
dissociation step. (E) Renaturation of heat-aggregated GFP by Ssa1-Sis1 (1 µM Ssa1, 1 µM Sis1) in the
presence of Sse1 or Sse1-2 at the indicated concentrations. Right panel shows the plot of the recovered
GFP activity after 2 h of incubation with the Hsp70 system in the presence of Sse1 (orange) or Sse1-2
(red). (D-E) Dashed lines show the fitting of the *[Agonist] vs response* model to the data from the

1 stimulation and inhibition phases separately using the *GraphPrism* Software. (A-E) Error bars and shades
2 indicate SD from 3 repeats.

4
5
Figure 4. Hsp110 and class B JDP show apparent competition for Hsp70. (A) Upper panel shows the
scheme of the BLI experiment. Binding of Ssa1 (1µM) in the presence of increasing concentrations of Sse1
to Sis1 immobilized on the BLI sensor through the His_6–SUMO tag. Dashed lines indicate the moment of
addition of chaperones to the sensor-bound Sis1 and the dissociation step. (B) Renaturation of
heat-aggregated GFP by Ssa1-Sis1 (1µM Ssa1, 0.1 µM or 1 µM Sis1) at increasing concentrations of
Sse1. (C) Plot of GFP activity after 3 h recovered from aggregates by Ssa1 at different concentrations of
Sis1 and Sse1 (left). IC50 was determined by fitting the [*Inhibitor] versus response* model to the data from
three experiments using the *GraphPrism* Software (dashed lines). Two-tailed t test: *p < 0.05, **p < 0.01.
Lines are the average of three replicates and the error bars and shades designate standard deviation.

Figure 5. Impact of Hsp110 on Hsp70-dependent disaggregation. Hsp110 stimulates aggregate disassembly by Hsp70 with class B JDPs and Hsp104 (green arrow), contrary to the stage of final folding of the solubilized polypeptide (light purple arrow), at which Hsp70 with class A JDP are most effective. Hsp110 improves the disaggregation by increasing Hsp70 binding to the aggregate (upper right panel) and mediates aggregate remodelling into smaller assemblies (middle right panel). By facilitating partial Hsp70 dissociation from the substrate and/or JDP, at the optimal, sub-stoichiometric concentration, Hsp110 might gradually uncover new Hsp70-binding sites (lower right panel, small black arrows), potentially leading to the more abundant and effective Hsp70 recruitment to the aggregate. At higher Hsp110 levels, the destabilisation of Hsp70 interactions with protein substrates and JDPs leads to the inhibition across all stages of protein disaggregation (lower right panel).