Integrated single-nuclei and spatial transcriptomic analysis reveals propagation of early acute vein harvest and distension injury signaling pathways following arterial implantation

Marina E. Michaud*,[1], Lucas Mota*,[4], Mojtaba Bakhtiari[1], Beena E. Thomas[1], John Tomeo[4], William Pilcher[3], Mauricio Contreras[4], Christiane Ferran [4,5], Swati Bhasin[1,2], Leena Pradhan-Nabzdyk[4], Frank W. LoGerfo[4,] Patric Liang*,[4#], Manoj K. Bhasin[1,2,3,#,$]

[1] Department of Pediatrics, Emory School of Medicine, Atlanta, GA 30322, USA

[2] Aflac Cancer and Blood Disorders Center, Children Healthcare of Atlanta, Atlanta, GA

[3] Department of Biomedical Engineering, Emory University, Atlanta, GA 30322, USA

[4] Department of Surgery, Division of Vascular and Endovascular Surgery, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA 02215, USA

[5] Center for Vascular Biology Research and the Division of Nephrology Department of Medicine, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA 02215, USA

* Co-First authors

Senior Authors

$ Corresponding author

Manoj K. Bhasin, MS, PhD

Aflac Cancer and Blood Disorders Center

Children Healthcare of Atlanta

Woodruff Memorial Research Building, Room 4107

101 Woodruff Circle, 4th Floor East

Emory School of Medicine

Atlanta, GA 30322

Email: manoj.bhasin@emory.edu

Telephone: (404) 712-9849

Abstract

Background: Vein graft failure (VGF) following cardiovascular bypass surgery results in significant patient morbidity and cost to the healthcare system. Vein graft injury can occur during autogenous vein harvest and preparation, as well as after implantation into the arterial system, leading to the development of intimal hyperplasia, vein graft stenosis, and, ultimately, bypass graft failure. While previous studies have identified maladaptive pathways that occur shortly after implantation, the specific signaling pathways that occur during vein graft preparation are not well defined and may result in a cumulative impact on VGF. We, therefore, aimed to elucidate the response of the vein conduit wall during harvest and following implantation, probing the key maladaptive pathways driving graft failure with the overarching goal of identifying therapeutic targets for biologic intervention to minimize these natural responses to surgical vein graft injury.

Methods: Employing a novel approach to investigating vascular pathologies, we harnessed both single-nuclei RNA-sequencing (snRNA-seq) and spatial transcriptomics (ST) analyses to profile the genomic effects of vein grafts after harvest and distension, then compared these findings to vein grafts obtained 24 hours after carotid-cartoid vein bypass implantation in a canine model (n=4).

Results: Spatial transcriptomic analysis of canine cephalic vein after initial conduit harvest and distension revealed significant enrichment of pathways ($P < 0.05$) involved in the activation of endothelial cells (ECs), fibroblasts (FBs), and vascular smooth muscle cells (VSMCs), namely pathways responsible for cellular proliferation and migration and platelet activation across the intimal and medial layers, cytokine signaling within the adventitial layer, and extracellular matrix (ECM) remodeling throughout the vein wall. Subsequent snRNA-seq analysis supported these findings and further unveiled distinct EC and FB subpopulations with significant upregulation ($P < 0.00001$) of markers related to endothelial injury response and cellular activation of ECs, FBs, and VSMCs. Similarly, in vein grafts obtained 24 hours after arterial bypass, there was an increase in myeloid cell, protomyofibroblast, injury-response EC, and mesenchymal-transitioning EC subpopulations with a concomitant decrease in homeostatic ECs and fibroblasts. Among these markers were genes previously implicated in vein graft injury, including *VCAN* (versican), *FBN1* (fibrillin-1), and *VEGFC* (vascular endothelial growth factor C), in addition to novel genes of interest such as *GLIS3* (GLIS family zinc finger 3) and *EPHA3* (ephrin-A3). These genes were further noted to be driving the expression of genes implicated in vascular remodeling and graft failure, such as *IL-6*, *TGFBR1*, *SMAD4*, and *ADAMTS9.* By integrating the ST and snRNA-seq datasets, we highlighted the spatial architecture of the vein graft following distension, wherein activated and mesenchymal-transitioning ECs, myeloid cells, and FBs were notably enriched in the intima and media of distended veins. Lastly, intercellular communication network analysis unveiled the critical roles of activated ECs, mesenchymal transitioning ECs, protomyofibroblasts, and VSMCs in upregulating signaling pathways associated with cellular proliferation (MDK, PDGF, VEGF), transdifferentiation (Notch), migration (ephrin, semaphorin), ECM remodeling (collagen, laminin, fibronectin), and inflammation (thrombospondin), following distension.

Conclusions: Vein conduit harvest and distension elicit a prompt genomic response facilitated by distinct cellular subpopulations heterogeneously distributed throughout the vein wall. This response was found to be further exacerbated following vein graft implantation, resulting in a cascade of maladaptive gene regulatory networks. Together, these results suggest that distension initiates the upregulation of pathological pathways that may ultimately contribute to bypass graft failure and presents potential early targets warranting investigation for targeted therapies. This work highlights the first applications of single-nuclei and spatial transcriptomic analyses to investigate venous pathologies, underscoring the utility of these methodologies and providing a foundation for future investigations.

Keywords: Vein graft distension; implantation injury; endothelial injury; spatial transcriptomics; single-nuclei transcriptomics; coronary artery bypass grafting (CABG)

1 Introduction

Coronary artery and peripheral arterial disease affect 20.1 million and 6.5 million adults in the United States,
respectively.[1,2] Both disease processes can lead to life-threatening ischemia, resulting in the need for surgical bypass
surgery to redirect blood flow around blocked vessels. Autogenous vein, namely the greater saphenous vein, is the most
widely used conduit for coronary artery bypass grafting (CABG) and is the gold-standard conduit for peripheral arterial
bypass surgery.[3,4] However, the 1-year primary patency rate of coronary and peripheral vein grafts is as low as 60%.[5,6]
The biological mechanisms of early vein graft failure (VGF) remain poorly understood but have been classically
attributed to a maladaptive vascular wall remodeling response and the development of intimal hyperplasia (IH).
Previously, to understand the molecular mechanism of IH in vein grafts, we performed a temporal analysis of bypass
graft smooth muscle and endothelial cells following graft implantation.[7] The innovative network-based analysis depicted
that unresolved inflammation within 12 hours of graft implantation results in inflammatory and immune responses,
apoptosis, proliferation, and extracellular matrix (ECM) reorganization in both cell types, leading to graft failure. Vein
graft (VG) stenoses can occur anywhere along the vein conduit, suggesting that important injury response pathways
may occur even earlier than previously described, namely in the preparation of autogenous vein grafts for implantation
in the arterial system, as the conduit is manipulated and subjected to several sources of injury. Vein graft harvest injury
includes ischemia of the vein wall during dissection from surrounding tissues with loss of vasa vasorum function in the
vessel, pressure distension that induces hoop stress of the smooth muscle and endothelial cells, and exposure to an
electrolyte solution, all of which may contribute to the upregulation of maladaptive pathways that persist and propagate
following implantation into the arterial system. This response could be further exacerbated with distention of smaller
diameter or poorly compliant phlebosclerotic veins.[8] Accordingly, we aimed to identify the timeline associated with the
upregulation of maladaptive pathways from initial distension during graft harvesting to implantation to set the groundwork
for developing gene-therapy-based strategies to prevent or treat early vein graft injury and improve overall long-term
vein bypass patency.

To accomplish this, we leveraged advances in spatial and single-cell technologies to provide a unique insight into the
heterogeneity of the vascular landscape. Through the integrated use of single-nuclei RNA-sequencing (snRNA-seq)
and spatial transcriptomics (ST) analysis, we examined the genomic effects of vein graft harvest, distention, and
implantation across the regions of the vein, revealing distinct regulatory programs and cellular subpopulations mediating
the response to acute distension injury.

29 Materials and Methods

30 Data availability

31 The sequencing data that supports the findings of this study are available via the Gene Expression Omnibus database
32 under accession number (pending).

33 Animals

34 All animal studies were approved by the Institutional Animal Care and Use Committee (IACUC) at Beth Israel Lahey
35 Health and conducted in accordance with the National Institute of Health (NIH) Guide for the Care and Use of Laboratory
36 Animals. Canines were obtained from Marshall Bioresources (North Rose, NY 14516, US), a registered class A vendor
37 and AAALAC-accredited canine breeder for preclinical research purposes.

38 Canine Surgery

Bilateral upper limb cephalic vein harvest was performed on 25 kg male or female mongrel dogs (n=4). General
anesthesia was established and maintained with an initial sodium pentothal injection and subsequent 1% isofluorane

inhalation after orotracheal intubation. Vein harvest was performed in conventional fashion. The cephalic vein was

identified and exposed along the anterior surface from the wrist to the upper forearm. The distal portion of the vein was

then ligated and transected. The end of the transected vein was cannulated with a metal cannula and secured with a

silk suture. The vein was then sequentially distended by injecting a saline solution (0.9% sodium chloride, papaverine

hydrochloride 60 mg, and 43 mg of bivalirudin) while occluding the outflow. Intervening tributaries were identified and

ligated with silk sutures. A total length of 15 cm of vein was harvested. Total vein distension and harvest time was

approximately 30 minutes. The cephalic vein following distention, along with a separate segment of the undistended

cephalic vein, which served as the experimental control, were excised. Samples were collected and immediately snap-

frozen in cryovials using liquid nitrogen for single-nuclei analysis or embedded in O.C.T. blocks and flash-frozen in an

isopentane and liquid nitrogen bath for spatial transcriptomics analysis. Systemic bivalirudin was then infused before

the creation of the anastomoses. Bilateral end-to-side anastomoses were created on the vein with 7-0 Prolene sutures.

The intervening segment of the carotid artery was then ligated, creating preferential flow through the graft. After

hemostasis was achieved, the wounds were closed in three layers with 3-0 vicryl sutures. The patency of the graft was

confirmed through direct palpation and ultrasound imaging. Early (24 hours) and late (30 days) time points were chosen,

and harvest was performed with the dog under general anesthesia. Vein graft samples were then collected and

processed as previously described for cryopreservation and OCT embedding.

Immunohistochemistry

Formalin-fixed and paraffin-embedded tissue sections from each sample were cut to a 5-µm thickness and air-dried.

The staining for GLIS Family Zinc Finger 3 (GLIS3), Versican (VCAN), and fibrillin 1 (FBN1) was performed using the

Leica Bond RXM automated immunohistochemistry staining platform (Leica Biosystems). Before staining, the slides

were baked for 30 minutes in a 60 ºC oven. The slides were then labeled and loaded onto the Bond RXM. The slides

were deparaffinized with the Bond Dewax Solution and rinsed with Leica Wash Buffer. Following deparaffination, the

slides were heated to 100 ºC, and antigens were retrieved for 20 minutes with ER2 (high pH) antigen retrieval buffer

and then rinsed with Leica Wash Buffer. The peroxidase block was applied at room temperature for 5 minutes, and the

sections were washed with three rinses of wash buffer. The diluted antibodies (GLIS3 at 1:40, VCAN at 1:200, and

FBN1 at 1:500) were applied to the slides for 30 min at room temperature, followed by three rinses of wash buffer. Leica

Bond anti-Rb HRP secondary was applied and incubated for 8 min, and the detection was completed in combination

with the Leica Refine DAB kit, as per manufacturer recommendations. Slides were counterstained with hematoxylin for

5 min. Slides were then dehydrated, cover-slipped, and evaluated by light microscopy with scanned images of the slides.

The slides were scanned on a Hamamatsu Nanozoomer HT 2.0 at 40x.

Single-molecule RNA in situ hybridization

RNA in situ hybridization experiments were performed using the RNAscope® technology,[9] which has been previously

described.[9] Paired double-Z oligonucleotide probes were designed against target RNA using custom software. The

following probes were used: Cl-LMCD1-C1 (Part ID: 1317341-C1), Cl-ELN-C1 (Part ID: 1317351-C1), and Cl-FBN1-C1

(Part ID: 1317361-C1). The RNAscope Multiplex Fluorescent Detection Kit v2 (cat. no. 323110) (Advanced Cell

Diagnostics) was used according to the manufacturer's instructions. Slides of flash-frozen tissue samples of control and

distended veins were prepared according to the manufacturer's recommendations. Each sample was quality-controlled

for RNA integrity with a probe specific to the housekeeping genes *Polr2a, PPIB,* and *UBC*. Negative control background

staining was evaluated using a probe specific to the bacterial *dapB* gene. Fluorescent images were acquired using a

Leica Stellaris X5 confocal microscope with a 40x objective.

Single-nuclei transcriptomics sample preparation

We used the sample prep user guide (10x Genomics, CG000505, Rev A) for the Chromium Nuclei Isolation Kit with RNase Inhibitor (10x Genomics, 1000494) to isolate nuclei from flash-frozen tissues. The various buffers (lysis, debris removal, wash, and resuspension buffers) were prepared using kit components according to user guide instructions (10x Genomics, CG000505, Rev A). Briefly, ~50 mg of tissue, cut into smaller pieces to enable better lysis of cells, was added to pre-chilled sample dissociation tubes and dissociated by adding lysis buffer followed by multiple strokes with the pestle against the dissociation tubes. Following incubation on ice for 10 minutes, the dissociated tissue was added to the nuclei isolation column, placed in a pre-chilled collection tube, and spun at 16,000 g for 20 sec at 4°C. The flowthrough was collected, vortexed to resuspend the nuclei, and spun at 500g, 3 min, 4°C. The pellet was then resuspended in debris removal buffer, pipet mixed, and spun at 700 g, 10 min, 4°C. After discarding the supernatant, the pellet was washed with wash and resuspension buffer by spinning at 500 g, 5min, 4°C. For counting, the nuclei were stained with acridine orange/propidium iodide stain (Logos Biosystems) and counted using a LUNA-FX7 cell counter (Logos Biosystems). The nuclei pellet was resuspended in wash and resuspension buffer to get the desired nuclei concentration (700-1200 nuclei/ml). The snRNA-seq libraries were prepared immediately from the isolated nuclei with the Chromium Next GEM Single Cell 3' v3.1 reagent kits (10x Genomics, 1000268). Briefly, GEMs were prepared using the 10x Genomics controller followed by RT-PCR and cDNA amplification. The cDNA was then used to prepare snRNA-seq libraries according to the user guide (10x Genomics, CG000315 rev. B). The quality of the cDNA and final library preps was assessed using DNA HS bioanalyzer kits (Agilent), and concentration was estimated using a Qubit fluorometer (Thermo Fisher Scientific). Sequencing was performed using the massively parallel sequencing on the Novaseq 6000 platform using Novaseq S4 PE 100 kits (Illumina) to capture the expression of ~200-5,000 genes/nuclei.

Visium tissue permeabilization optimization, gene expression library construction, and sequencing

Frozen samples embedded in O.C.T. (TissueTek Sakura) were cryosectioned (10 μm sections) at –20 °C and transferred onto pre-chilled Visium Tissue Optimization Slides (10x Genomics, 3000394) and Visium Spatial Gene Expression Slides (10x Genomics, 2000233). H&E staining was performed according to the Visium Spatial Gene Expression User Guide (10x Genomics, CG000239 rev. F). Optimal permeabilization time was determined using the tissue optimization slide, which was processed according to the optimization protocol (10x Genomics, CG000239 rev. F). The tissue was permeabilized with permeabilization enzyme for different periods of time (6, 9, 12, 18, 24, and 30 minutes). Brightfield H&E images were taken using a 40X objective on Nanozoomer 2.0 HT. For tissue permeabilization optimization experiments, fluorescent images were taken using a 20X objective (0.5 μm/pixel) and CY3 and CY5 MSI filters at 177.75 ms and 73.31 ms exposure time in an Akoya Vectra Polaris whole slide scanner. The permeabilization time with maximum fluorescent signal was taken as the optimal time for permeabilization when processing gene expression slides. Brightfield H&E images were taken using a 40X objective on Nanozoomer 2.0 HT. For tissue permeabilization optimization experiments, fluorescent images were taken using a 20X objective (0.5 μm/pixel) and CY3 and CY5 MSI filters at 177.75 ms and 73.31 ms exposure time in an Akoya Vectra Polaris whole slide scanner. For the Visium spatial transcriptomics gene expression assay, tissue was permeabilized for the optimal time (9 minutes) followed by reverse transcription, second strand synthesis and denaturation, cDNA amplification and QC, and gene expression library construction according to 10x Genomics user guide (CG000239 rev. F). The gene expression libraries were quantified using a Qubit fluorometer (Thermo Fisher Scientific) and checked for quality using DNA HS bioanalyzer chips (Agilent). Sequencing depth was calculated based on the percent capture area covered by the tissue, and the 10x Genomics recommended sequencing parameters were used to run on the Novaseq S4 PE 100 kits (Illumina).

Transcriptional profiling and data analysis

FASTQ files for each sample were aligned to the Dog10K_Boxer_Tasha_1.0 canine reference genome[10] using Space

Ranger (v2.0.0, 10x Genomics) for spatial transcriptomics samples (n=7) and Cell Ranger (v7.0.0, 10x Genomics) for

single-nuclei samples (n=4). For spatial samples, microscopy images of the H&E stained samples were aligned to the

spatial voxels through the Space Ranger pipeline. The filtered feature-barcode matrices for the single-nuclei samples

were imported into Seurat (v4.3.0),[11] then low-quality nuclei with <200 unique genes, <500 UMI reads, or ≥30%

mitochondrial transcripts were filtered out. Due to the increased number of nuclei retrieved from the distended vein of

Dog 2, this sample was downsampled to 1,000 nuclei that were used for subsequent analysis, keeping both the control

vein and distended vein groups similar in total nuclei (1,233 and 1,188 nuclei, respectively) for comparison. For both the

single-nuclei and spatial datasets, the samples were normalized with SCTransform and then integrated using integration

anchors-based batch correction via Seurat. Principal component analysis (PCA) was then performed on the integrated

spatial and single-nuclei datasets using all non-ribosomal features, followed by construction of a K-nearest neighbors

graph (using cosine similarity) and Louvain clustering, then non-linear dimensional reduction via Uniform Manifold

Approximation and Projection (UMAP). The resolution and number of principal components used for clustering were

selected using the clustree R package (v0.5.0).[12] Differential gene expression analysis was performed via Seurat using

the FindAllMarkers function, comparing expression between clusters or sample groups (distended veins versus control

veins).

Statistical analysis and data visualization

All statistical analysis was performed using R. Significantly differentially expressed genes were determined based on

the Wilcoxon rank test with Bonferroni multiple test correction ($P < 0.05$). Significantly enriched pathways were

calculated using Benjamini-Hochberg adjusted p-values (*P adj (BH)* < 0.05). Spatial plots, UMAP plots, and differential

gene expression heatmaps/dot plots were all generated using Seurat. Bar plots of the cluster distributions and the

deconvolution matrix were generated using ggplot2 (v3.4.1).[13] Volcano plots based on differential gene expression were

made using the EnhancedVolcano R package (v3.16). Dot plots of enriched pathways were generated via the dotplot

function in clusterProfiler. Networks of enriched pathways were generated via the cnetplot function in the clusterProfiler

Bioconductor package.[14]

Results

Optimization of tissue permeabilization techniques for the preparation of vein samples for spatial and single-nuclei transcriptomic analyses

Bilateral upper limb cephalic vein harvest was performed on canine specimens to yield three matched pairs of control

and distended veins (Figure 1A). Following distension of the cephalic vein with a saline solution containing papaverine

and bivalirudin, segments of the vein were excised along with separate undistended cephalic vein segments. These

tissue samples were subsequently cryopreserved in liquid nitrogen for single-nuclei analysis or embedded in O.C.T. and

flash-frozen for spatial transcriptomic assays. Single-nuclei methodologies uniquely enable the transcriptomic analysis

of challenging vascular tissues by eliminating the need to maintain whole-cell integrity during tissue dissociation while

still preserving individual cellular resolution at the transcriptomic level. Therefore, to overcome the challenges associated

with obtaining viable single-cell suspensions from fibrous vascular tissues, [15] we endeavored to harness snRNA-seq to

analyze the vein sample transcriptomes. Through this approach, we were successfully able to capture diverse cell types

within the vein (i.e., smooth muscle cells, fibroblasts, endothelial cells, and leukocytes) with 200-1,000 nuclei expressing

~200-5,000 genes per sample, illustrating single-nuclei isolation as a viable alternative to single-cell isolation for the

transcriptomic analysis of difficult-to-process vascular tissue (Table S1).

As this work presents the first spatial transcriptomic investigations of superficial vein samples, we first sought to optimize

the tissue permeabilization time for spatial transcriptome profiling. Using the Visium platform from 10x Genomics, tissue

sections are transferred and fixed onto capture areas containing ~5,000 barcoded spots; cDNA synthesis subsequently

incorporates these barcodes to ultimately facilitate the transcriptomic analysis of tissue by spot (1-10 cells captured per

spot) (Figure 1A). Tissue permeabilization presents a crucial step in the sample preparation, as insufficient

permeabilization will reduce the amount of RNA released and captured, thus reducing the transcriptomic resolution and

affecting downstream analyses, while over-permeabilization results in cDNA diffusion, compromising the spatial

localization.[16] Examining six different permeabilization times (6–30 minutes) via fluorescence microscopy (Figure 1B),

we identified 9 minutes as the optimal permeabilization time for 10 μM vein tissue based on the fluorescent signal.

Applying this technique, we successfully performed spatial transcriptomics on the vein samples, yielding 125-550 spots

captured per sample with an average expression of ~870 genes per spot (Table S1).

Lastly, to facilitate targeted spatial transcriptomic analysis of the veins, we developed a facile computational method for

annotating the spatial plot voxels according to histological analysis (Figure 1C). To accomplish this, we utilized the open-

source imaging software ImageJ to produce image masks for each layer of the vein wall, then developed an R-based

function for adding these labels to the metadata of each spatial sample (Figure 1D). This function, which will be available

through the STannotate package, enables researchers to readily add histological annotations to vein samples for

downstream analysis.

Integration of single-nuclei and spatial transcriptomics illustrates a significant impact of distension on

enrichment and profiles of endothelial and fibroblast subpopulations

To investigate how distension alters the spatial gene expression landscape of the vein, we first performed spatial

transcriptomic (ST) analysis on cephalic vein samples from four canines, including three vein samples retrieved following

distension with saline and four untouched control vein samples (Table S1). Applying our STannotate tool, we labeled

the spatial voxels of each sample into intimal, medial, and adventitial layers based on the tissue morphology observed

through histological analysis (Figure S1). After this, we performed a comparative analysis of the transcriptomic profiles

of the intimal, medial, and adventitial layers between distended and control veins (Figure 2A).

Following distension, significant overexpression of genes involved in extracellular matrix (ECM)-remodeling, cell

proliferation, and migration, as well as cytokine production, were found to be overall significantly upregulated

($P < 0.00001$) in the vein conduit. Specifically, within the intima, regulators of endothelial cell (EC) proliferation and

migration, fibronectin (*FN1*) and insulin-like growth factor-binding protein 7 (*IGFPB7*),[17,18] were transcriptionally

upregulated. Similarly, in the adventitia, genes associated with fibroblast (FB) activation and cytokine production, c-Fos

(*FOS*) and vimentin (*VIM*), were upregulated. Within the media, transcription of the contractile VSMC markers, α-SMA

(*ACTA2*) and myosin light chain proteins (*MYL6* and *MYL9*), were increased. Expression of contractile VSMC markers

is typically associated with a quiescent SMC phenotype; however, a significant decrease in the contractile markers

transgelin (*TAGLN*) and cysteine-rich protein 1 (*CSRP1*) was concurrently observed, conversely signifying VSMC-

preprogramming to a synthetic (activated) phenotype (Figure 2A, bottom panel). Furthermore, among the downregulated

genes in the distended veins, negative regulators of oxidative stress, metallothionein 2A (*MT2A*) and angiopoietin-like

4 (*ANGPTL4*), were decreased in the adventitia, suggesting that oxidative stress response may be dysregulated in

distended veins (Figure 2A, bottom panel).

We next examined the pathways enriched within each layer of the distended vein by performing pathway enrichment

analysis (Figure 2B) on the differentially expressed genes ($P < 0.05$) and mapping the network of enriched pathways

between distended and control veins (Figure 2C). The results of this pathway analysis further illustrate significantly

increased ($P < 0.05$) ECM remodeling, cell proliferation and migration, and cytokine production following distension,
wherein integrin- and syndecan-mediated interactions and collagen degradation and formation, as well as increased
Rho GTPase-, ephrin-, and MET-mediated cell motility, were elevated in the intima and media. Additionally, the media
displayed significant enrichment of platelet activation and profibrotic IL-4 and IL-13 cytokine signaling ($P < 0.05$),[19] the
latter of which was also enriched in the adventitia (Figure 2B).

Collectively, these initial results of our ST analysis suggest that distension of the vein initiates the upregulation of
pathways implicated in the activation of ECs, VSMCs, and FBs, namely ECM remodeling across the inner vein wall,
platelet activation driven by the media, cytokine signaling within the outer vein wall, and the increased migratory
signaling across all three layers.

Clustering of spatial voxels reveals the heterogeneous distribution of transcriptional profiles driving cellular
activation and vascular remodeling within the media and adventitia following distension
To expand upon these findings and investigate the spatial transcriptomic landscape through an unbiased approach, we
next clustered the spatial voxels (containing 1-10 cells each) based on their cell-type composition and transcriptomic
profiles via K-nearest neighbor (KNN) graph construction and Louvain clustering, visualized by Uniform Manifold
Approximation and Projection (UMAP) (Figure 3A). Seven clusters formed, which we subsequently labeled according
to their primary localization to the intima (e.g., I1), media (e.g., M1), or adventitia (e.g., A1). Markedly, the spatial
distribution of these clusters within the distended vein (Figure 3B, Figure S2) was nonuniform compared to the control
vein (Figure 3C, Figure S2), highlighting the heterogeneous effect of sheer stress during distension that may contribute
to dysregulated healing in segments of the vein, leading to IH progression observed in graft failure.[20]

Particularly, while the intimal cluster (I1) evenly surrounded the lumen (Figure 3B) in the control vein, this cluster
significantly extended into portions of the media following distension (Figure 3C). Notably, this cluster depicted
significant upregulation ($P < 0.00001$) of EC activation and endothelial-mesenchymal transition (EndMT) associated
genes, namely fibronectin (*FN1*),[21] von Willebrand factor (*VWF*),[22] tropomyosin 1 (*TPM1*),[23] α-SMA (*ACTA2*),[24] γ-actin
(*ACTG1*),[25] and thrombospondin 1 (*THBS1*),[26] suggesting that distension promotes the emergence of activated EC and
mesenchymal phenotypes (Figure 3E).

Furthermore, distended veins became enriched in their proportion of clusters A1 and M1 and correspondingly decreased
in clusters A2 and M2 (Figure 3D), indicating a shift in genomic expression within the media and adventitia following
distension. While clusters M1 and M2 shared similar expression profiles, the M1 cluster (enriched in distended veins)
displayed modest upregulation ($P < 0.00001$) of contractile VSMC marker genes (Figure 3E). Further pathway
enrichment analysis showed significant enrichment ($P \, adj \, (BH) < 0.005$) of cell activation and migration pathways,
including ECM remodeling, MAPK signaling, L1CAM-mediated cell adhesion, and platelet degranulation (Figure 3F). In
contrast, the M2 cluster (enriched in control veins) primarily showed significant upregulation ($P < 0.00001$) of genes
negatively regulating proliferation, such as cysteine-rich protein (*CRIP1*)[27], antioxidant glutathione peroxidase-1
(*GPX1*),[28] adipogenesis regulatory factor (*ADIRF*), and gelsolin (*GSN*) (Figure 3E). Notably, gelsolin was found to be
the most significantly downregulated gene ($P < 0.00001$) overall in distended veins (Figure 2A). While previous work
has demonstrated that gelsolin inhibits angiotensin II-mediated activation of myofibroblasts in mycardiofibrosis,[29] its role
in vascular injury has not been elucidated and warrants future investigation. Within the adventitia, the A1 cluster
displayed significant upregulation ($P < 0.00001$) for key regulators of ECM remodeling, including collagen I (*COL1A*)
and collagen III (*COL3A*), fibulin-1 (*FBLN*1), elastin (*ELN*), and extracellular matrix protein 1 (*ECM1*) (Figure 3E). In
contrast, the A2 cluster showed significant over-expression ($P < 0.00001$) of genes implicated in fibroblast-mediated
healing, including negative regulator of cell migration apolipoprotein E (*APOE*),[30] profibrotic collagen VI (*COL6A2*),[31]

and vascular-remodeling regulator aortic carboxypeptidase-like protein (*ACLP/AEBP1*) (Figure 3E).[32] Furthermore,
pathway enrichment analysis illustrated significantly increased activation (*P adj (BH)* < 0.01) of ECM remodeling and
cytokine signaling pathways in the A1 cluster, while the A2 cluster exhibited significant activation (*P adj (BH)* < 0.01) of
ROCKs, PAKs, and ephrin kinases involved in modulating cell migration, potentially aiding in fibroblast-mediated healing
(Figure 3G).

Together, these results of spatial clustering, coupled with the initial ST analysis, demonstrate that distension alters the
transcriptomic landscape of the vein, coordinating a distinct response to the imparted endothelial injury across each
layer of the vein wall. Specifically, within the intima and media, markers of EC activation, cellular proliferation and
migration, platelet activation, and EndMT are primarily upregulated following distension, while in the adventitia, key
markers of fibroblast-mediated ECM remodeling are upregulated.

254 Single-nuclei RNA-seq analysis identifies distinct fibroblast and endothelial subpopulations mediating
255 homeostasis, injury response, and mesenchymal-transition following distension

While sequencing-based ST approaches provided unique insights into the spatial context of genome-wide transcription,
the technology remains limited in cellular resolution.[33] Therefore, we further investigated the effects of distension at the
single-cell level through single-nuclei RNA sequencing (snRNA-seq) analysis as a complementary approach. Using four
paired cephalic vein samples from two canines, including two vein samples retrieved following distension with saline
and two control vein samples (Table S1), we performed snRNA-seq analysis on a total of 1,188 nuclei derived from
control veins and 1,223 nuclei derived from distended veins. Clustering of the nuclear transcriptomes yielded ten clusters
visualized by UMAP embedding, representing endothelial, fibroblast, vascular smooth muscle, and myeloid cell types
(Figure 4A) based on the expression of canonical markers (Figure S3). Owing to the plasticity of endothelial cells and
fibroblasts, four distinct subpopulations also appeared within both the EC and FB clusters. Examining the distribution of
these cell types and subpopulations showed an overall decrease in the proportion of ECs and SMCs following distension
with an overall increase in FB and myeloid cell populations (Figure 4B). These alterations in cellular composition are
noteworthy, as endothelial cell loss and increased immune cell infiltration, coupled with dedifferentiation and necrosis
of contractile VSMCs, are associated with the early development of intimal hyperplasia in VGF.[20]

Analysis of the effects of distension on the gene expression profile further reflected these differences in cellular
composition (Figure 4B). Specifically, within distended veins, key transcription factors (TFs), including Krüppel-like factor
4 (*KLF4*), GLIS Family Zinc Finger 3 (*GLIS3*), early growth response factor 1 (*EGR1*), and zinc finger E-box-binding
homeobox 2 (*ZEB2*), were all significantly upregulated (*P* < 0.00001), in addition to the AP-1 subunits, c-Fos (*FOS*) and
c-Jun (*JUN*) (Figure 4C). While these TFs all have well-documented roles in cellular reprogramming and
transdifferentiation, only *KLF4, EGR1, ZEB2,* and *AP-1* have been documented in vascular endothelial-to-mesenchymal
and myofibroblast transitions. In contrast, *GLIS3* has not been previously reported to play a role in vascular remodeling,
yet was found to be significantly upregulated in the Fibroblast B, C, and D clusters, as well as the Endothelial B and C
clusters. Intriguingly, this TF was recently found to promote the transdifferentiation of fibroblasts into retinal pigmented
epithelial cells,[34] suggesting that it may play a significant role in fibroblast reprogramming observed in response to
endothelial injury following distension.

In addition to these transcription factors, genes encoding ECM proteins, including the EGF-family proteins integrin
subunit beta like 1 (*ITGBL1*), epidermal growth factor receptor (*EGFR*), fibrillin (*FBN1*), and versican (*VCAN*), as well
as the secreted matrix proteins slit guidance ligand 2 (*SLIT2*), collagen I (*COL1A2*), peroxidasin (*PXDN*), and coiled-
coil domain-containing protein 80 (*CCDC80*) were significantly upregulated (*P* < 0.00001) across the fibroblast
subpopulations within the distended vein, highlighting the upregulation of ECM remodeling following distension (Figure

4D).

Notably, significantly downregulated genes ($P < 0.00001$) within the distended veins were primarily associated with
maintaining endothelial identity and integrity, such as von Willebrand factor (*VWF*), vascular endothelial protein tyrosine
phosphatase receptor type B (*PTPRB*),[35] and MDS1 and EVI1 complex locus (*MECOM*)[36] (Figure 4C). Regulators of
endothelial injury response myosin VIIA and Rab interacting protein (*MYRIP*),[37] forkhead box C1 protein (*FOXC1*),[38]
and LIM and cysteine-rich domains 1 (*LMCD1*)[39] were also found to be significantly downregulated ($P < 0.00001$),
particularly among the endothelial subpopulations (Figure 4C). Overall, the downregulation of these homeostatic and
regulatory genes indicates that distension induces a transcriptomic shift towards the activation of ECs and the positive
regulation of endothelial injury response.

Among the differentially expressed genes within the SMC population of distended veins, chromatin remodeler *ATRX*
and VSMC-specific voltage-dependent calcium channel *CACNA1C* were significantly upregulated ($P < 0.00001$).
Although the epigenetic role of ATRX has not been elucidated in the context of VSMC function, its upregulation among
the SMC population may point toward the reprogramming of VSMCs following distension, as observed through the ST
analysis. In contrast, the metalloprotease protein MMP-28 gene expression decreased in VSMCs following distension.
Importantly, MMP-28 is considered a protective metalloprotease, wherein its depletion has been demonstrated to
impede M2 macrophage activation, leading to impaired inflammatory response and cardiac dysfunction.[40]

We next explored the differentially expressed genes within each cluster (Figure 4D) and their associated pathways
(Figure 4E), revealing similar trends between several cell types and subpopulations. First, Endothelial D, Fibroblast D,
and myeloid cell clusters were significantly enriched ($P < 0.05$) for pathways related to cellular proliferation and
migration, including AKT, NOTCH, and HIF signaling pathways, RHO and RAC GTPase cycle activation, and cell
division, suggesting that these endothelial and fibroblast subpopulations represent activated phenotypes (Figure 4E).
Upregulated genes among the fibroblast clusters overlapped in their enrichment of ECM remodeling pathways, integrin-
and syndecan-mediated interactions, and MET-mediated cellular migration ($P < 0.05$), mirroring the pathways enriched
within the media and adventitia observed via ST analysis (Figure 2B). Within the SMC cluster, gene expression was
primarily associated with positive regulation of platelet homeostasis and negative regulation of proliferation and
migration through nitric oxide modulation of cGMP, congruent with the increased proportion of this cluster observed in
control veins compared to distended veins. However, similar to the contrasting SMC gene expression between control
and distended veins observed through differential gene expression analysis (Figure 4C), several pathways, including
ECM remodeling, RHO and RAC GTPase cycle activation, and platelet activation, were partially enriched ($P < 0.005$)
within the SMC cluster. This data suggests that the portion of the SMC population associated with distended veins
displays altered transcriptional regulation toward a synthetic phenotype following distension.

To elucidate the different subpopulations present among the fibroblast and endothelial cell clusters, we isolated both
the fibroblast (Fibroblast A-D) and endothelial cell (Endothelial A-D) clusters, then conducted differential gene
expression analysis between each cluster and identified key markers and biological themes differentiating each
subpopulation (Figures 5–6). First, upon examining the effects of distension on the overall gene expression within the
fibroblast population (Figure 5A), we found that downregulated genes in the fibroblast population after distension were
primarily associated with the Fibroblast A subpopulation. These downregulated genes ($P < 0.00001$) overlapped with
genes previously identified in control veins (Figure 4C) and also included the antiapoptotic regulator secreted frizzled-
related protein 1 (*SFRP1*),[41] the negative regulator of angiogenesis multimerin-2 (*MMRN2*),[42] and negative regulator of
vascular EC proliferation (*SELENOP*).[43] In contrast, significantly upregulated genes ($P < 0.00001$) within the fibroblast
population following distension included transcriptional promoters of cell proliferation and ECM proteins previously

identified amongst the set of upregulated genes within the distended veins (Figure 4C), in addition to ECM-related genes

laminin 2 subunit (*LAMA2*), dystonin (*DST*), and BicC family RNA binding protein 1 (*BICC1*). Interestingly, the roles of

dystonin and BICC1 in vascular injury have not been previously investigated, but their function as an anchor within the

actin cytoskeleton and negative regulator of Wnt signaling, respectively, suggest that they may play an important role

in modulating fibroblast motility and transdifferentiation following vascular injury.

A closer look at the top gene markers associated with each fibroblast cluster (target fibroblast cluster versus other

fibroblast clusters) facilitated the characterization of each fibroblast subpopulation (Figure 5B). First, expression of

negative regulators of cell proliferation and migration (*ARHGAP31*, *SFRP1*, *MMRN2*, and *SELENOP*)[41–44] within the

Fibroblast A cluster and its increased proportion within control veins (Figure 5C) indicate that this fibroblast

subpopulation is likely responsible for promoting vascular homeostasis following endothelial injury (Figure 5D). Similarly,

within the Fibroblast B cluster (Figure 5B), the proapoptotic TF nuclear receptor 4A2 (*NR4A2*)[45] was significantly

upregulated with the profibrotic TF early growth response protein (*EGR1*), as well as the ECM proteins integrin subunit

β-like 1 (*ITGBL1*), periplakin (*PPL*), and elastin (*ELN*) ($P < 0.00001$), signaling that these fibroblasts are likely involved

in promoting fibroblast-mediated healing following endothelial injury (Figure 5D). In contrast, the Fibroblast C and D

clusters both exhibited increased expression of genes associated with ECM remodeling; however, Fibroblast D

specifically displayed significant upregulation of genes associated with myofibroblast transition ($P < 0.00001$), such as

differentiation markers Krüppel-like factor 6 (*KLF6*) and EBF transcription factor 1 (*EBF1*), and production of type III

collagen (*COL3A1*), type I collagen (*COL1A2*), laminin 2 subunit (*LAMA2*), fibrillin (*FBN1*), and fibronectin (*FN1*),

indicating that Fibroblast C cells are responsible for ECM remodeling while Fibroblast D cells are protomyofibroblasts

(Figure 5D). Collectively, the fibroblast subpopulations Fibroblast B (injury response fibroblasts), Fibroblast C (ECM

remodeling fibroblasts), and Fibroblast D (protomyofibroblasts) were all enriched in their proportions within distended

veins compared to control veins (Figure 5C), in addition to the overall increase in fibroblasts within the vein following

distension (Figure 4B), revealing that distinct fibroblast subpopulations may play a crucial role in mediating the acute

response to distension injury.

Within the endothelial cell subpopulations (Figure 6), many similarities appeared, as with the fibroblast population.

However, distension notably resulted in the significant upregulation ($P < 0.00001$) of numerous genes within the

endothelial population, including thrombospondin 1 (*THBS1*), multimerin 1 (*MMRN1*), and cadherins 7 (*CHD7*) and 13

(*CDH13*) (Figure 6A). Importantly, these genes have been reported to play significant roles in endothelial injury response

and, moreover, the development of thrombosis, intimal hyperplasia, and atherosclerosis.[46–49] The largest endothelial

cluster, Endothelial A, was largely characterized by the downregulation of genes that were significantly overexpressed

across the other endothelial clusters (Figure 6B). This cluster depicted an increase in proportion within control veins

compared to distended veins (Figure 6C). Furthermore, this cluster displayed subtle expression of promoters of EC

homeostasis, including inositol 1,4,5-trisphosphate receptor type 3 (*ITPR3*)[50] and negative regulators of EC migration

RAS protein activator like 2 (*RASAL2*) and EGF-like domain multiple 7 (*EGFL7*), suggesting that this subpopulation may

be responsible for preventing EC activation and maintaining endothelial homeostasis (Figure 6D). Similarly, within the

Endothelial B cluster (Figure 6B), genes previously implicated in mediating healing following endothelial injury were

significantly upregulated ($P < 0.00001$), namely *LMCD1*, forkhead box protein P1 (*FOXP1*), plexin domain containing 2

(*PLXCD2*),[39,51,52] and navigator-3 (*NAV3*). Intriguingly, navigator-3 was previously shown to suppress growth factor

signaling, promote apoptosis, and decrease cell migration in epithelial cell lines and breast cancer models,[53] and thus

may play a similar role in maintaining endothelial homeostasis (Figure 6D).

In contrast to the Endothelial A and Endothelial B clusters, the Endothelial C and Endothelial D clusters exhibited

markers of EC activation and EndMT, respectively. Specifically, the Endothelial C subpopulation (Figure 6B), which was
the only endothelial population displaying overall enrichment in the distended vein (Figure 4B), exhibited significantly
higher expression ($P < 0.00001$) of platelet adhesion ligand multimerin-1 (*MMRN1*)[47] and mediator of activated EC
migration A-kinase anchor protein 12 (*AKAP12*),[54] as well as elevated expression of proliferation and migration genes:
serine/threonine kinase 39 (*STK39*) and *KLF4*. Moreover, this subpopulation exhibited increased expression ($P <$
0.00001) of kalirin (*KALRN*), reelin (*RELN*), and peroxidasin (*PXDN*); notably, kalirin promotes neointimal hyperplasia
by activating RAC expression,[55] while reelin increases leukocyte adhesion molecules ICAM-1, VCAM-1, and E-selectin
and the progression of atherosclerosis,[56] and peroxidasin has been correlated with vascular endothelial dysfunction by
disrupting eNOS function.[57] Together, this indicates that the Endothelial C subpopulation represents an activated EC
phenotype expressing markers associated with poor prognosis following graft implantation (Figure 6D).

Lastly, the Endothelial D subpopulation displayed increased markers of EndMT ($P < 0.00001$), including vimentin (*VIM*),
phosphoglycerate Mutase 1 (*PGAM1*), S100 calcium-binding protein A6 (*S100A6*), and cysteine-rich protein 1 (*CRIP1*)
(Figure 6D). Intriguingly, while EndMT is often associated with vascular dysfunction, the proportion of this mesenchymal
subpopulation was decreased in distended veins compared to control veins (Figure 6C), possibly reflecting the
alternative role of mesenchymal cells as mediators of fibrosis and inflammation in proper vascular healing, such as that
observed in type II epithelial-mesenchymal transition. Moreover, the concomitant increase in activated ECs (Endothelial
C) and decrease in mesenchymal-transitioning cells (Endothelial D) in distended veins also supports recent findings
that EndMT plays a marginal role in the contribution to neointimal formation leading to VGF and rather appears to
represent an adaptive response capable of suppressing VSMC reprogramming and promoting quality vascular
remodeling following graft implantation.[58]

387 Integration of single-nuclei and spatial transcriptomics illustrates a significant impact of distension on the
388 enrichment of activated endothelial and fibroblast subpopulations within the intima
To achieve the most granular view of the cellular transcriptome landscape, we integrated our single-nuclei and spatial
transcriptomics data by deconvoluting the spatial voxels of control and distended veins based on the transcriptional
signatures of the cellular subpopulations elucidated via snRNA-seq (Figure 7A).

First, employing a scoring-based integration approach, we generated a set of representative and differentially expressed
genes for each of the identified subpopulations, then utilized these gene signatures to assign an enrichment score of
each cell type to the spatial voxels of control and distended veins (Method S3, Figure S4). To examine the spatial
distribution of the subpopulations across the vein wall regions, we then averaged the cell type enrichment scores of
voxels within the intima, media, and adventitia, generating enrichment score matrices for distended and control veins.
Next, a differential enrichment matrix was computed to compare the relative spatial distribution of cell types between
the distended and control veins, displaying corresponding increases and decreases in cell types across layers of the
distended veins relative to control veins (Figure 7B). The resulting differential enrichment score matrix illustrated that
Endothelial A and B subpopulations are reduced in the intimal layer of distended veins, pointing to decreased
maintenance of EC homeostasis and mitigation of EC-mediated response at the site of the endothelial injury.
Conversely, Endothelial C and D subpopulations appear enriched in the intima, reflecting the activation and
dedifferentiation of ECs following distension. Furthermore, these mesenchymal-transitioning ECs (Endothelial D) were
also enriched in the media of distended veins. Despite the overall decrease in the proportion of Endothelial D cells in
distended veins captured by snRNA-seq (Figure 4B), the observed enrichment of the Endothelial D signature within the
intima and media suggests that the limited Endothelial D subpopulation within distended veins may functionally differ
from that of control veins, which were enriched in the adventitia. Increased fibroblast populations were also observed in

the intima and media of distended veins, suggesting that fibroblasts may also localize to the site of endothelial injury

following distension. Importantly, migration of fibroblasts to the intima plays a well-documented role in the development

of intimal hyperplasia and atherosclerosis.[59] Within the intima of distended veins, myeloid cells are also enriched, further

underscoring the recruitment and infiltration of immune cells initiated by the effects of distension on the endothelium.

Lastly, SMCs display increased enrichment in the adventitia of distended veins, supporting our hypothesis that while

the overall proportion of SMCs decreases following distension, SMCs present in the distended veins may undergo

VSMC-reprogramming toward a synthetic phenotype facilitating their migration to the adventitia.

As a complementary integration approach, we additionally performed model-based spatial deconvolution via

Cell2Location with subsequent neighborhood enrichment analysis via Squidpy. Through neighborhood enrichment

analysis, we examined the proximity probabilities of each cellular subpopulation in relation to one another throughout

the vein (Figure 7C) based on the most abundant cellular subpopulation within each voxel (Figure 7D). This analysis

showed the increased proximity of each subpopulation to other members of the respective subpopulation, particularly

for SMCs, myeloid cells, and mesenchymal-transitioning cells (Endothelial D). This increased colocalization of the

Endothelial D subpopulation aligns with the documented enrichment of intra-cell type signaling promoting EC activation

and EndMT.[60] Furthermore, the Endothelial D and SMC populations exhibited increased proximity, highlighting the

crosstalk between ECs and VSMCs to mediate EC and VSMC activation in vascular dysfunction.[61] To illustrate the

spatial distribution of these cellular subpopulations, we constructed spatial plots displaying the most abundant

population within each voxel, predicted using a Bayesian model generated from the snRNA-seq dataset via

Cell2Location (Figure 7D). Congruent with the previous findings, the intima of the control vein displayed an increased

Endothelial B (injury response) subpopulation, while the intima and media of the distended vein displayed enrichment

of the Endothelial D population. Concurrent with this increase in Endothelial D abundance, a decrease in the overall

proportion of VSMC-rich regions within the media is observed, mirroring the reduction in overall SMC proportion

observed via single-nuclei analysis. Similarly, within the distended vein, enrichment of the protomyofibroblast (Fibroblast

D) population is observed within the adventitia, demonstrating the increase in fibroblast activation following distension.

To validate the integration of our single-nuclei and spatial datasets, we lastly performed RNAscope on control and

distended vein samples (Figure 7E, Figure S5), using fluorescent antibodies for representative marker genes (*LMCD*

and *FBN1*) corresponding to key subpopulations differentially enriched between control and distended veins (Endothelial

B and Fibroblast D, respectively). The upregulation of *LMCD1* within the intima of control veins detected via RNAscope

(Figure 7E, right) aligns with the predicted enrichment of the Endothelial B population within the intima of control veins

observed by our scoring-based (Figure 7E, left) and model-based (Figure 7D) integration approaches. Similarly, the

upregulation of *FBN1* within the adventitia of distended veins (Figure 7E, right) aligns with the predicted enrichment of

the Fibroblast D population within the adventitia of distended veins, also observed by our integration approaches.

Taken together, the results of the integrated single-nuclei and spatial transcriptomic analysis elucidate the effects of

distension on the cellular architecture of the vein. Most notably, the enrichment of activated ECs within the intima and

migratory EndMT cells extending toward the media is observed following distension, along with the migration of

fibroblasts to the intima and a decrease in homeostatic and injury response in luminal ECs. Importantly, these changes

are also implicated in the development of IH and atherosclerosis, further suggesting that distension may initiate the

pathways implicated in these VGF-related pathologies.

Cellular communication analysis elucidates key signaling molecules promoting inflammation and vascular

injury responses following distension

Finally, to elucidate the intercellular communication network between the subpopulations present in the veins following

distension, we analyzed the number and strength of incoming and outgoing signaling interactions for each cell type, as
well as the key signaling molecules mediating these interactions, via cellular communication analysis.[62] Overall, the total
number of intercellular signaling interactions within distended veins increased in addition to the strength of these
interactions (Figure S6). Most notably, signaling to the Endothelial B (injury response ECs), Endothelial D
(mesenchymal-transitioning ECs), Fibroblast C (ECM remodeling FBs), Fibroblast D (protomyofibroblasts), myeloid
cells, and SMCs increased significantly following distension (Figure 8A-B). Furthermore, the strength of signaling to the
mesenchymal-transitioning and myeloid cells in distended veins greatly increased (Figure S6), highlighting the roles of
these populations in response to distension.

To characterize the intercellular communication among these subpopulations, we first examined which signaling
pathways were enhanced in distended versus control veins (Figure 8C), revealing several pathways also observed in
our ST and single-nuclei analyses, including ephrin (EPHA), reelin (RLN), collagen, laminin, and fibronectin (FN1). In
addition to these previously implicated pathways, several interesting signaling pathways emerged. First, within the
control veins, signaling pathways associated with vascular healing and angiogenesis were enriched, namely fibroblast
growth factor (FGF),[63] cell adhesion molecule (CADM),[64] and bone morphogenic protein (BMP)[65] mediated signaling
pathways. In contrast, distended veins were enriched for disparate angiogenic signaling pathways, particularly those
mediated by vascular endothelial growth factors (VEGF), platelet-derived growth factors (PDGF), midkine neurite
growth-promoting factor 2 (MK), and semaphorins (SEMA6, SEMA3). Notably, the upregulation of VEGF, PDGF, and
MK have all been well-documented as key contributors to intimal hyperplasia, contributing to graft failure.[66–68]

Next, to illuminate which subpopulations are modulating these key signaling pathways in distended veins compared to
control veins, we generated a heatmap displaying the differential signaling of each pathway within the cellular
subpopulations (Figure 8D) and elucidated the intercellular communication networks mediated by these pathways
(Figure 8E). Markedly, signaling within the Endothelial D (mesenchymal-transitioning ECs) and Fibroblast D
(protomyofibroblasts) subpopulations were most significantly affected by distension, closely followed by the Endothelial
C (activated ECs) and SMC populations.

Within the Endothelial D (mesenchymal-transitioning ECs) population, signaling pathways associated with cell growth
and differentiation (MK, NOTCH, CSPG4),[69,70] neointimal formation and hyperplasia (THBS),[46] and leukocyte adhesion
and atherosclerosis (RELN)[56] were specifically enhanced following distension. Similarly, within the SMC population,
signaling associated with VSMC activation, namely chondroitin sulfate proteoglycan 4 (*CSPG4*) and fibronectin (*FN1*)
pathways, were enriched, indicating that VSMC-reprogramming is initiated by distension injury. Inspecting the
intercellular communication networks for these pathways unveiled that Endothelial D and SMC populations are primarily
responsible for producing ligands associated with these pathways, whose surface receptors were displayed on other
EC and FB subpopulations. For example, the outgoing MK signal (*MDK*) produced by the Endothelial D population
interacted with the low-density lipoprotein (*LPR1*), nucleolin (*NCL*), and integrin (*ITGA6*, *ITGB1*) receptors present
across the four fibroblast populations, activated EC subpopulation (Endothelial C), and myeloid cells, suggesting a
potential role of the Endothelial D subpopulation in promoting the proliferation of fibroblasts, activated ECs, and immune
cells following distension (Figure 8E). Likewise, the SMC population of the distended veins predominantly produced
fibronectin (*FN1*), which was received by integrin surface receptors (*ITGA5*, *ITGB1*) on Endothelial D cells, as well as
CD44 receptors on Fibroblast C (ECM remodeling FBs), Fibroblast D (protomyofibroblast), and myeloid cells (Figure
8E); stimulation by fibronectin is known to promote EndMT and myofibroblast transdifferentiation, suggesting that
differentiation of the cell types is strongly influenced by the SMC population within distended veins. Collectively, these
findings further support our hypothesis that while the proportions of mesenchymal-transitioning ECs and VSMC

populations are decreased in distended veins, the residing populations of these cell types are phenotypically different
from those of the control veins, promoting cell proliferation, migration, and activation that may initiate the development
of intimal hyperplasia in VGF.

Within the Fibroblast D (protomyofibroblasts) population, increased signaling via the ephrin (EPHA), platelet-derived
growth factor (PDGF), and vascular endothelial growth factor (VEGF) pathways were observed (Figure 8D). These
pathways all have well-documented roles in angiogenesis, including the implication of VEGF and PDGF in the
development of intimal hyperplasia.[66,67] Interestingly, ephrin signaling in the distended vein was mediated by ephrin-A5
(*EFNA5*), produced by Fibroblast D cells, with ephrin-A3 (*EFNA3*) receptors displayed on both Fibroblast D and
Fibroblast B (injury response FBs) cells. While ephrin-B has been thoroughly investigated for its role in angiogenesis as
a VEGF receptor,[71] the role of ephrin-A signaling remains understudied and may warrant future investigation. Notably,
VEGF signaling was also upregulated among the Endothelial C (activated ECs) population, which was uniquely enriched
in distended veins. A closer examination of the VEGF signaling network subsequently revealed that VEGF signaling is
mediated by VEGFC, produced by Fibroblast D, Endothelial A (homeostatic ECs), and Endothelial B (injury response
ECs) cells, and received exclusively by VEGFR3 receptors on the Endothelial C subpopulation (Figure 8E). This finding
suggests that the Fibroblast D, Endothelial A, and Endothelial B cells induce the activation of ECs in response to
distension. Likewise, PDGF signaling via the PDGFD ligand was produced by Fibroblast D, Fibroblast C (ECM
remodeling FBs), Fibroblast B (injury response FBs), and VSMC populations and received exclusively by Fibroblast D
cells by PDGFRB expression, facilitating the activation of these fibroblasts following distension.

Together, the results of our intercellular communication analysis provide greater insight into the distinct roles of specific
EC and FB subpopulations in combination with VSMCs and immune cells in modulating the response to endothelial
injury initiated by distension. Combining the key signaling molecules with prominent differentially expressed genes within
distended veins, we generated a network of the key pathways implicated in distension injury, illustrating the effects of
distension on genomic expression within the vein (Figure 8F). The results of these analyses cumulatively demonstrate
that distension elicits a coordinated response throughout the vein wall, driving the activation of endothelial cells,
fibroblasts, and vascular smooth muscle cells and stimulating proinflammatory and immune responses.

Vascular remodeling, proinflammatory pathways, and key genes (FBN1, VCAN, GLIS3) initiated by distension
persist post-bypass
Lastly, to determine the relation of the above findings to the pathophysiology of vein grafts post-implantation, we
extended our investigation to a vein graft sample retrieved 24 hours post-bypass, examining the cellular composition
and transcriptomic profile. At 24 hours post-bypass, an increase in myeloid cell, protomyofibroblast, injury-response EC,
and mesenchymal-transitioning EC subpopulations was observed with a concomitant decrease in homeostatic ECs and
fibroblasts (Figure 9A). These findings are congruent with a response to increased cellular activation, proinflammatory,
and immune signaling pathways upregulated at the time of distension. Furthermore, spatial transcriptomic analysis of
the implanted graft illustrated the distribution of these subpopulations with protomyofibroblasts extending from the
adventitia to the media and injury response and EndMT ECs throughout the intima (Figure 9B). Next, to examine how
the genomic response initiated by distension extends to the transcriptomic profile observed 24 hours post-implantation,
we constructed a temporally-resolved multilayer gene network that illustrates the trajectory of the genomic response
from control veins to distended and implanted veins. Notably, there were several significantly upregulated genes at the
time of distension, including fibrillin-1 (*FBN1*) and versican (*VCAN*),[72,73] that were subsequently driving the expression
of genes implicated in vascular remodeling and graft failure, such as *IL-6*, *TGFBR1*, *SMAD4*, and *ADAMTS9*,[74–76] within
the implanted vein (Figure 9C). Additionally, novel genes of interest identified through the above single-nuclei

transcriptomic analysis, such as *GLIS3*, were present among the drivers of post-implantation gene expression,
underscoring the advantage of this approach in identifying potential therapeutic targets. Specifically, the transcription
factor *GLIS3*, which was significantly upregulated at the time of distension and has been previously demonstrated to
play a role in epithelial cell reprogramming,[34] was found to drive expression of TAZ (*WWTR1*), a central regulator in
HIPPO signaling pathway driving EndMT,[77] within the grafted vein (Figure 9C). Finally, to validate the expression of
these genes at the protein level, we performed IHC analysis on control, distended, and grafted vein samples, staining
for the expression of VCAN, FBN-1, and GLIS3 (Figure 9D, Figure S7). Mirroring the results of transcriptomic analyses,
an increase in the expression of these proteins was observed at the time of distension, which was further increased 24
hours post-implantation by IHC staining.

Discussion

Vein bypass graft failure and thrombosis following peripheral and coronary artery bypass grafting can result in limb loss,
heart failure, and death. Thus, to overcome this challenge, there is a need to elucidate the molecular mechanisms
underlying graft failure and identify potential targets for therapeutic development. VGF is primarily attributed to IH, a
process that causes thickening and intraluminal narrowing of a bypass graft, obstructing blood flow and ultimately
leading to graft thrombosis.[78] It has been posited that IH may occur in response to bypass implantation, where repeated
exposure to arterial hemodynamics, ischemia-reperfusion injury, and biomechanical stressors results in damage to the
endothelium, triggering changes in vascular gene expression and concomitant transition of quiescent ECs to an
activated proinflammatory and prothrombotic state and VSMCs from contractile to an activated synthetic state.[78–81]
Subsequent VSMC proliferation and migration, coupled with excessive ECM production and fibroblast migration,
ultimately generates obstructive IH lesions and VG atherosclerosis, contributing to graft failure. While the development
of IH following graft implantation has been investigated, we have sought to elucidate the onset of these maladaptive
pathways, beginning at the time of graft harvesting and its connections with the changes seen post-implantation.
Through this study, we unveiled that conduit distension during graft harvesting induces endothelial injury and initiates
the dysregulation of genes and pathways implicated in graft failure, which partially drives injury-related pathways
identified following implantation. Harnessing an integrated single-nuclei and spatial transcriptomics approach, we
provide unprecedented insights into the genomic effects of distension, the unique cellular subpopulations mediating this
response, and the spatial heterogeneity of acute distension injury.

By leveraging recent advances in spatial transcriptomic technologies, we performed the first reported spatial
transcriptomic investigation of vein tissue samples. This methodology uniquely facilitates unbiased genomic analysis
with spatial resolution, capturing the expression of over 35,000 genes localized across the layers of the vein wall.
Through this approach, we illustrated the effects of distension on the genomic and cellular landscape of the vein graft.
Aligning with previous investigations of IH in VGF,[78] we found that distension alters gene expression heterogeneously
throughout the layer of the vein, reflecting the traumatic sheer stress resulting from distention to diameters greater than
the existing cellular architecture permits. Specifically, examining each layer individually, we identified markers of EC,
VSMC, and fibroblast activation within the intima, media, and adventitia, respectively. Among these markers, we
identified the upregulation of several genes implicated in IH and atherosclerosis, including fibronection (*FN1*) within the
intima and vimentin (*VIM*) within the adventitia.[82,83] Concomitant with the differential expression of these genes in
distended veins, we identified corresponding enriched pathways within the intima, media, and adventitia; pathways
associated with ECM remodeling, cell proliferation, and migration (collagen, integrin, syndecan, and MET signaling)
were upregulated in the intima and media, while platelet activation pathways were significantly upregulated in the media
and profibrotic IL-4 and IL-13 signaling were upregulated in both the adventitia and media. To validate these findings,

we further performed unsupervised clustering of the spatial voxels based on their transcriptomic profiles, identifying

clusters that heterogeneously localized to distinct regions of the vein. Interestingly, we identified a cluster (I1) exhibiting

significant upregulation of markers of EC activation and EndMT, including fibronectin (*FN1*), VWF, tropomyosin 1

(*TPM1*), α-SMA (*ACTA2*), γ-SMA (*ACTG1*), and thrombospondin 1 (*THBS1*), which protruded from the intima into the

media of the distended vein. Furthermore, within distended veins, we identified enriched medial and adventitial clusters

displaying markers of cell activation and migration, including MAPK and L1CAM signaling within the media, and markers

of inflammation, including IL-4 and IL-13 signaling in the adventitia. Collectively, our spatial transcriptomic analyses

reveal that distension results in the upregulation of ECM remodeling, cellular activation, and inflammatory pathways

across the intima, media, and adventitia of the vein, highlighting the sensitivity of veins to manipulation and dissection.

While these pathways play critical roles in the physiological response to endothelial injury, their dysregulation leads to

IH development. Therefore, the upregulation of these at the time of distension may illustrate the initiation of pathways

that later become dysregulated and lead to IH in the context of graft failure. Vasodilatory agents have been suggested

as potential therapeutics to reduce endothelial injury, but even with the use of papaverine-based solution for intraluminal

vasodilation in this study, distension was found to impart significant endothelial injury, underscoring the need for

alternative approaches.

These findings are further supported by the results of our single-nuclei transcriptomic analysis, wherein we identified

distinct populations of ECs, FBs, VSMCs, and immune cells mediating the genomic response to distension. Overall, we

observed decreased EC and VSMC populations following distension, with a concurrent increase in FB and myeloid cell

populations. We posit that these changes to the cellular architecture reflect injury and desquamation of the endothelial

layer. Moreover, EC loss, leukocyte recruitment, and dedifferentiation of contractile VSMCs are documented in the early

phases of IH development in VGF,[20] supporting the idea that IH leading to VGF may be initiated during graft harvesting.

Subsequent analysis of the nuclear transcriptomes further enabled the identification and characterization of unique

endothelial cell and fibroblast subpopulations associated with maintaining vascular homeostasis (Endothelial A,

Fibroblast A), endothelial injury response and repair (Endothelial B, Fibroblast B), ECM remodeling (Fibroblast C), and

cellular activation (Endothelial C) and transdifferentiation (Endothelial D, Fibroblast D). Following distension, the

proportion of injury response and activated ECs increased among the EC population, and within the FB population, the

proportion of injury response FBs, ECM remodeling FBs, and protomyofibroblasts increased. Intriguingly, the proportion

of mesenchymal transitioning ECs (Endothelial D) was increased in the control veins compared to distended veins

despite the documented role of EndMT in vascular dysfunction. Notably, this finding supports the recent work by Wu

and coworkers, which illuminated the pro-fibrotic and pro-inflammatory role of mesenchymal cells in proper wound

healing,[58] similar to that of type II epithelial-mesenchymal transition. Furthermore, differential gene expression, pathway

enrichment, and intercellular communication analyses together suggested that the EndMT (Endothelial D)

subpopulation exhibits a disparate transcriptomic profile in distended veins compared to control veins. Specifically,

Endothelial D cells within the distended veins were upregulated in cell growth, proliferation, and migration (MK, CSPG4),

leukocyte adhesion and stenosis (RELN), and neointimal formation and hyperplasia (THBS) associated pathways. While

the population of VSMCs was reduced in distended veins, the residing VSMCs of distended veins exhibited a distinct

expression profile compared to those of control veins, wherein the VSMCs of distended veins produced greater collagen

and fibronectin, promoting EndMT and transdifferentiation-associated pathways. Additionally, through the single-nuclei

transcriptomic analyses, we also identified the upregulation of several key genes previously implicated in VGF, including

versican (*VCAN*), fibrillin (*FBN1*), and vascular endothelial growth factor-C (*VEGFC*). Moreover, we unveiled several

novel genes of interest that were significantly upregulated following distension and have yet to be investigated in the

context of graft implantation and failure, including neuron navigator 3 (*NAV3*), semaphorin-6A (*SEMA6A*), BicC family

RNA binding protein 1 (*BICC1*), and dystonin (*DST*). Furthermore, owing to the increased resolution of nuclear

transcripts via single-nuclei analysis, we also identified significant upregulation of several key transcriptional regulators

following distension, including Krüppel-like factor 4 (*KLF4*), early growth response factor 1 (*EGR1*), and zinc finger E-

box-binding homeobox 2 (*ZEB2*), and GLIS Family Zinc Finger 3 (*GLIS3*), the latter of which has previously been

implicated in fibroblast to epithelial cell transdifferentiation[34] but warrants investigation for its role in regulating

transdifferentiation during vascular remodeling. Taken together, the results of our single-nuclei analysis elucidate

differentially expressed key genes and enriched pathways in distended veins, underscoring the acute effects of

distension and providing a foundation for future analyses of these mechanisms. Integrating the ST and snRNA-seq

datasets, we illustrated that EndMT-associated (Endothelial D) gene expression was increased in the intima and media

of distended veins, along with homeostatic and injury response FB subpopulations and myeloid cells. Collectively, these

findings illustrate that distension results in acute injury to the endothelium, triggering EC, VSMC, and FB activation with

concomitant increases in the associated pathways involving cell proliferation and migration, ECM remodeling,

inflammatory and immune responses, and transdifferentiation. This switch from quiescent to activated, pro-inflammatory

phenotypes, coupled with the upregulation of prothrombogenic factors such as PDGF, Tumor Necrosis Factor (TNF),

cytokines, and other damage-associated molecular patterns (DAMPs), as part of the initial responses to injury

contributing to VGF, has long been noted.[84,85] However, through this work, we have unveiled that these mechanisms

may originate during the initial vein harvest dissection and distension process, highlighting the potential advantage of

alternative harvesting techniques or early gene-targeted therapeutic interventions in improving vein graft patency.

Significantly, the pro-inflammatory changes identified immediately following distension, both in terms of cellular

population and gene expression, persist post-implantation and, in some instances, appear to be driving graft failure

mechanisms in the implanted vein. One of the key mechanisms of pathological intimal hyperplasia formation is

excessive myofibroblast differentiation from quiescent fibroblast and migration from adventitia to the neointima.[86,87] This

excessive myofibroblast migration to the neointima has been previously implicated in graft failure following

implantation,[20,88,89] and the increase in protomyofibroblast seen in our samples 24 hours post-implantation indicates a

similarly prominent role in the cellular response to injury post-implantation. However, while this phenotypical shift in

fibroblasts has been primarily attributed to the arterialization of the vein following implantation,[78,90] our results indicate

that this transition is initiated during harvest and distension of the vein, as demonstrated by the increase in the proportion

of injury response and ECM-remodeling FBs and protomyofibroblasts following vein distension, as well as their migration

to the intima illustrated by spatial transcriptomics. Additionally, our temporally-resolved multilayer gene network showing

the relationship between the distended and post-implantation transcriptomic profiles indicates that several genes

previously implicated in both early and late graft failure are present at 24 hours post-implantation and driven by the

upregulation of key genes during graft distension. Namely, the upregulation of IL-6, responsible for promoting leukocyte

adhesion and vascular wall infiltration leading to thrombosis, and TGF-ß receptor 1, which plays a key role in mediating

SMC differentiation as part of the TGF-ß signaling pathway, were demonstrated to be directly influenced by upregulated

genes at the time of distension, including fibrillin-1 (*FBN1*) and versican (*VCAN*),[91–93] which were validated by IHC.

These findings further illustrate the putative role of vein harvest and distension as the initial insult that results in the

cascade of physiopathological processes leading to VGF and highlights pre-implantation as a promising time point for

intervention.

While the study presented herein pioneers the investigation of venous pathophysiologies, specifically acute distension

injury through an integrated spatial and single-nuclei transcriptomics approach, it has several limitations. First, the study

was not conducted in human vessels, and although we have previously utilized canine end-to-side cephalic vein

interposition VG bypass models to investigate VGF after bypass grafting due to the animal model's established

relevance to pre-clinical human disease of IH and VG atherosclerosis, the difference in injury pathways between canine
and human may exist.[94] Additionally, as this dataset does not include failed graft samples, we cannot determine whether
the genomic changes demonstrated in our work would ultimately result in graft failure. Furthermore, the study sample
size limits the statistical power of our results, specifically when comparing cellular populations between distended and
control veins. Population comparisons are also inherently confounded by the ability of different transcriptomic assays to
capture specific cell types. For example, through our investigation, we found that the ST methodology was superior in
capturing SMC-associated gene expression, while snRNA-seq provided high resolution of EC and FB population gene
expression. While this limitation can impair statistical significance, it also underscores the utility of an integrated spatial
and single-nuclei transcriptomics approach for efficiently capturing the heterogeneous architecture of the vein.

Conduit distension during vein graft harvesting is canonically performed during peripheral artery bypass surgery to
prevent vasospasms and ensure adequate vein graft diameter and compliance.[95] During this process, the saphenous
vein is exposed along its entire length, followed by the insertion of a cannula at the distal aspect of the vein and
subsequent distension. However, the sheer stress imparted on the vascular wall from elevated distension pressure has
been illustrated to cause damage to the endothelium and media.[95] Thus, to circumvent these effects, recent efforts to
minimize manipulation and avoid distention of the vein graft have been employed, including the use of warmed
papaverine solution to vasodilate the vein graft during distention. This process has been shown to reduce endothelial
loss by scanning electron microscopy.[96,97] Furthermore, in-situ vein bypass techniques, wherein the vein graft is left
along its anatomic course without complete explanation, were performed to minimize dissection and manipulation of the
vein graft. However, this type of in-situ bypass can only be utilized for certain distal arterial targets and has not been
found to be superior to previously described techniques.[98] Others have advocated for a "no-touch" vein harvest
technique, in which the adventitia and perivascular tissue of the vein are kept intact during dissection while employing
minimal to no distension.[99] This "no-touch" harvest technique has been recently popularized in the cardiac literature for
vein harvest during coronary artery bypass grafting, with a randomized control trial showing a significantly reduced risk
of VG occlusion at one year.[8] The results of our study support the use of these alternative graft harvesting approaches,
highlighting the adverse effects of distensions on the vein and providing a possible explanatory mechanism for the
significant improvement in graft patency seen in grafts implanted with minimal distension compared to those harvested
with the conventional technique.

In conclusion, the results of this work present the first investigations of the acute genomic responses to distension during
vein graft harvesting by leveraging single-nuclei and spatial transcriptomics. These responses, mediated by distinct
populations of endothelial, fibroblast, smooth muscle, and myeloid cells throughout the layers of the vein, initiate EC,
FB, and SMC activation through upregulation of cell proliferation, differentiation, and migration, ECM remodeling,
proinflammatory, and angiogenic pathways. Furthermore, the upregulation of genes implicated in IH, endothelial
dysfunction, and atherosclerosis, all contributing to VGF, were observed following graft distension, in addition to novel
genes of interest that may play a role in VGF. As a whole, this work provides a foundation for future investigations of
vein graft harvesting, graft implantation, and graft failure, exemplifies the utility of spatial and single-nuclei
transcriptomics for the investigation of vascular pathologies, and identifies potential targets for the development of
therapeutics that limit graft failure.

Novelty and Significance

What Is Known?

- Autogenous vein, namely the greater saphenous vein, is the most widely used conduit for coronary artery bypass grafting (CABG) and is the gold-standard conduit for peripheral arterial bypass surgery, however, the 1-year primary patency rate of coronary and peripheral vein grafts is as low as 60%, primarily due to maladaptive vascular wall remodeling response and the development of intimal hyperplasia (IH).
- While the late mechanisms leading to vein graft failure (VGF) have received great attention, less attention has been given to how these mechanisms may be initiated, particularly the role of graft harvesting techniques in the upregulation of maladaptive pathways implicated in VGF.

What New Information Does This Article Contribute?

- This study elucidated the acute genomic response following distension during vein graft harvesting, including the upregulation of genes and pathways previously implicated in VGF following implantation, and lays a foundation for future single-nuclei and spatial transcriptomics approaches to studying the mechanisms of vein graft failure and other vascular pathologies.
- Leveraging single-nuclei and spatial transcriptomics, this work unveils the mechanisms of endothelial cell (EC), fibroblast (FB), and vascular smooth mucle cells (VSMC) activation following distension via the upregulation of cell proliferation, differentiation, and migration, ECM remodeling, proinflammatory, and angiogenic pathways by distinct populations of ECs, FBs, VSMCs, and myeloid cells.

This work sheds light on the acute genomic effects of distension during vein graft harvesting, in which EC, FB, and VSMC activation is initiated through the upregulation of cell proliferation, differentiation, and migration, ECM remodeling, proinflammatory, and angiogenic pathways. We identified distinct subpopulations of Ecs, FBs, and VSMCs mediating this response, which were localized to specific regions of the vascular wall following distension, highlighting the effects of the traumatic sheer stress via distension on the pathophysiology of the vein. Furthermore, our analyses collectively suggest that the mechanisms implicated in VGF, including IH and endothelial dysfunction that are seen post-implantation, may originate at the time of distension during graft harvesting, underscoring the underscoring the importance of careful vein harvesting techniques and providing a basis for designing gene-targeted therapies to minimize maladaptive distention injury responses.

Article Information

Acknowledgments

The RNA sequencing in this work was supported by the Emory Integrated Genomics Core at Emory University, subsi-
dized by the School of Medicine and Winship Cancer Institute. Dr. Vasantha L. Kolachala (Department of Pediatrics,
Emory University) aided in the Visium assay standardization. Vaunita Parihar performed H&E imaging of the Visium
slides and immunohistochemistry staining at the Cancer Tissue and Pathology shared resource of Winship Cancer
Institute of Emory University and NIH/NCI under award number P30CA138292. The content is solely the authors' re-
sponsibility and does not necessarily represent the official views of the National Institutes of Health. Figure illustrations
were made with BioRender.com.

Sources of Funding

This work was funded in part by the following grants from the National Institutes of Health: 1) R01 5R01HL086741 to
M.K.B, C.J.F., and F.W.L.; 2) 5 T32 HL007734 to L.P-N., C.J.F., and F.W.L.; 3) 5R01HL021796 to C..J.F., and F.W.L.

Disclosures

The authors declare no competing interests.

Supplemental Materials

Methods S1-S5

Tables S1

Figures S1-S7

References

1. Tsao CW, Aday AW, Almarzooq ZI, Alonso A, Beaton AZ, Bittencourt MS, Boehme AK, Buxton AE, Carson AP, Commodore-Mensah Y, Elkind MSV, Evenson KR, Eze-Nliam C, Ferguson JF, Generoso G, et al. Heart Disease and Stroke Statistics-2022 Update: A Report from the American Heart Association. *Circulation*. 2022;145(8):E153–E639.

2. Virani SS, Alonso A, Aparicio HJ, Benjamin EJ, Bittencourt MS, Callaway CW, Carson AP, Chamberlain AM, Cheng S, Delling FN, Elkind MSV, Evenson KR, Ferguson JF, Gupta DK, Khan SS, et al. Heart Disease and Stroke Statistics - 2021 Update: A Report From the American Heart Association. *Circulation*. 2021;143(8):E254–E743.

3. Almasri J, Adusumalli J, Asi N, Lakis S, Alsawas M, Prokop LJ, Bradbury A, Kolh P, Conte MS, Murad MH. A systematic review and meta-analysis of revascularization outcomes of infrainguinal chronic limb-threatening ischemia. *Journal of Vascular Surgery*. 2019;69(6):126S-136S.

4. Martínez-González B, Reyes-Hernández CG, Quiroga-Garza A, Rodríguez-Rodríguez VE, Esparza-Hernández CN, Elizondo-Omaña RE, Guzmán-López S. Conduits used in coronary artery bypass grafting: A review of morphological studies. *Annals of Thoracic and Cardiovascular Surgery*. 2017;23(2):55–65.

5. Widimsky P, Straka Z, Stros P, Jirasek K, Dvorak J, Votava J, Lisa L, Budesinsky T, Kolesar M, Vanek T, Brucek P. One-year coronary bypass graft patency: A randomized comparison between off-pump and on-pump surgery angiographic results of the PRAGUE-4 trial. *Circulation*. 2004;110(22):3418–3423.

6. Conte MS, Bandyk DF, Clowes AW, Moneta GL, Seely L, Lorenz TJ, Namini H, Hamdan AD, Roddy SP, Belkin M, Berceli SA, DeMasi RJ, Samson RH, Berman SS. Results of PREVENT III: A multicenter, randomized trial of edifoligide for the prevention of vein graft failure in lower extremity bypass surgery. *Journal of Vascular Surgery*. 2006;43(4):742–751.

7. Bhasin M, Huang Z, Pradhan-Nabzdyk L, Malek JY, LoGerfo PJ, Contreras M, Guthrie P, Csizmadia E, Andersen N, Kocher O, Ferran C, LoGerfo FW. Temporal network based analysis of cell specific Vein graft transcriptome defines key pathways and hub genes in implantation injury. *PLoS ONE*. 2012;7(6):e39123.

8. Tian M, Wang X, Sun H, Feng W, Song Y, Lu F, Wang L, Wang Y, Xu B, Wang H, Liu S, Liu Z, Chen Y, Miao Q, Su P, et al. No-Touch Versus Conventional Vein Harvesting Techniques at 12 Months After Coronary Artery Bypass Grafting Surgery: Multicenter Randomized, Controlled Trial. *Circulation*. 2021;144(14):1120–1129.

9. Wang F, Flanagan J, Su N, Wang LC, Bui S, Nielson A, Wu X, Vo HT, Ma XJ, Luo Y. RNAscope: a novel in situ RNA analysis platform for formalin-fixed, paraffin-embedded tissues. *The Journal of molecular diagnostics : JMD*. 2012;14(1):22–29.

10. Jagannathan V, Hitte C, Kidd JM, Masterson P, Murphy TD, Emery S, Davis B, Buckley RM, Liu YH, Zhang XQ, Leeb T, Zhang YP, Ostrander EA, Wang GD. Dog10k_boxer_tasha_1.0: A long-read assembly of the dog reference genome. *Genes*. 2021;12(6):847.

11. Hao Y, Hao S, Andersen-Nissen E, Mauck WM, Zheng S, Butler A, Lee MJ, Wilk AJ, Darby C, Zager M, Hoffman P, Stoeckius M, Papalexi E, Mimitou EP, Jain J, et al. Integrated analysis of multimodal single-cell data. *Cell*. 2021;184(13):3573-3587.e29.

12. Zappia L, Oshlack A. Clustering trees: a visualization for evaluating clusterings at multiple resolutions. *GigaScience*. 2018;7(7):1–9.

13. Wickham H. Ggplot2. *Ggplot2*. 2009.

14. Yu G, Wang L-G, Han Y, He Q-Y. clusterProfiler: an R Package for Comparing Biological Themes Among Gene Clusters. *OMICS: A Journal of Integrative Biology*. 2012;16(5):284–287.

15. Chavkin NW, Hirschi KK. Single Cell Analysis in Vascular Biology. *Frontiers in Cardiovascular Medicine*. 2020;7:42.

16. Hudson WH, Sudmeier LJ. Localization of T cell clonotypes using the Visium spatial transcriptomics platform. *STAR Protocols*. 2022;3(2):101391.

17. Mariko B, Ghandour Z, Raveaud S, Quentin M, Usson Y, Verdetti J, Huber P, Kielty C, Faury G. Microfibrils and fibrillin-1 induce integrin-mediated signaling, proliferation and migration in human endothelial cells. *American Journal of Physiology - Cell Physiology*. 2010;299(5):977–987.

18. Tamura K, Hashimoto K, Suzuki K, Yoshie M, Kutsukake M, Sakurai T. Insulin-like growth factor binding protein-7 (IGFBP7) blocks vascular endothelial cell growth factor (VEGF)-induced angiogenesis in human vascular endothelial cells. *European Journal of Pharmacology*. 2009;610(1–3):61–67.

19. Wynn TA. Cellular and molecular mechanisms of fibrosis. *Journal of Pathology*. 2008;214(2):199–210.

20. Owens CD, Gasper WJ, Rahman AS, Conte MS. Vein graft failure. *Journal of Vascular Surgery*. 2015;61(1):203–216.

21. Al-Yafeai Z, Yurdagul A, Peretik JM, Alfaidi M, Murphy PA, Orr AW. Endothelial FN (Fibronectin) deposition by α5β1 integrins drives atherogenic inflammation. *Arteriosclerosis, Thrombosis, and Vascular Biology*. 2018;38(11):2601–2614.

22. Holthenrich A, Gerke V. Regulation of von-willebrand factor secretion from endothelial cells by the annexin a2-s100a10 complex. *International Journal of Molecular Sciences*. 2018;19(6).

23. Ajami NE, Gupta S, Maurya MR, Nguyen P, Li JYS, Shyy JYJ, Chen Z, Chien S, Subramaniam S. Systems biology analysis of longitudinal functional response of endothelial cells to shear stress. *Proceedings of the National Academy of Sciences of the United States of America*. 2017;114(41):10990–10995.

24. Xu S, Ilyas I, Little PJ, Li H, Kamato D, Zheng X, Luo S, Li Z, Liu P, Han J, Harding IC, Ebong EE, Cameron SJ, Stewart AG, Weng J. Endothelial dysfunction in atherosclerotic cardiovascular diseases and beyond: From mechanism to pharmacotherapies. *Pharmacological Reviews*. 2021;73(3):924–967.

25. Dugina VB, Shagieva GS, Shakhov AS, Alieva IB. The cytoplasmic actins in the regulation of endothelial cell function. *International Journal of Molecular Sciences*. 2021;22(15):7836.

26. Yamashiro Y, Thang BQ, Ramirez K, Shin SJ, Kohata T, Ohata S, Vu Nguyen TA, Ohtsuki S, Nagayama K, Yanagisawa H. Matrix mechanotransduction mediated by thrombospondin-1/integrin/YAP in the vascular remodeling. *Proceedings of the National Academy of Sciences of the United States of America*. 2020;117(18):9896–9905.

27. Latonen L, Järvinen PM, Laiho M. Cytoskeleton-interacting LIM-domain protein CRP1 suppresses cell proliferation and protects from stress-induced cell death. *Experimental Cell Research*. 2008;314(4):738–747.

28. Takapoo M, Chamseddine AH, Bhalla RC, Miller FJ. Glutathione peroxidase-deficient smooth muscle cells cause paracrine activation of normal smooth muscle cells via cyclophilin A. *Vascular pharmacology*. 2011;55(5–6):143–148.

29. Jana S, Aujla P, Hu M, Kilic T, Zhabyeyev P, McCulloch CA, Oudit GY, Kassiri Z. Gelsolin is an important mediator of Angiotensin II-induced activation of cardiac fibroblasts and fibrosis. *FASEB Journal*. 2021;35(10).

30. Hui DY, Basford JE. Distinct signaling mechanisms for apoE inhibition of cell migration and proliferation. *Neurobiology of Aging*. 2005;26(3):317–323.

31. Tracy LE, Minasian RA, Caterson EJ. Extracellular Matrix and Dermal Fibroblast Function in the Healing Wound. *Advances in Wound Care*. 2016;5(3):119–136.

32. Wang D, Rabhi N, Yet SF, Farmer SR, Layne MD. Aortic carboxypeptidase-like protein regulates vascular adventitial progenitor and fibroblast differentiation through myocardin related transcription factor A. *Scientific Reports*. 2021;11(1):1–13.

33. Choe K, Pak U, Pang Y, Hao W, Yang X. Advances and Challenges in Spatial Transcriptomics for Developmental Biology. *Biomolecules*. 2023;13(1):156.

34. Scoville DW, Kang HS, Jetten AM. GLIS1-3: Emerging roles in reprogramming, stem and progenitor cell differentiation and maintenance. *Stem Cell Investigation*. 2017;4(9).

35. Winderlich M, Keller L, Cagna G, Broermann A, Kamenyeva O, Kiefer F, Deutsch U, Nottebaum AF, Vestweber D. VE-PTP controls blood vessel development by balancing Tie-2 activity. *Journal of Cell Biology*. 2009;185(4):657–671.

36. McCracken IR, Dobie R, Bennett M, Passi R, Beqqali A, Henderson NC, Mountford JC, Riley PR, Ponting CP, Smart N, Brittan M, Baker AH. Mapping the developing human cardiac endothelium at single-cell resolution identifies MECOM as a regulator of arteriovenous gene expression. *Cardiovascular Research*. 2022;118(14):2960–2972.

37. Nightingale TD, Pattni K, Hume AN, Seabra MC, Cutler DF. Rab27a and MyRIP regulate the amount and multimeric state of VWF released from endothelial cells. *Blood*. 2009;113(20):5010–5018.

38. Norden PR, Sabine A, Wang Y, Demir CS, Liu T, Petrova T V., Kume T. Shear stimulation of FOXC1 and FOXC2 differentially regulates cytoskeletal activity during lymphatic valve maturation. *eLife*. 2020;9:1–35.

39. Govatati S, Pichavaram P, Janjanam J, Zhang B, Singh NK, Mani AM, Traylor JG, Orr AW, Rao GN. NFATc1-E2F1-LMCD1-Mediated IL-33 Expression by Thrombin Is Required for Injury-Induced Neointima Formation. *Arteriosclerosis, Thrombosis, and Vascular Biology*. 2019;39(6):1212–1226.

40. Ma Y, Halade G V., Zhang J, Ramirez TA, Levin D, Voorhees A, Jin YF, Han HC, Manicone AM, Lindsey ML. Matrix metalloproteinase-28 deletion exacerbates cardiac dysfunction and rupture after myocardial infarction in mice by inhibiting M2 macrophage activation. *Circulation Research*. 2013;112(4):675–688.

41. Li CH, Amar S. Role of secreted frizzled-related protein 1 (SFRP1) in wound healing. *Journal of Dental Research*. 2006;85(4):374–378.

42.   Pellicani R, Poletto E, Andreuzzi E, Paulitti A, Doliana R, Bizzotto D, Braghetta P, Colladel R, Tarticchio G, Sabatelli P, Bucciotti F, Bressan G, Iozzo R V., Colombatti A, Bonaldo P, et al. Multimerin-2 maintains vascular stability and permeability. *Matrix Biology*. 2020;87:11–25.

43.   Yang Y, Li D, Wu W, Huang D, Zheng H, Aihaiti Y. A pan-cancer analysis of the role of selenoprotein p mrna in tumorigenesis. *International Journal of General Medicine*. 2021;14:7471–7485.

44.   Ouadda ABD, He Y, Calabrese V, Ishii H, Chidiac R, Gratton JP, Roux PP, Lamarche-Vane N. CdGAP/ARHGAP31 is regulated by RSK phosphorylation and binding to 14-3-3β adaptor protein. *Oncotarget*. 2018;9(14):11646–11664.

45.   Liu H, Liu P, Shi X, Yin D, Zhao J. NR4A2 protects cardiomyocytes against myocardial infarction injury by promoting autophagy. *Cell Death Discovery*. 2018;4(1):1–11.

46.   Zhang K, Li M, Yin L, Fu G, Liu Z. Role of thrombospondin–1 and thrombospondin–2 in cardiovascular diseases (Review). *International Journal of Molecular Medicine*. 2020;45(5):1275–1293.

47.   Leatherdale A, Parker D, Tasneem S, Wang Y, Bonna A, Gross PL, Ni H, Doble B, Farndale RW, Hayward CPM. Multimerin-1 Binds to Specific Motifs in Vessel Wall Collagens and Contributes to Thrombosis: Novel Insights Regarding the Mechanisms That Support Human and Mouse Platelet Adhesion. *Blood*. 2017;130(Supplement 1):549.

48.   Jones M, Sabatini PJB, Lee FSH, Bendeck MP, Langille BL. N-cadherin upregulation and function in response of smooth muscle cells to arterial injury. *Arteriosclerosis, Thrombosis, and Vascular Biology*. 2002;22(12):1972–1977.

49.   Philippova M, Suter Y, Toggweiler S, Schoenenberger AW, Joshi MB, Kyriakakis E, Erne P, Resink TJ. T-cadherin is present on endothelial microparticles and is elevated in plasma in early atherosclerosis. *European Heart Journal*. 2011;32(6):760–771.

50.   He M, Huang TS, Li S, Hong HC, Chen Z, Martin M, Zhou X, Huang HY, Su SH, Zhang J, Wang WT, Kang J, Huang H Da, Zhang J, Chien S, et al. Atheroprotective Flow Upregulates ITPR3 (Inositol 1,4,5-Trisphosphate Receptor 3) in Vascular Endothelium via KLF4 (Krüppel-Like Factor 4)-Mediated Histone Modifications. *Arteriosclerosis, Thrombosis, and Vascular Biology*. 2019;39(5):902–914.

51.   Wang D, Liu B, Xiong T, Yu W, Yang H, Wang J, Jing X, She Q. Transcription factor Foxp1 stimulates angiogenesis in adult rats after myocardial infarction. *Cell Death Discovery*. 2022;8(1):1–9.

52.   Guan Y, Du Y, Wang G, Gou H, Xue Y, Xu J, Li E, Chan DW, Wu D, Xu P, Ni P, Xu D, Hu Y. Overexpression of PLXDC2 in Stromal Cell-Associated M2 Macrophages Is Related to EMT and the Progression of Gastric Cancer. *Frontiers in Cell and Developmental Biology*. 2021;9:1353.

53.   Cohen-Dvashi H, Ben-Chetrit N, Russell R, Carvalho S, Lauriola M, Nisani S, Mancini M, Nataraj N, Kedmi M, Roth L, Köstler W, Zeisel A, Yitzhaky A, Zylberg J, Tarcic G, et al. Navigator-3, a modulator of cell migration, may act as a suppressor of breast cancer progression. *EMBO Molecular Medicine*. 2015;7(3):299–314.

54.   Benz PM, Ding Y, Stingl H, Loot AE, Zink J, Wittig I, Popp R, Fleming I. AKAP12 deficiency impairs VEGF-induced endothelial cell migration and sprouting. *Acta Physiologica*. 2020;228(1).

55.   Wu JH, Fanaroff AC, Sharma KC, Smith LS, Brian L, Eipper BA, Mains RE, Freedman NJ, Zhang L. Kalirin promotes neointimal hyperplasia by activating rac in smooth muscle cells. *Arteriosclerosis, Thrombosis, and Vascular Biology*. 2013;33(4):702–708.

56.   Ding Y, Huang L, Xian X, Yuhanna IS, Wasser CR, Frotscher M, Mineo C, Shaul PW, Herz J. Loss of Reelin protects against atherosclerosis by reducing leukocyte-endothelial cell adhesion and lesion macrophage accumulation. *Science Signaling*. 2016;9(419).

57.   Jing Cao, Zhang G, Liu Z, Xu Q, Li C, Cheng G, Shi R. Peroxidasin promotes diabetic vascular endothelial dysfunction induced by advanced glycation end products via NOX2/HOCl/Akt/eNOS pathway. *Redox Biology*. 2021;45:102031.

58.   Wu W, Wang C, Zang H, Qi L, Azhar M, Nagarkatti M, Nagarkatti P, Cai G, Weiser-Evans MCM, Cui T. Mature vascular smooth muscle cells, but not endothelial cells, serve as the major cellular source of intimal hyperplasia in vein grafts. *Arteriosclerosis, Thrombosis, and Vascular Biology*. 2020;40(8):1870–1890.

59.   Déglise S, Bechelli C, Allagnat F. Vascular smooth muscle cells in intimal hyperplasia, an update. *Frontiers in Physiology*. 2023;13:2694.

60.   Ricard N, Bailly S, Guignabert C, Simons M. The quiescent endothelium: signalling pathways regulating organ-specific endothelial normalcy. *Nature Reviews Cardiology*. 2021;18(8):565–580.

61.   Li M, Qian M, Kyler K, Xu J. Endothelial–Vascular Smooth Muscle Cells Interactions in Atherosclerosis. *Frontiers in Cardiovascular Medicine*. 2018;5:151.

62.   Jin S, Guerrero-Juarez CF, Zhang L, Chang I, Ramos R, Kuan CH, Myung P, Plikus M V., Nie Q. Inference and analysis of cell-cell communication using CellChat. *Nature Communications 2021 12:1*. 2021;12(1):1–20.

63. Kumar P, Kumar S, Udupa EP, Kumar U, Rao P, Honnegowda T. Role of angiogenesis and angiogenic factors in acute and chronic wound healing. *Plastic and Aesthetic Research*. 2015;2(5):243.

64. Li H, Gao J, Zhang S. Functional and Clinical Characteristics of Cell Adhesion Molecule CADM1 in Cancer. *Frontiers in Cell and Developmental Biology*. 2021;9:2007.

65. Pulkkinen HH, Kiema M, Lappalainen JP, Toropainen A, Beter M, Tirronen A, Holappa L, Niskanen H, Kaikkonen MU, Ylä-Herttuala S, Laakkonen JP. BMP6/TAZ-Hippo signaling modulates angiogenesis and endothelial cell response to VEGF. *Angiogenesis*. 2021;24(1):129–144.

66. Ohtani K, Egashira K, Hiasa KI, Zhao Q, Kitamoto S, Ishibashi M, Usui M, Inoue S, Yonemitsu Y, Sueishi K, Sata M, Shibuya M, Sunagawa K. Blockade of vascular endothelial growth factor suppresses experimental restenosis after intraluminal injury by inhibiting recruitment of monocyte lineage cells. *Circulation*. 2004;110(16):2444–2452.

67. Lebleu VS, Kalluri R. Blockade of PDGF receptor signaling reduces myofibroblast number and attenuates renal fibrosis. *Kidney International*. 2011;80(11):1119–1121.

68. Banno H, Takei Y, Muramatsu T, Komori K, Kadomatsu K. Controlled release of small interfering RNA targeting midkine attenuates intimal hyperplasia in vein grafts. *Journal of Vascular Surgery*. 2006;44(3):633–641.

69. Van Der Horst EH, Frank BT, Chinn L, Coxon A, Li S, Polesso F, Slavin A, Ruefli-Brasse A, Wesche H. The growth factor midkine antagonizes VEGF signaling in vitro and in vivo. *Neoplasia*. 2008;10(4):340–346.

70. Rippe C, Morén B, Liu L, Stenkula KG, Mustaniemi J, Wennström M, Swärd K. NG2/CSPG4, CD146/MCAM and VAP1/AOC3 are regulated by myocardin-related transcription factors in smooth muscle cells. *Scientific Reports*. 2021;11(1).

71. Wang Y, Nakayama M, Pitulescu ME, Schmidt TS, Bochenek ML, Sakakibara A, Adams S, Davy A, Deutsch U, Lüthi U, Barberis A, Benjamin LE, Mäkinen T, Nobes CD, Adams RH. Ephrin-B2 controls VEGF-induced angiogenesis and lymphangiogenesis. *Nature*. 2010;465(7297):483–486.

72. Badila E. *Atherothrombotic risk factors and graft disease*.; 2016.

73. Kahle B, Schmidtke C, Hunzelmann N, Bartels C, Sievers HH, Steenbock H, Reinhardt DP, Brinckmann J. The Extracellular Matrix Signature in Vein Graft Disease. *Canadian Journal of Cardiology*. 2016;32(8):1008.e11-1008.e17.

74. Guida GA, Angelini GD. Pathophysiology and Mechanisms of Saphenous Vein Graft Failure. *Brazilian Journal of Cardiovascular Surgery*. 2022;37(Special Issue 1):32–37.

75. He C, Ye P, Zhang X, Esmaeili E, Li Y, Lü P, Cai C. The Role of TGF-β Signaling in Saphenous Vein Graft Failure after Peripheral Arterial Disease Bypass Surgery. *International Journal of Molecular Sciences*. 2023;24(12):10381.

76. Kenagy RD, Min SK, Clowes AW, Sandy JD. Cell death-associated ADAMTS4 and versican degradation in vascular tissue. *Journal of Histochemistry and Cytochemistry*. 2009;57(9):889–897.

77. Wesseling M, Sakkers TR, de Jager SCA, Pasterkamp G, Goumans MJ. The morphological and molecular mechanisms of epithelial/endothelial-to-mesenchymal transition and its involvement in atherosclerosis. *Vascular Pharmacology*. 2018;106:1–8.

78. De Vries MR, Simons KH, Jukema JW, Braun J, Quax PHA. Vein graft failure: From pathophysiology to clinical outcomes. *Nature Reviews Cardiology*. 2016;13(8):451–470.

79. Allaire E, Clowes AW. Endothelial cell injury in cardiovascular surgery: The intimal hyperplastic response. *Annals of Thoracic Surgery*. 1997;63(2):582–591.

80. Davies MG, Hagen P -O. Pathobiology of intimal hyperplasia. *British Journal of Surgery*. 1994;81(9):1254–1269.

81. Newby AC, Zaltsman AB. Molecular mechanisms in intimal hyperplasia. *Journal of Pathology*. 2000;190(3):300–309.

82. Fernandez DM, Giannarelli C. Immune cell profiling in atherosclerosis: role in research and precision medicine. *Nature Reviews Cardiology*. 2022;19(1):43–58.

83. Mylonaki I, Allain E, Strano F, Allémann E, Corpataux JM, Meda P, Jordan O, Delie F, Rougemont AL, Haefliger JA, Saucy F. Evaluating intimal hyperplasia under clinical conditions. *Interactive Cardiovascular and Thoracic Surgery*. 2018;27(3):427–436.

84. Pober JS, Sessa WC. Evolving functions of endothelial cells in inflammation. *Nature Reviews Immunology*. 2007;7(10):803–815.

85. Gimbrone MA, García-Cardeña G. Endothelial Cell Dysfunction and the Pathobiology of Atherosclerosis. *Circulation Research*. 2016;118(4):620–636.

86. Sartore S, Chiavegato A, Faggin E, Franch R, Puato M, Ausoni S, Pauletto P. Contribution of Adventitial Fibroblasts to Neointima Formation and Vascular Remodeling. *Circulation Research*. 2001;89(12):1111–1121.

87. Tinajero MG, Gotlieb AI. Recent Developments in Vascular Adventitial Pathobiology: The Dynamic Adventitia as a Complex Regulator of Vascular Disease. *The American journal of pathology*. 2020;190(3):520–534.

88. Shi Y, O'Brien JE, Mannion JD, Morrison RC, Chung W, Fard A, Zalewski A. Remodeling of autologous saphenous vein grafts. The role of perivascular myofibroblasts. *Circulation*. 1997;95(12):2684–93.

89. Roy-Chaudhury P, Wang Y, Krishnamoorthy M, Zhang J, Banerjee R, Munda R, Heffelfinger S, Arend L. Cellular phenotypes in human stenotic lesions from haemodialysis vascular access. *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association*. 2009;24(9):2786–91.

90. de Vries MR, Quax PHA. Inflammation in Vein Graft Disease. *Frontiers in cardiovascular medicine*. 2018;5:3.

91. Cha JK, Jeong MH, Bae HR, Han JY, Jeong SJ, Jin HJ, Lim YJ, Kim SH, Kim JW. Activated platelets induce secretion of interleukin-1beta, monocyte chemotactic protein-1, and macrophage inflammatory protein-1alpha and surface expression of intercellular adhesion molecule-1 on cultured endothelial cells. *Journal of Korean medical science*. 2000;15(3):273–8.

92. He C, Ye P, Zhang X, Esmaeili E, Li Y, Lü P, Cai C. The Role of TGF-β Signaling in Saphenous Vein Graft Failure after Peripheral Arterial Disease Bypass Surgery. *International journal of molecular sciences*. 2023;24(12).

93. Yuan S-M, Wang Y-Q, Shen Y, Jing H. Transforming growth factor-β in graft vessels: histology and immunohistochemistry. *Clinics (Sao Paulo, Brazil)*. 2011;66(5):895–901.

94. Schachner T, Laufer G, Bonatti J. In vivo (animal) models of vein graft disease. *European Journal of Cardiothoracic Surgery*. 2006;30(3):451–463.

95. Gurkan S, Gur O, Yuksel V, Tastekin E, Huseyin S, Gur DO, Canbaz S. The effect of distension pressure on endothelial injury and vasodilatation response in saphenous vein grafts: Conversion of a bypass graft to a dead pipe. *Kardiochirurgia i Torakochirurgia Polska*. 2014;11(2):119–125.

96. LoGerfo FW, Quist WC, Crawshaw HM, Haudenschild C. An improved technique for preservation of endothelial morphology in vein grafts. *Surgery*. 1981;90(6):1015–1024.

97. Baumann FG, Catinella FP, Cunningham JN, Spencer FC. Vein contraction and smooth muscle cell extensions as causes of endothelial damage during graft preparation. *Annals of Surgery*. 1981;194(2):199–211.

98. Harris PL, Veith FJ, Shanik GP, Nott D, Wengerter KR, Moore DJ. Prospective randomized comparison of in situ and reversed infrapopliteal vein grafts. *British Journal of Surgery*. 1993;80(2):173–176.

99. Souza D. A new no-touch preparation technique: Technical notes. *Scandinavian Cardiovascular Journal*. 1996;30(1):41–44.

Figure 1. Overview of the workflow used for sample analysis. (A) Canine vein graft samples were retrieved following distension, then analyzed via single-nuclei RNA-seq (snRNA-seq) and spatial transcriptomics (ST) through the 10x Genomics platform. Single-nuclei-based transcriptomics enables the sequencing of difficult-to-dissociate samples while preserving detailed individual cellular resolution. As a complementary approach, ST provides insight into the spatial gene expression of the tissue, with the transcriptomes of 1-10 cells captured per barcoded spot of the sample slide. Combined, these approaches illuminate the unique cellular architecture and transcriptomic landscape of the tissue. **(B) Varying permeabilization times were assessed for fluorescent cDNA footprint to optimize tissue permeabilization**. The left panel shows the brightfield images (H&E), and the right panel shows fluorescent images (Cy3, Cy5 Multispectral Imaging (MSI)). **(C)** Images of H&E strained veins. Notably, the lower panel illustrates the expansion of the lumen following distension. **(D)** Overview of the developed method for adding histological annotations to ST samples for targeted downstream analyses. Capture spots are projected onto the H&E image as voxels, generating a spatial plot that can be annotated with STannotate and used for analysis.

Figure 2. Spatial transcriptomic analysis of distended veins. (A) Top: Spatial plots displaying capture spots (voxels) projected onto H&E images of control (left) and distended (right) veins from the same canine, annotated using our ST annotation tool. Bottom: Volcano plot of significantly differentially expressed genes ($P < 0.00001$) within the distended vein compared to the control vein, wherein each gene is colored according to the vein wall layer with the highest expression of the marker. **(B)** Enriched pathways (via Reactome database) within each layer of the distended veins compared to control veins, based on significantly upregulated genes between each layer ($P < 0.05$). **(C)** Network of key differentially expressed genes and associated pathways ($P < 0.05$) between control and distended veins, derived from clusterProfiler analysis using the Reactome pathways database.

Figure 3. Spatial clustering of control and distended veins. (A) UMAP displaying the clustering of spatial transcriptomic voxels (containing 1-10 cells each) derived from control and distended veins. Clusters are derived based on similar transcriptomic profiles correlating to similar cell type compositions. Cluster labels were designated by the predominant localization of associated voxels to either the intima (I), media (M), or adventitia (A) of the vein samples. The spatial distribution of these clusters was then mapped onto the vein samples. **(B)** Mapping of spatial clusters onto a control vein sample. **(C)** Mapping of spatial clusters onto a distended vein sample from the same canine. **(D)** Relative proportions of each spatial cluster present within control vein and distended vein groups. **(E)** Heatmap of marker genes associated with different spatial clusters. Columns represent individual voxels grouped by spatial cluster, while rows display individual genes. Horizontal colored bars above the heatmap indicate the different spatial cellular clusters. Relative gene expression is shown in pseudo color, where blue represents low expression and red represents high expression. Enriched biological pathways (via clusterProfiler using Reactome database) based on differentially upregulated genes between the **(F)** M1 and M2 clusters, as well as the **(G)** A1 and A2 clusters (*P adjusted* < 0.05). Gene ratio is defined as the proportion of upregulated genes present in the cluster that overlap with the respective pathway. The multiple test correction for the P value has been performed using the Benjamini-Hochberg (BH) approach.

Figure 4. Single-nuclei transcriptomic analysis of control and distended veins. (A) UMAP displaying the clustering of nuclei derived from control and distended veins. Canonical cell types are based on the expression of marker genes (Figure S3). **(B)** Relative proportions of each cluster present within control vein and distended vein groups. **(C)** Volcano plot of differentially expressed genes within the distended vein compared to the control vein ($P < 0.00001$), wherein each gene is colored according to the cluster with the highest expression of the marker. **(D)** Heatmap of top marker genes associated with different cell types. Columns represent individual cells grouped by cell types, while rows display individual genes. Horizontal colored bars above the heatmap indicate the different cell types. Relative gene expression is shown in pseudo color, where blue represents low expression and red represents high expression. Top marker genes for clusters are defined by the average $\log_2$ fold-change of genes expressed in greater than 30% of nuclei within a cluster and $P < 0.05$. **(E)** Network of key differentially enriched pathways between clusters, derived from clusterProfiler analysis using the Reactome database ($P < 0.05$). Each node depicts the proportion of each cluster displaying enrichment of the associated pathway and the cumulative number of genes enriched in the pathway between the represented clusters.

Figure 5. Characterization of fibroblast subpopulations. (A) Volcano plot of differentially expressed genes within fibroblast subclusters of the distended vein compared to the control vein ($P < 0.001$), wherein each gene is colored according to the fibroblast subpopulation with the highest expression of the marker. **(B)** Heatmap of top marker genes associated with different fibroblast subclusters. Columns represent individual cells grouped by cell types, while rows display individual genes. Horizontal colored bars above the heatmap indicate the different cell types. Relative gene expression is shown in pseudo color, where blue represents low expression and red represents high expression. Top markers for each fibroblast cluster are defined by the average log$_2$ fold-change and $P < 0.05$. **(C)** Relative proportions of each fibroblast subpopulation present within control vein and distended vein groups. **(D)** Dot plot of the top markers representing the phenotype of each fibroblast subpopulation. The relative expression and percent of cells expressing specific markers are shown by shades of blue and the dot size, respectively.

Figure 6. Characterization of endothelial cell subpopulations. (A) Volcano plot of differentially expressed genes within endothelial cells of the distended vein compared to the control vein, wherein each gene is colored according to the endothelial cell subpopulation with the highest expression of the marker. **(B)** Heatmap of top marker genes associated with different endothelial sub-clusters. Columns represent individual cells grouped by cell types, while rows display individual genes. Horizontal colored bars above the heatmap indicate the different cell types. Relative gene expression is shown in pseudo color, where blue represents low expression and red represents high expression. Top marker genes between endothelial clusters are defined by the average log₂ fold-change and $P <$ 0.05. **(C)** Relative proportions of each endothelial cell subpopulation present within control and distended vein groups. **(D)** Dot plot of the top markers representing the phenotype of each endothelial cell subpopulation. The relative expression and percent of cells expressing specific markers are shown by shades of blue and the dot size, respectively.

Figure 7. Deconvolution of spatial transcriptomic data through the integration of single-nuclei data. (A) Summary of distinct fibroblast and endothelial subpopulations identified in control and distended veins via snRNA-seq analysis along with expression ok key markers. **(B)** Heatmap displaying the differential enrichment scores of cellular subpopulations across layers of vein wall following distension. An increased differential score (red) indicates increased gene signature expression associated with the respective sub-population in the distended veins relative to control veins, and blue indicates increased enrichment in control veins. **(C)** Heatmap of neighborhood enrichment Z-scores generated via Squidpy, illustrating the proximity of cellular populations to one another based on the dominant cell type assigned to each voxel. Red indicates enriched proximity between two populations, whereas blue indicates depleted proximity. **(D)** Spatial plots of representative control and distended vein samples displaying the dominant cell type within each voxel predicted by model-based deconvolution of the spatial transcriptomic data using the single-nuclei dataset via Cell2loca-tion. **(E)** Validation of the enrichment score (left) and model-based (panel D) integration approaches using RNAscope (right).

Figure 8. Intercellular communication analysis. Network displaying the number of intercellular communication interactions between subpopulations for control **(A)** and distended veins **(B)**. **(C)** The proportion of key signaling molecules active in control versus distended veins. **(D)** Heatmap of the differential signaling pattern (both incoming and outgoing signaling per cell type or subpopulation) in distended veins (red) compared to control veins (blue). **(E)** Chord diagrams illustrate intercellular communication networks for distended veins, with the exception of BMP, which is illustrated for the control vein. The outer ring and arrows of the diagram represent the signaling produced by each cellular subpopulation, while the inner ring represents which subpopulations are receiving the outgoing signals. **(F)** Network depicting the interactions of key signaling molecules and differentially expressed genes across pathways implicated in distension injury.

Figure 9. Relationship of pre- and post-implantation transcriptomic profiles and select protein expression. (A) Relative proportions of each fibroblast subpopulation present within control vein and distended vein groups. **(B)** Spatial plot of grafted vein sample displaying the dominant cell type within each voxel based on deconvolution of the spatial transcriptomic data using the single-nuclei dataset via Cell2location. **(C)** Temporally resolved multilayer gene network analysis illustrating the effects of key genes (circled) at the time of vein harvesting and distension (left) to 24 hours post-implantation (right). **(D)** Immunohistochemistry staining of control, distended, and grafted veins with VCAN, GLIS3, or FBN with quantification values (left).