Title: An immunohistochemical atlas of necroptotic pathway expression

Authors: Shene Chiou[1,2,^], Aysha H. Al-Ani[1,2,3,^], Yi Pan[1], Komal M. Patel[1], Isabella Y. Kong[4], Lachlan W. Whitehead[1,2], Amanda Light[1], Samuel N. Young[1], Marilou Barrios[1,2], Callum Sargeant[1,2], Pradeep Rajasekhar[1,2], Leah Zhu[1], Anne Hempel[1], Ann Lin[1], James A. Rickard[1,6], Cathrine Hall[1], Pradnya Gangatirkar[1], Raymond K.H. Yip[1,2], Wayne Cawthorne[1,2], Annette V. Jacobsen[1,2], Christopher R. Horne[1,2], Katherine R. Martin[7], Lisa J. Ioannidis[1,2], Diana S. Hansen[1,2,5], Jessica Day[1,2,3], Ian P. Wicks[1,2], Charity Law[1,2], Matthew E. Ritchie[1,2], Rory Bowden[1,2], Joanne M. Hildebrand[1,2], Lorraine A. O'Reilly[1,2], John Silke[1,2], Lisa Giulino-Roth[4], Ellen Tsui[1], Kelly L. Rogers[1,2], Edwin D. Hawkins[1,2], Britt Christensen[1,2,3], James M. Murphy[1,2,7,*] and André L. Samson[1,2,*].

Affiliations: [1]Walter and Eliza Hall Institute of Medical Research, Parkville, Australia

[2]University of Melbourne, Parkville, Australia

[3]Royal Melbourne Hospital, Parkville, Australia

[4]Pediatric Hematology/Oncology, Weill Cornell Medical College, New York, USA

[5]Monash Biomedicine Discovery Institute, Department of Microbiology, Monash University, Clayton, Australia

[6]Austin Hospital, Heidelberg, Australia

[7]Drug Discovery Biology, Monash Institute of Pharmaceutical Sciences, Parkville, Australia

[^]These authors contributed equally to this work

*Joint senior and corresponding authors: jamesm@wehi.edu.au (JMM) and samson.a@wehi.edu.au (ALS).

Highlights:

- 13 automated immunohistochemistry protocols for detecting the necroptotic pathway
- Necroptotic pathway expression is confined to fast-cycling immune barriers
- Necroptotic pathway expression changes at sites of immunoinflammatory challenge
- Immunodetection of necrosomes in IBD patients is a putative new diagnostic tool

Abstract: Necroptosis is a lytic form of regulated cell death reported to contribute to inflammatory diseases of the gut, skin and lung, as well as ischemic-reperfusion injuries of the kidney, heart and brain. However, precise identification of the cells and tissues that undergo necroptotic cell death *in vivo* has proven challenging in the absence of robust protocols for immunohistochemical detection. Here, we provide automated immunohistochemistry protocols to detect core necroptosis regulators – Caspase-8, RIPK1, RIPK3 and MLKL – in formalin-fixed mouse and human tissues. We observed surprising heterogeneity in protein expression within tissues, whereby short-lived immune barrier cells were replete with necroptotic effectors, whereas long-lived cells lacked RIPK3 or MLKL expression. Local changes in the expression of necroptotic effectors occurred in response to insults such as inflammation, dysbiosis or immune challenge, consistent with necroptosis being dysregulated in disease contexts. These methods will facilitate the precise localisation and evaluation of necroptotic signaling *in vivo*.

Main text:

Introduction

The necroptotic cell death pathway leads to cell lysis and expulsion of cellular contents into the extracellular milieu, which in turn provokes an innate immune response. Necroptosis is considered to be an altruistic cell death pathway whose principal role is to protect the host from pathogens (Fletcher-Etherington *et al*, 2020; Liu *et al*, 2021; Palmer *et al*, 2021; Pearson *et al*, 2017; Petrie *et al*, 2019; Upton *et al*, 2010; Yeap & Chen, 2022; Zhang *et al*, 2020). Despite this, it is the aberrant functions of necroptosis associated with inflammatory diseases that have spurred interest in its underlying mechanisms and therapeutic prospects (Choi *et al*, 2019; Fang *et al*, 2021). Studies of mice lacking the terminal effectors of the pathway – RIPK3 (Receptor-interacting protein kinase-3) or MLKL (Mixed-lineage kinase domain-like) – have led to the concept that excess necroptosis drives a range of inflammatory pathologies in organs including the skin, gut, brain, heart, lung, kidney and testes (Devos *et al*, 2020; Gunther *et al*, 2011; Ito *et al*, 2016; Li *et al*, 2017; Linkermann *et al*, 2013; Lu *et al*, 2021; Luedde *et al*, 2014; Naito *et al*, 2020). However, many of these attributions have been disputed (Dominguez *et al*, 2021; Newton *et al*, 2016; Wang *et al*, 2020b), likely reflecting an evolving understanding of the pathway and the limited availability of validated reagents to interrogate necroptosis in pathological specimens.

The core signaling axis of the necroptotic pathway has been well-defined and can be activated in response to a variety of inflammatory cues including ligation of Death, Toll-like or Pathogen-Pattern receptors (Chen *et al*, 2022b; Cho *et al*, 2009; Degterev *et al*, 2008; He *et al*, 2011; He *et al*, 2009; Kaiser *et al*, 2013; Kaiser *et al*, 2011). Caspase-8 is a critical negative regulator of necroptotic signaling (Kaiser *et al*., 2011), whereby its deletion or loss-of-function promotes oligomerisation of RIPK1 (Receptor-interactor protein kinase-1), TRIF (TIR domain-containing adapter molecule 1) and/or ZBP1 (Z-DNA-binding protein

1) (Kaiser *et al.*, 2013). This oligomeric structure, otherwise known as the necrosome, promotes activation of the downstream effectors RIPK3 and MLKL (Samson *et al*, 2021b). RIPK3 recruits the MLKL pseudokinase to the necrosome, where it phosphorylates MLKL to provoke a conformational change, release from the necrosome, oligomerization and trafficking to the plasma membrane (Garnish *et al*, 2021; Murphy *et al*, 2013; Samson *et al*, 2020; Sun *et al*, 2012; Wang *et al*, 2014; Zhao *et al*, 2012). At the plasma membrane, accumulation of activated MLKL to a critical threshold level is required for membrane permeabilization via a poorly understood mechanism that brings about the cell's demise (Chen *et al*, 2014; Hildebrand *et al*, 2014; Samson *et al.*, 2020).

As our understanding of the necroptosis pathway has grown, new tools and protocols have been developed to study necroptotic signaling in fixed cultured cells (Rodriguez *et al*, 2016; Samson *et al*, 2021a; Wang *et al.*, 2014; Webster *et al*, 2018). However, robust procedures for assessing the necroptotic pathway in tissues are still lacking, often leading to contradictory reports in the literature and misattributions of necroptotic pathologies. Here, we report automated immunostaining protocols for detecting Caspase-8, RIPK1, RIPK3 and MLKL, in mouse formalin-fixed paraffin-embedded tissues. These procedures have enabled the assembly of an atlas of necroptotic pathway expression in mouse tissues under basal conditions and during innate immune challenge. While the necroptosis machinery is rarely expressed in cell types other than short-lived barrier cells, sterile inflammation increased RIPK3 expression in the gut and liver, broadly predisposing multiple cell types to necroptotic death. In contrast, elimination of the intestinal microflora diminished ~~RIPK3~~ expression of RIPK3 and MLKL to reduce necroptotic propensity in the gut. RIPK3 is also uniquely upregulated in splenic germinal centres suggesting it may have a non-necroptotic role in humoral immunity. Furthermore, we present robust protocols for detecting human Caspase-8, RIPK1, RIPK3 and MLKL and illustrate their utility for detecting dysregulated necroptosis in biopsies from patients

101   with inflammatory bowel disease (IBD). Collectively, these protocols will empower the

102   definitive evaluation of where and when necroptosis occurs in vivo in health and disease.

Results:

Standardised immunohistochemical detection of the necroptotic pathway in mouse tissues

We recently compiled a toolbox of immunofluorescence assays to detect necroptotic signaling in cells (Samson *et al.*, 2021a). This toolbox requires use of: 1) non-crosslinking fixatives and 2) gene knockouts to account for non-specific signals; requirements that often cannot be met when immunostaining tissues. Here we aimed to develop robust immunohistochemistry protocols to detect the necroptotic pathway in formalin-fixed paraffin-embedded mouse tissues. Embedding and immunostaining was performed in an automated manner (see *Methods*) to allow reliable and scalable detection of the necroptotic pathway, and to lessen the future need to account for non-specific immunosignals using appropriate gene knockout controls. The specificity of thirteen monoclonal antibodies against Caspase-8, RIPK1, RIPK3 or MLKL was first tested by immunoblotting spleen homogenates from wild-type versus knockout mice (Fig. EV1). The intensity and specificity of antibodies for immunohistochemistry was then iteratively optimised across 21 conditions (see *Methods* and Fig. EV1). At each optimisation step, immunohistochemical signals from the spleen of wild-type versus knockout mice were quantified (Fig. 1Ai), ratioed (Fig. 1Aii) and integrated to yield an index of performance (Fig. 1Aiii). For example, this pipeline improved the detection of RIPK1 with the monoclonal antibody D94C12 by approximately three orders of magnitude (Fig. EV1). In total, seven automated immunohistochemistry protocols to detect mouse Caspase-8, RIPK1, RIPK3 or MLKL were developed (Fig. 1B). The detection of Caspase-8, RIPK1 and RIPK3 using these immunohistochemistry protocols (Fig. 1B) closely aligned with the abundance of these proteins across multiple tissues as measured by high-resolution quantitative mass spectrometry (Fig. 1C), indicating both specificity and sensitivity. Despite many rounds of optimisation with three specific anti-MLKL antibodies, mouse MLKL

remained difficult to detect via immunohistochemistry in all tissues except the spleen (Fig.

1B,D and **EV1**).

Basal expression of the necroptotic pathway is restricted to fast-cycling immune

barriers.

The immunohistochemical profile of Caspase-8, RIPK1, RIPK3 and MLKL across seven

different organs suggested that expression of the necroptotic pathway is heavily restricted

in unchallenged mice (Fig. **1D**). For example, RIPK3$^+$ cells were scarce in the kidney and

heart, and RIPK3 was undetectable in the brain (Fig. **1D**). By comparison, co-expression

of Caspase-8, RIPK1 and RIPK3 was evident in intestinal epithelial cells, some splenic

regions and Kupffer cells (Fig. **2A**). The *Tabula Muris* single cell RNA sequencing dataset

supports the conclusion that expression of the necroptotic pathway is highly restricted in

mice (Fig. **EV2A**; (Tabula Muris *et al*, 2018)). Transcript expression of the necroptotic

effectors, MLKL and RIPK3, was below detection limits in kidney epithelial, cardiac muscle

and resident brain cells, but was frequently detected in progenitor and immune barrier cell

populations (Fig. **EV2A**; (Tabula Muris *et al.*, 2018)).

Close inspection of sites where the necroptotic pathway was constitutively

expressed showed unexpected layers of spatial regulation (Fig. **2A-B**). In the epithelial

barrier of the ileum, Caspase-8 expression was lower in crypts and higher at villi tips,

whereas RIPK3 levels peaked in the transit amplifying region and decreased towards the

villus tip (Fig. **2Ai, 2Bi**). In the colonic epithelial barrier, both Caspase-8 and RIPK3

exhibited higher expression in the base of the crypt and decreased towards the tip of the

crypt (Fig. **2Aii, 2Bii**). It is noteworthy that expression patterns of Caspase-8 and RIPK3

differ between the small and large intestine because these organs exhibit distinct cell

death responses to the same inflammatory stimuli (e.g. TNF) or genetic deficiency (e.g.

deletion of *Casp8* or *Fadd*) (Bader *et al*, 2023; Schwarzer *et al*, 2020; Tisch *et al*, 2022;

Zelic *et al*, 2018). In liver, Kupffer cells expressed Caspase-8, RIPK1 and RIPK3

(arrowheads Fig. **2Aiii, 2Biii**), whereas hepatocytes expressed Caspase-8 and RIPK1, but

not RIPK3. Interestingly, Caspase-8 levels were higher in pericentral hepatocytes than in

periportal hepatocytes (arrow Fig. **2Aiii, 2Biii**). Zonation of the necroptotic pathway was

also evident in the spleen, with Caspase-8, RIPK1, RIPK3 and MLKL levels peaking in the

marginal zone, where circulating antigens are trapped for immune presentation (Fig. **2Aiv,**

EV2B). Prior spatial transcriptomics data support the conclusion that necroptotic potential

is zoned along the intestinal crypt-to-villus axis (Fig. **EV2C**; (Moor *et al*, 2018)) and along

the hepatic central-to-portal axis (Fig. **EV2D**; (Ben-Moshe *et al*, 2019)). We also used

spatial transcriptomics on mouse spleen to confirm that necroptotic potential peaks in the

marginal zone (Fig. **EV2E-G**).

The expression of RIPK3 appears to be under particularly strict spatial control. For

example, in the ileum, RIPK3 levels were high in fast-cycling epithelial progenitors, but low

in adjacent, terminally differentiated Paneth cells (open arrowhead Fig. **2Ai**). Published

single cell transcriptomics data supports the conclusion that Paneth cells express low

levels of RIPK3 under basal conditions (Haber *et al*, 2017). As another example of

differential expression, RIPK3 levels were high in fast-cycling colonic epithelial cells, but

undetectable in slow-cycling renal epithelial cells (Fig. **1D**). These observations suggest

that RIPK3 expression is linked to cell turnover. Indeed, across 103 cell ontologies in the

Tabula Muris dataset, gene expression of cell cycle markers *Top2a* and *Mki67* correlated

with the expression of *Ripk3,* but not *Ripk1* (Fig. **2C** and **EV2A**; (Tabula Muris *et al.*,

2018)). Prior cell cultures studies further suggest that the expression and function of

RIPK3 fluctuates during the mitotic cell cycle (Gupta & Liu, 2021; Liccardi *et al*, 2019).

Altogether, by applying a set of optimised immunohistochemistry protocols to

multiple organs, we have found that the necroptotic pathway is preferentially expressed at

fast-cycling immune barriers under basal conditions. Such targeted expression is

consistent with the evolutionary origin of necroptosis being an anti-pathogen defence

measure (Palmer *et al.*, 2021; Petrie *et al.*, 2019; Upton *et al.*, 2010). We further find that

necroptotic potential is spatially graded along barriers such as the intestinal mucosa.

These gradations in the availability of cell death mediators along barriers likely allow

multiple cell death programs to be flexibly deployed against invading pathogens (Cook *et*

al, 2014; Doerflinger *et al*, 2020).

Inflammation, dysbiosis or immune challenge trigger local changes in RIPK3

expression

To demonstrate scalability, we used automated immunohistochemistry to characterise the

expression of Caspase-8, RIPK1 and RIPK3 across six tissues during TNF-induced

Systemic Inflammatory Response Syndrome (SIRS) – a widely-used model of RIPK-

dependent pathology (Fig. **3A**; (Duprez *et al*, 2011; Harris *et al*, 2017; Newton *et al.*, 2016;

Newton *et al*, 2014; Zelic *et al.*, 2018)). Littermate wild-type mice were intravenously

administered TNF, or vehicle, and tissues were harvested 9 hours later when symptoms

such as hypothermia were manifesting (Fig. **3B**). No major changes to Caspase-8 or

RIPK1 expression were observed after TNF administration, except for an unidentified

population of RIPK1-expressing cells appearing at the onset of apoptosis in lymphoid

tissues (Fig. **3C-D**; arrowhead). By comparison, RIPK3 was upregulated in intestinal

epithelial cells (Fig. **3E-F**), certain vascular beds (Fig. **3G-H**) and in liver (Fig. **3I-J**); the

main sites where RIPK1- and RIPK3-mediated signaling during SIRS has been implicated

by knock-in and knockout mouse studies (Duprez *et al.*, 2011; Newton *et al.*, 2016; Zelic *et*

al., 2018). In contrast, RIPK3 levels were not increased in resident cells of the kidney or

heart in TNF-treated mice. Our data therefore suggest that targeted upregulation of RIPK3

in resident cells of the gut and liver underlies RIPK-mediated pathology in SIRS. TNF-

treatment also changed the pattern of RIPK3 expression in the intestine, potentially

skewing cell death responses in the inflamed gut (Fig. **3E-F**). It was surprising that RIPK3

was detected in peri-portal hepatocytes after TNF administration, given that RIPK3 is

epigenetically silenced in hepatocytes under basal conditions (Preston *et al*, 2022).

Collectively, our immunohistochemical characterisation of the SIRS mouse model leads us

to propose that RIPK3 is regulated akin to a positive acute phase reactant, with hepatic

and intestinal expression that rapidly increases in response to inflammation. In support of

this notion, we find that TNF-treatment increased the levels of RIPK3 in serum (Fig. **3K-L**).

Moreover, RIPK1-inhibition prevented both TNF-induced hypothermia and the release of

RIPK3 into blood (Fig. **3K-L**). These data raise the exciting possibilities that RIPK3 is a

novel acute phase reactant, and that circulating levels of RIPK3 are a surrogate measure

of RIPK1-mediated signaling.

Next, we addressed whether microbiota-depletion affects necroptotic pathway

expression. This question was prompted by studies showing that antibiotics offer

protection in various models of intestinal necroptosis (Bader *et al.*, 2023; Gunther *et al*,

2015; Li *et al*, 2020; Xie *et al*, 2020). As shown in Fig. **4A**, a litter of wild-type mice was

split into two cages and the water for one cage was supplemented with antibiotics for 6

days. As expected, the cecum of antibiotic-treated mice was enlarged and canonical anti-

microbial factors such as lysozyme and REG-3β were reduced in the ileum, but not the

spleen, of antibiotic-treated mice (Fig. **4B-C**). These predictable responses to microbiota-

depletion also coincided with a lowering of RIPK3 and MLKL gene and protein expression

in the ileum, but not the spleen (Fig. **4B-C** and **EV3**). In contrast, Caspase-8 gene and

protein expression in the ileum were unaffected by microbiota-depletion, whereas ileal

RIPK1 protein levels were increased by antibiotic treatment (Fig. **4B-C** and **EV3**). Similar

trends were observed by immunohistochemistry, with epithelial Caspase-8 expression

remaining constant, while RIPK1 levels were elevated and RIPK3 expression reduced in

the crypt and transit amplifying regions of the ileum in antibiotic-treated mice (Fig. **4D**).

Unexpectedly, immunohistochemistry also showed that microbiota-depletion triggered

cytoplasmic accumulations of RIPK1 and RIPK3 in enterocytes at villi tips (Fig. **4D**;

arrowheads). These RIPK1$^+$ RIPK3$^+$ Caspase-8$^-$ clusters are unlikely to be necrosomes,

as no corresponding phospho-activation of RIPK1 or MLKL was observed (Fig. **EV3**).

Instead, these clusters may be due to a microbe-related function, such as

lipopolysaccharide handling, that is preferentially performed by enterocytes at villi tips

(Berkova *et al*, 2023; Ge *et al*, 2000). These changes to RIPK1/3 could also be related to

the reduced epithelial turnover that accompanies microbiota-depletion (Park *et al*, 2016).

Overall, we find that expression of the necroptotic pathway responds locally to changes in

the microbiome. This response is spatially restricted to the small intestine, zoned along the

crypt-to-villus axis, and warrants further investigation given that dysbiosis often occurs in

cell death-associated gut disorders such as Crohn's disease (Gevers *et al*, 2014).

Lastly, we used automated immunohistochemistry to uncover a potential non-

necroptotic role for RIPK3 in adaptive immunity. We immunised wild-type mice with the

model ligand, NP-KLH (4-hydroxy-3-nitrophenylacetyl hapten conjugated to keyhole limpet

hemocyanin), and harvested tissues 5 or 14 days later when antigen-specific antibody

responses were detectable (Fig. **EV4A**). No changes in the levels or zonation of Caspase-

8, RIPK1 or RIPK3 were noted in the ileum of immunised mice and no marked differences

in the expression of Caspase-8, RIPK1 or MLKL were observed in the spleen 14 days after

immunisation. However, RIPK3 levels were markedly elevated in Ki67$^+$ germinal centres

(arrowheads Fig. **EV4B**). This finding suggests that RIPK3 may have a non-necroptotic

role in antibody production. To investigate this possibility, *Ripk3$^{+/+}$* and *Ripk3$^{-/-}$* littermate

mice were immunised with NP-KLH and humoral immune responses were measured in

blood and spleen (Fig. **EV4C-J**). RIPK3-deficiency did not alter the circulating levels of

antigen-specific antibodies 5 days after immunisation (Fig. **EV4C-D**). Similarly, RIPK3-

deficiency did not influence the number of class-switched B cells in the spleen (Fig. **EV4E-**

F), the number of antigen-specific plasma cells in the spleen (Fig. **EV4G**), or the amount of

circulating antigen-specific antibodies 14 days after immunisation (Fig. **EV4H-J**). Thus,

consistent with prior studies (Newton *et al*, 2004), RIPK3 does not overtly affect early

antigen-specific antibody responses. Future studies should explore a role of RIPK3 in

splenic germinal centres, especially given that RIPK3 has a mechanistically undefined

non-necroptotic role during lymphoproliferative disease (Alvarez-Diaz *et al*, 2016).

In summary, by employing a toolbox of automated immunohistochemical stains, we

find that expression of the necroptotic pathway, in particular RIPK3, is responsive to

inflammation, dysbiosis or immunisation. These context-specific changes are tightly

regulated across space and time, underscoring the need for robust, scalable, *in situ*

assays to pinpoint necroptotic pathway expression and activation.

Automated immunohistochemical detection of the human necroptotic pathway

Important differences exist between the human and mouse necroptotic pathways (Chen *et*

al, 2013; Davies *et al*, 2020; Horne *et al*, 2023; Petrie *et al*, 2018; Samson *et al.*, 2021b;

Sun *et al.*, 2012; Tanzer *et al*, 2016). For instance, the primary sequence of RIPK3 and

MLKL are poorly conserved between species (Horne *et al.*, 2023), and humans uniquely

express Caspase-10 which likely negates necroptotic signaling (Ramirez & Salvesen,

2018; Tanzer *et al*, 2017). Thus, in parallel to developing assays for the murine necroptotic

pathway, seventeen antibodies against Caspase-8, Caspase-10, RIPK1, RIPK3 or MLKL

were tested on wild-type versus knockout formalin-fixed paraffin-embedded HT29 human

cells via immunoblot, and then iteratively optimised for immunohistochemistry (see

Methods and Fig. **EV5**). While Caspase-10 remained refractory to immunohistochemical

detection, six automated immunohistochemistry protocols were developed for human

Caspase-8, RIPK1, RIPK3 or MLKL (Fig. **EV5**).

These immunohistochemistry protocols detected diffuse cytoplasmic signals for Caspase-

8, RIPK1, RIPK3 and MLKL in cells under resting conditions (Fig. 5), and in cells

undergoing-TNF-induced apoptosis (via co-treatment with TNF (T) and a Smac mimetic

(S); Fig. 5). Conversely, immunohistochemistry detected intracellular clusters of Caspase-

8, RIPK1, RIPK3 or MLKL in cells undergoing TNF-induced necroptosis (via co-treatment

with T and S and the caspase inhibitor IDN-6556 (I); arrowhead Fig. 5). These intracellular

clusters are presumed to be necrosomes because they resemble prior images of

necrosomes (Chen *et al.*, 2022b; Samson *et al.*, 2021a; Samson *et al.*, 2020; Sun *et al.*,

2012) and because orthogonal approaches show that Caspase-8, RIPK1, RIPK3 and

MLKL are recruited to necrosomes during TNF-induced necroptosis (de Almagro *et al*,

2017; He *et al.*, 2009; Li *et al*, 2021; Sun *et al.*, 2012). Notably, the translocation of

Caspase-8 and RIPK1, but not MLKL, to necrosomes could also be detected in mouse

dermal fibroblasts undergoing TNF-induced necroptosis (Fig. EV6). This species-

dependent difference is likely due to dissimilarities in the interaction between RIPK3 and

MLKL, which is thought to be more transient in mouse than in human cells (Petrie *et al.*,

2018). Next, by combining automated immunohistochemistry with high-resolution digital

slide scanning (~250nm resolution) and customised image segmentation, we show that

necrosomes can be detected and quantified across a large population of cells in an

unbiased manner (Fig. 5B-C). We observed that the accuracy of segmenting human

Caspase-8 or MLKL at necrosomes is higher than that of RIPK1, because the small puncta

formed by necrosomal RIPK1 are near the resolution limit of existing brightfield slide

scanners (Fig. 5C). Nonetheless, since necrosomes are a pathognomonic feature of

necroptotic signaling, we propose that machine-based detection of necrosomes could be

developed into a diagnostic assay for pinpointing necroptosis in formalin-fixed human

patient biopsies. This proposal assumes that immunostaining protocols developed on cell

pellets retain their specificity when applied to tissues. To test this assumption, we used two

antibodies - one specific for mouse RIPK3 and one specific for human RIPK3 - on spleens

from $Ripk3^{-/-}$ mice, $Ripk3^{+/+}$ mice or knock-in mice expressing human RIPK3 (see

Methods). As shown in Fig. EV7, immunoblotting and immunohistochemistry with the

respective antibodies accurately discriminated between the expression of mouse RIPK3 or

human RIPK3 in spleen. These data suggest that immunohistochemistry protocols

optimised on cell pellets can also be used to specifically stain tissues.

Necrosome immunodetection in patients with IBD

Ulcerative colitis (UC) and Crohn's disease (CD) are the main types of IBD (Kobayashi *et*

al, 2020; Roda *et al*, 2020). The causes of adult onset IBD are multifactorial

(Ananthakrishnan, 2015; Graham & Xavier, 2020). While many studies show that excess

necroptosis promotes IBD-like pathology in mice (Gunther *et al.*, 2011; Matsuzawa-

Ishimoto *et al*, 2017; Schwarzer *et al.*, 2020; Vlantis *et al*, 2016; Wang *et al*, 2020a; Xie *et*

al., 2020; Xu *et al*, 2023), few studies have examined the prevalence of necroptosis in IBD

patients (Negroni *et al*, 2017; Pierdomenico *et al*, 2014; Shi *et al*, 2020). One Phase II trial

of a RIPK1 inhibitor in UC has failed to demonstrate clinical efficacy (Weisel *et al*, 2021),

but several other clinical and preclinical trials of RIPK1 inhibitors in IBD are underway.

Thus, the role of necroptosis in IBD requires further investigation. We collected intestinal

biopsies from adults with UC, CD and non-IBD patients (Fig. 6A together with clinical

information in Table 1). To capture the chronology of disease, biopsies were collected from

endoscopically 'non-inflamed', 'marginally inflamed' and 'inflamed' intestinal tissue from

patients with IBD. The grading of inflammation was verified by blinded histopathology

scores (Fig. 6B). Cell death signaling in biopsies from each location and endoscopic grade

was assessed by immunoblot (Fig. 6C and Table 1). To assist interpretation, patient

samples (blue annotations in Fig. 6C) were immunoblotted alongside lysates from HT29

cells undergoing apoptosis or necroptosis (red annotations in Fig. 6C). Apoptotic signaling

was inferred from increases in the conversion of pro-Caspase-3, -8 and -10 into their active cleaved forms (open arrowheads in Fig. 6C). Necroptotic signaling was inferred from increases in the abundance of phosphorylated RIPK3 and MLKL, relative to their non-phosphorylated forms (asterisks in Fig. 6C). This approach showed that cell death signaling is elevated in intestinal tissue from patients with IBD relative to patients without IBD, especially in inflamed intestinal biopsies from IBD patients (Fig. 6C and EV8). However, marked heterogeneity in the prevailing form of cell death was apparent in both UC and CD patients; with apoptosis dominant in some IBD cases (patients B and F in Fig. 6C and patient I in Fig. EV8), and necroptosis dominant in others (patients D and H in Fig. 6C). Given the ongoing development of RIPK1 inhibitors, it is noteworthy that phosphorylated MLKL coincided with phosphorylated RIPK1 in some, but not all patients with IBD (patients D versus H in Fig. 6C). Why cell death mechanisms vary between patients is currently unknown. Collectively, we find that cell death signaling increases in the inflamed gut, supporting the idea that cell death inhibitors are a potential treatment option for IBD. Whether apoptosis or necroptosis manifests in an individual IBD patient appears to be highly variable, highlighting the need for diagnostic approaches, such as automated immunohistochemistry, to identify patients who may benefit from anti-necroptotic therapy.

Having identified biopsies with differing modes of intestinal cell death, we next applied our panel of automated immunohistochemical stains to matched biopsies (collected from the same patients at the same time and from the same location; Table 1; note that a biopsy for immunohistochemistry was not collected from patient F). Consistent with immunoblotting data showing increased apoptosis in patients B and I (Fig. 6C and EV8), immunohistochemistry detected substantially more cleaved Caspase-3$^+$ epithelial cells in patients B and I (open arrowheads Fig. EV9 and EV10) compared to other patients in the cohort. No obvious changes to epithelial RIPK1, RIPK3 and MLKL were evident

between patients (Fig. **EV9** and **EV10**). By comparison, cytoplasmic clusters of Caspase-8

were evident in the epithelial layer of the inflamed biopsy from patient D and H (closed

arrowheads in **Fig. 6D,** closed arrowheads in **EV9** and **EV10**). Since these cytoplasmic

clusters of Caspase-8 were reminiscent of the Caspase-8$^+$ necrosomes in necroptotic

HT29 cells (**Fig. 5**), we used the same high-resolution digital slide scanning and unbiased

image segmentation approach as before. This quantitation showed that the number of

intraepithelial Caspase-8$^+$ clusters was low in non-IBD patient C and increased with

inflammation in IBD patient D (**Fig. 6E**). This trend mirrors the levels of necroptotic

signaling in IBD patient D (**Fig. 6B**), suggesting that cytoplasmic clusters of Caspase-8$^+$

may represent bona fide necrosomes. Why the immunohistochemical detection of

Caspase-8 distinguishes between necroptosis and apoptosis is unclear, but this likely

relates to the observation that Caspase-8-containing complexes that form during

necroptosis are qualitatively different to those that form during apoptosis (de Almagro *et*

al., 2017). This case study suggests that automated immunohistochemical detection of

Caspase-8$^+$ clusters in patient biopsies is feasible and could be developed into a

diagnostic assay for pinpointing necroptosis in clinical practice. To this end, future studies

with a larger number of biopsies are needed to determine whether the immunoblot

detection of necroptotic signaling significantly correlates with the immunohistochemical

detection of Caspase-8$^+$ clusters.

Discussion:

The difficulty of reliably detecting necroptotic signaling in fixed tissues has been a longstanding issue, generating confusion and conflicting results in the literature. To address this problem, we optimised the immunohistochemical detection of Caspase-8, RIPK1, RIPK3 and MLKL in formalin-fixed paraffin-embedded samples. While our prior studies immunostained non-crosslinked fixed monolayers (Samson *et al.*, 2021a; Samson *et al.*, 2020), here we used formalin-fixed paraffin-embedded specimens consistent with standard practice in clinical pathology and research departments around the world. In total, over 300 different immunostaining conditions were tested, yielding 13 automated immunohistochemistry protocols that we anticipate will be of broad utility to the cell death community and drive new insight into the causes, circumstances, and consequences of necroptosis. To assess the reliability of our automated protocols, we benchmarked our immunostaining results against data obtained using other methodologies. For instance, our automated immunohistochemistry protocols produced results that were comparable to public resources generated using proteomic, single cell transcriptomic, and spatial transcriptomic approaches (Ben-Moshe *et al.*, 2019; Geiger *et al*, 2013; Moor *et al.*, 2018; Tabula Muris *et al.*, 2018). Confidence was also taken from the similar staining patterns produced between three Caspase-8 antibodies (clones D53G2, 3B10 and 1G12), between two RIPK3 antibodies (clones 8G7 and 1H12), and between mouse and human intestinal tissue using the same RIPK1 antibody (clone D94C12). Thus, multiple lines of evidence suggest that the automated protocols described herein are specific and sensitive.

Our approach for optimising and interpreting immunohistochemical signals relied upon quantitative analyses that carries important technical considerations. First and foremost, the signals produced by immunohistochemistry are non-linear (Bobrow & Moen, 2001). Moreover, before quantitation, we digitally unmixed immunosignals from the haematoxylin counterstain, which is another non-linear transformation of signal intensity

(Landini *et al*, 2021). Thus, only relative changes in expression levels were inferred from

changes in immunohistochemical signal intensity. Because of this caveat, we only

compared and quantified immunosignals between closely matched specimens, such as

corresponding wild-type and knockout samples where both samples were sectioned at the

same time, mounted on the same slide, and stained and imaged contemporaneously. To

aid quantitation, all our immunohistochemistry protocols were developed to produce

unsaturated signals. One final salient point is that this study used automated embedding

and immunostaining procedures, with all quantification using macros that analyse a high

number (typically thousands) of cells per sample. Thus, the automated

immunohistochemistry protocols described herein can be used for quantitative purposes,

but only when comparing closely matched specimens, and ideally with supporting data

from alternative methodologies such as spatial transcriptomics. These recommendations

reduce, but do not eliminate, the need to use knockout samples as a control for specificity.

The phosphorylated forms of RIPK1, RIPK3 and MLKL are the most widely used

markers of necroptotic signaling (Horne *et al.*, 2023). Accordingly, prior attempts to detect

necroptotic signaling in fixed specimens have focussed on phospho-RIPK1, -RIPK3 and -

MLKL (Dominguez *et al.*, 2021; He *et al*, 2021; Li *et al.*, 2017; Li *et al*, 2022; Rodriguez *et

al*, 2022; Rodriguez *et al.*, 2016; Samson *et al.*, 2021a; Samson *et al.*, 2020; Wang *et al.*,

2014; Webster *et al.*, 2018; Zhang *et al.*, 2020). However, there are drawbacks with this

approach: 1) antibodies against phosphorylated epitopes in RIPK1/3 and MLKL exhibit

much poorer signal-to-noise properties than do antibodies against unphosphorylated

epitopes in RIPK1/3 and MLKL (Samson *et al.*, 2021a); 2) while knockout samples are

sufficient for verifying non-phospho-signals, multiple controls are needed to authenticate

phospho-signals (e.g. resting, knockout and phosphatase pre-treated samples); and 3)

because only a small fraction of RIPK1/3 and MLKL is phosphorylated during necroptosis

these phospho-species are inherently more difficult to detect than their unphosphorylated

counterparts. Indeed, as is standard practice when detecting necroptosis via immunoblot,

the immunohistochemical detection of phospho-RIPK1, -RIPK3 and -MLKL can only be

interpretated when their unphosphorylated forms can also be detected. For these reasons,

we focused on the immunohistochemical detection of unphosphorylated RIPK1, RIPK3

and MLKL, and used the immunodetection of necrosomes as a marker of necroptotic

signaling.

It was surprising that expression of the necroptotic pathway was heavily restricted

under steady-state conditions in mice. That fast-cycling progenitors and immune barrier

cells are the dominant expressors of RIPK3 and/or MLKL suggests that the existential role

of necroptosis is to protect the host from invading pathogens. The phenotypes and cell

types affected in mice carrying activation-prone polymorphisms in *Mlkl* also supports the

view that the ancestral role of necroptosis lies in innate immunity (Garnish *et al*, 2023;

Hildebrand *et al*, 2020; Zhu *et al*, 2022). The absence of necroptotic pathway expression in

slow-cycling cell populations was equally striking, with RIPK3 and/or MLKL undetectable in

resident cells of the heart (except for certain fibroblasts), the brain (except for

leptomeningeal vessels), the kidney and the liver (except for Kupffer cells) in mice under

basal conditions. These observations challenge a vast body of literature, highlighting the

need for robust well-controlled methodologies. These results lead us to propose that slow-

cycling cell populations ensure longevity by avoiding inadvertent necroptotic signaling. The

observation that RIPK3 expression is rapidly derepressed in hepatocytes during

inflammation supports the notion that long-lived cells actively suppress necroptotic

signaling, but only in the absence of challenge. The derepression of RIPK3 may reconcile

the contribution of necroptosis in inflammatory disorders such as hepatocellular carcinoma

(Vucur *et al*, 2023), acute myocardial infarction (Luedde *et al*., 2014), acute ischemic

stroke (Degterev *et al*, 2005) and kidney ischemia-reperfusion injury (Linkermann *et al*.,

2013). Notably, the Human Protein Atlas lacks immunohistochemical data for RIPK3 and

MLKL, and therefore it remains unknown whether the necroptotic pathway is similarly

restricted in humans (Uhlen *et al*, 2015).

It is noteworthy that certain non-mitotic cells, such as bone marrow-derived

macrophages, are highly sensitive to necroptotic stimuli (Zelic & Kelliher, 2018). Thus,

necroptosis is not reliant upon mitosis. Nonetheless, given the striking correlation between

RIPK3 and mitotic marker expression, and since RIPK3's activity and interactome vary

considerably during the cell cycle (Gupta & Liu, 2021; Liccardi *et al.*, 2019), future studies

should investigate how cell proliferation influences necroptotic susceptibility.

To exemplify the utility of our approach, we applied our full immunohistochemistry

panel to determine whether inflammation, dysbiosis or immunisation alter necroptotic

pathway expression in mice. Each of these challenges altered necroptotic pathway

expression in a manner that chiefly involved local shifts in RIPK3 expression. This profiling

of the necroptotic pathway yielded many unexpected observations, including: 1) RIPK3

expression is disinhibited in hepatocytes after TNF administration, with contemporaneous

increases in circulating RIPK3 levels; 2) RIPK3 expression is suppressed in the gut during

antibiotic administration with RIPK1/3 coalescing into unidentified cytoplasmic clusters in

epithelial cells at the villus tip; and 3) RIPK3 expression is uniquely upregulated in splenic

germinal centres after immunisation. While the mechanistic basis for these findings

warrants future attention, their initial description here illustrates the benefits of studying the

necroptotic pathway using automated immunohistochemistry, and raises questions about

whether RIPK3 expression is also regulated similarly in humans. This line of enquiry

circles back to the idea that variations in RIPK3 expression are a key determinant of

necroptotic potential (Cook *et al.*, 2014; He *et al.*, 2009; Najafov *et al*, 2018).

Our data raise the possibility that the immunodetection of intracellular Caspase-8[+]

clusters necrosomes could be used as an *in situ* marker of necroptotic signaling in IBD

patients.   To   reach   this   conclusion,   we   developed   a   suite   of   automated

immunohistochemical protocols to detect the relocation of necroptotic effectors into

necrosomes and validated the presence of necroptotic signaling in closely matched

biopsies from patients with IBD using immunoblot. Collectively, these experiments show

that intracellular clusters of Caspase-8, RIPK1, RIPK3, and MLKL are readily detectable

under idealised cell culture conditions, that elevated necroptotic signaling occurs in a

subset of IBD, and that cytoplasmic clustering of Caspase-8 correlated with necroptotic

signaling across a set of biopsies from two patients with IBD. Why clusters of Caspase-8,

but not clusters of RIPK1 or MLKL, were detectable in biopsies with active necroptotic

signaling is unknown but may be due to technical limitations (e.g. resolution limit) or gaps

in our understanding of how necroptosis manifests *in vivo*. Another important issue is

whether the immunohistochemical detection of Caspase-8^{+} clusters can be used to

quantify necroptotic signaling in larger cohorts of patients with IBD and in patients with

other clinical indications. Notwithstanding these issues, the detection of *in situ* changes in

necroptotic pathway expression in a scalable, quantitative, and automated manner

represents a major leap forward in the capacity to pinpoint when and where necroptosis

arises in both health and disease.

Materials and Methods:

Materials

Primary antibodies were rat anti-mouse Caspase-8 (clone 3B10; RRID:AB_2490519; 1g/L produced in-house (O'Reilly *et al*, 2004) and available from AdipoGen Cat#AG-20T-0138), rat anti-mouse Caspase-8 (clone 1G12; RRID:AB_2490518; 1g/L produced in-house (O'Reilly *et al.*, 2004) and available from AdipoGen Cat#AG-20T-0137), rabbit anti-Caspase-8 (clone D35G2; RRID:AB_10545768; Cell Signaling Technology Cat#4790), rabbit anti-phospho-RIPK1 (clone D813A; RRID:AB_2799268; Cell Signaling Technology Cat#44590S), mouse anti-RIPK1 (clone 38/RIP; RRID:AB_397831; 0.25g/L BD Biosciences Cat#610459), mouse anti-RIPK1 (clone 334640; RRID:AB_2253447; 0.5g/L; R&D Systems Cat#MAB3585), rabbit anti-RIPK1 (clone D94C12; RRID:AB_2305314; Cell Signaling Technology Cat#3493), rabbit anti-RIPK3 (clone 18H1L23; RRID: AB_2866471; 0.5g/L; Thermo Fisher Scientific Cat#703750), rabbit anti-phospho-RIPK3 (clone D6W2T; RRID:AB_2800206; Cell Signaling Technology Cat#93654), rat anti-RIPK3 (clone 1H12; 2g/L produced in-house (Samson *et al.*, 2021a)), rat anti-RIPK3 (clone 8G7; RRID: RRID:AB_2940810; 2g/L produced in-house (Petrie *et al.*, 2019) and available from Millipore Cat#MABC1595), rabbit anti-phospho-MLKL (clone D6E3G; RRID:AB_2799112; Cell Signaling Technology Cat# 37333), rabbit anti-phospho-MLKL (clone EPR9514; RRID:AB_2619685; Abcam Cat#ab187091; (Wang *et al.*, 2014)), mouse anti-MLKL (clone 3D4C6; RRID:AB_2882029; 1.957g/L; Proteintech Cat#66675-1-IG), rabbit anti-mouse MLKL (clone D6W1K; RRID:AB_2799118; Cell Signaling Technology Cat#37705), rat anti-mouse MLKL (clone 5A6; RRID:AB_2940800; 50g/L produced in-house (Samson *et al.*, 2021a) and available from Millipore Cat#MABC1634), rat anti-MLKL (clone 3H1; RRID:AB_2820284; 2g/L produced in-house (Murphy *et al.*, 2013) and available from Millipore Cat# MABC604), mouse anti-Caspase-10 (clone 4C1; RRID:AB_590721; 1g/L; MBL International Cat# M059-3), mouse anti-Caspase-8 (clone B.925.8;

RRID:AB_10978471; 0.619g/L Thermo Fisher Scientific Cat# MA5-15226), mouse anti-

Caspase-8 (clone 5D3; RRID:AB_590761; 1g/L; MBL International Cat#M058-3), rat anti-

human RIPK3 (clone 1H2; RRID:AB_2940816; 2g/L produced in-house (Petrie *et al.*,

2019) and available from Millipore Cat# MABC1640), rabbit anti-human RIPK3 (clone

E1Z1D; RRID:AB_2687467; Cell Signaling Technology Cat# 13526), rabbit anti-human

RIPK3 (clone E7A7F; RRID:AB_2904619; Cell Signaling Technology Cat# 10188), rat anti-

human MLKL (clone 7G2; RRID:AB_2940818; 2g/L produced in-house (Samson *et al.*,

2020) and available from Millipore Cat# MABC1636), rat anti-human MLKL (clone 10C2;

RRID:AB_2940821; 2g/L produced in-house (Samson *et al.*, 2020) and available from

Millipore Cat# MABC1635), rabbit anti-MLKL (clone 2B9; RRID:AB_2717284; 1g/L;

Thermo Fisher Scientific Cat#MA5-24846), rabbit anti-MLKL (clone EPR17514;

RRID:AB_2755030; 1.9g/L; Abcam Cat# ab184718), mouse anti-GAPDH (clone 6C5;

RRID:AB_2107445; 1g/L; Millipore Cat# MAB374), rabbit anti-smooth muscle actin (clone

D4K9N; RRID:AB_2734735; Cell Signaling Technology Cat#9245S), rabbit anti-lysozyme

(clone EPR2994(2); RRID:AB_10861277; Abcam Cat#108508), cleaved-caspase 3 (Cell

Signaling Technology, #9661), Ki67 (Cell Signaling Technology, #12202). The

concentration of antibodies from Cell Signaling Technology is often not provided and thus

was not listed here.

Secondary antibodies for immunoblotting were horseradish peroxidase (HRP)-conjugated

goat anti-rat immunoglobulin (Ig) (Southern BioTech Cat#3010-05), HRP-conjugated goat

anti-rabbit Ig (Southern BioTech Cat#4010-05) and HRP-conjugated goat anti-mouse Ig

(Southern BioTech Cat#1010-05).

Reagents for immunohistochemistry were HRP-conjugated anti-rabbit Ig (Agilent Cat#

K400311-2), HRP-conjugated anti-mouse Ig (Agilent Cat# K400111-2), MACH4 universal

HRP-Polymer (Biocare Medical Cat#M4U534L), ImmPRESS HRP-conjugated anti-rat IgG

for human samples (Vector Laboratories CatcVEMP740450), HRP-conjugated anti rat IgG

for mouse samples (R&D Systems Cat#VC005-125), Rabbit Linker (Agilent Cat#GV80911-

2). Epitope Retrieval Solution 1 (Leica Cat#AR9961), Epitope Retrieval Solution 2 (Leica

Cat#AR9640), Retrieval Solution Low pH (Agilent Cat#GV80511-2), Retrieval Solution

High pH (Agilent Cat#GV80411-2), 3,3'-diaminobenzidine (DAB) substrate (Agilent

Cat#GV82511-2 or GV92511-2), Dako REAL Peroxidase-blocking reagent (Agilent

S202386-2), bluing reagent (Leica, 3802915), Background Sniper (Biocare Medical

Cat#BS966L), Dako Protein Block (Agilent Cat#X0909), 'Normal' block (Agilent

Cat#S202386-2), EnVision FLEX TRS High pH (Agilent Cat# GV80411-2), EnVision FLEX

TRS Low pH (Agilent Cat# GV80511-2), MACH4 universal HRP polymer (Biocare Medical

Cat#M4U534L), and DPX (Trajan Cat#EUKITT), Opal TSA-Dig (Akoya Cat#OP-001007),

Opal Polaris 780 reagent (Akoya Cat#OP-001008), and HRP-conjugated anti-digoxigenin

antibody (Cell Signaling Technology, clone D8Q9J, Cat#49620).

Mice, research ethics and housing

All experiments were approved by The Walter and Eliza Hall Institute (WEHI) Animal

Ethics Committee, Australia, in accordance with the Prevention of Cruelty to Animals Act

(1986), with the Australian National Health and Medical Research Council Code of

Practice for the Care and Use of Animals for Scientific Purposes (1997), and with the

ARRIVE guidelines (Percie du Sert *et al*, 2020). Mice were housed at the WEHI animal

facility under specific pathogen-free, temperature- and humidity-controlled conditions and

subjected to a 12 h light/dark cycle with ad libitum feeding. Mice without functional MLKL

alleles (*Mlkl*$^{-/-}$) have been described previously (Murphy *et al.*, 2013). Mice without

functional RIPK3 alleles (*Ripk3*$^{-/-}$) have been described previously (Tovey Crutchfield *et al*,

2023). Mice without functional alleles of RIPK1, RIPK3 and Caspase-8 (*Ripk1*$^{-/-}$

Ripk3$^{-/-}$Casp8$^{-/-}$ triple knockout mice) and *Ripk3$^{-/-}$Casp8$^{-/-}$* double knockout mice have

been described previously (Rickard *et al*, 2014) and were derived from reported mouse

strains (Kelliher *et al*, 1998; Newton *et al.*, 2004; Salmena *et al*, 2003). Human RIPK3

knock-in mice in which the human RIPK3 coding sequence was inserted into the mouse

Ripk3 locus of C57BL/6J mice were provided by Anaxis Pharma Pty Ltd.

Mouse tissue lysate preparation

Mouse tissues were homogenised with a stainless steel ball bearing in a Qiagen

TissueLyzer II (30 Hz, 1min) in ice-cold RIPA buffer (10mM Tris-HCl pH 8.0, 1mM EGTA,

2mM MgCl2, 0.5% v/v Triton X100, 0.1% w/v sodium deoxycholate, 0.5% w/v sodium

dodecyl sulfate (SDS) and 90mM NaCl) supplemented with 1x Protease & Phosphatase

Inhibitor Cocktail (Cell Signaling Technology Cat#5872) and 100 U/mL Benzonase (Sigma-

Aldrich Cat#E1014). 1mL of RIPA buffer per 25mg of tissue was used for homogenisation.

Immunoblot

For Fig. EV1, EV5 and EV7, RIPA lysates were boiled for 10min in Laemmli sample buffer

(126 mM Tris-HCl, pH 8, 20% v/v glycerol, 4% w/v sodium dodecyl sulfate, 0.02% w/v

bromophenol blue, 5% v/v 2-mercaptoethanol) and fractionated by 4-15% Tris-Glycine gel

(Bio-Rad Cat#5678084) using Tris-Glycine running buffer (0.2M Tris-HCl, 8% w/v SDS,

0.15M glycine). After transfer onto nitrocellulose, membranes were blocked in 1% w/v

bovine serum albumin (BSA; for Caspase-8 antibody clone B.925.8) or 5% w/v skim cow's

milk (for all other antibodies) in TBS+T (50mM Tris-HCl pH7.4, 0.15M NaCl, 0.1 v/v

Tween20), probed with primary antibodies (1:2000 dilution for rat primary antibodies or

1:1000 for other primary antibodies in the above blocking buffers; supplemented with

0.01% w/v sodium azide; see *Materials* above) overnight at 4^{o}C, washed twice in TBS+T,

probed with an appropriate HRP-conjugated secondary antibody (see *Materials* above),

washed four times in TBS+T and signals revealed by enhanced chemiluminescence

(Merck Cat#WBLUF0100) on a ChemiDoc Touch Imaging System (Bio-Rad). Between

probing with primary antibodies from the same species, membranes were incubated in

stripping buffer (200 mM glycine pH 2.9, 1% w/v SDS, 0.5 mM TCEP) for 30 min at

room temperature then re-blocked.

For Fig. **4**, **6**, **EV3**, and **EV8**, RIPA lysates were boiled for 10min in Laemmli sample buffer

(126 mM Tris-HCl, pH 8, 20% v/v glycerol, 4% w/v SDS, 0.02% w/v bromophenol blue,

5% v/v 2-mercaptoethanol) and fractionated by 4–12% Bis-Tris gel (Thermo Fisher

Scientific Cat#NP0335BOX) using MES running buffer (Thermo Fisher Scientific

Cat#NP000202). After transfer onto polyvinylidene difluoride (Merck Cat# IPVH00010),

gels were Coomassie-stained as per manufacturer's instructions (Thermo Fisher Scientific

Cat#LC6060) and membranes were blocked in 5% w/v skim cow's milk in TBS+T and then

probed as above.

Mouse tissue fixation

Tissues were immediately harvested after euthanasia and placed in 10% v/v Neutral

Buffered Formalin (NBF) (Confix Green; Australian Biostain Cat#AGFG.5L). A ratio of one

part tissue to >10 parts formalin was used. Unless stipulated, tissues were incubated in

formalin at room temperature for 24-72 hours.

Paraffin embedding, microtomy and immunostaining

Unless stipulated, formalin-fixed cells/tissues were paraffin embedded using the standard

8 hour auto-processing protocol of the Tissue-Tek VIP® 6 AI Tissue Processor (Sakura

Finetek USA). Paraffin embedded cells/tissues were cut in 4μm-thick sections onto

adhesive slides (Menzel Gläser Superfrost PLUS). Haematoxylin and eosin staining was

done on the Autostainer XL (Leica ST5010). Immunohistochemistry was performed on an

automated system: the Bond RX (Leica) or the DAKO OMNIS (Agilent). While the

automated immunohistochemistry protocols used in this manuscript are fully stipulated in

File EV1, a brief overview of these methods is as follows: Step 1 was deparaffinisation

(Phase 1: Clearify Clearing Agent; Phase 2: deonised water), Step 2 was heat-induced

antigen retrieval (EnVision FLEX TRS, High pH or Low pH retrieval buffer. Treatment time

was antibody-dependent), Step 3 was endogenous peroxidase blocking   (Dako   REAL

Peroxidase-blocking reagent. Incubation time was antibody-dependent), Step 4 was

protein blocking (Background Sniper, Dako Protein Block or Normal block. Incubation 10

minutes), Step 5 was primary antibody incubation (dilutions and incubation times were

antibody-dependent), Step 6 was option signal amplification (amplification technique was

antibody-dependent. Rabbit linker. Incubation 15 minutes), Step 7 was secondary reagent

with horseradish peroxidase (anti-rabbit HRP polymer, anti-mouse HRP polymer, MACH4

universal HRP polymer, ImmPRESS anti-rat HRP polymer, anti-rat HRP polymer and in-

house HRP signal amplification detection. Reagent and incubation was antibody-

dependent), Step 8 was signal detection (chromogen-substrate DAB. Onboard mixing and

incubation 10 minutes), Step 9 was counterstaining   (in-house   made   Mayer's

Haematoxylin for 1 minute followed by Bluing reagent for 1 minute), Step 10 was

dehydration, mounting in DPX and coverslipping using Leica CV5030 platform.

Image acquisition

Unless stipulated, slides were scanned on: 1) 3D Histech Pannoramic Scan II (objective:

magnification 20x, numerical aperture 0.8, media dry; software: Pannoramic SCAN 150

1.23 SP1 RTM and SlideViewer 2.8.178749), or 2) Olympus VS200 (objective: 20x,

numerical aperture 0.8, media dry; software: Olympus VS200 ASW 3.41). Where higher

resolution was required, slides were scanned on the Olympus VS200 using the 60x

objective (numerical aperture 1.42, media oil).

Post-acquisition processing of displayed micrographs

All displayed micrographs were acquired with VS200 Research Slide Scanner (Olympus).

Representative full-resolution 8-bit RGB micrographs of the WT and KO tissues/cells were

imported into ImageJ 1.53t (Schindelin *et al*, 2012). Brightness-and-contrast was adjusted

to 0-235 units and then gamma levels adjusted by 1-2.5-fold. Capture settings and post-

acquisition image transformations were held constant between any micrographs that were

being compared.

Immunohistochemistry signal-to-noise ratio (S/N) analysis

Sections of wild-type (WT) and corresponding knockout (KO) tissues/cells on the same

slide were immunostained and imaged. Representative full-resolution 8-bit RGB

micrographs of the WT and KO tissues/cells were imported into ImageJ 1.53t (Schindelin

et al., 2012) and the 'H-DAB' function of the 'Colour Deconvolution 2' plugin (Landini *et al.*,

2021) was used to unmix the DAB and hematoxylin channels. A dilated mask of the auto-

thresholded hematoxylin channel was applied to the corresponding DAB channel to select

an area-of-interest, and then a histogram of pixel intensities from the DAB channel for both

the WT and KO micrographs was determined. The 'WT histogram' was then divided by the

'KO histogram' to yield a S/N curve (see Fig.1A for example). A weighted integral of the

S/N curve was calculated as a numerical index of specificity (red box in Fig. 1A). The

representative micrographs used to calculate S/N curves were of equivalent size, included

>1000 cells per sample and their gamma levels unchanged.

Quantitation of zonation via immunohistochemistry

Representative full-resolution 8-bit RGB micrographs of the WT tissues were imported into

ImageJ 1.53t (Schindelin *et al.*, 2012). Brightness-and-contrast was adjusted to 0-235

units and the 'H-DAB' function of the 'Colour Deconvolution 2' plugin (Landini *et al.*, 2021)

was used to unmix the DAB and hematoxylin channels. The look-up-table for the DAB

channel was converted to grayscale and pixel values inverted. The 'Segmented Line tool'

was used to draw a line (27.4μm wide to approximate 2 cell widths) along the main axis of

the following zones: 1) crypt base to villi tip in the ileum, 2) crypt base to crypt tip in the

colon, 3) peri-central hepatocytes to peri-portal hepatocytes, and 4) white matter to red

pulp in the spleen. The 'Plot Profile tool' was used to quantify the immunosignal along the

drawn axis, which was then averaged across N=10-20 representative zones per mouse.

Quantitation of endothelial RIPK3 levels via immunohistochemistry

Sections were subjected to automated immunohistochemistry for smooth muscle actin

(SMA) and RIPK3 (see File **EV1**). SMA signals were detected with a brown DAB product.

RIPK3 signals were detected with a pink DAB product. Sections were not counterstained

with hematoxylin. Full-resolution 8-bit RGB micrographs (taken with 60x objective;

Olympus VS200) were imported into Image J 1.53t (Schindelin *et al.*, 2012) and

endothelial RIPK3 levels analysed using a custom semi-automated macro in a non-blinded

manner. In brief, cross-sections of individual SMA[+] vessels were chosen, SMA and RIPK3

signals were unmixed using the 'Colour Deconvolution 2' plugin (Landini *et al.*, 2021). An

auto-thresholded mask of the SMA[+] intima was then used to segment the endothelial area.

The total RIPK3 signal in this endothelial area was expressed per unit area. This

procedure was repeated for N=50 vessels per mouse.

Quantitation of hepatocyte RIPK3 levels via immunohistochemistry

Sections were subjected to automated immunohistochemistry for RIPK3 (see File **EV1**).

RIPK3 signals were detected with brown DAB product and sections were counterstained

with hematoxylin. Representative full-resolution 8-bit RGB micrographs (taken with the 60x

objective; Olympus VS200) were imported into Image J 1.53t (Schindelin *et al.*, 2012) and

RIPK3 levels within individual hepatocytes analysed using a custom fully-automated

macro. In brief, haematoxylin-stained nuclei were unmixed from the RIPK3 signal using the

'Colour Deconvolution 2' plugin (Landini *et al.*, 2021). Nuclei were segmented using the

'Analyze Particles' function of Image J and hepatocyte nuclei distinguished from Kupffer

cell nuclei on the basis of their larger size and circularity. The total RIPK3 signal in the

cytosolic area surrounding hepatocyte nuclei was segmented, measured and expressed

per unit area. N=90 hepatocytes per mouse were measured.

Quantitation of Kupffer cell RIPK3 levels via immunohistochemistry

Sections were subjected to automated immunohistochemistry for RIPK3 (see File **EV1**).

RIPK3 signals were detected with brown DAB product and sections counterstained with

hematoxylin. Representative full-resolution 8-bit RGB micrographs (taken with the 60x

objective; Olympus VS200) were imported into Image J 1.53t (Schindelin *et al.*, 2012) and

RIPK3 levels within individual Kupffer cells were analysed using a custom fully-automated

macro. In brief, the RIPK3 signal was unmixed from Hematoxylin using the 'Colour

Deconvolution 2' plugin (Landini *et al.*, 2021). An auto-thresholded mask of cells with

relatively high expression of RIPK3 was created. This mask, together with the

comparatively small and non-circular shape of Kupffer cells, was used to segment

individual Kupffer cells. The total RIPK3 signal within each Kupffer cell was then measured

and expressed per unit area. N=90 Kupffer cells per mouse were measured.

Quantitation of apoptosis in splenic white pulp via immunohistochemistry

Sections were subjected to automated immunohistochemistry for cleaved Caspase-3 (see

File **EV1**). Cleaved Caspase-3 signals were detected with a brown DAB product and

sections counterstained with hematoxylin. Representative full-resolution 8-bit RGB

micrographs (taken with the 60x objective; Olympus VS200) were imported into Image J

1.53t (Schindelin *et al.*, 2012) and percent area with cleaved Caspase-3 signal was

analysed using a custom fully-automated macro. In brief, the cleaved Caspase-3 signal

was unmixed from Hematoxylin using the 'Colour Deconvolution 2' plugin (Landini *et al.*,

2021). A predefined threshold (50-255 units) was applied, and its area expressed as a

percentage of the total white pulp area. N=20 white pulp lobules per mouse were

measured.

Systemic Inflammatory Response Syndrome (SIRS)

For experiments in Fig. **3A-J**, co-housed 8-week-old female C57BL/6J wildtype mice were

administered either 300 μg/kg TNF (R&D Systems Cat#410-MT/CF Lot CS152103) in

endotoxin-free DPBS or endotoxin-free DPBS alone (Merck Cat# TMS-012-A) via bolus tail

vein injection. Core body temperature was measured hourly with a rectal probe. Mice were

euthanised via carbon dioxide inhalation 9 hours after injection.

For experiments in Fig. **3K-L**, co-housed 6-month-old female C57BL/6J wildtype mice

were administered either: 1) 6mg/kg of Necrostatin1s (Cell Signaling Technologies

Cat#17802) in dimethyl sulfoxide via intraperitoneal injection then 15-30 minutes later

given 150 μg/kg TNF (BioTimes Inc. Cat#CF09-1MG), or 150 μg/kg TNF (R&D Systems

Cat#410-MT/CF Lot CS152103) in DPBS, or DPBS alone via bolus tail vein injection. Core

body temperature was measured hourly with a rectal probe. Mice were euthanised via

carbon dioxide inhalation 8 hours after injection or if their temperature dropped below

30ºC.

Enzyme-linked ImmunoSorbent Assay (ELISA) for mouse RIPK3

96-well ELISA plates (Sigma-Aldrich, Cat#CLS3795) were coated with rat anti-RIPK3

(clone IH12; 5μg/ml diluted in PBS) and incubated overnight at room temperature in humid

conditions. Plates were then washed in PBS + 0.005% v/v Tween-20, then PBS, then

distilled water. Mouse serum samples were diluted and titrated on the plate in block

solution (PBS containing 1% v/v FCS, 0.002% v/v Tween-20 and 0.6% w/v skim milk

powder). Plates were incubated at room temperature in humid conditions overnight. Plates

were washed as before. Detection antibody, biotin-conjugated rat anti-RIPK3 (clone 8G7;

5μg/ml diluted in block solution) was added to each well. Plates were incubated for 4 hours

at room temperature in humid conditions. Plates were washed as before. Streptavidin-HRP

(SouthernBiotech, Cat#7100-05) was diluted in block solution and added to each well.

Plates were incubated for 1 hour at room temperature in humid conditions. Plates were

washed as before, and ABTS substrate solution (water containing 0.54mg/mL w/v 2'2-

Azinobis (3-ethylbenzthiazoline Sulfonic Acid) diammonium salt (Sigma-Aldrich, Cat#A-

1888), and 0.1M Citric Acid and 0.03% v/v Hydrogen Peroxide (H_2O_2)) was added to each

well. Plates were left to develop for 30-45 minutes at room temperature protected from

light. Color development was analysed on VersaMax ELISA microplate reader (Molecular

Devices) using wavelengths λ=415nm minus λ=492nm.

Tabula muris **analysis**

Robject files of the FACS-based T*abula Muris* (Tabula Muris *et al.*, 2018) dataset were

downloaded from Figshare. Expression values were normalised and analysed using

Seurat v4.3.0 (Butler *et al*, 2018; Hao *et al*, 2021; Satija *et al*, 2015; Stuart *et al*, 2019).

Within each cell ontology and tissue origin, the percentage of cells with expression values

>0 for *Mlkl, Ripk3, Ripk1, Casp8, Mki67 and Top2a* was tabulated and colorised using

Excel v16.74 (Microsoft).

Spatial transcriptomics of mouse spleen

To boost germinal centre numbers and size, one adult C57BL/6J mouse was infected

intravenously with 1×10^5 *Plasmodium berghei* parasitised red blood cells, and then drug-

cured at the onset of disease symptoms as described in (Ly *et al*, 2019). Twelve days later

the mouse was euthanised, spleen dissected, fixed in 10% v/v Neutral Buffered Formalin

and paraffin embedded (as above). Sections of formalin-fixed paraffin-embedded spleen

were cut onto slides, then spatial enhanced resolution omics-sequencing (stereo-seq) data

was generated using pre-release chemistry and MGI sequencers as in (Chen *et al*, 2022a)

by BGI, China. The spot-to-spot distance is 500nm and the data is binned to 50x50 spots

($25\mu m^2$). Binning was performed in stereopy (https://github.com/BGIResearch/stereopy)

before being converted to an anndata file (https://github.com/scverse/anndata). The

stereo-seq data was first loaded and pre-processed using the standard Scanpy workflow

(https://github.com/scverse/scanpy) and then principal component analysis and Leiden

clustering was performed at the bin (50x50) level. Leiden clusters were then plotted using

Uniform Manifold Approximation and Projection (UMAP) and on spatial coordinates using

the Squidpy package (https://github.com/scverse/squidpy). Scanpy's 'rank_genes_groups'

method (https://scanpy-tutorials.readthedocs.io) was used to generate a matrix of gene by

cluster populated by corresponding log-fold changes from 1 versus all t-tests. For each

zone/cell type of interest a matrix is constructed by subsetting the full gene by cluster

matrix generated from 'rank_genes_groups' to just the specific genes of interest. These

matrices were reduced to a single column sum aggregated vector. From the vectors of

scores for each 'zone' of the spleen (White Pulp, Red Pulp, Germinal Centres, and

Marginal Zones), the top 5 scores within the 50^{th} or 75^{th} (depending on expected

transcriptional variance within zone) percentile of the maximum are selected and the

corresponding Leiden clusters are aggregated hierarchically. After this aggregation of

clusters, the 'rank_genes_groups' method is rerun to calculate a new set of gene rankings

and log-fold changes for zones rather than clusters. The aggregated scores for genes of

interest in each group are calculated for each zone. These scores are the log-fold changes

for each gene within each zone. Scores for the *Casp8*, *Ripk1*, *Ripk3*, and *Mlkl* genes are

then extracted from the matrices for each 'zone' of the spleen, and each individual cluster

for side-by-side comparison in heatmaps. Software version used were Anndata (v0.7.5),

Stereopy (v0.12.0), Scanpy (v1.9.2), Squidpy (v1.2.2), Numpy (v1.21.6), Pandas (v1.5.3),

Matplotlib (v3.5.2), and Seaborn (v0.12).

Antibiotics administration

Co-housed littermates were split across two cages. The drinking water for one cage was

supplemented with 1g/L ampicillin, 1g/L neomycin, 1g/L metronidazole, 0.5g/L

enrofloxacin, 2.5g/L meropenem as in (Bader *et al.*, 2023). Antibiotic-supplemented water

was replaced after 3 days. To minimise weight loss, both cages were given ad libitum

access to Di-Vetelact supplement (Lillelund Pty Ltd). Mice were euthanised after 6 days of

treatment.

RNA extraction, library preparation and sequencing

Immediately following dissection, a ~2mm^3 piece of tissue was placed in 0.5mL RNA*later*

(Thermo Fisher Scientific Cat#4427575) then stored at -80°C. Samples were thawed,

RNA*later* was removed then tissues were transferred into screw-capped tube pre-filled

with 350 µl of RA1 buffer of NucleoSPin RNAXS kit (Macherey-Nagel Cat#

SKU: 740902.250). Tissues were homogenised with 10 pcs of 3 mm Acid- Washed

Zirconium Beads (OPS diagnostics Cat# BAWZ 3000-300-23) in a Qiagen TissueLyzer II

(30 Hz, 5 minutes). Homogenised samples were centrifuged (1 minute, 11,000 *g*) to

remove tissue debris then RNA was purified using Nucleospin RNAXS column kit as per

manufacturer's instructions without adding a carrier RNA. The purified RNA was quantified

using Qubit™ RNA HS Assay kit (Thermo Fisher Scientific Cat#Q32852) and RNA integrity

was visualised in high sensitivity RNA ScreenTape (Agilent Cat# 5067- 5579) using

TapeStation 4200 (Agilent Cat# G2991BA). Ten nanograms of RNA were used for

preparing indexed libraries using SMARTer Stranded Total RNA-Seq Pico-Input

Mammalian kit v.3 (Takara Bio. Cat# SKU: 634487) as per manufacturer's instructions

(except that fragmentation was performed for 3 minutes of fragmentation at 94 °C and 13

cycles was used for PCR2). The library concentration was quantified by Qubit™ dsDNA

Assay kit (Thermo Fisher Cat#Q32851) and library size was determined using D1000

ScreenTape (Agilent Cat# 5067-5582) and visualised in TapeStation 4200. Equimolar

amounts of the libraries were pooled and diluted to 750pM for 150-bp paired-end

sequencing on a NextSeq2000 instrument (Illumina) using the P2 300- cycle kit v3

chemistry (Illumina Cat# 20046813) as per manufacturer's instructions. To produce the

sequences, the base calling and quality scoring was performed by the Real Time Analysis

(v2.4.6) software. The FASTQ file generation and de-multiplexing for the samples was

performed by the bcl2fastq conversion software (v2.15.0.4).

Murine bulk RNA sequence analysis

The single-end 75 bp were demultiplexed using CASAVAv1.8.2 and Cutadapt (v1.9) was

used for read trimming (Martin, 2011). The trimmed reads were subsequently mapped to

the mouse genome (mm10) using HISAT2 (Kim *et al*, 2019). FeatureCounts from the

Rsubread package (version 1.34.7) was used for read counting after which genes <2

counts per million reads (CPM) in at least 3 samples were excluded from downstream

analysis (Liao *et al*, 2014, 2019). Count data were normalised using the trimmed mean of

M-values (TMM) method and differential gene expression analysis was performed using

the limma-voom pipeline (limma version 3.40.6) (Law *et al*, 2014; Liao *et al.*, 2014;

Robinson & Oshlack, 2010). Adjustment for multiple testing was performed per comparison

using the false discovery rate (FDR) method (Benjamini & Hochberg, 1995). Heatmaps of

logCPM were generated using pheatmap.

NP-KLH immunization and analysis

Immunisation was performed as previously described (Kong *et al*, 2022). 8–10-week-old

Ripk3$^{+/+}$ or *Ripk3*$^{-/-}$ littermate mice received a single intraperitoneal injection of 100 µg 4-

hydroxy-3-nitrophenylacetyl hapten coupled to keyhole limpet hemocyanin (NP-KLH;

produced in-house) at a ratio of 21:1 with alum. Fourteen days after immunisation mice

were euthanised via CO_2 inhalation and spleens harvested. To determine immune

response to NP immunization, single-cell splenic suspensions were stained as described

using antibodies to the following surface molecules: CD38 (clone:NIMR-5, in-house),

CD19 (clone:1D3, cat #BD552854), IgM (clone:331.12, in-house), IgD (clone:11–26 C, in-

house), Gr-1 (clone:RB6-8C5, in-house), CD138 (clone:281.2, cat #BD564068) and IgG1

(clone:X56, cat #BD550874). NP-binding was detected as described (Smith *et al*, 2000)

Mouse serum analyses

Blood was collected via cardiac puncture and immediately transferred to an EDTA-coated

tube (Sarstedt AG Cat#20.1341). Blood was transferred into a clot activator tube (Sarstedt

AG Cat#20.1344) and serum prepared as per the manufacturer's instructions.

Human research ethics

Ethical approval for intestinal tissue collection from participants undergoing endoscopy

procedures through the Gastroenterology Department at the Royal Melbourne Hospital

was attained from the Human Research Ethics Committee (HREC): HREC 2021.074. This

was in accordance with the National Health and Medical Research Council (NHMRC)

National Statement on Ethical Conduct in Human Research (2008) and the Note for

Guidance on Good Clinical Practice (CPMP/ICH-135/95). Site-specific governance was

sought for each of the collaborating sites: WEHI and the University of Melbourne.

Collaboration amongst all three participating institutions was officiated through the

Melbourne Academic Centre for Health Research Collaboration Agreement (Non-

Commercial). The human research in this study was performed in accordance with the

principles expressed in the Helsinki Declaration. The human materials used in this study

were obtained with informed consent from all subjects.

Human intestinal biopsy collection

Adult patients with or without IBD scheduled for endoscopic evaluation of the lower

gastrointestinal tract (flexible sigmoidoscopy or colonoscopy) at the Gastroenterology

Department at the Royal Melbourne Hospital were screened for study inclusion/exclusion.

Patients were excluded based on the following criteria: active systemic (gastrointestinal

and non-gastrointestinal) infection; active (solid-organ or haematological) malignancy or

treatment with anti-tumour therapies; non-steroidal anti-inflammatory drug use in the last

month; hereditary or familial polyposis syndromes; non-IBD forms of colitis including

microscopic colitis, ischaemic colitis, diversion colitis, or diverticulitis. Eligible patients (see

Table 1) were recruited and consented by the gastroenterologist (signed authorised

Participant Information Sheet/Consent Form). For patients with IBD, intestinal biopsies

were retrieved endoscopically from: 1) inflamed, 2) non-inflamed, or 3) marginal areas of

inflammation. The same exclusion criteria as above were applied to patients without IBD

(referred to as non-IBD control patients), with biopsies retrieved endoscopically from non-

inflamed segments of the intestine. Boston Scientific Radial Jaw^{TM} biopsy forceps and

Olympus EVIS EXERA III endoscopes were routinely utilised for intestinal biopsy

collections. Collected biopsies were immediately placed in 10% v/v Neutral Buffered

Formalin. A ratio of one part tissue to >10 parts formalin was used. Tissues were incubated

in formalin at room temperature for 24-72 hours before paraffin-embedding.

Blinded histopathological scoring of intestinal inflammation

Hematoxylin and eosin-stained slides from formalin-fixed paraffin-embedded biopsies were

blindly scored by an anatomical pathologist using the Robarts Histopathology Index (RHI),

a validated tool for assessing IBD activity, using previously described methodology (Mosli

et al, 2017).

Cell lines

HT29 cells were originally sourced from the American Type Culture Collection. The

$RIPK1^{-/-}$, $RIPK3^{-/-}$, $MLKL^{-/-}$ and $CASP8^{-/-}CASP10^{-/-}MLKL^{-/-}$ HT29 cells have been

previously reported (Jacobsen *et al*, 2022; Petrie *et al*., 2018; Tanzer *et al*., 2017) . Mouse

dermal fibroblasts were generated in-house from the tails of wild-type C57BL/6J mice and

immortalised by SV40 large T antigen as reported previously (Hildebrand *et al*., 2014). The

sex and precise age of these animals were not recorded, although our MDFs are routinely

derived from tails from 8-week-old mice. Mouse dermal fibroblast lines were generated in

accordance with protocols approved by the Walter and Eliza Hall Institute of Medical

Research Animal Ethics Committee. The origin of cell lines was not further verified,

although their morphologies and responses to necroptotic stimuli were consistent with their

stated origins. Cell lines were monitored via polymerase chain reaction every ~6 months to

confirm they were mycoplasma-free.

Cell culturing

HT29 cells and mouse dermal fibroblasts were maintained in Dulbecco's Modified Eagle

Medium (DMEM; Life Technologies) with 8% v/v fetal calf serum (FCS), 2mM L-Glutamine,

50 U/mL penicillin and 50 U/mL streptomycin (G/P/S). Cells were incubated under

humidified 10% CO_2 at $37^{o}C$.

Cell treatment

HT29 cells were treated in DMEM containing 1% v/v FCS and G/P/S. Mouse dermal

fibroblasts were treated in DMEM containing 8% v/v FCS and G/P/S. Media for treatment

was supplemented with: 100 ng/mL recombinant human TNF-α-Fc (produced in-house as

in (Bossen *et al*, 2006)), 500 nM Smac mimetic/Compound A (provided by Tetralogic

Pharmaceuticals as in (Vince *et al*, 2007)) 5 µM IDN-6556 (provided by Idun

Pharmaceuticals). HT29 cells were treated for 7.5 hours. Mouse dermal fibroblasts were

treated for 2 hours.

Cell lysate preparation

961 HT29 cells were homogenised in ice-cold RIPA buffer (10mM Tris-HCl pH 8.0, 1 mM

962 EGTA, 2mM MgCl2, 0.5% v/v Triton X100, 0.1% w/v Na deoxycholate, 0.5% w/v SDS and

963 90mM NaCl) supplemented with 1x Protease & Phosphatase Inhibitor Cocktail (Cell

964 Signaling Technology Cat#5872S) and 100 U/mL Benzonase (Sigma-Aldrich Cat#E1014).

Cell pellet preparation

Trypsinised cells were centrifuged at 671*g* (2000rpm) for 3 minutes at room temperature.

The supernatant was discarded, cell pellets resuspended in 10% v/v Neutral Buffered

Formalin, incubated for 15 minutes at room temperature, and centrifuged at 671*g*

(2000rpm) for 3 minutes at room temperature. Cell pellets were resuspended in 50-70µL of

HistoGel (Epredia Cat#HG-4000-012) pre-warmed to 56°C and then pipetted onto ice-cold

glass coverslips to set. Set pellets were stored in 70% (v/v) ethanol until paraffin

embedding.

Human tissue lysate preparation

Two intestinal biopsies (each ~1-2mm^3) were pooled, immediately washed in ice-cold

DPBS (Thermo Fisher Scientific Cat#14190144) supplemented with protease inhibitors

(Merck Cat#4693132001) and phosphatase inhibitors (Merck Cat#4906837001), and then

homogenised with a stainless steel ball bearing in a Qiagen TissueLyzer II (30 Hz, 1min) in

0.4mL of ice-cold RIPA buffer (10mM Tris-HCl pH 8.0, 1mM EGTA, 2mM MgCl2, 0.5% v/v

Triton X100, 0.1% w/v sodium deoxycholate, 0.5% w/v SDS and 90mM NaCl)

supplemented with 1x Protease & Phosphatase Inhibitor Cocktail (Cell Signaling

Technology Cat#5872) and 100 U/mL Benzonase (Sigma-Aldrich Cat#E1014).

Necrosome quantitation

Necrosome detection was performed with a custom pipeline developed in Fiji (Schindelin

et al., 2012). Cells were segmented using CellPose (Stringer *et al*, 2021) and puncta

detected using a difference-of-gaussian algorithm developed for graphics processing units

(Haase *et al*, 2020). To avoid false-positives, puncta were filtered to be more than twice

the signal of the average signal of the cell on which it appears. Number of cells with

puncta, along with number of puncta per cell was recorded. Prior to detecting necrosomes

in patient biopsies, the epithelial regions were segmented using FastPathology (v1.1.2)

with a deep learning model trained on CD3-stained colon biopsy whole slide images

(Pedersen *et al*, 2021; Pettersen *et al*, 2021). Briefly, the whole slide images were

imported into FastPathology and the default model pipeline was executed using the

attribute "patch-level 4". The binary mask was imported into QuPath (v0.4.3) using the

997 "importPyramidalTIFF.groovy" script provided at https://github.com/andreped/NoCodeSeg

998 (Pedersen *et al.*, 2021).

1000 **Statistical tests**

1001 The number of independent experiments and the employed statistical test for each dataset

1002 is stipulated in the respective figure legend. Statistical tests were performed using Prism

1003 v.9.5.1 (GraphPad). Statistical analyses were only performed on datasets collated from at

1004 least three independent experiments. The number of independent replicates or mice is

1005 stipulated by 'n', and the number of dependent replicates for each dataset is stipulated by

1006 'N' in the respective Fig. legend.

1007

1008 **Data Availability Statement:**

1009 Customised image analysis macros used in this study are available upon request to the

1010 corresponding authors. The automated immunohistochemical staining protocols used in

1011 this study are available as File **EV1**. The RNA sequencing datasets generated during this

1012 study have been submitted to the Gene Expression Omnibus repository.

1013

1014

1041

1042   Table **1.** Patient and clinical data related to Fig. **6** and **EV8-10**.

1043

1044   File **EV1** – Autostainer immunohistochemistry protocols used in this manuscript.

1045

References:

1046

Alvarez-Diaz S, Dillon CP, Lalaoui N, Tanzer MC, Rodriguez DA, Lin A, Lebois M, Hakem R, Josefsson

EC, O'Reilly LA *et al* (2016) The Pseudokinase MLKL and the Kinase RIPK3 Have Distinct Roles in

Autoimmune Disease Caused by Loss of Death-Receptor-Induced Apoptosis. *Immunity* 45: 513-526

Ananthakrishnan AN (2015) Epidemiology and risk factors for IBD. *Nat Rev Gastroenterol Hepatol*

12: 205-217

Bader SM, Preston SP, Saliba K, Lipszyc A, Grant ZL, Mackiewicz L, Baldi A, Hempel A, Clark MP,

Peiris T *et al* (2023) Endothelial Caspase-8 prevents fatal necroptotic hemorrhage caused by

commensal bacteria. *Cell Death Differ* 30: 27-36

Ben-Moshe S, Shapira Y, Moor AE, Manco R, Veg T, Bahar Halpern K, Itzkovitz S (2019) Spatial

sorting enables comprehensive characterization of liver zonation. *Nat Metab* 1: 899-911

Benjamini Y, Hochberg Y (1995) Controlling the False Discovery Rate: A Practical and Powerful

Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)* 57:

289-300

Berkova L, Fazilaty H, Yang Q, Kubovciak J, Stastna M, Hrckulak D, Vojtechova M, Dalessi T, Brugger

MD, Hausmann G *et al* (2023) Terminal differentiation of villus tip enterocytes is governed by

distinct Tgfbeta superfamily members. *EMBO Rep* 24: e56454

Bobrow MN, Moen PT, Jr. (2001) Tyramide signal amplification (TSA) systems for the enhancement

of ISH signals in cytogenetics. *Curr Protoc Cytom* Chapter 8: Unit 8 9

Bossen C, Ingold K, Tardivel A, Bodmer JL, Gaide O, Hertig S, Ambrose C, Tschopp J, Schneider P

(2006) Interactions of tumor necrosis factor (TNF) and TNF receptor family members in the mouse

and human. *J Biol Chem* 281: 13964-13971

Butler A, Hoffman P, Smibert P, Papalexi E, Satija R (2018) Integrating single-cell transcriptomic data

across different conditions, technologies, and species. *Nat Biotechnol* 36: 411-420

Chen A, Liao S, Cheng M, Ma K, Wu L, Lai Y, Qiu X, Yang J, Xu J, Hao S *et al* (2022a) Spatiotemporal

transcriptomic atlas of mouse organogenesis using DNA nanoball-patterned arrays. *Cell* 185: 1777-

1792 e1721

Chen W, Zhou Z, Li L, Zhong CQ, Zheng X, Wu X, Zhang Y, Ma H, Huang D, Li W *et al* (2013) Diverse

sequence determinants control human and mouse receptor interacting protein 3 (RIP3) and mixed

lineage kinase domain-like (MLKL) interaction in necroptotic signaling. *J Biol Chem* 288: 16247-

16261

Chen X, Li W, Ren J, Huang D, He WT, Song Y, Yang C, Li W, Zheng X, Chen P *et al* (2014)

Translocation of mixed lineage kinase domain-like protein to plasma membrane leads to necrotic

cell death. *Cell Res* 24: 105-121

Chen X, Zhu R, Zhong J, Ying Y, Wang W, Cao Y, Cai H, Li X, Shuai J, Han J (2022b) Mosaic

composition of RIP1-RIP3 signalling hub and its role in regulating cell death. *Nat Cell Biol*

Cho YS, Challa S, Moquin D, Genga R, Ray TD, Guildford M, Chan FK (2009) Phosphorylation-driven

assembly of the RIP1-RIP3 complex regulates programmed necrosis and virus-induced

inflammation. *Cell* 137: 1112-1123

Choi ME, Price DR, Ryter SW, Choi AMK (2019) Necroptosis: a crucial pathogenic mediator of

human disease. *JCI Insight* 4

Cook WD, Moujalled DM, Ralph TJ, Lock P, Young SN, Murphy JM, Vaux DL (2014) RIPK1- and RIPK3-

induced cell death mode is determined by target availability. *Cell Death Differ* 21: 1600-1612

Davies KA, Fitzgibbon C, Young SN, Garnish SE, Yeung W, Coursier D, Birkinshaw RW, Sandow JJ,

Lehmann WIL, Liang LY *et al* (2020) Distinct pseudokinase domain conformations underlie

divergent activation mechanisms among vertebrate MLKL orthologues. *Nat Commun* 11: 3060

de Almagro MC, Goncharov T, Izrael-Tomasevic A, Duttler S, Kist M, Varfolomeev E, Wu X, Lee WP,

Murray J, Webster JD *et al* (2017) Coordinated ubiquitination and phosphorylation of RIP1

regulates necroptotic cell death. *Cell Death Differ* 24: 26-37

Degterev A, Hitomi J, Germscheid M, Ch'en IL, Korkina O, Teng X, Abbott D, Cuny GD, Yuan C,

Wagner G *et al* (2008) Identification of RIP1 kinase as a specific cellular target of necrostatins. *Nat*

Chem Biol 4: 313-321

Degterev A, Huang Z, Boyce M, Li Y, Jagtap P, Mizushima N, Cuny GD, Mitchison TJ, Moskowitz MA,

Yuan J (2005) Chemical inhibitor of nonapoptotic cell death with therapeutic potential for ischemic

brain injury. *Nat Chem Biol* 1: 112-119

Devos M, Tanghe G, Gilbert B, Dierick E, Verheirstraeten M, Nemegeer J, de Reuver R, Lefebvre S,

De Munck J, Rehwinkel J *et al* (2020) Sensing of endogenous nucleic acids by ZBP1 induces

keratinocyte necroptosis and skin inflammation. *J Exp Med* 217

Doerflinger M, Deng Y, Whitney P, Salvamoser R, Engel S, Kueh AJ, Tai L, Bachem A, Gressier E,

Geoghegan ND *et al* (2020) Flexible Usage and Interconnectivity of Diverse Cell Death Pathways

Protect against Intracellular Infection. *Immunity* 53: 533-547 e537

Dominguez S, Varfolomeev E, Brendza R, Stark K, Tea J, Imperio J, Ngu H, Earr T, Foreman O,

Webster JD *et al* (2021) Genetic inactivation of RIP1 kinase does not ameliorate disease in a mouse

model of ALS. *Cell Death Differ* 28: 915-931

Duprez L, Takahashi N, Van Hauwermeiren F, Vandendriessche B, Goossens V, Vanden Berghe T,

Declercq W, Libert C, Cauwels A, Vandenabeele P (2011) RIP kinase-dependent necrosis drives

lethal systemic inflammatory response syndrome. *Immunity* 35: 908-918

Fang Z, Wei H, Gou W, Chen L, Bi C, Hou W, Li Y (2021) Recent progress in small-molecule inhibitors

for critical therapeutic targets of necroptosis. *Future Med Chem* 13: 817-837

Fletcher-Etherington A, Nobre L, Nightingale K, Antrobus R, Nichols J, Davison AJ, Stanton RJ,

Weekes MP (2020) Human cytomegalovirus protein pUL36: A dual cell death pathway inhibitor.

Proc Natl Acad Sci U S A 117: 18771-18779

Garnish SE, Martin KR, Kauppi M, Jackson VE, Ambrose R, Eng VV, Chiou S, Meng Y, Frank D, Tovey

Crutchfield EC *et al* (2023) A common human MLKL polymorphism confers resistance to negative

regulation by phosphorylation. *Nat Commun* 14: 6046

Garnish SE, Meng Y, Koide A, Sandow JJ, Denbaum E, Jacobsen AV, Yeung W, Samson AL, Horne CR,

Fitzgibbon C *et al* (2021) Conformational interconversion of MLKL and disengagement from RIPK3

precede cell death by necroptosis. *Nat Commun* 12: 2211

Ge Y, Ezzell RM, Warren HS (2000) Localization of endotoxin in the rat intestinal epithelium. *J Infect*

Dis 182: 873-881

Geiger T, Velic A, Macek B, Lundberg E, Kampf C, Nagaraj N, Uhlen M, Cox J, Mann M (2013) Initial

quantitative proteomic map of 28 mouse tissues using the SILAC mouse. *Mol Cell Proteomics* 12:

1709-1722

Gevers D, Kugathasan S, Denson LA, Vazquez-Baeza Y, Van Treuren W, Ren B, Schwager E, Knights D,

Song SJ, Yassour M *et al* (2014) The treatment-naive microbiome in new-onset Crohn's disease. *Cell*

Host Microbe 15: 382-392

Graham DB, Xavier RJ (2020) Pathway paradigms revealed from the genetics of inflammatory

bowel disease. *Nature* 578: 527-539

Gunther C, Buchen B, He GW, Hornef M, Torow N, Neumann H, Wittkopf N, Martini E, Basic M,

Bleich A *et al* (2015) Caspase-8 controls the gut response to microbial challenges by Tnf-alpha-

dependent and independent pathways. *Gut* 64: 601-610

Gunther C, Martini E, Wittkopf N, Amann K, Weigmann B, Neumann H, Waldner MJ, Hedrick SM,

Tenzer S, Neurath MF *et al* (2011) Caspase-8 regulates TNF-alpha-induced epithelial necroptosis

and terminal ileitis. *Nature* 477: 335-339

Gupta K, Liu B (2021) PLK1-mediated S369 phosphorylation of RIPK3 during G2 and M phases

enables its ripoptosome incorporation and activity. *iScience* 24: 102320

Haase R, Royer LA, Steinbach P, Schmidt D, Dibrov A, Schmidt U, Weigert M, Maghelli N, Tomancak

P, Jug F *et al* (2020) CLIJ: GPU-accelerated image processing for everyone. *Nat Methods* 17: 5-6

Haber AL, Biton M, Rogel N, Herbst RH, Shekhar K, Smillie C, Burgin G, Delorey TM, Howitt MR,

Katz Y *et al* (2017) A single-cell survey of the small intestinal epithelium. *Nature* 551: 333-339

Hao Y, Hao S, Andersen-Nissen E, Mauck WM, 3rd, Zheng S, Butler A, Lee MJ, Wilk AJ, Darby C,

Zager M *et al* (2021) Integrated analysis of multimodal single-cell data. *Cell* 184: 3573-3587 e3529

Harris PA, Berger SB, Jeong JU, Nagilla R, Bandyopadhyay D, Campobasso N, Capriotti CA, Cox JA,

Dare L, Dong X *et al* (2017) Discovery of a First-in-Class Receptor Interacting Protein 1 (RIP1) Kinase

Specific Clinical Candidate (GSK2982772) for the Treatment of Inflammatory Diseases. *J Med Chem*

60: 1247-1261

He P, Ai T, Yang ZH, Wu J, Han J (2021) Detection of necroptosis by phospho-MLKL

immunohistochemical labeling. *STAR Protoc* 2: 100251

He S, Liang Y, Shao F, Wang X (2011) Toll-like receptors activate programmed necrosis in

macrophages through a receptor-interacting kinase-3-mediated pathway. *Proc Natl Acad Sci U S A*

108: 20054-20059

He S, Wang L, Miao L, Wang T, Du F, Zhao L, Wang X (2009) Receptor interacting protein kinase-3

determines cellular necrotic response to TNF-alpha. *Cell* 137: 1100-1111

Hildebrand JM, Kauppi M, Majewski IJ, Liu Z, Cox AJ, Miyake S, Petrie EJ, Silk MA, Li Z, Tanzer MC *et*

al (2020) A missense mutation in the MLKL brace region promotes lethal neonatal inflammation

and hematopoietic dysfunction. *Nat Commun* 11: 3150

Hildebrand JM, Tanzer MC, Lucet IS, Young SN, Spall SK, Sharma P, Pierotti C, Garnier JM, Dobson

RC, Webb AI *et al* (2014) Activation of the pseudokinase MLKL unleashes the four-helix bundle

domain to induce membrane localization and necroptotic cell death. *Proc Natl Acad Sci U S A* 111:

15072-15077

Horne CR, Samson AL, Murphy JM (2023) The web of death: the expanding complexity of
necroptotic signaling. *Trends Cell Biol* 33: 162-174

Ito Y, Ofengeim D, Najafov A, Das S, Saberi S, Li Y, Hitomi J, Zhu H, Chen H, Mayo L *et al* (2016)
RIPK1 mediates axonal degeneration by promoting inflammation and necroptosis in ALS. *Science*
353: 603-608

Jacobsen AV, Pierotti CL, Lowes KN, Au AE, Zhang Y, Etemadi N, Fitzgibbon C, Kersten WJA, Samson
AL, van Delft MF *et al* (2022) The Lck inhibitor, AMG-47a, blocks necroptosis and implicates RIPK1
in signalling downstream of MLKL. *Cell Death Dis* 13: 291

Kaiser WJ, Sridharan H, Huang C, Mandal P, Upton JW, Gough PJ, Sehon CA, Marquis RW, Bertin J,
Mocarski ES (2013) Toll-like receptor 3-mediated necrosis via TRIF, RIP3, and MLKL. *J Biol Chem*
288: 31268-31279

Kaiser WJ, Upton JW, Long AB, Livingston-Rosanoff D, Daley-Bauer LP, Hakem R, Caspary T,
Mocarski ES (2011) RIP3 mediates the embryonic lethality of caspase-8-deficient mice. *Nature* 471:
368-372

Kelliher MA, Grimm S, Ishida Y, Kuo F, Stanger BZ, Leder P (1998) The death domain kinase RIP
mediates the TNF-induced NF-kappaB signal. *Immunity* 8: 297-303

Kim D, Paggi JM, Park C, Bennett C, Salzberg SL (2019) Graph-based genome alignment and
genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* 37: 907-915

Kobayashi T, Siegmund B, Le Berre C, Wei SC, Ferrante M, Shen B, Bernstein CN, Danese S, Peyrin-
Biroulet L, Hibi T (2020) Ulcerative colitis. *Nat Rev Dis Primers* 6: 74

Kong IY, Trezise S, Light A, Todorovski I, Arnau GM, Gadipally S, Yoannidis D, Simpson KJ, Dong X,
Whitehead L *et al* (2022) Epigenetic modulators of B cell fate identified through coupled
phenotype-transcriptome analysis. *Cell Death Differ* 29: 2519-2530

Landini G, Martinelli G, Piccinini F (2021) Colour deconvolution: stain unmixing in histological
imaging. *Bioinformatics* 37: 1485-1487

Law CW, Chen Y, Shi W, Smyth GK (2014) voom: Precision weights unlock linear model analysis

tools for RNA-seq read counts. *Genome Biol* 15: R29

Li D, Meng L, Xu T, Su Y, Liu X, Zhang Z, Wang X (2017) RIPK1-RIPK3-MLKL-dependent necrosis

promotes the aging of mouse male reproductive system. *Elife* 6

Li L, Huang K, Ruan C, Han J, Zhang Y (2022) Immunostaining of phospho-RIPK3 in L929 cells,

murine yolk sacs, ceca, and small intestines. *STAR Protoc* 3: 101517

Li X, Zhang M, Huang X, Liang W, Li G, Lu X, Li Y, Pan H, Shi L, Zhu H *et al* (2020) Ubiquitination of

RIPK1 regulates its activation mediated by TNFR1 and TLRs signaling in distinct manners. *Nat*

Commun 11: 6364

Li X, Zhong CQ, Wu R, Xu X, Yang ZH, Cai S, Wu X, Chen X, Yin Z, He Q *et al* (2021) RIP1-dependent

linear and nonlinear recruitments of caspase-8 and RIP3 respectively to necrosome specify distinct

cell death outcomes. *Protein Cell* 12: 858-876

Liao Y, Smyth GK, Shi W (2014) featureCounts: an efficient general purpose program for assigning

sequence reads to genomic features. *Bioinformatics* 30: 923-930

Liao Y, Smyth GK, Shi W (2019) The R package Rsubread is easier, faster, cheaper and better for

alignment and quantification of RNA sequencing reads. *Nucleic Acids Res* 47: e47

Liccardi G, Ramos Garcia L, Tenev T, Annibaldi A, Legrand AJ, Robertson D, Feltham R, Anderton H,

Darding M, Peltzer N *et al* (2019) RIPK1 and Caspase-8 Ensure Chromosome Stability

Independently of Their Role in Cell Death and Inflammation. *Mol Cell* 73: 413-428 e417

Linkermann A, Brasen JH, Darding M, Jin MK, Sanz AB, Heller JO, De Zen F, Weinlich R, Ortiz A,

Walczak H *et al* (2013) Two independent pathways of regulated necrosis mediate ischemia-

reperfusion injury. *Proc Natl Acad Sci U S A* 110: 12024-12029

Liu Z, Nailwal H, Rector J, Rahman MM, Sam R, McFadden G, Chan FK (2021) A class of viral inducer

of degradation of the necroptosis adaptor RIPK3 regulates virus-induced inflammation. *Immunity*

54: 247-258 e247

Lu Z, Van Eeckhoutte HP, Liu G, Nair PM, Jones B, Gillis CM, Nalkurthi BC, Verhamme F, Buyle-
Huybrecht T, Vandenabeele P *et al* (2021) Necroptosis Signaling Promotes Inflammation, Airway
Remodeling, and Emphysema in Chronic Obstructive Pulmonary Disease. *Am J Respir Crit Care Med*
204: 667-681

Luedde M, Lutz M, Carter N, Sosna J, Jacoby C, Vucur M, Gautheron J, Roderburg C, Borg N,
Reisinger F *et al* (2014) RIP3, a kinase promoting necroptotic cell death, mediates adverse
remodelling after myocardial infarction. *Cardiovasc Res* 103: 206-216

Ly A, Liao Y, Pietrzak H, Ioannidis LJ, Sidwell T, Gloury R, Doerflinger M, Triglia T, Qin RZ, Groom JR
et al (2019) Transcription Factor T-bet in B Cells Modulates Germinal Center Polarization and
Antibody Affinity Maturation in Response to Malaria. *Cell Rep* 29: 2257-2269 e2256

Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads.
2011 17: 3

Matsuzawa-Ishimoto Y, Shono Y, Gomez LE, Hubbard-Lucey VM, Cammer M, Neil J, Dewan MZ,
Lieberman SR, Lazrak A, Marinis JM *et al* (2017) Autophagy protein ATG16L1 prevents necroptosis
in the intestinal epithelium. *J Exp Med* 214: 3687-3705

Moor AE, Harnik Y, Ben-Moshe S, Massasa EE, Rozenberg M, Eilam R, Bahar Halpern K, Itzkovitz S
(2018) Spatial Reconstruction of Single Enterocytes Uncovers Broad Zonation along the Intestinal
Villus Axis. *Cell* 175: 1156-1167 e1115

Mosli MH, Feagan BG, Zou G, Sandborn WJ, D'Haens G, Khanna R, Shackelton LM, Walker CW,
Nelson S, Vandervoort MK *et al* (2017) Development and validation of a histological index for UC.
Gut 66: 50-58

Murphy JM, Czabotar PE, Hildebrand JM, Lucet IS, Zhang JG, Alvarez-Diaz S, Lewis R, Lalaoui N,
Metcalf D, Webb AI *et al* (2013) The pseudokinase MLKL mediates necroptosis via a molecular
switch mechanism. *Immunity* 39: 443-453

Naito MG, Xu D, Amin P, Lee J, Wang H, Li W, Kelliher M, Pasparakis M, Yuan J (2020) Sequential

activation of necroptosis and apoptosis cooperates to mediate vascular and neural pathology in

stroke. *Proc Natl Acad Sci U S A* 117: 4959-4970

Najafov A, Zervantonakis IK, Mookhtiar AK, Greninger P, March RJ, Egan RK, Luu HS, Stover DG,

Matulonis UA, Benes CH *et al* (2018) BRAF and AXL oncogenes drive RIPK3 expression loss in

cancer. *PLoS Biol* 16: e2005756

Negroni A, Colantoni E, Pierdomenico M, Palone F, Costanzo M, Oliva S, Tiberti A, Cucchiara S,

Stronati L (2017) RIP3 AND pMLKL promote necroptosis-induced inflammation and alter

membrane permeability in intestinal epithelial cells. *Dig Liver Dis* 49: 1201-1210

Newton K, Dugger DL, Maltzman A, Greve JM, Hedehus M, Martin-McNulty B, Carano RA, Cao TC,

van Bruggen N, Bernstein L *et al* (2016) RIPK3 deficiency or catalytically inactive RIPK1 provides

greater benefit than MLKL deficiency in mouse models of inflammation and tissue injury. *Cell*

Death Differ 23: 1565-1576

Newton K, Dugger DL, Wickliffe KE, Kapoor N, de Almagro MC, Vucic D, Komuves L, Ferrando RE,

French DM, Webster J *et al* (2014) Activity of protein kinase RIPK3 determines whether cells die by

necroptosis or apoptosis. *Science* 343: 1357-1360

Newton K, Sun X, Dixit VM (2004) Kinase RIP3 is dispensable for normal NF-kappa Bs, signaling by

the B-cell and T-cell receptors, tumor necrosis factor receptor 1, and Toll-like receptors 2 and 4.

Mol Cell Biol 24: 1464-1469

O'Reilly LA, Divisekera U, Newton K, Scalzo K, Kataoka T, Puthalakath H, Ito M, Huang DC, Strasser A

(2004) Modifications and intracellular trafficking of FADD/MORT1 and caspase-8 after stimulation

of T lymphocytes. *Cell Death Differ* 11: 724-736

Palmer SN, Chappidi S, Pinkham C, Hancks DC (2021) Evolutionary Profile for (Host and Viral) MLKL

Indicates Its Activities as a Battlefront for Extensive Counteradaptation. *Mol Biol Evol* 38: 5405-

5422

Park JH, Kotani T, Konno T, Setiawan J, Kitamura Y, Imada S, Usui Y, Hatano N, Shinohara M, Saito Y

et al (2016) Promotion of Intestinal Epithelial Cell Turnover by Commensal Bacteria: Role of Short-

Chain Fatty Acids. *PLoS One* 11: e0156334

Pearson JS, Giogha C, Muhlen S, Nachbur U, Pham CL, Zhang Y, Hildebrand JM, Oates CV, Lung TW,

Ingle D *et al* (2017) EspL is a bacterial cysteine protease effector that cleaves RHIM proteins to

block necroptosis and inflammation. *Nat Microbiol* 2: 16258

Pedersen A, Valla A, Bofin M, De Frutos JP, Reinersten I, Smistad E (2021) FastPathology: An Open-

Source Platform for Deep Learning-Based Research and Decision Support in Digital Pathology. *IEEE*

Access 9: 58216-58229

Percie du Sert N, Hurst V, Ahluwalia A, Alam S, Avey MT, Baker M, Browne WJ, Clark A, Cuthill IC,

Dirnagl U *et al* (2020) The ARRIVE guidelines 2.0: Updated guidelines for reporting animal research.

PLoS Biol 18: e3000410

Petrie EJ, Sandow JJ, Jacobsen AV, Smith BJ, Griffin MDW, Lucet IS, Dai W, Young SN, Tanzer MC,

Wardak A *et al* (2018) Conformational switching of the pseudokinase domain promotes human

MLKL tetramerization and cell death by necroptosis. *Nat Commun* 9: 2422

Petrie EJ, Sandow JJ, Lehmann WIL, Liang LY, Coursier D, Young SN, Kersten WJA, Fitzgibbon C,

Samson AL, Jacobsen AV *et al* (2019) Viral MLKL Homologs Subvert Necroptotic Cell Death by

Sequestering Cellular RIPK3. *Cell Rep* 28: 3309-3319 e3305

Pettersen HS, Belevich I, Royset ES, Smistad E, Simpson MR, Jokitalo E, Reinertsen I, Bakke I,

Pedersen A (2021) Code-Free Development and Deployment of Deep Segmentation Models for

Digital Pathology. *Front Med (Lausanne)* 8: 816281

Pierdomenico M, Negroni A, Stronati L, Vitali R, Prete E, Bertin J, Gough PJ, Aloi M, Cucchiara S

(2014) Necroptosis is active in children with inflammatory bowel disease and contributes to

heighten intestinal inflammation. *Am J Gastroenterol* 109: 279-287

Preston SP, Stutz MD, Allison CC, Nachbur U, Gouil Q, Tran BM, Duvivier V, Arandjelovic P, Cooney JP, Mackiewicz L *et al* (2022) Epigenetic Silencing of RIPK3 in Hepatocytes Prevents MLKL-mediated Necroptosis From Contributing to Liver Pathologies. *Gastroenterology* 163: 1643-1657 e1614

Ramirez MLG, Salvesen GS (2018) A primer on caspase mechanisms. *Semin Cell Dev Biol* 82: 79-85

Rickard JA, O'Donnell JA, Evans JM, Lalaoui N, Poh AR, Rogers T, Vince JE, Lawlor KE, Ninnis RL, Anderton H *et al* (2014) RIPK1 regulates RIPK3-MLKL-driven systemic inflammation and emergency hematopoiesis. *Cell* 157: 1175-1188

Robinson MD, Oshlack A (2010) A scaling normalization method for differential expression analysis of RNA-seq data. *Genome Biol* 11: R25

Roda G, Chien Ng S, Kotze PG, Argollo M, Panaccione R, Spinelli A, Kaser A, Peyrin-Biroulet L, Danese S (2020) Crohn's disease. *Nat Rev Dis Primers* 6: 22

Rodriguez DA, Quarato G, Liedmann S, Tummers B, Zhang T, Guy C, Crawford JC, Palacios G, Pelletier S, Kalkavan H *et al* (2022) Caspase-8 and FADD prevent spontaneous ZBP1 expression and necroptosis. *Proc Natl Acad Sci U S A* 119: e2207240119

Rodriguez DA, Weinlich R, Brown S, Guy C, Fitzgerald P, Dillon CP, Oberst A, Quarato G, Low J, Cripps JG *et al* (2016) Characterization of RIPK3-mediated phosphorylation of the activation loop of MLKL during necroptosis. *Cell Death Differ* 23: 76-88

Salmena L, Lemmers B, Hakem A, Matysiak-Zablocki E, Murakami K, Au PY, Berry DM, Tamblyn L, Shehabeldin A, Migon E *et al* (2003) Essential role for caspase 8 in T-cell homeostasis and T-cell-mediated immunity. *Genes Dev* 17: 883-895

Samson AL, Fitzgibbon C, Patel KM, Hildebrand JM, Whitehead LW, Rimes JS, Jacobsen AV, Horne CR, Gavin XJ, Young SN *et al* (2021a) A toolbox for imaging RIPK1, RIPK3, and MLKL in mouse and human cells. *Cell Death Differ* 28: 2126-2144

Samson AL, Garnish SE, Hildebrand JM, Murphy JM (2021b) Location, location, location: A compartmentalized view of TNF-induced necroptotic signaling. *Sci Signal* 14

Samson AL, Zhang Y, Geoghegan ND, Gavin XJ, Davies KA, Mlodzianoski MJ, Whitehead LW, Frank D,

Garnish SE, Fitzgibbon C *et al* (2020) MLKL trafficking and accumulation at the plasma membrane

control the kinetics and threshold for necroptosis. *Nat Commun* 11: 3151

Satija R, Farrell JA, Gennert D, Schier AF, Regev A (2015) Spatial reconstruction of single-cell gene

expression data. *Nat Biotechnol* 33: 495-502

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C,

Saalfeld S, Schmid B *et al* (2012) Fiji: an open-source platform for biological-image analysis. *Nat*

Methods 9: 676-682

Schwarzer R, Jiao H, Wachsmuth L, Tresch A, Pasparakis M (2020) FADD and Caspase-8 Regulate

Gut Homeostasis and Inflammation by Controlling MLKL- and GSDMD-Mediated Death of Intestinal

Epithelial Cells. *Immunity* 52: 978-993 e976

Shi Y, Cui X, Sun Y, Zhao Q, Liu T (2020) Intestinal vitamin D receptor signaling ameliorates dextran

sulfate sodium-induced colitis by suppressing necroptosis of intestinal epithelial cells. *FASEB J* 34:

13494-13506

Smith KG, Light A, O'Reilly LA, Ang SM, Strasser A, Tarlinton D (2000) bcl-2 transgene expression

inhibits apoptosis in the germinal center and reveals differences in the selection of memory B cells

and bone marrow antibody-forming cells. *J Exp Med* 191: 475-484

Stringer C, Wang T, Michaelos M, Pachitariu M (2021) Cellpose: a generalist algorithm for cellular

segmentation. *Nat Methods* 18: 100-106

Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, 3rd, Hao Y, Stoeckius M,

Smibert P, Satija R (2019) Comprehensive Integration of Single-Cell Data. *Cell* 177: 1888-1902

e1821

Sun L, Wang H, Wang Z, He S, Chen S, Liao D, Wang L, Yan J, Liu W, Lei X *et al* (2012) Mixed lineage

kinase domain-like protein mediates necrosis signaling downstream of RIP3 kinase. *Cell* 148: 213-

227

Tabula Muris C, Overall c, Logistical c, Organ c, processing, Library p, sequencing, Computational

data a, Cell type a, Writing g *et al* (2018) Single-cell transcriptomics of 20 mouse organs creates a

Tabula Muris. *Nature* 562: 367-372

Tanzer MC, Khan N, Rickard JA, Etemadi N, Lalaoui N, Spall SK, Hildebrand JM, Segal D, Miasari M,

Chau D *et al* (2017) Combination of IAP antagonist and IFNgamma activates novel caspase-10- and

RIPK1-dependent cell death pathways. *Cell Death Differ* 24: 481-491

Tanzer MC, Matti I, Hildebrand JM, Young SN, Wardak A, Tripaydonis A, Petrie EJ, Mildenhall AL,

Vaux DL, Vince JE *et al* (2016) Evolutionary divergence of the necroptosis effector MLKL. *Cell Death*

Differ 23: 1185-1197

Tisch N, Mogler C, Stojanovic A, Luck R, Korhonen EA, Ellerkmann A, Adler H, Singhal M,

Schermann G, Erkert L *et al* (2022) Caspase-8 in endothelial cells maintains gut homeostasis and

prevents small bowel inflammation in mice. *EMBO Mol Med*: e14121

Tovey Crutchfield EC, Garnish SE, Day J, Anderton H, Chiou S, Hempel A, Hall C, Patel KM,

Gangatirkar P, Martin KR *et al* (2023) MLKL deficiency protects against low-grade, sterile

inflammation in aged mice. *Cell Death Differ* 30: 1059-1071

Uhlen M, Fagerberg L, Hallstrom BM, Lindskog C, Oksvold P, Mardinoglu A, Sivertsson A, Kampf C,

Sjostedt E, Asplund A *et al* (2015) Proteomics. Tissue-based map of the human proteome. *Science*

347: 1260419

Upton JW, Kaiser WJ, Mocarski ES (2010) Virus inhibition of RIP3-dependent necrosis. *Cell Host*

Microbe 7: 302-313

Vince JE, Wong WW, Khan N, Feltham R, Chau D, Ahmed AU, Benetatos CA, Chunduru SK, Condon

SM, McKinlay M *et al* (2007) IAP antagonists target cIAP1 to induce TNFalpha-dependent

apoptosis. *Cell* 131: 682-693

Vlantis K, Wullaert A, Polykratis A, Kondylis V, Dannappel M, Schwarzer R, Welz P, Corona T,

Walczak H, Weih F *et al* (2016) NEMO Prevents RIP Kinase 1-Mediated Epithelial Cell Death and

Chronic Intestinal Inflammation by NF-kappaB-Dependent and -Independent Functions. *Immunity*

44: 553-567

Vucur M, Ghallab A, Schneider AT, Adili A, Cheng M, Castoldi M, Singer MT, Buttner V, Keysberg LS,

Kusgens L *et al* (2023) Sublethal necroptosis signaling promotes inflammation and liver cancer.

Immunity 56: 1578-1595 e1578

Wang H, Sun L, Su L, Rizo J, Liu L, Wang LF, Wang FS, Wang X (2014) Mixed lineage kinase domain-

like protein MLKL causes necrotic membrane disruption upon phosphorylation by RIP3. *Mol Cell*

54: 133-146

Wang R, Li H, Wu J, Cai ZY, Li B, Ni H, Qiu X, Chen H, Liu W, Yang ZH *et al* (2020a) Gut stem cell

necroptosis by genome instability triggers bowel inflammation. *Nature* 580: 386-390

Wang T, Perera ND, Chiam MDF, Cuic B, Wanniarachchillage N, Tomas D, Samson AL, Cawthorne W,

Valor EN, Murphy JM *et al* (2020b) Necroptosis is dispensable for motor neuron degeneration in a

mouse model of ALS. *Cell Death Differ* 27: 1728-1739

Webster JD, Solon M, Haller S, Newton K (2018) Detection of Necroptosis by Phospho-RIPK3

Immunohistochemical Labeling. *Methods Mol Biol* 1857: 153-160

Weisel K, Scott N, Berger S, Wang S, Brown K, Powell M, Broer M, Watts C, Tompson DJ, Burriss SW

et al (2021) A randomised, placebo-controlled study of RIPK1 inhibitor GSK2982772 in patients

with active ulcerative colitis. *BMJ Open Gastroenterol* 8

Xie Y, Zhao Y, Shi L, Li W, Chen K, Li M, Chen X, Zhang H, Li T, Matsuzawa-Ishimoto Y *et al* (2020) Gut

epithelial TSC1/mTOR controls RIPK3-dependent necroptosis in intestinal inflammation and cancer.

J Clin Invest 130: 2111-2128

Xu J, Li S, Jin W, Zhou H, Zhong T, Cheng X, Fu Y, Xiao P, Cheng H, Wang D *et al* (2023) Epithelial

Gab1 calibrates RIPK3-dependent necroptosis to prevent intestinal inflammation. *JCI Insight* 8

Yeap HW, Chen KW (2022) RIPK1 and RIPK3 in antibacterial defence. *Biochem Soc Trans* 50: 1583-

1594

Zelic M, Kelliher MA (2018) Analyzing Necroptosis Using an RIPK1 Kinase Inactive Mouse Model of

TNF Shock. *Methods Mol Biol* 1857: 125-134

Zelic M, Roderick JE, O'Donnell JA, Lehman J, Lim SE, Janardhan HP, Trivedi CM, Pasparakis M,

Kelliher MA (2018) RIP kinase 1-dependent endothelial necroptosis underlies systemic

inflammatory response syndrome. *J Clin Invest* 128: 2064-2075

Zhang T, Yin C, Boyd DF, Quarato G, Ingram JP, Shubina M, Ragan KB, Ishizuka T, Crawford JC,

Tummers B *et al* (2020) Influenza Virus Z-RNAs Induce ZBP1-Mediated Necroptosis. *Cell* 180: 1115-

1129 e1113

Zhao J, Jitkaew S, Cai Z, Choksi S, Li Q, Luo J, Liu ZG (2012) Mixed lineage kinase domain-like is a

key receptor interacting protein 3 downstream component of TNF-induced necrosis. *Proc Natl*

Acad Sci U S A 109: 5322-5327

Zhu X, Yang N, Yang Y, Yuan F, Yu D, Zhang Y, Shu Z, Nan N, Hu H, Liu X *et al* (2022) Spontaneous

necroptosis and autoinflammation are blocked by an inhibitory phosphorylation on MLKL during

neonatal development. *Cell Res* 32: 407-410

1403

1404

1405

Acknowledgements:

We thank James Vince, Jiyi Pang, David Tarlington, Angus Stock, Najoua Lalaoui, Asha Jois, Marcel Doerflinger, Quentin Gouil, Eric Hanssen and Bruce Rosengarten for constructive feedback and/or reagents during the preparation of this manuscript. We thank WEHI Monoclonal antibody facility for producing several antibodies used in this study. We thank the WEHI Histology team for high level support with immunohistochemistry. We thank the WEHI Bioservices team for high level support for animal experimentation, welfare, and ethics. We thank Kim Newton and Vishva Dixit for sharing the $Ripk3^{-/-}$ mice that were used to make the $Casp8^{-/-}Ripk3^{-/-}$ mice, and thank Michelle Kelliher for the $Ripk1^{-/-}$ mice that were used to generate the $Casp8^{-/-}Ripk1^{-/-}Ripk3^{-/-}$ mice. We thank Anaxis Pharma Pty Ltd for providing human RIPK3 expressing mice for our control experiments, and thank Andrew Kueh and Ueli Nachbur for their roles in generating this strain. We are grateful to the Department of Gastroenterology at the Royal Melbourne Hospital for performing endoscopies and to the patients who consented to donating their material to be used in this study.

Funding:

This work was supported by National Health and Medical Research Council of Australia (grants 1172929 to JMM, 2008652 to EDH, 2002965 to ALS, and the Independent Research Institutes Infrastructure Support Scheme 9000719), by the Kenneth Rainin Foundation (award to JMM, BC, AHA, ALS), and by the Victorian State Government Operational Infrastructure Support scheme. SC is supported by the WEHI Handman PhD scholarship, and AHA by the Avant Foundation, Crohn's and Colitis Australia and the University of Melbourne Scholarships. JMM and JS received research funding from Anaxis Pharma Pty Ltd, from which the salaries of KMP, AH, AL and PG were paid.

Conflict of interests:

KMP, SNY, AH, AL, PG, CRH, JD, IPW, JMH, JS, JMM and ALS have contributed to the development of necroptosis pathway inhibitors in collaboration with Anaxis Pharma Pty Ltd. YP is a co-holder of patent for material arising from this study. All other authors have no additional financial interests.

Abbreviations:

C8 (Caspase-8), C10 (Caspase-10), cl. C3 (cleaved Caspase-3), TNF (Tumour Necrosis Factor), RIPK1 (Receptor-interacting serine/threonine-protein kinase-1), pRIPK1 (phosphorylated RIPK1), RIPK3 (Receptor-interacting serine/threonine-protein kinase-3), pRIPK3 (phosphorylated RIPK3), MLKL (Mixed lineage kinase domain-like protein), pMLKL (phosphorylated MLKL), R1 (Receptor-interacting serine/threonine-protein kinase-1), R3 (Receptor-interacting serine/threonine-protein kinase-1), ML (Mixed lineage kinase domain-like protein), Nec1s (Necrostatin1s), WEHI (Walter and Eliza Hall Institute of Medical Research), WT (wild-type), KO (knockout), EV (Expanded View), IBD (inflammatory bowel disease), UC (ulcerative colitis), CD (Crohn's disease), SMA (smooth muscle actin), Kpf (Kupffer cell), endo (endothelial cell), p.o. (oral administration), i.p. (intraperitoneal administration), CV (central vein), PV (portal vein), BD (bile duct), CA (splenic central artery), WP (splenic white pulp), MZ (splenic marginal zone), RP (splenic red pulp), CRP (C-reactive protein), TS (Tumour Necrosis Factor and Smac mimetic), TSI (Tumour Necrosis Factor and Smac mimetic and IDN-6556), Ab (antibody), Uniform Manifold Approximation and Projection (UMAP), NP-KLH (4-hydroxy-3-nitrophenylacetyl hapten conjugated to keyhole limpet hemocyanin), CDS (coding sequence).

Figure legends:

Fig. 1. Automated immunohistochemistry shows constitutive necroptotic pathway expression is restricted. A To gauge immunohistochemistry performance, immunosignals from wild-type (WT) versus knockout (KO) tissue were deconvoluted, (i) pixel intensities plotted, (ii) ratioed to yield a signal-to-noise (S/N) histogram, and then (iv) integrated. **B** Heatmap shows relative integrated S/N values from 7 automated immunohistochemistry protocols across 7 tissues. Column headers indicate the antibody target clone name. Data are representative of n≥3 for each target and tissue. **C** Heatmap depicts relative protein abundance values as measured by (Geiger *et al.*, 2013). **D** Immunosignals of Caspase-8, RIPK1, RIPK3 and MLKL in wild-type versus the appropriate knockout (KO) tissue from *Mlkl*$^{-/-}$ or *Casp8*$^{-/-}$*Ripk3*$^{-/-}$ or *Casp8*$^{-/-}$*Ripk1*$^{-/-}$*Ripk3*$^{-/-}$. Data are representative of n≥3 for each target and tissue. Scale bars are 500μm. Related to Fig. **EV1**.

Fig. 2. Necroptotic potential is spatially graded across tissue zones. A Immunosignals of Caspase-8, RIPK1 and RIPK3 from wild-type mouse ileum (i), colon (ii), liver (iii) and spleen (iv). The crypt base (crypt), villi/crypt tip (tip), central vein (CV), portal vein (PV), bile duct (BD), central artery (CA), white pulp (WP), marginal zone (MZ) and red pulp (RP) are annotated. Inset of immunostaining in the ileum shows lower RIPK3 expression in Paneth cells (open arrowhead) relative to neighbouring cells. Arrow shows peri-central hepatocytes that express higher levels of Caspase-8. Closed arrowheads show Caspase-8^{+} RIPK1^{+} RIPK3^{+} Kupffer cells. Scale bars are 50μm, except for 10μm scale bar in inset. Data are representative of n≥3 for each target and tissue. **B** Relative expression levels of Caspase-8, RIPK1, RIPK3 (and splenic MLKL; Fig. **EV2B**) along the indicated tissue axes. Red datapoints indicate immunosignal intensities, and the overlaid dark blue line indicating the LOWESS best-fit along N=20 axes per tissue. Best-fit curves

are superimposed in the left-most column. Dashed line indicates the boundary between

splenic white pulp and marginal zone. Data are representative of n>3 mice per target per

tissue. **C** Scatterplots where each dot represents a different cell ontology from the *Tabula*

Muris dataset *(Tabula Muris et al., 2018)*. The percent of cells within each ontology that

expressed *Mki67*, *Ripk1* or *Ripk3* was plotted against that of *Top2a*. Pearson correlation

coefficient values are shown. Related to Fig. **EV2**.

1489

Fig. 3. RIPK3 expression is rapidly altered during systemic inflammation. A

Experimental design. **B** Core temperatures of vehicle- and TNF-injected mice (n=3 mice

per group). **C** Immunosignals for cleaved Caspase-3 (cl. C3), Caspase-8, RIPK1 or RIPK3

from the spleen of vehicle- or TNF-injected mice. Insets show unidentified RIPK1^{high} cells

that associate with apoptotic bodies in splenic white pulp. **D** Graph of white pulp area

occupied by cleaved Caspase-3$^+$ material in vehicle- and TNF-treated mice. Each red

datapoint represents one white pulp lobule (N=20 lobules/mouse). Blue datapoints indicate

the median value per mouse (n=3 mice/treatment). Black bars represent the mean value

per group. *p<0.05 by unpaired 2-tailed t-test. **E** RIPK3 immunosignals in colon of vehicle-

or TNF-treated mice. **F** Best-fit curves of RIPK3 immunosignals along the crypt-to-tip axis

from N=10 axes per mouse (n=3 mice/group). ***p<0.001 by multiple unpaired 2-tailed t-

test. **G** RIPK3 (pink) and smooth muscle actin (brown) immunosignals in intestinal

submucosa of vehicle- or TNF-treated mice. Insets show vessel cross-sections.

Arrowheads show RIPK3$^+$ endothelial cells (endo). **H** Plot of RIPK3 signals per vessel.

Each red datapoint represents one vessel (N=50 vessels/mouse). Blue datapoints indicate

the median value per mouse (n=3 mice/treatment). Black bars represent the mean value

per group. *p<0.05 by unpaired 2-tailed t-test. **I** RIPK3 immunosignals in liver of vehicle- or

TNF-treated mice. Central vein (CV), portal vein (PV) and Kupffer cell (Kpf). **J** Plot of

RIPK3 signals per hepatocyte or Kupffer cell. Each transparent datapoint represents one

cell (N=90 cells/mouse). Opaque datapoints indicate the median value per mouse (n=3

mice/treatment). Black bars represent the mean value per group. *p<0.05 and **p<0.01 by

unpaired 2-tailed t-test. **K** Core temperatures of vehicle-, TNF- and Nec1s+TNF-injected

wild-type mice (n=4-5 mice/treatment; one dot/mouse/time). Line indicates mean. X

indicates a euthanised mouse due to its body temperature being <30°C. **L** RIPK3 levels in

serum from the mice in panel **K** or from untreated *Ripk3$^{-/-}$* mice. Data expressed as

arbitrary optical density units (A.U.). One dot per mouse. Mean ± SEM is shown. *p<0.05,

**p<0.01 by 1-way ANOVA with Tukey's post-hoc correction.

1517

Fig. 4. RIPK3 expression changes in response to dysbiosis. A Experimental design. **B**

Bulk RNA sequencing was performed on indicated tissues. Heatmap depicts the log fold-

change in gene expression for antibiotic- versus water-treated mice. Each row represents

a different mouse. Legend shows the colour-to-value scale. **C** Immunoblots for the

indicated proteins in the ileum and spleen of water- versus antibiotic-treated mice.

Arrowheads indicate full-length proteins of interest. Coomassie staining of total protein

content was used as a loading control. Data are representative of n=7 mice per tissue per

group. **D** Caspase-8, RIPK3 and RIPK3 immunosignals in the ileum of water- or antibiotic-

treated mice. Arrowheads to cytosolic accumulations of RIPK1 and RIPK3 in epithelial

cells at villi tips. Scale bars in lower magnification micrographs are 100μm. Scale bars in

insets are 10μm. Data are representative of n=7 mice per group. Related to Fig. **EV3**.

1529

1530 **Fig. 5. Automated immunohistochemistry quantifies necroptotic signaling in human**

1531 **cells. A** Immunosignals of cleaved Caspase-3, Caspase-8, RIPK1, RIPK3 and MLKL in

1532 wild-type versus *MLKL$^{-/-}$* or *RIPK1$^{-/-}$* or *CASP8$^{-/-}$CASP10$^{-/-}$MLKL$^{-/-}$* HT29 cells. Arrowheads

1533 indicate Caspase-8$^+$, RIPK1$^+$, RIPK3$^+$ and MLKL$^+$ puncta that are presumed to be

1534 necrosomes. Data are representative of n≥2 for each protein and treatment. Scale bars in

lower magnification micrographs are 10μm. Scale bars in insets are 2μm. **B** The percent of cells per treatment group that contain cytosolic necrosome-like puncta immunostained by the stipulated antibody. N=1051-5630 cells were analysed per condition per stain. Data representative of n=2 experiments. **C** The number of puncta per cell. N=1000 cells per treatment group were analysed. Each datapoint represents one cell. Black bar indicates mean value. ****p<0.0001 by one-way ANOVA with Krukal-Wallis post-hoc correction. Data representative of n=2 experiments. Related to Fig. **EV5**, **EV6** and **EV7**.

Fig. 6. Case study for the detection of necroptotic signaling in inflammatory bowel disease. A Study design. **B** Blinded histopathological (Robarts Histopathology Index; RHI) scores of disease activity in intestinal biopsies relative to their endoscopic grading of inflammation. Diamond indicates a sample that could not be formally scored, as it was solely comprised of neutrophilic exudate, but was given a pseudo-score of 15 that likely underrepresents the extent of disease activity in this biopsy. Biopsies scored in Panel B were matched to those used in Panels C-E and Fig. **EV8** (see Table **1** for details). **C** Immunoblot of lysates from HT29 cells (red annotations) and intestinal biopsies from patients A-H (blue annotations). The fifth lane of each gel contained lysates from TSI-treated $RIPK3^{-/-}$ or TSI-treated $MLKL^{-/-}$ cells (see source data for details). Patients A,C,E,G were non-IBD controls. Patients B and D had ulcerative colitis (UC). Patients F and H had Crohn's disease (CD). The endoscopic grading of the biopsy site as 'non-inflamed', 'marginally inflamed' or 'inflamed' is stipulated. Closed arrowheads indicate full-length form of proteins. Asterisks indicate active, phosphorylated forms of RIPK3 (pRIPK3) and MLKL (pMLKL). Open arrowheads indicate active, cleaved forms of Caspase-8, Caspase-10 and Caspase-3. GAPDH was used as a loading control. **D** Immunohistochemistry for Caspase-8 (clone B.925.8) on intestinal biopsies. Insets a-e show diffuse epithelial Caspase-8 in patient C. Insets f-j show mild clustering of epithelial Caspase-8 and insets k-o show more

pronounced clustering of epithelial Caspase-8 in patient D (arrowheads). Scale bars in

lower magnification micrographs are 500µm. Scale bars in insets are 10µm. **E** The number

of Caspase-8$^+$ puncta per 100 cells. Each datapoint represents one crypt. Whole slide

scans with N=20246 cells from the 'non-IBD patient C' biopsy, N=10416 cells from the

'non-inflamed patient D' biopsy, and N=30799 cells from the 'inflamed patient D' biopsy

were analysed. Black bar indicates mean value. **p<0.01 by one-way ANOVA with Tukey's

post-hoc correction. Related to Table **1** and Fig. **EV8-10**.

1568

EV Figures and Tables:

Fig. EV1. Optimisation pipeline for the immunohistochemical detection of the mouse

necroptotic pathway. Summary of the steps used to test the specificity and to optimise

the immunohistochemistry performance of monoclonal antibodies against mouse MLKL,

RIPK3, RIPK1 and Caspase-8. Step 1: Immunoblot signals for each antibody on spleen

homogenates from wild-type mice (WT) and *Mlkl*$^{-/-}$ or *Casp8*$^{-/-}$*Ripk3*$^{-/-}$ or *Casp8*$^{-/-}$*Ripk1*$^{-/-}$

Ripk3$^{-/-}$ mice (KO). Arrowheads indicate the full-length protein of interest. Representative

GAPDH immunoblots are shown as loading controls. Data are representative of n=1-2

immunoblots per antibody. Steps 2-4: Heatmaps depict the integrated signal-to-noise

values derived from immunohistochemical signals for each antibody on WT versus KO

spleen sections. Legend shows the heatmap color-to-value scale. Step 2 varied the

primary antibody concentration and the antigen retrieval pH. Step 3 varied the blocking

reagent and the antigen retrieval time. Step 4 varied the amplification technique,

amplification time, peroxidase treatment time and whether peroxidase treatment

preceded/superceded primary antibody incubation. For each antibody, the optimal

condition at Step 2 (white stars) was the starting point for Step 3. Similarly, the optimal

condition for Step 3 was the starting point for Step 4. Yellow stars indicate the final

automated immunohistochemistry protocol stipulated in File **EV1**. Data in Steps 2-4 are representative of n=1-5 experiments per antibody per condition.

Fig. EV2. Constitutive co-expression of necroptotic effectors is confined to fast-cycling cells within progenitors, immune and barrier populations. A Heatmap of cell ontologies from the *Tabula Muris* dataset (Tabula Muris *et al.*, 2018). Left-most column depicts the tissue origin of each cell ontology. Other columns indicate the percent of cells within each ontology that expressed *Top2a, Mki67*, *Casp8*, *Ripk1, Ripk3* or *Mlkl*. Legend shows the color-to-tissue and the color-to-frequency scales. Cell ontologies of interest are annotated. **B** Micrograph of MLKL immunosignals from the wild-type mouse spleen. The white pulp (WP), marginal zone (MZ) and red pulp (RP) are annotated. Scale bar is 50µm. Scatterplot shows relative expression levels of MLKL along the white pulp-to-red pulp axis. Red datapoints show immunosignal intensities and the overlaid dark blue line indicates the LOWESS best-fit along N=20 axes from n=1 mouse. Dashed line indicates the boundary between splenic white pulp and marginal zone. Data are representative of n>3 mice. **C-D** Spatial transcriptomic data from (Moor *et al.*, 2018)) and (Ben-Moshe *et al.*, 2019) showing the relative expression levels (arbitrary units; A.U.) of Caspase-8, RIPK1, RIPK3 or MLKL along the ileal crypt-to-villus axis (Panel C) or the hepatic central vein-to-portal vein axis (Panel D). **E-G** Spatial transcriptomic data on mouse spleen 12 days after *Plasmodium berghei*-infection. Panel E shows a Uniform Manifold Approximation and Projection (UMAP) of cell populations distinguished by unsupervised leiden clustering. Legend shows the color assigned to each population. Panel F shows the location of each cell cluster. Scale bar is 500µm. Panel G shows the normalised expression for each gene product. Expression values for *Casp8*, *Ripk1*, *Ripk3* and *Mlkl* were summated to provide an index of 'cluster pathway expression', which was averaged to provide an index of 'zone pathway expression'. Data are from n=1 mouse.

1612

Fig. EV3. Antibiotic-induced clusters of epithelial RIPK1/3 are not necrosomes.
Immunoblots for the indicated proteins in homogenates of the ileum and spleen of water-
versus antibiotic-treated mice or in mouse dermal fibroblasts (MDF) undergoing TNF-
induced necroptosis (TSI). Arrowheads indicate full-length proteins of interest. Coomassie
staining of total protein content was used as a loading control. Data are representative of
n=7 mice per tissue per group.

1619

Fig. EV4. RIPK3 is uniquely upregulated in splenic germinal centres. A Experimental
design. **B** Ki67, Caspase-8, RIPK1, RIPK3 and MLKL immunosignals from adjacent
sections of the naïve or NP-KLH-immunised mouse spleen. Arrowheads show a Ki67$^+$
germinal centre that co-stains for RIPK3, but not other members of the pathway.
Representative of n>3 mice per group. Scale bars in lower magnification micrographs are
500µm. Scale bars in insets are 100µm. Data representative of n>3 mice per group. **C-H**
Ripk3$^{-/-}$ or *Ripk3*$^{+/+}$ mice were immunised with NP-KLH and circulating NP-specific IgG$_1$
(Panel C), circulating NP-specific IgM (Panel D), splenic mature B cells (Panel E and F),
splenic NP-specific plasma cells (Panel G), circulating low affinity NP-specific IgG$_1$
antibody (Panel H), circulating high affinity NP-specific IgG$_1$ (Panel H) and the ratio
between circulating low-and-high affinity NP-specific antibody (Panel J) were measured at
the indicated day after immunisation. Each datapoint represents one mouse. ns = non-
significant by 2-sided t-test with Welch's correction.

1633

**Fig. EV5. Standardised quality control and optimisation of the immunohistochemical
detection of human necroptotic proteins.** Summary of the optimisation for
immunodetection with monoclonal antibodies against human MLKL, RIPK3, RIPK1,
Caspase-8 and Caspase-10 (C10). Step 1: Immunoblot signals for each antibody on

lysates from wild-type (WT) versus *MLKL*$^{-/-}$ or *RIPK1*$^{-/-}$ or *CASP8*$^{-/-}$*CASP10*$^{-/-}$*MLKL*$^{-/-}$ (KO)

HT29 cells after treatment with the indicated stimuli. Arrowheads indicate the full-length

protein of interest. Representative GAPDH immunoblots are shown as loading controls.

Data are representative of n=1-2 immunoblots per antibody. Steps 2-4: Heatmaps depict

the integrated signal-to-noise values derived from immunohistochemical signals for each

antibody on WT versus KO HT29 cells. Legend shows the heatmap colour-to-value scale.

Step 2 varied the primary antibody concentration and the antigen retrieval pH. Step 3

varied the blocking reagent and the antigen retrieval time. Step 4 varied the amplification

technique, amplification time, peroxidase treatment time and whether peroxidase

treatment preceded/superceded primary antibody incubation. For each antibody, the

optimal condition at Step 2 (white stars) was the starting point for Step 3. Similarly, the

optimal condition for Step 3 was the starting point for Step 4. Yellow stars indicate the final

automated immunohistochemistry protocol stipulated in File **EV1**. Data in Steps 2-4 are

representative of n=1-5 experiments per antibody per condition. The annotation 'see Fig.

EV1' indicates optimisations that were only done using mouse tissues because these

antibodies recognise both human and mouse orthologs.

1654

Fig. EV6. Automated immunohistochemistry to detect necroptotic signaling in

mouse cells. Immunosignals of cleaved Caspase-3, Caspase-8, RIPK1, RIPK3 and MLKL

in wild-type mouse dermal fibroblasts. Arrowheads indicate Caspase-8^{+} or RIPK1^{+} puncta

that are presumed to be necrosomes. Data are representative of n=1 for each protein and

treatment. Scale bars in lower magnification micrographs are 10μm. Scale bars in insets

are 1μm.

1661

1662 **Fig. EV7. Assessing the specificity of immunostaining using wild-type, knockout**

1663 **and human RIPK3 knock-in mice. A** Approach used to insert the human RIPK3 coding

sequence (CDS) into the mouse *Ripk3* locus. **B** Immunoblot of spleen homogenates from *Ripk3*$^{+/+}$, *Ripk3*$^{-/-}$, human RIPK3 (*RIPK3*$^{KI/KI}$) or hemizygous human RIPK3 (*Ripk3*$^{+}$*RIPK3*^{KI}) mice. GAPDH immunoblots are shown as loading controls. Each lane represents a different mouse. **C** Immunosignals produced by the anti-human RIPK3 (clone 37A7F) or anti-mouse RIPK3 (clone 8G7) antibodies on spleen sections from *RIPK3*$^{KI/KI}$*, Ripk3*$^{+/+}$ or *Casp8*$^{-/-}$*Ripk1*$^{-/-}$*Ripk3*$^{-/-}$. Data are representative of n≥3 for each target and tissue. Scale bars are 500µm.

Fig. EV8. Another instance of elevated intestinal apoptosis in a patient with ulcerative colitis. Immunoblot of lysates from HT29 cells (red annotations) and intestinal biopsies from patients A-H (blue annotations). The fifth lane of each gel contained lysates from TSI-treated *RIPK3*$^{-/-}$ or TSI-treated *MLKL*$^{-/-}$ cells (see source data for details). Patient J was a non-IBD control. Patient I had ulcerative colitis (UC). The endoscopic grading of the biopsy site as 'non-inflamed', 'marginally inflamed' or 'inflamed' is stipulated. Closed arrowheads indicate full-length form of proteins. Asterisks indicate active, phosphorylated forms of RIPK3 (pRIPK3) and MLKL (pMLKL). Open arrowheads indicate active, cleaved forms of Caspase-10 and Caspase-3. GAPDH was used as a loading control.

Fig. EV9. Atlas of necroptotic pathway expression in human intestinal crypts. Immunohistochemistry for cleaved Caspase-3, Caspase-8 (clone B.925.8), RIPK1, RIPK3 and MLKL (clone EPR171514) on intestinal biopsies from the stipulated patients. Four representative micrographs per biopsy are shown (i-v). Open arrowheads indicate instances of epithelial apoptosis. Closed arrowheads indicate instances of epithelial Caspase-8 clustering. Scale bars are 10µm. The location for each micrograph within the biopsy is indicated in Fig. EV10.

Fig. EV10. Atlas of necroptotic pathway expression in a cohort of human intestinal biopsies. Immunohistochemistry for cleaved Caspase-3, Caspase-8 (clone B.925.8), RIPK1, RIPK3 and MLKL (clone EPR171514) on intestinal biopsies from the stipulated patients. The insets show the location of each micrographs in Fig. EV9 (annotated as i-v per biopsy). Scale bars are 100μm.

Table 1. Patient and clinical data related to **Fig. 6**.

File EV1. The automated immunohistochemical staining protocols used in this study will be provided upon final publication.

A Quantitation of IHC specificity

i)

ii)

iii) integration of S/N
1.5E+06

B IHC signal-to-noise values

	C8 D35G2	C8 1G12	C8 3B10	R1 D94C12	R3 8G7	R3 1H12	MLKL 5A6
ileum							
colon							
liver							
spleen							
kidney							
heart							
brain							

low amount of specific signal — high amount of specific signal

C protein abundance

	C8	R1	R3	MLKL
sm. int.				
colon				
liver				
spleen				
kidney				
heart				
brain				

below detection — max

D

	Caspase-8	RIPK1	RIPK3	MLKL
ileum				
colon				
liver				
spleen				
kidney				
heart				
brain				