Targeted Deletion of All Known Thyroid Hormone Receptors Causes Maturation Retardation and Early-onset Degeneration of Cochlear Outer Sulcus

XianHua Ma[a, #], Fei Jiang[b, #], Chunchun Wei[a], Shuang Han[c], Yuqing Zhang[b], Lianhua Sun[b], Jiaxi Qu[d], Hao Ying[e], Yuxia Chen[a], Jie Tang[d], David Z. He[f], Weiping J. Zhang[a,c,*], Zhifang Xie[b,*]

[a]Department of Pathophysiology, Naval Medical University, Shanghai 200433, China

[b]Ministry of Education-Shanghai Key Laboratory of Children's Environmental Health, Institute of Early Life Health, Xinhua Hospital Affiliated to Shanghai Jiao Tong University School of Medicine, Shanghai 200092, China

[c]NHC Key Laboratory of Hormones and Development, Chu Hsien-I Memorial Hospital & Tianjin Institute of Endocrinology, Tianjin Medical University, Tianjin 300134, China

[d]Department of Physiology, Southern Medical University, Guangzhou, Guangdong 510515, China

[e]CAS Key Laboratory of Nutrition, Metabolism and Food Safety, Shanghai Institute of Nutrition and Health, University of Chinese Academy of Sciences, Chinese Academy of Sciences, and Shanghai Jiao Tong University Affiliated Sixth People's Hospital, Shanghai 200031, China.

[f]Department of Biomedical Sciences, Creighton University School of Medicine, Omaha, NE 68178, USA

[#]They contributed equally to the work.

Corresponding authors: xiezhifang@xinhuamed.com.cn, zfxie70@hotmail.com (Z.X., lead contact); Weiping J. Zhang, wzhang@smmu.edu.cn (W.J.Z)

Abstract

Thyroid hormone (TH) and its receptors (TRs) are crucial for cochlear late-stage

development and endocochlear potential (EP) maintenance. However, the mechanism

underlying EP reduction in the absence of TH or TRs remains elusive. Cochlear outer sulcus

root cells undergo significant morphological changes during late-stage cochlear development

and are believed to play a role in maintaining endolymph homeostasis and EP. Yet, it is unknown

whether TH and/or TRs are necessary for root cell differentiation and function. Here, we

elucidate the essential role of TH or TRs in postnatal root cell development and survival in mice.

TH deficiency significantly delayed root cell differentiation. Otocyst-selective deletion of both

Thra and *Thrb*, but not *Thrb* alone, leads to a similar impairment, accompanied by early

degeneration of root cells, with the stria vascularis remaining unaffected. Furthermore, a 22%

reduction in mean EP magnitudes was observed in conditional *TRs* double knockout mice at 4

months of age, less pronounced than in global *TRs* knockout mice. Transcriptome analysis

reveals that TH deficiency downregulates a significant portion of root cell-enriched genes.

These findings underscore the redundant roles of TRα and TRβ in promoting the late-stage

differentiation and survival of root cells. Additionally, they suggest that the expression of TRs in

cochlear epithelium is crucial for maintaining an optimal EP magnitude, while TRs expressed

in areas outside cochlear epithelium, particularly in spiral ligament fibrocytes, may also

significantly contribute to EP maintenance. This study advances our understanding of TH in

cochlear outer sulcus development and EP maintenance.

Keywords: cochlea, thyroid hormone receptor, development, root cell, endocochlear potential

Introduction

Thyroid hormone (TH) is crucial for the normal development of the cochlea and hearing (Ng

et al., 2013). Both early-onset congenital hypothyroidism and environmentally induced iodine

deficiency can lead to hearing loss in both humans and rodents (Christ et al., 2004; Deol, 1973a;

Mustapha et al., 2009). TH primarily exerts its effects by binding to two distinct thyroid hormone

receptors (TRs), namely TRα and TRβ, which are ligand-dependent transcription factors

encoded by *Thra* and *Thrb*, respectively (Forrest et al., 1996; Rusch et al., 2001; Winter et al.,

2009). TRα is individually dispensable for hearing(Rüsch et al., 1998), while the absence of

TRβ alone leads to deafness accompanied by malformation of the tectorial membrane (TM)

(Forrest et al., 1996; Winter et al., 2009). Global deletion of both *Thra* and *Thrb* in mice

exacerbates cochlear defects, including more severe distortion of the TM, significant retardation

in the regression of the great epithelial ridge (GER, also known as Kolliker's organ), and

remodeling of the organ of Corti, as well as a reduction in endocochlear potential (EP) (Rusch

et al., 2001). The mechanism underlying the reduction in EP upon deletion of TRs remains

unclear, as no significant alterations in the stria vascularis (SV) have been reported in these

mice, suggesting that defects beyond the SV may contribute to the reduction in EP.

The cochlear outer sulcus epithelial cells, also known as root cells, are poorly understood

cell types that are considered to be important for the maintenance of endolymph homeostasis

and EP (Spicer and Schulte, 1996; Cazals et al., 2015; Galic and Giebel, 1989; Jagger et al.,

2010; Jagger and Forge, 2013). Positioned on the cochlear lateral wall between the outer

border Claudius cells and the spiral prominence, the outer sulcus undergoes significant

morphological changes during the postnatal developmental window when the cochlea is most

sensitive to TH (Deol, 1973b; Karolyi et al., 2007; Uziel et al., 1985). At birth, the outer sulcus

consists of a simple layer of cuboidal-shaped epithelial cells lining the scala media. Shortly after

birth, these cells begin lateral and superior infiltration into the spiral ligament beneath the spiral

prominence, forming extensive tree-like branched processes, hence termed "root cells."

Concurrently, Claudius cells enlarge significantly, meeting the spiral prominence epithelium and

covering the underlying root cell bodies, thereby preventing root cells from direct contact with

endolymph in basal turns. However, in the apical turn, root cells directly face the endolymph

and exhibit fewer processes (Duvall III, 1969; Galic and Giebel, 1989; Jagger and Forge, 2013;

Lim and Anniko, 1985; Shodo et al., 2017; Spicer and Schulte, 1996).

There is limited understanding of the mechanisms that regulate root cell differentiation. Our

recent study has highlighted the importance of the transcription factor ZBTB20 in cochlear late-

stage development, particularly in root cell differentiation. Selective deletion of *Zbtb20* in the

cochlear epithelium results in deafness reduced EP, and cochlear malformations, such as a

malformed tectorial membrane (TM) and organ of Corti, along with impaired maturation of the

outer border and outer sulcus region (Xie et al., 2023). Intriguingly, *Thrb* expression has been

detected in both developing and adult root cells, as demonstrated by a lacz gene reporter (Ng

et al., 2015) or single-cell RNA sequencing (Gu et al., 2020), while *Thra* mRNA is widely

expressed in the lateral wall, including root cells (Bradley et al., 1994; Jean et al., 2023).

However, it has long been neglected whether TH and the expression of TRs in the cochlea play

essential roles in the remodeling and maturation of the outer sulcus.

Here we find that TRs expression in otocyst-derived cochlear epithelium is necessary for

postnatal root cell development and survival. Root cell development is significantly delayed in

mice with antithyroid drug-induced developmental hypothyroidism or those with otocyst-

selective deletion of both *Thra* and *Thrb*, while no apparent root cell defects are observed upon

individual deletion of *Thrb*. Moreover, the absence of both TRα and TRβ results in early-onset

degeneration of root cells and a compromised EP magnitude.

Results

Hypothyroidism causes severe retardation in the outer sulcus development

To investigate the role of thyroid hormone (TH) in outer sulcus development, we induced

developmental hypothyroidism through antithyroid treatment, following a previously established

protocol(Sawant et al., 2015). Antithyroid treatment commenced on gestation day 17 and

extended through lactation until postnatal day (P) 21. This duration covers a critical

developmental window during which the cochlea exhibits heightened sensitivity to TH

deprivation (Deol, 1973b; Uziel et al., 1985). Histological analysis was conducted on cochleae

harvested from euthyroid or hypothyroid mice at P21, a stage when the cochlea attains both

morphological and functional maturity in normal mice (Lim and Anniko, 1985; Walters and Zuo,

2013).

To better preserve cochlear histological features, we employed a resin-embedding

technique. In euthyroid cochleae, the sensory epithelium exhibited typical mature morphology,

characterized by an upright organ of Corti, a well-developed inner sulcus, and a normal tectorial

membrane (TM). Within the outer sulcus region, enlarged Claudius cells covered the underlying

root cells, displaying numerous bright branched processes extending deep into the spiral

ligament and upwards behind the spiral prominence (Fig. 1A). In line with previous

findings(Deol, 1973b), hypothyroid cochlear epithelium exhibited marked retardation in sensory

epithelium remodeling, evidenced by the abnormal persistence of the greater epithelial ridge

(GER), closure of the tunnel of Corti, and severe distortion of the TM. Notably, the outer sulcus

displayed an immature appearance, characterized by a simple layer of cells with few short

processes bordering the scala media, along with flattened Claudius cells (Fig. 1B). This

observation suggests severe retardation in both root cell and Claudius cell differentiation.

Our previous work, along with studies by others, has demonstrated that root cell

cytoplasm and their processes exhibit enrichment of acetyl-α-tubulin (Jagger and Forge, 2013;

Liu et al., 2018; Xie et al., 2023). To gain further insight into root cell morphology, we conducted

immunofluorescence staining for acetyl-α-tubulin. Consistent with the histological findings,

euthyroid cochleae exhibited robust and extensive acetyl-α-tubulin signals extending deep into

the spiral ligament (Fig. 1C). In contrast, hypothyroid cochleae displayed limited signals,

primarily distributed around the luminal surface of the spiral ligament, indicative of root cell

differentiation defects (Fig. 1D). These findings underscore the essential role of TH signaling in

the timely development of the outer sulcus.

Otocyst-selective deletion of all known *TRs* causes deafness and a compromised EP

As TH effects are mainly mediated by TRα and TRβ(Ng et al., 2013), both of which are

expressed in the developing outer sulcus(Bradley et al., 1994; Gu et al., 2020; Ng et al., 2015),

we therefore asked if the two receptors are essential for root cell differentiation. To address this

question, we generated mice with either individual deletion of the *Thrb* gene or combined

deletion of the *Thra* and *Thrb* genes selectively in the otic vesicle using the Cre-loxP system.

These mice are hereafter referred to as OV-TRβKO and OV-TRα/βKO, respectively. Individual

Thra-deleted mice were not included in this study, as *Thra*-null mice have been reported to

exhibit normal hearing thresholds (Rüsch et al., 1998). We crossed *Thrb-floxed* (*Thrb^flox*)

mice(Yan et al., 2022) either individually or in combination with *Thra ^flox* mice(Liu et al., 2022) to

Foxg1-Cre transgenic mice, a strain capable of mediating recombination selectively in the otic

vesicle by embryonic day 8.5(Hebert and McConnell, 2000). We verified the tissue specificity

of *Cre*-mediated recombination by mating *Foxg1-Cre* mice with *Rosa26*-lacZ reporter mice.

Consistently, *Cre*-recombinase activity, as indicated by X-gal staining, was confined to otocyst-

derived cochlear epithelium at P5. This included the greater epithelial ridge (GER), lesser

epithelial ridge (LER), outer sulcus epithelium (root cells), stria vascularis (SV) marginal cells,

and spiral ganglion. As anticipated, no signal was detected in cochlear fibrocytes derived from

the periotic mesenchyme surrounding the otocyst (Song et al., 2011). No background signal

was observed in cre-negative control mice (Fig. 2A). The efficiency of deletion was further

evaluated through qRT-PCR analysis, which revealed that the expression levels of *Thrb* mRNA

were significantly downregulated in OV-TRβKO cochleae at P10, while both *Thrb* and *Thra*

mRNA were significantly downregulated in OV-TRα/βKO cochleae (Fig. 2B).

To evaluate the auditory function of the mutant mice, we performed ABR (auditory

brainstem response) tests. Both OV-TRβKO and OV-TRα/βKO mice did not show typical ABR

responses to either click (Fig. 2C, D) or frequency-specific tone burst stimuli at 90 dB (the

highest intensity tested) around 2 months (Fig. 2D), in sharp contrast to normal thresholds seen

in wild-type (WT) control mice. These data demonstrate that cochlear expression of TRβ or all

known TRs is essential for hearing, in agreement with previous reports(Forrest et al., 1996;

Rusch et al., 2001).

The global deletion of all known *TRs* has previously been associated with a roughly 50%

reduction in the EP magnitude in adult mice (Rusch et al., 2001). To investigate if the expression

of TRα and TRβ within the cochlear epithelium is essential for EP maintenance, we measured

EP in OV-TRα/βKO mice. At around 1.5 months of age, OV-TRα/βKO mice exhibited a slight

but statistically significant decrease in EP magnitude compared to WT controls (WT = 107.2 ±

8.0 mV vs OV-TRα/βKO = 92.8 ± 9.0 mV, p = 0.01). By 4 months of age, the mean EP magnitude

in OV-TRα/βKO mice further declined to 85.1 ± 15.6 mV, while no significant changes were

observed in the WT group (108.8 ± 4.6 mV) (Fig. 2E). These findings underscore the role of

TRs expressed within the cochlear epithelium in maintaining optimal EP magnitude. Additionally,

they suggest that TRs expression in areas outside of the cochlear epithelium may also

contribute to EP maintenance.

Otocyst-selective deletion of all *TRs* retards root cell development

We next investigated cochlear morphological changes in *TRs* knockout mice, with an

emphasis on the development of the outer sulcus region. At P10, 2 days before the onset of

hearing in mice, root cells with branched processes had already invaded deep into the spiral

ligaments in WT pups (Fig. 3A). Comparable root cell development was also observed in OV-

TRβKO counterparts (Fig. 3B). The outer sulcus in OV-TRα/βKO mice, however, was largely

composed of a single cuboidal epithelial layer bordering the endolymphatic space (Fig. 3C),

suggesting a delay in the root cell differentiation. Consistent with the histological observations,

OV-TRα/βKO showed fewer acetyl-α-tubulin positive cell processes within the spiral ligament

compared with those in WT and OV-TRβKO counterparts (Fig. 3D-F). To further demonstrate

root cell differentiation, we employed in situ hybridization (ISH) to investigate expression

patterns of *Slc26a4* (also known as *Pendrin*) and *Epyc* (*Epiphycan*) mRNA. These genes are

known to be abundantly and selectively expressed by root cells(Gu et al., 2020; Royaux et al.,

2003; Xie et al., 2023).) Strong signals of *Slc26a4* (Fig. 3G-I) or *Epyc* (Supplementary Fig.

S1A-C) mRNA could be detected deep inside the spiral ligament in both WT and OV-TRβKO

cochleae, but in OV-TRα/βKO cochleae, expression domains of both molecules were adjacent

to the luminal surface of the spiral ligament, consistent with a delay in the root cell development. These data suggest that root cell development is normally initiated in OV-TRβKO but disrupted in OV-TRα/βKO pups.

By P21, mature morphological features were comparable between OV-TRβKO and wild-type (WT) outer sulci, with elaborate root cell branches extending deep into the inferior region of the spiral ligament and upward behind the spiral prominence in the basal turn (Fig. 3J and K). Consistent with previous descriptions (Duvall III, 1969; Kimura, 1984), the surface of root cell bodies was covered by greatly enlarged Claudius cells and spiral prominence epithelium, no longer directly exposed to the endolymph at this stage. In contrast, the double knockout mice (OV-TRα/βKO) exhibited only initial signs of root cell differentiation at P21. Root cells with a few processes were mainly located along the scala media, separating the flattened Claudius cell and the spiral prominence epithelium, indicating a severe retardation in the differentiation of root cells and Claudius cells (Fig. 3L). Additionally, some cells with pale-staining cytoplasm and nucleus were occasionally detected in the OV-TRα/βKO outer sulcus at this age, suggesting degenerative changes (Fig. 3L). This observation was further supported by reduced acetyl-α-tubulin-positive root cell processes in the OV-TRα/βKO outer sulcus compared to WT or OV-TRβKO counterparts (Supplementary Fig. S1D-F).

By 1.5 months, the development of root cells and Claudius cells in OV-TRα/βKO mice had progressed to some extent but still lagged behind the developmental status observed in WT controls (Supplementary Fig. S2A-B, A' and B'). Additionally, degenerative changes were noted in root cells and cochlear outer border cells (Boettcher's cells) in some OV-TRα/βKO samples (Supplementary Fig. S2C and C').

Taken together, these data demonstrate that root cell development is largely unaffected in

OV-TRβKO mice but significantly delayed in OV-TRα/βKO mice. This suggests that while TRβ

may not be individually indispensable, TRα and TRβ collectively play redundant roles in root

cell development.

Otocyst-selective deletion of all *TRs* causes early-onset root cell degeneration

Although the degeneration of cochlear hair cells and supporting cells has been reported

previously in developmental hypothyroidism, and in mice with global deletion of TH transporters

(Uziel et al., 1983; Sharlin et al., 2018; Mustapha et al., 2009), it remains unclear whether the

survival of root cells depends on TH signaling. As described above, sporadic degenerative cells

were occasionally detected in OV-TRα/βKO outer sulcus at P21 (Fig. 3L), or ~1.5 months of

age (Supplementary Fig. S2C and C'). By 2 months, WT control or OV-TRβKO outer sulcus

showed no obvious degenerative changes, whereas OV-TRα/βKO exhibited reduced root cells

with pale-staining processes and nuclei in both basal and apical turns (Fig. 4A-C, A'-C', and

Supplementary Fig. S3), suggesting progressive degenerations. Concomitant with the root cell

degeneration, variable degenerations of the organ of Corti were observed in OV-TRα/βKO

cochlea (Fig. 4C, Supplementary Fig. S3C), resembling those described previously in

developmental hypothyroidism, or in mice with global deletion of TH transporters (Mustapha et

al., 2009; Sharlin et al., 2018; Uziel et al., 1983).

To assess root cell viability, we quantified the number of root cell nuclei at 2 months of

age. Previous studies, including ours, have shown that Sox10 is localized in the nucleus of

cochlear non-sensory cells, including outer border cells, root cells, and spiral prominence

epithelium, but is excluded from fibrocytes in the lateral wall (Hao et al., 2014; Xie et al., 2023).

Therefore, by performing double immunostaining for Sox10 and acetyl-α-tubulin (intensely

expressed in the root cell cytosol), root cells can be easily identified as double-positive cells in

the spiral ligament (Fig. 4D). Consistent with the histological observations, the number of root

cell's nucleus was significantly decreased in OV-TRα/βKO but not in OV-TRβKO, compared to

that in WT (Fig. 4D-E). This supports the occurrence of root cell degeneration in the absence

of both TRs.

Furthermore, we conducted a TUNEL analysis to detect apoptotic cells in the outer sulcus

region. We observed sporadic yet limited TUNEL-positive cells in OV-TRα/βKO mice, but not

in WT controls, at 2 months of age (Fig. 4F). Quantitative analysis revealed a statistically

significant increase in TUNEL-positive cells in OV-TRα/βKO mice compared to the WT control

group (Fig. 4G), suggesting apoptosis occurs at a low rate upon TRs loss. These data

collectively indicate that TRα and TRβ function redundantly for root cell survival.

Stria vascularis remains intact in OV-TRα/βKO mice

Given that OV-TRα/βKO mice exhibited a reduced EP compared to control mice (refer to

Fig. 2E), we then examined the structural and functional integrity of SV, which plays an essential

role in generating EP and maintaining K^+ homeostasis in the endolymph(Hibino et al., 2010;

Kim et al., 2013; Wangemann, 2006). At P21, no obvious histological abnormalities were

observed in either OV-TRβKO or OV-TRα/βKO SV (Fig. 5A). The area of SV in both mouse

lines was comparable to that in WT mice (Fig. 5B). Consistently, immunostaining analysis

revealed normal expression patterns of EP-generating ion-transport molecules, including

KCNJ10 (Kir4.1, a marker for intermediate cells)(Ando and Takeuchi, 1999; Marcus et al., 2002),

Na^+/K^+-ATPase α(Kerr et al., 1982), NKCC1(Dixon et al., 1999) and KCNQ1 (Kv7.1)(Casimiro

et al., 2001; Sakagami et al., 1991) (markers for marginal cells) (Fig. 5C). These data suggest

that SV remains intact upon all *TRs* deleted in cochlear epithelium. Indeed, potential primary

defects can be excluded from SV intermediate cells and basal cells in either OV-TRβKO or OV-

TRα/βKO mice, as these cell types do not originate from the otic epithelium (Steel and Barkway,

1989) and thus should display no Cre recombination activity (refer to Fig. 2A).

As gap junctions are also essential for EP generation and hearing(Cohen-Salmon et al.,

2002; Sun et al., 2005). We then analyzed the expression of Connexin26 (Cx26) and Cx30

proteins, which are the dominant gap-junction channel-forming proteins expressed in the

cochlear epithelium(Ahmad et al., 2003). Comparable expression patterns of Cx26 and Cx30

were observed between control and OV-TRα/βKO cochleae at P21 (Supplementary Fig. S4).

This suggests the gap junction system remains intact despite TRs deficiency.

TH deficiency leads to the downregulation of a considerable number of root cell-

enriched genes

To unravel the molecular mechanisms underlying outer sulcus development regulated by

TH, we employed a microdissection technique to isolate outer sulcus regions from both

euthyroid and hypothyroid mice at P10. Subsequently, we conducted a whole transcriptome

analysis using RNA sequencing (RNA-seq). Each experimental group comprised five biological

replicates. We identified 863 differentially expressed genes (DEGs) with a fold change of ≥ 2

and a Q-value of ≤ 0.05 (Supplementary Fig. S5, Supplementary Dataset S1). Genes with

Fragments per kilobase per million mapped reads (FPKM) < 10 were excluded from the analysis.

Given that the microdissected region primarily consists of outer sulcus epithelial cells

and fibrocytes, and our research focus is on root cell development, we conducted an analysis

to identify DEGs preferentially expressed by root cells. This involved intersecting our dataset

with a previously reported list of root cell-enriched genes (Jean et al., 2023). We identified a

total of 178 root cell-enriched DEGs, consisting of 11 upregulated and 167 downregulated

genes (Fig. 6A). Notably, several known marker genes for root cells, such as *Slc26a4* (Royaux

et al., 2003), *Lgr5*, *Epyc* (Gu et al., 2020; Xie et al., 2023), and *P2rx2* (Lee et al., 2001), were

among the downregulated DEGs identified, consistent with a delayed root cell differentiation in

hypothyroid mice. Interestingly, 11 of the downregulated genes, including *Slc26a4*, *Tmprss3*,

Gas2, *S100a1*, *Qsox1*, *Tmprss9*, *Prss36*, and *P2rx2*, have been associated with hearing loss

in humans and/or mice when mutated (Groza et al., 2023) (https://hereditaryhearingloss.org;

https://www.mousephenotype.org) (Fig. 6B).

It is well established that the biological activities of TH are mainly mediated by TRs, which

bind to TH response elements (TREs) to regulate target gene expression. The most frequently

characterized TRE motif is a direct hexameric repeat of 5'(A/G)GGT(C/A)A3' with 4 base spacer

(called DR4 elements) (Chatonnet et al., 2013; Umesono et al., 1991). We, therefore, utilized

the RcisTarget tool (https://bioconductor.org) to search for DEGs containing proximal TRE

motifs to identify potential direct target genes of TH. We identified 21 potential direct target

genes of TRs, with all but one showing downregulation in the hypothyroid outer sulcus (see Fig.

6C). Notably, among these genes, *Thrsp* has been previously established as a direct target

gene of TRs in the liver (Zilz et al., 1990). Moreover, a significant number of these genes have

been found to harbor endogenous TR binding sites in their proximal regions in the liver or

nervous system, including *Thrsp, Usp2* (Grøntved et al., 2015), *Stat5a* (Chatonnet et al., 2013),

Nr1d1, *Tbc1d10a*, *Qsox1*, *Zhx2*, *Atp1b1*, *Lbh*, *Tns1*, and *Plekha6* (Mendoza et al., 2021),

supporting a potential direct regulation by TH. Further investigation is needed to determine if

this also applies to the cochlea.

Fifteen DEGs were chosen for additional validation through qRT-PCR (Fig. 6D) or ISH (Fig.

6E). Selection criteria included significant fold changes or potential relevance to auditory

function. Notably, the expression of *Prss36* in the cochlea has not been previously

characterized. Together, these data underscore the significance of TH signaling in regulating

the gene expression profile associated with root cell differentiation.

Discussion

This study highlights the pivotal role of TH and TRs in postnatal root cell development and

survival. The absence of both TRα and TRβ, but not TRβ alone, in cochlear epithelium led to

significant retardation in root cell differentiation and early-onset root cell degeneration. These

findings suggest that TRα and TRβ function redundantly in promoting late-stage root cell

differentiation and maintaining cellular homeostasis.

The critical period when TH is essential for cochlear development and normal hearing

covers from late embryonic stages to around the second postnatal week in rodents(Deol, 1973b;

Uziel et al., 1985; Uziel, 1986). During this developmental window, the cochlear sensory

epithelium undergoes dramatic morphological changes including regression of GER leading to

the formation of the inner sulcus, and opening of the tunnel of Corti (Deol, 1973b; Roth and

Bruns, 1992). Concomitantly, the outer sulcus epithelium transforms from a simple layer of

cuboidal-shaped cells lining the scala media into root cells with large, branched processes

residing deep in the spiral ligament(Lim and Anniko, 1985; Roth and Bruns, 1992; Xie et al.,

2023). This study uncovers that TH or TRs are necessary for the outer sulcus remodeling, in

addition to previously reported GER regression and the organ of Corti maturation(Rusch et al.,

2001). Although both TRα and TRβ are expressed by the developing outer sulcus(Bradley et

al., 1994; Gu et al., 2020; Jean et al., 2023; Ng et al., 2015), severe retardation in root cell

differentiation was observed only in the absence of both receptors indicating that TRα and TRβ

compensate functionally for each other in controlling root cell development. Our findings extend

those data obtained from the mouse model with global deletion of TRs, which suggest

synergistic effects of TRα and TRβ in controlling the remodeling of the organ of Corti and the

correct formation of the tectorial membrane (Rusch et al., 2001).

Another intriguing finding of this study is the early-onset progressive degeneration of root

cells observed only in OV-TRα/βKO but not OV-TRβKO cochlea, suggesting TRα and TRβ play

a combined role in root cell survival. Interestingly, degeneration of root cells and the organ of

Corti initiated concomitantly in juvenile OV-TRα/βKO cochlea, indicating potential coordinated

mechanisms underlying these degenerations. As root cells, with their extensive processes, are

closely associated with capillaries and fibrocytes within the spiral ligament, it has been

proposed that they may provide a cellular route for transporting nutrients, blood-borne

hormones (such as T3), metabolites, and ions between the vasculature in the spiral ligament

into the inferior cochlear sensory epithelium(Duvall III, 1969; Jagger and Forge, 2013; Shodo

et al., 2017), which lacks direct blood flow. This hypothesis is supported by an epithelial gap

junction network formed by the syncytium of supporting cells in the organ of Corti and extending

through the Claudius' cells and root cells(Jagger and Forge, 2015, 2006). Hence, it's

reasonable to assume that the proper timely development of root cells may contribute to

maintaining the health of the cochlear sensory epithelium. The early-onset degeneration of root

cells might compromise the supply of nutrients and hormones or disrupt ion and metabolite

homeostasis in the organ of Corti, thereby promoting its degeneration. Alternatively, the health

of root cells might be influenced by the degeneration of the organ of Corti. Severe early-onset

degenerations of cochlear hair cells have been reported previously in congenital

hypothyroidism or mice with global deletion of thyroid hormone transporters (Mustapha et al.,

2009; Sharlin et al., 2018), although the status of root cells in these animal models has not

been described. It would be valuable to investigate any alterations in root cell structure or

function in these mice.

Several lines of evidence including our recently published report suggest that root cells

may play a significant role in the regulation of endolymph homeostasis and EP (Spicer and

Schulte, 1996; Jagger et al., 2010; Cazals et al., 2015; Xie et al., 2023). For instance, the

absence of the KCNK5 channel, primarily located in the outer sulcus, leads to severe hearing

loss accompanied by a more than 60% decrease in EP magnitudes (Cazals et al., 2015).

Furthermore, our recent study revealed that the disruption of outer sulcus development due to

Zbtb20 deletion results in a reduction of over 50% in EP values (Xie et al., 2023). In this study,

however, OV-TRα/βKO mice showed only a reduction of approximately 13% and 22% in mean

EP values at 1.5 months and 4 months, respectively. This discrepancy could be attributed to

varying degrees of disruption in root cell development and function among different animal

models. Unlike *Zbtb20* knockout mice, where there is a permanent disruption in outer sulcus

differentiation, the outer sulcus development in OV-TRα/βKO mice had progressed to some

extent by approximately 1.5 months. Subsequently, root cells experienced gradual

degeneration, but approximately 50% of them persisted in OV-TRα/βKO mice. The mild EP

reduction in OV-TRα/βKO mice may indicate a robust compensatory ability of root cells in

regulating endolymph homeostasis and EP. Further investigation through targeted ablation of

root cells could provide insight into the extent of their contribution to EP magnitude and clarify

their role in regulating cochlear function.

It is noteworthy that the reduction in EP observed in OV-TRα/βKO mice is less pronounced

than that documented in global *Thra* and *Thrb* double knockout mice (Rusch et al., 2001). This

discrepancy indicates that TRs expression outside of the cochlear epithelium, particularly in

spiral ligament fibrocytes, may also contribute to EP maintenance. This hypothesis is supported

by the postnatal expression of both *Thra* and *Thrb* mRNA by fibrocytes (Jean et al., 2023),

which are known to play a role in EP maintenance (Adachi et al., 2013; Furness, 2019). Our

initial observations indicate a disruption in fibrocyte development in hypothyroid mice. However,

in OV-TRα/βKO mice, primary fibrocyte defects can be ruled out as Cre recombination activity

was confined in the otocyst-derived cochlear epithelium, as shown in Fig. 2A.

Although SV defects have been observed in cases of secondary hypothyroidism caused by

pituitary transcription factor 1 (Pit1) mutation (Mustapha et al., 2009), there have been no

reported SV alterations in cases of deficiency in TH transporters　(Sharlin et al., 2018), or in

global *TRs* double knockout mice(Rusch et al., 2001), both of which exhibit a significant

reduction in EP. Consistently, our findings in this study show normal SV upon deletion of both

Thra and *Thrb* in cochlear epithelium. Furthermore, previous studies utilizing a sensitive *LacZ*

gene reporter system have shown that *Thrb* mRNA is not expressed in the developing SV (Ng

et al., 2015). Therefore, the SV is unlike a primary defect site responsible for a reduced EP,

which occurred only in mice lacking all TRs(Rusch et al., 2001).

In summary, this study highlights the redundant role of TH receptors (TRα and TRβ) in

promoting the late-stage differentiation and survival of root cells, likely through the activation of

a significant portion of genes enriched in root cells. These findings suggest that the expression

of TRs in the cochlear epithelium is essential for maintaining an optimal EP magnitude and that

TRs expressed in areas outside of the cochlear epithelium, such as spiral ligament fibrocytes,

may also play a significant role in EP maintenance. These findings advance our understanding

of how TH regulates cochlear remodeling and maturation and provide new insights into the

mechanisms governing the development and function of cochlear outer sulcus regions.

Materials and Methods

Animal models

All animal maintenance and experimental procedures were conducted in compliance with

the guidelines issued by the Animal Ethics Committee of Naval Medical University. Protocols

(XIEC-NSFC-2022-033) were approved by the Ethics Committee of Xinhua Hospital affiliated

with Shanghai Jiaotong University School of Medicine. *Thra* flox(Liu et al., 2022), *Thrb*flox (Yan et

al., 2022), or ROSA26-*LacZ* reporter (Soriano, 1999)mice have been described previously. The

OV-TRβKO mice were generated by cross-breeding of *Thrb* flox mice with *Foxg1-Cre* mice (Jax,

Stock No. 004337)(Hebert and McConnell, 2000). The OV-TRα/βKO mice were then generated

by cross-breeding of heterozygous OV-TRβKO mice with *Thra* flox mice. All mice were

maintained on a C57BL/6J background. Floxed/*Cre*-negative littermate mice served as wild-

type controls. PCR primers for genotyping are listed in Supplementary Table S1. To induce

hypothyroidism, timed-pregnant C57BL/6J mice were treated with antithyroid agents (0.05%

methimazole, 1.0% sodium perchlorate monohydrate, and 5.0% sucrose in drinking water)

(Sawant et al., 2015) or vehicle from embryonic day 17. Treatment continued during lactation

and ceased on the designated day. Data from both genders were pooled together because no

gender differences in the phenotypes were observed.

Histological analysis

Mice were transcardially perfused with 4% paraformaldehyde (pH 7.4) to fix the tissues.

Cochleae were then dissected and perilymphatically perfused with 2.5% glutaraldehyde,

followed by overnight fixation. Post-fixation was performed with 1% osmium tetroxide.

Subsequently, the cochleae were decalcified, dehydrated, and embedded in Spurr's low-

viscosity resin. Mid-modiolar sections (1 μm) were stained with 1.25% (w/v) toluidine blue.

For quantitative analysis, images of the upper basal turns from three non-consecutive

sections were captured at 200x magnification. The stria vascularis (SV) areas were measured

using the Image-Pro Plus 6.0 program. Each analysis included a single cochlea from each

animal. The number of animals analyzed at a given age was indicated in figure legends.

Immunohistochemistry and X-Gal staining

Immunostaining was conducted following previously established protocols(Xie et al., 2023).

X-gal staining was performed according to published methods (Xie et al., 2008). Details

regarding primary antibodies are provided in Supplementary Table S2. To quantify root cells,

non-consecutive mid-modiolar sections per cochlea underwent double immunostaining for

Sox10 and acetyl-α-tubulin. The nuclei of double-positive cells within the outer sulcus region

were manually counted from at least 5 sections for each animal, with the genotype blinded

during counting. A single ear was analyzed for each animal.

Auditory brainstem response (ABR) test

Auditory brainstem responses (ABRs) were recorded using a TDT RZ6/BioSigRZ system

(Tucker-Davis Technologies Inc., USA), following established procedures (Xie et al., 2023).

Briefly, anesthetized mice were placed on a heated mat within a sound-proof chamber. Click or

pure tone stimuli ranging from 4 to 32 kHz were generated. The sound intensity was gradually

decreased from 90 to 10 dB sound pressure level (SPL) in 5 dB steps, and auditory thresholds

were determined as the lowest sound intensity that elicited reproducible and recognizable

waves.

Endocochlear potential (EP) measurement

EP recording via the cochlear lateral wall was conducted following previously described

methods (Li et al., 2020; Xie et al., 2023). Both ears' basal turns were analyzed for each animal.

In brief, mice were anesthetized, and a tracheotomy was performed. The tympanic bulla was

opened to expose the cochlea. A small hole was drilled in the cochlear lateral wall of the basal

turn, through which a micropipette electrode filled with 150 mM KCl was carefully advanced into

the scala media using a micromanipulator. The voltage changes during the insertion were

continuously recorded using an amplifier (MultiClamp 700B, Molecular Devices) combined with

a digitizer (Digidata 1440A, Molecular Devices). The EP magnitude of each ear was recorded

and presented individually on the graph.

In situ hybridization (ISH)

ISH was carried out according to previously published methods(Xie et al., 2023). In brief,

mice were transcardially perfused with 4% paraformaldehyde (pH 9.0) and cochleae were

dissected and perfused perilymphatically with the same fixative. The fixed and decalcified

cochleae were cryostat sectioned (14 μm thickness). Sections were then subjected to ISH as

described previously(Xie et al., 2010). Information on antisense cRNA probes is listed in

Supplementary Table S3. Cochleae from n ≥ 4 mice per genotype at a given age were analyzed.

TUNEL (Terminal deoxynucleotidyl transferase dUTP Nick-End labeling) assay

Semi-serial cochlear sections from mice at indicated ages were digested with 10 μg/mL

proteinase K for 30 minutes at 37°C，followed by incubation with a TUNEL reaction mixture

at 37°C for 2 hours. Images of the outer sulcus regions spanning from the basal to apical turns

were captured at 400x magnification. The TUNEL positive nuclei were manually counted on 8

non-consecutive sections per ear. Analysis was conducted on one ear per animal.

Real-time RT-PCR (qRT-PCR)

Total RNA extraction, cDNA synthesis, and qRT-PCR were performed as described

previously(Xie et al., 2023). Data were normalized to 36B4 (Rplp0, internal control) and

presented as fold changes of gene expression in the mutant mice compared to the WT Control.

Each sample was analyzed in duplicate. The primers used are shown in Supplementary Table

S1.

Outer sulcus region isolation and RNA-seq analysis

To isolate the outer sulcus regions from either euthyroid or hypothyroid C57BL/6J mice at

P10, the spiral ligament and attached basilar membrane were isolated in cold Hanks' balanced

salt solution buffer. Subsequently, the outer sulcus region was microdissected by carefully

removing the adjacent basilar membrane and the reddish stria vascularis, along with the

underlying spiral ligament, using two 1 ml syringes equipped with 26G needles. This dissection

method ensured that the isolated subregion primarily contained the outer sulcus with its

underlying fibrocytes, as well as the directly connected spiral prominence epithelium and

Claudius cells.

The isolated outer sulcus regions from four ears were pooled together, and total RNA was

extracted using the Qiagen RNeasy Micro kit. RNA-seq experiments were conducted with five

biological replicates for each of the two conditions, following a previously established

protocol(Xie et al., 2023). Clean reads obtained were aligned to the Mus musculus reference

genome (GCF_000001635.26_GRCm38.p6) using HISAT2 (v2.0.4). Significant differentially

expressed genes (DEGs) between euthyroid and hypothyroid outer sulcus samples were

identified using DESeq2 (v1.4.5), applying criteria of |log2FC|≥1 and Q-value≤0.05, and filtering

out genes with FPKM<10 in both conditions. DEGs that were preferentially expressed by root

cells were identified by intersecting the DEG dataset with the top 800 root cell-enriched genes

at P12, as identified in a previous study using single-cell RNA-seq (Jean et al., 2023).

To identify DEGs containing proximal TH response elements (TREs), the region spanning

10 Kb up and downstream of the transcription start site for each DEG (with reference to the

Mus musculus mm10 genome) was scanned with a classical TRE motif

([THA_MOUSE.H11MO.0.C](#), [THB_MOUSE.H11MO.0.D)](#), which is accessible at

https://hocomoco11.autosome.org. The RcisTarget tool 1.14.0, available at

[https://bioconductor.org,](https://bioconductor.org) was employed for this analysis. The TRE employed is the most

extensively characterized TRs binding consensus sequence. It comprises a direct hexameric

repeat of 5'(A/G)GGT(C/A)A3' with a 4-base spacer, commonly referred to as DR4 elements

(Chatonnet et al., 2013; Umesono et al., 1991).

Statistical analysis

All statistical analyses were conducted using GraphPad Prism 7 software. Data normality

tests were initially performed. Pairwise comparisons were performed using either an unpaired

Student's t-test or a non-parametric Mann-Whitney test. For multiple group comparisons, a

Kruskal-Wallis test (a non-parametric alternative to ANOVA) with Dunn's multiple comparisons

test or a two-way ANOVA followed by post-hoc pairwise tests (Bonferroni method) was utilized

when applicable. Values are presented as mean ± SD or as medians with interquartile ranges

(25-75%) for non-parametric tests exclusively. Tests and groups are noted in figure legends. In

all instances, a two-tailed p-value < 0.05 was considered statistically significant. Non-significant

differences between samples are denoted as "ns," while p-values for significant differences are

displayed in the graphs.

Author Contributions: ZX and WZ conceived the idea, and wrote the manuscript; ZX, WZ,

DH, and JT analyzed data and revised the manuscript; HY provided animal models; XM, FJ, JT,

LS, YC, and SH performed most of the experiments. CW analyzed RNA-seq data. All authors

were involved in the final approval of the submission of the manuscript.

Declarations of interest: The authors declare that there are no conflicts of interest regarding

the publication of this paper.

Acknowledgments and fundings: This work was supported by grants from the National Key

R&D Program (2019YFA0802500, 2018YFA0800602), the National Natural Science

Foundation of China (32271162), and the Collaborative Innovation Program of Shanghai

Municipal Health Commission (2020CXJQ01).

Data availability: RNA-seq data are accessible in Gene Expression Omnibus (GEO) under the

accession number GSE268897 with the hyperlinks

https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE268897 (Xie, et al, 2024). Other data

supporting the findings of this study are available within the manuscript and its Supplementary

information files.

Reference

Xie Z, Zhang W. Gene expression profiles of microdissected cochlear outer sulcus regions from eu- and hypothyroid mice at P10. GEO. https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE268897. Deposited 3 Jun 2024.

Adachi N, Yoshida T, Nin F, Ogata G, Yamaguchi S, Suzuki T, Komune S, Hisa Y, Hibino H, Kurachi Y. 2013. The mechanism underlying maintenance of the endocochlear potential by the K+ transport system in fibrocytes of the inner ear. *J Physiol* **591**:4459–4472. doi:10.1113/jphysiol.2013.258046

Ahmad S, Chen S, Sun J, Lin X. 2003. Connexins 26 and 30 are co-assembled to form gap junctions in the cochlea of mice. *Biochemical and Biophysical Research Communications* **307**:362–368. doi:10.1016/S0006-291X(03)01166-5

Ando M, Takeuchi S. 1999. Immunological identification of an inward rectifier K+ channel (Kir4.1) in the intermediate cell (melanocyte) of the cochlear stria vascularis of gerbils and rats. *Cell Tissue Res* **298**:179–183. doi:10.1007/s004419900066

Bradley DJ, Towle HC, Young WS. 1994. Alpha and beta thyroid hormone receptor (TR) gene expression during auditory neurogenesis: evidence for TR isoform-specific transcriptional regulation in vivo. *Proc Natl Acad Sci U S A* **91**:439–443. doi:10.1073/pnas.91.2.439

Casimiro MC, Knollmann BC, Ebert SN, Vary JC, Greene AE, Franz MR, Grinberg A, Huang SP, Pfeifer K. 2001. Targeted disruption of the Kcnq1 gene produces a mouse model of Jervell and Lange-Nielsen Syndrome. *Proc Natl Acad Sci U S A* **98**:2526–2531. doi:10.1073/pnas.041398998

Cazals Y, Bevengut M, Zanella S, Brocard F, Barhanin J, Gestreau C. 2015. KCNK5 channels mostly expressed in cochlear outer sulcus cells are indispensable for hearing. *Nat Commun* **6**:8780. doi:10.1038/ncomms9780

Chatonnet F, Guyot R, Benoît G, Flamant F. 2013. Genome-wide analysis of thyroid hormone receptors shared and specific functions in neural cells. *Proc Natl Acad Sci U S A* **110**:E766-775. doi:10.1073/pnas.1210626110

Christ S, Biebel UW, Hoidis S, Friedrichsen S, Bauer K, Smolders JWT. 2004. Hearing loss in athyroid pax8 knockout mice and effects of thyroxine substitution. *Audiol Neurootol* **9**:88–106. doi:10.1159/000076000

Cohen-Salmon M, Ott T, Michel V, Hardelin J-P, Perfettini I, Eybalin M, Wu T, Marcus DC, Wangemann P, Willecke K, Petit C. 2002. Targeted Ablation of Connexin26 in the Inner Ear Epithelial Gap Junction Network Causes Hearing Impairment and Cell Death. *Current Biology* **12**:1106–1111. doi:10.1016/S0960-9822(02)00904-1

Deol MS. 1973a. Congenital deafness and hypothyroidism. *Lancet* **2**:105–6. doi:10.1016/s0140-6736(73)93310-2

Deol MS. 1973b. An Experimental Approach to the Understanding and Treatment of Hereditary Syndromes with Congenital Deafness and Hypothyroidism. *Journal of Medical Genetics* **10**:235. doi:10.1136/jmg.10.3.235

Dixon MJ, Gazzard J, Chaudhry SS, Sampson N, Schulte BA, Steel KP. 1999. Mutation of the Na-K-Cl Co-Transporter Gene Slc12a2 Results in Deafness in Mice. *Human Molecular Genetics*

8:1579–1584. doi:10.1093/hmg/8.8.1579

Duvall III AJ. 1969. The ultrastructure of the external sulcus in the guinea pig cochlear duct. *The*
Laryngoscope **79**:1–29. doi:10.1288/00005537-196901000-00001

Forrest D, Erway LC, Ng L, Altschuler R, Curran T. 1996. Thyroid hormone receptor beta is essential
for development of auditory function. *Nat Genet* **13**:354–7. doi:10.1038/ng0796-354

Furness DN. 2019. Forgotten Fibrocytes: A Neglected, Supporting Cell Type of the Cochlea With
the Potential to be an Alternative Therapeutic Target in Hearing Loss. *Front Cell Neurosci*
13:532. doi:10.3389/fncel.2019.00532

Galic M, Giebel W. 1989. An electron microscopic study of the function of the root cells in the
external spiral sulcus of the cochlea. *Acta Otolaryngol Suppl* **461**:1–15.

Grøntved L, Waterfall JJ, Kim DW, Baek S, Sung M-H, Zhao L, Park JW, Nielsen R, Walker RL, Zhu
YJ, Meltzer PS, Hager GL, Cheng S. 2015. Transcriptional activation by the thyroid
hormone receptor through ligand-dependent receptor recruitment and chromatin
remodelling. *Nat Commun* **6**:7048. doi:10.1038/ncomms8048

Groza T, Gomez FL, Mashhadi HH, Muñoz-Fuentes V, Gunes O, Wilson R, Cacheiro P, Frost A,
Keskivali-Bond P, Vardal B, McCoy A, Cheng TK, Santos L, Wells S, Smedley D, Mallon A-
M, Parkinson H. 2023. The International Mouse Phenotyping Consortium: comprehensive
knockout phenotyping underpinning the study of human disease. *Nucleic Acids Research*
51:D1038–D1045. doi:10.1093/nar/gkac972

Gu S, Olszewski R, Taukulis I, Wei Z, Martin D, Morell RJ, Hoa M. 2020. Characterization of rare
spindle and root cell transcriptional profiles in the stria vascularis of the adult mouse
cochlea. *Sci Rep* **10**:18100. doi:10.1038/s41598-020-75238-8

Hao X, Xing Y, Moore MW, Zhang J, Han D, Schulte BA, Dubno JR, Lang H. 2014. Sox10 expressing
cells in the lateral wall of the aged mouse and human cochlea. *PLoS One* **9**:e97389.
doi:10.1371/journal.pone.0097389

Hebert JM, McConnell SK. 2000. Targeting of cre to the Foxg1 (BF-1) locus mediates loxP
recombination in the telencephalon and other developing head structures. *Dev Biol*
222:296–306. doi:10.1006/dbio.2000.9732

Hibino H, Nin F, Tsuzuki C, Kurachi Y. 2010. How is the highly positive endocochlear potential
formed? The specific architecture of the stria vascularis and the roles of the ion-transport
apparatus. *Pflugers Arch* **459**:521–33. doi:10.1007/s00424-009-0754-z

Jagger DJ, Forge A. 2015. Connexins and gap junctions in the inner ear--it's not just about K(+)
recycling. *Cell Tissue Res* **360**:633–44. doi:10.1007/s00441-014-2029-z

Jagger DJ, Forge A. 2013. The enigmatic root cell - emerging roles contributing to fluid
homeostasis within the cochlear outer sulcus. *Hear Res* **303**:1–11.
doi:10.1016/j.heares.2012.10.010

Jagger DJ, Forge A. 2006. Compartmentalized and signal-selective gap junctional coupling in the
hearing cochlea. *J Neurosci* **26**:1260–8. doi:10.1523/JNEUROSCI.4278-05.2006

Jagger DJ, Nevill G, Forge A. 2010. The Membrane Properties of Cochlear Root Cells are Consistent
with Roles in Potassium Recirculation and Spatial Buffering. *J Assoc Res Otolaryngol*
11:435–48. doi:10.1007/s10162-010-0218-3

Jean P, Wong Jun Tai F, Singh-Estivalet A, Lelli A, Scandola C, Megharba S, Schmutz S, Roux S,
Mechaussier S, Sudres M, Mouly E, Heritier A-V, Bonnet C, Mallet A, Novault S, Libri V,
Petit C, Michalski N. 2023. Single-cell transcriptomic profiling of the mouse cochlea: An

atlas for targeted therapies. *Proceedings of the National Academy of Sciences*
120:e2221744120. doi:10.1073/pnas.2221744120

Karolyi IJ, Dootz GA, Halsey K, Beyer L, Probst FJ, Johnson KR, Parlow AF, Raphael Y, Dolan DF,
Camper SA. 2007. Dietary thyroid hormone replacement ameliorates hearing deficits in
hypothyroid mice. *Mamm Genome* **18**:596–608. doi:10.1007/s00335-007-9038-0

Kerr TP, Ross MD, Ernst SA. 1982. Cellular localization of Na+,K+-ATPase in the mammalian
cochlear duct: significance for cochlear fluid balance. *Am J Otolaryngol* **3**:332–338.
doi:10.1016/s0196-0709(82)80006-9

Kim HJ, Gratton MA, Lee J-H, Perez Flores MC, Wang W, Doyle KJ, Beermann F, Crognale MA,
Yamoah EN. 2013. Precise toxigenic ablation of intermediate cells abolishes the "battery"
of the cochlear duct. *J Neurosci* **33**:14601–14606. doi:10.1523/JNEUROSCI.2147-13.2013

Kimura RS. 1984. Sensory and accessory epithelia of the cochlea. *Friedmann, I And J Ballantyne*
(Ed) Ultrastructural Atlas Of The Inner Ear Ix++329p Butterworths London, England;
Boston, Mass , Usa Illus P101-132.

Lee JH, Chiba T, Marcus DC. 2001. P2X2 receptor mediates stimulation of parasensory cation
absorption by cochlear outer sulcus cells and vestibular transitional cells. *J Neurosci*
21:9168–9174. doi:10.1523/JNEUROSCI.21-23-09168.2001

Li Y, Liu H, Zhao X, He DZ. 2020. Endolymphatic Potential Measured From Developing and Adult
Mouse Inner Ear. *Front Cell Neurosci* **14**:584928. doi:10.3389/fncel.2020.584928

Lim DJ, Anniko M. 1985. Developmental morphology of the mouse inner ear. A scanning electron
microscopic observation. *Acta Otolaryngol Suppl* **422**:1–69.

Liu S, Shen S, Yan Y, Sun C, Lu Z, Feng H, Ma Y, Tang Z, Yu J, Wu Y, Gereben B, Mohácsik P, Fekete
C, Feng X, Yuan F, Guo F, Hu C, Shao M, Gao X, Zhao L, Li Y, Jiang J, Ying H. 2022.
Triiodothyronine (T3) promotes brown fat hyperplasia via thyroid hormone receptor α
mediated adipocyte progenitor cell proliferation. *Nat Commun* **13**:3394.
doi:10.1038/s41467-022-31154-1

Liu W, Wang C, Yu H, Liu S, Yang J. 2018. Expression of acetylated tubulin in the postnatal
developing mouse cochlea. *European journal of histochemistry : EJH* **62**.
doi:10.4081/ejh.2018.2942

Marcus DC, Wu T, Wangemann P, Kofuji P. 2002. KCNJ10 (Kir4.1) potassium channel knockout
abolishes endocochlear potential. *Am J Physiol Cell Physiol* **282**:C403-407.
doi:10.1152/ajpcell.00312.2001

Mendoza A, Tang C, Choi J, Acuña M, Logan M, Martin AG, Al-Sowaimel L, Desai BN, Tenen DE,
Jacobs C, Lyubetskaya A, Fu Y, Liu H, Tsai L, Cohen DE, Forrest D, Wilson AA, Hollenberg
AN. 2021. Thyroid hormone signaling promotes hepatic lipogenesis through the
transcription factor ChREBP. *Sci Signal* **14**:eabh3839. doi:10.1126/scisignal.abh3839

Mustapha M, Fang Q, Gong T-W, Dolan DF, Raphael Y, Camper SA, Duncan RK. 2009. Deafness
and Permanently Reduced Potassium Channel Gene Expression and Function in
Hypothyroid Pit1dw Mutants. *The Journal of Neuroscience* **29**:1212.
doi:10.1523/JNEUROSCI.4957-08.2009

Ng L, Cordas E, Wu X, Vella KR, Hollenberg AN, Forrest D. 2015. Age-Related Hearing Loss and
Degeneration of Cochlear Hair Cells in Mice Lacking Thyroid Hormone Receptor beta1.
Endocrinology **156**:3853–65. doi:10.1210/en.2015-1468

Ng L, Kelley MW, Forrest D. 2013. Making sense with thyroid hormone--the role of T(3) in auditory

development. *Nat Rev Endocrinol* **9**:296–307. doi:10.1038/nrendo.2013.58

Roth B, Bruns V. 1992. Postnatal development of the rat organ of Corti. I. General morphology, basilar membrane, tectorial membrane and border cells. *Anat Embryol (Berl)* **185**:559–69. doi:10.1007/BF00185615

Royaux IE, Belyantseva IA, Wu T, Kachar B, Everett LA, Marcus DC, Green ED. 2003. Localization and functional studies of pendrin in the mouse inner ear provide insight about the etiology of deafness in pendred syndrome. *J Assoc Res Otolaryngol* **4**:394–404. doi:10.1007/s10162-002-3052-4

Rüsch A, Erway LC, Oliver D, Vennström B, Forrest D. 1998. Thyroid hormone receptor β-dependent expression of a potassium conductance in inner hair cells at the onset of hearing. *Proceedings of the National Academy of Sciences* **95**:15758–15762. doi:10.1073/pnas.95.26.15758

Rusch A, Ng L, Goodyear R, Oliver D, Lisoukov I, Vennstrom B, Richardson G, Kelley MW, Forrest D. 2001. Retardation of cochlear maturation and impaired hair cell function caused by deletion of all known thyroid hormone receptors. *J Neurosci* **21**:9792–9800. doi:10.1523/JNEUROSCI.21-24-09792.2001

Sakagami M, Fukazawa K, Matsunaga T, Fujita H, Mori N, Takumi T, Ohkubo H, Nakanishi S. 1991. Cellular localization of rat Isk protein in the stria vascularis by immunohistochemical observation. *Hear Res* **56**:168–172. doi:10.1016/0378-5955(91)90166-7

Sawant O, Horton AM, Shukla M, Rayborn ME, Peachey NS, Hollyfield JG, Rao S. 2015. Light-Regulated Thyroid Hormone Signaling Is Required for Rod Photoreceptor Development in the Mouse Retina. *Invest Ophthalmol Vis Sci* **56**:8248–8257. doi:10.1167/iovs.15-17743

Sharlin DS, Ng L, Verrey F, Visser TJ, Liu Y, Olszewski RT, Hoa M, Heuer H, Forrest D. 2018. Deafness and loss of cochlear hair cells in the absence of thyroid hormone transporters Slc16a2 (Mct8) and Slc16a10 (Mct10). *Sci Rep* **8**:4403. doi:10.1038/s41598-018-22553-w

Shodo R, Hayatsu M, Koga D, Horii A, Ushiki T. 2017. Three-dimensional reconstruction of root cells and interdental cells in the rat inner ear by serial section scanning electron microscopy. *Biomed Res* **38**:239–248. doi:10.2220/biomedres.38.239

Song MH, Choi S-Y, Wu L, Oh S-K, Lee HK, Lee DJ, Shim D-B, Choi JY, Kim U-K, Bok J. 2011. Pou3f4 deficiency causes defects in otic fibrocytes and stria vascularis by different mechanisms. *Biochemical and Biophysical Research Communications* **404**:528–533. doi:10.1016/j.bbrc.2010.12.019

Soriano P. 1999. Generalized lacZ expression with the ROSA26 Cre reporter strain. *Nat Genet* **21**:70–71. doi:10.1038/5007

Spicer SS, Schulte BA. 1996. The fine structure of spiral ligament cells relates to ion return to the stria and varies with place-frequency. *Hear Res* **100**:80–100. doi:10.1016/0378-5955(96)00106-2

Steel KP, Barkway C. 1989. Another role for melanocytes: their importance for normal stria vascularis development in the mammalian inner ear. *Development* **107**:453–463. doi:10.1242/dev.107.3.453

Sun J, Ahmad S, Chen S, Tang W, Zhang Y, Chen P, Lin X. 2005. Cochlear gap junctions coassembled from Cx26 and 30 show faster intercellular Ca2+ signaling than homomeric counterparts. *American Journal of Physiology-Cell Physiology* **288**:C613–C623. doi:10.1152/ajpcell.00341.2004

Umesono K, Murakami KK, Thompson CC, Evans RM. 1991. Direct repeats as selective response
elements for the thyroid hormone, retinoic acid, and vitamin D3 receptors. *Cell* **65**:1255–
1266. doi:10.1016/0092-8674(91)90020-Y
Uziel A. 1986. Periods of sensitivity to thyroid hormone during the development of the organ of
Corti. *Acta Otolaryngol Suppl* **429**:23–27. doi:10.3109/00016488609122726
Uziel A, Legrand C, Ohresser M, Marot M. 1983. Maturational and degenerative processes in the
organ of Corti after neonatal hypothyroidism. *Hear Res* **11**:203–218. doi:10.1016/0378-
5955(83)90079-5
Uziel Alain, Legrand C, Rabie A. 1985. Corrective effects of thyroxine on cochlear abnormalities
induced by congenital hypothyroidism in the rat. I. Morphological study. *Developmental*
Brain Research **19**:111–122. doi:10.1016/0165-3806(85)90236-6
Walters BJ, Zuo J. 2013. Postnatal development, maturation and aging in the mouse cochlea and
their effects on hair cell regeneration. *Hear Res* **297**:68–83.
doi:10.1016/j.heares.2012.11.009
Wangemann P. 2006. Supporting sensory transduction: cochlear fluid homeostasis and the
endocochlear potential. *J Physiol* **576**:11–21. doi:10.1113/jphysiol.2006.112888
Winter H, Rüttiger L, Müller M, Kuhn S, Brandt N, Zimmermann U, Hirt B, Bress A, Sausbier M,
Conscience A, Flamant F, Tian Y, Zuo J, Pfister M, Ruth P, Löwenheim H, Samarut J, Engel
J, Knipper M. 2009. Deafness in TRbeta mutants is caused by malformation of the tectorial
membrane. *J Neurosci* **29**:2581–2587. doi:10.1523/JNEUROSCI.3557-08.2009
Xie Z, Ma X, Ji W, Zhou G, Lu Y, Xiang Z, Wang YX, Zhang L, Hu Y, Ding Y-Q, Zhang WJ. 2010.
Zbtb20 is essential for the specification of CA1 field identity in the developing
hippocampus. *Proc Natl Acad Sci U S A* **107**:6510–6515. doi:10.1073/pnas.0912315107
Xie Z, Ma X-H, Bai Q-F, Tang J, Sun J-H, Jiang F, Guo W, Wang C-M, Yang R, Wen Y-C, Wang F-
Y, Chen Y-X, Zhang H, He DZ, Kelley MW, Yang S, Zhang WJ. 2023. ZBTB20 is essential
for cochlear maturation and hearing in mice. *Proceedings of the National Academy of*
Sciences **120**:e2220867120. doi:10.1073/pnas.2220867120
Xie Z, Zhang H, Tsai W, Zhang Y, Du Y, Zhong J, Szpirer C, Zhu M, Cao X, Barton MC, Grusby MJ,
Zhang WJ. 2008. Zinc finger protein ZBTB20 is a key repressor of alpha-fetoprotein gene
transcription in liver. *Proc Natl Acad Sci U S A* **105**:10859–10864.
doi:10.1073/pnas.0800647105
Yan Y, Niu Z, Sun C, Li P, Shen S, Liu S, Wu Y, Yun C, Jiao T, Jia S, Li Yuying, Fang Z-Z, Zhao L,
Wang J, Xie C, Jiang C, Li Yan, Feng X, Hu C, Jiang J, Ying H. 2022. Hepatic thyroid hormone
signalling modulates glucose homeostasis through the regulation of GLP-1 production
via bile acid-mediated FXR antagonism. *Nat Commun* **13**:6408. doi:10.1038/s41467-022-
34258-w
Zilz ND, Murray MB, Towle HC. 1990. Identification of multiple thyroid hormone response elements
located far upstream from the rat S14 promoter. *J Biol Chem* **265**:8136–8143.

Figures and figure legends

Fig. 1. Hypothyroidism disrupts the outer sulcus development. (A and B) Representative
Toluidine Blue-stained sections of cochlear upper basal turn from euthyroid (**A**) or methimazole
(MMI)-induced hypothyroid mice (**B**) at P21. (**A**) Euthyroid cochlea showing a mature outer
sulcus characterized by root cells (Rc. indicated by red arrows) with numerous bright branched
processes extending deep into the spiral ligament and upward to behind the spiral prominence
(Sp). The surfaces of root cell bodies were covered by enlarged Claudius cells (Cc, red bar
indicating its height). Co, the organ of Corti; Is, inner sulcus; Sm, scala media; Sv, stria
vascularis; Tc, tunnel of Corti, Tm, tectorial membrane. (**B**) The hypothyroid outer sulcus
consists of a simple layer of cells with a few short processes bordering the scala media,
continued with the flattened Claudius cells. Red arrowheads indicate defects including a
distorted Tm, an abnormal persistence of GER, closure of tunnel of Corti, and a flattened border
region. (**C** and **D**) Immunostaining shows extensive acetyl-α-tubulin positive (red) root cell
processes extending into the inferior region of the euthyroid spiral ligament at P21 (**C**) but only
a few short processes observed in the luminal surface of hypothyroid counterparts (indicated
by white arrowhead) (**D**). Bars = 50 μm for all panels. Nuclei were stained with DAPI (turquoise).
n = 4 mice/ group. Scale bars in all panels = 50 μm. Abbreviations are consistent across figures
in this manuscript.

Fig.2 Otocyst-selective deletion of all known TRs causes deafness and a compromised

EP. (A) X-gal staining showing *Foxg1-Cre* recombinase activity was confined in the cochlear

epithelium lining the scala media (Sm) at P5, including GER, LER (the lesser epithelial ridge),

root cells (Rc), stria vascularis (Sv), as well as spiral ganglions (Sg). n = 3 mice/group. Bar =

100 μm. **(B)** QRT-PCR analysis showing relative mRNA expression levels of *Thra* and *Thrb* in

three groups of cochlea at P10. n = 5 mice/group. **(C)** Representative ABR waveforms obtained

to broadband click stimuli from three groups of mice at 2 months (mon) of age. Sound pressure

levels (SPL) were indicated in decibels (dB). (**D**) Averaged thresholds of Click- and Pure tone-

evoked ABRs from three groups of mice at 2 mon. n = 5 mice/group. (**E**) OV-TRα/βKO mice

showed a reduced EP magnitude in the basal turns compared with control mice at around 1.5

or 4 mon. The number of ears tested was indicated in the graph. Values are shown as medians

with 25-75% interquartile ranges for (**B**), mean ± SD for (**D**), and (**E**). A Mann-Whitney test and

an unpaired Student's t-test were used to compare differences in (**B**) and (**E**), respectively. A

Kruskal-Wallis test with Dunn's multiple comparisons test and a two-way ANOVA followed by

post-hoc pairwise tests (Bonferroni method) were used to compare differences in Click- and

Pure tone-evoked ABR thresholds, respectively. *P* Values were indicated. ns, not significant.

Fig. 3. Otocyst-selective deletion of all known TRs retards root cell development. (A-C)

Representative Toluidine Blue-stained sections showing delayed root cell (RC) development in

OV-TRα/βKO pups at P10 (**C**), compared with that in control **(A)** or OV-TRβKO **(B),** Claudius

cells (Cc), scala media (Sm), spiral prominence (Sp). **(D-F)** Immunostaining showing decreased

acetyl-α-tubulin positive (red) root cell processes within OV-TRα/βKO outer sulcus **(F)**,

compared with that in control **(D)** or OV-TRβKO counterparts **(E)** at P10. Nuclei were stained

with DAPI (turquoise). n = 4 mice/ group. **(G-I)** ISH revealed a reduced expression domain of

Slc26a4 mRNA in OV-TRα/βKO (**I**), compared with that in control **(G)** or OV-TRβKO **(H)** at P10.

(J-L) Representative sections showing disruption of root cell development in OV-TRα/βKO **(L),**

compared with a mature outer sulcus in control **(J)** or OV-TRβKO **(K)** mice at P21. Cc is outlined

by black dots, and Sp is outlined by magenta dots. A degenerated cell is indicated by "De" in

(**L**). The upper basal turns were shown for all panels. Arrowheads indicate defects. Scale bars:

50 μm for all panels. n = 4 mice/group. See also Supplementary Fig. S1-2.

Fig. 4. Otocyst-selective deletion of all TRs causes early-onset root cell degeneration.

(A-C) Representative Toluidine Blue-stained sections of upper basal turn showing malformed tectorial membrane (Tm, Magenta arrowheads), a collapsed organ of Corti (Co, red arrow) in OV-TRα/βKO cochleae at 2 months **(C)**. **(A')**, **(B')** and **(C')** are magnified views of outer sulcus regions indicated by red brackets in **(A)**, **(B)**, and **(C)**, respectively. OV-TRα/βKO showed reduced root cells with pale-staining processes **(C')**. Sp, Spiral prominence. n = 6 mice/group.

(D-E) Representative sections (**D**) and a bar graph (**E**) show a decreased number of Sox10

(magenta) and acetyl-α-tubulin (green) double-positive root cells in OV-TRα/βKO outer sulcus,

compared with that in control or OV-TRβKO at 2 mon. Nuclei were stained with DAPI (blue). n

= 6 mice/group. (**F-G**) Representative sections (**F**) and a bar graph (**G**) show increased

TUNEL-positive cells (magenta) in the OV-TRα/βKO outer sulcus at 2 mon, compared with

that in the control group. The right column in (**F**) is a magnified view of the boxed area in the

middle column, with separated color channels to better show the colocalization of TUNEL-

positive signals with DAPI-stained (blue) nuclei (indicated by arrowheads). n = 8 mice/group.

All values are represented as mean ±SD. A two-way ANOVA and an unpaired T-test were used

to compare differences in (**E**) and (**G**), respectively. *P* Values were indicated. ns, not significant.

Scale bars: 50 μm for all panels. See also Supplementary Fig. S3.

Fig. 5. Stria vascularis (SV) remains intact in OV-TRα/βKO mice. (**A**) Representative Toluidine Blue-stained sections of cochlear upper basal turns showing comparable SV histology among Control, OV-TRβKO, and OV-TRα/βKO mice at P21. (**B**) Bar graph showing comparable SV areas among Control, OV-TRβKO, and OV-TRα/βKO mice at P21. n = 6 mice/group. Values are represented as mean ±SD. A one-way ANOVA followed by post-hoc pairwise tests (Bonferroni method) was used to compare differences between the two groups. ns: not significant. (**C**) Immunostaining revealing comparable expressions of Kir4.1, Na+/K+-ATPase α, NKCC1, and KCNQ1 (red) among Control, OV-TRβKO, and OV-TRα/βKO mice at P21. n = 3 mice/group for each molecular marker. Scale bars: 20 µm for all panels. Nuclei were stained with DAPI (turquoise). See also Supplementary Fig. S4.

Fig. 6. Deficiency of TH leads to the downregulation of a significant portion of root cell-

enriched genes (A) Heatmap displaying 11 upregulated and 167 downregulated DEGs

preferentially expressed by root cells in hypothyroid compared with those in euthyroid mice at

P10 revealed by RNA-seq (n = 5 pairs). **(B-C)** Heatmap displaying expression levels of 11

DEGs associated with hearing loss when mutated in humans or mice (**B**) and 21 DEGs carrying

proximal TRE (**C**). (**D**) Validation of DEGs by qRT-PCR analysis. n = 8 pairs. Values are

represented as mean ±SD. An unpaired T-test was used to compare differences. (**E**) Validation

of DEGs by ISH analysis. Representative sections showing decreased mRNA expression

domains of indicated genes at outer sulcus regions of hypothyroid mice (indicated by red

arrowheads) compared with euthyroid counterparts at P10. n = 3 pairs. Rc: root cell, Sm: scala

media, Sp: spiral prominence. Scale bar: 50 μm. See also Supplementary Fig. S5 and Dataset

S1.

Supplementary information for

Targeted Deletion of All Known Thyroid Hormone Receptors Causes Maturation Retardation and Early-onset Degeneration of Cochlear Outer Sulcus

XianHua Ma[a, #], Fei Jiang[b, #], Chunchun Wei[a], Shuang Han[c], Yuqing Zhang[b], Lianhua Sun[b], Jiaxi Qu[d], Hao Ying[e], Yuxia Chen[a], Jie Tang[d], David Z. He[f], Weiping J. Zhang[a,c,*], Zhifang Xie[b,*]

[#]They contributed equally to the work.

*Corresponding authors: xiezhifang@xinhuamed.com.cn (Z.X., lead contact); Weiping J. Zhang, wzhang@smmu.edu.cn (W.J.Z)

This PDF file includes:

Tables S1 to S3

Figures S1 to S5

Legend for Dataset S1

The other supporting material for this manuscript is Dataset S1.

837 **Table S1. Sequence information for PCR primers**

Gene target	GeneBank accession No.	Primers	Note
Thra-Flox		F: 5'gtatgtgcgccttagagcag R:5'ttatgcccttattctacccc	Genotyping
Thrb-Flox		F: 5'aggcacaggcaggaagtag R:5'tgaagacatcggctctaagga	
Cre		F:5'ggcggatccgaaaagaaaa R:5'cagatggcgcggcaacacc	
Thra	NM_001313983.1	F: 5' atggaacagaagccaagcaa R:5' ttccatctggtgacctggca	RT-PCR
Thrb	NM_001113417.1	F: 5' gtaccactatcgctgcatcacctg R:5' ccgccgcctcttctcacg	
Rplp0 (36B4)	NM_007475.5	F: 5'aagcgcgtcctggcattgtctgtg R:5'tggttgctttggcgggattagtcg	
Tmprss9	NM_001081688.2	F:5' cggaagtgtatgatctcaggctg R:5' atgcggtcagtgagggagaagt	
Ces1d	NM_008084	F:5' catcactgccacccagaagactg R:5' atgccagtgagcttcccgttcag	
Bcam	NM_020486	F:5' agagtggaggattacgatgccg R:5' tgctgttcaggaatacgaagagc	
Atp1b1	NM_009721	F:5' ctggcatcttcatcgggaccat R:5' ggtcattaggacggaaggagatc	
Tmprss3	NM_080727	f:5' ggcgacttcgtgtccatcaatc r:5' gctgagcacttcaaggtgacca	
P2rx2	NM_153400	F:5' actacgagacgcccaaggtgat R:5' cctggtagcttttctgcacgatg	
Atp1a1	NM_144900	F:5' ccgtggataacctctgcttcgt R:5' cgctgtgattggatggtctcct	
S100a1	NM_011309	F:5' agaactgaaagacctgctacaaac R:5' ctccgttttcatccagttccttc	
Cbx6	NM_028763	F:5' aagggaacgtgagctgtatggg R:5' gctcggcttgacagagaaatgc	
Qsox1	NM_001024945	F:5' tggactgtgctgaggagaccaa R:5'gatgagcctcatacgcagagtc	
Padi2	NM_008812	F:5' cgagtttggctacattgaggcac R:5'ctggtgacagtctcaaagaggg	

838
839

840 **Table S2. The information for primary antibodies**

Antibody	Resource	Dilution
anti-alpha 1 Sodium Potassium ATPase	Abcam, ab7671	1:2000
anti-KCNQ1	Santa Cruz, SC-10646	1:2000
anti-Kir4.1	Alomone labs, APC-035	1:2000
anti-NKCC1	Invitrogen, PA5-98154	1:2000
anti-Acetylated Tubulin	Sigma, T7451	1:300
anti-Sox10	Abclonal, A15100	1:2000
anti-CX26	Abcam, ab65969	1:2000
anti-CX30	Invitrogen, 33-2500	1:2000

843 **Table S3. Sequence information for cRNA probes**

Gene target	Genebank accession No.	PCR Primers	Probe length (bp)
Slc26a4#	NM_011867.4	F: 5'cagagatgcggcccgagtgttg R: 5'tcccgttcttggttgccctgagtt	1271
Epyc	NM_001379456. 1	F: 5' aagcattctccaggttaggtttt R:5' tcttgggcagtggaggaatag	596
Gas2	NM_001330607 .2	F: 5'gaccttcgctggctatct R:5'taaaaggtacctgctattcc	801
Prss36	NM_001319147 .1 425-1589	F: 5'gggggctcggacgctcatc R:5'ggctcccgggcagaaaatagtgtt	1165

(#) This cRNA probe has been reported previously (reference: PMID 37279265). *The other* 3
probes were prepared using a PCR-based technique (reference: PMID 25624266). The reverse
primers include the T7 promoter sequence (GGATCCTAATACGACTCACTATAGGGAG) at the
5' end. The antisense-RNA strand was produced by transcription of the PCR product using T7
RNA polymerase.

Fig. S1. Cochlea-selective deletion of all known TRs retards root cell development. (A-C)

ISH showing a decreased expression domain of *Epyc* mRNA (a root cell marker) in OV-

TRα/βKO outer sulcus **(C)** at P10, compared with that in control **(A)** or OV-TRβKO **(B)**

counterparts. n = 4 mice/group. **(D-F)** Representative immunostained sections showing

reduced acetyl-α-tubulin positive (red) root cell processes within the inferior region of the spiral

ligament from OV-TRα/βKO basal turns **(F)** compared with those from control **(D)** or OV-TRβKO

counterparts **(E)** at P21. Scale bars: 50 μm for all panels. Nuclei were stained with DAPI

(turquoise). n = 4 mice/group. Rc, root cell; Sm, scala media; Sv, stria vascularis; Sp, spiral

prominence. Related to Fig.3.

Fig. S2. Development retardation compound with degeneration in root cell from OV-TRα/βKO mice at 1.5 months of age. (A-C) Representative Toluidine Blue-stained sections of upper basal turns from control **(A)**, and OV-TRα/βKO **(B and C)** cochleae at 1.5 months of age. Magenta arrowheads in **(B and C)** indicate tectorial membrane (Tm) malformations in OV-TRα/βKO cochleae. Red arrows indicate either delayed differentiation **(B)** or degeneration **(C)** of Boettcher cells (Boc). **(A')**, **(B')** and **(C')** are magnified views of outer sulcus regions indicated by red brackets in **(A)**, **(B)**, and **(C)**, respectively. **(A')** Control mice showed numerous branched root cells (Rc) processes extending deep into the inferior region of the spiral ligament and upward to behind the spiral prominence (Sp), **(B')** whereas OV-TRα/βKO displayed reduced root cell processes within the superficial region of the spiral ligament. **(C')** Some samples from OV-TRα/βKO mice showed degenerative changes, displaying reduced root cells with pale-staining processes (indicated by red arrowhead). Cc, Claudius cells; Spiral prominence. Scale bars: 50 μm for all panels. n = 4 mice/group. Related to Fig. 3.

Fig. S3. Absence of all known TRs in cochlear epithelium causes early-onset root cell degeneration. (A-C) Representative Toluidine Blue-stained sections of apical turns show an upright organ of Corti (Co) in control **(A)** or OV-TRβKO **(B)**, whereas only sparse cells (indicated by the red arrow) in OV-TRα/βKO epithelium **(C)** at 2 months of age. Tectorial membrane (Tm, indicated by magenta arrowheads) malformations were observed in both OV-TRβKO and OV-TRα/βKO. **(A')**, **(B')** and **(C')** are magnified views of boxed region in **(A)**, **(B)**, and **(C)**, respectively, showing reduced root cells (Rc) in OV-TRα/βKO outer sulcus (indicated by a red bracket) **(C')**, compared with those in control **(A')** or OV-TRβKO **(B')**. Sp, Spiral prominence. n = 6 mice/group. Scale bars: 50 μm for all panels. Related to Fig. 4.

Fig. S4. Ablation of TRs does not affect the expression patterns of connexins in cochleae.

Representative immuno-stained sections of upper basal turn showing comparable expression

patterns of CX26 and CX30 (red) between Control and OV-TRα/βKO mice at P21. Nuclei were

stained with DAPI (turquoise). Co, organ of Corti; Os, outer sulcus; Sl, spiral ligament; Sv, stria

vascularis. Scale bars: 50 μm. n = 3 mice/group. Related to Fig. 5.

Fig. S5. Volcano plot showing the significantly differentially expressed genes (DEGs) detected

in microdissected outer sulcus regions from hypothyroid cochlea at P10 compared to euthyroid

counterparts by RNA-seq analysis (n = 5 pairs, fold change ≥ 2, Q value ≤ 0.05). Related to

Fig. 6 and Supplementary Dataset 1.

Dataset S1 (separate file): List of total DEGs and DEGs preferentially expressed by root

cells in microdissected outer sulcus regions from hypothyroid cochlea at P10 compared to

euthyroid counterparts by RNA-Seq analysis. DEGs were identified based on a fold change ≥

2, Q value ≤ 0.05 (Genes of FPKM (Fragments per kilobase per million mapped reads) < 10

were filtered out). Related to Fig. 6 and Supplementary Fig. S5.

899