Targeted Deletion of All Known Thyroid Hormone Receptors Causes Maturation Retardation and Early-onset Degeneration of Cochlear Outer Sulcus

Xian-Hua Ma[1#], Fei Jiang[2#], Shuang Han[3], Yuqing Zhang[1,2], Lianhua Sun[2], Jiaxi Qu[4], Hao Yin[5], Yuxia Chen[1], Jie Tang[4], David Z. He[6], Weiping J. Zhang[1,3,*], Zhifang Xie[2,*]

[1]Department of Pathophysiology, Naval Medical University, Shanghai 200433, China
[2]Ministry of Education-Shanghai Key Laboratory of Children's Environmental Health, Institute of Early Life Health, Xinhua Hospital, Shanghai Jiaotong University School of Medicine, Shanghai 200092, China
[3]NHC Key Laboratory of Hormones and Development, Chu Hsien-I Memorial Hospital & Tianjin Institute of Endocrinology, Tianjin Medical University, Tianjin 300134, China
[4]Department of Physiology, Southern Medical University, Guangzhou, Guangdong 510515, China
[5]CAS Key Laboratory of Nutrition, Metabolism and Food Safety, Shanghai Institute of Nutrition and Health, University of Chinese Academy of Sciences, Chinese Academy of Sciences, and Shanghai Jiao Tong University Affiliated Sixth People's Hospital, Shanghai 200031, China.
[6]Department of Biomedical Sciences, Creighton University School of Medicine, Omaha, NE 68178, USA

[#]They contributed equally to the work.

*Corresponding authors: Zhifang Xie (zfxie70@hotmail.com); Weiping Zhang, zbtb20@aliyun.com;

Author Contributions: ZX and WZ conceived the idea, and wrote the manuscript; ZX, WZ, DH and JT analyzed data and revised the manuscript; HY provided animal models; XM, FJ, JT, LS, YC, and SH performed most of the experiments. All authors were involved in final approval of the submission of the manuscript.

Competing Interest Statement: All the authors claim there is no interest of conflict about this work.

Classification: Developmental biology, Neuroscience

Keywords: Cochlea, Thyroid hormone receptor, development, root cell, transcription factor

Abstract

Thyroid hormone (TH) and/or its receptors (TRs) are essential for hearing, cochlear

late-stage development and maintaining endocochlear potential (EP). The mechanism

underlying the EP reduction upon deficiency of TH or TRs remains unknown. The cochlear

outer sulcus (root cell) is considered to be important for the maintenance of endolymph

homeostasis and EP, and undergoes dramatic morphological changes during cochlear

late-stage developmental window. It has been long neglected whether TH and/or TRs are

required for root cell differentiation and function. Here we demonstrate a key role for TH or

TRs in postnatal root cell development and survival in mice. Anti-thyroid drug-induced

developmental hypothyroidism results in severe retardation in root cell differentiation.

Targeted deletion of both TRα and TRβ but not individual TRβ in cochlear epithelium

causes a similar impairment, compounded with an early-onset degeneration of root cells

and a reduced EP, without obvious alterations in stria vascularis. Furthermore, we

demonstrate that TH signaling acts in parallel with ZBTB20 in the outer sulcus, a

transcription factor which we have recently reported to be essential for the outer sulcus

differentiation. Our results support that TRα and TRβ act redundantly in promoting root cell

late-stage differentiation and maintaining its cellular homeostasis, and root cell dysfunction

at least partially contribute to a reduced EP upon deficiency of TH or TRs. These findings

imply that root cells may play an important role in maintaining the health of cochlear

sensory epithelium and normal EP.

Significance Statement

Thyroid hormone (TH) and/or its receptors (TRs) are essential for hearing and

cochlear development. The cochlear outer sulcus (root cell) is considered to be important for

cochlear ionic and metabolic homeostasis. We found that combined deletion of both TRα and

TRβ but not individual TRβ results in severe retardation in root cell differentiation accompanied

by an early-onset degeneration and a reduced endocochlear potential (EP), suggesting TRα

and TRβ act redundantly in promoting root cell differentiation and maintaining its cellular

homeostasis. Our findings suggest that root cell dysfunction may at least partially contribute

to a reduced EP upon deficiency of TRs, and root cells may play an important role in

maintaining the health of cochlear sensory epithelium and EP.

Introduction

Thyroid hormone (TH) is necessary for normal development of cochlea and of hearing[1]. Both early-onset congenital hypothyroidism and environmentally based iodine deficiency can cause hearing loss in humans and rodents[2–4]. TH exerts its function mainly by binding to two different thyroid hormone receptors (TRs), namely TRα and TRβ, which are ligand-dependent transcription factors encoded by *Thra* and *Thrb*, respectively [5–7]. TRα is individually dispensable for hearing[8], while lack of TRβ alone causes deafness accompanied by tectorial membrane (TM) malformation [5, 7]. Deficiency of all known TRs by combined global deletion of *Thra* and *Thrb* in mice results in exacerbated cochlear defects including a more severe distorted TM, a severe retardation in the regression of great epithelial ridge (GER, also known as Kolliker's organ) and remodeling of the organ of Corti, as well as a reduced endocochlear potential (EP) [6]. The mechanism underlying EP reduction upon TRs deficiency remains enigmatic, as no obvious stria vascularis (SV) alterations have been reported in these mice, indicating defects beyond SV might contributing to the EP reduction.

Cochlear outer sulcus epithelial cells (root cells) are poorly understood cell types that are considered to be important for the maintenance of endolymph homeostasis and EP [9–13]. The outer sulcus is located on cochlear lateral wall between the outer boarder Claudius cells and the spiral prominence, and undergoes dramatic morphological changes during postnatal developmental window, when the cochlea displays the greatest sensitivity to TH [14–16]. At birth, the outer sulcus is composed of a simple layer of cuboidal-shaped epithelial cells lining the scala media. Shortly after birth, these cells start to infiltrate laterally and superiorly into the spiral ligament underlying the spiral prominence and develop extensive tree-like branched

processes, and thus be designated as "root cells". During the same period, Claudius cells

become greatly enlarged and meet the spiral prominence epithelium covering the underlying

root cell bodies, so that root cells are no longer directly in contact with the endolymph in basal

turns. In the apical turn, however, root cells face the endolymph directly and have fewer

processes(9, 11, 13, 17–19).

Little is known about how root cell differentiation is controlled. We have recently reported

that transcription factor ZBTB20 is essential for cochlear late-stage development including root

cell differentiation. Cochlear epithelium selective deficiency of ZBTB20 causes deafness, EP

reduction and cochlear malformation including a deformed TM and the organ of Corti , as well

as a failure in the maturation of the outer border and outer sulcus region (20). Very interestingly,

Thrb is expressed by the developing and adult root cells as revealed by using a *lacz* gene

reporter(21) or single-cell RNA sequencing (22), and *Thra* mRNA is widely expressed by the

lateral wall including root cells (23, 24). It remains unknown whether TH and the expression of

TRs in cochlea are essential for remodeling and maturation of the outer sulcus.

Here we find that TRs expression in otocyst-derived cochlear epithelium is necessary for

postnatal root cell development and survival. We show that root cell development was severely

retarded in antithyroid drug-induced developmental hypothyroidism or otocyst-selective *Thra*

and *Thrb* double knockout mice, while no obvious root cell defects were observed upon

individual *Thrb* deletion. Absence of both TRα and TRβ also led to an early-onset degeneration

of root cells and a compromised EP value. We further show that TH signaling acts in parallel

with ZBTB20 in controlling outer sulcus development.

Results

Hypothyroidism results in the disruption in the outer sulcus development.

To determine the role of thyroid hormone in the outer sulcus development, developmental

hypothyroidism was induced by antithyroid treatment as previously described (25). The

antithyroid treatment was initiated on day 17 of gestation and continued during lactation until

postnatal day (P) 21, which covers the developmental window when cochlea shows high

sensitivity to TH deprivation (14, 26). We first performed histological analysis on the cochleae

from eu- or hypothyroid mice at P21, when cochlea becomes mature morphologically and

functionally in normal mice(18, 27). We applied a resin-embedding technique to better preserve

cochlear histological features. The semi-thin sections of cochlear basal turns revealed that

euthyroid cochlear sensory epithelium displayed a typical mature morphology: an upright organ

of Corti, a well-developed inner sulcus and a normal TM. In the outer sulcus region, enlarged

Claudius cells covered the underlying root cells, which showed numerous bright branched

processes extending deep into the spiral ligament and upward to behind the spiral prominence

(Fig. 1*A*). Consistent with previous report(14), the hypothyroid cochlear epithelium displayed a

severe retardation in the sensory epithelium remodeling manifested by an abnormal

persistence of GER, closure of tunnel of Corti, and a severe distorted TM. Interestingly, the

outer sulcus also displayed an immature appearance: a simple layer of cells with a few short

processes bordering the scala media, continued with the flattened Claudius cells, suggesting a

severe retardation in the root cell and Claudius cell differentiation (Fig. 1*B*).

We and others have previously shown that root cell cytoplasm and their processes are

enriched with acetyl-α-tubulin (13, 20, 28). To better visualize root cells, we then performed

immunofluorescence staining for acetyl-α-tubulin. Consistent with the above histological

observations, euthyroid cochleae displayed robust and extensive acetyl-α-tubulin signals deep

into the spiral ligament (Fig. 1C), whereas the hypothyroid counterparts only showed limited

signals distributed mainly around the luminal surface of the spiral ligament, suggesting root cell

differentiation defects (Fig. 1D). Thus, TH signaling is necessary for the timely development of

the outer sulcus

Cochlea-selective deletion of all known TRs causes deafness and a compromised EP.

As TH effects are mainly mediated by TRα and TRβ(1), both of which are expressed in the

developing outer sulcus(21–23), we therefore asked if the two receptors are essential for root

cell differentiation. To answer this question, we generated mice with individual *Thrb* or combined

with *Thra* gene selectively deleted in otic vesicle (hereafter referred as OV-TRβKO and OV-

TRα/βKO, respectively) using the Cre-loxP system. Individual *Thra*-deleted mice were excluded

in this study as *Thra*-null mice have normal hearing threshold(8). We crossed *Thrb-floxed*

(*Thrb^flox^*) mice(29) individually or combined with *Thra ^flox^* mice(30) to *Foxg1-Cre* transgenic

mice, a line that can mediate recombination selectively in the otic vesicle by E8.5(31). We

confirmed the tissue specificity of Cre-mediated recombination by mating *Foxg1-Cre* mice with

Rosa26-lacZ reporter mice. Accordingly, Cre-recombinase activity (displayed by X-gal staining)

was confined to otocyst-derived cochlear epithelium at P5, including GER, less epithelial ridge

(LER), outer sulcus epithelium (root cells) and stria vascularis (SV) marginal cells, as well as

spiral ganglions. As expected, no signal was detected in cochlear fibrocytes which are derived

from the periotic mesenchyme surrounding the otocyst (32). No background signal was

detected on *cre*-negative control mice (Fig. 2*A*). The deletion efficiency was further determined

by real-time RT-PCR analysis showing that expression levels of *Thrb* but not *Thra* mRNA were

significantly down regulated in OV-TRβKO cochleae at P10, while both of *Thrb* and *Thra* mRNA

were significantly down regulated in OV-TRα/βKO cochleae (Fig. 2*B*).

To evaluate the auditory function of the mutant mice, we performed ABR (auditory

brainstem response) tests. Both of OV-TRβKO and OV-TRα/βKO mice did not show typical

ABR responses to either click （Fig. 2*C, D*） or frequency-specific tone burst stimuli at 90 dB

(the highest intensity tested) around 2 months （Fig. 2*D*), in sharp contrast to normal thresholds

seen in wild-type (WT) control mice. These data demonstrate that cochlear expression of TRβ

or all known TRs is essential for hearing, in agreement with previous reports(5, 6).

As global deletion of all known TRs results in a ~50% reduction in EP magnitude in adult

mice (6), we thus measure EP from OV-TRα/βKO mice. The OV-TRα/βKO mice showed a slight

but statistically significant reduction in EP magnitude when compared with WT controls at ~1.5

months of age （Fig. 2*E*）, suggesting expression of TRs in cochlear epithelium contributes to

EP maintenance at this age.

Cochlea-selective deficiency of all known TRs retards root cell development.

We next investigated cochlear morphological changes in TRs mutant mice, with an

emphasis on the development of outer sulcus region. At P10, 2 days prior to the onset of hearing

in mice, root cells with branched processes had already invaded deep into the spiral ligaments

in WT pups (Fig. 3*A*). Comparable root cell development was also observed in OV-TRβKO

counterparts (Fig. 3*B*). The outer sulcus in OV-TRα/βKO mice, however, was largely composed

of a single cuboidal epithelial layer bordering the endolymphatic space (Fig. 3*C*), suggesting a

delay in the root cell differentiation. Consistent with the histological observations, OV-TRα/βKO

showed apparently less acetyl-α-tubulin positive cell processes within the spiral ligament

compared with those in WT and OV-TRβKO counterparts (Fig. 3*D-F*). To further demonstrate

root cell differentiation, we examined expression patterns of *Pendrin* (*Slc26a4*) and *Epyc*

(*Epiphycan*), which are abundantly and selectively expressed by root cells(20, 22, 33).) Strong

signals of *Pendrin* or *Epyc* mRNA could be detected deep inside the spiral ligament in both WT

and OV-TRβKO cochleae, but in OV-TRα/βKO cochleae, expression domains of both

molecules were adjacent to the luminal surface of the spiral ligament (Fig. 3*G-I, and SI*

Appendix Fig. S1*A-C*), consistent with a delay in the root cell development. These data

demonstrate that root cell development appeared to be normally initiated in OV-TRβKO but was

delayed in OV-TRα/βKO pups.

By P21, comparable mature morphological features were observed between OV-TRβKO

and WT outer sulcus, showing elaborate root cell branches extending deep into the inferior

region of spiral ligament and upward to behind the spiral prominence in the basal turn (Fig. 3*J*

and *K*). As previously described(17, 34), the surface of root cell bodies was covered by greatly

enlarged Claudius cells and spiral prominence epithelium, thus no longer directly exposed to

the endolymph at this stage. The double knockout mice, however, showed only an initial sign

of root cell differentiation. Root cells with a few processes, were mainly bordering the scala

media, separating the flattened Claudius cell and the spiral prominence epithelium, suggesting

a severe retardation in the differentiation of root cells and Claudius cells (Fig. 3*L*). It should be

noted that some cells with a pale-staining cytoplasm and nucleus could be occasionally

detected in OV-TRα/βKO outer sulcus at this age, suggesting degenerative changes (Fig. 3*L*).

The defects were further demonstrated by reduced acetyl-α-tubulin positive root cell processes

in OV-TRα/βKO outer sulcus, compared with WT or OV-TRβKO counterparts (*SI Appendix* Fig.

S1*D-F*). By 1.5 months, the development of root cells and Claudius cells in OV-TRα/βKO had

progressed to some extent but still lagged behind the developmental status in WT controls (*SI*

Appendix Fig. S2*A-B*, *A'* and *B'*). Degenerative changes in root cells and cochlear outer border

cells (Boettcher's cells) were noticed in some OV-TRα/βKO samples (*SI Appendix* Fig. S2*C*

and *C'*).

Taken together, these data show that the root cell development appeared intact in OV-

TRβKO but was severely retarded in OV-TRα/βKO mice, suggesting that TRβ is individually

dispensable and TRs may play redundant roles in root cell development.

Deficiency of all known TRs causes early-onset root cell degeneration.

Although the degeneration of cochlear hair cells and supporting cells has been reported

previously in developmental hypothyroidism, and in mice with global deletion of TH transporters

(35, 36, 4), it remains unknown if root cells survival depends on TH signaling. As described

above, sporadic degenerative cells were occasionally detected in OV-TRα/βKO outer sulcus at

P21 (Fig. 3*L*), or ~1.5 months of age (*SI Appendix* Fig. S2*C* and *C'*). By 2 months, WT control

or OV-TRβKO outer sulcus showed no obvious degenerative changes, whereas OV-TRα/βKO

exhibited apparently reduced root cells with pale-staining processes and nuclei in both basal

and apical turns (Fig. 4*A-C*, *A'-C'*, and *SI Appendix* Fig. S3), suggesting progressive

degenerations. Concomitant with the root cell degeneration, variable degenerations of the

organ of Corti were observed in OV-TRα/βKO cochlea, resembling those described previously

in developmental hypothyroidism, or in mice with global deletion of TH transporters (4, 35, 36).

To further evaluate root cell viability, we then quantified the number of root cell's nucleus

at 2 months. We and others have previously reported that Sox10 is localized in the nucleus of

cochlear non-sensory cells including outer border, root cells, spiral prominence epithelium but

is excluded from fibrocytes in the lateral wall (20, 37). Therefore, by performing double immune-

staining, root cells could be easily distinguished as Sox10 and acetyl-α-tubulin (intensely

expressed in the root cell cytosol) double positive cells in the spiral ligament (Fig. 4D).

Consistent with the histological observations, the number of root cell's nucleus was significantly

decreased in OV-TRα/βKO but not in OV-TRβKO, compared to that in WT, supporting root cell

degeneration upon the absence of both TRs (Fig. 4D-E). Thus, TRs are necessary and also

function redundantly for root cell survival at this age.

Stria vascularis remains intact upon deletion of all known TRs in cochlea.

As OV-TRα/βKO showed a reduced EP compared with control mice (See Fig. 2E), we then

examined the structural and functional integrity of SV, which plays an essential role in

generating EP and maintaining K^+ homeostasis in the endolymph(38–40). No obvious

histological abnormality was observed in either OV-TRβKO or OV-TRα/βKO SV at P21 (Fig.

5A). The area of SV from both mice lines was comparable to that from WT (Fig. 5B).

Consistently, immuno-staining analysis showed that both groups showed normal expression

patterns of EP-generating ion-transport molecules including KCNJ10 (Kir4.1, a marker for

intermediate cells)(41, 42), Na^+/K^+-ATPase α(43), NKCC1(44) and KCNQ1 (Kv7.1)(45, 46)

(markers for marginal cells) (Fig. 5C). These data suggest that SV remains intact upon all TRs

deletion in cochlear epithelium. Indeed, potential primary defects can be excluded from SV

intermediate cells and basal cells in either OV-TRβKO or OV-TRα/βKO mice, as these two cell

types are not originated from the otic epithelium(47) and thus should display no Cre

recombination activity (see Fig. 2*A*).

As gap junctions are also essential for EP generation and hearing(48, 49). We then

analyzed the expression of Connexin26 (Cx26) and Cx30 proteins, which are the dominant

gap-junction channel-forming proteins expressed in cochlear epithelium(50). Comparable

expression patterns of Cx26 and Cx30 were observed between control and OV-TRα/βKO

cochleae at P21 (*SI Appendix* Fig. S4), suggesting gap junction system remains intact upon

TRs deficiency.

TH signaling acts in parallel with ZBTB20 in the outer sulcus development.

We have recently reported that transcription factor ZBTB20 is essential for root cell

differentiation and maturation in mice(20). We therefore asked whether TH acts upstream of

ZBTB20 in regulating root cell differentiation. To test if TH regulates ZBTB20 expression in the

outer sulcus, we compared ZBTB20 protein levels in the drug-induced hypothyroid cochlea with

that in euthyroid during postnatal development stage. Western blot showed that ZBTB20 protein

levels were significantly decreased in the extracts of hypothyroid cochleae at P4 compared with

those in euthyroid controls (Fig. 6*A*, and *SI Appendix* Fig. S5). Immunostaining revealed that

the decrease in ZBTB20 expression was mainly detected in the GER epithelial cells of

hypothyroid cochleae at P5, while its expression in the rest of the cochlear epithelium including

supporting cells and the outer sulcus/root cells was comparable with those in euthyroid controls

(Fig. 6*B-D*). Consistently, comparable ZBTB20 expression was observed between OV-

TRα/βKO and WT lateral wall at P10 (*SI Appendix* Fig. S6). These data suggest that ZBTB20

expression in outer sulcus/root cells is not TH-dependent although its expression in the GER

may be regulated by TH.

We then tested if ZBTB20, instead, functions upstream of TH signaling pathway. OV-

ZB20KO mice had normal serum levels of tri-iodothyronine (T3), and thyroxine (T4) at P10 (*SI*

Appendix Fig. S7A-B). Quantitative RT-PCR analyses showed that OV-ZB20KO cochleae

expressed comparable mRNA levels of thyroid hormone transporters at P3 (*SI Appendix* Fig.

S7C), as well as TH receptors including *TRα1*, *TRβ1* and *TRβ2* at P10 (*SI Appendix* Fig. S7D).

In situ hybridization of cochleae revealed no significant difference in the mRNA expression

pattern of genes encoding type 2 and type3 deiodinases (*Dio2* and *Dio3*) (*SI Appendix* Fig.

S7E), or *TRα1* and *TRβ1* between the two genotypes (*SI Appendix* Fig. S7F). Thus ZBTB20

unlike acts upstream of TH signaling pathway.

Together, these results suggest that TH signaling may acts in parallel with ZBTB20 in

regulating the outer sulcus remodeling.

Discussion

This study demonstrates a key role for TH and TRs in postnatal root cell development and

survival. Absence of both TRα and TRβ but not TRβ alone in cochlear epithelium resulted in

severe retardation in root cell differentiation and an early-onset root cell degeneration,

suggesting that TRα and TRβ act redundantly in promoting root cell late-stage differentiation

and maitaining cellular homeostasis.

The critical period when TH is essential for cochlear development and normal hearing

covers from late embryonic stages to around the second postnatal week in rodents(14, 26, 51).

During this developmental window, cochlear sensory epithelium undergoes dramatic

morphological changes including regression of GER leading to the formation of inner sulcus,

and opening of the tunnel of Corti (14, 52). Concomitantly, the outer sulcus epithelium

transformes from a simple layer of cuboidal-shaped cells lining the scala media into root cells

with large, branched processes resided deep in the spiral ligament(18, 20, 52). This study

uncovers that TH or TRs is nessesary for the outer sulcus remodeling, in addition to previous

reported GER regression and the organ of Corti maturation(6). Although both TRα and TRβ are

expressed by the developing outer sulcus(21–24), severe retardation in root cell differentiation

was observed only in the absence of both receptors indicating that TRα and TRβ compensate

functionally for each other in controlling root cell development. Our findings extend those data

obtained from the mouse model with global deletion of TRs, which suggest synergistic effects

of TRα and TRβ in controlling the remodeling of the organ of Corti and the correct formation of

tectorial membrane (6).

Another intrigueing finding of this study is the early-onset progressive degeneration of root

cells observed only in OV-TRα/βKO but not OV-TRβKO cochlea, suggesting TRα and TRβ play

a combined role in root cell survival. Interestingly, degeneration of root cells and the organ of

Corti initiated concomitantly in juvenile OV-TRα/βKO cochlea, indicating potential coordinated

mechanisms underlying these degenerations. As root cells with their extensive processes are

closely associated with capillaries and fibrocytes within the spiral ligament, it has been sugested

that root cells may provide a cellular route for transporting nutrients, blood-borne hormones

(such as T3), metabolites and ions between the vasculature in the spiral ligament into the infeior

cochlear sensory epithelium(13, 17, 19), which lacks direct blood flow. This hypothesis is

supported by an epithelial gap junction network formed by the syncytium of supporting cells in

the organ of Corti and extending through the Claudius' cells and root cells(53, 54). It is thus

reasonable to assume that the proper timely development of root cells may help to maintain the

health of cochlear sensory epithelium. The early-onset degeneration of root cells might

comprimise the nutrients and hormones supplement, or ion and metabolite homeostasis in the

organ of Corti, promoting its degeneration. Alternatively, root cells health might be affected by

the degeneration of organ of Corti. Severe early-onset degenerations of cochlear hair cells have

been reported previously in congenital hypothyroidism, or in mice with global deletion of thyroid

hormone transporters (4, 36), though the status of root cells in these animal models have not

been reported. It would be worth to explore any alterations in root cells structure or function in

these mice.

Several lines of evidence including our recently published report suggests that root

cells may play important roles in the maintenance of endolymph homeostasis and EP(9,

12, 10, 20). Our findings in this study also support this hypothesis and imply that root cell

dysfunction may at least partially contribute to a reduced EP upon deficiency of TH or all

known TRs during the critical period of cochlear development (4, 6, 36).

The EP is generated and maintained mainly by the SV (38, 55–57). Although SV

defects have been reported in a secondary hypothyroidism caused by pituitary transcription

factor 1 (Pit1) mutant (4), no obvious SV alterations have been reported in either TH

transporters deficiency (36), or global *Thra* and *Thrb* double knockout mice(6), both of which

display a significant reduction in EP, indicating defects beyond SV might contributing to the EP

reduction in these mice models. Consistently, our findings also show a reduced EP without

obvious defects in SV upon deficiency of both TRα and TRβ in cochlear epithelium. Indeed,

Thrb mRNA is not detected in the developing SV, as revealed previously using a sensitive *LacZ*

gene reporter system(21).Thus the SV is unlike a primary defect site responsible for a reduced

EP, which occurred only in mice lacking all TRs(6). As mentioned above, root cells are

located at the lateral border of cochlear epithelial gap junction network adjacent to type II

fibrocytes and have been suggested a role in K^+ recycling(58). This hypothesis is supported by

a report showing deficiency of KCNK5 channel, which is mostly expressed by the outer sulcus,

results in profound deafness associated with a decreased EP(10). Consistently, our recent

report showed that disruption of outer sulcus development upon ZBTB20 deletion results in EP

reduction(20). Therefore, our results in this study are consistent with a role for TH and TRs in

the development and function of root cells.

EP reduction observed in juvenile OV-TRα/βKO mice appears to be less severe than that

reported in adult global *Thra* and *Thrb* double knockout mice(6). This discrepancy may be due

to that EP was recorded at a younger age in this study, as root cell degeneration may be

progressive, causing further decrease in EP magnitude. Alternatively, global deletion of both

Thra and *Thrb* may cause defects in other cell types such as fibrocytes in the lateral wall, which

also play a role in EP maintenance(59, 60). In OV-TRα/βKO mice, however, primary fibrocyte

defects can be excluded as the Cre recombination activity was confined in otocyst-derived

cochlear epithelium, as shown in Fig.1A. On the other hand, the average ~15 mV of EP

reduction might be an underestimate of the real change in EP in these mice. Previous studies

have shown that the magnitude of EP also depends on the status of mechanotransduction of

hair cells since mechanotransduction in the stereocilia acts like an adjustable resistor(61, 62).

Reduction in mechanotransduction and/or loss of hair cells can lead to an increase in EP

value(63, 64). The severely distorted TM in OV-TRα/βKO mice would cause the detachment of

the TM from stereocilia bundles of outer hair cells, resulting in more mechanotransduction

channels to be closed at rest, and thus increasing EP value to some extent. Therefore, the 15-

mV reduction of EP seen in these mice is the result of two opposite actions. It is thus highly

likely that the contribution of the outer sulcus to EP may be underestimated in juvenile OV-

TRα/βKO mice.

Finally, our results reveal that TH signaling regulates root cell differentiation mostly by

acting in parallel with ZBTB20, which we have recently demonstrated as an essential

transcription factor governing the late-stage differentiation of the outer sulcus region(20). The

root cell differentiation was permanently disrupted upon *Zbtb20* ablation, while was severely

retarded but had progressed to some extent around 1.5 months in OV-TRα/βKO mice (*SI

Appendix* Fig. S2). This evidence let us to propose that the proper development of the outer

sulcus requires a coordination between TRs and ZBTB20 or other transcriptional pathways.

Materials and Methods

Animal models

All animal maintenance and experiments were conducted in accordance with the guidelines of the Naval Medical University Animal Ethics Committee. *Thra* [flox](30), *Thrb*[flox] (29), OV-ZB20KO(20), or ROSA26-*LacZ* reporter (65)mice have been described previously. The OV-TRβKO mice were generated by cross-breeding of *Thrb* [flox] mice with *Foxg1-Cre* mice (Jax, Stock No. 004337)(66). The OV-TRα/βKO mice were then generated by cross-breeding of heterozygous OV-TRβKO mice with *Thra* [flox] mice. All mice were maintained on a C57BL/6J background. Floxed/*Cre*-negative littermate mice were served as wild-type controls. PCR primers for genotyping are listed in *SI Appendix* Table S1. Hypothyroidism induction：Timed-pregnant C57BL/6J mice were treated with either antithyroid agents (0.05% methimazole, 1.0% sodium perchlorate monohydrate and 5.0% sucrose in drinking water)(25) or vehicle from embryonic day 17. The treatment was continued during lactation and stopped until the day indicated.

Histological analysis

Mice were transcardially perfused with 4% paraformaldehyde (pH 7.4). Dissected cochleae were then perfused perilymphatically with 2.5% glutaraldehyde and fixed overnight, followed by post-fixed with 1% osmium tetroxide. Cochleae were then decalcified, dehydrated and embedded in Spurr's low-viscosity resin. Mid-modiolar sections (1 μm) were stained with 1.25 % (w/v) toluidine blue. For quantitative analysis, the upper basal turns from three non-consecutive sections were photographed at a 200x magnification and SV areas were measured using

Image-Pro Plus 6.0 program. A single cochlea from each animal and n ≥ 5 mice per genotype

were analyzed at a given age.

Immunohistochemistry and X-Gal staining

Immunostaining was performed essentially as previously described (20). X-gal staining was

done as previously published (67). The information for primary antibodies are listed in *SI*

Appendix Table S2. To quantify root cells, non-consecutive mid-modiolar sections per cochlea

were subjected to double immunostaining for Sox10 and acetyl-α-tubulin. The nuclei of double

positive cells within the outer sulcus region from at least 5 sections were counted manually,

blind to genotype.

For quantitative analysis of ZBTB20 immunostaining intensity, more than three non-

consecutive mid-modiolar paraffin-embedded sections for each cochlea (single ear was

analyzed for each animal) from hypothyroid and euthyroid mice at P5 were stained in parallel

with anti-ZBTB20 antibody. The images were acquired at X400 magnification by a camera

(DP80; Olympus) equipped with a fluorescence microscope (BX53; Olympus). All imaging

parameters were kept at the same for each group. The averaged staining intensities in nuclei

of cells from defined regions (greater epithelial ridge (GER), supporting cells (SC), outer sulcus

epithelium (OS) and hair cells (HC)) at the upper-basal turn were numerically quantified by the

ImageJ program. The mean intensities of each sub-region were divided by those of HC at the

same turn to get normalized mean intensities, as HCs did not express ZBTB20 at this stage

thus could serve as the internal negative control.

Auditory brainstem response (ABR) test

ABRs were recorded using a TDT RZ6/BioSigRZ system (Tucker-Davis Tech. Inc., US) as

previously described (20). Briefly, anesthetized mice were kept on a heated mat in a sound-

proof chamber. Click or pure tone stimuli from 4 to 32 kHz were generated. The sound intensity

was decreased from 90 to 10 dB sound pressure level (SPL) in 5 dB steps and auditory

thresholds were defined as the lowest sound intensity evoking reproducible and recognizable

waves.

Endocochlear potential (EP) measurement

EP recording through cochlear lateral wall was previously described (20, 68). The basal

turns from both ears were analyzed for each animal. In brief, mice were anesthetized and

tracheotomy was performed. The tympanic bulla was opened to display the cochlea. A small

hole was drilled in cochlear lateral wall of the basal turn, through which a micropipette electrode

filled with 150 mM KCl was advanced into the scala media using a micromanipulator. The

voltage changes during the insertion were continuously recorded using an amplifier

(MultiClamp 700B, Molecular Devices) combined with a digitizer (Digidata 1440A, Molecular

Devices).

In situ hybridization.

In situ hybridization was performed as described(20). In brief, mice were transcardially

perfused with 4% paraformaldehyde (pH 9.0) and cochleae were dissected and perfused

perilymphatically with the same fixative. The fixed and decalcified cochleae were cryostat

sectioned (14 μm thickness). Sections were then subjected to in situ hybridization as described

previously(69). Information of antisense cRNA probes are listed in *SI Appendix* Table S3.

Cochleae from n ≥ 4 mice per genotype at a given age were analyzed.

Real-time RT-PCR

Total RNA extraction, cDNA synthesis and quantitative Real-time PCR was performed as described previously(20). Data were normalized to 36B4 (Rplp0, internal control) and presented as fold changes of gene expression in the mutant mice compared to the WT Control. Each sample was analyzed in duplicate. The primers used are shown in *SI Appendix* Table S1.

Measurement of thyroid hormones.

Blood samples were collected and separated into serum aliquots by centrifugation and stored at −80ºC until assayed. Serum levels of T3 and T4 were measured using T3 ELISA kit from MLBIO (Cat. #m1559442) and T4 CLIA kit from cloud-clone Corp. (Cat. #CCA452Ge), according to the manufacturers' instructions.

Western blot

Total proteins of cochleae from 2 ears per animal were electrophoretically separated on 4–15% gradient SDS–PAGE gels and transferred onto nitrocellulose membranes. Membranes were probed with the following primary antibodies: anti-Zbtb20 monoclonal antibody clone 9A10 (1:1000, home-made)(67); anti-histone H3 (1:5,000, Cell Signaling Technology). Secondary antibodies were either HRP-conjugated anti-rabbit or anti-mouse IgG (1:8,000, Vector Laboratories). Signals were then developed using a Chemiluminescent Detection Kit (Thermo Fisher, #34580). The blot images were acquired by Luminescent Image Analyzer (Las 4000, GE Healthcare) and the densities of ZBTB20 bands were quantified and normalized to those of internal control.

Statistical analysis

All statistical analysis was performed using GraphPad Prism 7 software. Values are shown

as means ±SEMs. The data normality tests were first carried out, and then a two-tailed

Student's t test and one-way or two-way ANOVA were used to compare the means between

two groups and among three groups, respectively. $p \leq 0.05$ was considered statistically

significant. The non-significant difference between samples is indicated as "ns"; p values of

significant differences are shown in the graphs.

Data availability: Data supporting the findings of this study are available within the article and

its Supplementary information files.

Acknowledgements. This work was supported by grants from the National Key R&D Program

(2019YFA0802500, 2018YFA0800602), National Natural Science Foundation of China

(32271162), and Collaborative Innovation Program of Shanghai Municipal Health Commission

(2020CXJQ01).

Reference

1.   L. Ng, M. W. Kelley, D. Forrest, Making sense with thyroid hormone--the role of T(3) in
auditory development. *Nat Rev Endocrinol* **9**, 296–307 (2013).
2.   S. Christ, *et al.*, Hearing loss in athyroid pax8 knockout mice and effects of thyroxine
substitution. *Audiol Neurootol* **9**, 88–106 (2004).
3.   M. S. Deol, Congenital deafness and hypothyroidism. *Lancet* **2**, 105–6 (1973).
4.   M. Mustapha, *et al.*, Deafness and Permanently Reduced Potassium Channel Gene
Expression and Function in Hypothyroid Pit1dw Mutants. *The Journal of Neuroscience*
29, 1212 (2009).
5.   D. Forrest, L. C. Erway, L. Ng, R. Altschuler, T. Curran, Thyroid hormone receptor beta is
essential for development of auditory function. *Nat Genet* **13**, 354–7 (1996).
6.   A. Rusch, *et al.*, Retardation of cochlear maturation and impaired hair cell function
caused by deletion of all known thyroid hormone receptors. *J Neurosci* **21**, 9792–9800
(2001).
7.   H. Winter, *et al.*, Deafness in TRbeta mutants is caused by malformation of the tectorial
membrane. *J Neurosci* **29**, 2581–2587 (2009).
8.   A. Rüsch, L. C. Erway, D. Oliver, B. Vennström, D. Forrest, Thyroid hormone receptor β-
dependent expression of a potassium conductance in inner hair cells at the onset of
hearing. *Proceedings of the National Academy of Sciences* **95**, 15758–15762 (1998).
9.   S. S. Spicer, B. A. Schulte, The fine structure of spiral ligament cells relates to ion return
to the stria and varies with place-frequency. *Hear Res* **100**, 80–100 (1996).
10.   Y. Cazals, *et al.*, KCNK5 channels mostly expressed in cochlear outer sulcus cells are
indispensable for hearing. *Nat Commun* **6**, 8780 (2015).
11.   M. Galic, W. Giebel, An electron microscopic study of the function of the root cells in
the external spiral sulcus of the cochlea. *Acta Otolaryngol Suppl* **461**, 1–15 (1989).
12.   D. J. Jagger, G. Nevill, A. Forge, The Membrane Properties of Cochlear Root Cells are
Consistent with Roles in Potassium Recirculation and Spatial Buffering. *J Assoc Res*
Otolaryngol **11**, 435–48 (2010).
13.   D. J. Jagger, A. Forge, The enigmatic root cell - emerging roles contributing to fluid
homeostasis within the cochlear outer sulcus. *Hear Res* **303**, 1–11 (2013).
14.   M. S. Deol, An Experimental Approach to the Understanding and Treatment of
Hereditary Syndromes with Congenital Deafness and Hypothyroidism. *Journal of*
Medical Genetics **10**, 235 (1973).
15.   I. J. Karolyi, *et al.*, Dietary thyroid hormone replacement ameliorates hearing deficits in
hypothyroid mice. *Mamm Genome* **18**, 596–608 (2007).
16.   A. Uziel, C. Legrand, A. Rabie, Corrective effects of thyroxine on cochlear abnormalities
induced by congenital hypothyroidism in the rat. I. Morphological study. *Brain research*
351, 111–22 (1985).
17.   A. J. Duvall III, The ultrastructure of the external sulcus in the guinea pig cochlear duct.
The Laryngoscope **79**, 1–29 (1969).
18.   D. J. Lim, M. Anniko, Developmental morphology of the mouse inner ear. A scanning
electron microscopic observation. *Acta Otolaryngol Suppl* **422**, 1–69 (1985).
19.   R. Shodo, M. Hayatsu, D. Koga, A. Horii, T. Ushiki, Three-dimensional reconstruction of
root cells and interdental cells in the rat inner ear by serial section scanning electron

microscopy. *Biomed Res* **38**, 239–248 (2017).

20.   Z. Xie, *et al.*, ZBTB20 is essential for cochlear maturation and hearing in mice.
Proceedings of the National Academy of Sciences **120**, e2220867120 (2023).

21.   L. Ng, *et al.*, Age-Related Hearing Loss and Degeneration of Cochlear Hair Cells in Mice
Lacking Thyroid Hormone Receptor beta1. *Endocrinology* **156**, 3853–65 (2015).

22.   S. Gu, *et al.*, Characterization of rare spindle and root cell transcriptional profiles in the
stria vascularis of the adult mouse cochlea. *Sci Rep* **10**, 18100 (2020).

23.   D. J. Bradley, H. C. Towle, W. S. Young, Alpha and beta thyroid hormone receptor (TR)
gene expression during auditory neurogenesis: evidence for TR isoform-specific
transcriptional regulation in vivo. *Proc Natl Acad Sci U S A* **91**, 439–443 (1994).

24.   P. Jean, *et al.*, Single-cell transcriptomic profiling of the mouse cochlea: An atlas for
targeted therapies. *Proceedings of the National Academy of Sciences* **120**,
e2221744120 (2023).

25.   O. Sawant, *et al.*, Light-Regulated Thyroid Hormone Signaling Is Required for Rod
Photoreceptor Development in the Mouse Retina. *Invest Ophthalmol Vis Sci* **56**, 8248–
8257 (2015).

26.   A. Uziel, C. Legrand, A. Rabie, Corrective effects of thyroxine on cochlear abnormalities
induced by congenital hypothyroidism in the rat. I. Morphological study. *Brain Res* **351**,
111–122 (1985).

27.   B. J. Walters, J. Zuo, Postnatal development, maturation and aging in the mouse cochlea
and their effects on hair cell regeneration. *Hear Res* **297**, 68–83 (2013).

28.   W. Liu, C. Wang, H. Yu, S. Liu, J. Yang, Expression of acetylated tubulin in the postnatal
developing mouse cochlea. *European journal of histochemistry : EJH* **62** (2018).

29.   Y. Yan, *et al.*, Hepatic thyroid hormone signalling modulates glucose homeostasis
through the regulation of GLP-1 production via bile acid-mediated FXR antagonism.
Nat Commun **13**, 6408 (2022).

30.   S. Liu, *et al.*, Triiodothyronine (T3) promotes brown fat hyperplasia via thyroid hormone
receptor α mediated adipocyte progenitor cell proliferation. *Nat Commun* **13**, 3394
(2022).

31.   J. M. Hebert, S. K. McConnell, Targeting of cre to the Foxg1 (BF-1) locus mediates loxP
recombination in the telencephalon and other developing head structures. *Dev Biol* **222**,
296–306 (2000).

32.   M. H. Song, *et al.*, Pou3f4 deficiency causes defects in otic fibrocytes and stria vascularis
by different mechanisms. *Biochemical and Biophysical Research Communications* **404**,
528–533 (2011).

33.   I. E. Royaux, *et al.*, Localization and functional studies of pendrin in the mouse inner ear
provide insight about the etiology of deafness in pendred syndrome. *J Assoc Res*
Otolaryngol **4**, 394–404 (2003).

34.   R. S. Kimura, Sensory and accessory epithelia of the cochlea. *Friedmann, I And J*
Ballantyne (Ed) Ultrastructural Atlas Of The Inner Ear Ix++329p Butterworths, London,
England; Boston, Mass , Usa Illus P101-132 (1984).

35.   A. Uziel, C. Legrand, M. Ohresser, M. Marot, Maturational and degenerative processes
in the organ of Corti after neonatal hypothyroidism. *Hear Res* **11**, 203–218 (1983).

36.   D. S. Sharlin, *et al.*, Deafness and loss of cochlear hair cells in the absence of thyroid

hormone transporters Slc16a2 (Mct8) and Slc16a10 (Mct10). *Sci Rep* **8**, 4403 (2018).

37.   X. Hao, *et al.*, Sox10 expressing cells in the lateral wall of the aged mouse and human
cochlea. *PLoS One* **9**, e97389 (2014).

38.   H. Hibino, F. Nin, C. Tsuzuki, Y. Kurachi, How is the highly positive endocochlear potential
formed? The specific architecture of the stria vascularis and the roles of the ion-
transport apparatus. *Pflugers Arch* **459**, 521–33 (2010).

39.   H. J. Kim, *et al.*, Precise toxigenic ablation of intermediate cells abolishes the "battery"
of the cochlear duct. *J Neurosci* **33**, 14601–14606 (2013).

40.   P. Wangemann, Supporting sensory transduction: cochlear fluid homeostasis and the
endocochlear potential. *J Physiol* **576**, 11–21 (2006).

41.   M. Ando, S. Takeuchi, Immunological identification of an inward rectifier K+ channel
(Kir4.1) in the intermediate cell (melanocyte) of the cochlear stria vascularis of gerbils
and rats. *Cell Tissue Res* **298**, 179–183 (1999).

42.   D. C. Marcus, T. Wu, P. Wangemann, P. Kofuji, KCNJ10 (Kir4.1) potassium channel
knockout abolishes endocochlear potential. *Am J Physiol Cell Physiol* **282**, C403-407
(2002).

43.   T. P. Kerr, M. D. Ross, S. A. Ernst, Cellular localization of Na+,K+-ATPase in the
mammalian cochlear duct: significance for cochlear fluid balance. *Am J Otolaryngol* **3**,
332–338 (1982).

44.   M. J. Dixon, *et al.*, Mutation of the Na-K-Cl Co-Transporter Gene Slc12a2 Results in
Deafness in Mice. *Human Molecular Genetics* **8**, 1579–1584 (1999).

45.   M. C. Casimiro, *et al.*, Targeted disruption of the Kcnq1 gene produces a mouse model
of Jervell and Lange-Nielsen Syndrome. *Proc Natl Acad Sci U S A* **98**, 2526–2531 (2001).

46.   M. Sakagami, *et al.*, Cellular localization of rat Isk protein in the stria vascularis by
immunohistochemical observation. *Hear Res* **56**, 168–172 (1991).

47.   K. P. Steel, C. Barkway, Another role for melanocytes: their importance for normal stria
vascularis development in the mammalian inner ear. *Development* **107**, 453–463 (1989).

48.   M. Cohen-Salmon, *et al.*, Targeted Ablation of Connexin26 in the Inner Ear Epithelial
Gap Junction Network Causes Hearing Impairment and Cell Death. *Current Biology* **12**,
1106–1111 (2002).

49.   J. Sun, *et al.*, Cochlear gap junctions coassembled from Cx26 and 30 show faster
intercellular Ca2+ signaling than homomeric counterparts. *American Journal of*
Physiology-Cell Physiology **288**, C613–C623 (2005).

50.   S. Ahmad, S. Chen, J. Sun, X. Lin, Connexins 26 and 30 are co-assembled to form gap
junctions in the cochlea of mice. *Biochemical and Biophysical Research Communications*
307, 362–368 (2003).

51.   A. Uziel, Periods of sensitivity to thyroid hormone during the development of the organ
of Corti. *Acta Otolaryngol Suppl* **429**, 23–27 (1986).

52.   B. Roth, V. Bruns, Postnatal development of the rat organ of Corti. I. General morphology,
basilar membrane, tectorial membrane and border cells. *Anat Embryol (Berl)* **185**, 559–
69 (1992).

53.   D. J. Jagger, A. Forge, Connexins and gap junctions in the inner ear--it's not just about
K(+) recycling. *Cell Tissue Res* **360**, 633–44 (2015).

54.   D. J. Jagger, A. Forge, Compartmentalized and signal-selective gap junctional coupling

in the hearing cochlea. *J Neurosci* **26**, 1260–8 (2006).

55.   F. Nin, *et al.*, The endocochlear potential depends on two K+ diffusion potentials and
an electrical barrier in the stria vascularis of the inner ear. *Proceedings of the National*
Academy of Sciences **105**, 1751–1756 (2008).

56.   G. Rickheit, *et al.*, Endocochlear potential depends on Cl- channels: mechanism
underlying deafness in Bartter syndrome IV. *The EMBO journal* **27**, 2907–17 (2008).

57.   P. Wangemann, J. Liu, D. C. Marcus, Ion transport mechanisms responsible for K+
secretion and the transepithelial voltage across marginal cells of stria vascularis in vitro.
Hearing Research **84**, 19–29 (1995).

58.   S. S. Spicer, B. A. Schulte, The fine structure of spiral ligament cells relates to ion return
to the stria and varies with place-frequency. *Hear Res* **100**, 80–100 (1996).

59.   N. Adachi, *et al.*, The mechanism underlying maintenance of the endocochlear potential
by the K+ transport system in fibrocytes of the inner ear. *J Physiol* **591**, 4459–4472
(2013).

60.   D. N. Furness, Forgotten Fibrocytes: A Neglected, Supporting Cell Type of the Cochlea
With the Potential to be an Alternative Therapeutic Target in Hearing Loss. *Front Cell*
Neurosci **13**, 532 (2019).

61.   H. Davis, A model for transducer action in the cochlea. *Cold Spring Harb Symp Quant*
Biol **30**, 181–190 (1965).

62.   I. J. Russell, Origin of the receptor potential in inner hair cells of the mammalian cochlea-
-evidence for Davis' theory. *Nature* **301**, 334–336 (1983).

63.   H. Liu, *et al.*, Organ of Corti and Stria Vascularis: Is there an Interdependence for Survival?
PLoS One **11**, e0168953 (2016).

64.   H. Liu, *et al.*, Molecular and cytological profiling of biological aging of mouse cochlear
inner and outer hair cells. *Cell Rep* **39**, 110665 (2022).

65.   P. Soriano, Generalized lacZ expression with the ROSA26 Cre reporter strain. *Nat Genet*
21, 70–71 (1999).

66.   J. M. Hebert, S. K. McConnell, Targeting of cre to the Foxg1 (BF-1) locus mediates loxP
recombination in the telencephalon and other developing head structures. *Dev Biol* **222**,
296–306 (2000).

67.   Z. Xie, *et al.*, Zinc finger protein ZBTB20 is a key repressor of alpha-fetoprotein gene
transcription in liver. *Proc Natl Acad Sci U S A* **105**, 10859–10864 (2008).

68.   Y. Li, H. Liu, X. Zhao, D. Z. He, Endolymphatic Potential Measured From Developing and
Adult Mouse Inner Ear. *Front Cell Neurosci* **14**, 584928 (2020).

69.   Z. Xie, *et al.*, Zbtb20 is essential for the specification of CA1 field identity in the
developing hippocampus. *Proc Natl Acad Sci U S A* **107**, 6510–6515 (2010).

Figures and figure legends

Fig. 1. Hypothyroidism causes the disruption in the outer sulcus development. (A and B) Representative Toluidine Blue-stained sections of cochlear upper basal turns from euthyroid (**A**) or methimazole (MMI)-induced hypothyroid mice (**B**) at P21. (**A**) Euthyroid cochlea showing a mature outer sulcus characterized by root cells (Rc. indicated by red arrows) with numerous bright branched processes extending deep into the spiral ligament and upward to behind the spiral prominence (Sp). The surfaces of root cell bodies were covered by enlarged Claudius cells (Cc, red bar indicating its height). Co, the organ of Corti; Is, inner sulcus; Sm, scala media; Sv, stria vascularis; Tc, tunnel of Corti, Tm, tectorial membrane. (**B**) The hypothyroid outer sulcus was consist of a simple layer of cells with a few short processes bordering the scala media, continued with the flattened Claudius cells. Red arrow heads indicate defects including a distorted Tm, an abnormal persistence of GER, closure of tunnel of Corti, and a flattened border region. (**C** and **D**) Immunostaing showing extensive acetyl-α-tubulin positive (red) root cell processes extending into the inferior region of euthyroid spiral ligament at P21 (**C**) but only a few short processes observed in the luminal surface of hypothyroid counterparts (indicated by white arrow head) (**D**). Bars = 50 μm for all panels. Nuclei were stained with DAPI (turquoise). n = 4 mice/ group. Scale bars in all panels = 50 μm. Abbreviations are consistent across figures in this manuscript.

Fig.2 Cochlea-selective deletion of all known TRs causes deafness and a compromised EP value. (A) X-gal staining showing *Foxg1-Cre* recombinase activity was specifically confined in cochlear epithelium lining the scala media (Sm) at P5, including GER, LER (the lesser epithelial ridge), root cells (Rc), stria vascularis (Sv), as well as spiral ganglions (Sg). n = 3 mice/group. Bar = 100 μm. **(B)** QRT-PCR analysis showing relative mRNA expression levels of *Thra* and *Thrb* in control, OV-TRβKO, or OV-TRα/βKO cochlea at P10. n = 4 mice/group. **(C)** Representative ABR waveforms obtained to broadband click stimuli from control, OV-TRβKO and OV-TRα/βKO mice at 2 months (mon) of age. Sound pressure levels (SPL) were indicated in decibels (dB). **(D)** Averaged thresholds of Click- and Pure tone-evoked ABRs from control, OV-TRβKO and OV-TRα/βKO mice at 2 mon. n = 5 mice/group. **(E)** OV-TRα/βKO mice showing a reduced endocochlear potential (EP) in the basal cochlear turns compared with control mice at around 1.5 mon. The number of animals tested was indicated. All values are represented as mean ± SEM. A two-tailed Student's t-test was used for compare differences in relative mRNA expression levels, or EP values. A Kruskal-Wallis test (a nonparametric version of the ANOVA) with Dunn's multiple comparisons test and a two-way ANOVA followed by post-hoc pairwise tests (Bonferroni method) were used to compare differences in Click- and Pure tone-evoked ABR thresholds, respectively. *P* Values were indicated in each graphs. ns, not significant.

Fig. 3. Cochlea-selective deletion of all known TRs retards root cell development. **(A-C)** Representative Toluidine Blue-stained sections showing developing root cells (Rc) infiltrating deep into the spiral ligament(indicated by red arrows) from control **(A)** or OV-TRβKO **(B)** but mainly present in the luminal surface of scala media (Sm) in OV-TRα/βKO pups at P10 **(C)**. **(D-F)** Immunostaining showing extensive acetyl-α-tubulin positive (red) root cell processes within control **(D)** or OV-TRβKO **(E)** spiral ligament at P10, whereas only a few detected in OV-TRα/βKO counterparts **(F).** Nuclei were stained with DAPI (turquoise). n = 4 mice/ group. **(G-I)** In situ hybridization revealing that expression domain of *Pendrin* mRNA was extended into the inferior region of the spiral ligament in control **(G)** or OV-TRβKO **(H)** but were confined in the luminal surface in OV-TRα/βKO counterparts at P10 **(I)** . **(J-L)** Representative sections showing a mature outer sulcus in control **(J)** or OV-TRβKO **(K)** but not in OV-TRα/βKO **(L)** mice at P21. Both of control and OV-TRβKO mice showed elaborate root cell branches within the spiral ligament, the surfaces of which covered by the enlarged Claudius cells (Cc, outlined by black dots) and the spiral prominence epithelium (Sp, outlined by magenta dots) **(J and K),** whereas OV-TRα/βKO root cells (indicated by red bracket) were mainly bordering the scala media, separating a flattened Claudius cell (indicated by arrow head) and the spiral prominence epithelium. A cell with a pale-staining cytoplasm and nucleus (indicated by De) was detected in OV-TRα/βKO outer sulcus. The upper basal turns were shown for all panels and defects were indicated by arrow heads. Scale bars: 50 μm for all panels. n = 4 mice/group. See also *SI Appendix* Fig. S1and S2.

682
Fig. 4. Absence of all known TRs in cochlear epithelium causes early-onset root cell
degeneration. (A-C) Representative Toluidine Blue-stained sections of upper basal turns
from control **(A)**, OV-TRβKO **(B)**, and OV-TRα/βKO **(C)** cochleae at 2 months. Tectorial
membrane (Tm, indicated by magenta arrow heads) malformations were observed in OV-
TRβKO and OV-TRα/βKO, whereas a partially collapsed organ of Corti (Co) was only
observed in OV-TRα/βKO mice, indicated by the red arrow in **(C)**. **(A')**, **(B')** and **(C')** are
magnified views of outer sulcus regions indicated by red brackets in **(A)**, **(B)**, and **(C)**,
respectively. Both of control and OV-TRβKO mice showed numerous branched root cells
(Rc) within the spiral ligament, whereas OV-TRα/βKO counterparts showed reduced root
cells with pale-staining processes and nuclei, indicated by red arrow head in (C'). Sp, Spiral
prominence. n = 6 mice/group. **(D)** Double immunostaining showing Sox10 (magenta) and
acetyl-α-tubulin (green) double-positive root cells within upper basal and apical turns from
control, OV-TRβKO, or OV-TRα/βKO mice at 2 months. Nuclei were stained with DAPI
(blue). n = 6 mice/group. (E) Bar graph showing reduced averaged number of Sox10 and
acetyl-α-tubulin double-positive root cells in OV-TRα/βKO compared with that in control
mice at 2 months. All values are represented as mean ±SEM. A two-way ANOVA followed
by post-hoc pairwise tests (Bonferroni method) was used to compare differences. *P* Values
were indicated in the graph. ns, not significant. Scale bars: 50 μm for all panels. See also
SI Appendix Fig. S3.

Fig. 5. Stria vascularis (SV) remains intact upon deletion of all known TRs in cochlea.
(**A**) Representative Toluidine Blue-stained sections of cochlear upper basal turns showing
comparable SV histology among Control, OV-TRβKO, and OV-TRα/βKO mice at P21. (**B**)
Bar graph showing comparable SV areas among Control, OV-TRβKO, and OV-TRα/βKO
mice at P21. n = 6 mice/group. Values are represented as mean ±SEM. A one-way ANOVA
followed by post-hoc pairwise tests (Bonferroni method) was used to compare differences
between the two groups. ns: not significant. (**C**) Immunostaining revealing comparable
expressions of Kir4.1, Na+/K+-ATPase α, NKCC1, and KCNQ1 (red) among Control, OV-
TRβKO, and OV-TRα/βKO mice at P21. n = 3 mice/group for each molecular marker. Scale
bars: 20 µm for all panels. Nuclei were stained with DAPI (turquoise). See also *SI Appendix*
Fig. S4.

Fig. 6. TH signaling acts in parallel with ZBTB20 in the outer sulcus development. (A)
Representative blot showing ZBTB20 protein was down regulated in cochleae from hypothyroid
mice at P4 (left panel). Three isoforms of ZBTB20 can be recognized in each sample. Histone
H3 is shown as a loading control. Quantitation of relative ZBTB20 expression is represented in
the right panel, n = 7 mice/group. **(B)** Immunostaining showing ZBTB20 expression was down-
regulated in GER (indicated by arrowhead) from hypothyroid cochlea, but was comparable in
other cochlear epithelia including supporting cells (Sc) and outer sulcus (Os)/root cell (Rc),
when compared to those from euthyroid counterparts at P5. Scale bar = 50 μm.   **(C)**
Representative image showing the quantification of ZBTB20 immunostaining intensity in
indicated regions of the upper basal turn using Image J program (left panel). **(D)** Bar graph
showing relative mean intensities of each sub-region normalized by those of hair cells (Hc,
serve as the internal negative control) at the same turn. n = 7 mice/group. Values are expressed
as mean ± SEM. Paired Student's t-test was used to compare the means of the two groups. ns,
not significant. Uncropped western blot/gel images are shown in *SI Appendix* Fig. S5. See also
SI Appendix Fig. S6 and S7.