Rapid evolution of promoters from germline-specifically expressed genes including transposon silencing factors

DAVID W. J. MCQUARRIE[1,2], AZAD ALIZADA[3], BENJAMIN CZECH NICHOLSON[3] AND MATTHIAS SOLLER[1,2*]

[1]School of Biosciences, College of Life and Environmental Sciences, University of Birmingham, Edgbaston, Birmingham, B15 2TT, United Kingdom

[2]Birmingham Centre for Genome Biology, University of Birmingham, Edgbaston, Birmingham, B15 2TT, United Kingdom

[3]Cancer Research UK Cambridge Institute, University of Cambridge, Li Ka Shing Centre, Cambridge CB2 0RE, United Kingdom

[*]Corresponding author: m.soller@bham.ac.uk

Running title: Evolution of promoters of germline transposon silencing factors

Abstract

Background: The piRNA pathway in animal gonads functions as an 'RNA-based

immune system', serving to silence transposable elements and prevent inheritance of novel

invaders. In *Drosophila*, this pathway relies on three gonad-specific Argonaute proteins

(Argonaute-3, Aubergine and Piwi) that associate with 23-28 nucleotide piRNAs, directing the

silencing of transposon-derived transcripts. Transposons constitute a primary driver of genome

evolution, yet the evolution of piRNA pathway factors has not received in-depth exploration.

Specifically, channel nuclear pore proteins, which impact piRNA processing, exhibit regions

of rapid evolution in their promoters. Consequently, the question arises whether such a mode

of evolution is a general feature of transposon silencing pathways.

Results: By employing genomic analysis of coding and promoter regions within genes

that function in transposon silencing in *Drosophila*, we demonstrate that the promoters of germ

cell-specific piRNA factors are undergoing rapid evolution. Our findings indicate that rapid

promoter evolution is a common trait among piRNA factors engaged in germline silencing

across insect species, potentially contributing to gene expression divergence in closely related

taxa. Furthermore, we observe that the promoters of genes exclusively expressed in germ cells

generally exhibit rapid evolution, with some divergence in gene expression.

Conclusion: Our results suggest that increased germline promoter evolution, in

partnership with other factors, could contribute to transposon silencing and evolution of species

through differential expression of genes driven by invading transposons.

Background

Eukaryotic organisms are continuously challenged from genomic parasites called transposable elements (TEs) [1-4]. Unchecked transposon activity often results in reduced reproductive fitness [5-8]. To negate the detrimental effects posed by TE mobilisation, vital regulatory pathways evolved that efficiently suppress transposon activity. At the centre of these are 23-28 nucleotide small RNAs, called PIWI-interacting RNAs (piRNAs). piRNAs are generated from transposons and from genomic clusters and bind to Argonaute proteins of the PIWI-clade, including P-element induced wimpy testis (Piwi). Aubergine (Aub) and Argonaute-3 (Ago3). The piRNA pathway is often referred to as an 'RNA-based immune system' thanks to its ability to adopt to new TE insertions and the memory of past transposon activity through piRNA clusters through RNA transgenerational inheritance [1, 2, 9, 10].

The piRNA pathway is arguably best understood in the ovary of *Drosophila melanogaster*. It was shown that ovaries feature two branches of the silencing machinery [5], one that is active in germ cells and a condensed version that is specific to somatic follicle cells that surround the germline [2, 11]. Somatic follicle cells only feature unistrand piRNA clusters, such as *flamenco* (*flam*), and exclusively produce piRNAs via a Zucchini (Zuc)-dependent biogenesis mechanism that takes place on the mitochondria surface involving Armitage (Armi), Papi, Vreteno (Vret), Gasz, Daedalus (Daed), Minotaur (Mino), Shutdown (Shu), and Sister of Yb (SoYb) (Fig. 1A) [3, 12, 13]. Licensing of *flam* for piRNA production takes place upstream, at so-called Yb-bodies [1, 12]. Somatic cells only express Piwi, which following the association with a mature piRNA shuttles to the nucleus. There, Piwi-piRNA complexes scan for nascent TE transcripts and instruct co-transcriptional gene silencing (cTGS) that requires components of the general chromatin silencing machinery and the piRNA pathway-specific factors Asterix

(Arx), Piwi, Panoramix (Panx), Nuclear export factor 2 (Nxf2) and Maelstrom (Mael) (Fig.

1A) [1, 2, 12, 14].

In germ cells, piRNAs predominantly originate from dual-strand clusters and are

produced by a different biogenesis mechanism called the ping-pong amplification cycle (Fig.

1B) [1-3, 5, 11, 12, 15]. Dual-strand clusters harbour remnants of transposons and are expressed

via a non-canonical transcription process that is centred on a complex containing the HP1a

homolog Rhino (Rhi), Deadlock (Del) and Cutoff (Cuff), and further requires Moonshiner

(Moon) (Fig. 1B). This Rhi-Del-Cuff complex also recruits a non-canonical export machinery

consisting of Bootlegger (Boot) and Nuclear export factor 3 (Nxf3), which transport piRNA

precursors for processing to a germ-cell specific perinuclear region termed "nuage" (Fig. 1B).

Here, the ping-pong cycle takes place and relies on the two PIWI proteins Aubergine (Aub)

and Argonaute 3 (Ago3), as well as several RNA binding proteins and Tudor-domain

containing factors including Tejas (Tej), Vasa (Vas), Qin, Squash (Squ), Tapas, Brother of Yb

(BoYb), Krimper (Krimp), and Spindle-E (Spn-E) (Fig. 1B) [1-3, 12, 15, 16].

Channel nuclear pore proteins (Nups) have been implicated in piRNA regulated

transposon silencing in *C. elegans* [17]. Interestingly, in *Drosophila*, Nups were shown to

affect transposon silencing in both the germline [16, 18], and in somatic cells of the ovary, here

specifically by contributing to the conversion of precursor RNAs from the *flam* locus into

piRNAs [19]. Recently, we found that inner and outer ring Nups in the nuclear pore complex

(NPC) evolve rapidly through indel accumulation in their promoters [20, 21]. Such promoter

indel variability in Nup54 has dominant, pleiotropic effects on sexual differentiation including

neuronal wiring important for female post-mating behaviours directed by male-derived sex

peptide, that as a result of sexual conflict could drive speciation by such mechanism [17, 20,

22]. Intriguingly, nuclear import/export pathways have been linked to speciation through

hybrid incompatibility, but are in addition important for many cellular processes beyond

piRNA processing [23, 24]. Hence, promoter evolution particularly of piRNA processing genes

could have wide impact on transposon silencing and evolution of species through differential

expression of genes, but whether regulators of germline transposon silencing generally evolve

rapidly through their promoters has not been determined.

Here, we analyse the promoters of genes involved in the ovarian piRNA pathway. We

compare piRNA factors required in somatic cells (Yb) or in both the somatic and germline

compartments (cTGS and piRNA biogenesis, Fig. 1A) with those that are germ cell-specific

(dual-strand clusters and the ping-pong cycle, Fig. 1B). We find that the promoters of germline-

specific piRNA factors are hot spots for rapid evolution compared to those of somatic and germ

cell piRNA pathway genes and the *Drosophila* genome average throughout insect species.

Analysis of genes with significant accumulation of promoter changes reveals a mix of indel

and base change accumulation. Further, our analysis reveals that rapid promoter evolution is a

general feature of genes specifically expressed in germ cells, while soma-expressed genes

evolve at a similar rate to the average of all *Drosophila* genes. Through cross-species

differential ovary RNA-seq analysis we reveal that germ cell-specific genes minimally diverge

in gene expression levels compared to genes expressed in somatic cells. Our findings highlight

that promoters of germline genes involved in transposon silencing evolve rapidly and are

accompanied by diverging gene expression, suggesting a possible mode of rapid speciation

through accumulation of changes in promoters in correlation with additional regulatory factor

divergence.

Results

Promoter regions of germline-specific piRNA factors are hot spots for rapid evolution

Since Nups have been shown to function in transposon silencing in the germline (12

from 14 tested are above the control) and display fast evolving promoters [16, 20, 21], we

analysed the rate of evolution in promoters and coding regions of piRNA pathway genes. We

separated all piRNA pathway genes based on their genetic requirement into two categories:

piRNA factors essential in somatic cells, including the soma-specific piRNA factor Yb and

piRNA factors expressed in both somatic and germline cells (cTGS and piRNA biogenesis,

Fig. 1A), and germ cell-specific piRNA proteins (dual-strand clusters and ping-pong cycle, Fig

1b). The piRNA factors essential in somatic cells include *armi*, *papi*, *vret*, *Gasz*, *daed*, *mino*,

zuc, *shu*, *SoYb*, *Yb*, *arx*, *piwi*, *panx*, *nxf2*, and *mael* (Fig. 1A). The group of piRNA factors that

are exclusively required in the germline are *aub*, *tej*, *Ago3*, *vas*, *qin*, *squ*, *tapas*, *BoYb*, *krimp*,

spn-E, Boot, *moon*, *nxf3*, *cuff*, *rhi*, and *del* (Fig. 1B) [3, 14, 25].

We defined the promoter containing region as a 350-nucleotide (-30 to -380) window

upstream of the predicted TATA box region 30 nucleotides upstream of the transcription start

site (TSS) [26]. In a first approach, we performed a genome-wide promoter evolution analysis

of germline-specific as well as somatic and germ cell piRNA factors and compared it to all

genes in the *Drosophila* genome (Fig 1C-D). We used publicly available PhyloP27way data a

conservation score. We observed a significant accumulation of changes in the germ cell-

specific piRNA factor promoter regions compared to the genome (Fig. 1C-F). Of note, the

somatic and germ cell piRNA factors evolved slower compared to the genome.

We next compared indel and base changes in the promoter regions between five closely

related *Drosophila* species (*D. melanogaster*, *D. simulans*, *D. sechellia*, *D. yakuba* and *D.*

erecta). Promoter scores (*d*) were calculated and compared to a control group comprised of the

m^6A methylation machinery genes (*Mettl3*, *Mettl14*, *fl(2)d*, *virilizer*, *flacc*, *nito* and *Hakai*,

Ythdc1 and *Ythdf*), chosen due to their high evolutionary conservation and requirement for strict

stoichiometry for functionality, making this an ideal control group to monitor promoter

evolution [21, 27, 28]. In this analysis, the group of somatic and germ cell piRNA factors

showed similar evolution rates compared to the control (Fig. 1E-J), while germ cell-specific

piRNA factors as a group evolved at a much higher rate (Fig. 1G and H), accruing a

significantly increased number of promoter-located indels (Fig. 1G and H) and base changes

(Fig. 1I and J). At the individual gene level, 7 out of 14 of germ cell-specific piRNA factors

(*Ago3* was omitted due to its low conservation between closely related *Drosophila* species [29-

31] showed significant accumulation of promoter changes (Supplementary Fig. 1A). Among

the somatic and germ cell piRNA factors, 3 out of 15 genes had significant accumulation of

promoter changes (Supplementary Fig. 1A).

To assess general evolution in the coding regions, we used McDonald-Kreitman tests

(MKTs) and analysed polymorphisms and divergence within *D. melanogaster* and between *D.*

melanogaster and *D. simulans* for two ancestral populations (Congo and Zambia) [32].

Analysis of grouped germ cell-specific piRNA factors flagged these factors as under positive

selection for both populations. Conversely, the somatic and germ cell piRNA factors were not

flagged as under positive selection, (Supplementary Fig. 1B).

Promoters of genes required for germline transposon silencing are fast evolving

Given the rapid evolution of germ cell-specific piRNA factors among closely related

Drosophila species, we wanted to determine (1) whether this is a general feature of insect

evolution, and (2) whether this is a general feature of genes required for transposon silencing

with pleiotropic roles that have been identified in a genetic screen (GTS100 genes), since this

provides a larger gene group [16]. To get a scale of the persistence of evolutionary changes in

promoters of GTS100 genes we used publicly available PhyloP27way data from UCSC

genome browser [33, 34]. Genes were ordered based on transposon de-repression scores [16]

and d^P was calculated for individual genes, the genome average as a control group, and for the

average of GTS100 gene promoter regions (Fig. 2A-C, 3A). Genes without full PhyloP

coverage were removed from the analysis (where ≥1 nucleotide of the analysed genomic region

contained the *D. melanogaster* sequence only).

The promoter score average of GTS100 genes showed a significant increase in

promoter nucleotide changes compared to the control (Fig. 2C). Although many of these genes

have additional roles, gene ontology (GO) analysis confirmed a primary role in piRNA binding

(Fig. 2D) [35-37].

To understand whether GTS100 genes were individually fast evolving, individual d^P

scores were calculated (Fig. 3A). Compared to the genome average, 49 genes (FEPG, Fast

Evolving Promoter Germline genes) were significantly increased in promoter change events,

while 12 were significantly decreased (Fig. 3A). Of note *Brother of Yb* (*BoYb*) which is

considered the germline replacement of *Yb* [3], also displayed a fast evolving promoter in

contrast to its somatic counterpart *Yb* (Fig. 3A). To specifically look at indels at a high

confidence level, we analysed FEPG genes for rapid promoter evolution between the

previously used five *Drosophila* species (Fig. 3B-D). Here, indels and base changes were

measured, showing a significant increase in both indels and base changes compared to the

control (Fig. 3C and D).

A subset of piRNA factors that are part of distinct protein complexes display hot spots

for rapid evolution in promoters

In a saturating genetic screen for regulators of transposon silencing in germ cells

members of the Nup complex were identified as regulators of germline piRNA silencing, and

later shown to rapidly evolve through their promoters [16, 20, 21]. In the same screen,

numerous genes whose products are part of distinct protein complexes were identified,

including the nuclear cap-binding complex (CBC), THO complex, the exon junction complex

(EJC) inner core, outer shell and EJC transient factors, the non-specific lethal (NSL) complex,

SUMOylation modifiers and the complex of proteins associated with Set1 (COMPASS) (Supplementary Fig. 2). Given the fast evolution of promoters in germline-specific piRNA pathway genes and the Nup complex, we analysed promoters of the additional protein complexes affecting germline piRNA silencing to see whether their rate of evolution matched that of other piRNA silencing factors. Compared to somatic and germ cell piRNA factors, promoter regions for the nuclear CBC, SUMO modifiers, COMPASS complex, NSL complex and EJC outer shell showed a significant accumulation of sequence changes (Fig. 3E and F). Analysis of individual genes from both significant and non-significant gene groups revealed that 43% of analysed genes have rapidly accumulated sequence changes in their promoters (Supplementary Fig. 2). Specifically looking at genes in fast evolving complexes (Fig. 3E and F) revealed that 56% of genes in these groups rapidly evolve through accumulation of mutations in their promoters (Supplementary Fig. 2).

When analysing the base changes and indels of rapid evolving promoters (Supplementary Fig. 3A-D), indel accumulation was observed for the SUMO modifiers, COMPASS complex, NSL complex and EJC outer shell (4 out of 5), while the nuclear CBC showed a significant decrease (Supplementary Fig. 3B). The nuclear CBC, SUMO modifiers and COMPASS complex also showed a significant accumulation of base changes in their promoter regions (3 out of 5) (Supplementary Fig. 3D). Individually, 4 out of 27 genes were significantly enriched in indel events (*Hcf*, *dgt1*, *nsl1*, and *Bin1*), while base change accumulation events indicated 8 out of 27 genes (*Ars2*, *cbp80*, *Ulp1*, *Set1*, *Hcf*, *Cfp1*, *MBD-R2*) being significantly increased (Supplementary Fig. 3E).

Next, we analysed the rate of evolution in the coding regions of these complexes using MKT tests [32]. This analysis flagged the EJC outer shell, EJC transient facts, NSL complex, and SUMO modifiers as under positive selection in both populations, while the Nuclear CBC was under positive selection in the Zambia population only (Supplementary Fig. 4).

Analysis of piRNA factor differential gene expression in ovaries of *D. melanogaster* and

D. yakuba

Since promoters of FEPG genes are fast evolving, we analysed publicly available RNA-seq data from ovaries between *D. melanogaster* and *D. yakuba* to assess gene expression divergence in the FEPG gene group compared to the somatic and germ cell piRNA factors (Fig. 4A) [38-40]. Here, many FEPG genes had a tendency for slight significant divergence in gene expression in both directions (*D. melanogaster* or *D. yakuba*), but this was true for only few somatic and germ cell piRNA factors genes including *arx* (Fig. 4A). Notably, we observed large expression divergence for *CG32152, MEP-1* and *Rbbp5* (Fig. 4A). Aligning the promoters of these three genes revealed an increased number of indels and base changes compared to the 5'UTR or the coding region (Fig. 4B-D), but the cohort was unfortunately too small to detect whether increased changes in promoters correlate with altered expression between the two species. Further limitations stem from the pleiotropic roles of the analysed genes, and therefore expression profiles likely differ across cell types in the analysed tissue.

Germ cell-specific genes show rapid evolution of promoters associated with divergent

gene expression between species

To investigate whether the rapid evolution of promoters in piRNA factors expressed in germ cells is a common feature of germline-expressed genes, we used publicly available RNA-seq data from FACS-sorted vasa-GFP ovaries to identify genes specifically expressed in either germ cells or somatic cells of the ovary. First, using differential RNA-seq analysis of FACS-sorted vasa-GFP+ (germline) and vasa-GFP- (somatic) cells from cultured *D.mel* ovary GSC line (vasa-GFP;hs-bam;bamΔ86) (GSE119862) [41], we defined germline-enriched (vasa-

GFP+; log2FC>0, p-val<0.01) and soma-enriched (vasa-GFP-; log2FC<0, p-val<0.01) genes

(Fig. 5A and B). Next, using RNA-seq data from ovarian somatic cells (OSCs) (GSE160860)

[42], we further filtered the soma-enriched genes that are highly expressed in OSCs.

Additionally, we further filtered the germline-enriched genes that show expression in early fly

embryos (0-2hr, modENCODE). In this way, we generated a stringent list of 107 genes

specifically expressed in germ cells and of 59 soma-expressed genes (Supplementary Dataset

1).

We then analysed sequence change accumulation in promoters using PhyloP data and

calculated d^P for all *D. melanogaster* genes and genes expressed specifically in somatic cells

or the germline (Fig. 5C and D) or the average (Fig 5E). Germ cell-specific expressed genes

significantly accumulated changes (d^P) in their promoters compared to soma-expressed genes

(Fig. 5F). Comparison of the average promoter PhyloP scores and d^P conservation scores

between the genome average and somatic or germ cell-specifically expressed genes equally

revealed a significant increase in germ cell gene promoter nucleotide changes compared to the

genome average (Fig. 5G and H).

Since rapid promoter evolution of germline piRNA silencing factors correlated with

gene expression divergence in our previous analysis, we hypothesized that this was a general

feature of germ cell genes. To test this, we compared gene expression divergence between

somatic and germ cell-specific genes using publicly available RNA-seq data of *D.*

melanogaster and *D. yakuba* ovaries (Fig. 6A) [38-40]. Notably, genes specifically expressed

in germ cells showed slightly higher divergence in expression compared to soma-expressed

genes (Fig. 6A). When d^P conservation scores were plotted against the gene expression change

between *D. melanogaster* and *D. yakuba*, the germ cell-specifically expressed genes showed

slightly higher divergence in expression especially in the higher d^P score groups (>50-75, >75-

100) compared to soma-expressed genes, which showed relatively equal expression divergence

across the different gene groups (Fig. 6C-E).

Discussion

Through genomic analysis of germline and somatic transposon silencing genes and

their regulators we identify hot spots for rapid evolution in promoter regions (-30 to -380

nucleotides upstream of the TSS) of germ cell-specific (dual-strand clusters and the ping-pong

cycle), but not soma-expressed piRNA pathway genes (cTGS and piRNA biogenesis). Further,

we show that rapid promoter evolution is a general feature of germ cell-specific genes

compared to those expressed only in the soma. Our analysis suggests that this mode of

evolution in the germline pathway could be a general feature of at least insect evolution.

Overall, our results point towards a key role for rapid evolution of gene promoters in the germ

cell-specific piRNA pathway which could, coupled with other drivers of expression

divergence, impact the expression of germline regulatory genes.

Transposon mobility has been attributed to the accumulation of sequence changes in

promoter regions because of the presence of open chromatin around the transcription start site

of genes [43]. Interestingly, in core NPC genes, rapid evolution mostly led to accumulation of

indels [20, 21], while piRNA pathway genes display equal accumulation of indels and base

changes. Likely, indels have more profound effects on changes in transcription than single

nucleotide alterations. This feature might reflect that compromised transposon silencing causes

sterility, hence not allowing for substantial changes in expression in piRNA processing genes.

Likewise, stoichiometric changes in protein complexes can drastically alter complex

functionality, e.g. the male-specific lethal (MSL) complex binds less to its targets when one

component of the complex is missing [44, 45]. Hence, changes in the protein levels of

individual piRNA factors likely have dominant effects on their capacity to silence transposon

activity.

The occurrence of novel transposons can fundamentally impact species reproductive

success through a phenomenon call hybrid dysgenesis [46]. Here, if a novel transposable

element is transmitted by the male, female fertility is severely compromised and can result in

sterility as a result of missing piRNA silencing of the novel invader. Essential to combat these

novel active transposable elements is the ping-pong cycle, primed by transcripts from the novel

transposon. Accordingly, females can be become resistant to such novel transposons over time

through RNA transgenerational inheritance of piRNAs to build up ping-pong cycle

amplification in germ cells [9, 10, 47].

Our examination of evolution in coding regions flagged germ cell-specific piRNA

factors, as well as some pleiotropic piRNA processing associated complexes (EJC outer shell,

EJC transient facts, NSL complex, SUMO modifiers, and Nuclear CBC) as under positive

selection. This is unsurprising in the case of primary piRNA processing factors in the germline,

as there are many examples of these factors being under selection [48-56]. For instance,

replacement of *D. melanogaster rhino* and *deadlock* genes with *D. simulans* homologues

results in non-functionality [49]. Here, *D. simulans* Rhino binding domains no longer bind to

D. melanogaster Deadlock, resulting in failed localisation to piRNA clusters [49]. Such effects

resulting from few amino acid changes act as powerful driving forces in diverging piRNA

processing. In this setting, rapid promoter evolution of piRNA silencing genes could act as an

additional factor driving changes in expression levels and factor stoichiometry for piRNA

processing divergence, which may be difficult to explore when coupled with functional

divergence. Intuitively, one would associate rapid promoter revolution with changes in

expression, however, we observed minimal effects on expression. Since transposon silencing

is essential, changes in cis regulatory elements of these regulatory factors is likely constrained

by compensatory mechanisms and could be accompanied by changes in enhancer regions.

Whether this is the case would required a detailed knowledge of relevant enhancers and

transcription factor binding.

The pattern of changes observed in fast evolving promoters like germline specific

piRNA factors and Nups resembles the outcome of a *P*-element mutagenesis experiment in the

egghead (*egh*) locus coding for a glycosphingolipid biosynthesis enzyme [57]. Here, multiple

base changes were observed in the region of the first promoter after *P*-element mutagenesis,

but not an actual *P*-element insert, resulting in a sex peptide insensitive allele *egh^{cm}*. Mutations

in the *egh* gene result in pleiotropic phenotypes and the *egh^{cm}* allele disrupts neuronal wiring

required for the female post-mating response and development of the optic lobe [22, 57, 58].

Since establishing transposon resistance involves forced selection, genes whose

expression changes result in pleiotropic effect, like channel Nup54, could combine adaptations

in piRNA processing with changes in neuronal wiring resulting in altered behavioural

preferences [20]. Of note, in *D. simulans*, a close relative of *D. melanogaster*, projections of

fruitless P1 sensory neurons that control courtship have changed and alter mate preference [59].

The severe impact of hybrid dysgenesis on fertility likely limits fast evolution to species

with high numbers of progeny. Perhaps this can explain the differences seen when comparing

insect and mammalian promoters that are generally more conserved [26]. However, rapid

promoter evolution has also been observed in primate promoters, suggesting a common mode

of evolution that has yet to be explored [20, 21, 26, 60].

Materials and methods

Sequence/data retrieval and alignment

Gene and promotor sequences were retrieved from UCSC Genome Browser using the

UCSC Table Browser sequence retrieval tool [33, 34]. A standardised region of 2000 bases

upstream of the annotated gene TSS was exported for each gene to ensure inclusion of promoter

regions. Sequences were imported and aligned with clustalW in MEGA11 [61]. PhyloP27way

data were sourced from UCSC genome browser (genome.ucsc.edu) through the Table Browser

tool [33, 34]. Data points were collected for a region of 1000 nucleotides upstream and 300

nucleotides downstream for each of the analysed genes. Genes without full species coverage

when analysing evolution between the 5 *Drosophila* species, where ≥ 1 species had no

conserved genomic sequence were removed from the analysis. Genes without full PhyloP

coverage were removed from the analysis (where ≥ 1 nucleotide of the analysed genomic region

contained the *D. melanogaster* sequence only).

Molecular evolution of open reading frames analysis

Analysis of adaptive protein evolution was performed via MKTs [32] using the PopFly

online database (imkt.uab.cat) which uses Drosophila Genome Nexus project sequence data

[62-64]. Polymorphism and divergence data were collected from the ancestral Congo

population, chosen since it is a sub-Saharan population with comparably high ancestral stability

compared to other populations. The 'Standard MKT' test was used to calculate results for

individual genes. Fisher's Exact Test with significance defined as Bonferroni corrected $p \leq 0.05$

was used to calculate significance between nonsynonymous to synonymous polymorphisms

within *D. melanogaster* or between *D. melanogaster* and *D. simulans*.

Identification of promotor hot spots and comparison of substitution rates

All gene models were manually inspected in FlyBase using the JBrowse browser and

dominant transcripts were chosen based on comparative modENCODE expression data [65].

Gene and promoter sequence alignments were trimmed to -1000/+300 nucleotides from the

TSS of each gene. To calculate the frequency of sequence change hotspots in promoter

sequences, we calculated the hotspot accumulation score d using alignments generated as

described, or PhyloP data. Alignments were translated into 'events' for each species compared

to *D. melanogaster*, where for each nucleotide in the sequence, 0 signified a conserved

sequence and 1 signified a sequence change (base change or indel event) [66]. Events were

calculated for all changes, as well as base changes and indels individually. The sum of events

was calculated for concatenated gene groups and individual genes at each nucleotide position.

A sliding event (*Se*) score was calculated from this using a sliding window of five bases along

the sequence, from which heatmaps were generated. To calculate the percentage of events

greater than the average control promoter *Se* score (*d*), a 350-nucleotide region upstream of the

estimated TATA box region was analysed where the total number of *Se* scores exceeding the

control group average sliding event score (Se^C) was divided by the total number of events in

that region (*N*).

378
$$d = \frac{\text{Total number of } Se \text{ events where } Se > Se^C}{N} \times 100$$

379   To calculate the promoter region d scores from PhyloP data, the total number of PhyloP

380   (*p*) scores in the 350 nucleotide regions less than the control group average promoter region

381   (p^C) took the place of the total number of *Se* events where *Se* is greater than Se^C (see equation

382   below).

383
$$d^P = \frac{\text{Total number of } p \text{ scores} < p^C}{N} \times 100$$

Significance was calculated using non-parametric chi-squared tests compared to the

control group d score. Significance values where $p \leq 0.05$ with Bonferroni correction were

considered statistically significant.

Identification of somatic and germ cell-specifically expressed genes in *D. melanogaster*

We defined somatic and germline genes using several steps. First, using processed

RNA-seq data from vasa-GFP+/- cells sorted from cultured ovary GSC line (vasa-GFP;hs-

bam;bamΔ86) (GSE119862) [41], we defined germline-enriched (vasa-GFP+) using logFC>0,

p-value<0.01 parameters and soma-enriched (vasa-GFP-) genes using logFC<0, p-value<0.01

parameters (Figures 4a and 4b). Next, using processed RNA-seq data from fly ovaries

(modENCODE), we filtered genes that are expressed in adult fly ovaries using RPKM>10

parameter for both soma and germline genes. To stringently define the germline genes, we

additionally used processed RNA-seq data from early fly embryos (0-2hr, modENCODE;

mRNA-seq fly embryo 0-2hr; RPKM; mE_mRNA_em0-2hr; FBlc0000086) to further filter

the germline-enriched genes using >9 RPKM parameter. To stringently define the somatic-

specific genes, we additionally used processed RNA-seq data from ovarian somatic cells

(OSCs) (n=4, GSE160860) [42] and filtered the soma-enriched genes using >20 RPKM to

select genes that are expressed in OSCs as well as parameter <2 RPKM for early fly embryo

(0-2hr, modENCODE) to filter out genes that are expressed in early embryos. This analysis

generated a list of 107 germ cell-specifically expressed genes and of 59 somatic cell-

specifically expressed genes (Fig. 4A and B, Supplementary Dataset 1).

Comparative gene expression analysis between *D. melanogaster* and *D. yakuba*

Using publicly available raw RNA-seq data from *D. melanogaster* (n=3) and *D. yakuba*

(n=4) ovaries, we trimmed adapter sequences from the raw reads using Cutadapt tool (v1.18,

default parameters) and aligned the trimmed reads to the genome assemblies (dm6 and

droYak2) using the RNA-seq aligner STAR (v2.7.3a). Gene counts were quantified using the

featureCounts tool (Subread package v1.5.3). Orthologues genes were identified using

blastDm2FB in UCSC Table Browser. Gene count normalization and differential gene

expression analysis between the species were performed using DESeq2 (v1.30.1) with an

additional species-specific gene lengths normalization step.

Supplementary information

417 Supplementary materials include Fig. S1-4 and Supplementary Dataset 1.

Authors' contributions

420 DWJM and MS conceived and directed the project. DWJM performed genomic and

421 evolutionary analysis. AA and DWJM analysed differential gene expression. DWJM wrote the

422 initial draft and all authors contributed to the final manuscript. All authors read and approved

423 the final manuscript.

Acknowledgements

426 We thank the Soller lab for helpful discussion.

Funding

429 This work was supported by the Medical Research Council to DWJM [MR/N013913/1] and

430 the Biotechnology and Biological Science Research Council to MS. Work by BCN and AA

431 was funded by Cancer Research UK [A21143] and the Wellcome Trust [110161/Z/15/Z].

Availability of data and materials

434 All data generated or analysed during this study are included in the supplementary information

435 files.

Ethics approval and consent to participate

438 Not applicable.

440 **Consent for publication**

441 Not applicable.

443 **Competing interests**

444 The authors declare no competing interests.

446 **References**

447 1. Iwakawa H-o, Tomari Y: **Life of RISC: Formation, action, and degradation of**

448  **RNA-induced silencing complex**. *Molecular Cell* 2022, **82**(1):30-43.

449 2. Czech B, Munafò M, Ciabrelli F, Eastwood EL, Fabry MH, Kneuss E, Hannon GJ:

450  **piRNA-Guided Genome Defense: From Biogenesis to Silencing**. *Annual Review of*

451  *Genetics* 2018, **52**(1):131-157.

452 3. Yamashiro H, Siomi MC: **PIWI-Interacting RNA in Drosophila: Biogenesis,**

453  **Transposon Regulation, and Beyond**. *Chemical Reviews* 2018, **118**(8):4404-4421.

454 4. Slotkin RK, Martienssen R: **Transposable elements and the epigenetic regulation**

455  **of the genome**. *Nature Reviews Genetics* 2007, **8**(4):272-285.

456 5. Malone CD, Brennecke J, Dus M, Stark A, McCombie WR, Sachidanandam R,

457  Hannon GJ: **Specialized piRNA Pathways Act in Germline and Somatic Tissues of**

458  **the Drosophila Ovary**. *Cell* 2009, **137**(3):522-535.

459 6. Khurana JS, Theurkauf W: **piRNAs, transposon silencing, and Drosophila**

460  **germline development**. *Journal of Cell Biology* 2010, **191**(5):905-913.

461 7. Senti K-A, Brennecke J: **The piRNA pathway: a fly's perspective on the guardian**

462  **of the genome**. *Trends in Genetics* 2010, **26**(12):499-509.

463   8.   Sakakibara K, Siomi MC: **The PIWI-Interacting RNA Molecular Pathway:**

464   **Insights From Cultured Silkworm Germline Cells**. *BioEssays* 2018,

465   **40**(1):1700068.

466   9.   Moelling K: **Epigenetics and transgenerational inheritance**. *J Physiol* 2023.

467   10.   Grentzinger T, Armenise C, Brun C, Mugat B, Serrano V, Pelisson A, Chambeyron S:

468   **piRNA-mediated transgenerational inheritance of an acquired trait**. *Genome Res*

469   2012, **22**(10):1877-1888.

470   11.   Wang X, Ramat A, Simonelig M, Liu M-F: **Emerging roles and functional**

471   **mechanisms of PIWI-interacting RNAs**. *Nature Reviews Molecular Cell Biology*

472   2023, **24**(2):123-141.

473   12.   Sato K, Siomi MC: **The piRNA pathway in Drosophila ovarian germ and somatic**

474   **cells**. *Proceedings of the Japan Academy, Series B* 2020, **96**(1):32-42.

475   13.   Signor S, Vedanayagam J, Kim BY, Wierzbicki F, Kofler R, Lai EC: **Rapid**

476   **evolutionary diversification of the flamenco locus across simulans clade**

477   **Drosophila species**. *PLOS Genetics* 2023, **19**(8):e1010914.

478   14.   Czech B, Hannon GJ: **One Loop to Rule Them All: The Ping-Pong Cycle and**

479   **piRNA-Guided Silencing**. *Trends in Biochemical Sciences* 2016, **41**(4):324-337.

480   15.   Haase AD: **An introduction to PIWI-interacting RNAs (piRNAs) in the context of**

481   **metazoan small RNA silencing pathways**. *RNA Biology* 2022, **19**(1):1094-1102.

482   16.   Czech B, Preall J, B. , McGinn J, Hannon G, J. : **A Transcriptome-wide RNAi**

483   **Screen in the Drosophila Ovary Reveals Factors of the Germline piRNA**

484   **Pathway**. *Molecular Cell* 2013, **50**(5):749-761.

485   17.   Brown JS, Zhang D, Chen W, Lee H-C: **Sensitized piRNA reporter identifies**

486   **multiple RNA processing factors involved in piRNA-mediated gene silencing**. In.

487   bioRxiv: Cold Spring Harbor Laboratory; 2023.

488   18.   Handler D, Meixner K, Pizka M, Lauss K, Schmied C, Franz, Brennecke J: **The**

489   **Genetic Makeup of the Drosophila piRNA Pathway**. *Molecular Cell* 2013,

490   **50**(5):762-777.

491   19.   Munafò M, Lawless VR, Passera A, Macmillan S, Bornelöv S, Haussmann IU, Soller

492   M, Hannon GJ, Czech B: **Channel nuclear pore complex subunits are required for**

493   **transposon silencing in Drosophila**. *eLife* 2021, **10**.

494   20.   Nallasivan MP, Haussmann IU, Civetta A, Soller M: **Channel nuclear pore protein**

495   **54 directs sexual differentiation and neuronal wiring of female reproductive**

496   **behaviors in Drosophila**. *BMC Biology* 2021, **19**(1).

497   21.   McQuarrie DWJ, Read AM, Stephens FHS, Civetta A, Soller M: **Indel driven rapid**

498   **evolution of core nuclear pore protein gene promoters**. *Scientific Reports* 2023,

499   **13**(1).

500   22.   Haussmann IU, Hemani Y, Wijesekera T, Dauwalder B, Soller M: **Multiple**

501   **pathways mediate the sex-peptide-regulated switch in female Drosophila**

502   **reproductive behaviours**. *Proc Biol Sci* 2013, **280**(1771):20131938.

503   23.   Tang S, Presgraves DC: **Evolution of the Drosophila Nuclear Pore Complex**

504   **Results in Multiple Hybrid Incompatibilities**. *Science* 2009, **323**(5915):779-782.

505   24.   Tang S, Presgraves DC: **Lineage-Specific Evolution of the Complex Nup160**

506   **Hybrid Incompatibility Between Drosophila melanogaster and Its Sister Species**.

507   *Genetics* 2015, **200**(4):1245-1254.

508   25.   Iwasaki YW, Siomi MC, Siomi H: **PIWI-Interacting RNA: Its Biogenesis and**

509   **Functions**. *Annu Rev Biochem* 2015, **84**:405-433.

510   26.   Taylor MS, Kai C, Kawai J, Carninci P, Hayashizaki Y, Semple CAM: **Heterotachy**

511   **in Mammalian Promoter Evolution**. *PLoS Genetics* 2006, **2**(4):e30.

512   27.   Balacco DL, Soller M: **The m(6)A Writer: Rise of a Machine for Growing Tasks**.

513   *Biochemistry* 2019, **58**(5):363-378.

514   28.   Bawankar P, Lence T, Paolantoni C, Haussmann IU, Kazlauskiene M, Jacob D,

515   Heidelberger JB, Richter FM, Nallasivan MP, Morin V *et al*: **Hakai is required for**

516   **stabilization of core components of the m6A mRNA methylation machinery**.

517   *Nature Communications* 2021, **12**(1):3778.

518   29.   Van Lopik J, Trapotsi M-A, Hannon GJ, Bornelov S, Czech Nicholson B: **Unistrand**

519   **piRNA clusters are an evolutionary conserved mechanism to suppress**

520   **endogenous retroviruses across the Drosophila genus**. *bioRxiv* 2023.

521   30.   Fridrich A, Moran Y: **Some flies do not play ping-pong**. *PLOS Biology* 2023,

522   **21**(6):e3002152.

523   31.   Chary S, Hayashi R: **Mechanistic divergence of piRNA biogenesis in Drosophila**.

524   In.: Cold Spring Harbor Laboratory; 2022.

525   32.   McDonald JH, Kreitman M: **Adaptive protein evolution at the Adh locus in**

526   **Drosophila**. *Nature* 1991, **351**(6328):652-654.

527   33.   Karolchik D: **The UCSC Table Browser data retrieval tool**. *Nucleic Acids Research*

528   2004, **32**(90001):493D-496.

529   34.   Kent WJ, Sugnet CW, Furey TS, Roskin KM, Pringle TH, Zahler AM, Haussler,

530   David: **The Human Genome Browser at UCSC**. *Genome Research* 2002,

531   **12**(6):996-1006.

532   35.   Consortium GO: **The Gene Ontology resource: enriching a GOld mine**. *Nucleic*

533   *Acids Res* 2021, **49**(D1):D325-d334.

534   36.   Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP,

535   Dolinski K, Dwight SS, Eppig JT *et al*: **Gene ontology: tool for the unification of**

536   **biology. The Gene Ontology Consortium**. *Nat Genet* 2000, **25**(1):25-29.

37.   Mi H, Muruganujan A, Ebert D, Huang X, Thomas PD: **PANTHER version 14:**

more genomes, a new PANTHER GO-slim and improvements in enrichment

analysis tools. *Nucleic Acids Research* 2018, **47**(D1):D419-D426.

38.   Sanfilippo P, Wen J, Lai EC: **Landscape and evolution of tissue-specific**

alternative polyadenylation across Drosophila species. *Genome Biology* 2017,

18(1).

39.   Smolko AE, Shapiro-Kulnane L, Salz HK: **The H3K9 methyltransferase SETDB1**

maintains female identity in Drosophila germ cells. *Nature Communications* 2018,

9(1).

40.   Vankuren NW, Vibranovski MD: **A Novel Dataset for Identifying Sex-Biased**

Genes in Drosophila. *Journal of Genomics* 2014, **2**:64-67.

41.   Story B, Ma X, Ishihara K, Li H, Hall K, Peak A, Anoja P, Park J, Haug J, Blanchette

M *et al*: **Defining the expression of piRNA and transposable elements in**

Drosophila ovarian germline stem cells and somatic support cells. *Life Science*

Alliance 2019, **2**(5):e201800211.

42.   Eastwood EL, Jara KA, Bornelöv S, Munafò M, Frantzis V, Kneuss E, Barbar EJ,

Czech B, Hannon GJ: **Dimerisation of the PICTS complex via LC8/Cut-up drives**

co-transcriptional transposon silencing in Drosophila. *eLife* 2021, **10**:e65557.

43.   Treiber CD, Waddell S: **Transposon expression in the Drosophila brain is driven**

by neighboring genes and diversifies the neural transcriptome. *Genome Research*

2020, **30**(11):1559-1569.

44.   Laverty C, Lucci J, Akhtar A: **The MSL complex: X chromosome and beyond**.

Current Opinion in Genetics & Development 2010, **20**(2):171-178.

45.   Straub T, Becker PB: **Dosage compensation: the beginning and end of**

generalization. *Nature Reviews Genetics* 2007, **8**(1):47-57.

46. Brennecke J, Malone CD, Aravin AA, Sachidanandam R, Stark A, Hannon GJ: **An**

Epigenetic Role for Maternally Inherited piRNAs in Transposon Silencing.

Science 2008, **322**(5906):1387-1392.

47. Luo Y, He P, Kanrar N, Fejes Toth K, Aravin AA: **Maternally inherited siRNAs**

initiate piRNA cluster formation. *Mol Cell* 2023.

48. Parhad SS, Yu T, Zhang G, Rice NP, Weng Z, Theurkauf WE: **Adaptive Evolution**

Targets a piRNA Precursor Transcription Network. *Cell Reports* 2020,

30(8):2672-2685.e2675.

49. Parhad SS, Tu S, Weng Z, Theurkauf WE: **Adaptive Evolution Leads to Cross-**

Species Incompatibility in the piRNA Transposon Silencing Machinery.

Developmental Cell 2017, **43**(1):60-70.e65.

50. Kelleher ES, Edelman NB, Barbash DA: **Drosophila Interspecific Hybrids**

Phenocopy piRNA-Pathway Mutants. *PLoS Biology* 2012, **10**(11):e1001428.

51. Vermaak D, Henikoff S, Malik HS: **Positive Selection Drives the Evolution of**

rhino, a Member of the Heterochromatin Protein 1 Family in Drosophila. *PLoS*

Genetics 2005, **1**(1):e9.

52. Obbard DJ, Finnegan DJ: **RNA Interference: Endogenous siRNAs Derived from**

Transposable Elements. *Current Biology* 2008, **18**(13):R561-R563.

53. Blumenstiel JP, Erwin AA, Hemmer LW: **What Drives Positive Selection in the**

Drosophila piRNA Machinery? The Genomic Autoimmunity Hypothesis. *Yale J*

Biol Med 2016, **89**(4):499-512.

54. Wang L, Barbash DA, Kelleher ES: **Adaptive evolution among cytoplasmic piRNA**

proteins leads to decreased genomic auto-immunity. *PLOS Genetics* 2020,

16(6):e1008861.

55.   Parhad SS, Theurkauf WE: **Rapid evolution and conserved function of the piRNA pathway**. *Open Biology* 2019, **9**(1):180181.

56.   Lawlor MA, Ellison CE: **Evolutionary dynamics between transposable elements and their host genomes: mechanisms of suppression and escape**. *Current Opinion in Genetics & Development* 2023, **82**:102092.

57.   Soller M, Haussmann IU, Hollmann M, Choffat Y, White K, Kubli E, Schäfer MA: **Sex-peptide-regulated female sexual behavior requires a subset of ascending ventral nerve cord neurons**. *Curr Biol* 2006, **16**(18):1771-1782.

58.   Fan Y, Soller M, Flister S, Hollmann M, Müller M, Bello B, Egger B, White K, Schäfer MA, Reichert H: **The egghead gene is required for compartmentalization in Drosophila optic lobe development**. *Dev Biol* 2005, **287**(1):61-73.

59.   Seeholzer LF, Seppo M, Stern DL, Ruta V: **Evolution of a central neural circuit underlies Drosophila mate preferences**. *Nature* 2018, **559**(7715):564-569.

60.   Main BJ, Smith AD, Jang H, Nuzhdin SV: **Transcription Start Site Evolution in Drosophila**. *Molecular Biology and Evolution* 2013, **30**(8):1966-1974.

61.   Tamura K, Stecher G, Kumar S: **MEGA11: Molecular Evolutionary Genetics Analysis Version 11**. *Molecular Biology and Evolution* 2021, **38**(7):3022-3027.

62.   Lack JB, Cardeno CM, Crepeau MW, Taylor W, Corbett-Detig RB, Stevens KA, Langley CH, Pool JE: **The Drosophila Genome Nexus: A Population Genomic Resource of 623 Drosophila melanogaster Genomes, Including 197 from a Single Ancestral Range Population**. *Genetics* 2015, **199**(4):1229-1241.

63.   Lack JB, Lange JD, Tang AD, Corbett-Detig RB, Pool JE: **A Thousand Fly Genomes: An Expanded Drosophila Genome Nexus**. *Mol Biol Evol* 2016, **33**(12):3308-3313.

64.   Mackay TFC, Richards S, Stone EA, Barbadilla A, Ayroles JF, Zhu D, Casillas S,

Han Y, Magwire MM, Cridland JM *et al*: **The Drosophila melanogaster Genetic**

Reference Panel. *Nature* 2012, **482**(7384):173-178.

65.   Gramates LS, Agapite J, Attrill H, Calvi BR, Crosby MA, dos Santos G, Goodman

JL, Goutte-Gattat D, Jenkins VK, Kaufman T *et al*: **FlyBase: a guided tour of**

highlighted features. *Genetics* 2022, **220**(4).

66.   Tang H, Lewontin RC: **Locating Regions of Differential Variability in DNA and**

Protein Sequences. *Genetics* 1999, **153**(1):485-495.

Figure Legends

Fig. 1: Promoter regions of germ cell-expressed piRNA processing factors are hot spots

for rapid evolution

A and B) Schematic depiction of somatic and germ cell piRNA factors involved in co-

transcriptional gene silencing and piRNA biogenesis (A) and germline-specific piRNA

pathway genes involved in dual-strand cluster regulation and the ping-pong cycle (B) in the

Drosophila ovary.

C and D) PhyloP27way conservation score averages (C) and PhyloP27way conservation

promoter d^P scores (D) for the 350 nucleotide promoter regions of the somatic and germ cell

piRNA factors and the germ cell-specific piRNA factors compared to all genes in the

Drosophila genome. Statistically significant differences from unpaired student t-tests (C) and

non-parametric chi-squared tests (D) are indicated by asterisks (*** $p \leq 0.001$, **** $p \leq 0.0001$

following Bonferroni correction).

E-J) Heatmaps indicating sequence (F, purple), indel (H, blue) or base change (J, green)

accumulation, and their quantification (E-I) depicted as *d* scores for each of the analysed gene

groups compared to the control among closely related *D. melanogaster*, *D. simulans*, *D.*

sechellia, *D. yakuba* and *D. erecta*. Regions of 1000 nucleotides upstream and 300 nucleotides

downstream of the TSS were analysed. Analysis was performed for the control group (m^{6}A

writer complex and readers), the somatic and germ cell piRNA factors, and the germ cell-

specific piRNA factors. The blue line indicates the promoter region used for quantification of

the substitution rate based on the gene and intergenic region schematic. Statistically significant

differences from non-parametric chi-squared tests are indicated by asterisks (**** p≤0.0001

following Bonferroni correction).

Fig. 2: Promoters of genes required for germline transposon silencing evolve fast

A) PhyloP27way nucleotide scores for the top 100 genes affecting transposon silencing in germ

cells based on transposon derepression z scores and ordered by promoter conservation d^P

scores. A region of 1000 nucleotides upstream and 300 nucleotides downstream are shown.

Red represents lower conservation while blue represents higher.

B) Average PhyloP27way nucleotide scores for the top 100 genes affecting transposon

silencing in germ cells compared to the genome average.

C) Comparison of PhyloP27way conservation promoter scores (d^P) for each of the 100 genes

affecting transposon silencing in germ cells compared to the genome average. Statistically

significant differences from non-parametric chi-squared tests are indicated by asterisks (***

p≤0.001 following Bonferroni correction).

D) Representative GO analysis for the top 100 gene candidates affecting transposon silencing.

Fig. 3: Promoters of FEPG genes and their associated complexes are hotspots for rapid

evolution

A) Individual gene comparison of PhyloP27way conservation promoter scores (d^P) for the top

100 genes affecting transposon silencing in germ cells compared to the genome average.

Statistically significant differences from non-parametric chi-squared tests are indicated by asterisks (* p≤0.05, ** p≤0.01, *** p≤0.001, **** p≤0.0001 following Bonferroni correction).

B-D) Sequence analysis for change accumulation among closely related *D. melanogaster*, *D. simulans*, *D. sechellia*, *D. yakuba* and *D. erecta*. Comparisons were performed for sequence change (B), indel (C), and base change (D) *d* scores for each of the positive 49 genes affecting transposon silencing in FEPG gene group compared to the somatic and germ cell piRNA factors. Statistically significant differences from non-parametric chi-squared tests are indicated by asterisks (**** p≤0.0001 following Bonferroni correction).

E) Comparison of *d* scores for protein complex genes involved in germ cell transposon silencing compared to the somatic and germ cell piRNA factors. Statistically significant differences from non-parametric chi-squared tests are indicated by asterisks (**** p≤0.0001 following Bonferroni correction).

F) Heatmaps indicating sequence change accumulation for protein complex genes involved in germ cell transposon silencing compared to the piRNA factors shown in purple among closely related *D. melanogaster*, *D. simulans*, *D. sechellia*, *D. yakuba* and *D. erecta*. Regions of 1000 nucleotides upstream and downstream of the TSS were analysed. The blue line indicates the promoter region used for quantification of the substitution rate.

Fig. 4: Analysis of differential gene expression of FEPG genes in *D. melanogaster* and *D. yakuba* ovaries

A) Volcano plot showing differential gene expression (DESeq2) between *D. melanogaster* and *D. yakuba* ovaries (RNA-seq data; Supplementary Dataset 1) for FEPG (blue) and the somatic and germ cell piRNA factors (red). Horizontal dashed lines indicate significance thresholds (adjusted p≤0.05).

B-D) Alignment of the promoter regions of *CG32152, MEP-1* and *Rbbp5* for *D. melanogaster*

and *D. yakuba* with indels (red) and base changes (blue) indicated as boxes. The black lines

indicate gene 5'UTRs.

Fig. 5: Germ cell-specifically expressed, but not somatic cell-specifically expressed gene

promoters evolve rapidly and are accompanied by divergence in gene expression

A) Ranking of genes using vasa-GFP signal from FACS-sorted vasa-GFP+/- cells of cultured

ovary GSC line (vasa-GFP;hs-bam;bamΔ86) (GSE119862) that defines germline-enriched

(vasa-GFP+) and soma-enriched (vasa-GFP-) genes.

B) Volcano plot showing differential gene expression between the FACS-sorted vasa-GFP+

(germline) and vasa-GFP- (somatic) cells from cultured ovary GSC line (vasa-GFP;hs-

bam;bamΔ86) (GSE119862).

C and D) PhyloP27way nucleotide scores for somatic cell-specific (C) and germ cell-specific

(D) expressed genes ordered by promoter conservation d^P scores. A region of 1000 nucleotides

upstream and 300 nucleotides downstream are shown. Red represents lower conservation while

blue represents higher.

E) Average PhyloP27way nucleotide scores for all genes in the *Drosophila* genome, somatic

cell and germ cell-specific genes. Purple represents lower conservation while blue represents

higher.

F) The percentage of somatic cell and germ cell-specific genes with significant promoter

nucleotide changes (d^P) versus all genes in the *Drosophila* genome average, divided by

significantly increased (low conservation) or decreased (high conservation). Statistically

significant differences from non-parametric chi-squared tests are indicated by asterisks (****

p≤0.0001 following Bonferroni correction).

G and H) PhyloP27way conservation score averages (G) and PhyloP27way conservation

promoter scores (d^P) (H) for the 350 nucleotide promoter regions compared for all genes in the

Drosophila genome, somatic cell-specifically expressed and germ cell-specifically expressed

genes. Statistically significant differences from unpaired student t-tests (G) and from non-

parametric chi-squared tests (H) are indicated by asterisks (** p≤0.01, **** p≤0.0001

following Bonferroni correction).

Fig. 6: Analysis of gene expression divergence between germ cell and somatic cell-

specifically expressed genes

A) Volcano plot of differential gene expression (DESeq2) between *D. melanogaster* and *D.*

yakuba ovaries (RNA-seq data; Supplementary Dataset 1) for all (grey), germ cell-specific

(blue) and somatic cell-specific (red) gene groups. Horizontal dashed lines indicate

significance (adjusted p≤0.05).

B) Scatterplot of individual gene d^P scores plotted against fold-change expression differences

between *D. melanogaster* and *D. yakuba* ovaries (RNA-seq data; Supplementary Dataset 1) for

germ cell-specific (blue) and somatic cell-specific (red) gene groups. The grey dotted lines

indicate log2Fold-change threshold values of ≥0.5 and ≤0.5.

C and E) Comparison of the expression distribution and percentage of genes with significant

changes in *D. melanogaster* log2FoldChange (≥0.5, D) or significant changes in *D. yakuba*

log2FoldChange (≤0.5, E) expression changes in PhyloP conservation promoter score (d^P)

ranges 25-50, >50-75, and >75-100. Expression changes for germ cell-specific and somatic

cell-specific gene groups were analysed between *D. melanogaster* and *D. yakuba*. Statistically

significant differences from Mann Whitney U tests (C) and non-parametric chi-squared tests

(D and E) are indicated by asterisks (** p≤0.01, *** p≤0.001, **** p≤0.0001 following

Bonferroni correction).

Supplementary Fig. 1: Germ cell-specific piRNA factors are under positive selection and accumulate rapid sequence changes in their promoters

A) Individual gene sequence change, indel, and base change *d* scores and statistically significant accumulation of sequence changes for the somatic piRNA factors and germ cell-specific piRNA factors. Statistically significant differences from non-parametric chi-squared tests are indicated by asterisks (* $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$ following FDR correction).

B) Evolution of coding regions analysed by MKT tests for polymorphisms and divergence between *D. melanogaster* and *D. simulans* for core genes in the somatic and germ cell transposon silencing pathways. R, replacement; S, synonymous. Statistically significant differences from Fisher's Exact Tests are indicated by asterisks (**** $p \leq 0.0001$ following FDR correction).

Supplementary Fig. 2: Genes coding for members of a subgroup of protein complexes implicated in piRNA silencing accumulate sequence changes in their promoters

Individual gene *d* scores and statistically significant accumulation of sequence changes for each gene separated by protein complex compared to the somatic piRNA factors. Statistically significant differences from non-parametric chi-squared tests are indicated by asterisks (* $p \leq 0.05$, ** $p \leq 0.01$, **** $p \leq 0.0001$ following Bonferroni correction).

Supplementary Fig. 3: Promoter evolution of protein complex genes involved in germ cell transposon silencing

A and C) Heatmaps indicating indel (A, blue) or base change (C, green) accumulation in protein complex genes involved in germ cell transposon silencing compared to Yb body genes among

closely related *D. melanogaster*, *D. simulans*, *D. sechellia*, *D. yakuba* and *D. erecta*. Regions

of 1000 nucleotides upstream and 300 nucleotides downstream of the TSS were analysed. The

blue line indicates the promoter region used for quantification of the substitution rate.

B and D) Comparison of indel (B, blue) or base change (D, green) d scores for each of the

protein complex gene groups compared to the somatic piRNA factors. Statistically significant

differences from non-parametric chi-squared tests are indicated by asterisks (* $p \leq 0.05$, **

$p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$ following Bonferroni correction).

E) Individual gene d scores and statistically significant accumulation of indels or base changes

for each gene separated by protein complex compared to the somatic piRNA factors.

Comparison of indel (B, blue) or base change (D, green) d scores for each of the protein

complex gene groups compared to the control group. Statistically significant differences from

non-parametric chi-squared tests are indicated by asterisks (* $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$,

**** $p \leq 0.0001$ following Bonferroni correction).

Supplementary Fig. 4: Coding sequences evolution of protein complex genes involved in

germ cell transposon silencing.

A) Impact of protein complex genes involved in germ cell transposon silencing on transposon

de-repression. Unprocessed data were taken from [1]. Derepression scores (z) were

concatenated for protein complexes in each transposon type (HeTA, TAHRE, blood, burdock)

and an average of all.

B) Evolution of coding regions analysed by MKT tests for polymorphisms and divergence

between *D. melanogaster* and *D. simulans* for individual genes from the selected germline

transposon silencing implicated protein complexes. R, replacement; S, synonymous.

Statistically significant differences from Fisher's Exact Tests are indicated by asterisks (*

$p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$ following FDR correction).

Figure 1

Figure 2

A

B

		Polymorphism			Divergence				
	Population	R	S	R/S	R	S	R/S	Fisher's exact *P*-value (FDR)	
Germ cell-specific piRNA factors	Congo	133	102	1.304	1271	519	2.449	0.000023	****
	Zambia	645	384	1.680	1204	473	2.545	0.000004	****
Somatic and germ cell piRNA factors	Congo	149	97	1.536	571	400	1.428	0.663171	
	Zambia	588	446	1.318	530	338	1.568	0.090960	

A

B

		Polymorphism			Divergence				
	Population	R	S	R/S	R	S	R/S	Fisher's exact *P*-value (FDR)	
Nuclear CBC	Congo	9	20	0.450	86	106	0.811	0.330182	
	Zambia	25	67	0.373	85	98	0.867	0.005210	**
THO complex	Congo	24	29	0.828	77	124	0.621	0.573867	
	Zambia	102	136	0.750	70	108	0.648	0.594249	
EJC inner core	Congo	7	11	0.636	26	52	0.500	0.895886	
	Zambia	32	68	0.471	24	46	0.522	0.926331	
EJC outer shell	Congo	15	26	0.577	274	135	2.030	0.000533	***
	Zambia	64	74	0.865	268	127	2.110	0.000042	****
EJC transient factors	Congo	18	66	0.273	110	124	0.887	0.000157	***
	Zambia	100	259	0.386	100	103	0.971	0.000003	****
NSL complex	Congo	42	62	0.677	366	253	1.447	0.000914	***
	Zambia	206	321	0.642	345	220	1.568	7.44E-12	****
SUMO modifiers	Congo	17	19	0.895	349	147	2.374	0.014222	*
	Zambia	94	97	0.969	337	133	2.534	0.000001	****
COMPASS complex	Congo	6	41	0.146	32	90	0.356	0.106720	
	Zambia	47	124	0.379	29	73	0.397	0.889564	