Promoters of germline transposon silencing genes evolve rapidly accompanied by diverging gene expression

DAVID W. J. MCQUARRIE[1,2], AZAD ALIZADA[3], BENJAMIN CZECH NICHOLSON[3] AND MATTHIAS SOLLER[1,2]*

[1]School of Biosciences, College of Life and Environmental Sciences, University of Birmingham, Edgbaston, Birmingham, B15 2TT, United Kingdom

[2]Birmingham Centre for Genome Biology, University of Birmingham, Edgbaston, Birmingham, B15 2TT, United Kingdom

[3]Cancer Research UK Cambridge Institute, University of Cambridge, Li Ka Shing Centre, Cambridge CB2 0RE, United Kingdom

*Corresponding author: m.soller@bham.ac.uk

Running title: Evolution of promoters of germline transposon silencing factors

Key Words: Piwi-interacting RNA (piRNA), germline transposon silencing, RNA transgenerational inheritance, rapid evolution, promoter evolution, speciation, nuclear pore complex, neuronal wiring, transposon silencing

Abstract

Background: The piRNA pathway in animal gonads functions as an 'RNA-based

immune system', serving to silence transposable elements and prevent inheritance of novel

invaders. In *Drosophila*, this pathway relies on three gonad-specific Argonaute proteins

(Argonaute-3, Aubergine and Piwi) that associate with 23-28 nucleotide piRNAs, directing the

silencing of transposon-derived transcripts. Transposons constitute a primary driver of genome

evolution, yet the evolution of piRNA pathway factors has not received in-depth exploration.

Specifically, channel nuclear pore proteins, which impact piRNA processing, exhibit regions

of rapid evolution in their promoters. Consequently, the question arises whether such a mode

of evolution is a general feature of transposon silencing pathways.

Results: By employing phylogenomic analysis of coding and promoter regions within

genes that function in transposon silencing in *Drosophila*, we demonstrate that the promoters

of germ cell-specific piRNA factors are undergoing rapid evolution. Our findings indicate that

rapid promoter evolution is a common trait among piRNA factors engaged in germline

silencing across insect species, potentially contributing to gene expression divergence in

closely related taxa. Furthermore, we observe that the promoters of genes exclusively

expressed in germ cells generally exhibit rapid evolution, which correlates with divergent gene

expression.

Conclusions: Our results suggest that promoter evolution could have wide impact on

transposon silencing and evolution of species through differential expression of genes driven

by invading transposons.

Background

Eukaryotic organisms are continuously challenged from genomic parasites called transposable elements (TEs) (1-4). Unchecked transposon activity often results in reduced reproductive fitness (5-8). To negate the detrimental effects posed by TE mobilisation, vital regulatory pathways evolved that efficiently suppress transposon activity. At the centre of these are 23-28 nucleotide small RNAs, called PIWI-interacting RNAs (piRNAs). piRNAs are generated from transposons and from genomic clusters and bind to Argonaute proteins of the PIWI-clade, including P-element induced wimpy testis (Piwi). Aubergine (Aub) and Argonaute-3 (Ago3). The piRNA pathway is often referred to as an 'RNA-based immune system' thanks to its ability to adopt to new TE insertions and the memory of past transposon activity through piRNA clusters through RNA transgenerational inheritance (1, 2, 9, 10).

The piRNA pathway is arguably best understood in the ovary of *Drosophila melanogaster*. It was shown that ovaries feature two branches of the silencing machinery (5), one that is active in germ cells and a condensed version that is specific to somatic follicle cells that surround the germline (2, 11). Somatic follicle cells only feature unistrand piRNA clusters, such as *flamenco*, and exclusively produce piRNAs via a Zucchini (Zuc)-dependent biogenesis mechanism that takes place on the mitochondria surface involving Armitage (Armi), Papi, Vreteno (Vret), Gasz, Daedalus (Daed), Minotaur (Mino), Shutdown (Shu), and Sister of Yb (SoYb) (Fig. 1A) (3, 12). Licensing of *flam* for piRNA production takes place upstream, at so-called Yb-bodies (1, 12). Somatic cells only express Piwi, which following the association with a mature piRNA shuttles to the nucleus. There, Piwi-piRNA complexes scan for nascent TE transcripts and instruct co-transcriptional gene silencing (cTGS) that requires components of the general chromatin silencing machinery and the piRNA pathway-specific factors Asterix (Arx), Piwi, Panoramix (Panx), Nuclear export factor 2 (Nxf2) and Maelstrom (Mael) (Fig. 1A) (1, 2, 12, 13).

In germ cells, piRNAs predominantly originate from dual-strand clusters and are

produced by a different biogenesis mechanism called the ping-pong amplification cycle (Fig.

1B) (1-3, 5, 11, 12, 14). Dual-strand clusters harbour remnants of transposons and are expressed

via a non-canonical transcription process that is centred on a complex containing the HP1a

homolog Rhino (Rhi), Deadlock (Del) and Cutoff (Cuff), and further requires Moonshiner

(Moon) (Fig. 1B). This Rhi-Del-Cuff complex also recruits a non-canonical export machinery

consisting of Bootlegger (Boot) and Nuclear export factor 3 (Nxf3), which transport piRNA

precursors for processing to a germ-cell specific perinuclear region termed "nuage" (Fig. 1B).

Here, the ping-pong cycle takes place and relies on the two PIWI proteins Aubergine (Aub)

and Argonaute 3 (Ago3), as well as several RNA binding proteins and Tudor-domain

containing factors including Tejas (Tej), Vasa (Vas), Qin, Squash (Squ), Tapas, Brother of Yb

(BoYb), Krimper (Krimp), and Spindle-E (Spn-E) (Fig. 1B) (1-3, 12, 14, 15).

Channel nuclear pore proteins (Nups) have been implicated in piRNA regulated

transposon silencing in *C. elegans* (16). Interestingly, in *Drosophila*, Nups were shown to

affect transposon silencing in both the germline (15, 17), and in somatic cells of the ovary, here

specifically by contributing to the conversion of precursor RNAs from the *flam* locus into

piRNAs (18). Recently, we found that inner and outer ring Nups in the nuclear pore complex

(NPC) evolve rapidly through indel accumulation in their promoters (19, 20). Such promoter

indel variability in Nup54 has dominant, pleiotropic effects on sexual differentiation including

neuronal wiring important for female post-mating behaviours directed by male-derived sex

peptide, that as a result of sexual conflict could drive speciation by such mechanism (16, 19,

21). Intriguingly, nuclear import/export pathways have been linked to speciation through

hybrid incompatibility, but are in addition important for many cellular processes beyond

piRNA processing (22, 23). Hence, promoter evolution particularly of piRNA processing genes

could have wide impact on transposon silencing and evolution of species through differential

expression of genes, but whether regulators of germline transposon silencing generally evolve

rapidly through their promoters has not been determined.

Here, we analyse the promoters of genes involved in the ovarian piRNA pathway. We

compare piRNA factors required in somatic cells (Yb) or in both the somatic and germline

compartments (cTGS and piRNA biogenesis, Fig. 1A) with those that are germ cell-specific

(dual-strand clusters and the ping-pong cycle, Fig. 1B). We find that the promoters of germline-

specific piRNA factors are hot spots for rapid evolution compared to those of somatic piRNA

genes and the *Drosophila* genome average throughout 27 insect species. Analysis of genes with

significant accumulation of promoter changes reveals a mix of indel and base change

accumulation. Further, our analysis reveals that rapid promoter evolution is a general feature

of genes specifically expressed in germ cells, while soma-expressed genes evolve at a similar

rate to the average of all *Drosophila* genes. Through cross-species differential ovary RNA-seq

analysis we reveal that germ cell-specific genes also acquired more divergent expression levels

compared to genes expressed in somatic cells. Our findings highlight that promoters of

germline genes involved in transposon silencing evolve rapidly and are accompanied by

diverging gene expression, suggesting a possible mode of rapid speciation through

accumulation of changes in promoters.

Results

Promoter regions, but not protein coding regions, of germline-specific piRNA factors are

hot spots for rapid evolution

Since Nups have been shown to function in transposon silencing in the germline (12

from 14 tested are above the control) and display fast evolving promoters (15, 19, 20), we

analysed the rate of evolution in promoters and coding regions of piRNA pathway genes. We

separated all piRNA genes based on their genetic requirement into two categories: piRNA

factors essential in somatic cells, including the soma-specific piRNA factor Yb and piRNA

factors expressed in both somatic and germline cells (cTGS and piRNA biogenesis, Fig. 1A),

and germ cell-specific piRNA proteins (dual-strand clusters and ping-pong cycle, Fig 1B). The

piRNA factors essential in somatic cells include *armi*, *papi*, *vret*, *Gasz*, *daed*, *mino*, *zuc*, *shu*,

SoYb, *Yb*, *arx*, *piwi*, *panx*, *nxf2*, and *mael* (Fig. 1A). The group of piRNA factors that are

exclusively required in the germline are *aub*, *tej*, *Ago3*, *vas*, *qin*, *squ*, *tapas*, *BoYb*, *krimp*, *spn-*

E, Boot, *moon*, *nxf3*, *cuff*, *rhi*, and *del* (Fig. 1B) (3, 13, 24).

We defined the promoter containing region as a 350-nucleotide (-30 to -380) window

upstream of the predicted TATA box region 30 nucleotides upstream of the transcription start

site (TSS) (25). In a first approach, we performed a genome-wide promoter evolution analysis

of germline-specific as well as somatic piRNA factors and compared it to all genes in the

Drosophila genome (Fig 1C-F). We used publicly available PhyloP27way data composed of

27 insect species as a large-scale conservation score. We observed a significant accumulation

of changes in the germ cell-specific piRNA factor promoter regions compared to the genome

(Fig. 1C-F). Of note, the somatic piRNA factors evolved slower compared to the genome.

We next compared indel and base changes in the promoter regions between five closely

related *Drosophila* species (*D. melanogaster*, *D. simulans*, *D. sechellia*, *D. yakuba* and *D.*

erecta). Promoter scores (*d*) were calculated and compared to a control group comprised of the

m^{6}A methylation machinery genes (*Mettl3*, *Mettl14*, *fl(2)d*, *virilizer*, *flacc*, *nito* and *Hakai*,

Ythdc1 and *Ythdf*), chosen due to their high evolutionary conservation and requirement for strict

stoichiometry for functionality, making this an ideal control group to monitor promoter

evolution (20, 26, 27). As previously reported, the m^{6}A machinery showed a slight increase in

sequence changes in promoter regions compared to coding regions (Fig. 1G-L) (20). In this

analysis, the group of somatic piRNA factors showed neutral evolution compared to the control

(Fig. 1G-L), while germ cell-specific piRNA factors as a group evolved at a much higher rate

(Fig. 1G and H), accruing a significantly increased number of promoter-located indels (Fig. 1I

and J) and base changes (Fig. 1K and L). At the individual gene level, 7 out of 14 of germ cell-

specific piRNA factors (*Ago3* was omitted due to its low conservation between closely related

Drosophila species (28-30) showed significant accumulation of promoter changes

(Supplementary Fig. 1B). Among the somatic piRNA factors, 3 out of 15 genes had significant

accumulation of promoter changes (Supplementary Fig. 1B).

To assess evolution in the coding region, we used McDonald-Kreitman tests (MKTs)

and analysed polymorphisms and divergence within *D. melanogaster* and between *D.*

melanogaster and *D. simulans* (31). None of the somatic piRNA factor genes were under

selection, while one of the germ cell-specific piRNA factors (*krimp*) was under selection

(Supplementary Fig. 1A).

Promoters of genes required for germline transposon silencing are fast evolving

Given the rapid evolution of germ cell-specific piRNA factors among closely related

Drosophila species, we wanted to determine (1) whether this is a general feature of insect

evolution, and (2) whether this is a general feature of genes required for transposon silencing

that have been identified in a genetic screen (GTS100 genes) (15). To get a scale of the

persistence of evolutionary changes in promoters of GTS100 genes we used publicly available

PhyloP27way data from UCSC genome browser (32, 33). Genes were ordered based on

transposon de-repression scores (15) and d^P was calculated for individual genes, the genome

average as a control group, and for the average of GTS100 gene promoter regions (Fig. 2A-E).

Genes without full PhyloP coverage (where ≥1 nucleotide in gene or upstream regions had no

homologous sequence in the 27 related species) were removed from the analysis.

The promoter score average of GTS100 genes showed a significant increase in

promoter nucleotide changes compared to the control (Fig. 2C). Although many of these genes

have additional roles, gene ontology (GO) analysis confirmed a primary role in piRNA binding (Fig. 2D) (34-36). To understand whether GTS100 genes were individually fast evolving, individual d^P scores were calculated (Fig. 2E). Compared to the genome average, 49 genes (FEPG, Fast Evolving Promoter Germline genes) were significantly increased in promoter change events, while 12 were significantly decreased (Fig. 2E). Of note *Brother of Yb* (*BoYb*) which is considered the germline replacement of *Yb* (3), also displayed a fast evolving promoter in contrast to its somatic counterpart *Yb* (Fig. 2E). To specifically look at indels at a high confidence level, we analysed FEPG genes for rapid promoter evolution between the previously used five *Drosophila* species (Fig. 2F-H). Here, indels and base changes were measured, showing a significant increase in both indels and base changes compared to the control (Fig. 2G and H).

A subset of piRNA factors that are part of distinct protein complexes display hot spots for rapid evolution in promoters

In a saturating genetic screen for regulators of transposon silencing in germ cells (15), numerous genes that are part of distinct protein complexes, including the nuclear cap-binding complex (CBC), THO complex, the exon junction complex (EJC) inner core, outer shell and EJC transient factors, the non-specific lethal (NSL) complex, SUMOylation modifiers and the complex of proteins associated with Set1 (COMPASS) were identified (Supplementary Fig. 2). Given the fast evolution of promoters in germline-specific piRNA genes, we analysed their promoters for the rate of evolution. Compared to soma piRNA factors, promoter regions for the nuclear CBC, SUMO modifiers, COMPASS complex, NSL complex and EJC outer shell showed a significant accumulation of sequence changes (Fig. 2I and J). Analysis of individual genes from both significant and non-significant gene groups revealed that 43% of analysed genes have rapidly accumulated sequence changes in their promoters (Supplementary Fig. 2).

Specifically looking at genes in fast evolving complexes (Fig. 2I-J) revealed that 56% of genes in these groups rapidly evolve through accumulation of mutations in their promoters (Supplementary Fig. 2).

When analysing the base changes and indels of rapid evolving promoters (Fig. S3A-D), indel accumulation was observed for the SUMO modifiers, COMPASS complex, NSL complex and EJC outer shell (4 out of 5), while the nuclear CBC showed a significant decrease (Fig. S3B). The nuclear CBC, SUMO modifiers and COMPASS complex also showed a significant accumulation of base changes in their promoter regions (3 out of 5) (Fig. S3D). Individually, 4 out of 27 genes were significantly enriched in indel events (*Hcf*, *dgt1*, *nsl1*, and *Bin1*), while base change accumulation events indicated 8 out of 27 genes (*Ars2*, *cbp80*, *Ulp1*, *Set1*, *Hcf*, *Cfp1*, *MBD-R2*) being significantly increased (Fig. S3E).

Next, we also analysed the rate of evolution in the coding region of these genes using MKT analysis (31). This analysis shows that most of the coding regions were not under selection (37 out of the 40 analysed genes), with the exception of cbc80 (a nuclear CBC protein), Aly (a factor of the EJC outer shell) and Upf2 (a EJC transient factor), which were under selection (Supplementary Fig. 4). In general, this indicates a neutral mode of evolution in the CDS of these gene complexes implicated in transposon silencing.

piRNA factors with rapidly evolving promoters show divergent gene expression in ovaries of *D. yakuba*

Since promoters of FEPG genes are fast evolving, we analysed publicly available RNA-seq data from ovaries between *D. melanogaster* and *D. yakuba* to assess gene expression divergence in the FEPG gene group compared to the somatic piRNA factors (Fig. 3A) (37-39). Here, many FEPG genes showed significant divergence in gene expression in both directions (*D. melanogaster* or *D. yakuba*), but this was true for only few somatic piRNA factors genes

including *arx* (Fig 3A). Notably, we observed large expression divergence for *CG32152, MEP-*

1 and *Rbbp5* (Fig. 3A). Aligning the promoters of these three genes revealed an increased

number of indels and base changes compared to the 5'UTR or the coding region (Fig. 3B-D),

but the cohort is too small to detect whether increased changes in promoters correlate with

altered expression between the two species.

Germ cell-specific genes show rapid evolution of promoters associated with divergent

gene expression between species

To investigate whether the rapid evolution of promoters in piRNA factors expressed in

germ cells is a common feature of germline-expressed genes, we used publicly available RNA-

seq data from FACS-sorted vasa-GFP ovaries to identify genes specifically expressed in either

germ cells or somatic cells of the ovary. First, using differential RNA-seq analysis of FACS-

sorted vasa-GFP+ (germline) and vasa-GFP- (somatic) cells from cultured *D.mel* ovary GSC

line (vasa-GFP;hs-bam;bamΔ86) (GSE119862) (40), we defined germline-enriched (vasa-

GFP+; log2FC>0, p-val<0.01) and soma-enriched (vasa-GFP-; log2FC<0, p-val<0.01) genes

(Fig. 4A and B). Next, using RNA-seq data from ovarian somatic cells (OSCs) (GSE160860)

(41), we further filtered the soma-enriched genes that are highly expressed in OSCs.

Additionally, we further filtered the germline-enriched genes that show expression in early fly

embryos (0-2hr, modENCODE). In this way, we generated a stringent list of 107 genes

specifically expressed in germ cells and of 59 soma-expressed genes (Supplementary Dataset

1).

We then analysed sequence change accumulation in promoters using PhyloP data and

calculated d^P for all *D. melanogaster* genes and genes expressed specifically in somatic cells

or the germline (Fig. 4C and D) or the average (Fig 4E). Germ cell-specific expressed genes

significantly accumulated changes (d^P) in their promoters compared to soma-expressed genes

(Fig. 4F). Comparison of the average promoter PhyloP scores and d^P conservation scores

between the genome average and somatic or germ cell-specifically expressed genes equally

revealed a significant increase in germ cell gene promoter nucleotide changes compared to the

genome average (Fig. 4G and H).

Since rapid promoter evolution of germline piRNA silencing factors correlated with

gene expression divergence in our previous analysis, we hypothesized that this was a general

feature of germ cell genes. To test this, we compared gene expression divergence between

somatic and germ cell-specific genes using publicly available RNA-seq data of *D.*

melanogaster and *D. yakuba* ovaries (Fig. 4I) (37-39). Notably, genes specifically expressed

in germ cells showed significantly higher divergence in expression compared to soma-

expressed genes (Fig. 4I). When d^P conservation scores were plotted against the gene

expression change between *D. melanogaster* and *D. yakuba*, the germ cell-specifically

expressed genes showed a higher divergence in expression especially in the higher d^P score

groups (>50-75, >75-100) compared to soma-expressed genes, which showed relatively

conserved expression between *D. melanogaster* and *D. yakuba* (Fig. 4J-L).

Discussion

Through analysis of polymorphisms within and between closely related *Drosophila*

species, we show that at the individual gene level, germline and somatic transposon silencing

genes and their regulators do not generally evolve through their coding regions. However, we

identify hot spots for rapid evolution in promoter regions (-30 to -380 nucleotides upstream of

the TSS) of germ cell-specific (dual-strand clusters and the ping-pong cycle), but not soma-

expressed piRNA genes (cTGS and piRNA biogenesis). Our analysis suggests that this mode

of evolution in the germline pathway could be a general feature of at least insect evolution.

Overall, our results point towards a key role for rapid evolution of gene promoters in the germ

cell-specific piRNA pathway. Intriguingly, the soma-specific Yb is replaced in germ cells with

BoYb and in contrast to its somatic cell counterpart, BoYb also displays a fast-evolving

promoter.

Transposon mobility has been attributed to the accumulation of sequence changes in

promoter regions because of the presence of open chromatin around the transcription start site

of genes (42). Interestingly, in core NPC genes, rapid evolution mostly led to accumulation of

indels (19, 20), while piRNA genes display equal accumulation of indels and base changes.

Likely, indels have more profound effects on changes in transcription than single nucleotide

alterations. This feature might reflect that compromised transposon silencing causes sterility,

hence not allowing for substantial changes in expression in piRNA processing genes. Likewise,

stoichiometric changes in protein complexes can drastically alter complex functionality, e.g.

the male-specific lethal (MSL) complex binds less to its targets when one component of the

complex is missing (43, 44). Hence, changes in the protein levels of individual piRNA factors

likely have dominant effects on their capacity to silence transposon activity.

The occurrence of novel transposons has fundamental impact on a species reproductive

success through a phenomenon call hybrid dysgenesis (45). Here, if a novel transposable

element is transmitted by the male, female fertility is severely compromised and can result in

sterility as a result of missing piRNA silencing of the novel invader. Essential to combat these

novel active transposable elements is the ping-pong cycle, primed by transcripts from the novel

transposon. Accordingly, females can be become resistant to such novel transposons over time

through RNA transgenerational inheritance of piRNAs to build up ping-pong cycle

amplification in germ cells (9, 10, 46).

The pattern of changes observed in fast evolving promoters like germline specific

piRNA factors and Nups resembles the outcome of a *P*-element mutagenesis experiment in the

egghead (*egh*) locus coding for a glycosphingolipid biosynthesis enzyme (47). Here, multiple

base changes were observed in the region of the first promoter after *P*-element mutagenesis,

but not an actual *P*-element insert, resulting in a sex peptide insensitive allele *egh^cm^*. Mutations

in the *egh* gene result in pleiotropic phenotypes and the *egh^cm^* allele disrupts neuronal wiring

required for the female post-mating response and development of the optic lobe (21, 47, 48).

Since establishing transposon resistance involves forced selection, genes whose

expression changes result in pleiotropic effect, like channel Nup54, could combine adaptations

in piRNA processing with changes in neuronal wiring resulting in altered behavioural

preferences (19). Of note, in *D. simulans*, a very close relative of *D. melanogaster* projections

of *fruitless* P1 sensory neurons that control courtship have changed and alter mate preference

(49).

The severe impact of hybrid dysgenesis on fertility likely limits fast evolution to species

with high numbers of progeny. Perhaps this can explain the differences seen when comparing

insect and mammalian promoters that are generally more conserved (25). However, rapid

promoter evolution has also been observed in primate promoters, suggesting a common mode

of evolution that has yet to be explored (19, 20, 25, 50).

Methods

Sequence/data retrieval and alignment

Gene and promotor sequences were retrieved from UCSC Genome Browser using the

UCSC Table Browser sequence retrieval tool (32, 33). A standardised region of 2000 bases

upstream of the annotated gene TSS was exported for each gene to ensure inclusion of promoter

regions. Sequences were imported and aligned with clustalW in MEGA11 (51). PhyloP27way

data were sourced from UCSC genome browser (genome.ucsc.edu) through the Table Browser

tool (32, 33). Data points were collected for a region of 1000 nucleotides upstream and 300

nucleotides downstream for each of the analysed genes. Genes without full species coverage

(where ≥1 species had no conserved genomic sequence) or PhyloP coverage (where ≥1

nucleotide in gene or upstream regions had no homologous sequence in the 27 related species)

were removed from the analysis.

Molecular evolution of open reading frames analysis

Analysis of adaptive protein evolution was performed via MKTs (31) using the PopFly

online database (imkt.uab.cat) which uses Drosophila Genome Nexus project sequence data

(52-54). Polymorphism and divergence data were collected from the ancestral Congo

population, chosen since it is a sub-Saharan population with comparably high ancestral stability

compared to other populations. The 'Standard MKT' test was used to calculate results for

individual genes. Fisher's Exact Test with significance defined as Bonferroni corrected $p \leq 0.05$

was used to calculate significance between nonsynonymous to synonymous polymorphisms

within *D. melanogaster* or between *D. melanogaster* and *D. simulans*.

Identification of promotor hot spots and comparison of substitution rates

All gene models were manually inspected in FlyBase using the JBrowse browser and

dominant transcripts were chosen based on comparative modENCODE expression data (55).

Gene and promoter sequence alignments were trimmed to -1000/+300 nucleotides from the

TSS of each gene. Alignments were translated into 'events' for each species compared to *D.*

melanogaster, where 0 signified no measurable sequence difference between all analysed

species and 1 signified sequence divergence (56). Events were calculated for base changes and

indels concatenated, as well as base changes and indels individually. The sum of events was

calculated for concatenated gene groups and individual genes at each nucleotide position. A

sliding event (*Se*) score was calculated from this using a sliding window of five bases along

the sequence, from which heatmaps were generated. To calculate the percentage of events

greater than the average control promoter *Se* score (*d*), a 350-nucleotide region upstream of the

estimated TATA box region was analysed where the total number of *Se* scores exceeding the

control group average sliding event score (Se^C) was divided by the total number of events in

that region (*N*).

$$d = \frac{\text{Total number of } Se \text{ events where } Se > Se^C}{N} \text{x}100$$

351   To calculate the promoter region *d* scores from PhyloP data, the total number of PhyloP

352   (*p*) scores in the 350 nucleotide regions less than the control group average promoter region

353   (p^C) took the place of the total number of *Se* events where *Se* is greater than Se^C (see equation

354   below).

$$d^P = \frac{\text{Total number of } p \text{ scores} < p^C}{N} \text{x}100$$

Significance was calculated using non-parametric chi-squared tests compared to the

control group *d* score. Significance values where $p \leq 0.05$ with Bonferroni correction were

considered statistically significant.

Identification of somatic and germ cell-specifically expressed genes in *D. melanogaster*

We defined somatic and germline genes using several steps. First, using processed RNA-seq

data from vasa-GFP+/- cells sorted from cultured ovary GSC line (vasa-GFP;hs-bam;bamΔ86)

(GSE119862) (40), we defined germline-enriched (vasa-GFP+) using logFC>0, p-value<0.01

parameters and soma-enriched (vasa-GFP-) genes using logFC<0, p-value<0.01 parameters

(Figures 4A and 4B). Next, using processed RNA-seq data from fly ovaries (modENCODE),

we filtered genes that are expressed in adult fly ovaries using RPKM>10 parameter for both

soma and germline genes. To stringently define the germline genes, we additionally used

processed RNA-seq data from early fly embryos (0-2hr, modENCODE; mRNA-seq fly embryo

0-2hr; RPKM; mE_mRNA_em0-2hr; FBlc0000086) to further filter the germline-enriched

genes using >9 RPKM parameter. To stringently define the somatic-specific genes, we

additionally used processed RNA-seq data from ovarian somatic cells (OSCs) (n=4,

GSE160860) (41) and filtered the soma-enriched genes using >20 RPKM to select genes that

are expressed in OSCs as well as parameter <2 RPKM for early fly embryo (0-2hr,

modENCODE) to filter out genes that are expressed in early embryos. This analysis generated

a list of 107 germ cell-specifically expressed genes and of 59 somatic cell-specifically

expressed genes (Fig. 4A and B, Supplementary Dataset 1).

Comparative gene expression analysis between *D. melanogaster* and *D. yakuba*

Using publicly available raw RNA-seq data from *D. melanogaster* (n=3) and *D. yakuba* (n=4)

ovaries, we trimmed adapter sequences from the raw reads using Cutadapt tool (v1.18, default

parameters) and aligned the trimmed reads to the genome assemblies (dm6 and droYak2) using

the RNA-seq aligner STAR (v2.7.3a). Gene counts were quantified using the featureCounts

tool (Subread package v1.5.3). Orthologues genes were identified using blastDm2FB in UCSC

Table Browser. Gene count normalization and differential gene expression analysis between

the species were performed using DESeq2 (v1.30.1) with an additional species-specific gene

lengths normalization step.

Supplementary information

Supplementary information includes Fig. S1-4, Table S1 and Supplementary Dataset 1.

Acknowledgements

We thank the Soller lab for helpful discussion.

Funding

This work was supported by the Medical Research Council Integrated Midlands Partnership

for Biomedical Training Program to D.W.J.M. and the Biotechnology and Biological Science

Research Council to M.S. Work by B.C.N and A.A. was funded by Cancer Research UK

(A21143) and the Wellcome Trust (110161/Z/15/Z).

Authors' contributions

D.W.J.M and M.S. conceived and directed the project. D.W.J.M. and A.A. analysed data.

D.W.J.M. wrote the initial draft and all authors contributed to the final manuscript. All authors

read and approved the final manuscript.

Availability of data and materials

All data generated or analysed during this study are included in the supplementary information

files.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

References

1.   Iwakawa H-o, Tomari Y. Life of RISC: Formation, action, and degradation of RNA-

induced silencing complex. Molecular Cell. 2022;82(1):30-43.

2.   Czech B, Munafò M, Ciabrelli F, Eastwood EL, Fabry MH, Kneuss E, et al. piRNA-

Guided Genome Defense: From Biogenesis to Silencing. Annual Review of Genetics.

2018;52(1):131-57.

3.   Yamashiro H, Siomi MC. PIWI-Interacting RNA in Drosophila: Biogenesis,

Transposon Regulation, and Beyond. Chemical Reviews. 2018;118(8):4404-21.

4.   Slotkin RK, Martienssen R. Transposable elements and the epigenetic regulation of

the genome. Nature Reviews Genetics. 2007;8(4):272-85.

5.   Malone CD, Brennecke J, Dus M, Stark A, McCombie WR, Sachidanandam R, et al.

Specialized piRNA Pathways Act in Germline and Somatic Tissues of the Drosophila Ovary.

Cell. 2009;137(3):522-35.

6.   Khurana JS, Theurkauf W. piRNAs, transposon silencing, and Drosophila germline

development. Journal of Cell Biology. 2010;191(5):905-13.

7.   Senti K-A, Brennecke J. The piRNA pathway: a fly's perspective on the guardian of

the genome. Trends in Genetics. 2010;26(12):499-509.

8.   Sakakibara K, Siomi MC. The PIWI-Interacting RNA Molecular Pathway: Insights

From Cultured Silkworm Germline Cells. BioEssays. 2018;40(1):1700068.

9.   Moelling K. Epigenetics and transgenerational inheritance. J Physiol. 2023.

10.   Grentzinger T, Armenise C, Brun C, Mugat B, Serrano V, Pelisson A, et al. piRNA-

mediated transgenerational inheritance of an acquired trait. Genome Res. 2012;22(10):1877-

88.

11.   Wang X, Ramat A, Simonelig M, Liu M-F. Emerging roles and functional

mechanisms of PIWI-interacting RNAs. Nature Reviews Molecular Cell Biology.

2023;24(2):123-41.

12.   Sato K, Siomi MC. The piRNA pathway in Drosophila ovarian germ and somatic

cells. Proceedings of the Japan Academy, Series B. 2020;96(1):32-42.

13.   Czech B, Hannon GJ. One Loop to Rule Them All: The Ping-Pong Cycle and piRNA-

Guided Silencing. Trends in Biochemical Sciences. 2016;41(4):324-37.

14.   Haase AD. An introduction to PIWI-interacting RNAs (piRNAs) in the context of

metazoan small RNA silencing pathways. RNA Biology. 2022;19(1):1094-102.

15.   Czech B, Preall J, B. , McGinn J, Hannon G, J. . A Transcriptome-wide RNAi Screen

in the Drosophila Ovary Reveals Factors of the Germline piRNA Pathway. Molecular Cell.

2013;50(5):749-61.

16.   Brown JS, Zhang D, Chen W, Lee H-C. Sensitized piRNA reporter identifies multiple

RNA processing factors involved in piRNA-mediated gene silencing. 2023.

17.   Handler D, Meixner K, Pizka M, Lauss K, Schmied C, Franz, et al. The Genetic

Makeup of the Drosophila piRNA Pathway. Molecular Cell. 2013;50(5):762-77.

18.   Munafò M, Lawless VR, Passera A, Macmillan S, Bornelöv S, Haussmann IU, et al.

Channel nuclear pore complex subunits are required for transposon silencing in Drosophila.

eLife. 2021;10.

19.   Nallasivan MP, Haussmann IU, Civetta A, Soller M. Channel nuclear pore protein 54

directs sexual differentiation and neuronal wiring of female reproductive behaviors in

Drosophila. BMC Biology. 2021;19(1).

20.   McQuarrie DWJ, Read AM, Stephens FHS, Civetta A, Soller M. Indel driven rapid

evolution of core nuclear pore protein gene promoters. Scientific Reports. 2023;13(1).

21.   Haussmann IU, Hemani Y, Wijesekera T, Dauwalder B, Soller M. Multiple pathways

mediate the sex-peptide-regulated switch in female Drosophila reproductive behaviours. Proc

Biol Sci. 2013;280(1771):20131938.

22.   Tang S, Presgraves DC. Evolution of the Drosophila Nuclear Pore Complex Results

in Multiple Hybrid Incompatibilities. Science. 2009;323(5915):779-82.

23.   Tang S, Presgraves DC. Lineage-Specific Evolution of the Complex Nup160 Hybrid

Incompatibility Between Drosophila melanogaster and Its Sister Species. Genetics.

2015;200(4):1245-54.

24.   Iwasaki YW, Siomi MC, Siomi H. PIWI-Interacting RNA: Its Biogenesis and

Functions. Annu Rev Biochem. 2015;84:405-33.

25.   Taylor MS, Kai C, Kawai J, Carninci P, Hayashizaki Y, Semple CAM. Heterotachy in

Mammalian Promoter Evolution. PLoS Genetics. 2006;2(4):e30.

26.   Balacco DL, Soller M. The m(6)A Writer: Rise of a Machine for Growing Tasks.

Biochemistry. 2019;58(5):363-78.

27.   Bawankar P, Lence T, Paolantoni C, Haussmann IU, Kazlauskiene M, Jacob D, et al.

Hakai is required for stabilization of core components of the m6A mRNA methylation

machinery. Nature Communications. 2021;12(1):3778.

28.   Van Lopik J, Trapotsi M-A, Hannon GJ, Bornelov S, Czech Nicholson B. Unistrand

piRNA clusters are an evolutionary conserved mechanism to suppress endogenous

retroviruses across the Drosophila genus. bioRxiv. 2023.

29.   Fridrich A, Moran Y. Some flies do not play ping-pong. PLOS Biology.

2023;21(6):e3002152.

30.   Chary S, Hayashi R. Mechanistic divergence of piRNA biogenesis in Drosophila.

2022.

31.   McDonald JH, Kreitman M. Adaptive protein evolution at the Adh locus in

Drosophila. Nature. 1991;351(6328):652-4.

32.   Karolchik D. The UCSC Table Browser data retrieval tool. Nucleic Acids Research.

2004;32(90001):493D-6.

33. Kent WJ, Sugnet CW, Furey TS, Roskin KM, Pringle TH, Zahler AM, et al. The

Human Genome Browser at UCSC. Genome Research. 2002;12(6):996-1006.

34. Consortium GO. The Gene Ontology resource: enriching a GOld mine. Nucleic Acids

Res. 2021;49(D1):D325-d34.

35. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene

ontology: tool for the unification of biology. The Gene Ontology Consortium. Nat Genet.

2000;25(1):25-9.

36. Mi H, Muruganujan A, Ebert D, Huang X, Thomas PD. PANTHER version 14: more

genomes, a new PANTHER GO-slim and improvements in enrichment analysis tools.

Nucleic Acids Research. 2018;47(D1):D419-D26.

37. Sanfilippo P, Wen J, Lai EC. Landscape and evolution of tissue-specific alternative

polyadenylation across Drosophila species. Genome Biology. 2017;18(1).

38. Smolko AE, Shapiro-Kulnane L, Salz HK. The H3K9 methyltransferase SETDB1

maintains female identity in Drosophila germ cells. Nature Communications. 2018;9(1).

39. Vankuren NW, Vibranovski MD. A Novel Dataset for Identifying Sex-Biased Genes

in Drosophila. Journal of Genomics. 2014;2:64-7.

40. Story B, Ma X, Ishihara K, Li H, Hall K, Peak A, et al. Defining the expression of

piRNA and transposable elements in Drosophila ovarian germline stem cells and somatic

support cells. Life Science Alliance. 2019;2(5):e201800211.

41. Eastwood EL, Jara KA, Bornelöv S, Munafò M, Frantzis V, Kneuss E, et al.

Dimerisation of the PICTS complex via LC8/Cut-up drives co-transcriptional transposon

silencing in Drosophila. eLife. 2021;10:e65557.

42. Treiber CD, Waddell S. Transposon expression in the Drosophila brain is driven by

neighboring genes and diversifies the neural transcriptome. Genome Research.

2020;30(11):1559-69.

43.   Laverty C, Lucci J, Akhtar A. The MSL complex: X chromosome and beyond.

Current Opinion in Genetics & Development. 2010;20(2):171-8.

44.   Straub T, Becker PB. Dosage compensation: the beginning and end of generalization.

Nature Reviews Genetics. 2007;8(1):47-57.

45.   Brennecke J, Malone CD, Aravin AA, Sachidanandam R, Stark A, Hannon GJ. An

Epigenetic Role for Maternally Inherited piRNAs in Transposon Silencing. Science.

2008;322(5906):1387-92.

46.   Luo Y, He P, Kanrar N, Fejes Toth K, Aravin AA. Maternally inherited siRNAs

initiate piRNA cluster formation. Mol Cell. 2023.

47.   Soller M, Haussmann IU, Hollmann M, Choffat Y, White K, Kubli E, et al. Sex-

peptide-regulated female sexual behavior requires a subset of ascending ventral nerve cord

neurons. Curr Biol. 2006;16(18):1771-82.

48.   Fan Y, Soller M, Flister S, Hollmann M, Müller M, Bello B, et al. The egghead gene

is required for compartmentalization in Drosophila optic lobe development. Dev Biol.

2005;287(1):61-73.

49.   Seeholzer LF, Seppo M, Stern DL, Ruta V. Evolution of a central neural circuit

underlies Drosophila mate preferences. Nature. 2018;559(7715):564-9.

50.   Main BJ, Smith AD, Jang H, Nuzhdin SV. Transcription Start Site Evolution in

Drosophila. Molecular Biology and Evolution. 2013;30(8):1966-74.

51.   Tamura K, Stecher G, Kumar S. MEGA11: Molecular Evolutionary Genetics

Analysis Version 11. Molecular Biology and Evolution. 2021;38(7):3022-7.

52.   Lack JB, Cardeno CM, Crepeau MW, Taylor W, Corbett-Detig RB, Stevens KA, et

al. The Drosophila Genome Nexus: A Population Genomic Resource of 623 Drosophila

melanogaster Genomes, Including 197 from a Single Ancestral Range Population. Genetics.

2015;199(4):1229-41.

53.   Lack JB, Lange JD, Tang AD, Corbett-Detig RB, Pool JE. A Thousand Fly Genomes:

An Expanded Drosophila Genome Nexus. Mol Biol Evol. 2016;33(12):3308-13.

54.   Mackay TFC, Richards S, Stone EA, Barbadilla A, Ayroles JF, Zhu D, et al. The

Drosophila melanogaster Genetic Reference Panel. Nature. 2012;482(7384):173-8.

55.   Gramates LS, Agapite J, Attrill H, Calvi BR, Crosby MA, dos Santos G, et al.

FlyBase: a guided tour of highlighted features. Genetics. 2022;220(4).

56.   Tang H, Lewontin RC. Locating Regions of Differential Variability in DNA and

Protein Sequences. Genetics. 1999;153(1):485-95.

Table and Figure Legends

Fig. 1: Promoter regions of germ cell-expressed piRNA processing factors are hot spots

for rapid evolution

A and B) Schematic depiction of somatic piRNA factors involved in co-transcriptional gene

silencing and piRNA biogenesis (A) and germline-specific piRNA pathway genes involved in

dual-strand cluster regulation and the ping-pong cycle (B) in the *Drosophila* ovary.

C and D) Evolutionary conservation per nucleotide around the TSS depicted by PhyloP27way

conservation score averages for the *D. melanogaster* dm6 genome (black) and the somatic

(grey, C) or the germ cell-specific piRNA factors (grey, D).

E and F) PhyloP27way conservation score averages (E) and PhyloP27way conservation

promoter d^P scores (F) for the 350 nucleotide promoter regions of the somatic piRNA factors

and the germ cell-specific piRNA factors compared to all genes in the *Drosophila* genome.

Statistically significant differences from unpaired student t-tests (E) and non-parametric chi-

squared tests (F) are indicated by asterisks (*** p≤0.001, **** p≤0.0001 following Bonferroni

correction).

G-L) Heatmaps indicating sequence (E, purple), indel (G, blue) or base change (I, green)

accumulation, and their quantification (F-J) depicted as d scores for each of the analysed gene

groups compared to the control among closely related *D. melanogaster*, *D. simulans*, *D.*

sechellia, *D. yakuba* and *D. erecta*. Regions of 1000 nucleotides upstream and 300 nucleotides

downstream of the TSS were analysed. Analysis was performed for the control group (m^6A

writer complex and readers), the somatic piRNA factors, and the germ cell-specific piRNA

factors. The blue line indicates the promoter region used for quantification of the substitution

rate based on the gene and intergenic region schematic. Statistically significant differences

from non-parametric chi-squared tests are indicated by asterisks (**** p≤0.0001 following

Bonferroni correction).

Fig. 2: Promoters of genes required for germline transposon silencing evolve fast

A) PhyloP27way nucleotide scores for the top 100 genes affecting transposon silencing in germ

cells based on transposon derepression z scores and ordered by promoter conservation d^P

scores. A region of 1000 nucleotides upstream and 300 nucleotides downstream are shown.

Red represents lower conservation while blue represents higher.

B) Average PhyloP27way nucleotide scores for the top 100 genes affecting transposon

silencing in germ cells compared to the genome average.

C) Comparison of PhyloP27way conservation promoter scores (d^P) for each of the 100 genes

affecting transposon silencing in germ cells compared to the genome average. Statistically

significant differences from non-parametric chi-squared tests are indicated by asterisks (***

p≤0.001 following Bonferroni correction).

D) Representative GO analysis for the top 100 gene candidates affecting transposon silencing.

E) Individual gene comparison of PhyloP27way conservation promoter scores (d^P) for the top

100 genes affecting transposon silencing in germ cells compared to the genome average.

Statistically significant differences from non-parametric chi-squared tests are indicated by

asterisks (* p≤0.05, ** p≤0.01, *** p≤0.001, **** p≤0.0001 following Bonferroni correction).

F-H) Sequence analysis for change accumulation among closely related *D. melanogaster*, *D.*

simulans, *D. sechellia*, *D. yakuba* and *D. erecta*. Comparisons were performed for sequence

change (F), indel (G), and base change (H) *d* scores for each of the positive 49 genes affecting

transposon silencing in FEPG gene group compared to the somatic piRNA factors. Statistically

significant differences from non-parametric chi-squared tests are indicated by asterisks (****

p≤0.0001 following Bonferroni correction).

I) Comparison of *d* scores for protein complex genes involved in germ cell transposon silencing

compared to the somatic piRNA factors. Statistically significant differences from non-

parametric chi-squared tests are indicated by asterisks (**** p≤0.0001 following Bonferroni

correction).

J) Heatmaps indicating sequence change accumulation for protein complex genes involved in

germ cell transposon silencing compared to the piRNA factors shown in purple among closely

related *D. melanogaster*, *D. simulans*, *D. sechellia*, *D. yakuba* and *D. erecta*. Regions of 1000

nucleotides upstream and downstream of the TSS were analysed. The blue line indicates the

promoter region used for quantification of the substitution rate.

Fig. 3: Divergent expression of FEPG genes in fly ovaries is accompanied by rapid

promoter evolution

A) Volcano plot showing differential gene expression (DESeq2) between *D. melanogaster* and

D. yakuba ovaries (RNA-seq data; Supplementary Dataset 1) for FEPG (blue) and the somatic

piRNA factors (red). Horizontal dashed lines indicate significance thresholds (adjusted

p≤0.05).

B-D) Alignment of the promoter regions of *CG32152, MEP-1* and *Rbbp5* for *D. melanogaster*

and *D. yakuba* with indels and base changes indicated as black boxes. The black lines indicate

gene 5'UTRs.

Fig. 4: Germ cell-specifically expressed, but not somatic cell-specifically expressed gene

promoters evolve rapidly and are accompanied by divergence in gene expression

A) Ranking of genes using vasa-GFP signal from FACS-sorted vasa-GFP+/- cells of cultured

ovary GSC line (vasa-GFP;hs-bam;bamΔ86) (GSE119862) that defines germline-enriched

(vasa-GFP+) and soma-enriched (vasa-GFP-) genes.

B) Volcano plot showing differential gene expression between the FACS-sorted vasa-GFP+

(germline) and vasa-GFP- (somatic) cells from cultured ovary GSC line (vasa-GFP;hs-

bam;bamΔ86) (GSE119862).

C and D) PhyloP27way nucleotide scores for somatic cell-specific (C) and germ cell-specific

(D) expressed genes ordered by promoter conservation d^P scores. A region of 1000 nucleotides

upstream and 300 nucleotides downstream are shown. Red represents lower conservation while

blue represents higher.

E) Average PhyloP27way nucleotide scores for all genes in the *Drosophila* genome, somatic

cell and germ cell-specific genes. Purple represents lower conservation while blue represents

higher.

F) The percentage of somatic cell and germ cell-specific genes with significant promoter

nucleotide changes (d^P) versus all genes in the *Drosophila* genome average, divided by

significantly increased (low conservation) or decreased (high conservation). Statistically

significant differences from non-parametric chi-squared tests are indicated by asterisks (****

p≤0.0001 following Bonferroni correction).

G and H) PhyloP27way conservation score averages (G) and PhyloP27way conservation

promoter scores (d^P) (H) for the 350 nucleotide promoter regions compared for all genes in the

Drosophila genome, somatic cell-specifically expressed and germ cell-specifically expressed

genes. Statistically significant differences from unpaired student t-tests (G) and from non-

parametric chi-squared tests (H) are indicated by asterisks (** p≤0.01, **** p≤0.0001

following Bonferroni correction).

I) Volcano plot of differential gene expression (DESeq2) between *D. melanogaster* and *D.*

yakuba ovaries (RNA-seq data; Supplementary Dataset 1) for all (grey), germ cell-specific

(blue) and somatic cell-specific (red) gene groups. Horizontal dashed lines indicate

significance (adjusted p≤0.05).

J) Scatterplot of individual gene d^P scores plotted against fold-change expression differences

between *D. melanogaster* and *D. yakuba* ovaries (RNA-seq data; Supplementary Dataset 1) for

germ cell-specific (blue) and somatic cell-specific (red) gene groups. The grey dotted lines

indicate log2Fold-change threshold values of ≥0.5 and ≤0.5.

K and L) Comparison of the percentage of genes with significant changes in *D. melanogaster*

log2FoldChange (≥0.5, K) or significant changes in *D. yakuba* log2FoldChange (≤0.5, L)

expression changes in PhyloP conservation promoter score (d^P) ranges 25-50, >50-75, and >75-

100. Expression changes for germ cell-specific (blue) and somatic cell-specific (red) gene

groups were analysed between *D. melanogaster* and *D. yakuba*. Statistically significant

differences from non-parametric chi-squared tests are indicated by asterisks (** p≤0.01, ***

p≤0.001, **** p≤0.0001 following Bonferroni correction).

Figure 1

Figure 2

<div align="center">**Supplementary Materials**</div>

Promoters of germline transposon silencing genes evolve rapidly

accompanied by diverging gene expression

DAVID W. J. MCQUARRIE[1,2], AZAD ALIZADA[3], BENJAMIN CZECH

NICHOLSON[3] AND MATTHIAS SOLLER[1,2]*

Supplementary Fig. 1: Coding region evolution rates for somatic and germline piRNA

processing genes.

A) Evolution of coding regions analysed by 'Standard MKT' tests for polymorphisms and

divergence between *D. melanogaster* and *D. simulans* for core genes in the somatic and germ

cell transposon silencing pathways. R, replacement; S, synonymous. Analyses were omitted

for genes without a Popfly record. Statistically significant differences from Fisher's Exact Tests

are indicated by asterisks (* $p \leq 0.05$ following FDR correction).

B) Individual gene sequence change, indel, and base change *d* scores and statistically

significant accumulation of sequence changes for the somatic piRNA factors and germ cell-

specific piRNA factors. Statistically significant differences from non-parametric chi-squared

tests are indicated by asterisks (* $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$ following

FDR correction).

Supplementary Fig. 2: Genes coding for members of a subgroup of protein complexes

implicated in piRNA silencing accumulate sequence changes in their promoters

Individual gene *d* scores and statistically significant accumulation of sequence changes for each

gene separated by protein complex compared to the somatic piRNA factors. Statistically

significant differences from non-parametric chi-squared tests are indicated by asterisks (*

$p \leq 0.05$, ** $p \leq 0.01$, **** $p \leq 0.0001$ following Bonferroni correction).

Supplementary Fig. 3: Promoter evolution of protein complex genes involved in germ cell

transposon silencing

A and C) Heatmaps indicating indel (A, blue) or base change (C, green) accumulation in protein

complex genes involved in germ cell transposon silencing compared to Yb body genes among

closely related *D. melanogaster*, *D. simulans*, *D. sechellia*, *D. yakuba* and *D. erecta*. Regions

of 1000 nucleotides upstream and 300 nucleotides downstream of the TSS were analysed. The

blue line indicates the promoter region used for quantification of the substitution rate.

B and D) Comparison of indel (B, blue) or base change (D, green) *d* scores for each of the

protein complex gene groups compared to the somatic piRNA factors. Statistically significant

differences from non-parametric chi-squared tests are indicated by asterisks (* $p \leq 0.05$, **

$p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$ following Bonferroni correction).

E) Individual gene *d* scores and statistically significant accumulation of indels or base changes

for each gene separated by protein complex compared to the somatic piRNA factors.

Comparison of indel (B, blue) or base change (D, green) *d* scores for each of the protein

complex gene groups compared to the control group. Statistically significant differences from

non-parametric chi-squared tests are indicated by asterisks (* $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$,

**** $p \leq 0.0001$ following Bonferroni correction).

Supplementary Fig. 4: Coding sequences evolution of protein complex genes involved in

germ cell transposon silencing.

A) Impact of protein complex genes involved in germ cell transposon silencing on transposon

de-repression. Unprocessed data were taken from (1). Derepression scores (z) were

concatenated for protein complexes in each transposon type (HeTA, TAHRE, blood, burdock)

and an average of all.

B) Evolution of coding regions analysed by 'Standard MKT' tests for polymorphisms and

divergence between *D. melanogaster* and *D. simulans* for individual genes from the selected

germline transposon silencing implicated protein complexes. R, replacement; S, synonymous.

Analyses were omitted for genes without a Popfly record. Statistically significant differences

from Fisher's Exact Tests are indicated by asterisks (** p≤0.01 following FDR correction).

References

1.	Czech B, Preall J, B. , McGinn J, Hannon G, J. . A Transcriptome-wide RNAi Screen

in the Drosophila Ovary Reveals Factors of the Germline piRNA Pathway. Molecular Cell.

2013;50(5):749-61.

A

	Gene	Polymorphism			Divergence			Fisher's exact *P*-value (FDR)	
		R	S	R/S	R	S	R/S		
Germ cell-specific piRNA factors	*aub*	10	9	1.111	96	33	2.909	0.35700	
	tej	10	6	1.667	73	32	2.281	1.01238	
	ago3	2	1	2.000	54	41	1.317	1.00000	
	qin	33	28	1.179	175	79	2.215	0.26175	
	squ	2	1	2.000	26	16	1.625	1.03448	
	tapas	14	12	1.167	122	55	2.218	0.59603	
	BoYb	3	1	3.000	102	50	2.040	1.07143	
	krimp	19	16	1.188	126	25	5.040	0.01500	*
	spn-E	6	14	0.429	82	65	1.262	0.34100	
	Boot	3	0	-	51	18	2.833	1.06520	
	moon	0	0	-	0	0	-	1.11111	
	Nxf3	7	2	3.500	69	23	3.000	1.15385	
	cuff	3	3	1.000	52	18	2.889	0.78053	
	rhi	14	4	3.500	91	20	4.550	1.24064	
	del	7	5	1.400	152	44	3.455	0.68032	
Somatic piRNA factors	*armi*	4	12	0.333	45	32	1.406	0.38400	
	papi	2	8	0.250	9	30	0.300	1.20000	
	vret	12	12	1.000	34	30	1.133	1.28700	
	Gasz	3	1	3.000	25	8	3.125	1.25000	
	daed	5	1	5.000	13	16	0.813	0.66532	
	mino	4	7	0.571	15	23	0.652	1.30435	
	zuc	0	1	0.000	28	16	1.750	0.80957	
	shu	3	2	1.500	22	22	1.000	1.36364	
	SoYb	76	27	2.815	86	52	1.654	0.35800	
	Yb	15	8	1.875	130	38	3.421	0.60810	
	arx	4	1	4.000	6	3	2.000	1.42857	
	piwi	9	12	0.750	23	46	0.500	0.89020	
	panx	7	2	3.500	62	26	2.385	1.50000	
	nxf2	3	1	3.000	28	44	0.636	0.74580	
	mael	2	2	1.000	45	14	3.214	0.72277	

B

A

B

		Polymorphism			Divergence				
	Gene	R	S	R/S	R	S	R/S	Fisher's exact *P*-value (FDR)	
Nuclear CBC	*Ars2*	9	6	1.500	29	47	0.617	0.77150	
	cbp80	0	14	0.000	56	49	1.143	0.00292	**
	cbp20	0	0	-	1	10	0.100	1.00000	
THO complex	*thoc5*	13	9	1.444	18	14	1.286	1.02564	
	thoc7	0	0	-	4	7	0.571	1.05263	
	thoc6	8	6	1.333	7	15	0.467	0.70120	
	tho2	2	7	0.286	34	58	0.586	1.21150	
	Hpr1	1	7	0.143	14	30	0.467	1.09893	
Exon junction complex — Inner core	*tsu*	0	0	-	2	6	0.333	1.08108	
	mago	0	3	0.000	0	2	0.000	1.11111	
	CG7483	0	0	-	14	12	1.167	1.14286	
	btz	7	8	0.875	10	32	0.313	0.75267	
Outer shell	*Pnn*	2	0	-	21	4	5.250	1.17647	
	Aly	7	20	0.350	39	18	2.167	0.00533	**
	Acn	6	5	1.200	28	26	1.077	1.21212	
	Bin1	0	0	-	1	8	0.125	1.25000	
	RnpS1	0	1	0.000	185	79	2.342	0.92892	
Transient factors	*Upf3*	4	6	0.667	19	24	0.792	1.29032	
	Upf2	7	32	0.219	36	32	1.125	0.00800	**
	SRm160	5	15	0.333	11	29	0.379	1.33333	
	Nxt1	0	0	-	4	2	2.000	1.37931	
	sbr	2	7	0.286	40	29	1.379	0.72600	
	Hel25E	0	6	0.000	0	8	0.000	1.42857	
NSL complex	*MBD-R2*	9	7	1.286	64	53	1.208	1.48148	
	mof	3	7	0.429	38	33	1.152	0.70364	
	Rcd5	4	10	0.400	9	29	0.310	1.45800	
	dgt1	6	14	0.429	9	12	0.750	1.09558	
	Rcd1	9	14	0.643	61	48	1.271	0.75911	
	nsl1	11	10	1.100	185	78	2.372	0.74000	
SUMO modifiers	*smt3*	0	1	0.000	0	5	0.000	1.53846	
	lwr	0	6	0.000	0	3	0.000	1.60000	
	Aos1	1	1	1.000	6	25	0.240	1.09857	
	Uba2	5	5	1.000	15	26	0.577	1.14400	
	Ulp1	11	6	1.833	328	88	3.727	0.74767	
COMPASS complex	*Set1* (not analysed)	-	-	-	-	-	-	-	
	wds	0	6	0.000	1	13	0.077	1.66667	
	Hcf (not analysed)	-	-	-	-	-	-	-	
	Rbbp5	0	6	0.000	3	9	0.333	1.14378	
	wdr82	0	9	0.000	0	16	0.000	1.73913	
	Cfp1	3	6	0.500	10	22	0.455	1.81818	
	Dpy-30L1	2	1	2.000	9	9	1.000	1.90476	
	ash2	1	13	0.077	9	21	0.429	0.75543	