Main Manuscript for

Genetic Reinstatement of RIG-I in Chickens Reveals Insights into
Avian Immune Evolution and Influenza Interaction

Hicham Sid[1*], Theresa von Heyl[1], Sabrina Schleibinger[1], Romina Klinger[1], Leah Heymelot Nabel[1],
Hanna Vikkula[1], Rodrigo Guabiraba[2], Vanaique Guillory[2], Mohanned Naif Alhussien[1], Brigitte
Böhm[3], Benjamin Schade[3], Daniel Elleder[4], Samantha Sives[5], Lonneke Vervelde[5], Sascha Trapp[2],
Benjamin Schusser[1,6]

[1] TUM School of Life Sciences, Weihenstephan, Department of Molecular Life Sciences,
Reproductive Biotechnology, Technical University of Munich, 85354, Freising, Germany
[2] UMR ISP, INRAE, Université de Tours. 37380, Nouzilly, France
[3] Bavarian Animal Health Service, Department of Pathology, 85586, Poing, Germany
[4] Institute of Molecular Genetics of the Czech Academy of Sciences, Prague, Czech Republic
[5] Division of Immunology, The Roslin Institute and Royal (Dick), School of Veterinary Studies,
University of Edinburgh, UK
[6] Center for Infection Prevention (ZIP), Technical University of Munich, 85354 Freising, Germany
16
*To whom correspondence may be addressed: Hicham Sid

Email: hicham.sid@tum.de

Author Contributions: H.S., T.v.H., S.Schleibinger, R.K., L.H.N., H.V., R.G., V.G., M.A., B.B., B.
Schade, D.E., S. Sives, L.V., S.T., and B. Schusser contributed to the production of experimental
data. H.S., T.v.H., S. Schleibinger, L.H.N., R.G., B.B., B. Schade, D.E., S. Sives, L.V., S.T., and
B. Schusser contributed to experimental design and interpretation of the data. H.S., S.T., and B.
Schusser wrote the manuscript.

Competing Interest Statement: The authors declare no competing interests.

Classification: Biological Sciences, Agricultural Sciences.

Keywords: avian influenza virus, RIG-I, RNF135, transgenic chicken, duck.

This PDF file includes:

Main Text
Figures 1 to 9

Abstract

Retinoic acid-inducible gene I (*RIG-I*) activates mitochondrial antiviral signaling proteins,

initiating the antiviral response. *RIG-I* and *RNF135*, a ubiquitin ligase regulator, are

missing in domestic chickens but conserved in mallard ducks. The chickens' *RIG-I* loss

was long believed to be linked to increased avian influenza susceptibility. We reinstated

both genes in chickens and examined their susceptibility to infection with an H7N1 avian

influenza virus. Uninfected *RIG-I*-expressing chickens exhibited shifts in T and B cells. At the same time, the H7N1 infection led to severe disease, persistent weight loss, and increased viral replication compared to wild-type chickens. The simultaneous expression of *RIG-I* and *RNF135* potentiated the *RIG*-I activity and was associated with exacerbated inflammatory response and increased mortality without influencing virus replication. Additional animal infection experiments with two other avian influenza viruses validated these findings. They confirmed that the harmful effects triggered by *RIG-I* or *RIG-I-RNF135*-expression require a minimum degree of viral virulence. Our data indicate that the loss of *RIG-I* in chickens has likely evolved to counteract deleterious inflammation caused by viral infection and highlight the outcome of restoring evolutionary lost genes in birds.

Significance Statement

The evolutionary loss of a crucial innate immune sensor like *RIG-I* in domestic chickens and its presence in closely related avian species such as ducks has long puzzled researchers. We genetically reinstated *RIG-I* in chickens, alongside its ubiquitination factor, *RNF135*, to uncover their roles in responding to influenza virus interactions in chickens. Our findings suggest that the loss of *RIG-I* in chickens may have occurred as an adaptive strategy to mitigate harmful inflammation associated with influenza infection. We shed light on the outcome of reinstating evolutionarily lost genes in birds and open new avenues for understanding immune responses in vertebrates.

Main Text

Introduction

Avian influenza virus (AIV) is an epizootic pathogen with zoonotic potential (1) that recently caused devastating outbreaks worldwide, leading to the loss of millions of birds due to animal death and culling (2). The ability of the virus to spread between mammals and humans is highly concerning due to the potential risk of pandemics (3, 4). The viral reservoir of AIVs are wild birds of the orders *Anseriformes* (ducks, geese, and swans) and *Charadriiformes* (gulls and terns) (5), which, compared to chickens or other galliform birds, exhibit milder clinical symptoms despite efficient viral replication (6). Certain genomic features of the duck, including a functional retinoic acid-inducible gene I *(RIG-I)* gene, were reported to be associated with their relative resistance to clinical avian influenza

infection (7). *RIG-I* is a cytosolic RNA sensor that recognizes and binds to the 5'
triphosphate end (5'-ppp) (8). It forms a first line of antiviral defense as a pathogen
recognition receptor (PRR) against RNA viruses (9). Upon activation, the *RIG-I* interacts
with the mitochondrial antiviral signaling proteins (MAVS), inducing a pro-inflammatory
antiviral response characterized by the upregulation of type I and type III interferons (IFNs)
followed by the expression of IFN stimulated genes (ISGs) (10). The activity of *RIG-I* is
believed to be controlled by post-translational modification of tripartite motif-containing
protein 25 (*TRIM25*) (11) and RING finger protein 135 (*RNF135*, also known as *Riplet* or
REUL). The latter was found to modify *RIG-I* by lysine 63-linked polyubiquitination of the
C-terminal region of the caspase activation and recruitment domain (CARD) (12), leading
to a stronger *RIG-I* signal transduction (13). The physiological role of mammalian *RIG-I*,
its function in viral infections, and its parallel evolution with other RLRs are well
documented (11, 14-16). Furthermore, the knockout (KO) of *RIG-I* in mice has shown that
this gene may have additional functions related to adaptive immunity (14). *RIG-I* KO mice
exhibited a defect in migratory dendritic cells in combination with a reduced frequency of
polyfunctional effector T cells upon influenza infection (17).

In ducks, *RIG-I* elicits a potent interferon (IFN) response within the first few hours after
infection, leading to survival against most AIV strains (18). In contrast, chickens lack *RIG-I*, which probably lost its function in a common ancestor of galliform birds (7, 19).
Interestingly, a recent study detected disrupted *RIG-I* pseudogenes in some Galliformes,
including the helmeted guineafowl (*N. meleagris*) and the northern bobwhite (*C. virginianus*) (20). Authors hypothesized a compensatory evolution of melanoma
differentiation-associated gene-5 (*MDA5*) that accompanied the gradual loss of *RIG-I* in
chickens (20). The evolutionary loss of *RIG-I* in different galliform birds correlated with the
simultaneous loss of its ubiquitin ligase *RNF135* (20), which has been described to be
critical for *RIG-I* ubiquitination in mammals, unlike *TRIM25*, which is increasingly believed
to be less important for *RIG-I* signaling (21). Reasons behind the loss of *RIG-I* and its
ubiquitination factor in chickens are still unknown and remain enigmatic, especially given
the virus-limiting effect of duck *RIG-I* overexpression in AIV-infected chicken DF-1 cells
(7).

To date, the effect of *in vivo* expression of *RIG-I* in chickens has never been studied, and
no transgenic chicken lines expressing duck *RIG-I* have been generated, likely due to the

lack of suitable biotechnological tools in avian research. Here, we used chicken primordial

germ cells (PGCs) to develop genetically modified chickens expressing duck *RIG-I* and

RNF135 under the control of their respective duck promoters. The generated birds were

healthy and developed normally compared to their WT siblings. In the absence of infection,

we observed an alternative phenotype of the adaptive immune system of *RIG-I*-expressing

chickens, particularly regarding the T cell population. In contrast, the co-expression of

RNF135 with *RIG-I* contributed to a balanced adaptive immune phenotype that appeared

to be similar to WT birds. Infection experiments with an H7N1 AIV led to severe clinical

disease associated with a strong inflammatory response, high *IFN-γ* expression, and

elevated viral replication in *RIG-I*-expressing chickens compared to other challenged

groups. In contrast, infected *RIG-I-RNF135*-expressing chickens presented with

inflammation and a differential expression of *IFN* and pro-inflammatory cytokines

compared to *RIG-I*-expressing chickens. The obtained data reveal the immunological

functions of *RIG-I* in chickens and the benefit of learning from less susceptible species to

influenza infection to improve the immune system's resilience towards infection.

118
119 **Results**
120
Generation of PGCs that express duck *RIG-I* and *RNF135* under the respective duck

promoters

The genetic modification of PGCs represents a crucial step for generating transgenic

chickens. PGCs are the precursors of sperm and eggs in adult animals, and therefore, we

used them to produce chimeric roosters paired with WT layer hens to obtain the desirable

transgene. To generate chickens that express *RIG-I* and *RNF135*, we started by cloning

duck *RIG-I* and *RNF135*, which were subsequently inserted into two different expression

vectors (Fig. 1b). *RIG-I* and *RNF135* were expressed under their respective duck

promoters and used to generate PGCs expressing both genes separately. While the duck

RIG-I promoter was previously described (22), we determined the activity of the duck

RNF135 promoter, which was examined by the generation of different deletion mutants

tested in the NanoDLR^TM Assay System. Therefore, different deletion mutants were

generated, including p1577, p1001, p349, and p197. Since we did not detect a core

promoter activity of the duck *RNF135*, we used the full-length sequence of the duck

RNF135 for the generation of duck *RNF135*-expressing PGCs (Fig. 1a). The assembled

expression vectors that were used to generate PGCs are presented in Fig. 1b.

Due to the unavailability of commercial antibodies for detecting the duck *RIG-I*, we inserted

a FLAG-Tag on the C-terminus to facilitate its detection using anti-FLAG antibodies. The

activity of the duck *RIG-I* was examined by differentiating the *RIG-I*-expressing PGCs into

PGC-derived fibroblasts (Fig. 1c, Supplementary Figure 7c) that were infected with a low

pathogenic avian influenza virus (LPAIV) H9N2 (Fig. 1c). Results showed that the infection

led to activation of *RIG-I* as shown by FLAG-Tag staining. At the same time, the MOCK-

infected cells remained negative for *RIG-I* expression (Fig. 1c).

The expression of *RIG-I* did not limit the viral replication under *in vitro* and *in ovo*

conditions

According to previously published data by Barber et al. (7), the overexpression of duck

RIG-I in chicken DF-1 cells reduced the replication of H5N2 or H5N1 viruses when cells

were infected at an MOI 1. Authors constitutively expressed *RIG-I* under the control of the

CMV promoter using expression vector pcDNA 3.1. Therefore, we wanted to examine if

the expression of *RIG-I* under the control of the duck RIG-I promoter can lead to a similar

effect in limiting virus replication. To this end, we isolated chicken embryonic fibroblasts

(CEFs) and produced embryonated eggs from the generated transgenic chickens and

infected them with LPAIV H9N2. After virus infection, supernatants and allantoic fluid were

collected from CEFs and embryonated eggs, respectively. We quantified the newly

produced viral particles using a focus-forming assay. Surprisingly, transgene expression

did not significantly affect the viral replication in both tested systems: CEFs (Fig. 1d,

Supplementary Figure 10) and embryonated chicken eggs (Fig.1e, Supplementary Figure

13), which was the opposite in the previously published study (7). This highlights the

importance of the chosen promoter, which may have affected the expression of *RIG-I* and

influenced the innate immune response towards influenza virus.

Phenotypic characterization of the genetically modified chickens

Upon the generation of *RIG-I-* and *RNF135*-expressing chicken lines, we wanted to ensure

that the genetic modification did not negatively affect the growth of the generated

transgenic birds. Therefore, we monitored their development by weekly measuring their

body weights (Fig. 2a). Both generated chicken lines showed comparable growth to their

WT siblings. The animals matured sexually, and no harmful phenotype was detected (Fig.

2a and 2b). The expression of both genes was examined via RT-PCR, revealing that both

genes are expressed differentially in various tissues (Fig. 2c). We also detected

comparable levels in the expression levels with the duck (Fig. 2d)

Previous studies show that mammalian *RIG-I* affects adaptive immunity, mainly T cells

(17). Therefore, we sought to examine whether the re-expression of *RIG-I* in chickens will

have similar effects. We analyzed the numbers of peripheral blood mononuclear cells

(PBMCs) using flow cytometry to investigate the possible impact of expressing *RIG-I*,

RNF135, or both on immune cell counts. No differences were observed in the number of

immune cells between *RNF135*-expressing chickens and their WT siblings

(Supplementary Figure 1). However, *RIG-I*-expressing chickens exhibited a significantly

higher number of αβ and γδ T cells as well as B cells in comparison to their WT siblings

(p<0.05) (Fig. 2e). This was not the case for monocyte counts, where no significant

differences were observed (Fig. 2e). In addition, no significant differences were observed

in *RIG-I*-expressing chickens compared to WT birds regarding the levels of IgM and IgY

at 12 and 14 weeks of age (Supplementary Figure 2).

Further analysis of different T cell subpopulations indicated that *RIG-I*-expressing

chickens had a significantly higher number of CD4+T cells, CD8α^{neg}T cells and a

significantly lower number of CD8α+^{high}T cells (p<0.05) (Fig. 3a and 3b). T-cell activation

was quantified after lectin activation using Concanavalin A (ConA). *RIG-I*-expressing

chickens had a higher level of T cell activation in TCR1+/CD25+ T cells, which was not

the case for TCR2,3+/CD25+ T cells (Fig. 3c).

Heterozygous birds expressing each gene separately were crossed to obtain birds that

simultaneously express both genes *RIG-I* and *RNF135*. The obtained *RIG-I-RNF135*-

expressing chickens were monitored weekly for weight gain and closely investigated at

eight and twelve weeks of age for possible alternate immune phenotypes similar to those

observed in *RIG-I*-expressing birds. Unlike *RIG-I*-expressing chickens, no significant

differences in T cells or B cell counts were detected in *RIG-I-RNF135*-expressing chickens

in comparison to their WT siblings (Fig. 4d). Our analysis also comprised the investigated

cell populations in *RIG-I*-expressing birds including αβ, γδ and B cells (Fig. 4a).

Furthermore, we quantified CD4+, CD8+ T cells and corresponding subpopulations (Fig.
4b and Fig.4c). This was also the case for body weight gain, where no significant
differences were observed in *RIG-I-RNF135*-expressing chickens in comparison to WT
birds (Supplementary Figure 3).

These data indicate that the *RIG-I*, an innate immune sensor, can influence adaptive
immunity by causing shifts in T and B cell populations. In contrast, co-expression of
RNF135 with *RIG-I* seems to balance the adaptive immune cell populations, comparable
to WT birds.

RNF135 is required for efficient antiviral activity of *RIG-I*

The role of *RNF135* as an essential ubiquitination factor that supports the antiviral activity
of *RIG-I* was previously determined in mammalian cells (13). Later on, *RNF135* was found
to be the obligatory ubiquitin for *RIG-I* (23), while so far, this was not known in birds. We
investigated the effect of expressing *RNF135* on the antiviral activity of *RIG-I* in genetically
modified chickens. We examined the susceptibility of the generated transgenic lines
towards an H7N1 AIV, a direct precursor of a highly pathogenic avian influenza virus
(HPAIV), known to cause severe acute respiratory disease in chickens (24). Birds were
challenged with the virus and monitored for clinical signs, weight gain, viral replication,
and lesion development. *RIG-I*-expressing chickens had the highest rate of morbidity, as
indicated in Fig. 5a. The clinical symptoms started within the first two days of infection,
where the morbidity rate reached a significant rate of 44% at one-day post-infection (dpi)
($p<0.05$) and increased to 67% at two dpi ($p<0.05$), which then decreased to 18% by three
dpi ($p<0.05$). The onset of clinical disease was accompanied by a significant weight loss
at two dpi that remained significantly low at six dpi ($p<0.05$) (Fig. 5c) and coincided with a
pulmonary lesion score that increased from 0.8 at two dpi to 2.3 at six dpi (Fig. 6a and 6b).
This was also confirmed by histological assessment of the caecum that revealed necrotic
lesions associated with pronounced epithelial hyperplasia (Fig. 6b). *RIG-I-RNF135*-
expressing chickens exhibited increasing clinical illness, from 13% of sick animals at one
dpi to 50% at three dpi ($p<0.05$). Severe and prolonged clinical symptoms associated with
mortality in this group lasted until six dpi.

The expression of *RNF135* alone resulted in a mortality rate comparable to WT birds. In
contrast, the clinical symptoms in *RNF135*-expressing chickens lasted till three dpi with a

total of 25% sick animals (Fig. 5b). This was not the case for WT chickens that exhibited clinical symptoms only for the first two days after infection and had a morbidity rate of 13% and 19% at one and two dpi, respectively (Fig. 5b).

The quantification of the virus genome copies via qRT-PCR showed that the expression of *RIG-I* together with *RNF135* resulted in an efficient antiviral effect, as demonstrated by the significantly reduced amount of viral genome copies in the caecum compared to *RIG-I* expressing chickens at two and six dpi ($p<0.05$) (Fig. 5d). In the lung, challenged *RIG-I-RNF135*-expressing chickens displayed a virus replication rate similar to WT birds and a significantly lower replication rate than *RIG-I*-expressing chickens at two dpi ($p<0.05$). Conversely, *RIG-I*-expressing chickens had the highest viral replication rate in the lung and the caecum among all investigated groups, with a significant difference compared to *RIG-I-RNF135*-expressing chickens ($p<0.05$) (Fig. 5d). The amount of viral nucleic acid remained at a high level at six dpi, indicating the lack of viral clearance in these birds, which was not the case for the other challenged chicken lines (Fig. 5d).

The observed low viral titer in *RIG-I-RNF135*-expressing chickens, compared to *RIG-I*-expressing chickens, confirms the crucial role of *RNF135* as a ubiquitin ligase at the early stage of infection. Also, it demonstrates that the chicken *TRIM25* does not overtake the part of *RNF135* in *RIG-I*-expressing chickens.

The expression of *RIG-I* and *RNF135* in H7N1-challenged birds coincided with the acute phase of infection

We aimed to quantify possible changes in the expression levels of both transgenes in the challenged animals after the H7N1 infection. We quantified *RIG-I* and *RNF135* expressions via qRT-PCR in the lung, and the caecum (Fig. 5e). The caecal expression of *RIG-I* in virus-infected *RIG-I-RNF135* chickens was increased by ~26 folds. At the same time, infected *RIG-I* chickens exhibited an increase of ~24-fold compared to *RIG-I-RNF135* MOCK controls. Infected *RIG-* and *RIG-RNF135*-expressing chickens had a comparable *RIG-I* expression to *RIG-RNF135*-MOCK controls with no significant upregulation (Fig. 5e). Overall, we observed lower expression of *RNF135* in the caecum and lungs compared to *RIG-I-RNF135*-MOCK controls. The expression increased by 1.4 folds in *RNF135*-infected birds and ~2 folds in *RIG-I-RNF135*-infected chickens. In the lung, the *RNF135* expression at two dpi was upregulated by ~6 folds and ~4 folds in

infected *RNF135* and *RIG-I-RNF135*-expressing chickens, respectively. At six dpi, the *RNF135* expression was at a comparable level with the *RIG-I-RNF135* MOCK controls (Fig. 5e). These data indicate that the infection led to a quick upregulation of *RIG-I* at the acute phase of infection, which quickly decreased afterward.

Differential regulation of innate immune genes in naïve as well as in H7N1-challenged *RIG-I, RNF135,* and *RIG-I-RNF135*-expressing chickens

We speculated that the observed acute inflammatory reaction could be due to a differential regulation of inflammatory genes related to *RIG-I* signaling. Therefore, we sought to quantify the differentially expressed genes (DEGs) between transgenic and WT birds. We used the Fluidigm qRT-PCR array to identify changes in the expression of selected innate immune genes in both naïve and H7N1-challenged birds (25). The results revealed that in the absence of infection, *RIG-I-RNF135*-expressing chickens already had a significantly higher expression of ISGs in the caecum, including *ISG12-2* and *OASL,* as well as the transcription factor Early Growth Response *1 (EGR1)* (Fig. 7a, Supplementary Figure 4b). In contrast, others, including *EGR1*, interleukin 4 induced 1 (*IL4I1)*, Interleukin-1 beta (*Il1β),* and Interleukin 8 (*IL8),* were significantly upregulated in the spleen (Fig. 7a and Supplementary Figure 4).

The comparison of challenged groups with non-infected controls (WT-MOCK) indicated that *RIG-I* chickens expressed several inflammatory genes in the caecum and lungs that were not expressed in the other groups. This included *IFITM1*, *IL6,* and *LIPI* (Fig. 7b, Supplementary Figure 4c and 4d). At two dpi, we observed the upregulation of genes involved in the JAK-STAT signaling pathway, including *IL-6* and *STAT3,* which was not the case at 6 dpi (Fig. 7b, 7c and 7d). The significant upregulation of inflammatory and immune regulatory genes in the *RIG-I*-expressing chickens persisted until six dpi, where over 20 genes were exclusively upregulated in the spleen (Fig. 7e and Supplementary Figure 4e). This was not the case for *RIG-I-RNF135*-expressing chickens and WT chickens with an exclusive expression of three and seven genes, respectively (Fig. 7e and Supplementary Figure 4e). In addition, the analysis of expressed genes in the spleen revealed that only *RIG-I-RNF135* chickens expressed *IRF7* and *TRIM25*, in contrast to *RIG-I*-expressing chickens (Fig. 7e and Supplementary Figure 4e).

The functional enrichment analysis of the genes involved in the biological processes of transgenic chickens indicated that the regulated genes in *RIG-I*-expressing chickens were highly involved in metabolic activities. In contrast, those in *RIG-I-RNF135*-expressing birds were primarily involved in regulatory mechanisms (Supplementary Figure 5). Overall, the obtained data confirm that *RIG-I*-expressing chickens exhibited a significant increase of inflammatory cytokines that were not observed in other challenged birds, which may explain the acute inflammatory reaction in these birds.

High expression of *IFN-γ* is associated with significant virus replication in *RIG-I*-expressing birds

Since we observed a significant reduction of viral genome copies in the *RIG-I-RNF135*-expressing chickens compared to *RIG-I*-expressing birds, we compared the DEGs between H7N1-infected groups. This helped us determine possible factors behind the increased H7N1 replication in *RIG-I*-expressing birds. Interestingly, these birds lacked interferon upregulation compared to infected WT birds or *RIG-I-RNF135*-expressing chickens (Fig. 8a and 8d). We also found that the viral infection led to a significant increase in the expression of *IFN-γ* at six dpi when compared to H7N1-infected *RIG-I-RNF135*-expressing chickens (14-fold change increase) and WT birds (9-fold change increase) (Fig. 8b and 8d) (*p*<0.05). In contrast, *RIG-I-RNF135*-expressing birds quickly exhibited a significant increase of *IFN-α* expression, already at two dpi, in comparison to *RIG-I*-expressing birds (12-fold change) as well as to WT birds (17-fold change) (Fig. 8c). We concluded that the co-expression of *RNF135* and *RIG-I* in chickens leads to a significant increase of *IFN-α* expression, which was not observed when *RIG-I* was expressed solely. In addition, *RIG-I*-expression caused an upregulation of *IFN-γ* (Fig. 8b and 8d) that was accompanied by a significant increase in virus genome copies.

The deleterious inflammatory response in *RIG-I* and *RIG-I-RNF135*-expressing chickens depends on the virulence of the influenza subtype

The infection with H7N1 revealed a unique phenotype that manifested in acute inflammation and increased mortality in *RIG-I* and *RIG-I-RNF135*-expressing chickens. Hence, we were interested in investigating the effect of reinstating *RIG-I* and *RNF135* in the chicken genome on the outcome of infection with two additional virus strains of high or low virulence. We conducted two additional *in vivo* infection studies using two viruses:

an H3N1 (A/chicken/Belgium/460/201) and an H9N2 (A/chicken/Saudi Arabia/CP7/1998). While both subtypes are classified as low pathogenic AIVs due to their monobasic hemagglutinin cleavage sites, H3N1 has been described as a highly virulent strain causing severe clinical infection and mortality in adult layers (26). Displaying a distinct tropism for the hen's oviduct, H3N1 causes salpingitis and peritonitis, which are associated with a severe drop in egg production. In contrast, the H9N2 virus is effectively low virulent and does not cause any detectable symptoms in experimentally infected chickens(27). The H9N2- and H3N1- infection experiments of *RIG-I* and *RIG-I-RNF135*-expressing chickens revealed major differences in the outcome of infection between the two viruses. While H9N2 infection did not cause any clinical disease in the *RIG-I* or *RIG-I-RNF135*-expressing chickens (Supplementary Figure 12), infection with H3N1 led to marked clinical/pathological disease signs and disease aggravation, similar to those observed in the H7N1 infection experiment. The infection with H3N1 led to early clinical disease and mortality onset that were more pronounced in *RIG-I*-expressing birds (Fig 9a, 9b and Supplementary Figure 11). However, we did not detect differences in viral RNA loads in tracheal swabs between the H3N1-infected groups except for 7dpi, where *RIG-I*-expressing birds exhibited significantly higher loads compared to WT birds (Fig. 9a). *RIG-I*-expressing chickens also manifested a significantly increased expression of *IL-1β*, *IL6*, *IFN-α* and *IFN-γ* compared to the other infected birds (Fig. 9c). The assessment of histological lesions of the reproductive system indicated that the infection with H3N1 led to pronounced atrophy in *RIG-I* and *RIG-I-RNF135*-expressing birds in comparison to WT birds (Fig. 9d), accompanied by fibrinous peritonitis, salpingitis and vasculitis (Fig. 9e). These data confirm the observations made with H7N1 and indicate that the deleterious clinical disease caused by *RIG-I* reinstatement depends on the degree of viral virulence.

Discussion

The constant arms race between pathogens and the host affects different aspects of the immune system, including innate sensors. The reason for the loss of *RIG-I* in Galliformes, which correlated with the disappearance of the ubiquitin ligase *RNF135*, remained a mystery (20), especially given the established contributing role of *RIG-I* in the resilience towards influenza in ducks and other species (7, 28). It was previously speculated that the gradual loss of *RIG-I* and *RNF135* in the chicken was possibly caused by the pathogen's resistance to the innate sensor or the disappearance of some relevant pathogens (20). At

the same time, preservation of the antiviral competence accompanied the loss of *RIG-I* and *RNF135* in chickens via the evolutionary selection of *MDA5* and *LGP2* (29). The beneficial effect of the duck *RIG-I*-overexpression in chicken DF-1 cells was demonstrated by the limited replication of AIV in these transgene-expressing cells (7). So far, no chickens expressing the duck *RIG-I* have been generated to investigate their antiviral response *in vivo*. In this study, we produced genetically modified chickens that express *RIG-I* with or without its ubiquitin ligase *RNF135* to examine the physiological role of both genes and their function during AIV infection. Both genes were cloned from the duck representing the most studied avian influenza reservoir with evolutionary conserved *RIG-I* and *RNF135* (6, 20, 30). The expression of both genes was controlled under their respective duck promoters since the duck *RIG-I* was previously tested in chicken cells(22). This strategy allows the strict control of gene expression that can be induced only upon infection, which prevents undesirable production of inflammatory cytokines that may cause autoimmune diseases (22). While the duck *RIG-I* promoter was previously identified (22), we described the activity of the duck *RNF135* promoter using chicken cells.

Without infection, *RIG-I* expression did not cause any harmful phenotype, but led to different adaptive immune cell counts compared to WT siblings. This suggests the role of the *RIG-I* in priming T cell immunity in the chicken, which can be similar to mice that exhibited a lack of T cell immunity associated with deficiencies in migratory dendritic cell activation, viral antigen presentation and CD8+ and CD4+ T cell priming (17) (14). In the case of *RIG-I*-*RNF135*-co-expression, the balanced adaptive immune phenotype implies a possible role of *RNF135* in modulating the T cell immune response in birds, which was previously described for Th1 response and cytotoxic T cells in mice (31). Moreover, naïve *RIG-I*-*RNF135*-expressing chickens had a significantly higher expression of interleukin 4 inducible gene 1 (*IL4I1*) as well as protein nuclear translocation 7 *(IRF7)* and tripartite motif-containing protein 29 *(TRIM29)*. This may explain the differential adaptive immune phenotype between *RIG-I* and *RIG-I*-*RNF135*-expressing chickens. Previous studies indicated the involvement of *IL4I1* in the signaling to T and B lymphocytes (32) and the effective role of translocation factors such as *IRF7* in chicken cells transfected with the duck *RIG-I* (22).

The lack of an antiviral effect after *in ovo* and *in vitro* experimental infections with LPAIV H9N2 was discordant with the previously described antiviral effect of duck *RIG-I* in chicken

DF-1 cells (7). Barber et al, observed low virus replication upon the overexpression of
RIG-I under the control of the CMV promoter. The difference between our results and
previously published data can be related to the chosen promoter used by Barber et al,
which may lead to a higher upregulation of *RIG-I* and, consequently, a strong production
of interferon-stimulated genes. Additional factors can stand behind the differences
between the previously published results by Barber et al(7), and our data, including
influenza subtypes and the type of cells.

H7N1-challenge experiments of the generated transgenic birds revealed that *RIG-I*-
expressing chickens had an early upregulation of proinflammatory cytokines such as *IL-*
6, known to support *IL1β* in suppressing regulatory T cells, which can lead to an
uncontrolled increase in the number of CD4+ T cells (33). Furthermore, the expression of
RIG-I without *RNF135* was not beneficial in limiting H7N1 replication since the *RIG-I*-
expressing chickens had the highest viral genome copies among all challenged groups.
The requirement of *RNF135* for an *RIG-I*-mediated antiviral effect in *RIG-I-RNF135*-
expressing chickens was also described in mammalian cells (13). In addition, we found
that the H7N1 infection caused a significant upregulation of *RIG-I* in the caecum compared
to the lungs. This can be because *RIG-I* expression rapidly occurs upon stimulation and
may reach a peak by three hours post-infection (34). Similar observations were described
by Cornelissen et al. that detected a significant upregulation of *RIG-I* expression in the
lungs of ducks at 8h post H7N1 infection that significantly decreased at two dpi (35). *RIG-*
I-expressing birds infected with H7N1 exhibited an acute inflammatory response and
weight loss at two dpi, lasting till six dpi. Pang et al, reported that influenza virus could
hijack the inflammatory reaction associated with RIG-I signaling for its replicative
advantage, particularly in the respiratory tract (15). H7N1-infected *RIG-I*-expressing birds
showed several regulated immune genes compared to other infected groups, including
upregulation of the suppressor of cytokine signaling 1 (*SOCS1*) in the caecum. This may
explain the significant increase in viral genome copies and the observed T cell phenotype
since *SOCS1* is a potent inhibitor of JAK/STAT signaling (36) and is involved in several
mechanisms that regulate the T cell maturation, differentiation, and function (37).
Furthermore, the assessment of the immunophenotype of *RNF135*-expressing chickens
revealed differential upregulation of various genes, which suggests that this gene, on its
own, may function independently from *RIG-I* (38).

Interestingly, we detected significant upregulation of *IFN-γ* in the lungs of infected *RIG-I-*
expressing chickens compared to infected *RIG-I-RNF135-* and WT-chickens, which can
be responsible for lung-mediated injury and acute death caused by respiratory distress.
Similarly, H3N1 infection caused a relative upregulation of *IFN-γ* in the spleen and the
duodenum of *RIG-I*-expressing chickens. The role of *IFN-γ* was previously described in
IFN-γ KO mice that were less susceptible to lung inflammation and pathology upon
influenza infection (39). The lack of an antiviral effect in *RIG-I*-expressing chickens may
also indicate that chicken *TRIM25* is dispensable for *RIG-I* efficient ubiquitination. Similar
findings were described using human lung adenocarcinoma epithelial cells, where *TRIM25*
did not participate in the endogenous *RIG-I*-dependent antiviral responses (21). However,
other studies done in mammalian models indicated that the deletion of *TRIM25* increases
the susceptibility to influenza infection (40), supporting the evidence that *TRIM25* may
bind directly to the viral RNA, thereby contributing to the restriction of influenza virus
infection (41). The exacerbated inflammatory reaction in *RIG-I*-expressing chickens could
be related to the absence of *RNF135*, which suggests a stabilizing role of *RNF135*
comparable to *TRIM25* (42), though this requires further investigation. The increased
mortality rates in *RIG-I-* and *RIG-I-RNF135*-expressing chickens may be explained by
immunopathology in both chicken lines, despite the differences in viral replication. The
additional *in vivo* experimental challenge with the highly virulent H3N1 virus confirmed the
deleterious immunophenotype observed for the H7N1 virus, as demonstrated by the
upregulation of various inflammatory genes, including *IFN-γ, IFN-α, IL1β, and IL6*. The
absence of a similar phenotype after infection with the mildly virulent H9N2 virus revealed
that disease exacerbation in *RIG-I-* and *RIG-I-RNF135*-expressing chickens requires a
certain degree of viral virulence.

Our data suggest that the evolutionary loss of *RIG-I* in chickens and other galliform birds
was advantageous in coping with viral infections caused by AIV or other avian RNA
viruses. This subsequently helped decrease the acute inflammation and the damage to
the host. A comparable hypothesis was made in the case of pangolins that lost the *MDA5*
as an evolutionary mechanism to cope with coronavirus-induced inflammation (43). Above
that, we propose that the antiviral role of *RIG-I* in ducks(7) is not exclusively related to this
gene. Still, it may involve the ubiquitination factor *RNF135* and possibly other unknown
regulatory factors that support *RIG-I* signaling and reduce the repercussions of acute

inflammation by negatively regulating the *RIG-I* signaling (44). We provided novel
information regarding the outcome of the re-introduction of *RIG-I* and its ubiquitination
factor *RNF135* in the chicken genome. Future work should focus on identifying factors that
can help reduce the acute inflammatory reaction in *RIG-I-RNF135*-expressing chickens
while maintaining potent antiviral activity, which can lead to the generation of avian
influenza virus-resistant chickens.

Materials and Methods

455
Cloning of the duck *RIG-I* and the duck *RIG-I* promoter

Total RNA was isolated from the spleen of the mallard duck (*Anas platyrhynchos*) using
ReliaPrep™ RNA Tissue Miniprep System (Promega, USA), followed by cDNA synthesis
using GoScript™ Reverse Transcription System (Promega, USA) according to the
manufacturer's instructions. The duck *RIG-I* was amplified using Q5® High-Fidelity DNA
Polymerase (New England Biolabs, USA) using the primers 562_RIG-I_for (5'-
ATGACGGCGGACGAGAAGCGGAGC-3') and 563_RIG-I_rev (5'-
CTAAAATGGTGGGTACAAGTTGGAC-3') that were previously described (45). The PCR
thermal conditions were as follows: 98 °C 30 sec, followed by 35 cycles of 98 °C 10 sec,
67 °C 20 sec, 72 °C 1:30 min, and a final extension step of 72 °C 2 min.

The entire length of the duck *RIG-I* promoter was amplified using primers 706_RIG-I_For
(5'-AGCTGATGACCTGCAAAAAGTT-3') and 661_RIG-I_Rev (5'-
GGCTGGGCTCTGCCGGCCG-3'), which were described elsewhere (22). This resulted
in an amplicon of 2017 bp that was fully sequenced and aligned with the duck genome
(*Anas platyrhynchos*, NC_040075.1). The PCR was conducted following the following
thermal conditions   98 °C 30 sec, followed by 35 cycles of 98 °C 10 sec, 70 °C 20 sec,
72 °C 1:30 min, and a final extension step of 72 °C 2 min.

Cloning of the duck *RNF135*

The genomic region containing duck *RNF135* was obtained from the GenBank contig
PEDO01000017.1 and corrected based on multiple publicly available duck RNAseq data.
Although the *RNF135* sequence was correctly predicted in many birds, we detected a
partially incomplete annotated sequence of the duck *RNF135*, specifically the missing 5'
part that contains the RING domain (previous accession number XM_013092775), while

the predicted version of the *RNF135* was made available (XM_027471415.2). The full-

length sequence of the duck *RNF135* was synthesized after codon optimization using the

IDT online tool (IDT™, USA) (Supplementary Figure 6). The obtained *RNF135* sequence

was subsequently cloned into the *RNF135*-expression vector driven by the duck *RNF135*

promoter (Fig. 1b).

Identification of the duck *RNF135* promoter via Nano-Glo® Dual-Luciferase®
484

Reporter Assay

The putative duck *RNF135* promoter was obtained by amplifying 1577 bp 5' of the ATG

start codon from duck gDNA cloned into pGEM vector (Promega, USA) and analyzed by

Sanger sequencing. The PCR was done using Q5® High-Fidelity DNA Polymerase with

primers: 707_RNF_Prom_For (5'-GA GCA GAG CCA GGC AGC TAT A-3'), 708 (5'-GGT

CCT GCT CGG GGC GGA GC-3') resulting in an amplicon of 1557 bp. The PCR thermal

conditions were conducted using Q5® High-Fidelity DNA Polymerase (New England

Biolabs, USA) at two step-PCR 98 °C 30 sec, 98 °C 10 sec, 72 °C 60 sec and a final

elongation step of 72 °C 2 min.

The promoter activity of duck *RNF135* was assessed by measuring promoter-driven

NanoLuc™ luciferase activity normalized to the luminescence of Firefly luciferase. To this

end, a total of 50.000 chicken DF-1 cells were seeded in 24 well plates and were co-

transfected 24h later with a vector plasmid containing the deletion mutant and a second

plasmid for the expression of Firefly under the PGK promoter (Promega, USA). 24h after

transfection, cells were washed with PBS, trypsinized, and resuspended in 250µl culture

medium. Firefly signal was detected by mixing 80µl of the cell suspension with the same

amount of ONE-Glo™ EX Reagent, prepared by combining ONE-Glo™ EX Luciferase

Assay Buffer with ONE-Glo™ EX Luciferase Assay Substrate in 1:1 ratio (Promega, USA).

After measuring the signal of the Firefly luciferase in the GloMax® 20/20 Luminometer

(Promega, USA), 80 µl NanoDLR™ Stop & Glo® reagent, prepared by adding

NanoDLR™ Stop & Glo® Substrate 1:100 into NanoDLR™ Stop & Glo® Buffer (Promega,

USA), were added to the samples. These were incubated for 10, 30, 60, and 120 min and

measured again in the GloMax® 20/20 Luminometer (Promega, USA) to detect the

NanoLuc™ Luciferase. The cell-free culture medium was used as a blank control.

Determination of the Transgene Copy Number by Droplet Digital PCR

Droplet digital PCR (ddPCR) was used to select PGC clones with a single genomic transgene integration and was performed as described previously with slight modifications (46). Briefly, 500ng of gDNA was digested with 20 units XbaI (New England Biolabs, Germany) for one h, followed by an inactivation step at 65°C for 20 min. The Taqman PCR reaction was set up using 10ng digested DNA, 2× ddPCR supermix for probes (no dUTP) final concentration 1× (Bio-Rad Laboratories, USA), a 20× target primer/FAM-labeled probe mix, and a 20× reference primer/HEX-labeled probe mix, which was followed by droplet generation using the QX200 Droplet Generator by mixing 20ul of the TaqMan PCR reaction with 70ul droplet generator oil in a DG8 Cartridge. The cycling conditions comprised a 95°C for 10min, followed by 40 cycles of 94°C for 30 sec, 59°C for 60 sec, and a final hold for 98°C for 10 min with a 2°C/s ramp rate at all steps. The copy number was determined by calculating the proportion of positive and negative droplets using a QX200 droplet reader, which was then analyzed using Quantasoft software (Bio-Rad Laboratories, USA) (Supplementary Figure 7). The hygromycin fluorescent labeled-probe ([5′FAM-TCGTGCACGCGGATTTCGGCTCCAA-3′] along with the primers: ddHygro_F [5′-CATATGGCGCGATTGCTGATC-3′] and ddHygro_R [5′-GTCAATGACCGCTGTTATGC-3′]). As a reference gene, we used the beta-actin probe ([5′HEX-GTGGGTGGAGGAGGCTGAGC-BHQ3′] along with the primer combination ddBeta_actin_F: [5′-CAGGATGCAGAAGGAGATCA-3′] and ddBeta_actin_R: [5′-TCC ACCACTAAGACAAAGCA- 3′]). The quantification was done using the QX100 system (Bio-Rad Laboratories, USA) (Supplementary Figure 7).

Stimulation of duck *RIG-I* expression in PGC-derived fibroblasts

The generated *RIG-I*-expressing PGCs were differentiated into fibroblasts (PGCFs) as previously described (47) and subsequently infected with a LPAIV H9N2 to examine the ability of *RIG-I* to detect its ligand. Briefly, 50.000 cells were seeded into 48 well plates and infected with an MOI of 0.1 for 18h before they were fixed and stained using immunofluorescence, as previously described (48). Cells were fixed with 4% PFA and kept on ice for 10 min. Subsequently, they were washed with PBS and permeabilized with 0.5% Triton X. The FLAG-Tag was detected by using a mouse anti-FLAG antibody (Supplementary Table 1) that was incubated for 1h, followed by staining with goat anti-mouse Alexa 568 (Supplementary Table 1). Slides were subsequently covered with a mounting medium that contains DAPI for staining the nucleus and covered with coverslips

542 (Vector Laboratories, Inc., USA). Fluorescence microscopy was performed with a
543 fluorescence microscope (ApoTome, Zeiss).

Generation of *RIG-I-*and *RNF135*-expressing chickens

White Leghorn layer chickens (Lohmann selected White Leghorn (LSL), Lohmann-
Tierzucht GmbH, Cuxhaven, Germany) were used to generate transgenic chicken lines.
Animal experiments were approved by the government of Upper Bavaria, Germany (ROB-
55.2-2532.Vet_02-18-9). Experiments were performed according to the German Welfare
Act and the European Union Normative for Care and Use of Experimental Animals. All
animals received a commercial standard diet and water *ad libitum*. PGCs that express
either *RIG-I* or *RNF135* were generated using the DNA constructs shown in Fig. 1b, as
previously described (49, 50). To ensure the stable integration of the transgene, we used
the phiC31 integrase-mediated integration (51). Briefly, LSL PGCs were derived from the
blood of male embryonic vasculature at stages 13-15, according to Hamburger and
Hamilton (52). They were cultured at 37°C in a 5% CO2 environment using modified KO-
DMEM as described previously (53). A total of $5x10^6$ cells per transgene were washed
with phosphate-buffered saline (PBS) and resuspended in 100µl Nucleofector™ Solution
V (Lonza, Germany) containing the expression vector (Fig. 1b) and the integrase
expression construct. Electroporation was performed using an ECM 830 Square Wave
Electroporation System (BTX, USA), applying eight square wave pulses (350V, 100µsec).
After clonal selection using puromycin for *RIG-I* and blasticidin for *RNF135*, PGCs were
genotyped and tested for clones with one single genomic integration, which were then
used to generate the chimeric roosters. A total of 3000 cells with the desired genetic
modification were injected into the vasculature of 65h old embryos, transferred into a
surrogate eggshell, and incubated until the hatch. Upon sexual maturity, sperm was
collected from chimeric roosters for genotyping. (50, 54). The germline-positive roosters
were bred with wild-type hens to obtain heterozygous animals (the germline transmission
rate is presented in Supplementary Table 2).

The examination of the inheritance of duck *RIG-I* and the genotyping of the offspring was
done via PCR using FIREPol DNA Polymerase (Solis Biodyne) using the primer
combination 613_RIG-I-for (5'-CCTAGGAGAAGCATTCAAGGAG-3') and 563_RIG-
I_Rev (5'-CTAAAATGGTGGGTACAAGTTGGAC-3'). The following PCR conditions were

used: initial denaturation at 95 °C 3 min, followed by 40 cycles of 95 °C 30 sec, 60 °C 30

sec, 72 °C 20 sec, and a final extension step of 72 °C 5 min, resulting in a fragment of

298bp. The inheritance of the duck *RNF135* was examined using the primers combination

1121_RNF_For (5'GCATGGGATCAACCGACAGCATC-3') and 1017_RNF_rev

(5'CCACACACCAACTTGACTCGGTC-3', using the following PCR conditions 95 °C 3

min, followed by 40 cycles of 95 °C 30 sec, 60 °C 30 sec, 72 °C for 1 min and a final

extension step of 72 °C 5 min, resulting in an amplicon of 931bp.

Following the generation of different transgenic lines, they were monitored till sexual

maturity for possible harmful phenotypes that could been reflected in weight gain or the

ability to produce sperm or eggs. In addition, the immunophenotype of the generated birds

was assessed at 2, 5, 8, and 12 weeks after hatch by flow cytometry.

Isolation, culture, and infection of chicken embryonic fibroblasts (CEFs)

CEFs were isolated from 10-day-old (ED10) embryos according to the protocol published

elsewhere (55). Before the isolation of CEFs, embryos were genotyped by collecting blood

at ED10 and preparing a window of 0.5 cm^2 in the eggshell that allowed access to the

embryonic vasculature. CEFs were cultured using Iscove's liquid medium containing

stable glutamine (Biochrom, Germany) that was supplemented with 5% fetal bovine serum

(FBS) Superior (Biochrom, Germany), 2% chicken serum (ThermoFisher Scientific, USA)

and 1% Penicillin-Streptomycin-Solution (Penicillin 10,000 U/ml and Streptomycin 10

mg/ml) (Biochrom. Germany). Subsequently, CEFs were incubated at 40°C in a 5% CO2

atmosphere until infection. The infection of CEFs with an H9N2 virus (A/chicken/Saudi

Arabia/CP7/1998) was done after overnight seeding 250.000 cells/well in 6 well plates and

infecting them in three independent experiments with multiplicities of infection (MOIs) of

0.1 and 1. Supernatants were collected 24h post-infection from the infected cells and

titrated on MDCK (kindly provided by Prof. Silke Rautenschlein, University of Veterinary

Medicine, Hannover). The virus titration was done as previously described (48). An

additional *in vitro* infection experiment with CEFs was conducted using the H1N1-WSN

strain (A/WSN/1933), kindly provided by Prof. Bernd Kaspers, Ludwig Maximilian

University of Munich. Briefly, cells of the four genotypes were seeded overnight and

infected with two different MOIs, 0.001 and 0.01. 40h later, they were fixed and stained

for plaque formation following the standard titration protocol(48).

In ovo infection of embryonated eggs

Fertilized eggs from a crossing of *RIG-I* (+/-) and *RNF135*(+/-) were collected and incubated till ED10 or ED14. In a blind study, eggs were infected randomly with 10^3 FFU of the H9N2 virus, as previously described (56). 24h post-infection, the allantois fluid and muscle tissue were collected respectively for viral titration and genotyping. The virus titration was done on MDCK cells as previously described (48).

RT-PCR for the detection of transgene expression in different tissues

RNA from chicken organs, including lung, trachea, heart, liver, duodenum, thymus, bursa, and brain, was isolated with Reliaprep RNA Tissue Miniprep System according to manufacturer instructions (Promega), followed by cDNA synthesis using GoScript Reverse transcription mix (Promega). The detection of *RIG-I*, as well as *RNF135* from various tissues, was done using the following primers. *RIG-I*: 613_RIG-I-for (5'-CCTAGGAGAAGCATTCAAGGAG-3') and 563_RIG-I_Rev 5'-(CTAAAATGGTGGGTACAAGTTGGAC-3'), while *RNF135* was detected using the following primers: 898_duRNF_for (5'-CTTGAGAGAGGTGGAGGGAGC-3') and 899_duRNF_rev (5'-GGGCTGGTGGGAATTGTTGAGG-3'). *RIG-I* and *RNF135* PCRs produced an amplicon of 298bp and 148bp, respectively. β-actin mRNA was detected with primers Beta_actin_F (5′-TACCACAATGTACCCTGGC-3′) and Beta_actin_R (5′-CTCGTCTTGTTTTATGCGC-3′) (57), resulting in a 300-bp amplicon. The PCR was performed using FIREPol DNA Polymerase (Solis Bio-dyne) according to the manufacturer's instructions. The following PCR conditions were used: initial denaturation at 98 °C 30 sec, followed by 40 cycles of 98 °C 10 sec, 59 °C 30 sec, 72 °C 30 sec, and a final extension step of 72 °C 2 min.

Metanalysis of *RIG-I* and *RNF135* expression in duck tissues

A metanalysis of publicly available RNA-seq data was performed to estimate the expression of *RIG-I* and *RNF135* in duck tissues (reads per kilobase per million mapped reads; RPKM), as previously described (58).

Enzyme-linked immunosorbent assay (ELISA)

ELISA was done to measure the total plasma IgM and IgY concentrations. Briefly, 96 well plates were coated overnight with anti-chicken antibodies IgM and IgY at a concentration of 2µg/mL (Supplementary Table 1). The next day, plates were washed three times with washing buffer, followed by a blocking step with 4% skim milk for one hour. The prediluted plasma samples in 1:3 serial dilution were pipetted in the plates and incubated for one hour. The detection was done using secondary HRP-conjugated antibodies at the concentrations mentioned in Tabe1, which were incubated for 1h at RT. This was followed by adding 100µl/well of TMB (3,3', 5,5;-tetramethylbenzidine) substrate solution for 10 min, which was stopped by 50µl per well of 1M sulfuric acid solution. The optical density (OD) was measured using FluoStar Omega via the measuring filter 450nm and the reference filter 620 nm (Version 5.70 R2 BMG LABTECH, Ortenberg, Germany).

Flow Cytometry

Peripheral blood mononuclear cells (PBMCs) were isolated using Histopaque®-1077 density gradient centrifugation (Sigma, Taufkirchen, Germany) and analyzed using flow cytometry. Extracellular staining was carried out to detect various chicken immune cell markers including T cell subpopulations, B cells and Monocytes (list of antibodies is mentioned in Supplementary Table 1). Briefly, $5x10^6$ cells were washed with 2% BSA diluted in PBS (FLUO-Buffer). To determine the living cell population, cells were incubated with Fixable Viability Dye eFluor 780 (eBioscience, Thermo Fisher Scientific, USA). After washing with FLUO-Buffer, primary antibodies (concentration shown in Supplementary Table 1) were applied for 20 min. Cells were washed in FLUO-Buffer to remove unbound antibodies and incubated with conjugated secondary antibodies for 20 min. Subsequently, cells were again washed and analyzed using an Attune flow cytometer (Thermo Fisher Scientific, USA). The obtained data were then analyzed with FlowJo 10.8.1 software (FlowJo, Ashland, USA). An example of the gating strategy used in this study is presented in Supplementary Figure 8.

Assessment of _in vitro_ T cell activation

PBMCs were isolated from 12 weeks old _RIG-I_ or WT birds and resuspended in RPMI medium supplemented with 10%FBS and 1% Penicillin/Streptomycin and distributed on a 48 well plate with a total of $5x10^6$ cells/well. Cells were subsequently stimulated with Concanavalin A (Con A) (eBioscience™) at a concentration of 25 µg per well. They were

663　monitored every 24h for T cell activation via flow cytometry. Activated T cells were stained

664　for surface expression of γδTCR1, CD25, or αβTCR2&3 using antibodies listed in

665　Supplementary Table 1.

Challenge infection experiment with H7N1

The challenge infection experiment was performed at the INRAE-PFIE platform (INRAE

Centre Val de Loire, Nouzilly, France) and approved by the local Ethics Committee Val de

Loire and the Ministère de l' Enseignement Supérieur et de la Recherche under the

number 2021120115599580. Transgenic birds required for the experiment were hatched

in dedicated hatching isolators and genotyped by PCR using DNA extracted from EDTA

blood samples collected at day 7 post hatch. At 3 weeks of age, transgenic and WT-

chickens were distributed into four different groups (Supplementary Table 3)

corresponding to four BSL-3 isolator units. Two other groups were kept together as MOCK

controls in a single isolator (Supplementary Table 3). The birds were fed a commercial

standard diet and provided water ad libitum throughout the experiment. Prior to infection

(0 dpi), body weights were recorded, and blood samples were taken by occipital sinus

puncture. The virus used for challenge infection was A/ Turkey/Italy/977/1999 (kindly

provided by Dr. Ilaria Capua, Istituto Zooprofilattico Sperimentale Delle Venezie, Legnaro,

Italy), an H7N1 subtype virus that, despite its classification as an LPAIV, causes up to

50% mortality in experimentally infected White Leghorn chickens (24, 59, 60).

Chickens were PBS/mock-treated (groups 5 and 6) or virus-infected (groups 1-4) by intra-

tracheal and intra-choanal cleft inoculation of 0.1mL PBS or 5×10^5 EID50/1×10^6 EID50

H7N1. Animal behavior and clinical disease signs were monitored twice daily during the

trial. Clinical signs were evaluated according to the following score: 0 (no clinical signs), 1

(mild clinical signs), 2 (severe clinical signs), or 3 (dead/euthanized). Birds were

euthanized by pentobarbital injection into the occipital sinus at the end of the experiment

or once humane endpoints were reached. Samples collected from euthanized birds

included lung, caeca, and spleen. In addition, a scoring system was used to evaluate

macroscopic lung lesions as follows: 1 (mild, localized edema and fibrinous exudate), 2

(moderate edema and with hemorrhage and fibrinous exudate over ~1/4 of the lung), or 3

(severe hemorrhage and extensive edema over ~1/2 of the lung) (Supplementary Figure

9). Data collected from the animal experiment were assessed at 2 and 6 dpi, representing

the time points when tissue samples for the Fluidigm assay were collected. Details regarding the number of birds, group distribution and number of analyzed samples are given in Supplementary Tables 3 and 4.

Quantification of the viral genome in H7N1-challenged chickens

Total RNA was isolated from the isolated organs and conserved in 1ml NucleoProtect RNA (Macherey-Nagel, Germany). Samples were later processed for total RNA isolation with NucleoSpin RNA (Macherey-Nagel, Germany) according to the manufacturer's instructions. The viral genome of the H7N1 virus was quantified using qRT-PCR that was conducted with the Bio-Rad iTaq™ Universal SYBR ® Green One-Step Kit (BioRad, California, USA) using primers designed for the detection of the M gene as published elsewhere (61).

Analysis of gene expression via Fluidigm Dynamic Array

Gene expression was analyzed as previously described (25). Briefly, total RNA was extracted from infected animals' lungs, caeca, and spleen and processed for quality control via nanodrop. Reverse transcription was performed using the High Capacity Reverse Transcription Kit (Applied Biosystems) according to manufacturer's instructions with random hexamers and oligo (dT)18 in a final volume of 10μl, containing 250 ng total RNA. Subsequently, the cDNA was pre-amplified using TaqMan PreAmp Master Mix (Applied Biosystems). Quantitative PCR was performed in the BioMark HD instrument with the 96.96 IFC Dynamic Array (Fluidigm). The reaction was prepared by mixing 2.5 μl TaqMan Gene Expression Master Mix (Applied Biosystems), 0.25 μl 20X DNA Binding Dye Sample Loading Reagent (Fluidigm), 0.25 μl 20X EvaGreen DNA binding dye (Biotum) and 2 μl of preamplified cDNA. The qPCR was run under the following thermal conditions: 50°C for 2 min, 70°C for 30 min, 25°C for 10 min, followed by hot start 50°C for 2 min, 95°C for 10 min, PCR (x30 cycles) 95°C for 15 sec, 60°C for 60 sec and melting curve analysis 60°C for 3 sec to 95°C.

Raw quantitation cycle (Cq) data were collated with the Real-time PCR Analysis software v3.1.3 (Fluidigm), setting parameters of quality Cq threshold to auto (global) and the baseline correction to derivative. Raw Cq values were processed with GenEx.v6 MultiD, with correction for primer efficiency and reference gene normalization. Stability of the

expression of reference genes: *TATA box binding protein (TBP), Tubulin alpha chain (TUBA8B), beta-actin (ACTB), beta-glucuronidase (GUSB), glyceraldehyde-3-phosphate dehydrogenase (GAPDH) and ribosomal 28S (r28S)* were evaluated via NormFinder (GenEx). The geometric mean of the most stable (*GAPDH*, *GUSB* and *TBP*) were used for normalization. Technical replicates were averaged, and relative quantification was to the maximum Cq value obtained per gene, transformed to logarithmic scale, which was then statistically analyzed using T-test.

Challenge infection experiments with H3N1 and H9N2

Challenge experiments were performed at the Animal Research Center (ARC) of the Technical University of Munich and approved by the government of Upper Bavaria under the number ROB- 55.2-2532.Vet_02-21-11. The H3N1 virus (A/Chicken/Belgium/460/2019) was kindly provided by Dr. Joris Pieter De Gussem (Poulpharm BV). Infection with H3N1 was done with 28-week-old chickens. Infection with H9N2 (A/chicken/Saudi Arabia/CP7/1998) was done with four weeks old chicks. According to the obtained genotype at hatch (WT, *RIG-I-* and *RIG-I-RNF135-*expressing chickens), birds were distributed to groups of four or six birds per group. An infectious dose of 10^6 FFU in 0.2mL PBS per bird was distributed via nasal and tracheal routes for both viruses. In both experiments, birds were monitored daily for clinical symptoms, and tracheal swabs were collected on days 0, 3, 7, 11 and 17 to analyze the viral RNA loads by RT-qPCR(48). Duodenum, lung, and spleen samples were collected when mortality occurred or on the final day of the experiment (17 dpi). The expression of *RIG-I, RNF135, IFN-γ, IFN-α, IL1β, and IL6* was assessed by RT-qPCR (62).

Histology

Caeca and lungs (H7N1 infection experiment) or oviduct samples (infundibulum, magnum − H3N1 infection experiment) were fixed in 10% neutral buffered formalin and processed routinely. Four-micrometer tissue sections were stained with hematoxylin and eosin for light microscopy. The sections' analysis addressed histopathological changes and signs of inflammatory reactions, including degeneration, necrosis, infiltration of inflammatory cells, fibrin exudation, and epithelial hyperplasia.

Statistical analysis

Statistical analysis was done using IBM SPSS Statistics (Version 28.0.1.1. IBM, Armonk, USA). The normality of the data was examined via Kolmogorov-Smirnov and Shapiro-Wilk tests. The comparison between two groups was made either with Two Samples T-test or Wilcoxon Signed Rank test. Multiple group comparison was made using Kruskal-Wallis Test or with One-Way ANOVA. The statistical test result was considered significant when the P value was less than 0.05.

Acknowledgments

Animal husbandry and experimental infrastructure was provided by the TUM Animal Research Center (ARC). The authors thank the animal keepers of the ARC as well as the team at the INRAE Plateforme d'Infectiologie Expérimentale (INRAE-PFIE, Nouzilly, France) for conducting the H7N1 challenge infection experiment. Additionally, the authors thank the technical staff of the involved laboratories for excellent assistance.

References

1. A. Mostafa, E. M. Abdelwhab, T. C. Mettenleiter, S. Pleschka, Zoonotic potential of influenza A viruses: a comprehensive overview. *Viruses* **10**, 497 (2018).
2. E. F. S. Authority *et al.*, Avian influenza overview December 2022-March 2023. *EFSA journal. European Food Safety Authority* **21**, e07917 (2023).
3. G. Granata, L. Simonsen, N. Petrosillo, E. Petersen, Mortality of H5N1 human infections might be due to H5N1 virus pneumonia and could decrease by switching receptor. *The Lancet Infectious Diseases* **24**, e544-e545 (2024).
4. M. Kang *et al.*, Changing epidemiological patterns in human avian influenza virus infections. *The Lancet Microbe* **5** (2024).
5. N. MacLachlan, Orthomyxoviridae. *Fenner's veterinary virology*, 353-370 (2011).
6. J. K. Kim, N. J. Negovetich, H. L. Forrest, R. G. Webster, Ducks: the "Trojan horses" of H5N1 influenza. *Influenza and other respiratory viruses* **3**, 121-128 (2009).
7. M. R. Barber, J. R. Aldridge Jr, R. G. Webster, K. E. Magor, Association of RIG-I with innate immunity of ducks to influenza. *Proceedings of the National Academy of Sciences* **107**, 5913-5918 (2010).
8. J. Rehwinkel *et al.*, RIG-I detects viral genomic RNA during negative-strand RNA virus infection. *Cell* **140**, 397-408 (2010).
9. E. Kowalinski *et al.*, Structural basis for the activation of innate immune pattern-recognition receptor RIG-I by viral RNA. *Cell* **147**, 423-435 (2011).
10. M. Yoneyama *et al.*, The RNA helicase RIG-I has an essential function in double-stranded RNA-induced innate antiviral responses. *Nature immunology* **5**, 730-737 (2004).

11. M. U. Gack *et al.*, TRIM25 RING-finger E3 ubiquitin ligase is essential for RIG-I-mediated
antiviral activity. *Nature* **446**, 916-920 (2007).
12. D. Gao *et al.*, REUL is a novel E3 ubiquitin ligase and stimulator of retinoic-acid-inducible
gene-I. *PloS one* **4**, e5760 (2009).
13. H. Oshiumi *et al.*, The ubiquitin ligase Riplet is essential for RIG-I-dependent innate
immune responses to RNA virus infection. *Cell host & microbe* **8**, 496-509 (2010).
14. Y. Wang *et al.*, Rig-I−/− mice develop colitis associated with downregulation of Gαi2. *Cell
research* **17**, 858-868 (2007).
15. I. K. Pang, P. S. Pillai, A. Iwasaki, Efficient influenza A virus replication in the respiratory
tract requires signals from TLR7 and RIG-I. *Proceedings of the National Academy of
Sciences* **110**, 13910-13915 (2013).
16. R. Cagliani *et al.*, RIG-I-like receptors evolved adaptively in mammals, with parallel
evolution at LGP2 and RIG-I. *Journal of molecular biology* **426**, 1351-1365 (2014).
17. M. Kandasamy *et al.*, RIG-I signaling is critical for efficient polyfunctional T cell responses
during influenza virus infection. *PLoS pathogens* **12**, e1005754 (2016).
18. H. A. Vanderven *et al.*, Avian influenza rapidly induces antiviral genes in duck lung and
intestine. *Molecular immunology* **51**, 316-324 (2012).
19. W. Zheng, Y. Satta, Functional evolution of avian RIG-I-like receptors. *Genes* **9**, 456 (2018).
20. V. Krchlíková *et al.*, Dynamic Evolution of Avian RNA Virus Sensors: Repeated Loss of RIG-
I and RIPLET. *Viruses* **15**, 3 (2022).
21. T. J. Hayman *et al.*, RIPLET, and not TRIM25, is required for endogenous RIG-I-dependent
antiviral responses. *Immunology and cell biology* **97**, 840-852 (2019).
22. Y. Xiao, M. B. Reeves, A. F. Caulfield, D. Evseev, K. E. Magor, The core promoter controls
basal and inducible expression of duck retinoic acid inducible gene-I (RIG-I). *Molecular
immunology* **103**, 156-165 (2018).
23. C. Cadena *et al.*, Ubiquitin-dependent and-independent roles of E3 ligase RIPLET in innate
immunity. *Cell* **177**, 1187-1200. e1116 (2019).
24. W. G. Dundon, A. Milani, G. Cattoli, I. Capua, Progressive truncation of the Non-Structural
1 gene of H7N1 avian influenza viruses following extensive circulation in poultry. *Virus
research* **119**, 171-176 (2006).
25. K. J. Bryson *et al.*, Comparative Analysis of Different Inbred Chicken Lines Highlights How
a Hereditary Inflammatory State Affects Susceptibility to Avian Influenza Virus. *Viruses* **15**,
591 (2023).
26. J. de Wit *et al.*, Major difference in clinical outcome and replication of a H3N1 avian
influenza strain in young pullets and adult layers. *Avian pathology* **49**, 286-295 (2020).
27. J. Guan, Q. Fu, S. Sharif, Replication of an H9N2 avian influenza virus and cytokine gene
expression in chickens exposed by aerosol or intranasal routes. *Avian Diseases* **59**, 263-
268 (2015).
28. H. Kato *et al.*, Differential roles of MDA5 and RIG-I helicases in the recognition of RNA
viruses. *Nature* **441**, 101-105 (2006).
29. L. Xu, D. Yu, Y. Fan, Y.-P. Liu, Y.-G. Yao, Evolutionary selection on MDA5 and LGP2 in the
chicken preserves antiviral competence in the absence of RIG-I. *Journal of Genetics and
Genomics= Yi Chuan xue bao* **46**, 499-503 (2019).
30. D. Evseev, D. Miranzo-Navarro, X. Fleming-Canepa, R. G. Webster, K. E. Magor, Avian
Influenza NS1 Proteins Inhibit Human, but Not Duck, RIG-I Ubiquitination and Interferon
Signaling. *Journal of Virology* **96**, e00776-00722 (2022).

31. A. Iwamoto, H. Tsukamoto, H. Nakayama, H. Oshiumi, E3 Ubiquitin Ligase Riplet Is
Expressed in T Cells and Suppresses T Cell–Mediated Antitumor Immune Responses. *The*
Journal of Immunology **208**, 2067-2076 (2022).
32. M. Elsheimer-Matulova *et al.*, Interleukin 4 inducible 1 gene (IL4I1) is induced in chicken
phagocytes by Salmonella Enteritidis infection. *Veterinary Research* **51**, 1-8 (2020).
33. S. A. Nish *et al.*, T cell-intrinsic role of IL-6 signaling in primary and memory responses.
elife **3**, e01949 (2014).
34. D. T. Thoresen, D. Galls, B. Götte, W. Wang, A. M. Pyle, A rapid RIG-I signaling relay
mediates efficient antiviral response. *Molecular cell* **83**, 90-104. e104 (2023).
35. J. Cornelissen, J. Post, B. Peeters, L. Vervelde, J. Rebel, Differential innate responses of
chickens and ducks to low-pathogenic avian influenza. *Avian Pathology* **41**, 519-529
(2012).
36. N. P. Liau *et al.*, The molecular basis of JAK/STAT inhibition by SOCS1. *Nature*
communications **9**, 1558 (2018).
37. D. C. Palmer, N. P. Restifo, Suppressors of cytokine signaling (SOCS) in T cell
differentiation, maturation, and function. *Trends in immunology* **30**, 592-602 (2009).
38. C. Vazquez, C. Y. Tan, S. M. Horner, Hepatitis C virus infection is inhibited by a
noncanonical antiviral signaling pathway targeted by NS3-NS4A. *Journal of Virology* **93**,
e00725-00719 (2019).
39. T. Schmit *et al.*, Interferon-γ promotes monocyte-mediated lung injury during influenza
infection. *Cell reports* **38** (2022).
40. M. G. Koliopoulos *et al.*, Molecular mechanism of influenza A NS1-mediated TRIM25
recognition and inhibition. *Nature communications* **9**, 1820 (2018).
41. N. R. Meyerson *et al.*, Nuclear TRIM25 specifically targets influenza virus
ribonucleoproteins to block the onset of RNA chain elongation. *Cell host & microbe* **22**,
627-638. e627 (2017).
42. M. Okamoto, T. Kouwaki, Y. Fukushima, H. Oshiumi, Regulation of RIG-I activation by K63-
linked polyubiquitination. *Frontiers in immunology* **8**, 1942 (2018).
43. H. Fischer, E. Tschachler, L. Eckhart, Pangolins lack IFIH1/MDA5, a cytoplasmic RNA sensor
that initiates innate immune defense upon coronavirus infection. *Frontiers in immunology*
11, 939 (2020).
44. H. Oshiumi, M. Matsumoto, T. Seya, Ubiquitin-mediated modulation of the cytoplasmic
viral RNA sensor RIG-I. *The journal of biochemistry* **151**, 5-11 (2012).
45. Y. Chen *et al.*, Duck RIG-I CARD domain induces the chicken IFN-β by activating NF-κB.
Biomed Research International **2015** (2015).
46. B. Rieblinger *et al.*, Strong xenoprotective function by single-copy transgenes placed
sequentially at a permissive locus. *Xenotransplantation* **25**, e12382 (2018).
47. J. S. Long *et al.*, Species specific differences in use of ANP32 proteins by influenza A virus.
Elife **8**, e45066 (2019).
48. H. Sid, S. Hartmann, C. Winter, S. Rautenschlein, Interaction of influenza A viruses with
oviduct explants of different avian species. *Frontiers in Microbiology* **8**, 1338 (2017).
49. M.-C. Van de Lavoir *et al.*, Interspecific germline transmission of cultured primordial germ
cells. *PLoS One* **7**, e35664 (2012).
50. M.-C. Van de Lavoir *et al.*, Germline transmission of genetically modified primordial germ
cells. *Nature* **441**, 766-769 (2006).

51. P. A. Leighton, M. C. van de Lavoir, J. H. Diamond, C. Xia, R. J. Etches, Genetic modification
of primordial germ cells by gene trapping, gene targeting, and φC31 integrase. *Molecular*
Reproduction and Development: Incorporating Gamete Research **75**, 1163-1175 (2008).
52. V. Hamburger, H. L. Hamilton, A series of normal stages in the development of the chick
embryo. *Journal of morphology* **88**, 49-92 (1951).
53. J. Whyte *et al.*, FGF, insulin, and SMAD signaling cooperate for avian primordial germ cell
self-renewal. *Stem cell reports* **5**, 1171-1182 (2015).
54. B. Schusser *et al.*, Immunoglobulin knockout chickens via efficient homologous
recombination in primordial germ cells. *Proceedings of the National Academy of Sciences*
110, 20170-20175 (2013).
55. R. Hernandez, D. T. Brown, Growth and maintenance of chick embryo fibroblasts (CEF).
Current protocols in microbiology **17**, A. 4I. 1-A. 4I. 8 (2010).
56. R. Brauer, P. Chen, Influenza virus propagation in embryonated chicken eggs. *JoVE*
(Journal of Visualized Experiments), e52421 (2015).
57. F.-k. Kong, C.-l. H. Chen, A. Six, R. D. Hockett, M. D. Cooper, T cell receptor gene deletion
circles identify recent thymic emigrants in the peripheral T cell pool. *Proceedings of the*
National Academy of Sciences **96**, 1536-1540 (1999).
58. N. B. Burkhardt *et al.*, The long pentraxin PTX3 is of major importance among acute phase
proteins in chickens. *Frontiers in Immunology* **10**, 124 (2019).
59. I. Monne *et al.*, Emergence of a highly pathogenic avian influenza virus from a low-
pathogenic progenitor. *Journal of virology* **88**, 4375-4388 (2014).
60. S. Trapp *et al.*, Major contribution of the RNA-binding domain of NS1 in the pathogenicity
and replication potential of an avian H7N1 influenza virus in chickens. *Virology journal* **15**,
1-12 (2018).
61. C. Ward *et al.*, Design and performance testing of quantitative real time PCR assays for
influenza A and B viral load measurement. *Journal of clinical virology* **29**, 179-188 (2004).
62. T. von Heyl *et al.*, Loss of αβ but not γδ T cells in chickens causes a severe phenotype.
European Journal of Immunology **53**, 2350503 (2023).

Figures

Figure 1. Generation of PGCs, and susceptibility to infection using *in vitro* and *in vivo* systems. a. Promoter activity of the duck *RNF135* was examined by generating different deletion mutants tested in chicken DF-1 cells; the promoter activity was then evaluated by measuring the NanoLuc/Firefly ratio (n=3). **b.** Diagram of both constructs used to generate *RIG-I*-expressing chicken (upper diagram), using the previously identified duck *RIG-I* promoter (22) and *RNF135*-expressing chickens (lower diagram), using the duck *RNF135* promoter, which activity was identified in this study. **c.** Primordial germ cells (PGCs) that express the duck *RIG-I* under the duck *RIG-I* promoter were derived into PGC fibroblasts (PGCFs) and were infected later with avian influenza virus H9N2 to stimulate *RIG-I* expression upon influenza infection; the cells were infected for 18h with low pathogenic avian influenza virus H9N2 and subsequently stained for FLAG-Tag (red staining); MOCK control represents uninfected cells, and were not positive for anti-FLAG-Tag; DF-1 cells that express the FLAG-tagged duck *RIG-I* under the chicken beta-actin promoter were used as a positive control (red staining). **d.** Quantification of newly produced viral particles after infection of CEFs; CEFs were isolated from different transgenic embryos and experimentally infected with LPAIV H9N2 at two different multiplicities of infection (MOI 0.1, and 1); Supernatants were collected at 24hpi and titrated on MDCK cells; no significant differences were observed between the groups ($p>0.05$). **e.** Quantification of newly produced viral particles after infection of embryonated eggs. 10 day-old embryonated eggs were infected with LPAIV H9N2 at 10^3 FFU/egg; allantois fluid was collected 24hpi and titrated on MDCK cells; no significant differences were observed between the groups ($p>0.05$). Error bars indicate the standard error of mean (SEM); Depending on the normal distribution of the data, multiple group comparison was done either with one-way ANOVA or Independent-Samples Kruskal-Wallis Test

Figure 2. Generation and immunophenotypic characterization of *RIG-I* and *RNF135*-expressing chickens. a. Weekly weight monitoring of the generated heterozygous birds (*p*>0.05) (n=10). **b.** Representative picture of the generated heterozygous birds from left to right at four weeks of age: WT, *RNF135*-expressing chicken, and *RIG-I*-expressing chicken. **c.** RT-PCR of the transgenic expression of *RIG-I* and *RNF135* in different organs. **d.** Analysis of duck RIG-I and duck RNF135 expression in various tissues using reads per kilobase per million mapped reads (RPKM). **e.** Assessment of different immune cell

populations in *RIG-I*-expressing chickens compared to their WT siblings. (∗) indicate

statistical differences between groups tested simultaneously (*p*<0.05). Depending on the

normal distribution of the data, two group comparison was done with the Wilcoxon rank-

sum test or two samples T-test

Figure 3. Assessment of different T cell subpopulations in *RIG-I*-expressing
chickens compared to their WT siblings

a. PBMCS were analyzed for γδTCR1+/CD8α+ T cells at 12 weeks. B) PBMCS were

analyzed for αβTCR2,3+T cells and CD4+ or CD8α+T cells at 12 weeks (*p*<0.05). **c.**

Activation of isolated T cells from 12-week-old *RIG-I*-expressing chickens compared to

their WT siblings ($p<0.05$); cells were sorted according to their TCR expression, stimulated

with Concanavalin A, and quantified by flow cytometry at different time-points. The results

indicate the number of positive cells per 100 viable TCR + cells. Error bars indicate the

standard error of mean (SEM); (*) indicate statistical differences between groups tested

simultaneously ($p<0.05$). Depending on the normal distribution of the data, two group

comparison was done with the Wilcoxon rank-sum test or two samples T-test, while

multiple group comparison was done either with one-way ANOVA or Independent-

Samples Kruskal-Wallis Test

Figure 4. Simultaneous expression of *RIG-I* and *RNF135* in the chicken does not

lead to differences in the adaptive immune phenotype compared to WT birds

Two-time points, eight and twelve weeks of age, were chosen based on the data obtained from *RIG-I*-expressing chickens to assess the immunophenotype of *RIG-I-RNF135*-expressing chickens. PBMCS were isolated and analyzed for B cells, αβTCR2,3+ or γδTCR1 + T cells (**a**) as well as for CD4+ (B) and CDα8+ T cells (**c**). No significant differences were detected between the analyzed groups (*p*>0.05). Depending on the normal distribution of the data, two group comparison was done with the Wilcoxon rank-sum test or two samples T-test

1010

Figure 5. H7N1-challenge experiment reveals the susceptibility of *RIG-I*-expressing chickens and the role of *RNF135* in effective *RIG-I* antiviral response.

The generated transgenic chickens were challenged with H7N1 and assessed at two days
post-infection (dpi) and six dpi for different parameters. **a.** Percentage of animals
presenting clinical symptoms. **b.** Probability of survival in the challenged groups. **c.** Weight
gain in challenged groups compared to WT- and *RIG-I-RNF135*-MOCK- controls ($p<0.05$).
d. Viral replication rate in two main target organs, caecum, and lung ($p<0.05$); the
horizontal line indicates the detection threshold of the PCR based on the signal obtained
from the uninfected controls, which is $\log_{10}(10^3)$. **e.** Expression levels of transgenes in the
caecum and the lung upon H7N1-challenge. Error bars indicate standard error of mean
(SEM); (*) or different letters indicate statistical differences between groups tested
simultaneously ($p<0.05$). Depending on the normal distribution of the data, multiple group
comparison was done either with one-way ANOVA or Independent-Samples Kruskal-
Wallis Test

1025

1026

1027

1028

1029

1030

1031

1032

1033

1034

1035

1036

1037

1038

Figure 6. Developed lesions in H7N1-challenged birds a. Macroscopical lesion score

of the lungs showing a significant increase of lesions in *RIG-I*-expressing birds at six dpi

(*p*<0.05) **b.** Histology of the caecum showing the typical structure of the epithelium in the

control groups. *RIG-I-RNF135* challenged birds showing epithelial hyperplasia with

necrosis that diminished by 6 dpi. *RIG-I*-expressing chickens did not show necrotic lesions

but pronounced epithelial hyperplasia that lessened by six dpi. Asterix indicates epithelial

hyperplasia with the mitotic figures, while the arrow indicates necrotic epithelial cells; 200x.

Error bars indicate standard error of mean (SEM); (∗) indicate statistical differences

between groups tested simultaneously (*p*<0.05). Depending on the normal distribution of

the data, multiple group comparison was done either with one-way ANOVA or

Independent-Samples Kruskal-Wallis Test

Figure 7. Infection of transgenic chickens with H7N1 leads to upregulated
inflammatory   genes.   Venn   diagram   of   fluidigm   qRT-
PCR array of naive and challenged birds with H7N1 showing significantly upregulated
genes in various organs. Significant DEGs were identified by comparing the relative
expression values for every chicken line to the WT-MOCK individually per timepoint, with
a significance level set at $p < 0.05$; fold change >1 (n ≥ 5-time point). **a.** Upregulated genes
in *RIG-I-RNF135*-expressing chickens without infection **b.** Upregulated genes in the
caecum between H7N1-challenged groups at 2dpi **c.** Upregulated genes in the lung
between H7N1-challenged groups at 2dpi **d.** Upregulated genes in the lung between
H7N1-challenged groups at 6dpi **e.** Upregulated genes in the spleen between H7N1-
challenged groups at 6dpi

1062

1063

1064

1065

1066

1067

Figure 8. Gene expression in the lung upon H7N1-infection

Significant DEGs were identified by via fluidigm qPCR array and comparison of the relative

expression values between H7N1-challenged groups. **a.** Significantly upregulated genes

after comparison of infected *RIG-I-* with *RIG-I-RNF135*-expressing chickens at two dpi. **b.**

Significantly upregulated genes after comparison of infected *RIG-I-*with *RIG-I-RNF135*-

expressing chickens at 6dpi. **c.** Significantly upregulated genes after comparison of

infected *RIG-I-* with *RIG-I-RNF135*-expressing chickens at two dpi. **d.** Significantly

upregulated genes after comparison of infected *RIG-I-*expressing with chickens WT-

chickens at six dpi. Significance levels were set at $p < 0.05$; fold change >1 (n ≥ 5-time

point)

1078

1079

1080

Figure 9. H3N1-challenge experiment confirms the exacerbated disease phenotype of *RIG-I* and *RIG-I-RNF135* chickens infected with virulent avian influenza viruses

The generated transgenic chickens were challenged at 28 weeks of age with H3N1 and assessed for different parameters. **a.** Viral shedding based on tracheal swabbing and viral RNA load analysis. b. Probability of survival in the challenged groups. **c.** Expression of *RIG-I*, *RNF135*, and influenza-regulated genes in the duodenum, lung, and spleen. **d.** Histological assessment of the reproductive tract; WT-MOCK: normal macroscopical and histological appearance of the salpinx; RNF-RIG-I-H3N1: severe atrophy with mild fibrinous peritonitis in the infundibulum (asterisk), severe lymphoplasmacytic salpingitis (arrowheads) (40x) the magnum (20x) and vasculitis in the magnum vessels (200x). **e.** Scoring of lesions in the reproductive tract in all challenged groups, starting with the upper row WT-MOCK, WT-H3N1, *RIG-I*-H3N1 and *RIG-I-RNF135*-H3N1; FP: fibrinous peritonitis, LPS: lymphoplasmacytic salpingitis, LPV: lymphoplasmacytic vasculitis, EYP: egg yolk peritonitis. Error bars indicate standard error of mean (SEM); (∗) indicate statistical differences between groups tested simultaneously ($p<0.05$). Depending on the normal distribution of the data, multiple group comparison was done either with one-way ANOVA or Independent-Samples Kruskal-Wallis Test