1   **Reinstatement of *RIG-I* in chickens via genetic modification reveals new insights into the**
2   **dynamic evolution of avian immune sensors**
3

4   Hicham Sid[1*], Theresa von Heyl[1], Sabrina Schleibinger[1], Romina Klinger[1], Rodrigo
5   Guabiraba[3], Vanaique Guillory[3], Benjamin Schade[4], Daniel Elleder[5], Samantha Sives[6],
6   Lonneke Vervelde[6], Sascha Trapp[3], Benjamin Schusser[1,2]
7

[1] TUM School of Life Sciences, Weihenstephan, Department of Molecular Life Sciences,
Reproductive Biotechnology, Technical University of Munich, 85354, Freising, Germany
[2] Center for Infection Prevention (ZIP), Technical University of Munich, 85354 Freising,
Germany
[3] UMR ISP, INRAE, Université de Tours. 37380, Nouzilly, France.
[4] Bavarian Animal Health Service, Department of Pathology, 85586, Poing, Germany;
[5] Institute of Molecular Genetics of the Czech Academy of Sciences, Prague, Czech Republic
[6] Division of Immunology, The Roslin Institute and Royal (Dick), School of Veterinary Studies,
University of Edinburgh, UK
17
18

19   *Correspondence: hicham.sid@tum.de

21   **Graphical abstract**
22

Abstract

Retinoic acid-inducible gene I (*RIG-I*) activates mitochondrial antiviral signaling proteins, initiating the antiviral response. *RIG-I* and *RNF135*, a ubiquitin ligase regulator, are missing in domestic chickens but conserved in mallard ducks. It was long believed that chickens' *RIG-I* loss was linked to increased avian influenza susceptibility. We reinstated both genes in chickens and examined their susceptibility to the avian influenza virus H7N1. Uninfected *RIG-I-*expressing chickens exhibited shifts in T and B cells, while the H7N1 infection led to severe disease, persistent weight loss, and increased viral replication. Conversely, the co-expression of *RIG-I* and *RNF135* reduced the viral replication and was associated with high inflammatory response. Our data indicate that the loss of *RIG-I* in chickens likely evolved to counteract deleterious inflammation caused by viral infection. We highlight the effects of restoring evolutionary lost genes in birds and suggest a new immunological approach to reduce viral replication and prevent infection.

Introduction:

Avian influenza virus (AIV) is an epizootic pathogen with zoonotic potential [1] that recently caused devastating outbreaks worldwide, leading to the loss of more than 131 million birds due to animal death and culling [2]. The viral reservoir of AIV are wild birds of the orders *Anseriformes* (ducks, geese, and swans) and *Charadriiformes* (gulls and terns) [3], which, compared to chickens or other galliform birds, exhibit milder clinical symptoms despite efficient viral replication [4]. Certain genomic features of the duck, including a functional *retinoic acid-inducible gene I (RIG-I)* gene, were reported to be associated with their relative resistance to clinical avian influenza infection [5]. *RIG-I* is a cytosolic RNA sensor that recognizes and binds to the 5' triphosphate end (5'-ppp) [6]. It forms a first line of antiviral defense as a pathogen recognition receptor (PRR) against RNA viruses [7]. Upon activation, the *RIG-I* interacts with the mitochondrial antiviral signaling proteins (MAVS), inducing a pro-inflammatory antiviral response characterized by the upregulation of type I and type III interferons (IFNs) followed by the expression of IFN stimulated genes (ISGs) [8]. The activity of *RIG-I* is believed to be controlled by post-translational modification of *tripartite motif-containing protein 25 (TRIM25)* [9] and *RING finger protein 135 (RNF135*, also known as *Riplet* or *REUL*). The latter was found to modify *RIG-I* by lysine 63-linked polyubiquitination of the C-terminal region of the *caspase activation and recruitment domain* (CARD) [10], leading to a stronger *RIG-I* signal transduction [11]. The physiological role of mammalian *RIG-I*, its function in viral infections, and its parallel evolution with other RLRs are well documented [9, 12, 13, 14]. Furthermore, the knockout (KO) of *RIG-I* in mice has shown that this gene may have additional functions related to adaptive immunity [12]. *RIG-I* KO mice exhibited a defect in migratory dendritic cells in combination with a reduced frequency of polyfunctional effector T cells upon influenza infection [15].

In ducks, *RIG-I* elicits a potent interferon (IFN) response within the first few hours after infection, leading to survival against most AIV strains [16]. In contrast, chickens lack *RIG-I*, which probably lost its function in a common ancestor of galliform birds [17, 18]. Interestingly, a recent study detected disrupted *RIG-I* pseudogenes in some Galliformes, including the helmeted guineafowl (*N. meleagris*) and the northern bobwhite (*C. virginianus*) [19]. Authors hypothesized a compensatory evolution of *melanoma differentiation-associated gene-5 (MDA5)* that accompanied the gradual loss of *RIG-I* in chickens [19]. The evolutionary loss of *RIG-I* in different galliform birds correlated with the simultaneous loss of its ubiquitin ligase *RNF135* [19], which has been described to be critical for *RIG-I* ubiquitination in mammals, unlike *TRIM25*, which is increasingly believed to be less important for *RIG-I* signaling [20]. Reasons behind the loss of *RIG-I* and its ubiquitination factor in chickens are still unknown and remain enigmatic,

105 especially given the virus-limiting effect of duck *RIG-I* overexpression in AIV-infected chicken
106 DF-1 cells [5].

107
To date, the effect of *in vivo* expression of *RIG-I* in chickens has never been studied, and no
transgenic chicken lines expressing duck *RIG-I* have been generated, likely due to the lack of
suitable biotechnological tools in avian research. Here, we used chicken primordial germ cells
(PGCs) to develop genetically modified chickens expressing duck *RIG-I* and *RNF135* under
the control of their respective duck promoters. The generated birds were healthy and developed
normally compared to their WT siblings. In the absence of infection, we observed an alternative
phenotype of the adaptive immune system of *RIG-I*-expressing chickens, particularly regarding
the T cell population. In contrast, the co-expression of *RNF135* with *RIG-I* contributed to a
balanced adaptive immune phenotype that appeared to be similar to WT birds. Infection
experiments with an H7N1 AIV led to severe clinical disease associated with a strong
inflammatory response, high *IFN-γ* expression, and elevated viral replication in *RIG-I*-
expressing chickens compared to other challenged groups. In contrast, infected *RIG-I-RNF135*-
expressing chickens presented with inflammation, limited viral replication, and increased *IFN-*
α expression levels. The obtained data reveal the immunological functions of *RIG-I* in chickens
and the benefit of learning from less susceptible species to influenza infection to improve the
resilience of the immune system towards infection.

124
125 **Results:**
126
Generation of PGCs that express duck *RIG-I* and *RNF135* under the respective duck
promoters
The genetic modification of PGCs represents a crucial step for generating transgenic chickens.
PGCs are the precursors of sperm and eggs in adult animals, and therefore, we used them to
produce chimeric roosters paired with WT layer hens to obtain the desirable transgene. To
generate chickens that express *RIG-I* and *RNF135*, we started by cloning duck *RIG-I* and
RNF135, which were subsequently inserted into two different expression vectors (Fig. 1A, 1B).
RIG-I and *RNF135* were expressed under their respective duck promoters and used to generate
PGCs expressing both genes separately. While the duck *RIG-I* promoter was previously
described [21], we determined the activity of the duck *RNF135* promoter, which was examined
by the generation of different deletion mutants tested in the NanoDLR™ Assay System.
Therefore, different deletion mutants were generated, including p1577, p1001, p349, and p197.
Since we did not detect a core promoter activity of the duck *RNF135*, we used the full-length
sequence of the duck *RNF135* for the generation of duck *RNF135*-expressing PGCs (Fig. 1A).
The assembled expression vectors that were used to generate PGCs are presented in Fig. 1B.

Due to the unavailability of commercial antibodies for detecting the duck *RIG-I*, we inserted a

FLAG-Tag on the C-terminus to facilitate its detection using anti-FLAG antibodies. The

activity of the duck *RIG-I* was examined by differentiating the *RIG-I*-expressing PGCs into

PGC-derived fibroblasts (Fig. 1C) that were infected with a low pathogenic avian influenza

virus (LPAIV) H9N2 (Fig. 1D). Results showed that the infection led to activation of *RIG-I* as

shown by FLAG-Tag staining. At the same time, the MOCK-infected cells remained negative

for *RIG-I* expression (Fig. 1D).

The expression of *RIG-I* did not limit the viral replication under *in vitro* and *in ovo*

conditions

According to previously published data by Barber et al. [5], the overexpression of duck *RIG-I* in

chicken DF-1 cells reduced the replication of avian influenza viruses. Authors constitutively

expressed *RIG-I* under the control of the CMV promoter using expression vector pcDNA 3.1.

Therefore, we wanted to examine if the expression of *RIG-I* under the control of the duck RIG-

I promoter can lead to a similar effect in limiting virus replication. We isolated CEFs and

embryonated eggs obtained from the generated transgenic chickens and infected them with

LPAIV H9N2. After virus infection, supernatants and allantoic fluid were collected from CEFs

and embryonated eggs, respectively. We quantified the newly produced viral particles using a

focus-forming assay. Surprisingly, transgene expression did not significantly affect the viral

replication in both tested systems (Fig. 1D and 1E), which was the opposite in the previously

published study [5]. This highlights the importance of the chosen promoter, which may have

affected the expression of *RIG-I* and influenced the innate immune response towards the

influenza virus.

Phenotypic characterization of the genetically modified chickens

Upon the generation of *RIG-I*- and *RNF135*-expressing chicken lines, we wanted to ensure that

the genetic modification did not negatively affect the growth of the generated transgenic birds.

Therefore, we monitored their development by weekly measuring their body weights (Fig. 2A).

Both generated chicken lines showed comparable growth to their WT siblings. The animals

matured sexually, and no harmful phenotype was detected (Fig. 2A and 2B). The expression of

both genes was examined via RT-PCR, revealing that both genes are expressed differentially

in various tissues (Fig. 2C).

According to previously published studies, mammalian *RIG-I* affects adaptive immunity,

mainly T cells [15]. Therefore, we sought to examine whether the re-expression of *RIG-I* in

chickens will have similar effects. We analyzed the numbers of peripheral blood mononuclear

cells (PBMCs) using flow cytometry to investigate the possible impact of expressing *RIG-I*,

RNF135, or both on immune cell counts. No differences were observed in the number of

immune cells between *RNF135*-expressing chickens and their WT siblings (Supplementary file

1). However, *RIG-I*-expressing chickens exhibited a significantly higher number of αβ and γδ

T cells as well as B cells in comparison to their WT siblings ($p<0.05$) (Fig. 2D). This was not

the case for monocyte counts, where no significant differences were observed (Fig. 2D). In

addition, no significant differences were observed in *RIG-I*-expressing chickens compared to

WT birds regarding the levels of IgM and IgY at 12 and 14 weeks of age (Supplementary file

2).

Further analysis of different T cell subpopulations indicated that *RIG-I*-expressing chickens had

a significantly higher number of CD4+T cells and a significantly lower number of CD8α+T

cells ($p<0.05$) (Fig. 3). T-cell activation was quantified after lectin activation using

Concanavalin A (ConA). *RIG-I*-expressing chickens had a higher level of T cell activation in

TCRγδ/CD25+ T cells, which was not the case for TCRαβ1, αβ2/CD25+ T cells (Fig. 3).

Heterozygous birds expressing each gene separately were crossed to obtain birds that

simultaneously express both genes *RIG-I* and *RNF135*. The obtained *RIG-I-RNF135*-

expressing chickens were monitored weekly for weight gain and closely investigated at eight

and twelve weeks of age for possible alternate immune phenotypes similar to those observed in

RIG-I-expressing birds. Unlike *RIG-I*-expressing chickens, no significant differences in T cells

or B cell counts were detected in *RIG-I-RNF135*-expressing chickens in comparison to their

WT siblings (Fig. 4). This was also the case for body weight gain, where no significant

differences were observed in comparison to WT birds (Supplementary file 3).

These data indicate that the *RIG-I*, an innate immune sensor, can influence adaptive immunity

by causing shifts in T and B cell populations. In contrast, co-expression of *RNF135* with *RIG-*

I seems to balance the adaptive immune cell populations, which was comparable to WT birds.

RNF135 is required for efficient antiviral activity of RIG-I

The role of *RNF135* as an essential ubiquitination factor that supports the antiviral activity of

RIG-I was previously determined in mammalian cells [11]. Later on, *RNF135* was found to be the

obligatory ubiquitin for *RIG-I* [22], while so far, this was not known in birds. We investigated the

effect of expressing *RNF135* on the antiviral activity of *RIG-I* in genetically modified chickens.

We examined the susceptibility of the generated transgenic lines towards an H7N1 AIV, a direct

precursor of a highly pathogenic avian influenza virus (HPAIV), known to cause severe acute

respiratory disease in chickens [23]. Birds were challenged with the virus and monitored for

clinical signs, weight gain, viral replication, and lesion development. *RIG-I*-expressing

chickens had the highest rate of morbidity, as indicated in Fig. 5A. The clinical symptoms

started within the first two days of infection, where the morbidity rate reached a significant rate of 44% at one-day post-infection (dpi) ($p<0.05$) and increased to 67% at two dpi ($p<0.05$), which then decreased to 18% by three dpi ($p<0.05$). The onset of clinical disease was accompanied by a significant weight loss at two dpi that remained significantly low at six dpi ($p<0.05$) (Fig. 5C) and coincided with a pulmonary lesion score that increased from 0.8 at two dpi to 2.3 at six dpi (Fig. 6A and 6B). This was also confirmed by histological assessment of the caecum that revealed necrotic lesions associated with pronounced epithelial hyperplasia (Fig. 6B). *RIG-I-RNF135*-expressing chickens exhibited increasing clinical illness, from 13% of sick animals at one dpi to 50% at three dpi ($p<0.05$). Severe clinical symptoms associated with mortality in this group lasted until six dpi.

The expression of *RNF135* alone resulted in a mortality rate comparable to WT birds. In contrast, the clinical symptoms in *RNF135*-expressing chickens lasted till three dpi with a total of 25% sick animals (Fig. 5). This was not the case for WT chickens that exhibited clinical symptoms only for the first two days after infection and had a morbidity rate of 13% and 19% at one and two dpi, respectively (Fig. 5).

The quantification of the virus genome copies via qRT-PCR showed that the expression of *RIG-I* together with *RNF135* resulted in an efficient antiviral effect, as demonstrated by the lowest amount of viral genome copies in the caecum compared to all the other challenged chicken lines (Fig. 5D), with a significant low viral replication rate compared to *RIG-I* expressing chickens at two and six dpi ($p<0.05$). In the lung, challenged *RIG-I-RNF135*-expressing chickens displayed a virus replication rate similar to WT birds and a significantly lower replication rate than *RIG-I*-expressing chickens at two dpi ($p<0.05$). Conversely, *RIG-I*-expressing chickens had the highest viral replication rate in the lung and the caecum among all investigated groups, with a significant difference compared to *RIG-I-RNF-135*-expressing chickens ($p<0.05$) (Fig. 5D). The amount of viral nucleic acid remained at a high level at six dpi, indicating the lack of viral clearance in these birds, which was not the case for the other challenged chicken lines (Fig. 5D).

The observed low viral titer in *RIG-I-RNF135*-expressing chickens confirms the crucial role of *RNF135* as a ubiquitin ligase at the early stage of infection. Also, it demonstrates that the chicken *TRIM25* does not overtake the part of *RNF135* in *RIG-I*-expressing chickens.

The expression of *RIG-I* and *RNF135* in H7N1-challenged birds coincided with the acute phase of infection.

We aimed to quantify possible changes in the expression levels of both transgenes in the challenged animals after the H7N1 infection. We quantified *RIG-I* and *RNF135* expressions via qRT-PCR in the lung, and the caecum (Fig. 5E). The caecal expression of *RIG-I* in virus-

infected *RIG-I-RNF135* chickens was increased by ~26 folds. At the same time, infected *RIG-I* chickens exhibited an increase of ~24-fold compared to *RIG-I-RNF135* MOCK controls. Infected *RIG-* and *RIG-RNF135*-expressing chickens had a comparable *RIG-I* expression to *RIG-RNF135*-MOCK controls with no significant upregulation (Fig. 5E). Overall, we observed lower expression of *RNF135* in the caecum and lungs compared to *RIG-I-RNF135*-MOCK controls. The expression increased by 1.4 folds in *RNF135*-infected birds and ~2 folds in *RIG-I-RNF135*-infected chickens. In the lung, the *RNF135* expression at two dpi was upregulated by ~6 folds and ~4 folds in infected *RIG-I* and *RIG-I-RNF135*-expressing chickens, respectively. At six dpi, the *RNF135* expression was at a comparable level with the *RIG-I-RNF135* MOCK controls (Fig. 5E). These data indicate that the infection led to a quick upregulation of *RIG-I* at the acute phase of infection, which quickly decreased afterward.

Differential regulation of innate immune genes in naïve as well as in H7N1-challenged *RIG-I*, *RNF135*, and *RIG-I-RNF135*-expressing chickens

We speculated that the observed acute inflammatory reaction could be due to a differential regulation of inflammatory genes related to *RIG-I* signaling. Therefore, we sought to quantify the differentially expressed genes (DEGs) between transgenic and WT birds. We used the Fluidigm qRT-PCR array to identify changes in the expression of selected innate immune genes in both naïve and H7N1-challenged birds [24]. The results revealed that in the absence of infection, *RIG-I-RNF135*-expressing chickens already had a significantly higher expression of ISGs in the caecum, including *ISG12-2* and *OASL,* as well as the transcription factor *EGR1*. In contrast, others, including *EGR1*, *IL41*, *Il1β,* and *IL8,* were significantly upregulated in the spleen (Fig. 7 and Supplementary file 4).

The comparison of challenged groups with non-infected controls (WT-MOCK) indicated that *RIG-I* chickens expressed several inflammatory genes in the caecum and lungs that were not expressed in the other groups. This included *IFITM1*, *IL6,* and *LIPI* (Fig. 7 and Supplementary file 4). At two dpi, we observed the upregulation of genes involved in the JAK-STAT signaling pathway, including *IL-6* and *STAT3*. The significant upregulation of inflammatory and immune regulatory genes in the *RIG-I*-expressing chickens persisted until six dpi, where over 20 genes were exclusively upregulated in the spleen (Fig. 7 and Supplementary file 4). This was not the case for *RIG-I-RNF135*-expressing chickens and WT chickens with an exclusive expression of three and seven genes, respectively (Fig. 7 and Supplementary file 4). In addition, the analysis of expressed genes in the spleen revealed that only *RIG-I-RNF135* chickens expressed *IRF7* and *TRIM25*, which was not the case for *RIG-I* chickens (Fig. 7 and Supplementary file 4).

The functional enrichment analysis of the genes involved in the biological processes of transgenic chickens indicated that the regulated genes in *RIG-I*-expressing chickens were highly

involved in metabolic activities. In contrast, those in *RIG-I-RNF135*-expressing birds were primarily involved in regulatory mechanisms (Supplementary file 5). Overall, the obtained data confirm that *RIG-I*-expressing chickens exhibited a significant increase of inflammatory cytokines that were not observed in other challenged birds, which may explain the acute inflammatory reaction in these birds.

High expression of *IFN-γ* is associated with significant virus replication in *RIG-I*-expressing birds

Since we observed a significant reduction of viral genome copies in the *RIG-I-RNF135*-expressing chickens compared to *RIG-I*-expressing birds, we compared the DEGs between H7N1-infected groups. This helped us determine possible factors behind the increased H7N1 replication in *RIG-I*-expressing birds. Interestingly, these birds lacked interferon upregulation compared to infected WT birds or *RIG-I-RNF135*-expressing chickens. We also found that the viral infection led to a significant increase in the expression of *IFN-γ* at six dpi when compared to H7N1-infected *RIG-I-RNF135*-expressing chickens (14-fold change increase) and WT birds (9-fold change increase) (Fig. 8) ($p<0.05$). In contrast, *RIG-I-RNF135*-expressing birds quickly exhibited a significant increase of *IFN-α* expression, already at two dpi, in comparison to *RIG-I*-expressing birds (12-fold change) as well as to WT birds (17-fold change) (Fig. 8). We concluded that the co-expression of *RNF135* and *RIG-I* in chickens leads to a significant increase of *IFN-α* expression, which was not observed when *RIG-I* was expressed solely. In addition, *RIG-I*-expression caused an upregulation of *IFN-γ* that was accompanied by a significant increase in virus genome copies.

Discussion:

The constant arms race between pathogens and the host affects different aspects of the immune system, including innate sensors. The reason for the loss of *RIG-I* in Galliformes, which correlated with the disappearance of the ubiquitin ligase *RNF135*, remained a mystery [19], especially given the established contributing role of *RIG-I* in the resilience towards influenza in ducks and other species [5, 25]. It was previously speculated that the gradual loss of *RIG-I* and *RNF135* in the chicken was possibly caused by the pathogen's resistance to the innate sensor or the disappearance of some relevant pathogens [19]. At the same time, preservation of the antiviral competence accompanied the loss of *RIG-I* and *RNF135* in chickens via the evolutionary selection of *MDA5* and *LGP2* [26]. The beneficial effect of the duck *RIG-I*-overexpression in chicken DF-1 cells was demonstrated by the limited replication of AIV in these transgene-expressing cells [5]. So far, no chickens expressing the duck *RIG-I* have been generated to investigate their antiviral response *in vivo*. In this study, we produced genetically modified chickens that express *RIG-I* with or without its ubiquitin ligase *RNF135* to examine

the physiological role of both genes and their function during AIV infection. Both genes were

cloned from the duck as it represents the most studied avian influenza reservoir with

evolutionary conserved *RIG-I* and *RNF135* [4, 19, 27]. The expression of both genes was controlled

under their respective duck promoters since the duck *RIG-I* was previously tested in chicken

cells (12). This strategy allows the strict control of gene expression that can be induced only

upon infection, which prevents undesirable production of inflammatory cytokines that may

cause autoimmune diseases [21]. While the duck *RIG-I* promoter was previously identified [21], we

described the activity of the duck *RNF135* promoter using chicken cells.

Without infection, *RIG-I* expression did not cause any harmful phenotype, but it led to different

adaptive immune cell counts compared to WT siblings. We observed that, unlike CD4+ T cells,

RIG-I-expressing chickens had a significantly lower number of CD8α+ T cells than WT-

chickens, which can be comparable to mice where *RIG-I* signaling limits the number of CD8$^+$

T cells [28]. This suggests the role of the *RIG-I* in priming T cell immunity in the chicken, which

can be similar to mice that exhibited a lack of T cell immunity associated with deficiencies in

migratory dendritic cell activation and viral antigen presentation [15] [12]. The effect of *RIG-I* on

adaptive immunity was further confirmed by others who reported defects in CD8+ and CD4+

T cells priming after *RIG-I* KO [29].

Conversely, *RIG-I-RNF135*-expressing chickens appeared to have a balanced adaptive immune

profile indistinguishable from WT birds. This observation demonstrates the possible role of

RNF135 in modulating the T cell immune response in birds, which was previously described

for Th1 response and cytotoxic T cells in mice [30]. Moreover, naïve *RIG-I-RNF135*-expressing

chickens had a significantly higher expression of *interleukin 4 inducible gene 1* (*IL4I1*) as well

as *protein nuclear translocation 7 (IRF7)* and *tripartite motif-containing protein 29 (TRIM29)*.

This may explain the differential adaptive immune phenotype between *RIG-I* and *RIG-I-*

RNF135-expressing chickens. Previous studies indicated the involvement of *IL4I1* in the

signaling to T and B lymphocytes [31] and the effective role of translocation factors such as *IRF7*

in chicken cells transfected with the duck *RIG-I* [21].

The lack of an antiviral effect after *in ovo* and *in vitro* experimental infections with LPAIV

H9N2 was discordant with the previously described antiviral effect of duck *RIG-I* in chicken

DF-1 cells [5]. Barber et al, observed low virus replication upon the overexpression of *RIG-I*

under control of the CMV promoter. The difference between our results and previously

published data can be related to the chosen promoter used by Barber et al, which may lead to a

higher upregulation of *RIG-I* and, consequently, a strong production of interferon-stimulated

genes.

H7N1-challenge experiments of the generated transgenic birds revealed that *RIG-I*-expressing

chickens had an early upregulation of proinflammatory cytokines such as *IL-6*, known to

support *IL1β* in suppressing regulatory T cells, which can lead to an uncontrolled increase in the number of CD4+ T cells [32]. Furthermore, the expression of *RIG-I* without *RNF135* was not beneficial in limiting H7N1 replication since the *RIG-I*-expressing chickens had the highest viral genome copies among all challenged groups. The requirement of *RNF135* for an efficient *RIG-I*-mediated antiviral effect was demonstrated by the significantly lower number of viral genome copies detected in *RIG-I-RNF135*-expressing chickens compared to *RIG-I*-expressing chickens, which was also previously described in mammalian cells [11]. In addition, we found that the H7N1 infection caused a significant upregulation of *RIG-I* in the caecum compared to the lungs. This can be because *RIG-I* expression rapidly takes place upon stimulation and may reach a peak by three hours post-infection [33]. Similar observations were described by Cornelissen et al. that detected a significant upregulation of *RIG-I* expression in lungs of ducks at 8h post H7N1 infection that significantly decreased at two dpi [34]. Infected *RIG-I*-expressing birds exhibited an acute inflammatory response and weight loss at two dpi that lasted till six dpi. Pang et al, reported that the inflammatory reaction associated with *RIG-I* signaling could be used by the influenza virus for its replicative advantage, particularly in the respiratory tract [13]. *RIG-I*-expressing birds showed several regulated immune genes compared to other infected groups, including upregulation of the *suppressor of cytokine signaling 1* (*SOC1*) in the caecum. This may explain the significant increase in viral genome copies and the observed T cell phenotype since *SOCS1* is a potent inhibitor of JAK/STAT signaling [35] and is involved in several mechanisms that regulate the T cell maturation, differentiation and function [36].

Interestingly, we detected significant upregulation of *IFN-γ* in the lungs of infected *RIG-I*-expressing chickens compared to infected *RIG-I-RNF135*- and WT-chickens, which can be responsible for lung-mediated injury and acute death caused by respiratory distress. The role of *IFN-γ* was previously described in *IFN-γ* KO mice that were less susceptible to lung inflammation and pathology upon influenza infection [37]. The lack of an antiviral effect in *RIG-I*-expressing chickens may also indicate that chicken *TRIM25* is dispensable for *RIG-I* efficient ubiquitination. Similar findings were described using human lung adenocarcinoma epithelial cells, where *TRIM25* did not participate in the endogenous *RIG-I*-dependent antiviral responses [20]. However, other studies done in mammalian models indicated that the deletion of *TRIM25* increases the susceptibility to influenza infection [38], supporting the evidence that *TRIM25* may bind directly to the viral RNA, thereby contributing to the restriction of influenza virus infection [39].

The low viral replication in *RIG-I-RNF135*-expressing chickens can be related to the effectiveness of *RNF135* in *RIG-I*-mediated antiviral activity. This was reflected by the upregulation of *IFN-α* expression in infected birds on one side and the absence of

immunophenotype in naïve birds on the other side, suggesting a stabilizing role of *RNF135* that can be comparable to the role *TRIM25* [40]; however, this requires further investigations. *RIG-I-RNF135*-expressing chickens also exhibited an acute upregulation of *early growth response 1* (*EGR1*). This transcription factor participates in several functions, including cell death and the induction of a large repertoire of inflammatory genes [41]. This was supported by the histological assessment that revealed the presence of multiple necrotic cells in the caecum of *RIG-I-RNF135*-expressing chickens. Overall, the increased death in *RIG-I-RNF-135*-expressing chickens, which was beneficial in terms of antiviral effect, was deleterious for the birds due to increased gene regulation and excessive immune response. Furthermore, the assessment of the immunophenotype of *RNF135*-expressing chickens revealed that this gene, on its own, may function independently from the *RIG-I* [42].

Our data suggest that the evolutionary loss of *RIG-I* in chickens and other galliform birds was advantageous in coping with viral infections caused by AIV or other avian RNA viruses. This subsequently helped decrease the acute inflammation and the damage to the host. A comparable hypothesis was made in the case of pangolins that lost the *MDA5* as an evolutionary mechanism to cope with coronavirus-induced inflammation [43]. Above that, we propose that the antiviral role of *RIG-I* in ducks [5] is not exclusively related to this gene. Still, it may involve the ubiquitination factor *RNF135* and possibly other unknown regulatory factors that support *RIG-I* signaling and reduce the repercussions of acute inflammation by negatively regulating the *RIG-I* signaling [44]. We provided novel information regarding the outcome of the re-introduction of *RIG-I* and its ubiquitination factor *RNF135* in the chicken genome. Future work should focus on identifying factors that can help reduce the acute inflammatory reaction in *RIG-I-RNF135*-expressing chickens while maintaining potent antiviral activity, which can lead to the generation of avian influenza virus-resistant chickens.

Materials and Methods

Cloning of the duck *RIG-I* and the duck *RIG-I* promoter

Total RNA was isolated from the spleen of the mallard duck (*Anas platyrhynchos*) using ReliaPrep™ RNA Tissue Miniprep System (Promega, USA), followed by cDNA synthesis using GoScript™ Reverse Transcription System (Promega, USA) according to the manufacturer's instructions. The duck *RIG-I* was amplified using Q5® High-Fidelity DNA Polymerase (New England Biolabs, USA) using the primers 562_RIG-I_for (5'-ATGACGGCGGACGAGAAGCGGAGC-3') and 563_RIG-I_rev (5'-CTAAAATGGTGGGTACAAGTTGGAC-3') that were previously described [45]. The PCR thermal conditions were as follows: 98 °C 30 sec, followed by 35 cycles of 98 °C 10 sec, 67 °C 20 sec, 72 °C 1:30 min, and a final extension step of 72 °C 2 min.

The entire length of the duck *RIG-I* promoter was amplified using primers 706_RIG-I_For (5'-

AGCTGATGACCTGCAAAAAGTT-3')   and   661_RIG-I_Rev   (5'-

GGCTGGGCTCTGCCGGCCG-3'), which were described elsewhere [21]. This resulted in an

amplicon of 2017 bp that was fully sequenced and aligned with the duck genome (*Anas*

platyrhynchos, NC_040075.1). The PCR was conducted following the following thermal

conditions 98 °C 30 sec, followed by 35 cycles of 98 °C 10 sec, 70 °C 20 sec, 72 °C 1:30 min,

and a final extension step of 72 °C 2 min.

Cloning of the duck *RNF135*

The genomic region containing duck *RNF135* was obtained from the GenBank contig

PEDO01000017.1 and corrected based on multiple publicly available duck RNAseq data.

Although the *RNF135* sequence was correctly predicted in many birds, we detected a partially

incomplete annotated sequence of the duck *RNF135*, specifically the missing 5' part that

contains the RING domain (previous accession number XM_013092775), while the predicted

version of the *RNF135* was made available (XM_027471415.2). The full-length sequence of

the duck *RNF135* was synthesized after codon optimization using the IDT online tool (IDT™,

USA) (Supplementary file 6). The obtained *RNF135* sequence was subsequently cloned into

the *RNF135*-expression vector driven by the duck *RNF135* promoter (Fig. 1B).

Identification of the duck *RNF135* promoter via Nano-Glo® Dual-Luciferase® Reporter
Assay

The potential duck *RNF135* promoter was obtained by amplifying 1577 bp 5' of the ATG start

codon from the gDNA of duck that was cloned into pGEM vector (Promega, USA) and

analyzed by Sanger sequencing. The PCR was done using Q5® High-Fidelity DNA Polymerase

with primers: 707_RNF_Prom_For (5'-GA GCA GAG CCA GGC AGC TAT A-3'), 708 (5'-

GGT CCT GCT CGG GGC GGA GC-3') resulting in an amplicon of 1557 bp. The PCR

thermal conditions were conducted using Q5® High-Fidelity DNA Polymerase (New England

Biolabs, USA) at two step-PCR 98 °C 30 sec, 98 °C 10 sec, 72 °C 60 sec and a final elongation

step of 72 °C 2 min.

The promoter activity of the duck *RNF135* was assessed by measuring promoter-driven

NanoLuc™ luciferase activity normalized to the luminescence of Firefly luciferase. To this

end, a total of 50.000 chicken DF-1 cells were seeded in 24 well plates and were co-transfected

24h later with a vector plasmid containing the deletion mutant and a second plasmid for the

expression of Firefly under the PGK promoter (Promega, USA). 24h after transfection, cells

were washed with PBS, trypsinized, and resuspended in 250µl culture medium. Firefly signal

was detected by mixing 80µl of the cell suspension with the same amount of ONE-Glo™ EX

Reagent, prepared by combining ONE-Glo™ EX Luciferase Assay Buffer with ONE-Glo™

EX Luciferase Assay Substrate in 1:1 ratio (Promega, USA). After measuring the signal of the

Firefly luciferase in the GloMax® 20/20 Luminometer (Promega, USA), 80 µl NanoDLR™ Stop & Glo® reagent, prepared by adding NanoDLR™ Stop & Glo® Substrate 1:100 into NanoDLR™ Stop & Glo® Buffer (Promega, USA), were added to the samples. These were incubated for 10, 30, 60, and 120 min and measured again in the GloMax® 20/20 Luminometer (Promega, USA) to detect the NanoLuc™ Luciferase. The cell-free culture medium was used as a blank control.

Determination of the Transgene Copy Number by Droplet Digital PCR

Droplet digital PCR (ddPCR) was used to select PGC clones with a single genomic transgene integration and was performed as described previously with slight modifications [46]. Briefly, 500ng of gDNA was digested with 20 units XbaI (New England Biolabs, Germany) for one h, followed by an inactivation step at 65°C for 20 min. The Taqman PCR reaction was set up using 10ng digested DNA, 2× ddPCR supermix for probes (no dUTP) final concentration 1× (Bio-Rad Laboratories, USA), a 20× target primer/FAM-labeled probe mix, and a 20× reference primer/HEX-labeled probe mix, which was followed by droplet generation using the QX200 Droplet Generator by mixing 20ul of the TaqMan PCR reaction with 70ul droplet generator oil in a DG8 Cartridge. The cycling conditions comprised a 95°C for 10min, followed by 40 cycles of 94°C for 30 sec, 59°C for 60 sec, and a final hold for 98°C for 10 min with a 2°C/s ramp rate at all steps. The copy number was determined by calculating the proportion of positive and negative droplets using a QX200 droplet reader, which was then analyzed using Quantasoft software (Bio-Rad Laboratories, USA) (Supplementary file 7). The hygromycin fluorescent labeled-probe ([5′FAM-TCGTGCACGCGGATTTCGGCTCCAA-3′] along with the primers: ddHygro_F [5′-CATATGGCGCGATTGCTGATC-3′] and ddHygro_R [5′-GTCAATGACCGCTGTTATGC-3′]). As a reference gene, we used the beta-actin probe ([5′HEX-GTGGGTGGAGGAGGCTGAGC-BHQ3′] along with the primer combination ddBeta_actin_F: [5′-CAGGATGCAGAAGGAGATCA-3′] and ddBeta_actin_R: [5′-TCCACC ACTAAGACAAAGCA- 3′]). The quantification was done using the QX100 system (Bio-Rad Laboratories, USA) (Supplementary file 7).

Stimulation of duck *RIG-I* expression in PGC-derived fibroblasts

The generated *RIG-I*-expressing PGCs were differentiated into fibroblasts (PGCFs) as previously described [47] and subsequently infected with a LPAIV H9N2 to examine the ability of *RIG-I* to detect its ligand. Briefly, 50.000 cells were seeded into 48 well plates and infected with an MOI of 0.1 for 18h before they were fixed and stained using immunofluorescence, as previously described [48]. Cells were fixed with 4% PFA and kept on ice for 10 min. Subsequently, they were washed with PBS and permeabilized with 0.5% Triton X. The FLAG-Tag was detected by using a mouse anti-FLAG antibody (Supplementary file 8) that was incubated for 1h, followed by staining with goat anti-mouse Alexa 568 (Supplementary file 8).

Slides were subsequently covered with a mounting medium that contains DAPI for staining the

nucleus and covered with coverslips (Vector Laboratories, Inc., USA). Fluorescence

microscopy was performed with a fluorescence microscope (ApoTome, Zeiss).

Generation of *RIG-I*-and *RNF135*-expressing chickens

White Leghorns (Lohmann selected White Leghorn (LSL), Lohmann-Tierzucht GmbH,

Cuxhaven, Germany) were used to generate transgenic chicken lines. Animal experiments were

approved by the government of Upper Bavaria, Germany (ROB-55.2-2532.Vet_02-18-9).

Experiments were performed according to the German Welfare Act and the European Union

Normative for Care and Use of Experimental Animals. All animals received a commercial

standard diet and water *ad libitum*. PGCs that express either *RIG-I* or *RNF135* were generated

using the DNA constructs shown in Fig. 1, as previously described [49, 50]. To ensure the stable

integration of the transgene, we used the phiC31 integrase-mediated integration [51]. Briefly, LSL

PGCs were derived from the blood of male embryonic vasculature at stages 13-15, according

to Hamburger and Hamilton [52]. They were cultured at 37°C in a 5% CO_2 environment using

modified KO-DMEM as described previously [53]. A total of $5x10^6$ cells per transgene were

washed with phosphate-buffered saline (PBS) and resuspended in 100μl Nucleofector^{TM}

Solution V (Lonza, Germany) containing the expression vector (Fig. 1B) and the integrase

expression construct. Electroporation was performed using an ECM 830 Square Wave

Electroporation System (BTX, USA), applying eight square wave pulses (350V, 100μsec).

After clonal selection using puromycin for *RIG-I* and blasticidin for *RNF135*, PGCs were

genotyped and tested for clones with one single genomic integration, which were then used to

generate the chimeric roosters. A total of 3000 cells with the desired genetic modification were

injected into the vasculature of 65h old embryos, transferred into a surrogate eggshell, and

incubated until the hatch. Upon sexual maturity, sperm was collected from chimeric roosters

for genotyping. [50, 54]. The germline-positive roosters were bred with wild-type hens to obtain

heterozygous animals (the germline transmission rate is presented in Supplementary File 9).

The examination of the inheritance of duck *RIG-I* and the genotyping of the offspring was done

via PCR using FIREPol DNA Polymerase (Solis Biodyne) using the primer combination

613_RIG-I-for (5'-CCTAGGAGAAGCATTCAAGGAG-3') and 563_RIG-I_Rev (5'-

CTAAAATGGTGGGTACAAGTTGGAC-3'). The following PCR conditions were used:

initial denaturation at 98 °C 30 sec, followed by 40 cycles of 98 °C 10 sec, 60 °C 30 sec, 72 °C

30 sec, and a final extension step of 72 °C 2 min, resulting in a fragment of 298bp. The

inheritance of the duck *RNF135* was examined using the primers combination 1121_RNF_For

(5'GCATGGGATCAACCGACAGCATC-3') and 1017_RNF_rev

(5'CCACACACCAACTTGACTCGGTC-3', using the following PCR conditions 98 °C 30

sec, followed by 40 cycles of 98 °C 10 sec, 60 °C 30 sec, 72 °C for 1 min and a final extension

step of 72 °C 2 min, resulting in an amplicon of 931bp.

Following the generation of different transgenic lines, they were monitored till sexual maturity

for possible harmful phenotypes that could been reflected in weight gain or the ability to

produce sperm or eggs. In addition, the immunophenotype of the generated birds was assessed

at 2, 5, 8, and 12 weeks after hatch by flow cytometry.

Isolation, culture, and infection of chicken embryonic fibroblasts (CEFs)

CEFs were isolated from 10-day-old (ED10) embryos according to the protocol published

elsewhere [55]. Before the isolation of CEFs, embryos were genotyped by collecting blood at

ED10 and preparing a window of 0.5 cm^2 in the eggshell that allowed access to the embryonic

vasculature. CEFs were cultured using Iscove's liquid medium containing stable glutamine

(Biochrom, Germany) that was supplemented with 5% fetal bovine serum (FBS) Superior

(Biochrom, Germany), 2% chicken serum (ThermoFisher Scientific, USA) and 1% Penicillin-

Streptomycin-Solution (Penicillin 10,000 U/ml and Streptomycin 10 mg/ml) (Biochrom.

Germany). Subsequently, CEFs were incubated at 40°C in a 5% CO2 atmosphere until

infection. The *in vitro* infection of CEFs with H9N2 was done after overnight seeding 250.000

cells/well in 6 well plates and infecting them in three independent experiments with

multiplicities of infection (MOIs) of 0.1 and 1. Supernatants were collected 24h post-infection

from the infected cells and titrated on MDCK (kindly provided by Prof. Silke Rautenschlein,

University of Veterinary Medicine, Hannover). The virus titration was done as previously

described [48].

***In ovo* infection of embryonated eggs**

559 Fertilized eggs from a crossing of *RIG-I* (+/-) and *RNF135*(+/-) were collected and incubated

560 till ED10. Eggs were blindly infected with 10^3 FFU of LPAIV H9N2, as previously described

561 [56]. 24h post-infection, the allantois fluid and muscle tissue were collected respectively for viral

562 titration and genotyping. The virus titration was done on MDCK cells as previously described

563 [48].

RT-PCR for the detection of transgene expression in different tissues

RNA from chicken organs, including lung, trachea, heart, liver, duodenum, thymus, bursa, and

brain, was isolated with Reliaprep RNA Tissue Miniprep System according to manufacturer

instructions (Promega), followed by cDNA synthesis using GoScript Reverse transcription mix

(Promega). The detection of *RIG-I*, as well as *RNF135* from various tissues, was done using

the following primers. *RIG-I*: 613_RIG-I-for (5'-CCTAGGAGAAGCATTCAAGGAG-3')

and 563_RIG-I_Rev 5'-(CTAAAATGGTGGGTACAAGTTGGAC-3'), while *RNF135* was

detected using the following primers: 898_duRNF_for (5'-

CTTGAGAGAGGTGGAGGGAGC-3') and 899_duRNF_rev (5'-

GGGCTGGTGGGAATTGTTGAGG-3'). *RIG-I* and *RNF135* PCRs produced an amplicon of

298bp and 148bp, respectively. β-actin mRNA was detected with primers Beta_actin_F (5′-

TACCACAATGTACCCTGGC-3′) and Beta_actin_R (5′-CTCGTCTTGTTTTATGCGC-3′)

[57], resulting in a 300-bp amplicon. The PCR was performed using FIREPol DNA Polymerase

(Solis Bio-dyne) according to the manufacturer's instructions. The following PCR conditions

were used: initial denaturation at 98 °C 30 sec, followed by 40 cycles of 98 °C 10 sec, 59 °C

30 sec, 72 °C 30 sec, and a final extension step of 72 °C 2 min.

Enzyme-linked immunosorbent assay (ELISA)

ELISA was done to measure the total plasma IgM and IgY concentrations. Briefly, 96 well

plates were coated overnight with anti-chicken antibodies IgM and IgY at a concentration of

2µg/mL (Supplementary file 8). The next day, plates were washed three times with washing

buffer, followed by a blocking step with 4% skim milk for one hour. The prediluted plasma

samples in 1:3 serial dilution were pipetted in the plates and incubated for one hour. The

detection was done by using secondary HRP-conjugated antibodies at the concentrations

mentioned in Tabe1, which were incubated for 1h at RT. This was followed by adding

100µl/well of TMB (3,3', 5,5;-tetramethylbenzidine) substrate solution for 10 min, which was

stopped by 50µl per well of 1M sulfuric acid solution. The optical density (OD) was measured

using FluoStar Omega via the measuring filter 450nm and the reference filter 620 nm (Version

5.70 R2 BMG LABTECH, Ortenberg, Germany).

Flow Cytometry

Peripheral blood mononuclear cells (PBMCs) were isolated using Histopaque®-1077 density

gradient centrifugation (Sigma, Taufkirchen, Germany) and analyzed using flow cytometry.

Extracellular staining was carried out to detect various chicken immune cell markers including

T cell subpopulations, B cells and Monocytes (list of antibodies is mentioned in Supplementary

file 8). Briefly, $5x10^6$ cells were washed with 2% BSA diluted in PBS (FLUO-Buffer). To

determine the living cell population, cells were incubated with Fixable Viability Dye eFluor

780 (eBioscience, Thermo Fisher Scientific, USA). After washing with FLUO-Buffer, primary

antibodies (concentration shown in Table 2) were applied for 20 min. Cells were washed in

FLUO-Buffer to remove unbound antibodies and incubated with conjugated secondary

antibodies for 20 min. Subsequently, cells were again washed and analyzed using an Attune

flow cytometer (Thermo Fisher Scientific, USA). The obtained data were then analyzed with

FlowJo 10.8.1 software (FlowJo, Ashland, USA). An example of the gating strategy used in

this study is presented in supplementary file 10.

Assessment of *in vitro* T cell activation

PBMCs were isolated from 12 weeks old *RIG-I* or WT birds and resuspended in RPMI medium

supplemented with 10%FBS and 1% Penicillin/Streptomycin and distributed on a 48 well plate

with a total of 5×10^6 cells/well. Cells were subsequently stimulated with Concanavalin A (Con

A) (eBioscience™) at a concentration of 25 μg per well. They were monitored every 24h for T

cell activation via flow cytometry. Activated T cells were stained for surface expression of

γδTCR1, CD25, or αβTCR2&3 using antibodies listed in Supplementary file 8.

Challenge infection experiment with H7N1

The challenge infection experiment was performed at the INRAE-PFIE platform (INRAE

Centre Val de Loire, Nouzilly, France) and approved by the local Ethics Committee Val de

Loire and the Ministère de l'Enseignement Supérieur et de la Recherche under the number

2021120115599580. Transgenic birds required for the experiment were hatched in dedicated

hatching isolators and genotyped by PCR using DNA extracted from EDTA blood samples

collected at day 7 post hatch. At 3 weeks of age, transgenic and WT-chickens were distributed

into four different groups (Supplementary file 11) corresponding to four BSL-3 isolator units.

Two other groups were kept together as MOCK controls in a single isolator (Supplementary

file 11). Throughout the experiment, the birds were fed a commercial standard diet and provided

with water *ad libitum*. Prior to infection (0 dpi), body weights were recorded, and blood samples

were taken by occipital sinus puncture. The virus used for challenge infection was A/

Turkey/Italy/977/1999 (kindly provided by Dr. Ilaria Capua, Istituto Zooprofilattico

Sperimentale Delle Venezie, Legnaro, Italy), an H7N1 subtype virus that, despite its

classification as an LPAIV, causes up to 50% mortality in experimentally infected White

Leghorn chickens [23, 58, 59].

Chickens were PBS/mock-treated (groups 5 and 6) or virus-infected (groups 1-4) by intra-

tracheal and intra-choanal cleft inoculation of 0.1mL PBS or 5×10^5 EID$_{50}$/1×10^6 EID$_{50}$ H7N1.

Animal behavior and clinical disease signs were monitored twice daily during the trial. Clinical

signs were evaluated according to the following score: 0 (no clinical signs), 1 (mild clinical

signs), 2 (severe clinical signs), or 3 (dead/euthanized). Birds were euthanized by pentobarbital

injection into the occipital sinus at the end of the experiment or once humane endpoints were

reached. Samples collected from euthanized birds included lung, caeca, and spleen. In addition,

a scoring system was used to evaluate macroscopic lung lesions as follows: 1 (mild, localized

edema and fibrinous exudate), 2 (moderate edema and with hemorrhage and fibrinous exudate

over ~1/4 of the lung), or 3 (severe hemorrhage and extensive edema over ~1/2 of the lung)

(Supplementary file 13). Data collected from the animal experiment were assessed at 2 and 6

dpi, which also represent the time points were the Fluidigm assay was performed. Details

regarding the number of birds, group distribution and number of analyzed samples are reported

in supplementary files 11 and 12.

Quantification of the viral genome in H7N1-challenged chickens

Total RNA was isolated from the isolated organs and conserved in 1ml NucleoProtect RNA (Macherey-Nagel, Germany). Samples were later processed for total RNA isolation with NucleoSpin RNA (Macherey-Nagel, Germany) according to the manufacturer's instructions. The viral genome of the H7N1 virus was quantified using qRT-PCR that was conducted with the Bio-Rad iTaq™ Universal SYBR ® Green One-Step Kit (BioRad, California, USA) using primers designed for the detection of the M gene as published elsewhere [60].

Histology

Caeca and lungs were fixed in 10% neutral buffered formalin and processed routinely. Four-micrometer tissue sections were stained with hematoxylin and eosin for light microscopy. The sections' analysis addressed histopathological changes and signs of inflammatory reactions, including degeneration, necrosis, infiltration of inflammatory cells, fibrin exudation, and epithelial hyperplasia.

Analysis of gene expression via Fluidigm Dynamic Array

Gene expression was analyzed as previously described [24]. Briefly, total RNA was extracted from infected animals' lungs, caeca, and spleen and processed for quality control via nanodrop. Reverse transcription was performed using the High Capacity Reverse Transcription Kit (Applied Biosystems) according to manufacturer's instructions with random hexamers and oligo (dT)18 in a final volume of 10μl, containing 250 ng total RNA. Subsequently, the cDNA was pre-amplified using TaqMan PreAmp Master Mix (Applied Biosystems). Quantitative PCR was performed in the BioMark HD instrument with the 96.96 IFC Dynamic Array (Fluidigm). The reaction was prepared by mixing 2.5 μl TaqMan Gene Expression Master Mix (Applied Biosystems), 0.25 μl 20X DNA Binding Dye Sample Loading Reagent (Fluidigm), 0.25 μl 20X EvaGreen DNA binding dye (Biotum) and 2 μl of preamplified cDNA. The qPCR was run under the following thermal conditions: 50°C for 2 min, 70°C for 30 min, 25°C for 10 min, followed by hot start 50°C for 2 min, 95°C for 10 min, PCR (x30 cycles) 95°C for 15 sec, 60°C for 60 sec and melting curve analysis 60°C for 3 sec to 95°C.

Raw quantitation cycle (Cq) data were collated with the Real-time PCR Analysis software v3.1.3 (Fluidigm), setting parameters of quality Cq threshold to auto (global) and the baseline correction to derivative. Raw Cq values were processed with GenEx.v6 MultiD, with correction for primer efficiency and reference gene normalization. Stability of the expression of reference genes: *TATA box binding protein (TBP), Tubulin alpha chain (TUBA8B), beta-actin (ACTB), beta-glucuronidase (GUSB), glyceraldehyde-3-phosphate dehydrogenase (GAPDH) and ribosomal 28S (r28S)* were evaluated via NormFinder (GenEx). The geometric mean of the most stable (*GAPDH*, *GUSB* and *TBP*) were used for normalization. Technical replicates were

680 averaged, and relative quantification was to the maximum Cq value obtained per gene,
681 transformed to logarithmic scale.

Statistical analysis

Statistical analysis was done using IBM SPSS Statistics (Version 28.0.1.1. IBM, Armonk,
USA). The normality of the data was examined via Kolmogorov-Smirnov and Shapiro-Wilk
tests. The comparison between two groups was made either with Two Samples T-test or
Wilcoxon Signed Rank test. Multiple group comparison was made using Kruskal-Wallis Test
or with One-Way ANOVA. The statistical test result was considered significant when the P
value was less than 0.05.

Acknowledgments

Animal husbandry and experimental infrastructure was provided by the TUM Animal Research
Center (ARC). The authors thank the animal keepers of the ARC as well as the team at the
INRAE Plateforme d'Infectiologie Expérimentale (INRAE-PFIE, Nouzilly, France) for
conducting the challenge infection experiment. In addition, we would like to thank Leah
Heymelot Nabel for her help with the cloning of *RIG-I* and *RNF135*. Additionally, the authors
thank the technical staff of the involved laboratories for excellent assistance.

Funding

The project was financed by a grant from the German Research Foundation to HS (DFG SI
2478/2-1). B. Schusser and TvH were supported by a grant from the German Research
Foundation to BS (DFG SCHU2446/6-1). In addition, the challenge infection experiment was
supported by the European infrastructure project VetBioNet (EU Horizon 2020, grant
agreement No 731014). LV was supported by the Biotechnology and Biological Sciences
Research Council Institute Strategic Program Grant funding (BBS/E/D/30002276).

709
Fig. 1 Generation of PGCs, and susceptibility to infection using *in vitro* and *in vivo*
systems. A. Promoter activity of the duck *RNF135* was examined by generating different
deletion mutants tested in chicken DF-1 cells; the promoter activity was then evaluated by
measuring the NanoLuc/Firefly ratio (n=3). B. Diagram of both constructs used to generate
RIG-I-expressing chicken (upper diagram), using the previously identified duck *RIG-I*
promoter [21] and *RNF135*-expressing chickens (lower diagram), using the duck *RNF135*

promoter, which activity was identified in this study. C. Primordial germ cells (PGCs) that

express the duck *RIG-I* under the duck *RIG-I* promoter were derived into PGC fibroblasts

(PGCFs) and were infected later with avian influenza virus H9N2 to stimulate *RIG-I* expression

upon influenza infection; the cells were infected for 18h with low pathogenic avian influenza

virus H9N2 and subsequently stained for FLAG-Tag; MOCK control represents uninfected

cells stained for anti-FLAG-Tag; DF-1 cells that express the FLAG-tagged duck *RIG-I* under

the chicken beta-actin promoter were used as a positive control. D. Quantification of newly

produced viral particles after infection of CEFs; CEFs were isolated from different transgenic

embryos and experimentally infected with LPAIV H9N2 at two different multiplicities of

infection (MOI 0.1, and 1); Supernatants were collected at 24hpi and titrated on MDCK cells;

no significant differences were observed between the groups ($p>0.05$). E. Quantification of

newly produced viral particles after infection of embryonated eggs. 10 day-old embryonated

eggs were infected with LPAIV H9N2 at 10^3 FFU/egg; allantois fluid was collected 24hpi and

titrated on MDCK cells; no significant differences were observed between the groups ($p>0.05$).

Fig. 2 Generation and immunophenotypic characterization of *RIG-I* and *RNF135*-expressing chickens. A. Weekly weight monitoring of the generated heterozygous birds ($p>0.05$) (n=10). B. Representative picture of the generated heterozygous birds from left to right at four weeks of age: WT, *RNF135*-expressing chicken, and *RIG-I*-expressing chicken. C. RT-PCR of the transgenic expression of *RIG-I* and *RNF135* in different organs. D. Assessment of different immune cell populations in *RIG-I*-expressing chickens compared to their WT siblings.

Fig. 3 Assessment of different T cell subpopulations in *RIG-I*-expressing chickens compared to their WT siblings

A) PBMCS were analyzed for γδTCR1+/CD8α+ T cells at 12 weeks. B) PBMCS were analyzed for αβTCR2,3+T cells and CD4+ or CD8α+T cells at 12 weeks ($p<0.05$). C) Activation of isolated T cells from 12-week-old *RIG-I*-expressing chickens compared to their WT siblings ($p<0.05$); cells were sorted according to their TCR expression, stimulated with Concanavalin

751   A, and quantified by flow cytometry at different time-points. The results indicate the number

752   of positive cells per 100 viable TCR + cells.

Fig. 4 Simultaneous expression of *RIG-I* and *RNF135* in the chicken does not lead to

differences in the adaptive immune phenotype compared to WT birds

Two-time points, eight and twelve weeks of age, were chosen based on the data obtained from

RIG-I-expressing chickens to assess the immunophenotype of *RIG-I-RNF135*-expressing

chickens. PBMCS were isolated and analyzed for B cells, αβTCR2,3+ or γδTCR1 + T cells (A)

as well as for CD4+ (B) and CDα8+ T cells (C). No significant differences were detected

between the analyzed groups ($p > 0.05$).

Fig. 5 H7N1-challenge experiment reveals the susceptibility of *RIG-I*-expressing chickens

and the role of *RNF135* in effective *RIG-I* antiviral response.

The generated transgenic chickens were challenged with H7N1 and assessed at two days post-

infection (dpi) and six dpi for different parameters. A) Percentage of animals presenting clinical

symptoms. B) Probability of survival in the challenged groups. C) Weight gain in challenged

groups compared to WT- and *RIG-I-RNF135*-MOCK- controls ($p<0.05$). D) Viral replication

rate in two main target organs, caecum, and lung ($p<0.05$); the horizontal line indicates the

detection threshold of the PCR based on the signal obtained from the uninfected controls, which

is $\log_{10}(10^3)$. E) Expression levels of transgenes in the caecum and the lung upon H7N1-

challenge ($p<0.05$).

785
786
Fig. 6 Developed lesions in H7N1-challenged birds A) Macroscopical lesion score of the

lungs showing significant increase of lesions in *RIG-I*-expressing birds at six dpi ($p<0.05$) B)

Histology of the caecum showing the typical structure of the epithelium in the control groups.

RIG-I-RNF135 challenged birds showing epithelial hyperplasia with necrosis that diminished

by 6 dpi. *RIG-I*-expressing chickens did not show necrotic lesions but pronounced epithelial

hyperplasia that lessened by six dpi. Asterix indicates epithelial hyperplasia with the mitotic

figures, while the arrow indicates necrotic epithelial cells; 200x.

Fig. 7 Infection of transgenic chickens with H7N1 leads to upregulated inflammatory genes. Venn diagram of fluidigm qRT-PCR array of naive and challenged birds with H7N1 showing significantly upregulated genes in various organs. Significant DEGs were identified by comparing the relative expression

values for every chicken line to the WT-MOCK individually per timepoint, with a significance level set at $p < 0.05$; fold change >1 (n ≥ 5-time point).

Fig. 8 Gene expression in the lung upon H7N1-infection

Significant DEGs were identified by via fluidigm qPCR array and comparison of the relative expression values between H7N1-challenged groups. A) Significantly upregulated genes after comparison of infected *RIG-I-* with *RIG-I-RNF135*-expressing chickens at two dpi. B) Significantly upregulated genes after comparison of infected *RIG-I*-with *RIG-I-RNF135*-expressing chickens at 6dpi. C) Significantly upregulated genes after comparison of infected *RIG-I-* with *RIG-I-RNF135*-expressing chickens at two dpi. D) Significantly upregulated genes after comparison of infected *RIG-I*-expressing with chickens WT-chickens at six dpi. Significance levels were set at $p < 0.05$; fold change >1 (n ≥ 5-time point).

References

819 1. Mostafa, A., Abdelwhab, E.M., Mettenleiter, T.C. & Pleschka, S. Zoonotic potential
820   of influenza A viruses: a comprehensive overview. *Viruses* **10**, 497 (2018).
821
822 2. World Health Organization. Ongoing avian influenza outbreaks in animals pose risk to
823   humans [cited 2023 July 12]. https://www.who.int/news/item/12-07-2023-ongoing-
824   avian-influenza-outbreaks-in-animals-pose-risk-to-humans.
825
826 3. MacLachlan, N. Orthomyxoviridae. *Fenner's veterinary virology*, 353-370 (2011).
827
828 4. Kim, J.K., Negovetich, N.J., Forrest, H.L. & Webster, R.G. Ducks: the "Trojan
829   horses" of H5N1 influenza. *Influenza and other respiratory viruses* **3**, 121-128 (2009).
830
831 5. Barber, M.R., Aldridge Jr, J.R., Webster, R.G. & Magor, K.E. Association of RIG-I
832   with innate immunity of ducks to influenza. *Proceedings of the National Academy of
833   Sciences* **107**, 5913-5918 (2010).
834
835 6. Rehwinkel, J. *et al.* RIG-I detects viral genomic RNA during negative-strand RNA
836   virus infection. *Cell* **140**, 397-408 (2010).
837
838 7. Kowalinski, E. *et al.* Structural basis for the activation of innate immune pattern-
839   recognition receptor RIG-I by viral RNA. *Cell* **147**, 423-435 (2011).
840
841 8. Yoneyama, M. *et al.* The RNA helicase RIG-I has an essential function in double-
842   stranded RNA-induced innate antiviral responses. *Nature immunology* **5**, 730-737
843   (2004).
844
845 9. Gack, M.U. *et al.* TRIM25 RING-finger E3 ubiquitin ligase is essential for RIG-I-
846   mediated antiviral activity. *Nature* **446**, 916-920 (2007).
847
848 10. Gao, D. *et al.* REUL is a novel E3 ubiquitin ligase and stimulator of retinoic-acid-
849   inducible gene-I. *PloS one* **4**, e5760 (2009).
850
851 11. Oshiumi, H. *et al.* The ubiquitin ligase Riplet is essential for RIG-I-dependent innate
852   immune responses to RNA virus infection. *Cell host & microbe* **8**, 496-509 (2010).
853
854 12. Wang, Y. *et al.* Rig-I−/− mice develop colitis associated with downregulation of Gαi2.
855   *Cell research* **17**, 858-868 (2007).
856
857 13. Pang, I.K., Pillai, P.S. & Iwasaki, A. Efficient influenza A virus replication in the
858   respiratory tract requires signals from TLR7 and RIG-I. *Proceedings of the National
859   Academy of Sciences* **110**, 13910-13915 (2013).
860
861 14. Cagliani, R. *et al.* RIG-I-like receptors evolved adaptively in mammals, with parallel
862   evolution at LGP2 and RIG-I. *Journal of molecular biology* **426**, 1351-1365 (2014).
863
864 15. Kandasamy, M. *et al.* RIG-I signaling is critical for efficient polyfunctional T cell
865   responses during influenza virus infection. *PLoS pathogens* **12**, e1005754 (2016).
866
867 16. Vanderven, H.A. *et al.* Avian influenza rapidly induces antiviral genes in duck lung
868   and intestine. *Molecular immunology* **51**, 316-324 (2012).
869

870   17.   Barber, M.R., Aldridge, J.R., Jr., Webster, R.G. & Magor, K.E. Association of RIG-I
871   with innate immunity of ducks to influenza. *Proceedings of the National Academy of*
872   *Sciences of the United States of America* **107**, 5913-5918 (2010).
873

874   18.   Zheng, W. & Satta, Y. Functional evolution of avian RIG-I-like receptors. *Genes* **9**,
875   456 (2018).
876

877   19.   Krchlíková, V. *et al.* Dynamic Evolution of Avian RNA Virus Sensors: Repeated Loss
878   of RIG-I and RIPLET. *Viruses* **15**, 3 (2022).
879

880   20.   Hayman, T.J. *et al.* RIPLET, and not TRIM25, is required for endogenous RIG-I-
881   dependent antiviral responses. *Immunology and cell biology* **97**, 840-852 (2019).
882

883   21.   Xiao, Y., Reeves, M.B., Caulfield, A.F., Evseev, D. & Magor, K.E. The core promoter
884   controls basal and inducible expression of duck retinoic acid inducible gene-I (RIG-I).
885   *Molecular immunology* **103**, 156-165 (2018).
886

887   22.   Cadena, C. *et al.* Ubiquitin-dependent and-independent roles of E3 ligase RIPLET in
888   innate immunity. *Cell* **177**, 1187-1200. e1116 (2019).
889

890   23.   Dundon, W.G., Milani, A., Cattoli, G. & Capua, I. Progressive truncation of the Non-
891   Structural 1 gene of H7N1 avian influenza viruses following extensive circulation in
892   poultry. *Virus research* **119**, 171-176 (2006).
893

894   24.   Bryson, K.J. *et al.* Comparative Analysis of Different Inbred Chicken Lines
895   Highlights How a Hereditary Inflammatory State Affects Susceptibility to Avian
896   Influenza Virus. *Viruses* **15**, 591 (2023).
897

898   25.   Kato, H. *et al.* Differential roles of MDA5 and RIG-I helicases in the recognition of
899   RNA viruses. *Nature* **441**, 101-105 (2006).
900

901   26.   Xu, L., Yu, D., Fan, Y., Liu, Y.-P. & Yao, Y.-G. Evolutionary selection on MDA5 and
902   LGP2 in the chicken preserves antiviral competence in the absence of RIG-I. *Journal*
903   *of Genetics and Genomics= Yi Chuan xue bao* **46**, 499-503 (2019).
904

905   27.   Evseev, D., Miranzo-Navarro, D., Fleming-Canepa, X., Webster, R.G. & Magor, K.E.
906   Avian Influenza NS1 Proteins Inhibit Human, but Not Duck, RIG-I Ubiquitination and
907   Interferon Signaling. *Journal of Virology* **96**, e00776-00722 (2022).
908

909   28.   Jiang, X. *et al.* Intrinsic RIG-I restrains STAT5 activation to modulate antitumor
910   activity of CD8+ T cells. *The Journal of Clinical Investigation* **133** (2023).
911

912   29.   Kandasamy, M. *et al.* RIG-I Signaling Is Critical for Efficient Polyfunctional T Cell
913   Responses during Influenza Virus Infection. *PLoS pathogens* **12**, e1005754 (2016).
914

915   30.   Iwamoto, A., Tsukamoto, H., Nakayama, H. & Oshiumi, H. E3 Ubiquitin Ligase
916   Riplet Is Expressed in T Cells and Suppresses T Cell–Mediated Antitumor Immune
917   Responses. *The Journal of Immunology* **208**, 2067-2076 (2022).
918

919   31.   Elsheimer-Matulova, M. *et al.* Interleukin 4 inducible 1 gene (IL4I1) is induced in
920   chicken phagocytes by Salmonella Enteritidis infection. *Veterinary Research* **51**, 1-8
921   (2020).
922

923   32.   Nish, S.A. *et al.* T cell-intrinsic role of IL-6 signaling in primary and memory
924   responses. *elife* **3**, e01949 (2014).
925

926   33.   Thoresen, D.T., Galls, D., Götte, B., Wang, W. & Pyle, A.M. A rapid RIG-I signaling
927   relay mediates efficient antiviral response. *Molecular cell* **83**, 90-104. e104 (2023).
928

929   34.   Cornelissen, J., Post, J., Peeters, B., Vervelde, L. & Rebel, J. Differential innate
930   responses of chickens and ducks to low-pathogenic avian influenza. *Avian Pathology*
931   **41**, 519-529 (2012).
932

933   35.   Liau, N.P. *et al.* The molecular basis of JAK/STAT inhibition by SOCS1. *Nature*
934   *communications* **9**, 1558 (2018).
935

936   36.   Palmer, D.C. & Restifo, N.P. Suppressors of cytokine signaling (SOCS) in T cell
937   differentiation, maturation, and function. *Trends in immunology* **30**, 592-602 (2009).
938

939   37.   Schmit, T. *et al.* Interferon-γ promotes monocyte-mediated lung injury during
940   influenza infection. *Cell reports* **38** (2022).
941

942   38.   Koliopoulos, M.G. *et al.* Molecular mechanism of influenza A NS1-mediated TRIM25
943   recognition and inhibition. *Nature communications* **9**, 1820 (2018).
944

945   39.   Meyerson, N.R. *et al.* Nuclear TRIM25 specifically targets influenza virus
946   ribonucleoproteins to block the onset of RNA chain elongation. *Cell host & microbe*
947   **22**, 627-638. e627 (2017).
948

949   40.   Okamoto, M., Kouwaki, T., Fukushima, Y. & Oshiumi, H. Regulation of RIG-I
950   activation by K63-linked polyubiquitination. *Frontiers in immunology* **8**, 1942 (2018).
951

952   41.   Woodson, C.M. & Kehn-Hall, K. Examining the role of EGR1 during viral infections.
953   *Frontiers in Microbiology* **13**, 1020220 (2022).
954

955   42.   Vazquez, C., Tan, C.Y. & Horner, S.M. Hepatitis C virus infection is inhibited by a
956   noncanonical antiviral signaling pathway targeted by NS3-NS4A. *Journal of Virology*
957   **93**, e00725-00719 (2019).
958

959   43.   Fischer, H., Tschachler, E. & Eckhart, L. Pangolins lack IFIH1/MDA5, a cytoplasmic
960   RNA sensor that initiates innate immune defense upon coronavirus infection.
961   *Frontiers in immunology* **11**, 939 (2020).
962

963   44.   Oshiumi, H., Matsumoto, M. & Seya, T. Ubiquitin-mediated modulation of the
964   cytoplasmic viral RNA sensor RIG-I. *The journal of biochemistry* **151**, 5-11 (2012).
965

966   45.   Chen, Y. *et al.* Duck RIG-I CARD domain induces the chicken IFN-β by activating
967   NF-κB. *Biomed Research International* **2015** (2015).
968

969   46.   Rieblinger, B. *et al.* Strong xenoprotective function by single-copy transgenes placed
970   sequentially at a permissive locus. *Xenotransplantation* **25**, e12382 (2018).
971

972   47.   Long, J.S. *et al.* Species specific differences in use of ANP32 proteins by influenza A
973   virus. *Elife* **8**, e45066 (2019).
974

48.   Sid, H., Hartmann, S., Winter, C. & Rautenschlein, S. Interaction of influenza A viruses with oviduct explants of different avian species. *Frontiers in Microbiology* **8**, 1338 (2017).

49.   Van de Lavoir, M.-C. *et al.* Interspecific germline transmission of cultured primordial germ cells. *PLoS One* **7**, e35664 (2012).

50.   Van de Lavoir, M.-C. *et al.* Germline transmission of genetically modified primordial germ cells. *Nature* **441**, 766-769 (2006).

51.   Leighton, P.A., van de Lavoir, M.C., Diamond, J.H., Xia, C. & Etches, R.J. Genetic modification of primordial germ cells by gene trapping, gene targeting, and φC31 integrase. *Molecular Reproduction and Development: Incorporating Gamete Research* **75**, 1163-1175 (2008).

52.   Hamburger, V. & Hamilton, H.L. A series of normal stages in the development of the chick embryo. *Journal of morphology* **88**, 49-92 (1951).

53.   Whyte, J. *et al.* FGF, insulin, and SMAD signaling cooperate for avian primordial germ cell self-renewal. *Stem cell reports* **5**, 1171-1182 (2015).

54.   Schusser, B. *et al.* Immunoglobulin knockout chickens via efficient homologous recombination in primordial germ cells. *Proceedings of the National Academy of Sciences* **110**, 20170-20175 (2013).

55.   Hernandez, R. & Brown, D.T. Growth and maintenance of chick embryo fibroblasts (CEF). *Current protocols in microbiology* **17**, A. 4I. 1-A. 4I. 8 (2010).

56.   Brauer, R. & Chen, P. Influenza virus propagation in embryonated chicken eggs. *JoVE (Journal of Visualized Experiments)*, e52421 (2015).

57.   Kong, F.-k., Chen, C.-l.H., Six, A., Hockett, R.D. & Cooper, M.D. T cell receptor gene deletion circles identify recent thymic emigrants in the peripheral T cell pool. *Proceedings of the National Academy of Sciences* **96**, 1536-1540 (1999).

58.   Monne, I. *et al.* Emergence of a highly pathogenic avian influenza virus from a low-pathogenic progenitor. *Journal of virology* **88**, 4375-4388 (2014).

59.   Trapp, S. *et al.* Major contribution of the RNA-binding domain of NS1 in the pathogenicity and replication potential of an avian H7N1 influenza virus in chickens. *Virology journal* **15**, 1-12 (2018).

60.   Ward, C. *et al.* Design and performance testing of quantitative real time PCR assays for influenza A and B viral load measurement. *Journal of clinical virology* **29**, 179-188 (2004).

 Supplementary Data

 Supplementary file 1: Immunophenotype of *RNF135*-expressing chickens

1030

1031

1032

1033

1034

1035

1036

1037

1038

1039

1040

1041

Supplementary file 2: Immunoglobulin levels in *RIG-I*-expressing birds compared to WT

Relative amounts of total plasma IgM and IgY in *RIG-I*-expressing birds and their WT siblings compared by ELISA at 12 weeks of age (n≥3)

1068 **Supplementary file 3: Body weight development over 14 weeks of *RIG-I-RNF135-***
1069 **expressing chickens in comparison to WT siblings (n ≥4)**
1070

1071

Supplementary file 4: Fluidigm qRT-PCR array of naive and challenged birds with H7N1
A) Principal component analysis (PCA) of global gene expression of viral innate immune-related genes. B) *RIG-I-RNF135* MOCK compared to WT-MOCK birds; positive values indicate high expression in the *RIG-I-RNF135* expressing chickens, while negative values indicate high expression in WT-MOCK. C) Expressed genes in the caecum of infected groups compared to WT-MOCK at 2dpi; positive values indicate high expression in the infected groups (1=WT-H7N1, 2=*RIG-I-RNF135*-H7N1, 3=*RIG-I*-H7N1, 4=*RNF135*-H7N1), while negative values indicate high expression in WT-MOCK. Significant DEGs were identified by comparing the relative expression values for every chicken line to the WT-MOCK individually per timepoint, with a significance level set at *p* < 0.05; fold change >1 (n ≥5/timepoint)

Supplementary file 5: Involvement of the regulated genes in the biological processes. Each
category is represented by a different color (Biological Process = red , Cellular Component =
blue and Molecular Function = green). A) regulated genes in the caecum and spleen of *RIG-I-*
RNF135-expresing birds. The height of the bar represents the number of IDs in the uploaded
list of genes and the category. B) Function of genes expressed in the lung and caecum of *RIG-*
I-expressing chickens C) Function of genes expressed in the spleen of *RIG-I*-expressing
chickens at 6dpi. Figures were generated using WebGestalt (WEB-based GEne SeT AnaLysis
Toolkit).

1120

Supplementary file 6: Sequence of duck *RNF135* promoter and the duck *RNF135*

Duck RNF135 promoter (*Anas platyrhynchos*)
GGAGCAGAGCCAGGCAGCTATAGTCTACCTGACTGCAGCTCCTATCCAGACAGGGACATGTTTCTTTTCCATGGAGTGGTTTTAATGCCTTTTACAGTGTTAGATTTCTG
TCACAGAAATTGTAAAAAAAATAAAAGTCAAGTTTGAATTGTGGAGAATACTGTAATGGAATCACTCTAATGACAGATGTCCCTACCTCCTTGCTTTCAACAGTCTCTGCTA
TTTGATTCTTTTATGAGTGGCAAATGAGACATGTAAAGACTTTTTTTTTCTTTTCCAAAGCAACCCCCACATACTGCCTCAGTCAAGACGAGTTCATGATGAAGACACACTG
ATTGGTATTACCTCCTAAAAACAGCTAAAGCTTTGGGTCATGAGTTCACTTACTGAATTTGAGTAGTCTGAGCAGTGAAAGGCAGAAGTCAGTTTTAAAAGTGCTGTCTTC
CTTGGCTTGTTTGAAGACACTCAATGTATGAGGACCAGCACGGAAAGCACAGAGTGAGTGCCTCCTGATCAGGTGCTACGTACATCCATTCTGCGCAGGAATGGGTGA
GGATTTGCACTGGCTCACAAAGGGCACTGAGTGTTTCTTGCCTCACAGCCAAGCCCAAGATTTGGTTCTACCAAGAAGTCAGTCATTCAGATATTTAATTCTTCTTTTCA
GATTCCCTTAGTTTGCAAATCAAGCGGAAGTCTTGCAAACTAAGCTTCTGGTAAAGCTTCCACCTTTTGCTGTTGCTTGCCAGCCAGTCTAAGGCACCCCAAATAGTTCTA
CAGAAAGAAGGAAGTACCAATATATGCAGGGATTAGTGTGATTAACTCTTTTTTTCCCACAACAGTACAGAAAATTTGACTTGGTTCATCATTCTGTCTGTCTTAGTCAACT
CATCATCTTCCAGCCTTGAACAAACTGCAGGAGTTTTGCTTCCTGACATTCTCCCATAAACTCCCTGATGTTTCTGTTCTTTCTTTCCTCTCCCCATACGATCTGGGGAATT
ACAGTTCACGTGCTGGTTCAGCAAATCTTGTAACAGCCTCATCCTGCAGCTCCTGAAATCAACAGCGAGGCACTTGTTGAAGATCATAGTCTGGTTTGAGATGCTTTTTT
TTATTATTTATTTTGAGGTGAGCTGAGCAATTCTGGATGATCTCAAAAAGTTTAGTCTACTCAGAAGCCTCAGCTTCACGGATGGGGTTCCCTGTGTGATGTCCGGGAAA
GGTCAGCAGACCAACCGTGTCAGCAACCCTGTTACAAAACCAGGAAATTCCCAGAAGAAAAAAAAAAATGCTACTGTGCGAAGACGACAACTTTTTGCAGACTGGAGGC
GAGGTTACGGAGAGCTCCGTGTGACGGTACCTTCCTACAGCCAACCCAAGTCCTGGCGCCGAGCTGAAGGCAGCAGCAGCACCGGAGCCCGGCTTCGCGGGTGCCC
CGGGCCTTCAGCTGCCGGTAGCCCCGCACCGGGCCCGGGTCCAGCACCGAGCCCAGCCCCACACCGGGCCCGGCCCCCCCGGCGGGAGGCGGCGCCCCGAGGAAA
CGAAACCGCGGCCGCGGCTCCGCTCCGCCCCGAGCAGGACCT

GC-optimised duck RNF135 sequence (*Anas platyrhynchos*)
ATGTATCCTTATGATGTTCCCGATTATGCTGCGGTTCCCGCTGAGCTTGGCCGGCTCCTCGCTGACGTTGAACTCAGCTGTTCATGCTGTTTGCAGTACTTTACCGAGCC
CGTACGGTTGGCGTCTTGCTCTCACAGCTTCTGTCGGAGCTGTATCGACACGTACTGTCGGGGTCGCCGGCGGGCACCCTGCCCTCTCTGCAGAGAGGATTTTGAGC
CGAAAGATTTGCGCCCGAATAGAGAACTTGCAGCCCTTGTCTCACTCATGTTGGGCGGAGGCAGAGGTGAGGGACTGGGTGCATGGGATCAACCGACAGCATCTGGT
GACGGGGCGGGCGGAGGGTGCAGCAGTGCCTGGAGACGCCCGGGGGAAAAGGAGGAGCAAATACGGGATATTAGCAAACAGCTTGAGATAACTAAGGAAACTATTA
ATGCGCTGCGGAAAGATTTGTCTAAGACGGAAAGAATATACAAGCCAAATACTCTCACAAATAACAGAGGACTTTTGCTGCATGAAAGAGTATATTGAGAGGCAAGAAGAA
AATACTTTGATGTTCATCGAACAGGAACAGAGGGCGGCCCGGCAAAAGATAGTACAAACTATCCACCAGCTTTGTGTTGAAAAATACAAATTGATAGACATAAAGGCGCA
AATGGAGAAGGGGTTGGAAAGTGACGAAATGGAATGGCAGACCTCTAACCTTCTTGAGAGAGGTGGAGGGAGCCCTAGTACCATGCATAAGTTCACAATCGATGAGAA
ATTCAACGTCGTTAGGAGCGCCGTAGGAGATCTGAAAAGAAAATTGGAGATCTTGCTGCTCGAAGAGTATCCTCAACAATTCCCACCAGCCCAAAGTCCCGACTTGCAT
CAGGAAACGAGTGTCTGTTCATTGTCATCCGAAAGCGCGGCTAAGTCACCCGAACCGTCTATCTCATCCCAATTTTCACGCTGGGCTGATAACGTCACCTTTGACTTGAC
GACCGCCTATGATAGACTTGCCATCACCGATCAGAATAGAAAAGTTATGGTGTCATCTAACCCAACCTATTACGAGCCATCCCTTAAACGGTTCTGTATAAGTCAGGTTCT
TTGCAGCCAAGGCTTCAGTACCGGGTGCCACTATTGGGAGGTAATCACCAAGGACAGCGACGGTTGGGCGGTTGGCGTTGCCAGAGGAACCATCGGGAGACGGGAT
CGCTTGGGACGGACCGAGTCAAGTTGGTGTGTGGAGTGGGTGGGACCTCAGAAGCAGCTGAGCGCCTGGCATCGCAATCAAGAGACACTGCTGCGGAACGACAAGC
CACTTAAAGTAGGCGTGTTCCTTGAACTGCAAAAGACCGTATCTTTTTATGCGATAACCGATCGCGAGATGCTCTTGCATACATTCGAAATCAATAACAGTAACCCTTTGT
ACCCGGCCCTTCTGGCTCTATAGCTTGGACAAGAACGGCTCTCTTACGATAAATCATATTAACAGGAAATAAG

Protein sequence of the duck RNF135 (*Anas platyrhynchos*)
M**YPYDVPDYAA**VPAELGRLLADVELSCSCCLQYFTEPVRLASCSHSFCRSCIDTYCRGRRRAPCPLCREDFEPKDLRPNRELAALVSLMLGGGRGEGLGAWDQPTASGDGA
GGGCSSAWRRPGEKEEQIRDISKQLEITKETINALRKDLSKTKEYTSQILSQITEDFCCMKEYIERQEENTLMFIEQEQRAARQKIVQTIHQLCVEKYKLIDIKAQMEKGLESDEME
WQTSNLLERGGGSPSTMHKFTIDEKFNVVRSAVGDLKRKLEILLLEEYPQQFPPAQSPDLHQETSVCSLSSESAAKSPEPSISSQFSRWADNVTFDLTTAYDRLAITDQNRKVM
VSSNPTYYEPSLKRFCISQVLCSQGFSTGCHYWEVITKDSDGWAVGVARGTIGRRDRLGRTESSWCVEWVGPQKQLSAWHRNQETLLRNDKPLKVGVFLELQKTVSFYAITD
REMLLHTFEINNSNPLYPAFWLYSLDKNGSLTINHINRK*

Blue colored letters indicate the inserted HA tag

1122

1123

1124

1125

1126

1127

1128

1129

1130

1131

1132

1133

1134

1135

1136

1137

1138

1139

1140

1141

1142

1143

1144

1145

1146

1147

Supplementary file 7: Results of Droplet Digital PCR (ddPCR) of PGCs that express the
duck *RIG-I* (A) and those who express the duck *RNF135* (B)
A) #253 indicates the construct with the full *RIG-I* length promoter used to express the duck
RNF135 in PGCs. Clone #1 was used for the generation of *RIG-I* chimeric roosters.
B) #258 indicates the construct number used to express the duck *RNF135* in PGCs. Clone #6
was used for the generation of *RNF135* chimeric roosters.
C) Selected clones were derived into fibroblasts (PGC-derived fibroblasts) and tested for the
RIG-I activity by the detection of FLAG-Tag
1156

1157

1158

1159

1160

1161

1162

1163

1164 **Supplementary file 8: clones and concentrations of used antibodies**

Antibody	Company	Clone	Assay	Concentration
Mouse Anti-Chicken TCRγδ-UNLB	Biozol	TCR1	Flow cytometry	0.625 µg/mL
mouse anti-chicken TCRαβ/Vb1-UNLB	Biozol	TCR2	Flow cytometry	2.5µg/mL
mouse anti-chicken TCRαβ/Vb2-UNLB	Biozol	TCR3	Flow cytometry	2.5µg/mL
mouse-IgG1_anti-chicken chB6_UNLAB	Biozol	AV20	Flow cytometry	2.5µg/mL
mouse IgG2a -anti-chicken KUL01_UNLAB	Biozol	KUL01	Flow cytometry	0.625 µg/mL
goat anti-mouse IgG (H+L)-APC	Biozol	Polyclonal	Flow cytometry	0.625 µg/mL
mouse-IgG1_anti-chicken CD8α_PacBlue	Biozol	CT-8	Flow cytometry	0.625 µg/mL
mouse IgG2a -anti-chicken CD8ß_UNLAB	Biozol	EP42	Flow cytometry	0.625 µg/mL
mouse IgG1-anti-chicken CD4_FITC	Biozol	CT-4	Flow cytometry	0.625 µg/mL
mouse IgG1 Anti-Chicken TCRγδ-BIOT	Biozol	TCR-1	Flow cytometry	0.625 µg/mL
mouse anti-chicken TCRαβ/Vb1-BIOT	Biozol	TCR-2	Flow cytometry	2.5µg/mL
mouse anti-chicken TCRαβ/Vb2-BIOT	Biozol	TCR-3	Flow cytometry	2.5µg/mL
mouse anti-chicken Bu1-FITC	Biozol	AV-20	Flow cytometry	2.5µg/mL
mouse anti- chicken MRC1L-B	Biozol	KUL01	Flow cytometry	0.625 µg/mL
rat-anti-mouse IgG2a_PE	Biozol	SB84a	Flow cytometry	0.125µg/mL
Streptavidin_APC	VWR	-	Flow cytometry	0.2µg/mL
goat anti-mouse IgG (H+L)-APC	Biozol	polyclonal	Flow cytometery	0.625 µg/mL
Human anti-Chicken CD25	Biorad	monoclonal	Flow cytometery (T cell activation)	2.5µg/mL
Fixable Viability Dye eFluor 780	eBioscience		Flow cytometery	1:1000
Goat anti-chicken IgM	Biomol	polyclonal	ELISA	2µg/mL
Rabbit anti-chicken IgY (used for coating ELISA plates)	Jackson	polyclonal	ELISA	2µg/mL
Goat anti-chicken IgM-HRP	Biomol	polyclonal	ELISA	0.05ng/mL
Rabbit Anti-Chicken IgY-HRP (used for detection)	Jackson	polyclonal	ELISA	0.02µg/mL

1165

1166   **Supplementary file 9: Germline transmission of the injected PGC clones**

Injected PGC clone	Germline transmission (%)
RIG-I	10.2
RNF135	6

1167

1168

1169

1170

1171

1172

1173

1174

1175

1176

1177

1178

1179

1180

1181

1182

1183

1184

1185

1186

1187

1188

 Supplementary file 10 Gating strategies for the detection of different immune cells. A)
Detection of B cells and monocytes. B) detection of T cell subpopulations. C)Live/Dead
staining and detection of TCRαβ1, αβ2/CD25+ T cells and TCRγδ/CD25+ T cells

Supplementary file 11: Group distribution and number of birds used in the experimental challenge with H7N1

Genotype	RIG-I-RNF135		WT		RIG-I		RNF135		WT		RIG-I-RNF135	
Inoculation	H7N1		H7N1		H7N1		H7N1		MOCK		MOCK	
Number	(n=16)		(n=16)		(n=18)		(n=18)		(n=9)		(n=6)	
animal # (sex)	204	male	201	female	202	male	205	male	210	male	242	male
animal # (sex)	206	female	209	male	208	female	216	female	233	male	248	female
animal # (sex)	213	male	214	female	211	male	220	male	252	female	260	male
animal # (sex)	227	male	215	female	218	male	235	female	255	male	268	female
animal # (sex)	222	female	226	male	225	male	239	female	264	female	276	female
animal # (sex)	238	male	228	male	230	female	257	male	270	female	311	male
animal # (sex)	246	female	234	female	231	female	273	male	288	female		
animal # (sex)	253	female	237	female	232	female	274	female	298	male		
animal # (sex)	258	female	243	male	240	male	277	male	290	female		
animal # (sex)	262	male	267	female	244	female	279	female				
animal # (sex)	271	male	284	male	247	female	285	female				
animal # (sex)	278	female	301	female	249	male	286	female				
animal # (sex)	280	female	302	female	259	female	287	male				
animal # (sex)	282	male	303	male	265	male	292	female				
animal # (sex)	291	female	306	male	275	female	295	male				
animal # (sex)	294	female	317	male	281	female	307	male				
animal # (sex)					283	male	310	female				
animal # (sex)					305	male	322	male				

1230 Supplementary file 12: Number of sampled birds for organs at different days post-
1231 infection (dpi)

Group	Number of sampled birds		
	2dpi	6dpi	21dpi
RIG-I-RNF135-H7N1	6	5	0
WT-H7N1	7	6	3
RIG-I-H7N1	7	6	4
RNF135-H7N1	6	6	6
WT-MOCK	3	3	3
RIG-I-RNF135-MOCK	3	3	0

1232

1233

1234

1235

1236

1237

1238

1239

1240

1241

1242

1243

1244

1245

1246

1247

1248

1249

1250

1251

1252

1253

1254 **Supplementary file 13: Score system used to evaluate influenza-induced lung lesions.**
1255

Score 1 (+)
Mild, localized edema and
fibrinous exudate

Score 2 (++)
Moderate edema with hemorrhage
and fibrinous exudate over about
1/4 of the lung

Score 3 (+++)
Severe hemorrhage and
extensive edema
over about 1/2 of the lung

1256

1257

1258

1259

1260

1261

1262

1263

1264

1265

1266

1267

1268

1269

1270

1271

1272

1273