Title: Manipulating condensation of thermo-sensitive SUF4 protein tunes flowering time in Arabidopsis thaliana

Authors: Heather M. Meyer[1*], Takashi Hotta[2,3], Andrey Malkovskiy[1], Yixian Zheng[2], and David W. Ehrhardt[1*]

Affiliations:
1. Department of Plant Biology, Carnegie Institution for Science, Stanford, CA 94305, USA.

2. Department of Embryology, Carnegie Institution for Science, 3520 San Martin Dr. Baltimore, MD 21218, USA.

3. Department of Cell and Developmental Biology, University of Michigan, Ann Arbor, 109 Zina Pitcher Pl. Ann Arbor MI 48109, USA

* Corresponding authors: Heather M. Meyer:hmeyer@ciw.edu, David W. Ehrhardt: ehrhardt@stanford.edu

Abstract (286 words): Intrinsically disordered proteins (IDP) lack stable tertiary structures, which allows them to change conformation and function under different physicochemical conditions. This may be highly advantageous for plants, which often use changes in their environment to elicit a variety of responses, including developmental events. For instance, some plants delay flowering in the fall and require exposure to winter temperatures as a cue to initiate flowering the following spring. Many of the genes involved in temperature-dependent flowering have been extensively studied in *Arabidopsis*, yet how plants coordinate their molecular states with seasonal temperature change is poorly understood. Here, we explore the role of temperature-sensitive phase separation of the IDP and flowering-time regulator, SUPPRESSOR OF FRIGIDA 4 (SUF4), in modulating flowering time. SUF4 has a well-defined role in regulating temperature-dependent flowering time by activating the master floral suppressor *FLOWERING LOCUS C (FLC)*. We show that in plant nuclei, SUF4 is a temperature-sensitive protein that assembles into biomolecular condensates in warm temperatures (20°C). When temperatures cool (4°C), SUF4 nuclear condensates disassemble, causing SUF4 to disperse within the nucleoplasm. Additionally, we demonstrate that the number of SUF4 condensates quantitatively correlates with flowering time. Progressive alterations to the amino acid composition of SUF4's disordered region cause likewise progressive changes in temperature-dependent condensation both *in vitro* and *in vivo*, *FLC* transcription, and the onset of flowering. Lastly, we observe that SUF4

condensates coincide with the accumulation of other key flowering-time proteins (FRIGIDA and ELF7). These findings indicate that condensation of SUF4 likely plays a pivotal role in promoting flowering, possibly by concentrating and stabilizing regulatory factors needed for the transcriptional activation of *FLC* through temperature-dependent phase separation. This research suggests that in plants, IDPs can sense environmental cues and regulate critical developmental processes.

One-Sentence Summary: Evidence suggest that *Arabidopsis* uses temperature-sensitive condensation to regulate *FLC* transcription and flowering time.

Main text:
INTRODUCTION:

The survival of all species depends on their ability to reproduce. For plants, reproductive success relies heavily on the onset of flowering, a phenological feature influenced by changes in seasons. Many plant species, including winter crops, have evolved to use long exposure to winter temperatures (e.g., vernalization) as a cue to flower in the following spring. Temperature-induced development plays a crucial role in the long-term reproductive success and yield of plants, enabling them to synchronize their flowering with seasonal pollinators and set seed during favorable climatic conditions. Thus, as climate change drives new spatial and seasonal temperature trends, it is increasingly important to understand how plants physically sense and integrate thermal shifts.

In the model plant *Arabidopsis thaliana*, temperature-dependent flowering is modulated by the epigenetic silencing of the *FLOWERING LOCUS C (FLC)*. *FLC* encodes a MADS-box transcription factor that represses flowering by silencing floral-inducing genes including *FT* and *SOC1*[1–3]. In warm temperatures, SUPPRESSOR OF FRIGIDA 4 (SUF4), FRIGIDA (FRI), and FRI-associated proteins transcriptionally activate *FLC* by recruiting chromatin modifiers, such as members of the PAF1 complex, which methylate histone H3 lysine 4 (H3K4me3) at *FLC*'s transcriptional start site [3,4]. Conversely, exposure to cold temperatures promotes antisense long non-coding RNAs and a second set of chromatin remodelers belonging to the POLYCOMB REPRESSIVE COMPLEX 2 (PRC2) to epigenetically silence *FLC*, ultimately priming plants to flower the following spring [5,6]. While many elements of the genetic and epigenetic regulation of temperature-dependent flowering are well established, the mechanisms by which plants sense and transduce thermal changes to elicit these responses remain a mystery.

Intrinsically disordered proteins (IDPs) are highly sensitive to changes in their physical environment [7,8]. Unlike canonical proteins, which assume relatively stable structures with limited conformational changes in response to different stimuli and physiological conditions, IDPs possess low sequence complexity and a reduced ability to form stable tertiary structures. This inherent flexibility enables them to readily alter their conformation and function under varying physical and solution conditions. For example, many IDPs undergo phase separation into liquid-like protein droplets or viscoelastic hydrogels, facilitating rapid concentration, formation, and/or stabilization of higher-order complexes in response to environmental changes[8,9]. The ability for IDPs to phase separate has led to the intriguing hypothesis that this mechanism could serve as a biological means for cells and organisms to perceive and adapt to environmental signals[,10]. However, verifying this hypothesis poses a challenge due to the difficulty of distinguishing functions derived from phase separation from other protein properties.

Here, we identified that one of the key proteins required for regulation of *FLC* transcription and temperature-dependent timing of flowering, SUPPRESSOR OF FRIGIDA 4 (SUF4), exhibits significant similarities with BUGZ, an IDP known for its temperature-dependent phase condensation and involvement in mitotic chromosome alignment in animal cells[10–12]. Both proteins are comparable in size and feature two highly conserved zinc finger domains at their N-termini, followed by intrinsically disordered domains with low amino acid complexity enriched in proline, valine, and glycine residues. Given these structural parallels, along with BuGZ's documented biophysical behavior and SUF4's genetic and molecular implications in flowering timing, we asked if SUF4 also undergoes temperature-dependent phase transitions, using changes in its phase behavior to regulate the seasonal onset of flowering time.

Using confocal microscopy, genetics, and biochemistry, we discovered that SUF4 is indeed a temperature-sensitive protein that undergoes phase separation *in vitro*, forming proteinaceous droplets, and *in planta*, promoting the formation of nuclear condensates. At warm temperatures (20°C), SUF4 dynamically forms nuclear condensates, whereas at low temperatures (4°C), these condensates fail to form, causing SUF4 to disperse throughout the nucleoplasm. Furthermore, through use of targeted mutagenesis, we established the significance of SUF4's disordered region in nuclear condensate formation and the regulation of flowering time. Plants expressing SUF4 variants with graded aromatic/hydrophobic-to-hydrophilic amino acid substitutions in this region exhibit graded alterations in condensate formation, *FLC* transcript levels, and flowering-time responses. To explore the relationship between SUF4 condensates and other proteins involved in flowering time, we labeled two significant *FLC* regulators, FRIDIGA (FRI), and the PAF1-complex member EARLY FLOWERING 7 (ELF7) with fluorescent markers. We observed

that both proteins often appear in nuclear condensates that co-localize with a subset
of SUF4 condensates, indicating that SUF4 condensation occurs where other
important regulators of flowering time also accumulate. Interestingly, while FRI
could assemble into condensates containing SUF4 in some tissues (lateral roots),
its accumulation was apparently antagonized by SUF4 condensation in other tissue
(primary roots). Our findings provide strong evidence supporting the hypothesis
that SUF4 regulates flowering time by utilizing temperature-dependent phase
separation, which likely functions to enhance the activity of regulatory factors,
including itself, that are necessary for the transcriptional activation of *FLC*.
128
129 **RESULTS:**
130 **SUF4 forms nuclear condensates *in planta*.**
131   To investigate the hypothesis that SUF4 phase separation is important for the
regulation of flowering time, we first needed to assess if SUF4 undergoes
temperature-dependent condensation *in vivo*. To do this, we engineered a
monomeric fluorescent mEGFP-SUF4 fusion protein expressed from the putative
SUF4 native promoter and 3' UTR (pSUF4::mEGFP-SUF4) and introduced this
construct into *suf4* (*FRI*) mutant plants. We identified several transgenic lines that
rescued both the early flowering phenotype of the null mutant and the
transcriptional activity of *SUF4* as measured by the transcripts level of its
downstream target *FLC* (Supplemental Fig. 1A and 1B), suggesting that the *SUF4*
transgene is fully functional and expressed at adequate levels. We picked one of
these lines to perform all subsequent analyses regarding SUF4.
142   To assess the subcellular localization and behavior of SUF4 within living
plant cells, we imaged live cells using confocal microscopy at room temperature.
In both the shoot and the root, we observed SUF4 to exhibit a diffuse localization
pattern within the nucleoplasm and accumulate in nuclear foci reminiscent of
biomolecular condensates (Supplemental Fig. 2). Additionally, we found that
SUF4 localized in the nucleolar cavity, an elusive membrane-less compartment
that forms within the nucleolus (Supplemental Fig. 2). The nucleolar cavity is a
thermo-responsive structure in plant cells that exchanges its contents with the
nucleus [13–15]. However, we directed our analysis toward the function of the
nucleoplasmic foci outside of the nucleoli where most genes, such as *FLC*, are
transcribed [16,17].
153   If SUF4 condensates function as thermosensors, their formation should
exhibit temperature sensitivity across a physiologically relevant temperature range
in vivo. To test this, we subjected SUF4 rescue plants to a 24-hour cold treatment
at 4°C and then live imaged epidermal and cortical cells in the division and growth
zones of primary roots as the plants rewarmed to 20°C. At the beginning of
rewarming, mEGFP-SUF4 appeared diffuse and no condensates were detected

(Fig. 1A and 1B). As plants began to warm up, SUF4 signal progressively
partitioned into nuclear condensates, with condensates forming within seconds to
minutes once exposed to room temperature conditions. To ensure that these
changes were temperature-induced, we also imaged plants over the same time
course under constant 20˚C (Fig. 1B). Although some condensates showed
evidence of forming and dissipating over the time scale of observation (<10mins),
a portion of SUF4 always remained partitioned in condensates. Together, our
results suggest that SUF4 is a temperature-sensitive protein that assembles into
biomolecular condensates in warm temperatures and remains diffuse in the
nucleoplasm in cold temperatures.
169
SUF4 phase separates *in vitro*.
SUF4 is orthologous to the animal protein BuGZ (Supplemental Fig. 3), an
IDP initially found to undergo temperature-dependent phase separation *in vitro* and
regulate spindle assembly *in vivo*[10–12]. Therefore, to assess the potential for SUF4
to undergo phase separation independently of any plant cell factors, we expressed
and purified 6xHis-mVenus-SUF4 (isoform 1, WT) from SF9 insect cells. Using
both fluorescence and DIC microscopy, we observed that 6xHis-mVenus-SUF4
underwent phase separation at high temperatures ($25\mu M$ at 37˚C) and remained
miscible at low temperatures ($25\underline{\mu M}$ at 4˚C) when suspended in physiologically
relevant buffers (Fig. 2A, Supplemental Fig. 4, see methods). These results suggest
that, similar to BuGZ, SUF4 is a phase-separating protein displaying low critical
solution temperature (LCST) behavior. To investigate further the phase behavior of
SUF4, we conducted a turbidity assay to determine the thermal setpoint required
for SUF4's condensation. We found that $25\mu M$ of 6xHis-mVenus-SUF4 was fully
miscible between 4-10˚C but became increasingly turbid as temperatures gradually
rose to 30˚C (Fig. 2B). Additionally, in line with other phase-separating proteins,
SUF4's phase transition is concentration-dependent, with lower concentrations of
SUF4 requiring higher temperature regimes to undergo phase separation (Fig. 2B,
Supplemental Fig. 4). These results indicate that, similar to BuGZ, SUF4 is a
temperature-sensitive protein that undergoes LCST phase separation *in vitro*.
190
SUF4's disordered region drives nuclear condensate formation *in vitro* and *in*
planta.
IDP phase separation is driven by multivalent interactions, such as those
facilitated by aromatic/hydrophobic residues like tyrosine and phenylalanine.
Moreover, these same residues are required for *in vitro* condensation of SUF4's
ortholog, BuGZ[11,12]. Despite being a proline-rich protein, SUF4 contains a total of
14 regularly spaced tyrosine and phenylalanine residues within its disordered
region, with two of these residues falling in putative splice sites (Supplemental Fig.

3). To test if aromatic/hydrophobic residues in SUF4's disorder region promote its
temperature-dependent condensation, we created a SUF4 variant, termed 12S
SUF4, in which we substituted 12 non-splice-interfering tyrosine and
phenylalanine residues with serine and asked if this variant had a different phase
separation thermal setpoint *in vitro*. We found 12S SUF4 had a large shift in its
thermosensitive capabilities to phase separate, such that 25μM of purified 12S
SUF4 now required temperatures nearing 50°C to begin phase separating as
opposed to 20°C for the native protein (Fig. 2B). These results suggest that
changing SUF4's aromatic/hydrophobic patterning is sufficient to disrupt
condensate formation *in vitro* by increasing the thermal set point at which SUF4
can phase separate.
To test if changes in SUF4 aromatic/hydrophobic patterning also influenced
condensate formation *in vivo*, we generated the same 12 mutations in the mEGFP-
SUF4 fusion protein expressed over a chromosomal null locus (pSUF4::mEGFP-
12S SUF4). Similar to our observations *in vitro,* we found that 12S SUF4 exhibited
a severe reduction in condensate formation *in vivo*. On average, 12S SUF4 plants
formed 0-1 nuclear condensates per nucleus in root epidermal and cortical nuclei at
20°C, as compared to 4-5 per nucleus formed in rescue SUF4 (Fig. 3A and 3B,
nuclei observed: 12S SUF4 n= 27, rescue SUF4 n=32). These results suggest that
the altered coacervation properties of 12S SUF4 are maintained in the complex
environment of the living cell.
Next, we investigated if SUF4 uses phase separation as a means to regulate
flowering time. As a first step in testing this hypothesis, we asked if 12S SUF4
variant plants displayed a phenotypic difference in its response as compared to the
rescue SUF4 plants. Using a leaf counting assay, we found that 12S SUF4 plants
transitioned to flowering much sooner, producing significantly fewer rosette leaves
before flowering (Long Day (LD): 35 leaves (n=30)) as compared to SUF4 rescue
plants (LD: 71 leaves (n=29)), but significantly more so than the *suf4* null mutant
(LD: 26 leaves (n=31)) (Fig. 2C-E, Supplemental Fig. 5). Previous research has
shown that high levels of *FLC* must be produced for the suppression of flowering
time. We therefore asked if the flowering time phenotype caused by the 12S
mutant protein was correlated with a reduction in *FLC* transcripts. We found plants
expressing the 12S variant to produce approximately 18-fold fewer *FLC* transcripts
than our SUF4 rescue lines but 2-fold more than *suf4* null mutants (Fig. 3C). Thus,
although *FLC* still undergoes some transcription in the 12S SUF4 variant, the
mutations that caused disruption of SUF4 temperature-sensitive condensation
greatly accelerated the onset of flowering.
236
Tuning SUF4 phase behavior causes graded changes in *FLC* transcription and
flowering time

Previous work from Jiang et al (2015) has shown that the *in vitro* phase
behavior of SUF4's ortholog, BuGZ, may be tuned by incrementally decreasing the
number of combined phenylalanine and tyrosine to serine substitutions[10]. This
finding, paired with the correlation between SUF4 phase separation and flowering
time, prompted us to further assess if SUF4's phase change properties themselves
were critical for dictating different flowering time outcomes. We reasoned that if
so, substituting a smaller number of the phenylalanine and tyrosine residues should
lead to a scalable intermediate phase separation phenotype and proportionally tune
flowering time outcomes via decreasing the transcriptional levels of *FLC*. To test
this prediction, we created an intermediate SUF4 variant that only had 5 combined
tyrosine and phenylalanine to serine substitutions termed, 5S SUF4, and assessed
its phase separating thermal setpoint *in vitro*. Consistent with what had been
observed for BuGZ, we found purified 5S SUF4 to have a thermal shift that ranges
between the phase change of rescue SUF4 and 12S SUF4 *in vitro* (Fig. 2B; 25µM
showed a steep phase transition at 40° C instead of 20°C like rescue SUF4 and
50°C like 12S SUF4).
To test the function of the 5S SUF4 variant *in vivo*, we introduced it into
suf4 null mutant plants and asked if these plants exhibited intermediate phenotypes
relating to SUF4 phase separation, *FLC* transcriptional output, and flowering time.
Remarkably, we found 5S SUF4 plants to display an intermediate condensate
phenotype *in vivo*, in which they formed significantly fewer nuclear condensates in
root epidermal and cortical cells than rescue SUF4 but more than 12S SUF4 (Fig.
3A and 3B; rescue SUF4:4-5, 5S SUF4:2-3, 12S SUF4:0-1). This outcome differed
from the results observed with 5S SUF4 purified protein, which showed no phase
separation at room temperature, suggesting that the complex cellular environment
may influence the phase behavior of 5S SUF4 (Figure 2A). Moreover, 5S SUF4
plants exhibited *FLC* transcript levels and flowering time responses that both fell
between rescue SUF4 and 12S (Fig. 3C-E, Supplemental Fig. 5; rescue SUF4
produced 20-fold/ LD: 71 leaves (n=29), 5S SUF4: 10-fold/LD: 43 leaves (n=39),
12S SUF4 3-fold/ LD: 35 leaves (n=30)). These results indicate that tuning the
LCST behavior of SUF4 is sufficient to predictably change the onset of flowering
through modulating transcript levels of *FLC*.
271
SUF4's exchange dynamics scale with changes in LCST behavior
Biomolecular condensates may assume a broad range of material properties,
spanning liquid-, gel-, or solid-like structures, with these respective properties
often being hypothesized as being important for their biological function[7,18–20].
Under *in vitro* conditions, purified and condensed SUF4 exhibited behavior
consistent with a liquid, similar to its ortholog BuGZ. For example, *in vitro*
condensates on glass assume a round outline, as expected at a liquid-liquid

interface, and are rapidly and repeatedly reversible with temperature change (Fig.
2A, Supplemental Fig. 4). However, whether SUF4's material properties remain
liquid-like during interactions with other biomolecules under native cellular
conditions remains unknown. Previous research has suggested that the exchange
dynamics of molecules within condensates are expected to correlate with their
material state, with faster exchange rates in liquid-like condensates and slower
rates in more solid-like structures[20,21]. Thus, to investigate the exchange dynamics
of SUF4 and whether these dynamics are affected in our SUF4 variants, we
conducted fluorescence recovery after photobleaching (FRAP) experiments on
rescue SUF4, 5S SUF4, and a hydrophobic variant termed 30I SUF4, which
contains 30 alanine-to-isoleucine substitutions in its disordered region
(Supplemental Fig. 6). We excluded the 12S SUF4 variant as it rarely formed
condensates. FRAP was performed in a small region of approximately 1.094
microns^2, centered on a prominent mEGFP-SUF4 nuclear body in each cell
assayed. Our findings revealed that changing the hydrophobicity of SUF4 greatly
affected its FRAP dynamics. 30I SUF4 condensates did not reach their half
recovery time within a 10-minute period, whereas both rescue SUF4 and 5S SUF4
rapidly reached their half recovery time within seconds (Fig. 4A, rescue SUF4:
t1/2=1.27 seconds, n=13; 5S SUF4 t1/2=0.85 seconds, n=12, 1-sided t-test
p<0.01). Additionally, 30I SUF4 condensates exhibited a higher "immobile"
fraction compared to SUF4 rescue and 5S SUF4, which both largely recovered (30I
SUF4 = 0.094 mobile fraction, SUF4 rescue = 0.85 mobile fraction, and 5S SUF4
= 0.76 mobile fraction; Fig 4A). Interestingly, despite 30I SUF4 having formed
many condensates that were relatively slow to recover, we did not observe any
rescue in plant phenotype concerning *FLC* transcription or flowering time
(Supplemental Fig. 6). Previous studies have suggested that the biophysical
properties of some condensates must fall within a fluidic window, in which being
overly solid or fluid-like can lead to a decrease in functionality [20–25]. Therefore, in
agreement with these findings, we propose that SUF4 needs to maintain a specific
state of fluidity in order to function optimally.
A notable finding from our FRAP experiments is that the recovery curves of
5S SUF4 signal exhibited a lower R^2 value on average when fitted to a single
exponential recovery model, along with a lower bleaching depth compared to those
observed in our rescue lines (rescue SUF4 R^2 = .73, 5S SUF4 R^2 = .53). This
observation prompted us to further investigate the behavior of condensates in this
background. In doing so, we noticed that 5S SUF4 condensates appeared to display
greater motility compared to rescue SUF4 condensates. An increase in particle
motility could explain the reduced model fit as movement of condensates out of the
focal plane would tend to reduce their brightness, thus affecting measurement of
signal recovery kinetics. To quantify the perceived difference in motility, we

measured the total distance traveled by each condensate during our FRAP analysis. On average, we found that 5S SUF4 condensates traveled significantly greater distances within the nucleoplasm compared to rescue SUF4 condensates (Fig. 4B, average distance of 1.802 microns for rescue SUF4 and 3.046 microns for 5S SUF4). These results indicate that 5S SUF4 condensates are less constrained in their positional behavior compared to their rescue SUF4 counterparts.

SUF4 colocalizes with FRI within nuclear condensates at room temperature in root cells recently derived from meristems

SUF4 has been shown previously to physically and genetically interact with FRIGIDA (FRI) and other members of the vernalization pathway, and this complex has been proposed to transcriptionally activate *FLC* [3,4]. Therefore, to ask if SUF4 condensation occurs where other flowering time regulators are present, and to explore the possibility that SUF4 condensation might function to regulate *FLC* by concentrating flowering time proteins, we tested if there was an observable spatial relationship between SUF4 and FRI *in vivo*. We generated a fluorescently-labeled FRI fusion gene (FRI-mScarletI) that expresses from the native upstream sequence of FRI (pFRI) and transformed it into a mEGFP-SUF4 rescue line. Under room temperature conditions, we found FRI-mScarletI to exist in one of two distinct phase states within the nucleoplasm of root epidermal and cortical cells within the division and growth zones— either uncondensed and diffuse within the nucleoplasm or assembled into one-to-two apparent condensates (Fig 5A and B).

Next, we tested if FRI-mScarletI condensates colocalized with mEGFP-SUF4 condensates. To measure co-localization, we considered condensates to be colocalized if the centroids of spot-detected condensates were closer together than the radius of the kernel used to detect the SUF4 condensates (<250nm). Out of 56 FRI-mScarletI condensates analyzed, 38 of them co-localized with a SUF4 condensate (Fig. 5A; 68% co-localization, n=47 nuclei, distance threshold=0.3μm). To ask whether this degree of colocalization was specific to the vernalization complex and not likely due to chance alone given the density of SUF4 nuclear condensates, we co-transformed the nuclear splicing speckle marker U2B"-mRFP into mEGFP-SUF4 rescue plants, a marker, like that for FRI, produces 1-2 bodies of larger size per nucleus. In the 18 nuclei examined, we found no instances of U2B" bodies co-localizing with SUF4 at room temperature conditions (0/19 U2B" speckles; Fig. 5A). These data demonstrate a remarkable degree of colocalization between SUF4 and FRI, suggest that the formation of FRI condensates under warm temperatures is not a random occurrence but rather occurs preferentially where condensation of SUF4 also occurs.

SUF4 and FRI colocalization depends on tissue type and temperature

FRI has recently been reported to have no quantifiable physical interactions
with SUF4 and to accumulate into nuclear condensates when exposed to extended
periods of low temperatures [26–28]. Thus, to test the spatial relationship between
SUF4 and FRI in the cold, we exposed mEGFP-SUF4 and FRI-mScarletI co-
expressing plants to 4°C for 24 hours and imaged their root cells as the plants
warmed up. Consistent with previous findings[26–28], we found FRI to form cold-
induced nuclear condensates. However, unlike the case at room temperature
(~21°C), these condensates did not co-localize with SUF4. Instead, SUF4
condensates would sometimes appear in the same locations where FRI condensates
previously existed as plants began to warm (observed in at least 7 nuclei; Fig 6A
and B). These results suggest an antagonistic or independent relationship between
cold-induced FRI condensates and SUF4. Additionally, they suggest that FRI
undergoes cold-induced phase separation that is independent of *FLC* activation (a
process reliant on SUF4).
Given the established relationship between the spatiotemporal organization
of proteins and their biological function, we investigated the root tissue types in
which both warm- and cold-induced FRI condensation occurred. Under warm
temperatures, we observed the presence of FRI condensates in the division and
growth zones of a subset of emerging and young lateral roots but not in primary
roots (Fig. 6C, Nplants=8/29 lateral roots, Nplants=0/34 primary roots of 6-13
DAG). This was different to what occurred under cold temperatures, in which we
found FRI condensing in all root types, including the primary roots (Supplemental
Fig. 7). Interestingly, the occurrence of warm-induced FRI condensates within a
given lateral root cell was highly variable among lateral roots, with detectable FRI-
mScarletI condensates found in anywhere from 30% to 94% of the epidermal and
cortical cells within the growth zone (Supplemental Fig. 7). The seemingly
stochastic patterns of nuclei containing FRI condensates among and within lateral
meristems resemble the previously described phenomenon of transcriptional and
translational "bursting," which has been associated with epigenetic and phenotypic
diversity in response to environmental changes[17,29,30].
Previous research has highlighted the influence of plant tissue types on the
state of auxin response factor (ARF) biomolecular condensates[31]. The observation
of co-condensation between SUF4 and FRI in lateral roots, while absent in primary
roots of 6-13 days after germination plants, suggests the potential for a tissue-
specific functional relationship between these proteins. If this is true, we might
observe a tissue-specific change in mScarletI-FRI condensate behavior in response
to changes in SUF4 condensation. To test this hypothesis, we introduced FRI-
mScarletI into *suf4* null, 12S SUF4, and 5S SUF4 variant plants. At room
temperatures, we observed the formation of FRI-mScarletI condensates in about
10%-25% of primary roots of *suf4* null and 12S SUF4 plants, respectively (Fig.

6C, Table. 1; suf4 4/41, 12S SUF4 4/5), but no FRI-mScarletI condensates were
observed in the primary roots of rescue SUF4 (0/34) or 5S SUF4 lines (0/19). FRI-
mScarletI condensates were observed in a portion of lateral roots in all lines.
Although the number of mGFP-SUF4 condensates varies between rescue SUF4
and 5S SUF4, both genotypes have the ability to form condensates, whereas 12S
and *suf4* (*FRI*) do not. Thus, our data indicate an inverse relationship exists
between SUF4 and FRI condensates in the primary root. Furthermore, they suggest
that FRI is capable of forming at least two distinct types of condensates that vary
by tissue type: one type that assembles with SUF4, likely functioning to help
establish *FLC* transcription in newly formed tissue, and another type where the
proteins possibly act antagonistically.
410
411
412 **Table 1.** FRI-mScarletI condensate formation in different root types within SUF4
413 variants

	FRI Condensate formation			
	Lateral		**Primary**	
Genotype	Warm	Cold	Warm	Cold
FRI-mScarletI (*mEGFP-SUF4*)	Yes (n=8/29 roots)	?	No (n=0/34 roots)	Yes
FRI-mScarletI (*mEGFP-5S SUF4*)	Yes (n=2/6 roots)	?	No (n=0/19 roots)	Yes
FRI-mScarletI (*mEGFP-12S SUF4*)	Yes (n=2/2 roots)	?	Yes (n=4/5 roots)	Yes
FRI-mScarletI (*suf4 null*)	Yes (n=5/22 roots)	?	Yes (n=4/41 roots)	Yes

414
SUF4 colocalizes with ELF7 in nuclear condensates at room temperature
While FRI facilitates transcriptional activation of *FLC* by establishing a
permissive chromatin environment, other mechanisms have been shown to be
involved in the maintenance of *FLC* expression, such as transcriptional regulation
associated with histone H3 lysine 4 (H3K4) modifications [32,33]. In light of this, we
sought to investigate whether SUF4 condensates might serve as a hub for multiple
proteins involved in the transcriptional regulation of *FLC* by fluorescently tagging
and expressing EARLY FLOWERING 7 (ELF7), a known member of the PAF1
complex. We introduced mTFP-ELF7 expressed from its own upstream sequence
(pELF7::mTFP-ELF7) into rescue lines expressing mEGFP-SUF4 and mEGFP-
SUF4/FRI-mScarletI. ELF7 was specifically chosen due to its previous genetic

interaction with SUF4 and its role in establishing H3K4 activation marks on *FLC*,
as well as its physical co-localization with FRI in nuclear condensates under
normal temperature conditions[26,32,33]. Within both transgenic lines, we observed
mTFP-ELF7 to be diffuse within the nucleoplasm of cortical and epidermal cells
within the transition zone of both lateral and primary roots. In a subset of these
cells, mTFP-ELF7 formed 1-2 prominent nuclear condensates and occasionally
multiple smaller condensates (Fig. 5B).
As the introduction of mTFP-ELF7 did not appear to induce silencing, nor
alter condensate behavior in either the rescue mEGFP-SUF4 or mEGFP-
SUF4/FRI-mScarletI lines, we performed colocalization analysis using lines
expressing all three transgenes (Fig. 5B). While mEGFP-SUF4 formed more
condensates compared to ELF7 and FRI in nuclei analyzed (nSUF4condensates=
414, nELF7condensates= 53, nFRIcondensates= 34), the presence of an ELF7
condensate almost always coincided with a SUF4 condensate (47 out of 53 cases).
Further, out of these 47 ELF7-SUF4-containing condensates, we observed the
additional presence of a FRI condensate in 8 cases (17%). The quantitative
correlation between SUF4 condensation and flowering time, coupled with its
ability to co-localize with both FRI and ELF7 in 1-2 condensates, supports the
notion that SUF4 condensation likely functions in regulating the transcriptional
activity of *FLC* under different temperature conditions (Fig. 7).
446

DISCUSSION:
In this study, we have discovered that the IDP, SUF4, undergoes thermally-
regulated condensation both *in vitro* and *in planta*. These findings, along with the
demonstrated ability of SUF4 sequence alterations to progressively tune phase
separation, *FLC* transcription, and predictable flowering time outcomes, and the
evidence that SUF4 condenses with *FLC* transcriptional regulators under
appropriate physiological conditions, provide strong support for the hypothesis that
SUF4 phase separation may serve as a physical mechanism for plant temperature
perception. Our work emphasizes the biological importance of protein phase
separation in a multicellular organism by establishing links between
environmentally-sensitive phase separation, gene expression, development and
physiology.
Plant cells require an ability to filter out noise and interpret meaningful
signals from their environment to make developmental decisions. The
conformational promiscuity and multivalency of IDPs enables them to engage in a
variety of specific yet transient interactions under defined environmental
conditions. IDPs are enriched within plant proteomes, with many participating in
plant regulatory pathways and as integrators of environmental inputs[34–36]. The
phase separation states of such IDPs and client proteins could serve as an effective

mechanism for distinguishing real signals from noise within plant cells, thereby enabling more efficient and effective developmental outcomes. Plants, along with other organisms, use epigenetic factors, such as changes in chromatin/histone state, signaling pathways, and RNA modifications to promote environmental-driven developmental changes. However, these changes often require integration over a significant period of time in order to induce a physiologically-relevant response. Pairing IDP sensing with slower-acting epigenetic mechanisms may allow plants to better distinguish genuine signals from noise by facilitating the promotion or maintenance of molecular memory under persistent environmental cues. In the case of SUF4, we found that changes in temperature cause it to phase transition between condensed and uncondensed forms. Moreover, manipulating the aromatic/hydrophobic properties of SUF4 proved to be sufficient to modulate its material characteristics and tune the onset of flowering. Given SUF4's known interaction with *FLC*-related chromatin remodelers and our cell biological evidence that at least one of these co-condenses with SUF4 *in vivo*, these findings suggest the possibility that SUF4 condensates could also associate with and alter the activity of epigenetic factors to modulate *FLC* transcription.

Extensive investigations have shed light on the intricate regulation of flowering time in *Arabidopsis*, with a specific emphasis on the transcriptional modulation of *FLC*[1]. Among the key players in this regulatory network is FRI, an interacting partner of SUF4[2,36]. Equipped with two coil-coil domains, FRI is proposed to act as a scaffold protein, facilitating interactions with various *FLC* transcriptional regulators and chromatin remodelers. Nevertheless, FRI itself does not directly interact with the *FLC* gene locus. Instead, SUF4 serves as a mediator of this interaction by physically binding both *FLC* and FRI, along with many other members associated with the FRI complex [3,37]. Through both condensation and its ability to recognize the *FLC* promoter region, SUF4 has the potential to enhance its own activity by raising its local concentration, and also possibly by recruiting and concentrating transcriptional activators (e.g. previously reported SUF4 interactors FLX and TAF14), and chromatin remodelers (e.g. ELF7, SWC6, and RVB1) at the *FLC* promoter. Meanwhile, we hypothesize that FRI transiently interacts with SUF4, contributing to the establishment of transcriptional regulation of *FLC* in newly formed tissues such as lateral roots. Newly developed and emerging molecular images and technologies, such as those that enable the visualization and quantitation of molecular events at a single genetic locus (e.g., endogenous fluorescent tagging of gene loci via dCAS9/CRISPR), will aid in further investigating the functional dynamics that govern these interactions at the *FLC* gene locus.

The coacervation of prion-like proteins (PrLPs), such as FLL2 and FCA, has recently emerged as a potential regulatory mechanism in flowering time [38].

However, it remains uncertain whether PrLP coacervation is thermally-driven, or
even contributes to the post-transcriptional regulation of *FLC*. In this study, we
demonstrate that a distinct proline-rich IDP undergoes LCST condensation
behavior, which likely serves as a means to modulate flowering. This behavior
aligns with the LCST observations of synthetic elastin-like polypeptides and Pab1
[22,39–42], which also possess a high proline content and repetitive hydrophobic
features. While plants carry numerous proline-rich proteins and hybrid proline-rich
proteins that are involved with abiotic- and biotic-associated pathways, the thermo-
induced coacervation behavior of these proteins remains underexplored. By
unraveling the role of proline-rich IDPs in phase separation and its relevance to
flowering time regulation, our study extends our understanding of proline-rich
IDPs in biological processes and sheds light on their potential participation in
eukaryotic thermo-responses through phase separation phenomena. Moreover, our
findings provide a strong basis for exploring the role of phase separation and
environmental sensing, providing a valuable foundation and new insights into how
we may be able to engineer plants and other organisms with tailored environmental
responses.
Despite IDPs being often poorly conserved [43,44], SUF4 shares substantial
homology with the animal protein BuGZ. In these organisms, BuGZ plays essential
roles in both mitotic and transcriptional regulation by interacting with various
cellular machineries, including the spindle assembly checkpoint protein (SAC)
Bub3, the spliceosome, and epigenetic factors[10–12]. While SUF4's primary function
in plants revolves around regulating flowering time, recent research has unveiled
its potential involvement in analogous mitotic and transcriptional regulatory
processes [45–47]. These newfound roles raise a series of questions, including whether
SUF4's phase separation has evolved distinct functionality specific to plant
flowering or if it serves as a universal mechanism within eukaryotic systems.
Interestingly, animal BuGZ has been observed to undergo phase separation *in*
vitro, and its role in phase separation is hypothesized to contribute to mitotic
spindle formation. Therefore, studying SUF4 may not only advance our
comprehension of how environmentally-induced phase separation impacts plant
development and physiology but also hold significant promise in deepening our
understanding of the broader evolutionary significance and functional implications
associated with such environmentally-induced phase separation.

MATERIALS AND METHODS:

Plant Material, accessions, and growth conditions

All plants used in this study were grown under controlled conditions in growth rooms with a light intensity of approximately 180 μmol m−2 s−1 at a temperature of 21°C. Plants were either exposed to long day conditions (16 hours of light followed by 8 hours of darkness).

Col (FRI-SF2) [48] was used as the wild-type accession for all genotypes analyzed in this study. These lines are characterized as winter annuals and exhibit a delayed flowering phenotype.

suf4 (FRI-SF2) plants [32] were used as the mutant accession for this study. These plants carry an ~6kb deletion that spans the N-terminus of SUF4 (deleted region: exons 1,2,3 and part of intron 3), resulting in a null mutation that causes an early flowering phenotype.

Transgenic lines generated in this study
pSUF4::mEGFP-SUF4 in *suf4 (FRI-SF2)*
pSUF4::mEGFP-5SSUF4 in *suf4 (FRI-SF2)*
pSUF4::mEGFP-12SUF4 in *suf4 (FRI-SF2)*
pFRI::FRI-mScarletI; pSUF4::mEGFP-SUF4 in *suf4 (FRI-SF2)*
pFRI::FRI-mScarletI; pSUF4::mEGFP-5SSUF4 in *suf4 (FRI-SF2)*
pFRI::FRI-mScarletI; pSUF4::mEGFP-12SSUF4 in *suf4 (FRI-SF2)*
pFRI::FRI-mScarletI; pSUF4::mEGFP-30ISUF4 in *suf4 (FRI-SF2)*
pELF7::mTFP-ELF7; pSUF4::mEGFP-SUF4 in *suf4 (FRI-SF2)*
pELF7::mTFP-ELF7; pFRI::FRI-mScarletI; pSUF4::mEGFP-SUF4 in *suf4 (FRI-SF2)*
U2B"-mRFP(pROK2); pSUF4::mEGFP-SUF4 in *suf4 (FRI-SF2)*

Flowering time quantification
To assess flowering time in SUF4 variants, seeds of each genotype were
individually sown in Pro Mix BX soil and stratified at 4°C for two days. Then,
planted pots were transferred to either short day or long day chambers. The total
number of true rosette leaves was recorded every 3-5 days until plants bolted. To
minimize position effects, all genotypes were spatially mixed within their
respective short- or long day growth chambers.
613
614 **Cloning schemes:**
615
pHM11 (rescue SUF4):
To create pHM11 (pSUF4::6HIS-mEGFP-SUF4), a 2,981 fragment containing the
genomic SUF4 sequence was amplified using oHM3 (TTA GGA TCC GGT GGA
GGC GGT TCA GGC GGA GGT GGC TCT GGC GGT GGC GGA TCG ATG
GGT AAG AAG AAG AAG AGA), a forward primer with an added BamHI site
and a 3XGGGS linker, and oHM4 (GGC CTC GAG CTA AAA CGC CAT CCG
CCC AGC AA), a reverse primer with a 3' XhoI site. This fragment was then
cloned into pGEM-T easy to generate pHM1 (genSUF4). Additionally, the mEGFP
gene was amplified from addgene's commercially available mEGFP plasmid
(#18696) using oHM1 (ATA AGA TCT AGT GAG CAA GGG CGA GGA G), a
primer with a BglII 5' site, and oHM2 (ATG GAT CCC TTG TAC AGC TCG
TCC ATG CC), a primer with a 3' BamHI site. The resulting fragment was cloned
into pGEM-T easy to create pHM2 (mEGFP). Subsequently, pHM2 was digested
with BglII and BamHI, and then inserted into the pET-30a(+) vector, resulting in
pHM3 (6HIS-mEGFP). Plasmids pHM1 and pHM3 were then cleaved with BamHI
and XhoI and ligated, yielding pHM4 (6HIS-mEGFP-SUF4). A 1059 bp fragment
upstream of the SUF4 coding region was PCR amplified using oHM28 (ATC TTT
AAC TTT TTT GTT GTT GCA) and oHM24 (GAT GGC GCG CCC ACT AAA
TCT TTC TCC TA), a primer with an ASCI site, and inserted into the pGEM-T
easy vector, generating pHM5 (proSUF4). Additionally, a 332 bp region
downstream of the SUF4 coding region was PCR amplified using oHM25 (GAT
GGC GCG CCC TCG AGA ATT TGG CAC CAA ACC AA), a primer with both
ASCI XHOI restriction sites, and oHM17 (gaacaacatggtgcaaacatt), and cloned into
pGEM-T easy to produce pHM6 (termSUF4). Subsequently, pHM5 was digested
with SacII and AscI, and the resulting fragment was cloned into pHM6 to create
pHM7 (proSUF4::AscI-XhoI-3'UTR). pHM4 (6HIS-mEGFP-SUF4) was reverse
amplified to add a kozak consensus sequence using primers ohm45
(TTCACCAAAATGCACCATCATCATCATCAt) and oHM46
(GGcgcgCCCACTAAATCTTTCTCC), and then digested with AscI and XhoI for
cloning into pHM7, resulting in pHM10 (proSUF4::6HIS-mEGFP-SUF4-3'UTR).

For downstream biochemical assays requiring removal of mEGFP, an enterokinase
sequence was introduced before the SUF4 coding region using primers oHM35
(CGATCCGCCACCGCCAGAGCC) and oHM36
(gacgacgacgacaagATGGGTAAGAAGAAGAAGAGA), generating
pHM10_entero (proSUF4::6HIS-mEGFP-entero-SUF4-3'UTR). Subsequently,
pHM10_entero was cloned into pBIN30 via the NotI restriction site to make
pHM11 (proSUF4::6HIS-mEGFP-entero-SUF4-3'UTR).

pHM32 (30I SUF4):
To create pHM32, the last 1969 bp of SUF4's disordered region was synthesized
(GenScript, USA) with base pair substitutions that replaced all 30 alanine residues
with isoleucine residues. Gibson assembly was used to fuse this fragment with the
remaining 1012 bp genomic sequence of SUF4 to reassemble the complete gene.
Once the 30I SUF4 fragment was assembled, pHM10_entero and the 30I fragment
were cut with BamHI and XhoI and ligated together to make pHM31
(pSUF4::mEGFP-30ISUF4). Then, pHM31 was cut with Not1 and cloned into the
binary vector pBIN30 (Basta resistant) to make pHM32 (pSUF4::mEGFP-
30ISUF4).

pHM14 (12S SUF4):
To create pHM14, the last 1969 bp of SUF4's disordered region was synthesized
(GenScript, USA) with base pair substitutions that replaced 12 of the 14
phenylalanine and lysine residues with serines. Gibson assembly was used to fuse
this fragment with the remaining 1012 bp genomic sequence of SUF4 to
reassemble the complete gene. Once the 12S SUF4 fragment was assembled,
pHm10_entero and the 12S fragment were cut with BamHI and XhoI and ligated
together to make pHM12 (pSUF4::mEGFP-12SSUF4). Then, pHM12 was cut with
Not1 and cloned into the binary vector pBIN30 (Basta resistant) to make pHM14
(pSUF4::mEGFP-12SSUF4).

pHM5S (5S SUF4):
To create pHM5S, the last 1969 bp of SUF4's disordered region was synthesized
(GenScript, USA) with base pair substitutions that replaced the last 5
phenylalanine and lysine residues with serines. Gibson assembly was used to fuse
the remaining 1012 bp genomic sequence of SUF4 to reassemble the complete
gene. Once the 5S SUF4 fragment was assembled, pHm10_entero and the 5S
fragment were cut with BamHI and XhoI, and ligated together to make pHM13
(pSUF4::mEGFP-5SSUF4). Then, pHM13 was cut with Not1 and cloned into the
binary vector pBIN30 (Basta resistant) to make the final vector pHM5S
(pSUF4::mEGFP-5SSUF4).

686
pFRI::FRI-mScarletI:
pFRI::FRI-mScarletI was custom-synthesized by GenScript and consists of the
following elements: The FRI 1243 bp upstream promoter region with an added
Kozak consensus sequence (CACCAAA), directly followed by the 1351 bp FRI
(SF2) genomic region, 3XGGGS linker (GGT GGA GGC GGT TCA GGC GGA
GGT GGC TCT GGC GGT GGC GGA TCG), mScarletI sequence derived from
Addgene 98839, and FRI's putative 169 bp 3'UTR. Both promoter and 3'UTR
sequences were obtained from TAIR Sequence Viewer, whereas the genomic
sequence was provided by Sang Yeol Kim. The aggregated synthesized DNA was
flanked with NotI cut sites and subsequently inserted into the pBIN40 (Hyg
resistant) binary vector through NotI digestion and ligation.

pELF7::mTFP-ELF7:
pELF7::mTFP-ELF7 was custom synthesized by GenScript, USA and comprises
the following components: The ELF7 405 bp upstream promoter region with an
added Kozak consensus sequence (CACCAAA), directly followed by the mTFP
sequence (Addgene 55503), 3XGGGS linker (GGT GGA GGC GGT TCA GGC
GGA GGT GGC TCT GGC GGT GGC GGA TCG), the 3228 bp ELF7 genomic
region (TAIR), and the ELF7 putative 1791 bp 3' UTR. All promoter, genomic,
and 3'UTR sequences were obtained from TAIR Sequence Viewer. The
synthesized DNA was flanked by NotI sites, which were used for subsequent
insertion into the pBIN40 (Hyg resistant) binary vector.

U2B"-mRFP (pROK2): U2B" -mRFP (pROK2) was previously created by Alie
Pendle and Peter Shaw and commercially available for order via ABRC
(https://abrc.osu.edu/stocks/number/CD3-795056).

**Live Cell Microscopy and Fluorescence Recovery After Photobleaching
(FRAP)**

Live-cell imaging was performed on roots of 6-7 day old seedlings using either
point scanning or a spinning disk scanning confocal microscopy. Point-scanning
was performed on a Leica SP8 microscope equipped with HyD detectors, a white
light supercontinuum laser, and AOBS (Acousto-Optical Beam Splitter) control of
shuttering, excitation and emission bands. All images were acquired with a Leica
60x 1.3 N.A Plan-Apo glycerin immersion objective. Leica Lightning
deconvolution was used to improve signal to background for all point-scanning
images. Spinning disk confocal microscopy was performed either with a CSU-
W110 confocal head (Yokogawa) combined with a MESA Field-flattening device

(Intelligent Imaging Innovations) mounted on an Olympus IX83 microscope
equipped with a 100x Plan-Apo 1.35 NA silicon oil immersion objective, or a fully
automated Leica DMI6000B microscope equipped with a CSU-X spinning disk
confocal head (Yokogawa Corp.). Each sample was excited using 488nm for GFP
(Coherent cube laser), 561nm for mScarletI (Coherent cube laser), or 442nm for
mTFP (Coherent cube laser). Emitted light was collected with a 525/50 nm filter
for GFP, 605/64 nm filter for mScarletI and a 482/35 nm filter for mTFP (all filters
from Semrock).
734
FRAP experiments were conducted using the CSU-X spinning disk microscope
described above, with aid of a Vector laser scanning unit (Intelligent Imaging
Innovations) attached to the right camera port to focus and steer laser energy on the
image plane. The camera port was switched from imaging to bleaching paths via
software control (SlideBook, Intelligent Imaging Innovations) to alternate between
imaging and FRAP light paths. mEGFP-SUF4 was excited using 3.5 mW of light
energy (as measured at the end of the laser fiber entering the confocal head) from a
solid state 488nm laser (Coherent Technologies). Emission light was collected
using a 488nm dichroic mirror and 500-550nm emission filter (Semrock). Single-
plane Images were collected every 200ms using a Leica 100x Plan-Apochromat
100x 1.40 NA oil immersion objective and FRAP was performed within a 1.094
micron^2 region of interest centered on each assessed condensate spot using 5mW of
light energy as measured at the end of the fiber entering the Vector unit. The focal
spot was moved by galvometers to bleach a grid of spots separated by 20 nm
within the FRAP region. Seven frames were collected before bleaching and 143
frames were collected after bleaching (1.75 seconds to conduct bleaching, 0.2
second delay to first post-bleach acquisition). Once all images were acquired, they
were aligned in ImageJ [49] using the MultiStackReg plugin [50]. The Trackmate
plugin [51] was used to segment mEGFP-SUF4 spots for measurement of both the
mean fluorescent intensity and mobility of condensates over time (DoG detector,
estimated object diameter 0.6µm, sub-pixel localization, simple LAP tracker:
Linking max distance 1µm, Gap-closing distance 1µm, and Gap-closing max frame
gap 1). Afterwards, easyFRAP [52] (https://easyfrap.vmnet.upatras.gr/) was used to
calculate double normalized (initial intensity, photobleaching) FRAP values among
the different variants, with each curve undergoing a double exponential curve fit
(Normalization method: Double, Exponential Equation: Double, Data collected: T-
Half, Mobile Fraction, R-Squared).
762
Live-cell imaging during temperature change
Temperature change experiments were performed on the Leica SP8 microscope.
Seedlings were placed in a 4C growth chamber for 24 hours. Then, they were

mounted in ice water and moved onto the microscope stage, where they underwent
continuous imaging for 10 minutes. As a control, room temperature seedlings were
mounted in room temperature water and continuously imaged using the same
amount of time.
770
Colocalization analysis
Co-localization of mEGFP-SUF4 condensates with FRI-mScarletI, U2B", and/or
mTFP-ELF7 was investigated by collecting multichannel 3D datasets on the
spinning disk microscope with the CSU-W1 confocal head described above.
Images were acquired with a 100x Plan-Apo 1.35 NA silicon oil immersion
objective at 0.3μm intervals on the z-axis. The Spots function and Matlab plugin
for colocalization analysis in Imaris software (RRID:SCR_007370,
URL:http://www.bitplane.com/imaris/imaris) were used to segment condensate
signal and assess co-localization. Spot diameters of 0.5μm were used to detect and
determine the locations of mEGFP-SUF4, whereas FRI and U2B" condensates
were detected with a spot diameter of 0.7μm. These diameters values were
determined by assessing the diameter that worked best for segmentation of
condensate signal using the Imaris Spots function and were largely indicative of
the typical condensate diameters. Spots were considered co-localized if the
centroids of the spots (determined at a sub-pixel scale), were closer than the radius
of the smaller Spot size used to detect and segment mEGFP-SUF4 signal (0.25
μm).
788
Puncta quantification
To quantify SUF4 nuclear condensates, we auto-cropped individual nuclei from 3D
stacks using the NucleusJ2.0 auto crop function in Fiji. Subsequently, the cropped
images were imported into LabView, where a custom signal algorithm was applied
to normalize the signal loss due to light scattering and spherical aberration from the
top to the bottom of the z-stacks. To create the dataset used for normalization, we
isolated the diffuse EGFP-SUF4 signal within the nucleoplasm away from the
brighter SUF4 puncta via a set of binary masks, one mask for total SUF4 signal
above background and second mask for the puncta. The threshold for the total
mask was set by eye, and that for the puncta was created to include only pixels
falling between 0.02% and 20% below the maximum pixel value in the section. For
each section, the puncta signal was then subtracted from the total signal to obtain
the nucleoplasm signal, and the mean nucleoplasm value per pixel calculated. The
nucleoplasm signal was roughly linear to the focal depth once the focal plane was
within the nuclear volume, enabling us to model signal loss with a linear fit to the
mean intensity in these focal sections. The pixels in planes conjugate the nuclear
volume were normalized to the inverse of the fit line. The code for TIFF file

reading was shared online by Brian Hoover(forums.ni.com), the code for TIFF
multipage saving was shared online by Nico (forums.ni.com) and used with
modifications.
809
Statistical analysis
For the analysis of flowering time and condensate numbers, a nonparametric
Mann-Whitney U test was used to test for statistically significant differences
among sets of measurements made in different genotypes. For qPCR, FRAP and
motility results, a Student's two-tailed T-test for independent means was used. To
assess the significance of the observed co-localization frequency for FRI-mScarletI
and mEGFP-SUF4 condensates, we compared this frequency to that observed for a
control protein U2B"-mRFP, which forms 1-2 bodies per nucleus of similar sizes
to those observed for FRI-mScarletI. This experiment assumes that all three
proteins are localized to the nucleosol and allows for a statistical comparison
without knowing the spatial structure and volume of the nucleosol. A two-sample
(binomial) test of proportions was used to analyze the results.
822
QPCR
qPCR was employed to assess the transcript levels of both *SUF4* and *FLC*. To
achieve this, total RNA was extracted from six-day-old seedlings (approximately
20 seedlings per genotype) using the NucleoSpin RNA plant kit (Macherey-Nagel
CO.). Subsequently, 1 microgram of total RNA was reverse transcribed using the
QuantiTect Reverse Transcription kit (Qiagen). Real-time RT-PCR was performed
on 1:10 diluted cDNA using the SensiFAST SYBR No-ROX Kit (Bioline). Three
biological and technical replicates were analyzed per genotype, and ROC1
(AT4G38740) was used as a reference gene to normalize gene expression. The
reactions were carried out with an initial denaturation step at 95°C for 2 min,
followed by 45 cycles of 95°C for 5 s, 60°C for 10 s, and a final step at 72°C for 5
s.
835
836

Primer name	Sequence	5' or 3'	Amplified gene
FLC Forward	TCTCCTCCGGCGATAACCT	5'	*FLC*
FLC Reverse	GCATGCTGTTTCCCATATCGAT	3'	*FLC*

SUF4 Forward	AGGAATTCCACCCCATGTCTTGAC	5'	SUF4
SUF4 Reverse	CTGAGATTCGTCTGTCTATCGC	3'	SUF4
Cyclo Forward	CGATAAGACTCCCAGGACTGCCGA	5'	ROC1
Cyclo Reverse	TCGGCTTTCCAGATGATGATCCAACC	3'	ROC1

837
838

Expression of His-mVenus-SUF4 in Sf9 cells

Cloning and expression of His-mVenus-SUF4 was based on the Bac-to-Bac
Baculovirus Expression system (ThermoFisher). The control pFastBac-mVenus
and pFastBac-mVenus-SUF4 isoform 1 plasmids were obtained from Dr. Alex
Chin (*15*). In these vectors, mVenus was cloned solely or with an open reading
frame of SUF4 isoform 1 (NP_564369.1; 367 amino acid) in the modified
pFastBac vector (ThermoFisher). Site-directed mutagenesis was carried out to
create the 5S and 12S mutants using PrimeSTAR Max DNA polymerase (Takara-
bio). All plasmids were sequenced and verified prior to the transformation of
DH10Bac. Bacmid preparation, transfection and infection of Sf9 cells were
performed according to the manufacturer's instruction. Large scale protein
expression was carried out by infecting and culturing the Sf9 cells with P3 viruses
for 2 days. Cells were centrifuged and either used immediately for protein
purification or frozen and stored at -80°C until use. Purification of His-mVenus-
SUF4 fusion protein from Sf9 cells. His-mVenus-SUF4 proteins were purified with
IMAC followed by gel filtration as published previously with some modifications.
All procedures were performed at 4°C. Proteins were extracted by sonicating the
cells in lysis buffer [50 mM Tris, 500 mM NaCl, 20 mM imidazole, 5 mM TCEP
(SIGMA, C4706), 5% glycerol, protease inhibitor cocktail (SIGMA, S8830) pH
7.44] supplemented with benzonase nuclease (SIGMA, E1014). The lysate was
cleared by centrifugation at 20,000 x g, for 20 min at 4°C, filtered with 0.45 μm
PVDF membrane filter (Millipore, SLHV033RS) and loaded onto Ni-NTA
columns (QIAGEN, 30230). After washing with the lysis buffer, proteins were
eluted with 500 mM imidazole (lysis buffer supplemented with 480 mM
imidazole). The eluate was concentrated using a centrifugal filter device, Amicon
Ultra-15 (Millipore, UFC901024) and loaded to a gel filtration column Superdex
200 Increase 10/300 GL operated with AKTA Pure 25L (GE Healthcare). Gel

filtration was performed in the Size Exclusion Chromatography (SEC) buffer (20
mM Hepes, 225 mM NaCl, 1 mM TCEP pH 7.4) with a constant flow rate of 0.4
ml/min. Typically, His-mVenus-SUF4 and His-mVenus proteins were eluted at
11.0 and 15.5-ml after the protein loading, respectively. Obtained fractions were
analyzed with SDS-PAGE, combined and concentrated. Protein concentration was
determined with Bradford protein assay kit (Thermo, 23200) using BSA as a
standard.
873

Turbidity assay
His-mVenus or His-mVenus-SUF4 proteins were thawed and ultracentrifuged to
remove denatured proteins. After protein concentration was determined, the protein
sample was diluted to 50, 25, 12.5, 6.25 and 3.13 µM in SEC buffer. Each sample
was mixed well with equal volume of turbidity assay buffer (20 mM Hepes, 75
mM NaCl, pH 7.4) to make a 250 µl reaction mix and transferred into a quartz
cuvette (Starna Cells, 28F-Q-10). Temperature-dependent phase transition was
detected by measuring the optical scattering at 800 nm while increasing the
temperature from 4 to 60°C (with a ramping rate of 0.5°C/min) using a
spectrophotometer Cary 300UV-VIS (Agilent Technologies). Nitrogen gas purging
was used to prevent the moisture from condensing on the cuvette surface. Three
replicates were performed for each sample.
886

Droplet formation assay
Five microliter of 50 µM proteins prepared as described above were mixed with 5
µl of turbidity assay buffer on ice and incubated for 5 min at 37°C. Each 3 µl of
protein reaction mix was loaded into a flow-chamber and immediately observed
with a confocal microscope SP5 (Leica) equipped with a 60x oil objective (NA
1.42). Imaging was performed with both DIC and fluorescence (excitation at 514
nm, emission 530 nm).
894
895

899
900
901
902

FIGURES:

Figure 1. SUF4 forms nuclear condensates that are sensitive to temperature *in planta*. A) mEGFP-SUF4 is expressed from native expression sequences in a null mutant background, *suf4* (FRI). mEGFP-SUF4 is diffuse within the nucleoplasm and forms nuclear condensates at room temperature. However, at 4°C, condensates do not form, leaving SUF4 diffused in the nucleoplasm. **B)** mEGFP-SUF4 rapidly forms condensates as temperatures increase. Scale bar: 10µm, yellow arrowheads point to the locations of nuclear condensates (filled) or locations where they will appear in later images (empty).

A

B

Figure 2. SUF4 forms temperature-sensitive condensates that may be tuned *in vitro*. A) DIC and confocal images of purified SUF4 variants. Only rescue SUF4 forms condensates at 37°C. Scale bar:5μm. B**)** Turbidity assay demonstrating the temperature sensitivity of purified SUF4 variants. The thermal set points shift depending on the severity of amino acid substitutions and the concentration used. Each graph includes three replicates for each sample, error bars depict standard errors.

959
960
961

Figure 3. Tuning SUF4 condensate behavior corresponds to changes in flowering time outcomes. A) Confocal images of SUF4 variants, illustrating varying condensation patterns. Rescue mEGFP-SUF4 forms an average of 4-5 nuclear condensates; 5S SUF4 forms 2-3 nuclear condensates on average; 12S SUF4 shows 0-1 condensates on average. Scale bar: 5μm. **B)** Quantification of panel A. Statistical analysis was performed by Mann-Whitney U test. **C)** Relative abundance of *FLC* transcripts in different SUF4 variants. Rescue SUF4 shows *FLC* transcript levels similar to WT, while various substitutions lead to graded reductions. Statistical analysis conducted using a two-tailed Student's t-test for independent means. Error bars represent the standard deviation. **D)** Image comparing the flowering time phenotypes of plants expressing various mEGFP-SUF4 variants with those of wildtype Col *(FRI)* plants. From left to right: Col *(FRI)* showing delayed flowering, *suf4* (FRI) mutant displaying early flowering, rescue mEGFP-SUF4 (functional copy of SUF4) expressed in *suf4 (FRI)* restoring the delayed flowering phenotype, 5S SUF4 (5 Y and F substitutions for S) expressed in *suf4 (FRI)* exhibiting a weak early flowering phenotype, and 12S SUF4 (12 F and Y substitutions for S) expressed in *suf4 (FRI)* showing a moderate early flowering phenotype. **E)** Quantification of panel D. Statistical analysis performed by Mann-Whitney U test. Asterisks denote specific p-values * <0.01, ** <0.001, ***<0.001, and ****<0.0001. Error bars are standard deviation.

Figure 4. SUF4 variants exhibit altered exchange rate and mobility dynamics
A) Fluorescence Recovery After Photobleaching (FRAP) analysis of SUF4
variants reveals different recovery rates. 5S SUF4 exhibits the fastest recovery rate
(t1/2=0.85 seconds), followed by rescue SUF4 (t1/2=1.27 seconds). 30I SUF4 does
not recover within the duration of the experiment. **B)** Quantification of the total
distance traveled for condensates used in panel A). On average 5S condensates
were more mobile. Asterisks denote P-values from Student's T- test * <0.01
FRAP distributions between rescue and 5S SUF4 were assessed for significance
using a two-tailed Student's t-test (p-value=0.007). Error bars= standard deviation.

A

Spots 3D detection

	FRI	U2B"
condensates	56	19
SUF4 colocalization	38	0
proportion	68%	0%
binomial test	p<0.001	

250 nm

B

SUF4+FRI+ELF7: 17% of nuclei (n=47)

1005

Figure 5. In warm temperature conditions, SUF4 co-localizes with members of
the vernalization complex in newly derived meristematic tissues. A) SUF4 and
FRI co-localize in nuclear condensates within developing lateral roots at room
temperatures. FRI is visualized by expression of mScarletI-FRI. mScarletI-FRI co-
localization appears to be selective, as U2B", a protein found in nuclear splicing
speckles, does not co-localize with SUF4. U2B" is visualized by expression of
U2B"-mRFP. Binomial tests of proportions used to test for statistical significance.
Arrowheads point to locations of tagged FRI or U2B" puncta. **B)** ELF7, a
chromatin remodeler that functions with SUF4 and FRI to transcriptionally activate
FLC, is visualized as mTFP-ELF7. mTFP-ELF7 also co-localizes within SUF4-
FRI condensates. Arrowheads point to a nuclear body with signal concentration in
all three channels. Scale bars: 5μm
1018

1019

1020

Figure 6. SUF4 and FRI have different localization patterns under cold temperatures A and B) Cold-induced FRI condensates do not co-localize with SUF4. However, these condensates can exchange positions with SUF4 nuclear condensates upon warming. **A)** The first 20 minutes of an imaging series, showcasing the phase change behavior of FRI and SUF4 as temperatures rise. Dotted circle is not the ROI used for quantification but used to mark diminishing and appearing condensates. **B)** Graph quantifying the opposing behavior of FRI and SUF4 condensation state at four locations from the same times series shown in A) (40 minutes, five time points). Error bars are standard errors. **C)** SUF4 condensates spatially limit the formation of FRI condensates in the primary root. In SUF4 variant lines lacking condensates (12S and *suf4* mutants), FRI-mScarletI condensates begin to emerge. Schematic representing the phase state of SUF4 (green circles) and FRI (magenta circles) in nuclei of each genotype. Scale bars in all panels: 5μm.

Figure 7: Proposed Model for Flowering-Time Regulation. A proposed model for the regulation of flowering time mediated by SUF4 biomolecular condensates. SUF4 (green) biomolecular condensates function to concentrate and/or recruit members of the vernalization complex, including FRI (in red), ELF7 (in blue), and other associated factors (in purple) to the *FLC* gene locus. Certain members, such as FRI, exhibit transient localization within these condensates dependent on cell state. Condensate localization ultimately leads to the transcriptional activation of the *FLC* gene, resulting in the delay in flowering time. In response to cold temperatures, SUF4 condensates undergo disassembly. This disassembly may facilitate the recruitment of a new group of chromatin remodelers and the involvement of lncRNAs, both of which work together to transcriptionally silence the *FLC* gene. Transcriptional silencing of FLC is a necessary step for plants to undergo the floral transition and flower in the subsequent spring season.

1080　**Supplementary Materials**

A

suf4 (FRI)   Col *(FRI)* $\dfrac{SUF4}{suf4\ (FRI)}$

B

1082
Supplemental Figure 1. Expression of the pSUF4::mEGFP-SUF4 transgene
rescues the *suf4* mutant phenotype **A)** Image of plants demonstrating the ability of
the pSUF4::mEGFP-SUF4 transgene to rescue the *SUF4* flowering time
phenotype. From left to right: *suf4 (FRI)* mutant plants displaying an early
flowering time phenotype, Col *(FRI)* plants displaying a WT delayed flowering
time phenotype, SUF4/*suf4 (FRI)* displaying a rescued WT delayed phenotype. **B)**
Relative abundance of *FLC* transcripts in *suf4* mutant, Col *(FRI)*, and *SUF4 (FRI)*
shown in A). Rescue SUF4 shows FLC transcript levels similar to WT. These data
are reshown in Figure 2E. Statistical analysis conducted using a Student's t-test.
Error bars represent the standard deviation. Asterisks denote the p-values* <0.01,
** <0.001, ***<0.001, and ****<0.0001.

1111

1112
1113
1114
1115 **Supplemental Figure 2.** At room temperatures, SUF4 forms biomolecular
1116 condensates in shoot cells, primary root cell, and localizes in the nucleolar cavity.
1117 Arrows point to nuclear condensates whereas stars mark the nucleolar cavity. Leaf
1118 and Root images are 3D rendered Z-stacks whereas the nucleolar cavity image is a
1119 slice from a 3D stack. Scalebar: 10 μm

1140

1141

Supplemental Figure 3. Sequence alignment of *Arabidopsis* SUF4, mouse and
Xenopus BuGZs. N-terminal domain that contains predicted NLS and zinc fingers
(highlighted in cyan), is conserved well among three SUF4/BuGZ sequences
whereas the central/C-terminal disordered domain of *Arabidopsis* SUF4 seems
diverged from animal BuGZs. F and Y residues in the predicted disordered region
are highlighted in black, blue, or purple. Other strong hydrophobic residues are
highlighted in green. Among F and Y in the predicted disordered region of SUF4,
the last 5 F and Y (highlighted in purple) were mutated to S in the 5S mutant and
additional 7 F and Y at the central region (highlighted in blue) were mutated to the
S in the 12S mutant.

1152

1153

1154

Supplemental Figure 4. Purified SUF4 forms round and reversible liquid-like
condensates *in vitro*. **A)** A representative elution profile of His-mVenus-SUF4 on
gel filtration chromatography. Recombinant proteins were expressed in Sf9 insect
cells, purified with IMAC, followed by gel filtration. Fractions were eluted around
11-ml after sample loading to enrich for the fusion proteins (indicated with red
letters in the inset of the gel image of fractions). **B)** SDS-PAGE analysis of the
purified His-mVenus-SUF4 and control HIS-mVenus proteins. Note that 12S
mutant protein has two bands on gel. **C)** Purified SUF4 forms round and reversible
liquid-like condensates *in vitro*. SUF4 can undergo multiple rounds of temperature
shifts without transitioning into a more stable gel or solid-like aggregate. Prior to
temperature shifts, SUF4 is incubated at either 20˚C to 4˚C for 2 minutes. Upon
temperature shifts every two minutes, SUF4 changes its phase state and maintains
this behavior until the next shift. Scalebar: 45μm.

1168
1169

1170

suf4 (FRI)   12S SUF4   5S SUF4   SUF4

suf4 (FRI)

1171
Supplemental Figure 5. Image of SUF4 variants in comparison to mutant
***suf4*, which flowers early.** Left to right: *suf4 (FRI)* mutant with an early flowering
time phenotype, 12S SUF4 (12 F and Y substitutions for S) with a moderate early
flowering phenotype, 5S SUF4 (5 Y and F substitutions for S) with weak early
flowering phenotype, rescue mEGFP-SUF4 (functional copy of SUF4) rescuing
the delayed flowering time phenotype.

1178

1179

A

B

Condensate # in SUF4 variants

C

D

Leaves prior to flowering in LD

D

FLC transcript levels

E

SUF4 transcript levels

Supplemental Figure 6. Characterization of 30I SUF4 in relation to the remaining SUF4 variants. **A)** Confocal images showing that 30I SUF4 assembles into more nuclear condensates than SUF4 rescue. Scale bar: 5µm. **B)** quantification of A). 30I SUF4 forms an average of ~18 condensates per nucleus). **C-D)** Leaf counting assay of different variants in long- and short-day conditions. Note that 30I SUF4 does not rescue the flowering time phenotype. Statistical analysis done using a Mann-Whitney U test. **E)** Relative abundance of *FLC* transcripts in 30I compared to different SUF4 variants. 30I SUF4 does not rescue *FLC* transcript levels. **F)** qPCR of *SUF4* in 30I and other transgenic *SUF4* variant lines. 30I SUF4 exhibits similar levels of transcripts as the rescue SUF4 line. Statistical analysis for E) and F) conducted using a two-tailed Student's t-test for independent means. Error bars represent the standard deviation. All data other than 30I SUF4 are reproduced from the main figures. Asterisks denote P-values * <0.01, ** <0.001, ***<0.001, and ****<0.0001

1211

1212

Supplemental Figure 7. A) FRI forms condensates in the primary root of 6 day old seedlings after 24 hours of 4˚C treatment. B) FRI forms condensates in a burst-like fashion in lateral roots under warm temperatures. Scalebars:10μm

1223
1224
1225

1226 **Acknowledgements:** We would like to thank Carlos Castañeda, Broder Schmidt,
1227 Adrienne Roeder, Rick Amasino, Doris Wagner, Ryan O'Hanlon, as well as the
1228 members of the Ehrhardt lab, IDPSIG, Carnegie Institution for Science, and
1229 Stanford University for their insightful comments and meaningful discussions.

1230

1231 **Funding:** H.M.M., T.H., Y.Z., and D.W.E. was supported on the Carnegie
1232 Venture Grant Fund. H.M.M. was also supported by the Simons Foundation-Life
1233 Science Research Foundation [Meyer LSRF18/Carnegie Fund 10861].
1234
1235

Author contributions:

Conceptualization: HMM, DWE, YZ

Methodology: HMM, DWE, YZ, TH, AM

Investigation: HMM, TH, DWE

Visualization: HMM, DWE

Funding acquisition: HMM, DWE, YZ

Supervision: DWE, YZ

Writing – original draft: HMM, DWE

Writing – review & editing: HMM, DWE, YZ
1245
1246

1247 **Competing interests:** Authors declare that they have no competing interests
1248

1249 **Data and materials availability:** All data are available in the main text or the
1250 supplementary materials

REFERENCES:

1. Helliwell, C.A., Wood, C.C., Robertson, M., Peacock, W.J., and Dennis, E.S. (2006). The Arabidopsis FLC protein interacts directly in vivo with SOC1 and FT chromatin and is part of a high-molecular-weight protein complex. Plant J. *46*, 183–192. 10.1111/j.1365-313x.2006.02686.x.

2. Searle, I., He, Y., Turck, F., Vincent, C., Fornara, F., Kröber, S., Amasino, R.A., and Coupland, G. (2006). The transcription factor FLC confers a flowering response to vernalization by repressing meristem competence and systemic signaling in Arabidopsis. Genes Dev. *20*, 898–912. 10.1101/gad.373506.

3. Choi, K., Kim, J., Hwang, H.-J., Kim, S., Park, C., Kim, S.Y., and Lee, I. (2011). The FRIGIDA complex activates transcription of FLC, a strong flowering repressor in Arabidopsis, by recruiting chromatin modification factors. Plant cell *23*, 289–303. 10.1105/tpc.110.075911.

4. Kim, S., Choi, K., Park, C., Hwang, H.-J., and Lee, I. (2006). SUPPRESSOR OF FRIGIDA4, Encoding a C2H2-Type Zinc Finger Protein, Represses Flowering by Transcriptional Activation of Arabidopsis FLOWERING LOCUS C. Plant Cell *18*, 2985–2998. 10.1105/tpc.106.045179.

5. Csorba, T., Questa, J.I., Sun, Q., and Dean, C. (2014). Antisense COOLAIR mediates the coordinated switching of chromatin states at FLC during vernalization. Proc. Natl. Acad. Sci. *111*, 16160–16165. 10.1073/pnas.1419030111.

6. Kim, D.-H., and Sung, S. (2017). Vernalization-Triggered Intragenic Chromatin Loop Formation by Long Noncoding RNAs. Dev. Cell *40*, 302-312.e4. 10.1016/j.devcel.2016.12.021.

7. Kik, S.K., Christopher, D., Glauninger, H., Hickernell, C.W., Bard, J.A.M., Ford, M., Sosnick, T.R., and Drummond, D.A. (2023). An adaptive biomolecular condensation response is conserved across environmentally divergent species. bioRxiv, 2023.07.28.551061. 10.1101/2023.07.28.551061.

8. Moses, D., Ginell, G.M., Holehouse, A.S., and Sukenik, S. (2023). Intrinsically disordered regions are poised to act as sensors of cellular chemistry. Trends Biochem. Sci. *48*, 1019–1034. 10.1016/j.tibs.2023.08.001.

9. Martin, E.W., and Mittag, T. (2018). Relationship of Sequence and Phase
Separation in Protein Low-Complexity Regions. Biochemistry *57*, 2478–2487.
10.1021/acs.biochem.8b00008.

10. Jiang, H., Wang, S., Huang, Y., He, X., Cui, H., Zhu, X., and Zheng, Y.
(2015). Phase Transition of Spindle-Associated Protein Regulate Spindle
Apparatus Assembly. Cell *163*, 108–122. 10.1016/j.cell.2015.08.010.

11. Wan, Y., Zheng, X., Chen, H., Guo, Y., Jiang, H., He, X., Zhu, X., and Zheng,
Y. (2015). Splicing function of mitotic regulators links R-loop–mediated DNA
damage to tumor cell killing. J Cell Biology *209*, 235–246.
10.1083/jcb.201409073.

12. Chin, A.F., Hilser, V.J., and Zheng, Y. (2019). The Disordered Protein Bugz
Conserves Mitotic Function and Liquid-Liquid Phase Separation across 1.6 Billion
Years of Evolution. Biophys J *116*, 179a–180a. 10.1016/j.bpj.2018.11.997.

13. Stępiński, D. (2014). Functional ultrastructure of the plant nucleolus.
Protoplasma *251*, 1285–1306. 10.1007/s00709-014-0648-6.

14. Shaw, P.J., and Brown, J.W. (2004). Plant nuclear bodies. Curr. Opin. Plant
Biol. *7*, 614–620. 10.1016/j.pbi.2004.09.011.

15. Kalinina, N.O., Makarova, S., Makhotenko, A., Love, A.J., and Taliansky, M.
(2018). The Multiple Functions of the Nucleolus in Plant Development, Disease
and Stress Responses. Front. Plant Sci. *9*, 132. 10.3389/fpls.2018.00132.

16. Rosa, S., Duncan, S., and Dean, C. (2016). Mutually exclusive sense–antisense
transcription at FLC facilitates environmentally induced gene repression. Nat.
Commun. *7*, 13031. 10.1038/ncomms13031.

17. Ietswaart, R., Rosa, S., Wu, Z., Dean, C., and Howard, M. (2017). Cell-Size-
Dependent Transcription of FLC and Its Antisense Long Non-coding RNA
COOLAIR Explain Cell-to-Cell Expression Variation. Cell Syst. *4*, 622-635.e9.
10.1016/j.cels.2017.05.010.

18. Kroschwald, S., and Alberti, S. (2017). Gel or Die: Phase Separation as a
Survival Strategy. Cell *168*, 947–948. 10.1016/j.cell.2017.02.029.

19. Alberti, S. (2017). The wisdom of crowds: regulating cell function through
condensed states of living matter. J. Cell Sci. *130*, 2789–2796. 10.1242/jcs.200295.

20. Wang, Z., Lou, J., and Zhang, H. (2022). Essence determines phenomenon: Assaying the material properties of biological condensates. J. Biol. Chem. *298*, 101782. 10.1016/j.jbc.2022.101782.

21. Shin, Y., and Brangwynne, C.P. (2017). Liquid phase condensation in cell physiology and disease. Science *357*. 10.1126/science.aaf4382.

22. Riback, J.A., Katanski, C.D., Kear-Scott, J.L., Pilipenko, E.V., Rojek, A.E., Sosnick, T.R., and Drummond, D.A. (2017). Stress-Triggered Phase Separation Is an Adaptive, Evolutionarily Tuned Response. Cell *168*, 1028-1040.e19. 10.1016/j.cell.2017.02.027.

23. Vazquez, D.S., Toledo, P.L., Gianotti, A.R., and Ermácora, M.R. (2022). Protein conformation and biomolecular condensates. Curr. Res. Struct. Biol. *4*, 285–307. 10.1016/j.crstbi.2022.09.004.

24. Dorone, Y., Boeynaems, S., Flores, E., Jin, B., Hateley, S., Bossi, F., Lazarus, E., Pennington, J.G., Michiels, E., Decker, M.D., et al. (2021). A prion-like protein regulator of seed germination undergoes hydration-dependent phase separation. Cell *184*, 4284-4298.e27. 10.1016/j.cell.2021.06.009.

25. Lasker, K., Boeynaems, S., Lam, V., Scholl, D., Stainton, E., Briner, A., Jacquemyn, M., Daelemans, D., Deniz, A., Villa, E., et al. (2022). The material properties of a bacterial-derived biomolecular condensate tune biological function in natural and synthetic systems. Nat. Commun. *13*, 5643. 10.1038/s41467-022-33221-z.

26. Zhu, P., Lister, C., and Dean, C. (2021). Cold-induced Arabidopsis FRIGIDA nuclear condensates for FLC repression. Nature *599*, 657–661. 10.1038/s41586-021-04062-5.

27. Zhu, P., and Dean, C. (2023). Reply to: Cold induction of nuclear FRIGIDA condensation in Arabidopsis. Nature *619*, E33–E37. 10.1038/s41586-023-06190-6.

28. Zhang, Z., Luo, X., Yang, Y., and He, Y. (2023). Cold induction of nuclear FRIGIDA condensation in Arabidopsis. Nature *619*, E27–E32. 10.1038/s41586-023-06189-z.

29. Du, M., Stitzinger, S.H., Spille, J.-H., Cho, W.-K., Lee, C., Hijaz, M., Quintana, A., and Cissé, I.I. (2024). Direct observation of a condensate effect on

super-enhancer controlled gene bursting. Cell *187*, 331-344.e17.
10.1016/j.cell.2023.12.005.

30. Beckman, W.F., Jiménez, M.Á.L., and Verschure, P.J. (2021). Transcription
bursting and epigenetic plasticity: an updated view. Epigenetics Commun. *1*, 6.
10.1186/s43682-021-00007-1.

31. Powers, S.K., Holehouse, A.S., Korasick, D.A., Schreiber, K.H., Clark, N.M.,
Jing, H., Emenecker, R., Han, S., Tycksen, E., Hwang, I., et al. (2019). Nucleo-
cytoplasmic Partitioning of ARF Proteins Controls Auxin Responses in
Arabidopsis thaliana. Mol. Cell *76*, 177-190.e5. 10.1016/j.molcel.2019.06.044.

32. Kim, S.Y., and Michaels, S.D. (2006). SUPPRESSOR OF FRI 4 encodes a
nuclear-localized protein that is required for delayed flowering in winter-annual
Arabidopsis. Development *133*, 4699–4707. 10.1242/dev.02684.

33. He, Y., Doyle, M.R., and Amasino, R.M. (2004). PAF1-complex-mediated
histone methylation of FLOWERING LOCUS C chromatin is required for the
vernalization-responsive, winter-annual habit in Arabidopsis. Genes Dev. *18*,
2774–2784. 10.1101/gad.1244504.

34. Pazos, F., Pietrosemoli, N., García-Martín, J.A., and Solano, R. (2013). Protein
intrinsic disorder in plants. Front. Plant Sci. *4*, 363. 10.3389/fpls.2013.00363.

35. Bugge, K., Staby, L., Kemplen, K.R., O'Shea, C., Bendsen, S.K., Jensen,
M.K., Olsen, J.G., Skriver, K., and Kragelund, B.B. (2018). Structure of Radical-
Induced Cell Death1 Hub Domain Reveals a Common αα-Scaffold for Disorder in
Transcriptional Networks. Structure *26*, 734-746.e7. 10.1016/j.str.2018.03.013.

36. Jiang, J., Wang, T., Wu, Z., Wang, J., Zhang, C., Wang, H., Wang, Z.-X., and
Wang, X. (2015). The Intrinsically Disordered Protein BKI1 Is Essential for
Inhibiting BRI1 Signaling in Plants. Mol. Plant *8*, 1675–1678.
10.1016/j.molp.2015.07.012.

37. Shindo, C., Aranzana, M.J., Lister, C., Baxter, C., Nicholls, C., Nordborg, M.,
and Dean, C. (2005). Role of FRIGIDA and FLOWERING LOCUS C in
Determining Variation in Flowering Time of Arabidopsis. Plant Physiol. *138*,
1163–1173. 10.1104/pp.105.061309.

38. Fang, X., Wang, L., Ishikawa, R., Li, Y., Fiedler, M., Liu, F., Calder, G.,
Rowan, B., Weigel, D., Li, P., et al. (2019). Arabidopsis FLL2 promotes liquid-

liquid phase separation of polyadenylation complexes. Nature *569*, 265–269.
10.1038/s41586-019-1165-8.

39. Roberts, S., Harmon, T.S., Schaal, J.L., Miao, V., Li, K. (Jonathan), Hunt, A.,
Wen, Y., Oas, T.G., Collier, J.H., Pappu, R.V., et al. (2018). Injectable tissue
integrating networks from recombinant polypeptides with tunable order. Nat.
Mater. *17*, 1154–1163. 10.1038/s41563-018-0182-6.

40. Quiroz, F.G., and Chilkoti, A. (2015). Sequence heuristics to encode phase
behaviour in intrinsically disordered protein polymers. Nat. Mater. *14*, 1164–1171.
10.1038/nmat4418.

41. Ruff, K.M., Roberts, S., Chilkoti, A., and Pappu, R.V. (2018). Advances in
Understanding Stimulus-Responsive Phase Behavior of Intrinsically Disordered
Protein Polymers. J. Mol. Biol. *430*, 4619–4635. 10.1016/j.jmb.2018.06.031.

42. Dzuricky, M., Rogers, B.A., Shahid, A., Cremer, P.S., and Chilkoti, A. (2020).
De novo engineering of intracellular condensates using artificial disordered
proteins. Nat Chem *12*, 814–825. 10.1038/s41557-020-0511-7.

43. Chen, J.W., Romero, P., Uversky, V.N., and Dunker, A.K. (2006).
Conservation of Intrinsic Disorder in Protein Domains and Families: I. A Database
of Conserved Predicted Disordered Regions. J. Proteome Res. *5*, 879–887.
10.1021/pr060048x.

44. Riley, A.C., Ashlock, D.A., and Graether, S.P. (2023). The difficulty of
aligning intrinsically disordered protein sequences as assessed by conservation and
phylogeny. PLOS ONE *18*, e0288388. 10.1371/journal.pone.0288388.

45. Bao, Z., Zhang, N., and Hua, J. (2014). Endopolyploidization and flowering
time are antagonistically regulated by checkpoint component MAD1 and immunity
modulator MOS1. Nat Commun *5*, 5628. 10.1038/ncomms6628.

46. Resentini, F., Cyprys, P., Steffen, J.G., Alter, S., Morandini, P., Mizzotti, C.,
Lloyd, A., Drews, G.N., Dresselhaus, T., Colombo, L., et al. (2016).
SUPPRESSOR OF FRIGIDA (SUF4) Supports Gamete Fusion via Regulating
Arabidopsis EC1 Gene Expression. Plant Physiol. *173*, 155–166.
10.1104/pp.16.01024.

47. Leene, J.V., Hollunder, J., Eeckhout, D., Persiau, G., Slijke, E.V.D., Stals, H.,
Isterdael, G.V., Verkest, A., Neirynck, S., Buffel, Y., et al. (2010). Targeted

interactomics reveals a complex core cell cycle machinery in Arabidopsis thaliana.
Mol. Syst. Biol. *6*, 397–397. 10.1038/msb.2010.53.

48. Lee, I., and Amasino, R.M. (1995). Effect of Vernalization, Photoperiod, and
Light Quality on the Flowering Phenotype of Arabidopsis Plants Containing the
FRIGIDA Gene. Plant Physiol. *108*, 157–162. 10.1104/pp.108.1.157.

49. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M.,
Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji:
an open-source platform for biological-image analysis. Nat. Methods *9*, 676–682.
10.1038/nmeth.2019.

50. Thevenaz, P., Ruttimann, U.E., and Unser, M. (1998). A pyramid approach to
subpixel registration based on intensity. IEEE Trans. Image Process. *7*, 27–41.
10.1109/83.650848.

51. Ershov, D., Phan, M.-S., Pylvänäinen, J.W., Rigaud, S.U., Blanc, L.L.,
Charles-Orszag, A., Conway, J.R.W., Laine, R.F., Roy, N.H., Bonazzi, D., et al.
(2022). TrackMate 7: integrating state-of-the-art segmentation algorithms into
tracking pipelines. Nat. Methods *19*, 829–832. 10.1038/s41592-022-01507-1.

52. Koulouras, G., Panagopoulos, A., Rapsomaniki, M.A., Giakoumakis, N.N.,
Taraviras, S., and Lygerou, Z. (2018). EasyFRAP-web: a web-based tool for the
analysis of fluorescence recovery after photobleaching data. Nucleic Acids Res.
46, W467–W472. 10.1093/nar/gky508.

1439