The bioreactor microbiome of mass cultivated marine diatoms for industrial carbon capture and utilization

Nerea Johanna Aalto[1,2], Ingeborg Hulda Giæver[1], Gunilla Kristina Eriksen[1], Linn Israelsen[1], Stina Krsmanovic[1], Sebastian Petters[1], Hans C. Bernstein[1,2]*

Affiliations:
1. Norwegian College of Fisheries Sciences, UiT - The Arctic University of Norway, Tromsø, Norway
2. The Arctic Centre for Sustainable Energy, UiT - The Arctic University of Norway, Tromsø, Norway

Correspondence: Hans C. Bernstein, UiT-The Arctic University of Norway, BFE-fak., Postboks 6050 Langnes, 9037 Tromsø, NORWAY; Phone: +47 776 46 114; Email:* Hans.C.Bernstein@uit.no

Running Title: Industrial microbial communities for decarbonization

Keywords: *Algae, Mass cultivation, Phycosphere, Community Succession, Porosira glacialis*

ABSTRACT

Marine microalgae are a promising innovation platform for carbon capture and utilization

(CCU) biotechnologies to mitigate industrial greenhouse gas emissions. However, industrial-

scale cultivation of algal mono-cultures is challenging and often unscalable. Non-axenic

microalgae in large semi-open CCU photobioreactors leads to the co-cultivation of diverse

microbial communities. There is limited knowledge about the "bioreactor ecology" involving

microalgae interacting with the microbiome and its subsequent impact on stability and

productivity. In this study, we describe the semi-continuous industrial mass cultivation of the

cold-adapted marine diatom, *Porosira glacialis* UiT201, by investigating the prokaryotic and

microeukaryotic communities. Data were collected in two consecutive time series experiments,

representing the initiation and operation of an preindustrial scale CCU photobioreactor

(300,000 liters). The first experiment experienced a culture "crash" of the focal strain after 39

days, while the second culture remained stable and "healthy" for 60 days. The results highlight

that this mass cultivation system represents a unique industrial marine microbial ecosystem.

The succession of the prokaryotic community was primarily driven by species replacement,

indicating turnover due to selective bioreactor conditions and/or biological interactions.

Nonetheless, the bioreactor consistently harbors a recurring and abundant core microbiome,

suggesting that the closely associated bacterial community is influenced by microalgae-specific

properties and can endure a dynamic and variable environment. The observed culture collapse

of *P. glacialis* coincided with changes in the core microbiome structure and different

environmental growth conditions compared to the stable and "healthy" experiment. These

findings imply that cohabiting microbial taxa within industrial microalgae cultivation likely

play a critical role in stabilizing the conversion of industrial CO_2 into marine biomass, and

changes in community structure serve as an indicator of process stability.

1 Introduction

Growing levels of atmospheric carbon have an impact on biodiversity, human health, the economy, and the global geopolitical landscape. As the world's climate nears new tipping points [1], it is imperative that we not only reduce emissions but also take new steps toward carbon capture and utilization (CCU). The state-of-the-art for CCU, however, is insufficient and constrained by finances, human capability, and limitations imposed by the requirement to replace outdated industial installations. Innovative biotechnologies that capitalize on photosynthetic carbon fixation, which naturally controls global carbon cycles, will contribute significantly to the solution [2]. Marine microalgae are specifically promising because they have a much higher carbon fixation capacity than plants. Living microalgae make up just 1-2% of global biomass, but they account for almost 50% of the world's primary production [3, 4]. The utility of microalgae for CCU is being demonstrated at an industrial installation in (Arctic) northern Norway, that harnesses cold-adapted marine diatoms to capture CO_2 at the source from an operational ferrosilicon smelter plant [5]. This is a prominent example of how innovative algal biotechnology can be deployed for CCU by taking an advantage of locally isolated diatom strains adapted to the seasonality and seawater resources available to localized large scale industry.

Economically feasible mass cultivation of microalgae for CCU must, however, overcome several barriers that include managing the microbial popluations within large industrial scale photobioreactors. Axenic mass cultivation of microalgae is both technically challenging and cost ineffective. Hence, large photobioreactors are often populated by multikingdom microbial communities and function as their own unique ecosystems [6, 7]. A better understanding of how these communities assemble and behave under stable and unstable – i.e., reactor crash – conditions is needed to scale bioreactors up to volumes capable of sufficient CO_2 fixation. Open airlift column bioreactors are effective for scaled-up industrial operation, land space utilization,

and improved CO_2 gas-to-liquid transfer [5]. However, the challenge linked to open-systems is

a constant exposure to the environment and thereby capacity of recruiting new microbial taxa

to the system [7]. The extent to which a bioreactor recruits microbial taxa from the environment

depends on how the inputs (e.g., water, nutrients, innocula) and control volume are exposed to

the environment. This presents a unique intersection between bioprocess engineering for CCU

and microbial ecology.

Coinhabiting microbial taxa within industrial photobioreactors are not necessarily harmful.

Infact, recent studies have demonstrated that the majority of the interspecific interactions

between microalgae and bacteria are mutualistic [8-10]. The beneficial associations are based

on exchange of synthesized compounds enhancing growth of both partners [11, 12]. The

invasion of other microeukaryotes instead can cause a rapid culture collapse of the focal strain

[7]. Some mass cultivation attempts used euryhaline and high pH tolerate focal strain to prevent

the contamination of competitive or predative species [13, 14]. However, extreme conditions

reduce the number of potential strains and might complicate the operational processes. To

uncover the potential and risks associated with the coexisting microbiome, a shift in focus from

focal strain-centered investigations to ones that encompass the entire biological component of

the production system is necessary [15].

Major knowledge gaps exist in our understanding of how to control or even measure the

complex interactions between microalgae and bacteria. Only few previous studies have

measured the microbial diversity of photobioreactors and none (to our knolwege) have

investigated the effects on the stability of CCU-specific operations. In order to address some of

these knowledge gaps, mass cultivation of the marine centric diatom *Porosira glacialis* UiT201

in a preindustrial scale CCU operation was combined with an investigation of the community

composition and temporal dynamics of the bioreactor microbiome. Specifically, we addressed

three questions: i) What are the most abundant prokaryotic and eukaryotic taxa mass cultivated

in cooccurrence with the focal strain? ii) What are the major succession patterns among

prokaryotic and microeukaryotic members of the bioreactor microbiome and how do the

identified temporal patterns of taxon-specific blooms relate to a variable bioreactor

environment? iii) How did the two time series experiments – one ending to an unplanned culture

crash after 39 days and one staying stable and healthy for 60 days – differ from each other and

what were the similarities with respect to microbiome structure and bioreactor growth

conditions?

2   Materials and Methods

2.1   Algae inoculum

The focal strain used to inoculate bioreactors was the centric diatom *Porosira glacialis* (UiT-

201). It was originally single-cell isolated from a sediment sample collected in the Barents Sea

(N 76° 27.54′, E 033° 03.54′) in 2014 [16]. The monoculture was maintained in a temperature

regulated incubator (Termaks KBP 6151) at 8 °C with a scalar irradiance of 65 μmol m^{-2} s^{-1}

and diluted weekly with filtered (0.45 μm) and autoclaved seawater containing inorganic

marco- and micronutrients from a plant nutrient mix YaraTera Kristalon Purple (Yara, Norway)

and natriummetasilikat 5H$_2$O (Permakem, Norway).

2.2   Photobioreactor operation

Mass cultivation of *P. glacialis* took place in northern Norway (69.22°N; 18.08°E) in two

consecutive time series experiments: 19.08-27.09 and 19.10-17.12 in 2021. Hereafter the first

time series recording is referred to as "T1-crash" due to unplanned discontinuation of

cultivation caused by contamination and collapse of *P. glacialis* culture and the second as "T2-

healthy" in which the focal strain was maintained at targeted abundance and growth rates until

the end of time series. The cultivation platform is integrated into an industrial ferrosilicon

smelting plant, Finnfjord AS, in collaboration with UiT – The Arctic University of Norway.

The platform contains four vertical column airlift photobioreactors with volumes of 6 m^3, 6 m^3,

14 m^3, and 300 m^3 [5]. This study deployed the largest 300 m^3 reactor operated with culture

volume of 150 m^3 under semicontinuous conditions. Seawater for cultivation medium was

obtained from an adjacent fjord at the depth of 25 m and filtered through a multichannel

filtration unit with 0.45 μm as a smallest cartridge pore size and through a UV-sterilization unit.

Compressed air and flue gas from the factory with CO_2 content of 3-7% were added into the

bottom of the reactor using a rotating gas dispersing device [5]. Flue gas was sparged

continuously during the cultivation periods except for three dates: 28.10., 08.12. and

14.12.2021. Illumination was provided for the reactors by ambient sunlight plus submerged

LED lights. Additional artificial illumination was provided by a collection of 200 x 50W LED

light units, placed above the culture surface of the bioreactor. Inorganic nutrients (nitrite, nitrate

and phosphate) a plant nutrient mix YaraTera Kristalon Purple (Yara, Norway), and

$Na_2SiO_3 \cdot 5H2O$ (Permakem, Norway) were added manually to the culture from the start of T1-

crash and following 9-days (19.10.-29.10.2021). Thereafter, the nutrients were automatically

added using a dosing pump to ensure continuous addition of nutrients.

2.3   Inoculation of the 300 m^3 photobioreactor

Prior to each time series experiment, the inoculation volume of *P. glaciali*s was increased in

two smaller (6 m^3 and 14 m^3) bioreactors. The 300 m^3 bioreactor was pre-filled with ~45 000 L

of filtered seawater before the inoculum was introduced. Thereafter, the volume was increased

for the next two days until a volume of 150 m^3 was reached as this was the maximum operation

capacity for the light delivery system. Semicontinuous operation was maintained by

continuously adding filtered seawater for four days a week (Monday – Friday morning) with a

mean inflow of 2100 L h^{-1}. The culture surplus was extracted from overflows (150 m^3) and

algae was harvested.

2.4   Sample collection and growth condition monitoring

Daily measurements were obtained during scheduled work shifts (Monday-Friday) from the

300 m^3 bioreactor for: temperature (C°; Endress-Hauser sensor iTHERMA ModuLine TM131

w) and dissolved oxygen (DO mg L^{-1}; WTWMulti 360 meter with CellOx 325 sensor, Xylem

Anlaytics, Welheim, Germany) from subsurface at ~1 m depth. Liquid samples for cell counts

were collected daily and for analysis of inorganic nutrients every second day. Samples were

collected twice weekly (Monday and Friday) for genomic DNA isolation and dissolved organic

carbon (DOC mg L^{-1}). These samples were also collected from the 300 m^3 bioreactor pre-filled

with filtered intake water (hereafter referred as "SW bioreactor"), from inoculum (sample from

6 m^3 and 14 m^3 was combined), and from intake water used to semicontinuously dilute the

reactor.

2.5   Sample processing and analyses

Concentration of inorganic nutrients of nitrite-nitrate (NO$_2^-$+NO$_3^-$), phosphate (PO$_4^{3-}$), and

silicate (Si(OH)$_4$) were analyzed immediately after filtration of sample through a GF/C

microfiber filter (Whatman). Nutrient specific Spectroquant kit (Merck Millipore, US) was used

and the concentration was determined spectrophotometrically using a 96-well plate with a

microplate reader (FilterMax F5, Molecular Devices, CA) in which the absorbance was

measured at 540 nm. DOC samples were filtered through 0.2 μm cellulose syringe filters

(Corning®) into acid washed and pre-combusted 60 mL glass vials and stored in dark at +4 °C

for 12 months before acidification with 250 μl of 2N HCl to determine the concentration with

TOC-VPH analyzer (Shimadzu, Japan). Modified Utermöhl method was used to obtain an

estimation of *P. glacialis* cell density. Cells from two separate transects were counted from a

4-well dish with inverted light microscope (Zeiss PrimoVert) after 2 h of settling (Nunc,

Thermo Fisher Scientific, US). Depending on the culture density up to 1 L of sample was

filtered through each (n = 5 per sampling day) 0.22 µm Sterivex polyethersulfone membrane

filter unit (Millipore) to obtain biomass for DNA extraction and following amplicon sequencing

of 16S and 18S small subunit ribosomal RNA (SSU rRNA) gene. The filters were immediately

stored at -80 °C for 3 months before DNA extraction took place at the UiT – The Arctic

University of Norway.

2.6   Microbiome – DNA extraction and Illumina amplicon sequencing

DNeasy Power Water Kit (Qiagen, Carlsbad, CA) was used for the purification of genomic

DNA. DNA was extracted from the collected biomass following the manufacturer's protocol

and stored at -80 °C until sequencing. DNA quality and quantity were assessed by the

NanoDrop™ Spectrophotometer and Qubit fluorometer (Invitrogen TM). Bacterial

communities were identified following the Earth Microbiome Project (EMP) [17] 16S Standard

Illumina library preparation protocol, using the forward-barcoded 515f (5′-3′

GTGYCAGCMGCCGCGGTAA) [18] and 806r (5′-3′ GGACTACNVGGGTWTCTAAT)

[19] primers targeting the V4 region of 16S SSU rRNA. Targeted gene sequencing of

microeukaryotes was determined following the EMP 18S Illumina Amplicon Protocol using

the universal Euk_1391f (5′-3′ AATGATACGGCGACCACCGAGATCTACAC) and the

barcoded EukBr (5′-3′ TGATCCTTCTGCAGGTTCACCTAC) primers targeting the V9

region of the 18S SSU rRNA [20, 21]. The PCR products were confirmed by gel electrophoresis

in 1% agarose gels and visualized with GelRed® staining. Sequencing was performed on an

Illumina MiSeq instrument (Illumina, San Diego, CA) at the Environmental Sample

Preparation and Sequencing Facility at Argonne National Laboratory (Lemont, IL, USA).

2.7   Microbiome – Bioinformatic pipeline

Forward and reverse Illumina reads were imported into QIIME2 and thereafter demultiplexed

using the complementary barcode file, for each sequencing run, respectively. These steps were

executed using the EMP paired-end flag. DADA2 V2021.0 algorithm within QIIME2 was used

for denoising and merging the reads [22]. During the denoising steps of the DADA2 package,

the reads were pooled, filtered and chimera-checked. Additionally, the replicated reads were

discarded. After removing the chimeric sequences and singletons, the reads of the 16S runs and

18S runs were merged, respectively. Subsequently, the DADA2 package was used for inferring

and filtering amplicon sequence variants (ASVs). Taxonomy was assigned by aligning these

ASV sequences against the SILVA V138.1 database, using a 16S and 18S rRNA self-trained

classifier [23, 24]. Classifiers were trained using RESCRIPt [25]. Although the 16S and 18S

based taxonomic classification within SILVA V138.1 database is a standardized approach, it

has limitations regarding classification and taxonomic resolution. It is noted that manual

correction of taxonomic annotation was beyond the scope of this study. Nevertheless, order

Enterobacterales was manually changed to order *Alteromonadales* as the main classified

families obtained within *Enterobacterales* in our 16S dataset were *Colwelliaceae*,

Alteromonadaceae and *Pseudoalteromonadaea* that are known to be members of

Alteromonadales [26]. The phylogenetic rooted tree, generated with QIIME2 using MAFFT

algorithm, was used for downstream diversity analysis.

2.8   Microbiome – Diversity analysis

All downstream analyses were completed using R [27]. The microbiome time-series data was

analyzed using the "microeco" package (v16.0) [28] unless another package is specifically

mentioned. Prior to community analysis, all reads that belonged to mitochondria, chloroplast

and kingdom *Eukaryota* or were unassigned (no assigned kingdom) were removed from the

16S dataset. Filtering from the 18S dataset removed ASVs assigned to the kingdoms of

Bacteria and *Archae*a and phyla of *Vertebrata*, *Arthropoda*, *Nematozoa*, *Annelida*, *Cnidaria*,

Mollusca, *Tunicate*, *Phragmoplastophyt*a and *Basidiomycota* as well as unassigned reads were.

Both datasets, 16S and 18S, contained a high number of ASVs with low number of reads. For

example, 2074 ASVs found just in 300 m^3 bioreactor comprised only 0.6% of the total reads

within 16S dataset. As we are only considering taxa with higher abundance, an abundance-filter

was used to remove rare and possibly ambiguous ASVs from the datasets. The filter-threshold

included only ASVs which mean relative abundance within the complete 16S and 18S dataset

was > 0.01%, respectively. As a result of this the number of ASVs decreased from 7736 to 430

and from 1577 to 58 within the 16S and 18S datasets, respectively. Counts of unique ASVs per

sample were used to measure richness – i.e., alpha-diversity [29]. The "multicomp" R package

was used to perform *post hoc* multiple comparison test with Tukey's HSD [30] on alpha-

diversity of different sample types between and within T1-crash and T2-healthy using

generalized linear models with Poisson distribution. Potential core prokaryotic taxa associated

with the focal strain from inoculum to the end of time series were identified applying an

occurrence filter with 75% presence treshold separately on T1-crash and T2-healthy 16S

dataset.

The compositional differences in prokaryotic and microeukaryotic communities – i.e., beta

diversity was measured with unweighted Jaccard distance metric to emphasize

presence/absence of community members. Three sample days, both from the beginning and the

end of 300 m^3 bioreactor time series, were included to infer how the bioreactor microbiome

composition in the start and end differs and how they relate to the communities in inoculum,

intake water and SW bioreactor (pre-filled 300 m^3 reactor with filtrated seawater). The

community differences were visualized in Principal Coordinate Analysis (PCoA) and the

pairwise permutational multivariate analysis of variance (PERMANOVA) was performed to

test the (dis)similarity associated to each sample type and chosen time series fraction within

and between T1-crash and T2-healthy. Beta diversity was also measured via Jaccard distances

between T1-crash and T2-healthy bioreactor time series samples, respectively. The Jaccard

distances were partitioned to account for the contributions of "nestedness" (species loss) and

"turnover" (species gain) towards the measure of total dissimilarity – i.e., $\beta_{Jac} = \beta_{Jne} + \beta_{Jtu}$ [31,

32]. This was obtained using the R package "betapart" [33]. The associated changes within total

beta diversity, turnover, and nestedness, respectively, by sampling day were tested using

"adonis" function (PERMANOVA) with 999 permutations from the R package "vegan" [34].

2.9   Microbiome – Clustering around temporal patterns

The K-medoids clustering method was depolyed following the previously described approach

applied to microbiome time series data [35, 36]. It was used to investigate shifts in microbiome

structure by examining community members that share similar succession patterns across the

crash and healthy time series. Prior to clustering the ASV counts were preprocessed through

normalization steps including variance stabilizing transformation (R package "DESeq2") [37],

detrending (R package "pracma") [38], and scaling via z-transformation to convert values on z-

score scale to ensure intercomparability between ASVs. As the K-medoids is a distance-based

clustering method pairwise Euclidean distances were calculated before each ASV was

partitioned into K clusters according to similarity criterion using the R package "cluster" [39].

The clustering efficacy – a balance between number of clusters and data overfitting – was

assessed via the Calinski-Harabasz index (Supplementary Figure S1) [40, 41].

2.10 Growth conditions of 300 m³ bioreactor – Statistics

The Mann-Whitney U test (also known as Wilcoxon rank sum test) was performed to reveal

differences in bioreactor physical and biogeochemical conditions between T1-crash and T2-

healthy. The potential association between the measured bioreactor conditions and their

relationship with identified temporal patterns (cluster medoids) was obtained using Spearman's

rank correlation analysis. Non-parametric analyses were conducted as half of the included

factors indicated non-normal distribution based on Shapiro-Wilk normality test (Supplementary

Table S1).

2.11 Data repository and reproducible analyses

The unprocessed genetic sequencing data for both SSU rRNA 16S and 18S gene amplicons is

available in ENA repository under project PRJEB66331. All data including ASV, taxonomy

and relative abundance tables, and bioreactor measurements used for analysis and graphing

along with R markdown script are available on the Open Science Framework (osf.io) as part of

this project: https://osf.io/k2egy/. All Supplementary Data referred to in this study is available

and found on https://osf.io/k2egy/.

3   Results

3.1   The bioreactor´s microbiome

Open mass cultivation of marine diatoms within the industrial CCU setting resulted in a

complex microbiome that accompanied growth of the target strain *P. glacialis*. The prokaryotic

and microeukaryotic community members were diverse and relatively dynamic during both

crashed and healthy cultivation periods. This was observed across two time series experiments

with lengths of 39 (crashed) and 60 days (healthy), respectively. The first time series – T1-crash

– ended with the loss of dominating abundance of the focal strain, *P. glacialis*, as determined

by decrease in growth and microscopy observations revealed severe increases of dead and lysed

cells. During the second time series from a mass cultivation run – T2-healthy – *P. glacialis* kept

its growth and abundance until the operation was ended and biomass was harvested after 60

days. Despite the major differences in conditions and outcomes, the observed richness,

especially for the prokaryotic component of the microbiome, across both time series

experiments shared major similarities (Figure 1). The total number of prokaryotic ASVs in T1-

crash was 339 and in T2-healthy 368 and the corresponding microeukaryotic richness was 47

and 56 ASVs, respectively. The prokaryotic richness measured in 300 m^3 bioreactor did not

differ between the T1-crash and T2-health as in total 291 and 256 unique ASVs were observed

across time (Tukey's HSD results given in Supplementary Table S2). However, only 54-128

ASVs in T1-crash and 49-113 in T2-healthy were present in each sample day which

corresponds to the diversity measured within each inoculum, but not within the SW bioreactor

(pre-filled 300 m^3 reactor with filtrated seawater) or intake water samples which comprised a

higher number of ASVs (Figure 1a, b). The microeukaryotic portion of the bioreactor

microbiome (measured via 18S rRNA gene amplicons) showed a contrasting result as compared

to the prokaryotic community (Figure 1c, d). The SW reactor and intake water samples

comprised less or corresponding number of unique ASVs than the inoculum and per day 300

m^3 bioreactor samples. Also, the mean number of observed ASVs in the bioreactor across time

was significantly different between T1-crash and T2-healthy (Tukey's HSD, $p < 0.001$;

Supplementary Table S3). In both time series richness and changes that occurred over time

were somewhat similar until day 32. Thereafter the number of observed ASVs stayed constant

in T2-healthy whereas in T1-crash a strong decrease was observed towards the end (Figure 1c,

d).

Figure 1. Alpha diversity measured as observed number of unique ASVs in each sample type and across bioreactor time series cultivations. Prokaryotic richness (16S) in time series experiment (A) crash and (B) healthy. Microeukaryotic richnes (18S) in time series experiment (C) crash and (D) healthy. The values of biological replicates (n = 4 or n = 5) are shown over boxplots. Note the different y-axis range between 16S and 18S datasets.

3.2   The reactor's prokaryotic community structure was distinct between crash versus healthy

production

When the 16S ASVs were allocated to the taxonomic level order, the prokaryotic community

in both time series studies exhibited a dominance of three groups (Figure 2a, b). Specifically,

in the T1-crash, the community was dominated by taxa classified within the order

Rhodobacterales (22%-75% of the relative abundance) until sample day-11, comprising mainly

ASVs assigned to genus *Planktotalea* (Figure 2a). In contrast, T2-healthy, *Rhodobacterales*

became the most abundant class in the end of the time series, after sample day 46 (28%-54%

of relative abundance) and was dominated by ASVs not assigned to a lower taxonomic level

(Figure 2b). Taxa indicative to the order *Flavobacteriales* were predominant, notably ASVs

assigned to genera NS9 marine group and *Winogradskyella*, for the first half of the T2-healthy

time series with exception of sample day-14, while in T1-crash the predominance of the same

genera was shorter and limited in the middle of the time series (Figure 2a, b). The third

predominant prokaryotic order was different between the time series experiments as taxa

classified within order *Chitinophagales*, mainly ASVs assigned to the family *Saprospiraceae*,

became the most abundant in the end of T1-crash (25%-64% of relative abundance) (Figure

2a). In contrast to that in T2-healthy taxa classified as *Opitutales*, comprised almost solely by

genus indicative to *Lentimonas*, had a high relative abundance in most of the sample days and

was the most abundant on sample day-14 (49% of the relative abundance) and in the middle of

the second half of the time series (Figure 2b).

3.3   Mass cultivation has an abundant core microbiome

Nearly the same number of ASVs and taxonomic assignments within prokaryotes were found

when ASVs were filtered on occurrence (i.e., presence) at a threshold of 75% among

the inoculum and 300 m^3 bioreactor samples. These were 27 and 21 in T1-crash and 25 and 22

in T2-healthy, respectively, that were present throughout the mass cultivation runs. The core

ASVs accounted for 20-50% of the unique ASVs observed each sample day in bioreactor of

both time series runs. Fourteen of the core ASVs, categorized into nine different bacterial

orders, were discovered in the bioreactor and inoculum of the two time series periods. Between

44% and 91% of the prokaryotic community's daily relative abundance in T1-crash and,

correspondingly, between 33% and 91% in T2-healthy, were the taxa that the core ASVs were

classified into (core and other ASV(s), henceforth referred to as core taxa). Supplementary

Table S4 contains a list of all the core ASVs together with the corresponding taxonomic

assignments.

3.4   The bioreactor's microeukaryotic community structure was similar until the culture

collapse in T1-crash

The focal strain, *P. glacialis*, is currently classified within class *Mediophyceae* [42], although

this classification was not given by the SILVA v138 database as 19 different ASVs indicative

to class *Mediophyceae* were assigned under the same annotation (*Mediophyceae*) down to

genus level. Among these ASVs, ASV *032943128192fa7f0a17b2d2991a72f4*, comprised 2-3

magnitude higher number of reads compared to the other ASVs. It is therefore considered that

most of the relative abundance within class *Mediophyceae* is composed by the focal strain, *P.*

glacialis in the 18S microeukaryotic communities.

The community structure of microeukaryotes at the taxonomic level class was similar between

T1-crash and T2-healthy samples until a rapid turn from *P. glacialis* dominance to

predominance of ASVs classified within the class *Bacillariophyceae* after sample day 29 (59%-

89% of relative abundance) in T1-crash (Figure 2c, d). It is notable that members of

Bacillariophyceae were present from the start in both time series with similar abundances until

strong occupation took place in T1-crash whereas in T2-healthy the relative abundance stayed

constant until the end of experiment. This contamination was caused by small pennate diatoms,

as confirmed by preliminary microscopic detection (Supplementary Figure S2). The most

abundant ASV within *Bacillariophyceae* was *f030aa085cd7c8b4bb6538425a95161c*. ASVs

assigned to the class *Bacillariophyceae* were also present in the community in the SW

bioreactor and intake water samples in the T1-crash, but to a lower degree in the T2-healthy

run, where taxa representative of the class *Chrysophyceae* were prevalent, particularly in the

intake water. However, *Chrysophyceae* did not show increasing occupation across time in the

T2-healthy microbiome (Figure 2c, d).

Figure 2. Comparison of community structure in each sample type and across bioreactor time
series cultivations. The most common prokaryotic taxa classified within taxonomic level order
in (A) crash and (B) healthy time series. The most common microeukaryotic taxa classified
within taxonomic level class in (C) crash and (D) healthy time series.

3.5   Beta diversity – microbiome community composition between sample types and across

time

The community composition of prokaryotes and microeukaryotes changed from inoculation to

the end-point in both time series experiments as inferred from pairwise PERMANOVA

comparison of the three first (start) and last (end) sample days (T1-crash: 16S, $R^2 = 0.50$ and p

< 0.01 & 18S, $R^2 = 0.61$ and $p < 0.01$; T2-healthy: 16S, $R^2 = 0.61$, $p < 0.01$ & 18S, $R^2 = 0.63$,

$p < 0.01$). The communities within start- and end-time fractions were significantly dissimilar

to those obtained in corresponding inoculum, intake water, and SW bioreactor samples and

these sample types were also dissimilar in relation to each other (Supplementary Table S5 and

S6). The Jaccard dissimilarity distances within 16S and 18S datasets, respectively, were

visualized by ordination with PCoA which showed that 39.7% and 44.0% of the total variation

among all included prokaryotic and microeukaryotic samples, respectively, were explained by

the two first principal coordinate axis (Figure 3). Closer inspection of the ordination results

indicated that in both time series experiments, the community composition in the early time

series were more similar with the parallel inoculum community, except for microeukaryotes in

T1-crash, than in the end as the communities exhibited strong differentiation across time

(Figure 3). The communities within SW bioreactor and intake water were clearly distinct from

those in inoculum and 300 m^3 bioreactor samples but also from each other, although the 300

m^3 bioreactor was pre-filled with intake water (Figure 3).

Figure 3. Beta diversity in prokaryotic and microeukaryotic commmunity composition between sample types and bioreactor time series start- and end-time fractions measured with unweighted Jaccard distances and visualized via Principal coordinate analysis. Dissimilarity within (A) prokaryotic and (B) microeukaryotic communities. Time series start-time fraction comprises cultivation days 1, 4 and 8 in crash and 1, 4 and 7 in healthy experiment. The corresponding end-time fractions are 32, 36 and 39 for crash and 53, 56 and 60 for healthy time series.

3.6 Beta diversity – contribution of species-loss and replacement to the bioreactor´s community succession patterns

Temporal variation in the community composition of the bacterial fraction of the microbiome

was driven to a greater degree by turnover (species replacement) than nestedness (species loss)

in both time series (Figure 4). This inference was obtained by analysis that disentangles the

additive contributions of these two antithetic processes to the total beta diversity (as measured

by the Jaccard distance) between all-time points across T1-crash and T2-healthy, respectively

[31]. The variance attributed to the sampling day was much lower from nestedness than from

turnover and total Jaccard in both time series as these explained over 84% and 78% of the

observed temporal variation associated with the time points, respectively, in both time series

runs (Figure 4 and Supplementary Table S7). Strong turnover is also indicated by the observed

prokaryotic richness which stayed rather constant across T1-crash and T2-healthy as under

nestedness some level of decrease would be expected through temproal species loss (Figure 1).

Figure 4. Changes in prokaryotic communities across time. Unweighted Jaccard distance measured between cultivation days in (A) crash and (B) healthy time series experiments. The total Jaccard distance (β_{jac}) is partiotioned by contributions from nestedness (β_{jne}; species loss) and turnover (β_{jtu}; species replacement) i.e., $\beta_{jac} = \beta_{jne} + \beta_{jtu}$. Lighter color represents cultivation days with high dissimilarity. Variance and correspondig significance associated with cultivation day are shown for each processes.

3.7   *Temporal succession patterns and cooccurring taxa in distinct bioreactor growth*

environments

The physical and chemical environment and cell density of *P. glacialis* in the large 300 m^3

photobioreactor was distinct with respect to measurements obtained during the T1-crash and

T2-healthy mass cultivation runs. This was established using the Mann-Whitney U test that

showed statistically significant differences between *P. glacialis* cell number and raw

fluorescence, concentration of inorganic nutrients, DOC, and temperature measured across both

time series experiments – i.e., most of the monitored factors except pH level (Supplementary

Table S9). The measured values of these factors were higher in T2-healthy compared to the T1-

crash apart from DOC concentration and bioreactor temperature which were lower (Figure 5

and Supplementary Figure S3).

Figure 5. Variation in bioreactor environment across time in crash and healthy cultivation. Daily or twice a week measurements of (A) *P. glacialis* cell density, (B) DOC concentration, (C) bioreactor temperature and (D) pH level. Note that the time series crash was ended after cultivation day-39.

The potential association of community members to the variable bioreactor environment that

prevailed across the time series experiments was evidenced through correlation with temporal

patterns. A K-medoid clustering algorithm was used to reduce the temporal complexity within

the bioreactor microbiome by identifying the major temporal patterns and revealing the related

community members within each time series (T1-crash and T2-healthy) and dataset (16S and

18S) driving these distinct patterns. The ASVs within both time series and datasets were divided

into three clusters based on assessment of the Calinski-Harabasz index which was used as a

quality metric to quantify the ratio for inter-intra-cluster variance for a different number of

clusters (Supplementary Figure S1) [41]. The observation of succession patterns among

prokaryotes as specified and represented by a medoid taxon, revealed that parts of the

community members from both time series datasets, specifically 61 ASVs in T1-crash (cluster

2) and 75 ASVs in T2-healthy (cluster 2), exhibited similar temporal dynamics with increase in

abundance in the middle of the time series (Figure 6a, b). The medoid of cluster 2 in both time

series indicated a significant positive relationship to one of the two factors measuring *P.*

glacialis cell density. Yet, the association to the other growth condition factors was not

consistent (Figure 6c, d). Also, most of the prokaryotic taxa within cluster 2 (T1-crash and T2-

healthy) belonged to the same orders *Flavobacteriales*, *Rhodobacterales*, *Opitutales* and

Enterobacterales (Figure 6e, f). Together these orders comprised the highest mean relative

abundance in both time series although cluster 2 contained only half of the number of ASVs

compared to the two other clusters in T1-crash and was not the richest cluster in T2-healthy

either. The K-medoid clustering revealed that a high number of prokaryotes within cluster 1

and 3 first decreased and thereafter increased in their abundance towards the culture crash of *P.*

glacialis in the end of T1-crash (Figure 6a). Especially in cluster 1 the temporal pattern of

medoid showed a strong inverse relationship with *P. glacialis* cell density and DOC

concentration and indicated an intensive proliferation of ASVs classified within

Rhodobacterales and *Chitinophagales* in the last two sample days as these orders comprised

the major taxonomic groups within cluster 1 (Figure 6c, d).

Figure 6. The major temporal succession patterns within prokaryotic communities as assessed via K-medoid clustering analysis with z-transformation and detrending of 16S ASVs. Temproal dynamics of prokaryotic taxa and identified number of ASVs in bioreactor (A) crash and (B)

healthy time series experiemnts. The black line indicates cluster specific medoid taxon, a
representative for cluster's temporal pattern. ASV specific z-score is a y-axis and a value 0
denotes the mean abundance. Heatmap of Spearman's correaltion between temporal dynamics
of cluster specific medoid taxon (z-scores) and measured bioreactor factors in time series (C)
crash and (D) healthy. The color indicates Spearmna's rank correlation coefficient (rho) and
the significance is denoted with ★ (★, $p \leq 0.05$; ★ ★, $p \leq 0.01$; ★ ★ ★, $p \leq 0.001$). Cells, *P.*
glacialis cell density L^{-1}; DOC, DOC mg L^{-1}; FL, raw fluorescence as a proxy for
increase/decrease of *P. glacialis* cell density; NO2-NO3, $NO_2^- + NO_3^-$ µM; pH, pH level; PO4,
PO_4^{3-} µM; Si, $Si(OH)_4$ µM, Temp, bioreactor temperature ºC. The most common prokaryotic
taxa at the taxonomic level order within each cluster and their mean relative abundance in a
whole community in time series (E) crash and (F) healthy.
504

The cluster medoids identified within 18S microeukaryotic community revealed a strong

fluctuation in relative abundances and weak correlation to bioreactor conditions across time in

both time series (Figure 7a-d). ASVs classified within class *Mediophyceae* and

Bacillariophyceae were the main groups driving distinct temporal patterns in T1-crash and T2-

healthy. In both time series experiments the class *Bacillariophyceae* belonged to cluster 3 in

which the cluster medoid exhibited strong decrease in relative abundance in the middle of time

series (Figure 7e, f).

Figure 7. The major temporal succession patterns within microeukaryotic communities as assessed via K-medoid clustering analysis with z-transformation and detrending of 16S ASVs. Temproal dynamics of prokaryotic taxa and identified number of ASVs in bioreactor (A) crash and (B) healthy cultivation experiemnts. The black line indicates cluster specific medoid taxon, a representative for cluster's temporal pattern. ASV specific z-score is a y-axis and a value 0 denotes the mean abundance. The orange line specifies temporal dynamics of the most abundant ASV classified as *Mediophyceae* representing an expected focal strain, and the light green line specifies the most abundant ASVs classified as *Bacillariophyceae*, representing a potential pennate contaminant. Heatmap of Spearman's correaltion between temporal dynamics of

cluster specific medoid taxon (z-scores) and measured bioreactor factors in time series (C) crash
and (D) healthy. The color indicates Spearmna's rank correlation coefficient (rho) and the
significance is denoted with ★ (★, p ≤ 0.05; ★★, p ≤ 0.01; ★★★, p ≤ 0.001). Cells, *P. glacialis*
cell density L^{-1}; DOC, DOC mg L^{-1}; FL, raw fluorescence as a proxy for increase/decrease of
P. glacialis cell density; NO2-NO3, NO$_2^-$ + NO$_3^-$ µM; pH, pH level; PO4, PO$_4^{3-}$ µM; Si,
Si(OH)$_4$ µM, Temp, bioreactor temperature °C. The most common microeukaryotic taxa at the
taxonomic level class within each cluster and their mean relative abundance in a whole
community in time series (E) crash and (F) healthy.
530

4   Discussion

This study was designed to measure and compare the microbial community succession patterns

of an industrial scale CCU photobioreactor during both successful and crashed operations.

While contamination control is important, it is not usually feasible to operate large scale, open

airlift photobioreactors as axenic cultures of microalgae. Hence, industrial scale production of

microalgal biomass in open systems is also mass cultivation of the associated microbial

community members. This is not necessarily a hindrance to the goals of algal CCU where pure

cultures are not required for success, if the bioreactor environment can effectively select for a

stable microbial community that promotes growth and abundant proliferation of a focal strain

capable of converting CO_2-enriched flue gas into biomass and bioproducts. This study

highlights an example of how a unique core microbiome, associated with cold-water marine

environments, stabilizes within a photobioreactor used for mass cultivation of diatoms and how

these types of microbiomes represent an important signature and possibly a controlling variable

associated with successful operation.

4.1   CCU bioreactors are unique industrial marine microbial ecosystems

This study characterized a diverse microbial ecosystem within an industrial scale

photobioreactor that was distinct from sea water inflow and inoculum (Figure 2 and 3). This is

despite the fact that the operation has been designed to promote growth of large marine diatoms

– i.e., *P. glacialis.* The microbial community structure is also sensitive to unique temporal

succession patterns that serve as signatures of success or crash. New taxa are recruited to the

system via inputs from the semicontinuous inflow of sea water from the adjacent costal bay

[43] and likely the air-water exchanges within the industrial environment. The majority of

taxonomic recruitment was observed within the prokaryotic portion of the microbiome but

observations were also made with respect to microeukaryotic taxa. Changes were observed on

the time scale of days, in a similar fashion to previous observations from temperate and coastal

cold-water microbial ecosystems [36, 44]. It is expected that the diverse community comprises

members with different ecological properties, which potentially have a strong impact on the

function of the production system in relation to CCU [15]. In addition, most of the major

bacteria groups *Flavobacteriales*, *Rhodobacterales* and orders within *Gammaproteobacteria*

found in the 300 m^3 bioreactor have frequently been reported to associate with diatom driven

phytoplankton blooms in natural ecosystems or in diatom monoculture experiments (Figure 2a,

b) [10, 29, 45, 46].

Bacteria that are commonly associated with marine diatoms are known to utilize different alga-

derived organic substrates − *i.e.,* DOC that is released across different bloom or growth stages

or generated through temporal substrate degradation [44, 47]. These dynamic processes at least

partially underpin the successional blooming patterns of the prokaryotic groups that were

identified in this study and other investigations [36, 45]. The orders *Opitutales*

(*Verrumicrobiota*) and *Chitinophagales* which also belonged to the most abundant community

ASVs from the current study are not considered either uncommon or specific to our production

system as they have also been documented in environmental and experimental studies.

Members within *Opitutales* are likely part of free-living community thriving in eutrophic

environments such as wastewater related microalgae cultivations, with a capacity to degrade

complex polysaccharides [48]. Members within the *Saprospiraceae* family (*Chitinophagales*)

have been reported to interact with diatoms [49].

4.2   Importance of a strong core microbiome in mass cultivation

The relatively long and consecutively repeated temporal coverage on microbial community composition provided us an excellent opportunity to infer reoccurring community succession processes. The bioreactor harbors a prominent core microbiome although the taxonomic composition underwent effective replacement of prokaryotes (strong turnover) between time points (Figure 4). We found that the core ASVs comprised a high portion of the total number of unique ASVs observed on each sample day in both time series. Many prokaryotic taxa that are introduced by intake water and the pre-filled bioreactor at the beginning are replaced after a while as they do not integrate into the core community, as evidenced by the strong contribution of turnover towards total community dissimilarity. Consequently, the bioreactor functions as a selective environment because of the very different physical and biogeochemical conditions that these taxa suddenly encounter in comparison to the coastal environment, such as low pH and elevated levels of light, nutrients, and DOC, or because of the core taxa's competition and antagonistic relationships, which inhibit the growth of the other taxa. It is also expected that *P. glacialis* imposes a selective force that shapes core microbiome as partially evinced by the observed changes in the prokaryotic community when small pennate diatom contaminates supplanted the *P. glacialis* focal strain. The relative strength of diatom-driven selection compared to the physical bioreactor environment remains unclear, although it is understood that the closely associated community is structured by microalgae-specific manner and can persist through environmental variation [11, 50, 51]. The high similarity in the core microbiome between the two cultivation experiments indicates for a certain level of specificity with the production system as it is designed for mass cultivation of *P. glacialis*. It is noted that at least part of the infiltered taxa may reside in the system below detection or were excluded from the analysis along with the applied abundance filter (0.01%). The core taxa (including

core and other ASVs classified within the same taxonomic assignment) accounted for an average of 69% and 77% of the relative abundance within bioreactor samples taken from T1-crash and T2-healthy, respectively. Nevertheless, on the majority of sample days, the high abundance was accounted for by a small number of core taxa, whose predominance fluctuated and was only permanent for a short while. Consequently, the temporal change in the prokaryotic community structure was significantly influenced by a small subset of core taxa. The effects of strong temporal variations in community structure and the interactions of the (core) microbiome on the productivity, stability, and/or controllability of the production system are not well understood. Increasing complexity, or the number of interactions within cooccurring relationships, might positively affect stability and, in turn, have an impact on the growth and biomass concentration of the focal strain. [52]. On the other hand, a previous study on the microbiome of marine sponges inferred that a dense core microbiome with few abundant taxa had a stronger positive impact on community level stability than more diverse assemblies of sparsely abundant taxa. [53].

4.3   *P. glacialis is a promising candidate for mass cultivation*

Despite the relatively diversified coinhabiting microbiome, *P. glacialis* grows robustly in the mass cultivation system from inoculation through semicontinuous operation of the 300 m^3 photobioreactor. While this work is the first to look at the whole community of the bioreactor, mass cultivation of *P. glacialis* for CCU has been successfully demonstrated in industrial scale trials for several years before [5]. Reactor crash events are rare and conversion of industrial CO_2 to *P. glacialis* biomass is typically successful at scale, regardless of cooccurring microeukaryotes identified in this study. This relatively novel marine diatom is a viable agent for large scale algal CCU and production of high-value biomass [16].

 4.4   Biological factors associated with reactor crash

Even with a well controlled inoculation and a robust start during mass cultivation, the reactor

can occasionally crash. One of the main obstacles when expanding the size of a bioreactor or

switching from a closed to an open cultivation system is the high input water volume delivered

into the reactor in production systems with (semi)continuous operation, which makes them

susceptible to contamination. This is a well-recognized challenge for open cultivation systems

even though bioreactor crash events are rarely documented in peer reviewed scientific literature.

The research reports on biomass production collapse or culture crash often attribute it to

contaminants exhibiting predatory behavior or resource competition [7, 14, 54]. In the present

study, the culture collapse in T1-crash was presumably not caused by predation but rather

competition or antagonist interactions from another marine diatom or a member from the

prokaryotic community, however our study design does not allow for a conclusive mechanistic

understanding of the factors that lead to the culture crash. Our findings suggest two taxa that

showed temporal blooming patterns in relation to the observed crash. Particularly, toward the

end of the T1-crash, the *Saprospiraceae* family (order *Chitinophagales*) and the order

Bacillariophyceae demonstrated strongly rising dominance within prokaryotic and

microeukaryotic communities, respectively, after sample days-22 and 29 (Figure 6a, e and 7a,

e). The only highly prevalent taxonomic order that was not a component of the T2-healthy core

microbiome was *Chitinophagales*, albeit it was only present in the second half of the T1-crash

(Figure 2a). Algicidal activity has been connected on filamentous members of *Saprospiraceae*

[55]. In a previous study *Saprospir*a-like bacteria were shown to exhibit a direct attack with the

help of its gliding motility on the diatom *Cheatoceros ceratosporum* cell causing complete lysis

of diatom cells [49]. If algicidal activity occurred against *P. glacialis* one could ask why the

pennate diatoms within *Bacillariophyceae* proliferated and seemed undegraded and why the

abundance of *Saprospiraceae* increased after three weeks of successful cultivation? As the

presence of algicidal member of *Saprospiraceae* will remain highly tentative without further

investigation another potential explanation is that *P. glacialis* in T1-crash was not able to

withstand a competition over resources against members of *Bacillariophyceae* whose

increasing presence was likely also supported by infiltration of these taxa as the community

measured in intake water was dominated by *Bacillariophyceae* (Figure 2c).

4.5 Variable environments and blooming patterns within the bioreactor

The large scale mass cultivation system is also unique in relation to the prevailing environment

created in the bioreactor and therefore difficult to replicate. This is primarily a result of seasonal

variations in the coastal environment and constant production optimization. Although the

measured growth conditions varied and were distinct between the time series experiments, the

K-medoids clustering did not reveal a clear correlation between the temporal abundance

patterns of microeukaryotes and bioreactor factors. Instead, the impact on the prokaryotic

community structure might have been stronger (Figure 6). For example, the direct impact of

lower and more variable temperature in T2-healthy than in T1-crash on the growth of focal

strain was probably minor as *P. glacialis* is shown to obtain a strong growth across the

temperature range of 2-12 ℃ [16]. Whereas the growth rate of marine bacteria is known to have

a pronounced positive relationship within the similar temperature range (White et al., 1991).

Moreover, major differences were observed on the cell density of *P. glacialis* and

biogeochemical environment. While the cell density of *P. glacialis* was clearly lower across

T1-crash than T2-healthy, the measured DOC concentration revealed the opposite (Figure 5).

The dataset does not provide a clear insight on why the DOC level was almost constantly 2.1

mg L^{-1} higher in T1-crash than in T2-healthy or how it potentially governed the cooccurring

microbial community. DOC excretion can reflect physiological state of microalgae as a

response to stress or it can be triggered by biological interactions [56-58]. The lower cell density

in this case does not directly relate to decreased growth (before culture crash) which in turn

could indicate stress, because the dilution rate was higher in T1-crash than T2-healthy.

However, the concentration of inorganic nutrients, especially silicate in the first half of the time

series, was clearly lower in T1-crash than in T2-healthy and therefore might have impacted the

cellular response of *P. glacialis* (Supplementary Figure S3).

5 Conclusion

This work shows how basic concepts of marine microbial ecology can be used to gain a better

understanding of, and perhaps even control over, a promising industrial biotechnology for CCU.

Nevertheless, it is difficult to determine the mechanistic influence of the microbiome on large-

scale biomass production in large industrial-scale photobioreactors since they are difficult and

expensive to modify and operate with high replication for only experimental purposes. Each

cultivation run of the large vertical photobioreactor is distinct in terms of the cooccurring

microbiome community dynamics and maintained growth conditions throughout the *P.*

glacialis production process because of its location within a highly seasonal coastal factory

setting. However, the reactor is home to a relatively diverse microbial ecosystem that

transforms into a core microbiome via turnover of taxa introduced by the inoculum and

seawater intake. Changes to the core microbiome are either affected by or causative crashing

populations of the *P. glacialis* strain. Hence, the reactors microbiome is a unique signature of

stable CCU operation that likely plays a critical role in stabilizing the conversion of industrial

CO_2 into marine biomass.

Author contributions

HCB and NJA conceptualized the study. IHG, GKE and LI carried out the cultivation and

sampling. SK and SP processed molecular samples and amplicon sequence data. NJA and HCB

conducted data analysis and wrote and the manuscript draft. All authors contributed to the final

version.

Aknowledgements

This project was made possible through tight collaboration and technical support from

Finnfjord AS, with specific acknolwdegements to Hans Christian Eilertsen, Jo Strømholt, John-

Steinar Bergum and Geir-Henning Wintervoll.

Funding

This work was supported by the project AlgScaleUp, funded by the Research Councile of

Norway as part of the Green Platform (NFR project number: 328654) and ABSORB – Arctic

Carbon Storage from Biomes, which is a strategic funding inniative from UiT – The Arctic

University of Norway (https://site.uit.no/absorb/).

Conflict of interest

The authors have no conflicts of interest to declare.

References

1.   Masson-Delmotte, V., et al., *IPCC Special Report: Global Warming of 1.5° C.*
Summary for Policymakers. https://www.ipcc.ch/sr15/chapter/spm/, 2018.
2.   Schweitzer, H., et al., *Innovating carbon-capture biotechnologies through ecosystem-*
inspired solutions. One Earth, 2021. **4**(1): p. 49-59.
3.   Field, C.B., et al., *Primary production of the biosphere: integrating terrestrial and*
oceanic components. science, 1998. **281**(5374): p. 237-240.
4.   Falkowski, P.G., *The role of phytoplankton photosynthesis in global biogeochemical*
cycles. Photosynthesis research, 1994. **39**: p. 235-258.
5.   Eilertsen, H.C., et al., *Mass cultivation of microalgae: I. Experiences with vertical*
column airlift photobioreactors, diatoms and CO2 sequestration. Applied Sciences,
2022. **12**(6): p. 3082.
6.   Fulbright, S.P., et al., *Bacterial community changes in an industrial algae production*
system. Algal research, 2018. **31**: p. 147-156.

7.   Davies, F.K., et al., *Microbiota associated with the large-scale outdoor cultivation of*
734   *the cyanobacterium Synechococcus sp. PCC 7002.* Algal Research, 2021. **58**: p.
735   102382.
736   8.   Amin, S.A., M.S. Parker, and E.V. Armbrust, *Interactions between Diatoms and*
737   *Bacteria.* Microbiology and Molecular Biology Reviews, 2012. **76**(3): p. 667-684.
738   9.   Durham, B.P., et al., *Cryptic carbon and sulfur cycling between surface ocean*
739   *plankton.* Proceedings of the National Academy of Sciences, 2015. **112**(2): p. 453-
740   457.
741   10.   Buchan, A., et al., *Master recyclers: features and functions of bacteria associated with*
742   *phytoplankton blooms.* Nature Reviews Microbiology, 2014. **12**(10): p. 686-698.
743   11.   Raina, J.-B., et al., *Chemotaxis shapes the microscale organization of the ocean's*
744   *microbiome.* Nature, 2022. **605**(7908): p. 132-138.
745   12.   Amin, S.A., et al., *Interaction and signalling between a cosmopolitan phytoplankton*
746   *and associated bacteria.* Nature, 2015. **522**(7554): p. 98-101.
747   13.   Meseck, S.L., *Controlling the growth of a cyanobacterial contaminant, Synechoccus*
748   *sp., in a culture of Tetraselmis chui (PLY429) by varying pH: Implications for outdoor*
749   *aquaculture production.* Aquaculture, 2007. **273**(4): p. 566-572.
750   14.   von Alvensleben, N., et al., *Salinity tolerance of Picochlorum atomus and the use of*
751   *salinity for contamination control by the freshwater cyanobacterium Pseudanabaena*
752   *limnetica.* PloS one, 2013. **8**(5): p. e63569.
753   15.   Lian, J., et al., *The effect of the algal microbiome on industrial production of*
754   *microalgae.* Microbial biotechnology, 2018. **11**(5): p. 806-818.
755   16.   Svenning, J.B., et al., *Temperature dependent growth rate, lipid content and fatty acid*
756   *composition of the marine cold-water diatom Porosira glacialis.* Algal Research,
757   2019. **37**: p. 11-16.
758   17.   Thompson, L.R., et al., *A communal catalogue reveals Earth's multiscale microbial*
759   *diversity.* Nature, 2017. **551**(7681): p. 457-463.
760   18.   Parada, A.E., D.M. Needham, and J.A. Fuhrman, *Every base matters: assessing small*
761   *subunit rRNA primers for marine microbiomes with mock communities, time series*
762   *and global field samples.* Environmental microbiology, 2016. **18**(5): p. 1403-1414.
763   19.   Apprill, A., et al., *Minor revision to V4 region SSU rRNA 806R gene primer greatly*
764   *increases detection of SAR11 bacterioplankton.* Aquatic Microbial Ecology, 2015.
765   **75**(2): p. 129-137.
766   20.   Amaral-Zettler, L.A., et al., *A method for studying protistan diversity using massively*
767   *parallel sequencing of V9 hypervariable regions of small-subunit ribosomal RNA*
768   *genes.* PloS one, 2009. **4**(7): p. e6372.
769   21.   Stoeck, T., et al., *Multiple marker parallel tag environmental DNA sequencing reveals*
770   *a highly complex eukaryotic community in marine anoxic water.* Molecular ecology,
771   2010. **19**: p. 21-31.
772   22.   Callahan, B.J., et al., *DADA2: high-resolution sample inference from Illumina*
773   *amplicon data.* Nature methods, 2016. **13**(7): p. 581-583.
774   23.   Quast, C., et al., *The SILVA ribosomal RNA gene database project: improved data*
775   *processing and web-based tools.* Nucleic acids research, 2012. **41**(D1): p. D590-D596.
776   24.   Yilmaz, P., et al., *The SILVA and "all-species living tree project (LTP)" taxonomic*
777   *frameworks.* Nucleic acids research, 2014. **42**(D1): p. D643-D648.
778   25.   Robeson, M.S., et al., *RESCRIPt: Reproducible sequence taxonomy reference*
779   *database management.* PLoS computational biology, 2021. **17**(11): p. e1009581.
780   26.   Parte, A.C., et al., *List of Prokaryotic names with Standing in Nomenclature (LPSN)*
781   *moves to the DSMZ.* International Journal of Systematic and Evolutionary
782   Microbiology, 2020. **70**(11): p. 5607.

27.   R Core Team, *R: A language and environment for statistical computing, R Foundation*
for Statistical Computing. 2021.

28.   Liu, C., et al., *microeco: an R package for data mining in microbial community*
ecology. FEMS Microbiology Ecology, 2021. **97**(2): p. fiaa255.

29.   Aalto, N.J., et al., *Diversity and selection of surface marine microbiomes in the*
Atlantic-influenced Arctic. Frontiers in Microbiology, 2022.

30.   Hothorn, T., F. Bretz, and P. Westfall, *Simultaneous inference in general parametric*
models. Biometrical Journal: Journal of Mathematical Methods in Biosciences, 2008.
50(3): p. 346-363.

31.   Baselga, A., *The relationship between species replacement, dissimilarity derived from*
nestedness, and nestedness. Global Ecology and Biogeography, 2012. **21**(12): p. 1223-
1232.

32.   Brislawn, C.J., et al., *Forfeiting the priority effect: turnover defines biofilm community*
succession. The ISME journal, 2019. **13**(7): p. 1865-1877.

33.   Baselga, A. and C.D.L. Orme, *betapart: an R package for the study of beta diversity.*
Methods in ecology and evolution, 2012. **3**(5): p. 808-812.

34.   Oksanen, J., et al., *Community ecology package.* R package version, 2013. **2**(0).

35.   Coenen, A.R., et al., *A primer for microbiome time-series analysis.* Frontiers in
genetics, 2020. **11**: p. 310.

36.   Aalto, N.J., et al., *Microbial Community Dynamics during a Harmful*
Chrysochromulina leadbeateri Bloom in Northern Norway. Applied and
Environmental Microbiology, 2023. **89**(1): p. e01895-22.

37.   Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and*
dispersion for RNA-seq data with DESeq2. Genome biology, 2014. **15**(12): p. 1-21.

38.   Borchers, H.W. and M.H.W. Borchers, *Package 'pracma'.* accessed on, 2022. **4**.

39.   Maechler, M., et al., *Package 'cluster'.* Dosegljivo na, 2013.

40.   Lord, E., et al., *Using the stability of objects to determine the number of clusters in*
datasets. Information Sciences, 2017. **393**: p. 29-46.

41.   Caliński, T. and J. Harabasz, *A dendrite method for cluster analysis.* Communications
in Statistics-theory and Methods, 1974. **3**(1): p. 1-27.

42.   Guiry, M.D. and G.M. Guiry, *AlgaeBase. World-wide electronic publication, National*
University of Ireland, Galway. http://www.algaebase.org, 2022.

43.   Aalto, N.J., et al., *Drivers of atmosphere-ocean CO2 flux in northern Norwegian*
fjords. Frontiers in Marine Science, 2021: p. 841.

44.   Teeling, H., et al., *Substrate-controlled succession of marine bacterioplankton*
populations induced by a phytoplankton bloom. Science, 2012. **336**(6081): p. 608-611.

45.   Teeling, H., et al., *Recurring patterns in bacterioplankton dynamics during coastal*
spring algae blooms. elife, 2016. **5**: p. e11888.

46.   Schäfer, H., et al., *Genetic diversity of 'satellite' bacteria present in cultures of marine*
diatoms. FEMS microbiology ecology, 2002. **42**(1): p. 25-35.

47.   Eigemann, F., et al., *Phytoplankton Producer Species and Transformation of Released*
Compounds over Time Define Bacterial Communities following Phytoplankton
Dissolved Organic Matter Pulses. Applied and Environmental Microbiology, 2023.
89(7): p. e00539-23.

48.   Orellana, L.H., et al., *Verrucomicrobiota are specialist consumers of sulfated methyl*
pentoses during diatom blooms. The ISME Journal, 2022. **16**(3): p. 630-641.

49.   Furusawa, G., et al., *Algicidal activity and gliding motility of Saprospira sp. SS98-5.*
Canadian journal of microbiology, 2003. **49**(2): p. 92-100.

50. Mönnich, J., et al., *Niche-based assembly of bacterial consortia on the diatom*
Thalassiosira rotula is stable and reproducible. The ISME Journal, 2020. **14**(6): p.
1614-1625.
51. Jackrel, S.L., et al., *Host specificity of microbiome assembly and its fitness effects in*
phytoplankton. The ISME journal, 2021. **15**(3): p. 774-788.
52. Paddock, M.B., J.D. Fernández-Bayo, and J.S. VanderGheynst, *The effect of the*
microalgae-bacteria microbiome on wastewater treatment and biomass production.
Applied microbiology and biotechnology, 2020. **104**(2): p. 893-905.
53. Björk, J.R., et al., *The dynamic core microbiome: Structure, dynamics and stability.*
BioRxiv, 2017: p. 137885.
54. Wang, Y., et al., *Identification of harmful protozoa in outdoor cultivation of Chlorella*
and the use of ultrasonication to control contamination. Algal research, 2018. **31**: p.
298-310.
55. McIlroy, S.J. and P.H. Nielsen, *The family saprospiraceae.* The Prokaryotes: Other
Major Lineages of Bacteria and The Archaea.[4th ed.], 2014: p. 863-889.
56. Mühlenbruch, M., et al., *Mini-review: Phytoplankton-derived polysaccharides in the*
marine environment and their interactions with heterotrophic bacteria. Environmental
microbiology, 2018. **20**(8): p. 2671-2685.
57. Gärdes, A., et al., *Effects of Marinobacter adhaerens HP15 on polymer exudation by*
Thalassiosira weissflogii at different N: P ratios. Marine Ecology Progress Series,
2012. **461**: p. 1-14.
58. Hulatt, C.J. and D.N. Thomas, *Dissolved organic matter (DOM) in microalgal*
photobioreactors: a potential loss in solar energy conversion? Bioresource
technology, 2010. **101**(22): p. 8690-8697.
855