Title: Transferability of ecological forecasting models to novel biotic conditions in a long-term

experimental study

Authors: Patricia Kaye T. Dumandan[1], Juniper L. Simonis[3], Glenda M. Yenni[2], S. K. Morgan

Ernest[2], and Ethan P. White[2]

Institutional Affiliations:

1. School of Natural Resources and Environment, University of Florida, Gainesville, FL,

USA

2. Department of Wildlife Ecology and Conservation, University of Florida, Gainesville,

FL, USA

3. DAPPER Stats, Portland, OR, USA

Corresponding author: S. K. Morgan Ernest (skmorgane@ufl.edu)

ABSTRACT

Ecological forecasting models play an increasingly important role for managing natural

resources and assessing our fundamental knowledge of processes driving ecological dynamics.

As global environmental change pushes ecosystems beyond their historical conditions, the utility

of these models may depend on their transferability to novel conditions. Because species

interactions can alter resource use, timing of reproduction, and other aspects of a species'

realized niche, changes in biotic conditions, which can arise from community reorganization

events in response to environmental change, have the potential to impact model transferability.

Using a long-term experiment on desert rodents, we assessed model transferability under novel

biotic conditions to better understand the limitations of ecological forecasts. We show that

ecological forecasts can be less accurate when the models generating them are transferred to

novel biotic conditions, and that the extent of model transferability can depend on the species

being forecast. We also demonstrate the importance of incorporating uncertainty in forecast

evaluation with transferred models generating less accurate and more uncertain forecasts. These

results also suggest that how a species perceives its competitive landscape can influence model

transferability, and that when uncertainties are properly accounted for, transferred models may

still be appropriate for decision making. Assessing the extent of the transferability of forecasting

models is a crucial step to increase our understanding of the limitations of ecological forecasts.

INTRODUCTION

Ecological forecasts - predictions for the future state of ecosystems - are increasingly

important for understanding, managing, and conserving natural and managed systems (Dietze et

al., 2018; Bodner et al., 2021; Lewis et al., 2023). Most ecological forecasts are based on models

that are fit to the dynamics of the ecosystem being studied. Making forecasts from these models

assumes that the behavior of the ecosystem will remain the same in the future. However, with

climate change, land use change, and the spread of invasive species, many ecosystems will be

experiencing environmental shifts making it unclear how forecasting models will perform as

altered conditions take effect (Yates et al. 2018). Deploying models in novel conditions is also

important in ecology because data limitations often require us to use information from one

ecosystem to make inferences and predictions for less well-studied ecosystems (e.g., Fitzpatrick

and Hargrove 2009, McCune 2016). Evaluating forecasting models in novel conditions can also

provide an assessment of the generality of ecological theory (Lewis et al. 2023), strengthening

our overall knowledge of how ecological systems operate. Therefore, a crucial step for

ecological forecasting, and ecology more broadly, is understanding how well models perform

under conditions that differ from those used for model development (Werkowska et al. 2017;

Yates et al., 2018; Charney et al. 2021; Lewis et al., 2023).

The effectiveness of models for making predictions under novel conditions is known as

model transferability (Randin et al. 2006). In ecology, novel conditions in space or time can

result from differences in abiotic conditions, the biotic context (e.g., the presence or abundance

of native or non-native species), or both (Radeloff et al 2015) While more studies of model

transfer are necessary in ecological forecasting (Lewis et al. 2023), initial analyses indicate that

model transferability is negatively influenced by model complexity (with more complex models

tending to generalize less successfully than simpler models; Wenger and Olden 2012, Liu et al.

2020, Chen et al. 2023), and the degree of ecological novelty (with larger differences in

environmental conditions resulting in poorer transfer; Sequeira et al 2018, Lewis et al. 2023).

While analyses related to ecological novelty often focus on abiotic conditions or coarse biotic

conditions such as habitat structure (e.g., Spence and Tingley 2020, Qiao et al 2019, Regos et al

2019), altered biotic conditions are also a potential concern for model transferability. Changes in

the biotic conditions can fundamentally alter the observed dynamics between a species and their

resources and environment (Casini et al. 2009, Tingley et al. 2014). For example, the loss of a

key species may impact the surrounding habitat and therefore the abundance of other species

(Power et al. 1996, Goheen et al. 2018), the loss of predators or competitors may relieve biotic

pressures on species allowing them to increase in abundance (e.g., Holt et al. 2008, Trewby et al

2007, Leal et al 1998), and the arrival of invasive species may dramatically depress abundances

through predation and competition that the resident species are not adapted to deal with (Wiles et

al. 2003, Gallardo et al 2016). Shifts in the strength and number of species interactions can also

impact the skill of forecasts (Daugaard et al. 2022). Thus, changes in biotic conditions can

potentially alter the transferability of forecasting models by impacting the relationship between a

species' population dynamics and its environment, even if other environmental conditions

remain unchanged. Because many environmental issues - i.e., climate-induced range shifts

(Williams and Jackson 2007), colonization of invasive species (David et al. 2017), and global

extinctions (Raffaelli 2004) - involve altered biotic conditions, understanding the impacts of

biotic changes on forecasts is critical for understanding the potential limitations of model

transferability for ecological forecasting.

Little is known about the impact of altered biotic conditions on model transferability in

forecasting because suitable data is limited (Paniw et al. 2023). Community change - caused by

extinction, colonization, or shifts in dominance - generally occurs with larger-scale changes in

abiotic environment, habitat structure, or other landscape-level alterations (Whittaker 1960,

Wiens and Rotenberry 1981, Gentry 1988). Thus, disentangling the effects of community change

on model transferability from other environmental changes requires experimental manipulations

that selectively manipulate species to generate different biotic communities experiencing the

same general environment. Most experiments are short-term, however, lasting on average one to

three years (Field et al. 2007, Magnusson 1990), which reduces the data available to both fit a

model and test the model outcomes, especially if assessing performance under natural

environmental variation is a goal. Therefore, to assess the impact of biotic shifts on model

transferability and forecast performance, long-term experimental manipulations are required.

Here, we assess model transferability under novel biotic conditions using a long-term

experiment on desert rodents in the southwestern US. For over 40 years, the Portal Project has

collected monthly data on natural and experimentally manipulated rodent communities all

experiencing the same abiotic environment. In this experiment a competitively dominant genus,

Dipodomys spp. (kangaroo rats), has been excluded resulting in significant impacts on other

species in the system (Brown 1998, Bledsoe and Ernest 2019, Diaz and Ernest 2022). Using this

unique dataset, we investigate how biotic context influences parameter estimation and prediction

accuracy when models are fit under one set of biotic conditions but used to forecast under a

novel biotic regime. Changes in model parameterizations in response to biotic conditions would

indicate that inferences about population dynamics under climate change are sensitive to the

other species present in the ecosystem, while changes in forecast accuracy would indicate

potentially important challenges for ecological forecasting.

METHODS

Rodent data

To examine whether shifting biotic conditions can impact model transferability we obtained data on rodent population dynamics from a long-term monitoring program in the Chihuahuan Desert near Portal, Arizona (Brown 1998, Ernest et al. 2018). The 20-ha study site consists of 24 50 m x 50 m plots, each enclosed with a 50 cm fence with different sized gates to manipulate rodent species access. Plots are randomly assigned to three levels of rodent community manipulation: controls (large gates, all rodents have full access to plots), kangaroo rat removals (small gates, behaviorally dominant seedeaters, *Dipodomys* spp., are excluded), and total rodent removals (no gates, all rodents excluded but occasional transient individuals occur). The rodent communities in each plot are censused monthly around the new moon using 49 Sherman traps, and basic information is collected for all trapped rodents. Further details about the experimental setup and sampling methods are discussed elsewhere (Ernest et al. 2016, Ernest et al. 2018). In this study, we only used data on the communities found in long-term (i.e., treatments maintained across all years) controls (plots 4, 11, 14, 17) and kangaroo rat removal (plots 3, 15, 19, 21). Data were obtained using the 'portalr' package (Christensen et al. 2019) and are also archived on Zenodo (10.5281/zenodo.8436468).

We used count data from long-term control and *Dipodomys* removal plots for the desert pocket mouse (*C. penicillatus*) and Bailey's pocket mouse (*C. baileyi*). We selected these species because there were extended time periods when they were relatively abundant in both control and kangaroo rat removal plots (i.e., fewer zeros which can complicate modeling) and both species respond strongly to the experimental removal of *Dipodomys* (Bledsoe and Ernest 2019,

Diaz and Ernest 2022). Previous modeling efforts (Christensen et al. 2018) found five different

community regimes at the site, so we selected the two regimes where each non-*Dipodomys*

species was highly abundant. Regime transitions are probabilistic, so we used the edge of the

range for the transition to ensure that the data was entirely within the regime and did not include

transitions between the regimes. Continued trapping at the site suggests that the 2010-2015

regime has continued and so we extended this time period to the end of 2019, shortly before an

extensive gap in data collection due to the COVID-19 pandemic. This resulted in data for *C.*

baileyi spanning from December 1999 to June 2009 (new moon number 278-396) and for *C.*

penicillatus from September 2010 to December 2019 (new moon number 411-526).

Environmental covariates data

We used environmental covariates known to be important drivers of ecological processes

in this desert ecosystem. We obtained site-level monthly data on mean air temperature (°C) and

cumulative precipitation (millimeters) that fell during warm or cool months (calculated as the

sum of precipitation that fell on days when minimum temperature was > or < 4 °C) through the

'portalr' package (Christensen et al. 2019). This data is collected by an on-site weather station

and any gaps are filled with modeled data from nearby regional weather stations (Ernest et al.

2018). Mean air temperature is a strong driver of seasonal abundance of *C. penicillatus*, and

potentially other smaller rodent species, as it influences foraging effort and seasonal activity (i.e.,

entering bouts of torpor or seasonal dormancy; Reynolds and Haskell, 1949, Meyer and Valone

1999). Lags are known to be important in the system, with one-month lags frequently capturing

the time it takes for individuals to behaviorally respond to changing environmental conditions

(Thibault et al. 2010,Clark et al., 2023)), so we used a one-month lag for the temperature

response. We used cumulative precipitation over the preceding 365-day as a covariate because

the size of granivore populations responds to precipitation-related changes in annual seed

production over the last year, with little carryover to subsequent years (Brown et al. 1979, Brown

and Heske 1990). In this ecosystem, winter and summer precipitation have different influences

on plant growth and seed production, with cool precipitation being important for the winter

annual plant community and shrub growth and establishment, and warm precipitation being

important for the summer annual and perennial plant community (for information on the two

mostly distinct annual plant communities at the site, see Ernest et al. 2018).

Modeling Approach

To assess how well forecasting models can transfer to different biotic conditions, we fit

models separately to the control plots (where kangaroo rats are present) and the *Dipodomys*

removal plots (where kangaroo rats are absent). We fit these treatment-specific models for each

species to allow us to compare the parameters of the models from the different treatments and

assess how well the models from one treatment could predict abundances on the other treatment.

The general model structure was an autoregressive model with 1 time-step and 1-year lags

plus the three environmental covariates. Each time-series model had the form:

$$Z_t \sim NegBin\,(\lambda_t\,,\phi) \qquad \text{(Eqn. 1)}$$

$\lambda_t = \beta_0 + \beta_1 \; log(Z_{t-1} + 1) + \beta_2\, log\,(Z_{t-12} + 1) + \eta_1 T_{t-1} + \eta_2 \sum_{i=t-12}^{t} \; P_{w,i} +$

$\eta_3 \sum_{i=t-12}^{t} \; P_{c,i}$ (Eqn. 2)

where Z_t, the species-specific count at time *t* is drawn from a negative binomial distribution with

parameters λ_t (the conditional mean of abundance at time *t*) and ϕ (dispersion parameter). The

negative binomial distribution was chosen to model discrete abundances while addressing

overdispersion and provides a better fit to this dataset than the Poisson (Appendix S1 Table 1).

The conditional mean was modeled as a function of an intercept (β_0), autoregressive terms for

the abundance of the previous observation ($\beta_1 \, log(Z_{t-1} + 1)$) and the abundance at the same

time in the previous year $\beta_2 \, log(Z_{t-12} + 1)$, i.e., 12 time steps), linear terms for the effects of

mean temperature of the previous month ($\eta_1 T_{t-1}$) and the annual cumulative values of warm

($\eta_2 \sum_{i=t-12}^{t} \quad P_{w,i}$) and cold ($\eta_3 \sum_{i=t-12}^{t} \quad P_{c,i}$) precipitation. One time-step and annual lags

were selected to capture short-term correlations and seasonal cycles. These choices are supported

by analyses of autocorrelation structure on both the empirical data and model residuals

(Appendix S1 Figures 6 and 7). The annual lag is less important for *C. baileyi* but was kept to

allow direct comparisons between species. The inclusion of weather data up to time *t* is realistic

in the forecasting context of this system since the weather data is collected in real-time and

automatically integrated into the dataset (White et al. 2019) making it available for predictions

for time *t*. Models were implemented in a frequentist framework using the 'tscount' package

(Liboschik et al. 2017) in R version 4.1.1 (R Core Team, 2021). We chose to only include time-

series terms based on the observed counts (using Z_{t-1} and Z_{t-12} and not using λ_{t-1} and

λ_{t-12}) to allow models to be effectively transferred. This means that "internal" and "external"

forms of the model (see Liboschik et al. 2017) are the same.

This modeling approach requires consistently sampled data (Liboschik et al. 2017), so

values for the small number of missing samples (n = 9/116 (7.8%) for *C. penicillatus* and n =

4/119 (3.4%) for *C. baileyi*) were imputed using linear interpolation. Because we trap as close as

possible to the new moon (Ernest et al. 2018) the annual periodicity of sampling is not exactly 12

periods. There are on average 12.37 new moons/year. Therefore, we also fit the models using a

13-period lag for comparison. We also evaluated models with both scaled and unscaled

covariates. The results were qualitatively similar in both cases (Appendix S2 Tables 2 and 3).

Code used to conduct analyses is archived on Zenodo (10.5281/zenodo.11053744).

To examine the relative importance of biotic conditions in driving variation in model

parameters across the time-series, sequential model fitting with rolling origins was performed to

generate a number of different forecasting models each with five years of training data (Simonis

et al. 2021). Models were fit separately for data on *C. penicillatus* and *C. baileyi* in control and

removal plots. We used the 'rsample' package (Frick et al. 2022) to conduct rolling origin

modeling on each dataset, with 60 data points (12 observations/year for 5 years) used for model

training and 12 data points (12 observations/year for 1 year) for model evaluation. This produced

45 sets of overlapping models and evaluations for *C. penicillatus* and 48 sets for *C. baileyi*.

Comparing model parameters

We compared the distribution of the coefficients estimated from each rolling origin model to

determine the degree of similarity in the coefficients between the treatment-specific datasets.

Differences in coefficients are a prerequisite for differences in forecast performance between the

two treatments when models are transferred, and quantifying any differences will provide insight

into the aspects of biology that are impacting model transfer. We compared the estimated

coefficients by quantifying the degree of overlap in the probability density estimates of each

parameter using the 'overlapping' package (Pastore et al. 2022). The resulting overlap index is

on a scale from 0 to 1, with 0 indicating distinct distributions of estimated coefficients indicating

a strong change and 1 indicating completely overlapping distributions indicating no change

(Pastore et al. 2022). This analysis combines variation between the original non-transferred and

transferred models for a single origin with variation within models among origins, providing

perspective on whether the influence of biotic conditions is sufficiently strong to be observable

even when temporal variation in parameter estimates is present. To also focus directly on the

shift in estimated coefficients in response to the experimental manipulation of biotic context, we

characterized the proportion of pairwise changes for each origin by calculating the difference in

estimated coefficients from each treatment.

We checked to make sure that the interpretability of the parameters associated with

individual environmental covariates was not unduly influenced by collinearity by performing

pairwise correlation and covariance assessments among the covariates and their parameters.

Environmental covariates used in the models had low correlations and the covariances and

correlation values of their coefficients were low (Appendix S1).

Model transfer

To assess model transferability to different biotic contexts, we generated forecasts for both

the treatment data to which the model was fit (non-transferred model) and to the data for the

other treatment (transferred model). Forecasts from transferred models (e.g., model parameters

for the removal model used to predict counts in the control plots) were made using the initial

conditions from time-series being forecast, and the model parameters for the data the model was

trained on. Similar steps were followed to generate forecasts for the non-transferred model

(where data and model were matched, e.g., control model used to predict control data).

Forecasting evaluation

We evaluated the models from each rolling origin using end-sample evaluation - where

forecasts are made for observations h time-steps past the end of the final observation in the

training time-series and evaluating on the observed test data (Simonis et al. 2021). We made

forecasts for three-time horizons (h = 1, approximately 1 month; h = 6, approximately half a

year; and h = 12, approximately 1 year) for each rolling origin. The test data for each model were

the subsequent 12 observations following each set of training data (following White et al. 2019).

We assessed accuracy of point forecasts using root mean squared error (RMSE) and forecast

uncertainty using Brier score, which is a proper scoring rule that extends the mean squared error

to distributional forecasts (Simonis et al. 2021). For each species, RMSE and Brier scores of

non-transferred and transferred models were calculated for each rolling origin model at each

forecast horizon. We then calculated the difference between the pairs of RMSE values and Brier

scores from each origin for the non-transferred and transferred models to assess the effect of

novel biotic conditions in driving forecast predictability. Negative values for RMSE and Brier

score differences indicate better forecast performance from the non-transferred model, and

positive values indicate better forecast performance from the transferred model.

RESULTS

Model parameter comparison

The two species differed in whether their model parameters were influenced by the biotic

context. For *C. penicillatus*, the estimated coefficients did not differ significantly for models fit

to data on control and removal plots, indicating similar associations between abundances and

environmental variables in both plot types (Fig 2). This is indicated by the relatively high overlap

in the distributions of most of the parameters (range of overlap coefficient: [0.83,0.90]; Fig. 2).

Pairwise comparisons of model parameters from the same origin show that most parameters did

not shift in a consistent direction (Appendix S2 Tables 1 and 2). In contrast, *C. baileyi* parameter

estimates tended to differ between models fit to data on control and removal plots, with

parameter estimates for the environmental covariates showing relatively low overlap (range of

overlap coefficient: [0.15-0.56]). Autoregressive terms, on the other hand, exhibited more

overlap (AR (1) = 0.69, AR (12) = 0.67; Fig. 2). *C. baileyi* also exhibited high proportions of

shifts in one direction for all environmental variables and the intercept (Appendix S2 Table 1 and

2). This suggests that the form of the forecasting model is dependent on the biotic context for

this species.

Model transferability under novel biotic conditions

The two species also differed in how well transferred models performed at forecasting

compared to the non-transferred models. For *C. penicillatus* the transferred models performed

similarly to the non-transferred models on both control and removal plot data (Figs. 3 and 4).

Both transferred and non-transferred models showed a consistent pattern of increasing prediction

error with increasing forecast horizon length for both RMSE and Brier score (Fig. 4). In contrast,

for *C. baileyi,* the transferred models generally performed worse than the non-transferred models

when making forecasts. Point forecast (RMSE) scores showed a clear pattern of better

performance for the non-transferred models for both control and removal data (Figs. 3 and 4).

Brier scores were also generally better for the non-transferred model, particularly on the removal

plots. However, the Brier score result was weaker when evaluating forecasts made for the control

data. While the majority of origins showed worse forecasts for the transferred model, the mode

of difference between the original and transferred model was near zero for all horizons (Fig. 4).

This suggests that there was higher uncertainty (as indicated by Brier score) in the less accurate

predictions (as indicated by RMSE) from the models transferred from the removal plots to the

control plots. Finally, similar to the *C. penicillatus* models, both models fit to *C. baileyi* data

exhibited decreasing model performance at increasing horizons (Fig. 4).

DISCUSSION

Ecological forecasts can be less accurate when models are transferred to novel biotic

contexts. In this study, we observed this effect even though the long-term experimental nature of

the Portal Project meant that plots with different species compositions were intermingled and

there was no difference in the environmental conditions between the different biotic contexts.

Decreased performance from models transferred to novel biotic conditions, however, depended

on the species being forecast, with *C. baileyi* exhibiting significant shifts in both model

parameters and forecast abundances, but *C. penicillatus* showing little change in either.

This highlights the need to account for biotic interactions in forecasting models,and the need to

understand when and why changes in the biotic context impact model transferability.

We expect changes in the biotic context to impact forecasting models if shifts in the

biotic context involve species with strong influences on the species being forecast. *C. baileyi*,

which colonized the site in 1995, increased in abundance when *Dipodomys* were experimentally

removed, demonstrating strong competitive impacts by *Dipodomys* (Ernest and Brown 2001,

Bledsoe and Ernest 2019). Our forecast models reflect this competitive impact with higher

intercepts for the removal models when compared to the controls, which causes some of the

divergence in forecasts when models are transferred. While *C. baileyi* is competitively inferior to

Dipodomys, it is larger and competitively dominant over its congener *C. penicillatus*. With the

removal of *Dipodomys*, *C. baileyi* shifts its stature in the community to that of the competitive

dominant, which increases not only its abundance but could allow *C. baileyi* to exploit regions of

its fundamental niche that overlap with *Dipodomys* (Thibault et al 2010, Diaz and Ernest 2022).

A shift in the realized niche could explain differences in environmental parameters between

control and removal plots as *C. baileyi* is no longer constrained by competition and can react

more directly to the environmental drivers. The dual effect of altered competition on the

intercept and environmental parameters then causes the transferred models to perform poorly

(Fig. 2).

Changes in biotic context do not always alter competitive hierarchies, however. Like *C.*

baileyi, *C. penicillatus* increases in abundance when *Dipodomys* spp. are removed, indicating a

strong competitive interaction between these species (Valone and Brown 1995, Bledsoe and

Ernest 2019, Diaz and Ernest 2022). With the establishment of *C. baileyi* on removal plots*,*

however, the competitive pressures on *Dipodomys* removal plots increased. In response, *C.*

penicillatus decreased their residency in the previously preferred plot (i.e., removals) and

increased their probability of dispersing to nearby control plots (Bledsoe and Ernest 2019).

Perhaps due to behavioral interactions between these territorial species, the less dominant *C.*

penicillatus exhibited shifts in abundance between plots even when *C. baileyi* abundances

decreased in 2010 (Bledsoe and Ernest 2019, Christensen et al 2019). Thus, *C. penicillatus* may

perceive competition with its close congener as being a similar competitive environment to plots

containing *Dipodomys* spp. This could explain the similarities in both the intercepts and the

environmental parameters because competitive pressures are never alleviated and *C. penicillatus*

has little opportunity to exploit unexpressed areas of its fundamental niche. Species with many

weak interactions seem to be more forecastable (Durgaard et al 2022) as changes in a single

competitor in the network are unlikely to result in a large shift in the expressed niche of the focal

species being forecast. The fact that *C. penicillatus* does not exhibit significantly different

dynamics despite the removal of *Dipodomys* highlights the challenges of understanding when

biotic context will influence ecological forecasting due to complex species networks in nature.

Declines in the accuracy of forecasts with increasing forecast horizon exhibited an

interesting interaction with model transfer to novel biotic contexts. Decreasing forecast

performance as forecasts are made further into the future is a common pattern in ecological

forecasts (Dietze et al. 2018, Harris et al. 2018) that is demonstrated by both *C. baileyi* and *C.*

penicillatus models. However, transferred models for *C. baileyi* decrease in forecast accuracy

more rapidly with the forecast horizon (as indicated by increasing deviations between the

original and transferred models, Fig. 3) . At short timescales, the strong short-term

autoregressive components in the models allow good predictions even when transferring the

model, but as the forecast horizon increases the differences in other model parameters become

more prevalent leading to greater decay in accuracy for the transferred models (Fig. 1). This

interaction suggests that estimates of decay in forecast accuracy may be overly optimistic if the

composition of the community is also shifting at the timescales of the forecasts. This lends

experimental support to the idea that estimates of model transferability need to consider multiple

aspects of transfer (Gavish et al. 2017), in this case including both transfer to novel biotic context

and transfer outside of the historical window used for fitting the models.

Differences between our two metrics for assessing forecast performance (RMSE and

Brier score) demonstrate the importance of incorporating uncertainty in forecast evaluation and

show an interesting interaction between uncertainty and model transfer to novel biotic contexts.

The RMSE, which only evaluates point estimates (not uncertainty), was worse for transferred *C.*

baileyi models on both control and removal plots, even at short horizons. The Brier score, which

integrates model uncertainty, exhibited a similar pattern for the removal plot data, but showed

reduced responses to model transfer on the control plots (Fig. 3). This difference between the

Brier score and RMSE response suggests that while the predictions from the transferred removal

models are less accurate, the uncertainty in those predictions is also higher, so the model is less

confident in the less accurate predictions. Potentially, models fit to the removals exhibit better

uncertainty under model transfer because these models are exposed to a wider range of variation

in abundance than models fit to the control plots. Due to the competitive release from *Dipodomys*

spp., *C. baileyi* abundances are typically higher and more variable in removal plots. This wider

range of variation is likely due to reduced constraints on population growth during good years

and potentially a shift in response to environmental drivers. If this increased variation is not fully

captured by the models' dynamics it will result in increased error terms and uncertainty, thus

resulting in predictions that are penalized less by evaluation metrics that include uncertainty.

This complex interaction between model transfer, uncertainty, and experimental treatment

suggests that it is important to incorporate uncertainty into the assessment of model

transferability because it can provide insights that are different from point estimates alone. It also

shows that, in some cases, transferred models may be appropriate for decision making even if

they make less accurate point forecasts, as long as the decision making properly incorporates

uncertainty. In general, evaluating uncertainty - either by using metrics that include it or by

measuring model transferability and associated forecast uncertainties - will be important for

assessing how effectively models can be transferred and their utility for implementing

conservation strategies on species or locations with limited data availability (Houlahan et al.

2017, Yates et al. 2018).

In this study, we focused on single species models to demonstrate and assess model

transferability under varying biotic conditions. Single species models are common in ecological

modeling, forecasting, and management, but because they do not attempt to model species

interactions these models are likely to be particularly susceptible to changes in the biotic context.

Multivariate community models, which can include species interactions, have the potential to

provide improved transfer to novel biotic conditions by incorporating information on processes

such as competition. For example, for the control plots this type of model could include the

interactions between *C. baileyi* and *Dipodomys* species, potentially allowing it to transfer more

effectively to the removal plots where *Dipodomys* abundance would influence predictions as an

observed value at or near zero. The use of these types of models in dynamic ecological

forecasting remains uncommon since the number of ecosystems with sufficiently long time-

series on all of the key species in the community is limited. Since explicitly modeling

interactions is important for modeling population dynamics (e.g., Lima et al. 2008), species

distributions (e.g., Pollock et al. 2014), and model transferability, further exploration of

multivariate community predictions will be an important next step for ecological forecasting.

We have shown that changes in the presence of other species can impact both the

parameters of ecological forecasting models and their predictions. This suggests that caution will

be necessary when making forecasts in new systems or over long enough periods of time that the

composition of other species in the community undergoes change. This is important because the

development of ecological forecasting models for new scenarios is often limited by data

availability. Transferring models to new scenarios, using either no new data or a small amount

of new data to fine-tune the model, is a common approach to addressing this challenge

(Houlahan et al. 2017, Yates et al. 2018, Lewis et al. 2023). Therefore, models that better

represent the complex dynamics of biological interactions, and effectively predict beyond the

conditions they were built on, are needed in an era of fast-paced environmental change (Yates et

al. 2018). Developing models that can be effectively transferred across space, time, and biotic

context, and effectively communicate the uncertainties in their predictions, are important

endeavors to facilitate the expanded development and use of ecological forecasts (Houlahan et al.

2017, Yates et al. 2018).

ACKNOWLEDGMENTS

We thank the volunteers, staff, and research assistants who collected the Portal Project data over

the years. This work was supported by research grants awarded to SKM Ernest and EP White by

the US National Science Foundation (NSF DEB-1929730), USDA National Institute of Food and

Agriculture, Hatch Project FLA-WEC-005983 (Ernest) and FLA-WEC-005944 (White).

CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest.

REFERENCES

Bledsoe, E.K., and S. K. M. Ernest. 2019. "Temporal Changes in Species Composition Affect a

Ubiquitous Species' Use of Habitat Patches." *Ecology* 100(11):e02869. doi:

10.1002/ecy.2869.

Bodner, K., C. R. Firkowski, J. R. Bennett, C. Brookson, M. Dietze, S. Green, J. Hughes, J.

Kerr,et. al. 2021. "Bridging the Divide between Ecological Forecasts and Environmental

Decision Making." *Ecosphere* 12(12):e03869. doi: 10.1002/ecs2.3869.

Brown, J. H., D. W. Davidson, and O. J. Reichman. 1979. "An Experiment Study of Competition

between Seed-eating Desert Rodents and Ants". *American Zoologist* 19(4): 1129-1143.

Brown, James H. 1998. "The desert granivory experiments at Portal." *Experimental Ecology: Issues and Perspectives. Oxford University Press, New York*: 71-95.

Brown, J. H., and E. J. Heske. 1990. Control of a desert-grassland transition by a keystone rodent guild. *Science* 250 (4988): 1705-1707.

Casini, M., J. Hjelm, J. Molinero, J. Lövgren, M. Cardinale, V. Bartolino, A. Belgrano, et al. 2009. "Trophic Cascades Promote Threshold-like Shifts in Pelagic Marine Ecosystems." *Proceedings of the National Academy of Sciences* 106(1):197–202. doi: 10.1073/pnas.0806649105.

Charney, N. D., S. Record, B. E. Gerstner, C. Merow, P. L. Zarnetske, and B. J. Enquist. 2021. "A Test of Species Distribution Model Transferability Across Environmental and Geographic Space for 108 Western North American Tree Species." *Frontiers in Ecology and Evolution* 9.

Chen, X., Y. Liang, and X. Feng. 2024. "Influence of Model Complexity, Training Collinearity, Collinearity Shift, Predictor Novelty and Their Interactions on Ecological Forecasting." *Global Ecology and Biogeography* 33 (3): 371–84. https://doi.org/10.1111/geb.13793.

Christensen, E. M., D. J. Harris, and S. K. M. Ernest. 2018. "Long-Term Community Change through Multiple Rapid Transitions in a Desert Rodent Community." *Ecology* 99(7):1523–29. doi: 10.1002/ecy.2373.

Christensen, E. M., Yenni, G. M., Ye, H., Simonis, J. L., Bledsoe, E. K., Diaz, R., ... & Ernest, S. M. (2019). portalr: an R package for summarizing and using the Portal Project Data. *Journal of Open Source Software*, *4*(33): 1098.

Clark, N. J., S. K. M. Ernest, H. Senyondo, J. Simonis, E. P. White, G. M. Yenni, and K. Karunarathna. 2023. "Multi-Species Dependencies Improve Forecasts of Population Dynamics in a Long-Term Monitoring Study," https://ecoevorxiv.org/repository/view/5143/.

Daugaard, U., S. B. Munch, D. Inauen, F. Pennekamp, and O. L. Petchey. 2022. "Forecasting in

the Face of Ecological Complexity: Number and Strength of Species Interactions Determine

Forecast Skill in Ecological Communities." *Ecology Letters* 25(9):1974–85. doi:

10.1111/ele.14070.

David, P., E. Thébault, O. Anneville, P.-F. Duyck, E. Chapuis, and N. Loeuille. 2017. "Impacts

of Invasive Species on Food Webs: A Review of Empirical Data." *Advances in Ecological*

Research 56 (January): 1–60. https://doi.org/10.1016/bs.aecr.2016.10.001.

Diaz, R.M., and S. K. M. Ernest. 2022. "Maintenance of Community Function through

Compensation Breaks down over Time in a Desert Rodent Community." *Ecology*

103(7):e3709. doi: 10.1002/ecy.3709.

Dietze, M. C., A. Fox, L. M. Beck-Johnson, J. L. Betancourt, M. B. Hooten, C. S. Jarnevich, T.

H. Keitt, et al. 2018. "Iterative Near-Term Ecological Forecasting: Needs, Opportunities, and

Challenges." *Proceedings of the National Academy of Sciences* 115(7):1424–32. doi:

10.1073/pnas.1710231115.

Ernest, S. K. M., and J. H. Brown. 2001. Delayed compensation for missing keystone species by

colonization. *Science* 292 (5514): 101-104.

Ernest, S. K. M., G. M. Yenni, G. Allington, E. M. Christensen, K. Geluso, J. R. Goheen, M. R.

Schutzenhofer, et al. 2016. "Long-Term Monitoring and Experimental Manipulation of a

Chihuahuan Desert Ecosystem near Portal, Arizona (1977–2013)." *Ecology* 97 (4): 1082–

1082. https://doi.org/10.1890/15-2115.1.

Ernest, S. K. M., G. M. Yenni, G. Allington, E. K. Bledsoe, E. M. Christensen, R. M. Diaz, K.

Geluso, et al. 2018. "The Portal Project: A Long-Term Study of a Chihuahuan Desert

Ecosystem." https://doi.org/10.1101/332783.

Field, S. A., P. J. O'Connor, A. J. Tyre, and H. P. Possingham. 2007. "Making Monitoring

Meaningful." *Austral Ecology* 32(5):485–91. doi: 10.1111/j.1442-9993.2007.01715.x.

Fitzpatrick, M. C., and W. W. Hargrove. 2009. "The Projection of Species Distribution Models

and the Problem of Non-Analog Climate." *Biodiversity and Conservation* 18(8):2255–61. doi:

10.1007/s10531-009-9584-8.

Frick H, F. Chow, K. M, M. Mahoney, S. J., and H. Wickham. 2022. "rsample: General

Resampling Infrastructure". R package version 1.1.1, <https://CRAN.R-

project.org/package=rsample>.

Gallardo, B., M. Clavero, M. I. Sánchez, and M. Vilà. 2016. "Global Ecological Impacts of

Invasive Species in Aquatic Ecosystems." *Global Change Biology* 22(1):151–63. doi:

10.1111/gcb.13004.

Gavish, Y., J. O'Connell, and T. G. Benton. 2018. "Quantifying and Modelling Decay in

Forecast Proficiency Indicates the Limits of Transferability in Land-Cover Classification."

Methods in Ecology and Evolution 9(2):235–44. doi: 10.1111/2041-210X.12870.

Gentry, A.H., 1988. Changes in plant community diversity and floristic composition on

environmental and geographical gradients. *Annals of the Missouri Botanical Garden*, pp.1-34.

Goheen, J. R., D. J. Augustine, K. E. Veblen, D. M. Kimuyu, T. M. Palmer, L. M. Porensky, R.

M. Pringle, et al. 2018. "Conservation Lessons from Large☐mammal Manipulations in East

African Savannas: The KLEE, UHURU, and GLADE Experiments." *Annals of the New York*

Academy of Sciences 1429(1):31–49. doi: 10.1111/nyas.13848.

Harris, D. J., S. D. Taylor, and E. P. White. 2018. "Forecasting biodiversity in breeding birds

using best practices." *PeerJ* 6:e4278.

Holt, A. R., Zoe G. Davies, C. Tyler, and S. Staddon. 2008. "Meta-Analysis of the Effects of

Predation on Animal Prey Abundance: Evidence from UK Vertebrates." *PLOS ONE*

3(6):e2400. doi: 10.1371/journal.pone.0002400.

Houlahan, Jeff E., Shawn T. McKinney, T. Michael Anderson, and Brian J. McGill. 2017. "The

Priority of Prediction in Ecological Understanding." *Oikos* 126(1):1–7. doi:

10.1111/oik.03726.

Leal, M., J. Rodríguez-Robles, and J. Losos. 1998. "An Experimental Study of Interspecific

Interactions between Two Puerto Rican Anolis Lizards." *Oecologia* 117:273–78. doi:

10.1007/s004420050658.

Lewis, A. S. L., C. R. Rollinson, A. J. Allyn, J. Ashander, S. Brodie, C. B. Brookson, E. Collins,

et al. 2023. "The Power of Forecasts to Advance Ecological Theory." *Methods in Ecology and*

Evolution 14(3):746–56. doi: 10.1111/2041-210X.13955.

Liboschik T, K. Fokianos, and K. Fried. 2017. "tscount: An R Package for Analysis of Count

Time Series Following Generalized Linear Models." *Journal of Statistical Software* 82 (5): 1-

51. doi: 10.18637/jss.v082.i05

Lima, M., S. K. M. Ernest, J. H. Brown, A. Belgrano, and N. C. Stenseth. 2008. "Chihuahuan

desert kangaroo rats: Nonlinear Effects of Population Dynamics, Competition, And Rainfall."

Ecology 89(9):2594–2603. doi: 10.1890/07-1246.1.

Liu, C., C. Wolter, W. Xian, and J. M. Jeschke. 2020. "Species Distribution Models Have

Limited Spatial Transferability for Invasive Species." *Ecology Letters* 23(11):1682–92. doi:

10.1111/ele.13577.

McCune, J. L. 2016. "Species Distribution Models Predict Rare Species Occurrences despite

Significant Effects of Landscape Context." *Journal of Applied Ecology* 53(6):1871–79. doi:

10.1111/1365-2664.12702.

Meyer, M. D., and T. J. Valone. 1999. "Foraging under multiple costs: the importance of

predation, energetic, and assessment error costs to a desert forager." *Oikos* 571:579.

Paniw, M., D. García-Callejas, F. Lloret, R. D. Bassar, J. Travis, and O. Godoy. 2023. "Pathways

to Global-Change Effects on Biodiversity: New Opportunities for Dynamically Forecasting

Demography and Species Interactions." *Proceedings of the Royal Society B: Biological*

Sciences 290(1993):20221494. doi: 10.1098/rspb.2022.1494.

Pastore, M, P. A. D. Loro, M. Mingione, A. Calcagni. 2022. "Overlapping: Estimation of

Overlapping in Empirical Distributions". R package version 2.1, <https://CRAN.R-

project.org/package=overlapping>.

Pollock, L. J., R. Tingley, W. K. Morris, N. Golding, R. B. O'Hara, K. M. Parris, P. A. Vesk,

and M. A. McCarthy. 2014. "Understanding Co-Occurrence by Modelling Species

Simultaneously with a Joint Species Distribution Model (JSDM)." *Methods in Ecology and*

Evolution 5(5):397–406. doi: 10.1111/2041-210X.12180.

Power, M. E., D. Tilman, J. A. Estes, B. A. Menge, W. J. Bond, L. S. Mills, G. Daily, et al. 1996.

"Challenges in the Quest for Keystones." *BioScience* 46(8):609–20. doi: 10.2307/1312990.

R Core Team. 2021. "R: A language and environment for statistical computing". *R Foundation*

for Statistical Computing. Vienna, Austria. http://www.R-project.org

Raffaelli, D., 2004. How extinction patterns affect ecosystems. *Science*, 306(5699), pp.1141-

1142.

Randin, C. F., T. Dirnböck, S. Dullinger, N. E. Zimmermann, M. Zappa, and A. Guisan. 2006.

"Are Niche☐based Species Distribution Models Transferable in Space?" *Journal of*

Biogeography 33(10):1689–1703. doi: 10.1111/j.1365-2699.2006.01466.x.

Regos, A., L. Gagne, D. Alcaraz-Segura, J. P. Honrado, and J. Domínguez. 2019. "Effects of

Species Traits and Environmental Predictors on Performance and Transferability of

Ecological Niche Models." *Scientific Reports* 9(1):4221. doi: 10.1038/s41598-019-40766-5.

Reynolds, H.G., and H. S. Haskell. 1949. "Life history notes on Price and Bailey pocket mice of

southern Arizona." *Journal of Mammalogy* 30 (2): 150-156.

Spence, A. R., and M. W. Tingley. 2020. "The Challenge of Novel Abiotic Conditions for

Species Undergoing Climate-Induced Range Shifts." *Ecography* 43(11):1571–90. doi:

10.1111/ecog.05170.

Thibault, K. M., S. K. M. Ernest, E. P. White, J. H. Brown, and J. R. Goheen. 2010. "Long-Term

Insights into the Influence of Precipitation on Community Dynamics in Desert Rodents."

Journal of Mammalogy 91(4): 787–97. https://doi.org/10.1644/09-MAMM-S-142.1.

Tingley, R., M. Vallinoto, F. Sequiera, and M. R. Kearney. 2014. "Realized Niche Shift during a

Global Biological Invasion." 2014. https://doi.org/10.1073/pnas.1405766111.

Trewby, I. D., G. J. Wilson, R. J. Delahay, N. Walker, R. Young, J. Davison, C. Cheeseman, P.

A. et al. 2008. "Experimental Evidence of Competitive Release in Sympatric Carnivores."

Biology Letters 4(2):170–72. doi: 10.1098/rsbl.2007.0516.

Valone, T. J., and J. H. Brown. 1995. "Effects of Competition, Colonization, and Extinction on

Rodent Species Diversity." *Science* 267(5199):880–83.

Wenger, S. J., and J. D. Olden. 2012. "Assessing Transferability of Ecological Models: An

Underappreciated Aspect of Statistical Validation." *Methods in Ecology and Evolution*

3(2):260–67. doi: 10.1111/j.2041-210X.2011.00170.x.

Werkowska, W., A. L. Márquez, R. Raimundo, and P. Acevedo. 2017. "A Practical Overview of

Transferability in Species Distribution Modeling." *Environmental Reviews* 25 (1): 127-133.

Wiens, J.A. and J.T Rotenberry. 1981. Habitat associations and community structure of birds in

shrubsteppe environments. *Ecological Monographs* 51(1):21-42.

White, E. P., G. M. Yenni, S. D. Taylor, E. M. Christensen, E. K. Bledsoe, J. L. Simonis, and S.

K. M. Ernest. 2019. "Developing an Automated Iterative Near-Term Forecasting System for

an Ecological Study." *Methods in Ecology and Evolution* 10(3):332–44. doi: 10.1111/2041-

210X.13104.

Whittaker, R.H., 1960. Vegetation of the Siskiyou mountains, Oregon and California. *Ecological*

Monographs 30(3):279-338.

Wiles, G. J., J. Bart, R. E. Beck, and C. F. Aguon. 2003. "Impacts of the Brown Tree Snake:

Patterns of Decline and Species Persistence in Guam's Avifauna." *Conservation Biology*

17(5):1350–60. doi: 10.1046/j.1523-1739.2003.01526.x.

Williams, J.W. and Jackson, S.T., 2007. Novel climates, no□analog communities, and ecological

surprises. *Frontiers in Ecology and the Environment* 5(9): 475-482.

Yates, K. L., P. J. Bouchet, M. J. Caley, K. Mengersen, C. F. Randin, S. Parnell, A. H. Fielding,

et al. 2018. "Outstanding Challenges in the Transferability of Ecological Models." *Trends in*

Ecology & Evolution 33(10):790–802. doi: 10.1016/j.tree.2018.08.001.

 FIGURE LEGENDS

Figure 1. Time-series data on *Dipodomys* spp. (yellow lines), *Chaetodipus baileyi* (purple lines),

and *C. penicillatus* (green lines) on control plots (top panel) and removal plots (bottom panel) in

a long-term experiment near Portal, AZ. The two species-specific periods used for modeling and

forecasting *Chaetodipus* spp. are indicated by brackets.

Figure 2. Probability density estimates of the distribution of estimated coefficients (point

estimates) obtained from the suite of rolling origin models fit to data on *C. penicillatus* (top row;

n = 45) and *C. baileyi* (bottom row; n = 48) on control plots (purple densities) and removal plots

(yellow densities).

Figure 3. Predictions for *C. penicillatus* (top two plots) and *C. baileyi* (bottom two plots)

abundances from models fit to non-transferred (blue lines) and transferred (red lines) data.

Figure 4. Root Mean Squared Error (RMSE, top plots) and Brier score (bottom plots) of non-

transferred and transferred models of *C. penicillatus* (left plots) and *C. baileyi* (right plots) on

control (purple) and removal plots (yellow) at different forecast horizons.

 FIGURES

 Figure 1.

585 **Figure 2.**

588   **Figure 3.**

591   **Figure 4.**