Melanophilin mediates the association of myosin-5a with melanosome via three distinct interactions

Jiabin Pan[‡§], Rui Zhou[‡§], Lin-Lin Yao[‡], Jie Zhang[‡], Ning Zhang[‡], Qin-Juan Cao[‡], Shaopeng Sun[‡§], and
Xiang-dong Li[‡§1]

[‡]Group of Cell Motility and Muscle Contraction, State Key Laboratory of Integrated Management of Pest
Insects and Rodents, Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, China

[§]University of Chinese Academy of Sciences, Beijing 100049, China

[1]To whom correspondence should be addressed: Xiang-dong Li, Institute of Zoology, Chinese Academy
of Sciences, Beijing 100101, China; Tel: +86-10-6480-6015; Email: lixd@ioz.ac.cn

Running title: Myosin-5a and melanophilin interaction

Key words: actin, melanophilin, melanosome, myosin-5a, Rab27a

ABSTRACT

Transport and localization of melanosome at the periphery region of melanocyte are depended on
myosin-5a (Myo5a), which associates with melanosome by interacting with its adaptor protein melanophilin
(Mlph). Mlph contains four functional regions, including Rab27a-binding domain, Myo5a GTD-binding
motif (GTBM), Myo5a exon F-binding domain (EFBD), and actin-binding domain (ABD). The association
of Myo5a with Mlph is known to be mediated by two specific interactions: the interaction between the
exon-F-encoded region of Myo5a and Mlph-EFBD and that between Myo5a-GTD and Mlph-GTBM. Here,
we identify a third interaction between Myo5a and Mlph, i.e., the interaction between the exon-G-encoded
region of Myo5a and Mlph-ABD. The exon-G/ABD interaction is independent from the exon-F/EFBD
interaction and is required for the association of Myo5a with melanosome. Moreover, we demonstrate that
Mlph-ABD interacts with either the exon-G or actin filament, but cannot interact with both of them
simultaneously. Based on above findings, we propose a new model for the Mlph-mediated Myo5a
transportation of melanosomes.

Introduction

Melanosome is a specialized organelle within which melanin pigments are synthesized and stored. The
peripheral accumulation of melanosomes in melanocytes is essential for the intercellular transfer of the
organelles from the dendrite tips of melanocytes to the adjacent keratinocytes (A. N. Hume & Seabra, 2011).
Microtubule and the associated motor proteins are responsible for the long-range and bidirectional transport
of melanosomes between the nucleus to cell periphery, and actin filament and the associated motor protein
myosin-5a (Myo5a) are essential for the short-distance transport and the capture of melanosomes at cell
periphery (X. Wu, Bowers, Rao, Wei, & Hammer, 1998; X. Wu & Hammer, 2014). The overall effect of
the two transport systems is the peripheral accumulation of melanosomes in melanocytes. Lack of the latter
causes the perinuclear accumulation of melanosomes in melanocytes, a phenotype called "dilute" (Provance,
Wei, Ipe, & Mercer, 1996; Q. Wei, Wu, & Hammer, 1997).

Myo5a is a processive motor that has been implicated in the transportation and the tethering of
organelles along actin filaments (Lindsay & McCaffrey, 2014; Mehta et al., 1999; Rudolf, Bittins, & Gerdes,
2011; Yoshimura et al., 2006; N. Zhang, Yao, & Li, 2018). Myo5a has two heavy chains, each containing
an N-terminal motor domain, six IQ motifs that act as the binding sites for calmodulin (CaM) or CaM-like
light chains, and a tail (Ikebe, 2008; N. Zhang et al., 2018). The tail of Myo5a can be further divided into
three segments. The proximal tail is a long coiled-coil with length about 200 residues. The distal tail is the
C-terminal globular tail domain (GTD). Between the proximal tail and the GTD is the middle tail, which is
comprised of several short coiled-coils and the connecting loops. At least three essential functions are
served by Myo5a tail. First, the coiled-coil regions enable Myo5a to form a dimer. Second, the GTD
inhibits the motor activity of myosin head. Third, the tail is the binding site for the cargo proteins. The
motor activity of Myo5a is tightly regulated (Krementsov, Krementsova, & Trybus, 2004; Li, Jung,
Mabuchi, Craig, & Ikebe, 2006; Li et al., 2008; Li, Mabuchi, Ikebe, & Ikebe, 2004; Liu, Taylor,
Krementsova, Trybus, & Taylor, 2006; Thirumurugan, Sakamoto, Hammer, Sellers, & Knight, 2006; Wang
et al., 2004). In the inhibited state, Myo5a forms a folded conformation, in which the GTDs fold back to
interact and inhibit the motor function. Upon being activated by high Ca^{2+}, cargo or adaptor proteins,
Myo5a transforms from the folded conformation to the extended conformation, in which the GTD
dissociates from the motor domain and the inhibition is relieved.

Melanophilin (Mlph) is one of the best-studied cargo proteins of Myo5a. Together with Rab27a, Mlph
mediates the attachment of Myo5a to melanosomes. Rab27a is a small GTPase that, in the GTP-bound form,
is anchored into the melanosome membrane via its lipid moiety (Ishida, Arai, Ohbayashi, & Fukuda, 2014;
Wandinger-Ness & Zerial, 2014) and Mlph bridges the interaction between Rab27a and Myo5a (Fukuda,
Kuroda, & Mikoshiba, 2002; Nagashima et al., 2002; X. S. Wu et al., 2002). The tripartite complex of
Rab27a, Mlph, and Myo5a connects the melanosome to the actin filament. Lacking any one of these three
components in melanocytes causes dilute phenotype (perinuclear accumulation of melanosomes) (Fukuda et
al., 2002; A. N. Hume et al., 2001; Strom, Hume, Tarafder, Barkagianni, & Seabra, 2002; Q. Wei et al.,
1997; X. Wu et al., 2001; X. S. Wu et al., 2002).

Mlph contains four functional regions, including RBD (Rab27a-binding domain), GTBM (Myo5a
GTD-binding motif), EFBD (Myo5a exon F-binding domain), and ABD (actin-binding domain)(Fukuda et
al., 2002; Geething & Spudich, 2007; Li, Ikebe, & Ikebe, 2005; Yao et al., 2015). The association of Myo5a
with Mlph involves two molecular interactions. One is the interaction between the exon-F-encoded region
(exon-F for simplification) of Myo5a and the EFBD of Mlph (Fukuda & Itoh, 2004; X. Wu, Wang, Rao,
Sellers, & Hammer, 2002; X. S. Wu et al., 2002). Exon-F is an alternatively spliced exon presents in the
middle tail of the melanocyte-spliced isoform of Myo5a (Huang et al., 1998). The exon-F/EFBD interaction
is absolutely essential for the localization of Myo5a to the melanosome (X. Wu et al., 2002). The other
interaction is between the GTD of Myo5a and the GTBM of Mlph (Geething & Spudich, 2007; Pylypenko
et al., 2013; Z. Wei, Liu, Yu, & Zhang, 2013). We previously found that GTBM activates Myo5a motor
function by relieving the GTD inhibition on the motor domain and inducing a folded-to-extended
conformational transition of Myo5a (Li et al., 2005; Yao et al., 2015). Recently, we demonstrated that the

activation of Myo5a by Mlph-GTBM is regulated by another Myo5a-binding protein RILPL2
(Rab-interacting lysosomal protein-like 2) and the small GTPase Rab36, a binding partner of RILPL2 (Cao,
Zhang, Zhou, Yao, & Li, 2019).

Mlph-ABD is essential to proper melanosome transport (Fukuda & Kuroda, 2002; T. S. Kuroda, H.
Ariga, & M. Fukuda, 2003; Taruho S. Kuroda, Hiroyoshi Ariga, & Mitsunori Fukuda, 2003). Besides being
able to bind to actin filament, Mlph-ABD could interact with end-binding protein 1, which might facilitate
the transfer of melanosomes from microtubules to actin (X. S. Wu, Tsan, & Hammer, 2005). *In vitro*
motility assays at single-molecular level show that Mlph prolongs and slows the processive movements of
Myo5a, presumably via the interaction between ABD and actin filament (Sckolnick, Krementsova,
Warshaw, & Trybus, 2013). A cluster of conserved positively charged residues in ABD is essential for actin
binding and Melanophilin localization in melanocytes (T. S. Kuroda et al., 2003). An *in vitro* study suggests
that Mlph-ABD is a target of protein kinase A and that the phosphorylated Mlph preferred binding to actin
even in the presence of microtubules, whereas dephosphorylated Mlph bind to microtubules (Oberhofer et
al., 2017).

Biochemical analysis of Mlph show that the ABD is not essential for the interaction with Myo5a nor
the activation of Myo5a motor function, suggesting that a truncated Mlph protein containing the RBD and
the two Myo5a-binding domains (the GTBM and the EFBD) but lacking the ABD is able to form a tripartite
complex with Rab27a and Myo5a and thus sufficient for the recruitment of Myo5a to melanosome
membrane (T. S. Kuroda et al., 2003; X. Wu et al., 2002). However, using melan-ln cells (Mlph-null
melanocytes), Hume et al. have found that the ABD-deleted Mlph cannot to promote the recruitment of
Myo5a onto melanosomes, indicating that the ABD is necessary *in situ* for the association of Myo5a with
Mlph (A. N. Hume, A.K. Tarafder, J.S. Ramalho, E.V. Sviderskaya, and M.C. Seabra, 2006). This
discrepancy between *in vitro* and cell culture results promoted us to investigate the interaction between
Mlph-ABD and Myo5a.

In this study, we have identified a third interaction between Myo5a and Mlph, i.e., the interaction
between the exon-G-encoded region (exon-G for simplification) of Myo5a and Mlph-ABD. We find that the
exon-G/ABD interaction is independent from the exon-F/EFBD interaction, and that, similar to
exon-F/EFBD, exon-G/ABD is essential for the dilute-like phenotype induced by Myo5a-tail
overexpression. Moreover, we demonstrate that Mlph-ABD interacts with either the exon-G of Myo5a or
actin filament, but cannot interact with both of them simultaneously. Based on above findings, we propose a
new model for the Mlph-mediated Myo5a transportation of melanosomes.

108
109

                                         RESULTS

The actin-binding domain of melanophilin (Mlph-ABD) contains a novel Myo5a-binding site.

To detect interaction between the ABD of Mlph with Myo5a, we performed GST pulldown assay
using GST-Mlph-ABD (the GST-tagged ABD of Mlph, residues 400-590) and Flag-Myo5a-MTD (the
Flag-tagged middle tail domain of Myo5a, residues 1106-1469) (Figure 1). GST-Mlph-ABD was expressed
in E. coli and purified by GSH-Sepharose chromatography, and Flag-Myo5a-MTD was expressed in Sf9
cells and purified by anti-Flag affinity chromatography. GST pulldown assay shows that GST-Mlph-ABD
specifically interacted with Flag-Myo5a-MTD (Figure 1C, lane 2). Deleting the residues 400-437 of
Mlph-ABD had little effect on the interaction with Myo5a-MTD (Figure 1C, lane 3 and 4), indicating the
N-terminal 37 residues of Mlph-ABD are not essential for the interaction. Further deletion of 437-446
slightly decreased the interaction and deletion of 446-455 eliminated the interaction (Figure 1C, lane 5 and
6), indicating that residues 446-455 are essential for the interaction with Myo5a-MTD. Similarly, we
performed C-terminal truncation analysis of Mlph-ABD on the interaction with Myo5a-MTD and identified
residues 560-571 essential for the interaction (Figure 1D). Taking together, we identified the middle portion
of Mlph-ABD (residues 446-571) as the core of the novel binding site for Myo5a-MTD.

We then produced a number of truncated Myo5a-MTD constructs and performed GST pulldown
assays to narrow down the region in Myo5a-MTD binding to Mlph-ABD (Figure 2A). To simplify the
experiments, we first compared the bacterial-expressed Myo5a-MTD with the Sf9 cell-expressed
Myo5a-MTD in the interaction with Mlph-ABD. GST pulldown assay shows that both versions of
Myo5a-MTD specifically interacted with Mlph-ABD (Figure 2-figure supplement A). All the following
pulldown experiments were performed using the bacterial-expressed Myo5a-MTD, unless otherwise
indicated. C-terminal truncation of Flag-Myo5a-MTD shows that the C-terminal half of exon-G (residues
1453-1467) is essential for binding to Mlph-ABD (Figure 2-figure supplement B). Mlph-ABD interacted
strongly with Flag-Trx-Myo5a (1411-1467) (a short peptide containing exon-F and exon-G), but weakly
with Flag-Trx-Myo5a (1436-1467) (a short peptide containing exon-G) (Figure 2-figure supplement C, lane
3 and 4), suggesting that exon-F might be required for strong interaction of exon-G with Mlph-ABD.
However, deletion of exon-F from Myo5a-MTD did not affect the interaction with Mlph-ABD (Figure 2B,
lane 2), indicating that exon-F is not essential for the interaction. Deletion of the C-terminal portion
(residues 1454-1467) of exon-G abolished the binding of Myo5a-MTD with Mlph-ABD (figure 2B, lane 3),
but did not affect the interaction of Myo5a-MTD with Mlph-EFBD (Figure 2C). Figure 2A summaries the
truncation analysis of the interaction between Myo5a-MTD and Mlph-ABD. Based on above results, we
conclude that the exon-G of Myo5a binds to Mlph-ABD.

Ionic interactions are critical for the exon-G/ABD interaction.

We characterized the exon-G/ABD interaction by investigating the effect of ionic strength on their
interaction. GST pulldown assays of GST-Mlph-ABD and Flag-Myo5a-MTD were conducted in the
presence different concentrations of NaCl. Compared with 100 mM NaCl, 300 mM and 500 mM NaCl
greatly decreased the amount of Flag-Myo5a-MTD pulled down with GST-Mlph-ABD (Figure 3A),
suggesting that ionic interactions are critical for the exon-G/ABD interaction.

Sequence alignment among the Mlph of mammals and birds reveals several highly conserved charged
residues in the two regions of Mlph-ABD essential for the interaction with Myo5a-MTD, including four
acidic residues (D447, E449, E452, and E454) in the residues 446-455 and two basic residues (R562 and
R568) in the residues 560-571 (Figure 3B). To determine the roles of those conserved charged residues of
Mlph on the exon-G/ABD interaction, we mutated them individually into alanine. GST pulldown assays
show that both E449A and E452A mutations abolished the interaction with Myo5a-MTD, whereas other
mutations (D447A, E454A, R562A, R568A) had little effect on the interaction (Figure 3C), indicating that
both E449 and E452 of Mlph are essential for the exon-G/ABD interaction.

We expected the conserved acidic residues E449 and E452 of Mlph to interact with the conserved basic residues in the exon-G of Myo5a. Sequence alignment of the three Myo5 isoforms of mammals and birds reveals two highly conserved basic residues (K1456 and K1460) in the C-terminal portion of exon-G of Myo5a (Figure 3D). GST pulldown assay shows that both K1456A and K1460A mutations substantially decreased the amount of Flag-Myo5a-MTD pulled down with GST-Mlph-ABD (Figure 3E), suggesting that both K1456 and K1460 of Myo5a play a key role in the interaction with Mlph-ABD.

The exon-F/EFBD interaction and the exon-G/ABD interaction are independent from each other.

Because exon-F and exon-G are in immediate vicinity, it is possible that the exon-F/EFBD interaction and the exon-G/ABD interaction interfere with each other. To test this possibility, we performed GST pulldown assays of GST-Mlph-ABD and Flag-Mlph-EFBD in the presence or absence of Flag-Myo5a-MTD. If Myo5a-MTD is able to interact with both Mlph-ABD and Mlph-EFBD simultaneously, a tripartite complex will be formed by Myo5a-MTD, Mlph-ABD, and Mlph-EFBD (Figure 4A, upper panel), and Flag-Mlph-EFBD will be pulled down with GST-Mlph-ABD in the presence of Flag-Myo5a-MTD. As shown in Figure 4A, Flag-Mlph-EFBD was pulled down with GST-Mlph-ABD in the presence of Flag-Myo5a-MTD, but not in the absence of Flag-Myo5a-MTD. This result is consistent with a scenario that Myo5a-MTD binds to both Mlph-EFBD and Mlph-ABD simultaneously. We therefore conclude that the exon-F/EFBD interaction and the exon-G/ABD interaction do not interfere with each other.

As the exon-F/EFBD interaction and the exon-G/ABD interaction are independent to each other, we expected that those two interactions act synergically for the binding of Myo5a-MTD to Mlph. As expected, GST pulldown assays show that GST-Mlph-ΔRBD, which contains both EFBD and ABD, strongly bound to Flag-Myo5a-MTD, and deletion of either exon-F or exon-G from Myo5a-MTD substantially weakened the interaction with GST-Mlph-ΔRBD (Figure 4B). Conversely, Flag pulldown shows that substantial amount of GST-Mlph-ΔRBD could be pulled down with Flag-Myo5a-MTD, and deleting ABD from GST-Mlph-ΔRBD strongly decreased the amount of protein pulled down with Flag-Myo5a-MTD (Figure 4C). Above results indicate that the exon-F/EFBD interaction and the exon-G/ABD interaction act synergically, resulting in a strong interaction between Myo5a-MTD and Mlph.

Mlph-ABD cannot interact with F-actin and exon-G simultaneously.

In addition to interacting with the exon-G of Myo5a, Mlph-ABD is able to bind to actin. To investigate whether Mlph-ABD interacts with actin and exon-G simultaneously, we performed actin co-sedimentation of Mlph-ABD and Myo5a-MTD. As expected, substantial amount of Mlph-ABD, but essentially no Myo5a-MTD, was co-sedimented with F-actin (Figure 5A). Interestingly, Mlph-ABD did not increase the amount of Myo5a-MTD co-sedimented with F-actin (Figure 5A), suggesting that Mlph-ABD cannot interact with F-actin and exon-G simultaneously.

The inability of Mlph-ABD to interact with F-actin and exon-G simultaneously suggests that F-actin and exon-G compete in binding to Mlph-ABD. It is possible that Myo5a-MTD might antagonize the interaction between Mlph-ABD and F-actin. To test this possibility, we performed F-actin co-sedimentations of Mlph-ABD in the presence of different concentrations of Myo5a-MTD. As expected, Myo5a-MTD substantially decreased the amount of Mlph-ABD co-sedimented with F-actin (Figure 5B). We therefore concluded that Mlph-ABD cannot bind to F-actin and exon-G simultaneously and those two interactions are mutually exclusive.

The exon-G/ABD interaction is essential for Myo5a-tail to induce dilute-like phenotype in melanocytes.

Myo5a associates with melanosome via its interaction with Mlph, which attaches to melanosome via the interaction between its RBD and the melanosome-bound Rab27a. Overexpression of the GFP fusion protein of Mlph-ΔRBD in the melanocyte cell line, melan-a, caused dilute-like phenotype, i.e., the perinuclear distribution of melanosomes in melanocyte (Figure 6A). Generation of dilute-like phenotype by

Mlph-ΔRBD was attributed to the ability of Mlph-ΔRBD to compete with the melanosome-bound Mlph in
interacting with Myo5a molecules, thereby dissociating Myo5a from melanosome. To investigate whether
the exon-G/ABD interaction is essential for the dominant negative effect of Mlph-ΔRBD in melanocyte, we
introduced two single pointed mutations, E449A and E452A, in Mlph-ΔRBD, both of which are essential
for Mlph-ABD to interact with the exon-G (Figure 3C). We found that both two mutations substantially
decreased the number of the transfected cells with dilute-like phenotype. While dilute-like phenotype was
presented in 48.78% of cells expressing the wild-type Mlph-ΔRBD (n=194), only in 20.97% and 19.07% of
cells expressing E449A (n=208) and E542A (n=267), respectively (Figure 6).

To further determine the role of the exon-G/ABD interaction on the tethering of Myo5a with Mlph in
melanocyte, we expressed the GFP fusion protein of Myo5a-tail (residues 1106-1877), which contains all
three Mlph-binding regions, i.e., exon-F, exon-F, and GTD (Figure 7A). Overexpressing the tail region of
Myo5a in melanocyte was expected to compete with endogenous Myo5a in interacting with Mlph, thus
causing dilute-like phenotype (Wu et al 2002). Deletion analysis of the tail region of Myo5a have shown
that both exon-F and the GTD are required and that neither is sufficient for generating the dominant
negative phenotype in melanocytes (Wu et al 2002). Similar to the previous report, Myo5a-tail is well
co-localized with both melanosome and Mlph, and generated dilute-like phenotype in ~50% of transfected
melan-a melanocytes (n=233). In contrast, Myo5a-TailΔG generated dilute-like phenotype in only ~10% of
transfected cells (n=194) (Figure 7). Therefore, we concluded that, similar to exon-F, exon-G is also
required for Myo5a-Tail to disrupt endogenous Myo5a function in transportation of melanosomes.

Myo5a-MTD colocalized with Mlph but was unable to induce dilute-like phenotype.

Unlike Myo5a-Tail, overexpression of Myo5a-MTD, which contains both exon-F and exon-G but
lacks the GTD, did not generate dilute-like phenotype in melan-a cells. However, Myo5a-MTD is well
colocalized with endogenous Mlph and partially colocalized with melanosome (Figure 8A). We reasoned
that the presence of both exon-F and exon-G is insufficient for binding to the Mlph occupied by Myo5a, but
sufficient for binding to the unoccupied Mlph.

Deletion of exon-F almost eliminated the co-localization of Myo5a-MTD and Mlph (Figure 8B),
consistent with its essential role for the binding of Myo5a with Mlph. Deletion of exon-G substantially
decreased, but did not eliminate, the co-localization of Myo5a-MTD and Mlph (Figure 8C), indicating that
exon-G plays an auxiliary role for binding of Myo5a with Mlph.
232
233

DISCUSSION

It is well-established that Myo5a associates with melanosome via two independent interactions with
Mlph, i.e., the interaction between exon-F of Myo5a and Mlph-EFBD and the interaction between the GTD
and Mlph-GTBM. In the current study, we identified the third interaction between Myo5a and Mlph, i.e.,
the interaction between the exon-G of Myo5a and the Mlph-ABD. We demonstrated that, similar to the
exon-F/EFBD interaction, the exon-G/ABD interaction is also required for the strong binding of Myo5a and
Mlph.

Among the three interactions between Myo5a and Mlph, the GTD/GTBM interaction majorly regulates
the motor function of Myo5a. At relative high ionic strength but not at physiological ionic strength,
Mlph-GTBM binds to the GTD, inducing Myo5a to form the extended conformation and activating Myo5a
motor activity (Li et al., 2005; Yao et al., 2015). At physiological ionic strength, the GTBM-mediated
activation of Myo5a depends on the interaction between the GTD and RilpL2, which is regulated by Rab36
(Cao et al., 2019). Those findings are consistent with the three-dimensional structure of Myo5a-GTD and
the folded structure of full-length Myo5a (Niu et al., 2022; Pylypenko et al., 2013; Z. Wei et al., 2013), in
both of which the GTBM-binding sites are buried at the GTD-GTD interface. Upon binding to the RH1
domain of RilpL2, the GTD exposes the GTBM-binding site (Z. Wei et al., 2013). We therefore proposed
that the binding of Rab36/RilpL2 to the GTD exposes the GTBM-binding site, thus facilitating GTBM to
bind to the GTD and to activate the motor function of Myo5a (Cao et al., 2019).

The GTD/GTBM interaction is also required for the stable association between Myo5a and Mlph. The
Myo5a tail construct containing all three Mlph-binding sites (the exon-F, the exon-G, and the GTD)
produced dominant negative effect in melanocytes, and deleting the GTD abolished the dominant negative
effect (Figure 7 and 8, and reference (X. Wu et al., 2002)).

The exon-F/EFBD and the exon-G/ABD interactions play a major role in the binding of Myo5a with
Mlph. Although exon-F and exon-G are adjacent to each other, the exon-F/EFBD interaction and the
exon-G/ABD interaction do not interfere with each other. Rather, those two interactions act synergically.
We observed that Myo5a-MTD, containing both exon-F and exon-G but lacking the GTD, did not generate
a dilute-like phenotype in melanocytes, but was co-localized well with Mlph and partially with
melanosomes. Deletion of either exon-F or exon-G substantially weakened the interaction between
Myo5a-MTD and Mlph and decreased the colocalization of Myo5a-MTD with Mlph, indicating the both
exon-F and exon-G are required for the colocalization.

Our observation that Myo5a-MTD, containing both exon-F and exon-G, was co-localized partially
with melanosomes seems to contradict the early work from Hammer laboratory, which showed that MC
STK (a Myo5a fragment contains both exon-F and exon-G) did not exhibit any tendency to co-localize with
melanosomes (X. Wu et al., 2002). However, careful examining their image (Figure 5, in reference(X. Wu
et al., 2002)) reveals a partial colocalization of MC STK with melanosomes, particularly at the perinuclear
region. It is likely that some melanosome-bound Mlph molecules are unoccupied, some interact with
Myo5a via the exon-F/EFBD and the exon-G/ABD interactions but not the GTD/GTBM interaction, and
some interact with Myo5a via three interactions. In the former two cases, Myo5a-MTD would be able to
bind to the melanosome via Mlph. We therefore conclude that presence of both exon-F and exon-G is
sufficient for Myo5a to associate with Mlph, but insufficient for substituting the melanosome-bound Myo5a
molecules which bind to Mlph via three distinct interactions.

As discussed above, the GTBM-binding site is buried at the GTD-GTD interface in the folded state of
Myo5a, thus is inaccessible for Mlph, and the opening of GTBM-site depends on the interaction with
Rab36/RilpL2. On the other hand, the exon-F and the exon-G are likely exposed. We therefore propose that
the folded Myo5a first attaches to Mlph via the exon-F/EFBD and exon-G/ABD interactions, then interacts
with Rab36/RilpL2 to open the GTBM-site, and finally interacts with the GTBM, forming the extended
conformation and being activated.

One interesting finding of this study is that Mlph-ABD was able to separately interact with the exon-G
of Myo5a or actin filament, but unable to interact with both of them simultaneously. In other words,
Mlph-ABD binds to either exon-G or actin filament. This is consistent with the geometry of Myo5a, in
which exon-G is located far from the motor domain, which interacts with actin filament to produce motility.
Given the small size of Mlph-ABD, it is unlikely that Mlph-ABD is able to bridge the exon-G at one end of
Myo5a and actin filament which associates with the motor domain located at the other end of Myo5a. An
unsolved issue is how Mlph-ABD's interactions with exon-G and actin filament are regulated. *In vitro*
studies suggest that the interaction between Mlph-ABD and actin filament might be regulated by
phosphorylation (Oberhofer et al., 2017).

Based on our current finding and previous studies on the interaction between Myo5a and its associated
proteins, we propose following model for the Mlph-mediated Myo5a transportation of melanosome (Figure
9). At stage 1, Mlph associates with melanosome via its interaction with Rab27a, which directly binds to the
membrane of melanosome; the unattached Myo5a is in a folded conformation, in which the GTD binds to
and inhibits the motor domain. At stage 2, Mlph interacts with the folded Myo5a via the interactions of
EFBD/exon-F and ABD/exon-G; the attached Myo5a is still in folded conformation, because the
GTBM-binding surface in the GTD is buried at the GTD-GTD interface. At stage 3, the buried
GTBM-binding surface between the GTD-GTD interface is exposed and thus facilitate the binding of
GTBM, causing the dissociation of the GTD from the motor domain and inducing the extended
conformation of Myo5a (This step is probably regulated by the binding of Rab36/RilpL2 to the GTD). At
stage 4, Mlph-ABD dissociates from exon-G and then binds to actin filament, thus enhancing the processive
movement of Myo5a (This step might be regulated by the phosphorylation of Mlph-ABD). The interaction
between Mlph-ABD and actin filament is regulated by phosphorylation (Oberhofer et al., 2017).

Melanosomes in melanocytes are matured through a serial of well-defined morphological steps, from
melanin-lacking premelanosomes to blackened melanosomes with melanin fully loaded (Raposo & Marks,
2007). During this maturation process, melanosomes undergo microtubule and actin-based transport
towards the cell periphery, mediated by different Rab proteins and their effectors (Fukuda, 2021). For
example, Rab27a mainly associates with intermediate and mature melanosomes (Jordens et al., 2006). The
melanosome-bound Rab27a in GTP-bound state binds to Mlph, which in turn recruits Myo5a (X. S. Wu et
al., 2002). We propose that the recruitment of Myo5a by Mlph consists of multiple stages. We expect that
Myo5a-MTD, containing both exon-F and exon-G, is able to bind to Mlph at stage 1 and 2, but not at stage
3 and 4, and that Myo5a-Tail, containing all three Mlph-binding sites, is able to bind to Mlph at all four
stages.

In conclusion, we demonstrate a direct interaction between the exon-G of Myo5a and the ABD of
Mlph, which is essential for the tight binding of Myo5a to Mlph both *in vitro* and *in vivo*. We expect that
the melanosomal recruitment and activation of Myo5a are a highly coordinated process mediated by three
interactions between Myo5a and Mlph, i.e., the exon-F/EFBD interaction, the exon-G/ABD interaction, and
the GTD/GTBM interactions.
318

 EXPERIMENTAL PROCEDURES

Materials

Restriction enzymes and modifying enzymes were purchased from New England Biolabs (Beverly, MA), unless indicated otherwise. Actin was prepared from rabbit skeletal muscle acetone powder according to Spudich and Watt (Spudich & Watt, 1971). Ni-NTA agarose was purchased from Qiagen (Hilden, Germany). Anti-Flag M2 affinity gel and HPR conjugated anti-Flag M2 antibody were from Sigma Co. (St. Louis, MO). Glutathione(GSH)-Sepharose 4 Fast Flow was from GE Healthcare. FLAG peptide (DYKDDDDK) was synthesized by Augct Co. (Beijing, China). Oligonucleotides were synthesized by Invitrogen Co. (Beijing, China).

Proteins

All Myo5a constructs in this study were created from a melanocyte type Myo5a (Li et al., 2005). The recombinant proteins of truncated Myo5a containing an N-terminal His-tag and Flag-tag were expressed in BL21(DE3) E. coli. The cDNA of truncated Myo5a were amplified by using high-fidelity FastPfu DNA polymerase (TransGen) with Myo5a cDNA as template (Li et al., 2005), and subcloned into pET30HFa (a modified pET30a vector encoding His-tag and Flag-tag). To improve the expression of short peptides of truncated Myo5a, including Myo5a(1411-1467) and Myo5a(1436-1467), the cDNA of Trx-1 (Genbank ID: P0AA25) was inserted between the Flag-tag sequence and the cDNA of truncated Myo5a in pET30HFa. Myo5a-MTDΔF were created by using overlap extension PCR with Myo5a-MTD/pFast FTb as template and inserted into pET30HFa. The recombinant proteins were expressed in BL21(DE3) *E. coli* as His-Flag tagged proteins were purified by Ni-agarose affinity chromatography using standard procedures. We also produced an N-terminal His-tagged and Flag-tagged Myo5a-MTD using baculovirus/Sf9 system. The cDNA of Myo5a-MTD was subcloned into the baculovirus transfer vector pFastHFTb (a modified pFastHTb vector containing the His-tag and Flag-tag sequence). Recombinant baculovirus was generated using Bac-to-Bac system. The Myo5a-MTD expressed in Sf9 insect cells was purified by Anti-FLAG M2 affinity chromatography (Li et al., 2008).

Three truncated Mlph constructs containing an N-terminal GST-tag, including Mlph-ABD (residues 401-590), Mlph-ΔRBD (residues 148-590), Mlph-ΔRBDΔABD (residues 148-400), were created by subcloning the cDNAs of truncated Mlph into pGEX4T1 vector using BamHI and XhoI sites. Mlph-EFBD (residues 241-400) having an N-terminal His-tag and Flag-tag was created by the cDNA of Mlph-EFBD subcloning into pET30HFa vector using EcoRI and HindIII sites. Point mutations were created using Fast Mutagenesis System (TransGen) according to the manufacturer's instructions. The recombinant proteins were expressed in BL21(DE3) *E. coli* as His-Flag tagged proteins (in pET30HFa vector) or GST tagged proteins (in pGEX4T1 vector), and purified by Ni-agarose affinity chromatography or GSH-Sepharose affinity chromatography using standard procedures.

Protein pulldown assay

GST pulldown assays were performed as described previously (W. B. Zhang, Yao, & Li, 2016). For GST pulldown of Myo5a-MTD with GST-Mlph-ABD, GSH-Sepharose beads (10 µl) were mixed with 95 µl of 2 µM GST-Mlph-ABD, 4 µM Myo5a-MTD truncations in Pulldown Buffer-I (5 mM Tris-HCl (pH 7.5), 100 mM NaCl, 1 mM DTT, and 1 mM EGTA, 0.1% NP-40) with rotation at 4 °C for 2 h. The GSH-Sepharose beads were then washed three times with 200 µl of Wash Buffer-I (10 mM Tris-HCl (pH 7.5), 100 mM NaCl, 1 mM DTT, and 1 mM EGTA, 0.1% NP-40), before eluted by Elution Buffer (10 mM GSH, 50 mM Tris-HCl (pH 8.0), 1 mM DTT, and 200 mM NaCl). For GST pulldown of Myo5a-MTD with GST-Mlph-ΔRBD, GSH-Sepharose beads (10 µl) were mixed with 95 µl of 2 µM GST-Mlph-ΔRBD, 4 µM Myo5a-MTD in Pulldown Buffer-II (5 mM Tris-HCl (pH 7.5), 100 mM NaCl, 1 mM DTT, and 1 mM EGTA) with rotation at 4 °C for 2 h. The GSH-Sepharose beads were washed three times with 200 µl of Wash Buffer-II (10 mM Tris-HCl (pH 7.5), 100 mM NaCl, 1 mM DTT, and 1 mM EGTA) and then eluted by Elution Buffer.

For Flag pulldown assay, Flag-Myo5a-MTD (0.5 μM) was incubated with 1 μM GST-tagged Mlph
truncations in Pulldown Buffer I and rotated at 4 °C for 2 h, mixed with 10 μl of anti-FLAG M2 gel. The
anti-FLAG M2 gel were washed three times with 200 μl of Wash Buffer I. The bound proteins were eluted
twice with 20 μl of 0.1 mg/ml FLAG peptide in Wash Buffer I.

The inputs and the eluted proteins were analyzed by SDS-PAGE and visualized by Coomassie Brilliant
Blue (CBB) staining or Western blotting using the indicated antibody. The amounts of pulldowned proteins
were quantified using ImageJ (version 1.42Q), and their molar ratios were calculated on the basis of their
molecular masses.

Actin cosedimentation assay

Rabbit skeletal actin (2 μM) and Mlph-ABD (4 μM) was incubated in a 60 μL solution of 20 mM Tris-HCl
(pH 7.5), 100 mM NaCl, 1 mM EGTA, and 1 mM DTT at 4 °C for 20 min with or without Myo5a-MTD (2
μM). In addition, Myo5a-MTD (0-8 μM) was incubated with rabbit skeletal actin (2 μM) and Mlph-ABD (1
μM) in a 60 μL solution of 20 mM Tris-HCl (pH 7.5), 100 mM NaCl, 1 mM EGTA, and 1 mM DTT at 4 °C
for 20 min. The mixtures were centrifuged at 85,000 rpm (Beckman Optima MAX-XP, TLA-120.1 rotor)
for 15 min at 4 °C. Equal portions of the pellet and the supernatant were subjected to SDS−PAGE and
Coomassie brilliant blue staining.

Melanophilin antibody

To generate a polyclonal antibody against melanophilin, the sequence encoding residues 147-428 of
melanophilin (Mlph-147-428) was amplified by PCR using a full-length mouse melanophilin cDNA as
template. The PCR product was cloned into pET30a vector and pGEX4T2 vector and expressed in E. coli as
His-tagged protein (His-Mlph-147-428) and GST-tagged protein (GST-Mlph-147-428), respectively.
His-Mlph-147-428 was purified using Ni-NTA agrose chromatography and used for immunizing rabbit.
GST-Mlph-147-428 was purified using GSH-Sepharose chromatography. Mlph antibody was
affinity-purified from immune sera over a column of GST-Mlph-147-428 coupled to cyanogen
bromide-activated Sepharose 4B (GE Healthcare) by standard procedure.

Plasmid constructions for melanocyte transfection

392   Myo5a-Tail/pEGFP-C1 was produced as described previously(X. Wu et al., 2002).
393 Myo5a-TailΔF/pEGFP-C1 and Myo5a-TailΔG/pEGFP-C1 were created by overlapping PCR using
394 Myo5a-Tail/pEGFP-C1 as template. Myo5a-MTD/pEGFP-C1 and Myo5a-MTDΔG/pEGFP-C1 were
395 produced by PCR amplification using Myo5a-Tail/pEGFP-C1 as template. Myo5a-MTDΔF/pEGFP-C1 was
396 created by subcloning from   Myo5a-MTDΔF/pET30a into pEGFP-C1.

Melanocyte culture and transfection

The immortal mouse melanocyte cell line melan-a, kindly provided by Dr. Dorothy Bennett, was
cultured as described (Bennett, Cooper, & Hart, 1987). Briefly, melan-a cells were cultured in RPMI 1640
medium (HyClone) supplemented with 10% fetal bovine serum, 200 nM phorbol 12-myristate 13-acetate
and penicillin/streptomycin (1% v/v) in humidified chamber (37 °C, 5% CO2 incubator). Melan-a cells
were passaged at 3–4 day intervals. Hieff Trans® Universal Transfection Reagent (Yeasen) was used for
transfection of pEGFP-C1 plasmids into melan-a cells according to the manufacturer's instructions.

Immunocytochemistry
For immunofluorescence staining, melan-a cells were plated on the coverslips. After 24 h culturing, cells
were transfected of pEGFP-C1 plasmids. Two days after transfection, the cells were fixed using 4%
paraformaldehyde for 20 min, then permeabilized with 0.4% Triton-X in PBS for 15 min. Subsequently, the
coverslips were blocked with 1% BSA for 1 h. Melan-a cells were incubated with rabbit antibody against
Mlph overnight. Coverslips were washed with PBS for 3 times and then incubated for 1 h at room
temperature with secondary antibodies (anti-rabbit IgG coupled with DyLight 549, Jackson). Nuclei were

counter-stained with DAPI. coverslips were fixed on glass slides with Fluoromount-GTM (Yeasen). The
images were captured by Leica STELLARIS 5 fluorescence microscope.
413
414

FOOTNOTES

416 The abbreviations used are: ABD, actin-binding domain of melanophilin; CaM, calmodulin; EFBD,
417 exon-F-binding domain of melanophilin; GSH, glutathione; GST, glutathione S-transferase; GTBM,
418 GTD-binding motif of melanophilin; GTD, globular tail domain of myosin-5a; Mlph, melanophilin; MTD,
419 middle tail domain of myosin-5a; Myo5a, myosin-5a.

ACKNOWLEDGEMENTS

422 This work was supported by the National Natural Science Foundation of China (31970657).

CONFLICT OF INTEREST

425 The authors declare that they have no conflicts of interest with the contents of this article.
426

AUTHOR CONTRIBUTIONS

428 XdL conceived the study. JP performed most experiments with help from RZ. LLY, NZ and JZ performed
429 the initial experiments demonstrating the interaction between Mlph-ABD and Myo5a tail. QJC prepared
430 melanophilin antibody. SP participated in preparation of recombinant proteins and pulldown assays. JP and
431 XdL designed the study, analyzed the data and wrote the manuscript.
432
433

REFERENCES

Bennett, D. C., Cooper, P. J., & Hart, I. R. (1987). A line of non-tumorigenic mouse melanocytes, syngeneic with the B16 melanoma and requiring a tumour promoter for growth. *Int J Cancer, 39*(3), 414-418.

Cao, Q. J., Zhang, N., Zhou, R., Yao, L. L., & Li, X. D. (2019). The cargo adaptor proteins RILPL2 and melanophilin co-regulate myosin-5a motor activity. *J Biol Chem, 294*(29), 11333-11341. doi:10.1074/jbc.RA119.007384

Fukuda, M. (2021). Rab GTPases: Key players in melanosome biogenesis, transport, and transfer. *Pigment Cell Melanoma Res, 34*(2), 222-235. doi:10.1111/pcmr.12931

Fukuda, M., & Itoh, T. (2004). Slac2-a/melanophilin contains multiple PEST-like sequences that are highly sensitive to proteolysis. *J Biol Chem, 279*(21), 22314-22321. doi:10.1074/jbc.401791200

Fukuda, M., & Kuroda, T. S. (2002). Slac2-c (synaptotagmin-like protein homologue lacking C2 domains-c), a novel linker protein that interacts with Rab27, myosin Va/VIIa, and actin. *J Biol Chem, 277*(45), 43096-43103. doi:10.1074/jbc.M203862200

Fukuda, M., Kuroda, T. S., & Mikoshiba, K. (2002). Slac2-a/melanophilin, the missing link between Rab27 and myosin Va: implications of a tripartite protein complex for melanosome transport. *J Biol Chem, 277*(14), 12432-12436. doi:10.1074/jbc.C200005200

Geething, N. C., & Spudich, J. A. (2007). Identification of a minimal myosin Va binding site within an intrinsically unstructured domain of melanophilin. *J Biol Chem, 282*(29), 21518-21528. doi:10.1074/jbc.M701932200

Huang, J. D., Mermall, V., Strobel, M. C., Russell, L. B., Mooseker, M. S., Copeland, N. G., & Jenkins, N. A. (1998). Molecular genetic dissection of mouse unconventional myosin-VA: tail region mutations. *Genetics, 148*(4), 1963-1972. doi:10.1093/genetics/148.4.1963

Hume, A. N., A.K. Tarafder, J.S. Ramalho, E.V. Sviderskaya, and M.C. Seabra. (2006). A Coiled-Coil Domain of Melanophilin Is Essential forMyosin Va Recruitment and Melanosome Transport inMelanocytes. *Molecular Biology of the Cell, 17*, 4720-4735. doi:10.1091/mbc.E06-05-0457)

Hume, A. N., Collinson, L. M., Rapak, A., Gomes, A. Q., Hopkins, C. R., & Seabra, M. C. (2001). Rab27a regulates the peripheral distribution of melanosomes in melanocytes. *J Cell Biol, 152*(4), 795-808.

Hume, A. N., & Seabra, M. C. (2011). Melanosomes on the move: a model to understand organelle dynamics. *Biochem Soc Trans, 39*(5), 1191-1196. doi:10.1042/BST0391191

Ikebe, M. (2008). Regulation of the function of mammalian myosin and its conformational change. *Biochem Biophys Res Commun, 369*(1), 157-164. doi:10.1016/j.bbrc.2008.01.057

Ishida, M., Arai, S. P., Ohbayashi, N., & Fukuda, M. (2014). The GTPase-deficient Rab27A(Q78L) mutant inhibits melanosome transport in melanocytes through trapping of Rab27A effector protein Slac2-a/melanophilin in their cytosol: development of a novel melanosome-targetinG tag. *J Biol Chem, 289*(16), 11059-11067. doi:10.1074/jbc.M114.552281

Jordens, I., Westbroek, W., Marsman, M., Rocha, N., Mommaas, M., Huizing, M., . . . Neefjes, J. (2006). Rab7 and Rab27a control two motor protein activities involved in melanosomal transport. *Pigment Cell Res, 19*(5), 412-423. doi:10.1111/j.1600-0749.2006.00329.x

Krementsov, D. N., Krementsova, E. B., & Trybus, K. M. (2004). Myosin V: regulation by calcium, calmodulin, and the tail domain. *J Cell Biol, 164*(6), 877-886.

Kuroda, T. S., Ariga, H., & Fukuda, M. (2003). The actin-binding domain of Slac2-a/melanophilin is required for melanosome distribution in melanocytes. *Mol Cell Biol, 23*(15), 5245-5255.

Kuroda, T. S., Ariga, H., & Fukuda, M. (2003). The Actin-Binding Domain of Slac2-a/Melanophilin Is Required for Melanosome Distribution in Melanocytes. *Molecular and Cellular Biology, 23*(15), 5245-5255. doi:10.1128/mcb.23.15.5245-5255.2003

Li, X. D., Ikebe, R., & Ikebe, M. (2005). Activation of myosin Va function by melanophilin, a specific docking partner of myosin Va. *J Biol Chem, 280*(18), 17815-17822. doi:10.1074/jbc.M413295200

Li, X. D., Jung, H. S., Mabuchi, K., Craig, R., & Ikebe, M. (2006). The globular tail domain of myosin Va functions as an inhibitor of the myosin Va motor. *J Biol Chem, 281*(31), 21789-21798. doi:10.1074/jbc.M602957200

Li, X. D., Jung, H. S., Wang, Q., Ikebe, R., Craig, R., & Ikebe, M. (2008). The globular tail domain puts on the brake to stop the ATPase cycle of myosin Va. *Proc Natl Acad Sci U S A, 105*(4), 1140-1145. doi:10.1073/pnas.0709741105

Li, X. D., Mabuchi, K., Ikebe, R., & Ikebe, M. (2004). Ca2+-induced activation of ATPase activity of myosin Va is accompanied with a large conformational change. *Biochem Biophys Res Commun, 315*(3), 538-545. doi:10.1016/j.bbrc.2004.01.084

Lindsay, A. J., & McCaffrey, M. W. (2014). Myosin Va is required for the transport of fragile X mental retardation protein (FMRP) granules. *Biol Cell, 106*(2), 57-71. doi:10.1111/boc.201200076

Liu, J., Taylor, D. W., Krementsova, E. B., Trybus, K. M., & Taylor, K. A. (2006). Three-dimensional structure of the myosin V inhibited state by cryoelectron tomography. *Nature, 442*(7099), 208-211. doi:10.1038/nature04719

Mehta, A. D., Rock, R. S., Rief, M., Spudich, J. A., Mooseker, M. S., & Cheney, R. E. (1999). Myosin-V is a processive actin-based motor. *Nature, 400*(6744), 590-593. doi:10.1038/23072

Nagashima, K., Torii, S., Yi, Z., Igarashi, M., Okamoto, K., Takeuchi, T., & Izumi, T. (2002). Melanophilin directly links Rab27a and myosin Va through its distinct coiled-coil regions. *FEBS Lett, 517*(1-3), 233-238.

Niu, F., Liu, Y., Sun, K., Xu, S., Dong, J., Yu, C., . . . Wei, Z. (2022). Autoinhibition and activation mechanisms revealed by the triangular-shaped structure of myosin Va. *Sci Adv, 8*(49), eadd4187. doi:10.1126/sciadv.add4187

Oberhofer, A., Spieler, P., Rosenfeld, Y., Stepp, W. L., Cleetus, A., Hume, A. N., . . . Okten, Z. (2017). Myosin Va's adaptor protein melanophilin enforces track selection on the microtubule and actin networks in vitro. *Proc Natl Acad Sci U S A, 114*(24), E4714-E4723. doi:10.1073/pnas.1619473114

Provance, D. W., Jr., Wei, M., Ipe, V., & Mercer, J. A. (1996). Cultured melanocytes from dilute mutant mice exhibit dendritic morphology and altered melanosome distribution. *Proceedings of the National Academy of Sciences of the United States of America, 93*(25), 14554-14558.

Pylypenko, O., Attanda, W., Gauquelin, C., Lahmani, M., Coulibaly, D., Baron, B., . . . Houdusse, A. M. (2013). Structural basis of myosin V Rab GTPase-dependent cargo recognition. *Proc Natl Acad Sci U S A, 110*(51), 20443-20448. doi:10.1073/pnas.1314329110

Raposo, G., & Marks, M. S. (2007). Melanosomes--dark organelles enlighten endosomal membrane transport. *Nat Rev Mol Cell Biol, 8*(10), 786-797. doi:nrm2258 [pii]
10.1038/nrm2258

Rudolf, R., Bittins, C. M., & Gerdes, H. H. (2011). The role of myosin V in exocytosis and synaptic plasticity. *J Neurochem, 116*(2), 177-191. doi:10.1111/j.1471-4159.2010.07110.x

Sckolnick, M., Krementsova, E. B., Warshaw, D. M., & Trybus, K. M. (2013). More than just a cargo adapter, melanophilin prolongs and slows processive runs of myosin Va. *J Biol Chem, 288*(41), 29313-29322. doi:10.1074/jbc.M113.476929

Spudich, J. A., & Watt, S. (1971). The regulation of rabbit skeletal muscle contraction. I. Biochemical studies of the interaction of the tropomyosin-troponin complex with actin and the proteolytic fragments of myosin. *J Biol Chem, 246*(15), 4866-4871.

Strom, M., Hume, A. N., Tarafder, A. K., Barkagianni, E., & Seabra, M. C. (2002). A family of Rab27-binding proteins. Melanophilin links Rab27a and myosin Va function in melanosome transport. *J Biol Chem, 277*(28), 25423-25430. doi:10.1074/jbc.M202574200

Thirumurugan, K., Sakamoto, T., Hammer, J. A., Sellers, J. R., & Knight, P. J. (2006). The cargo-binding domain regulates structure and activity of myosin 5. *Nature, 442*(7099), 212-215. doi:10.1038/nature04865

Wandinger-Ness, A., & Zerial, M. (2014). Rab proteins and the compartmentalization of the endosomal system. *Cold Spring Harb Perspect Biol, 6*(11), a022616. doi:10.1101/cshperspect.a022616

Wang, F., Thirumurugan, K., Stafford, W. F., Hammer, J. A., 3rd, Knight, P. J., & Sellers, J. R. (2004). Regulated conformation of myosin V. *J Biol Chem, 279*(4), 2333-2336.

Wei, Q., Wu, X., & Hammer, J. A., 3rd. (1997). The predominant defect in dilute melanocytes is in melanosome distribution and not cell shape, supporting a role for myosin V in melanosome transport. *J Muscle Res Cell Motil, 18*(5), 517-527.

Wei, Z., Liu, X., Yu, C., & Zhang, M. (2013). Structural basis of cargo recognitions for class V myosins. *Proc Natl Acad Sci U S A, 110*(28), 11314-11319. doi:1306768110 [pii]
10.1073/pnas.1306768110

Wu, X., Bowers, B., Rao, K., Wei, Q., & Hammer, J. A. r. (1998). Visualization of melanosome dynamics within wild-type and dilute melanocytes suggests a paradigm for myosin V function In vivo. *J Cell Biol, 143*(7), 1899-1918.

Wu, X., & Hammer, J. A. (2014). Melanosome transfer: it is best to give and receive. *Curr Opin Cell Biol, 29*, 1-7. doi:10.1016/j.ceb.2014.02.003

Wu, X., Rao, K., Bowers, M. B., Copeland, N. G., Jenkins, N. A., & Hammer, J. A., 3rd. (2001). Rab27a enables myosin Va-dependent melanosome capture by recruiting the myosin to the organelle. *Journal of Cell Science, 114*(Pt 6), 1091-1100.

Wu, X., Wang, F., Rao, K., Sellers, J. R., & Hammer, J. A., 3rd. (2002). Rab27a is an essential component of melanosome receptor for myosin Va. *Mol Biol Cell, 13*(5), 1735-1749. doi:10.1091/mbc.01-12-0595

Wu, X. S., Rao, K., Zhang, H., Wang, F., Sellers, J. R., Matesic, L. E., . . . Hammer, J. A., 3rd. (2002). Identification of an organelle receptor for myosin-Va. *Nat Cell Biol, 4*(4), 271-278. doi:10.1038/ncb760

Wu, X. S., Tsan, G. L., & Hammer, J. A., 3rd. (2005). Melanophilin and myosin Va track the microtubule plus end on
EB1. *J Cell Biol, 171*(2), 201-207. doi:10.1083/jcb.200503028
Yao, L. L., Cao, Q. J., Zhang, H. M., Zhang, J., Cao, Y., & Li, X. D. (2015). Melanophilin Stimulates Myosin-5a
Motor Function by Allosterically Inhibiting the Interaction between the Head and Tail of Myosin-5a. *Sci Rep,*
5, 10874. doi:10.1038/srep10874
Yoshimura, A., Fujii, R., Watanabe, Y., Okabe, S., Fukui, K., & Takumi, T. (2006). Myosin-Va facilitates the
accumulation of mRNA/protein complex in dendritic spines. *Curr Biol, 16*(23), 2345-2351.
doi:10.1016/j.cub.2006.10.024
Zhang, N., Yao, L. L., & Li, X. D. (2018). Regulation of class V myosin. *Cell Mol Life Sci, 75*(2), 261-273.
doi:10.1007/s00018-017-2599-5
Zhang, W. B., Yao, L. L., & Li, X. D. (2016). The Globular Tail Domain of Myosin-5a Functions as a Dimer in
Regulating the Motor Activity. *J Biol Chem, 291*(26), 13571-13579. doi:10.1074/jbc.M116.724328

FIGURE LEGENDS

Figure 1.   The actin-binding domain of melanophilin (Mlph-ABD) interacts with the middle tail
domain (MTD) of Myo5a. (A) Diagram of the melanocyte-spliced isoform of Myo5a. Myo5a-MTD,
Myo5a coiled-coils (residues 1106-1469). IQ, the CaM binding site; GTD, the C-terminal globular tail
domain. (B) Diagram of melanophilin (Mlph). RBD, Rab27a-binding domain; GTBM, globular tail
domain-binding motif; EFBD, exon-F binding domain; ABD, actin-binding domain. (C, D) GST pulldown
of GST-Mlph-ABD with the N-terminus truncated (C) or the C-terminus truncated (D) with
Flag-Myo5a-MTD. GST-Mlph-ABD constructs were bound to GSH-Sepharose and then incubated with
Flag-Myo5a-MTD. The GSH-Sepharose-bound proteins were eluted by GSH and analyzed by Western blot
using anti-Flag antibody. The inputs were analyzed with SDS-PAGE and visualized by Coomassie brilliant
blue (CBB) staining. GST was used as negative control. Note: The Sf9 cell-expressed Myo5a-MTD was
used in the GST pulldown assays shown in this figure.
Figure 1-Source data 1 Original and uncropped gels and blots for Figure 1C.
Figure 1-Source data 2 Original and uncropped gels and blots for Figure 1D.

Figure 2.   The exon-G of Myo5a interacts with Mlph-ABD. (A) Summary of the interactions between
the truncated Myo5a constructs and Mlph-ABD based on the GST pulldown assay shown in Figure S1. +,
strong interaction; +/-, weak interaction; -, no interaction. (B) Deletion of the C-terminal portion of exon-G
abolishes the interaction between Myo5a-MTD and Mlph-ABD. (C) Deletion of the C-terminal portion of
exon-G does not affect the interaction between Myo5a-MTD and Mlph-EFBD. GST pulldown assays were
performed using GST-Mlph-ABD and Flag-Myo5a-MTD variants. The GSH-Sepharose-bound proteins
were eluted by GSH and analyzed by Western blot using anti-FLAG antibody; the inputs were analyzed
with SDS-PAGE and visualized by Coomassie brilliant blue (CBB) staining.
Figure 2-Source data 1 Original and uncropped gels and blots for Figure 2B.
Figure 2-Source data 2 Original and uncropped gels and blots for Figure 2C.

Figure 3.   Identification of the key residues for the exon-G/ABD interaction. GST pulldown was
performed using GST-Mlph-ABD variants and Flag-Myo5a-MTD variants. (A) Ionic strength dependence
of the interaction between Myo5a-MTD and Mlph-ABD. GST pulldown was performed using
GST-Mlph-ABD and Flag-Myo5a-MTD in the presence of different concentrations of NaCl. (B) Sequence
alignments of the regions in Mlph-ABD essential for binding to Myo5a-MTD. Conserved charged residues
are indicated. (C) Effects of alanine mutations of the conserved charged residues in Mlph-ABD on the
interaction with Myo5a-MTD. E449A and E452A mutations in Mlph abolished the interaction between
Mlph-ABD and Myo5a-MTD. (D) Sequence alignments of the C-terminal portion of exon-G, an essential
region for binding to Mlph-ABD. Two conserved basic residues (K1456 and K1460) are indicated. (E)
Effects of K1456A and K1460A mutations of Myo5a-MTD on the interaction with Mlph-ABD.

Figure 3-Source data 1 Original and uncropped gels and blots for Figure 3A.
Figure 3-Source data 2 Original and uncropped gels and blots for Figure 3C.
Figure 3-Source data 3 Original and uncropped gels and blots for Figure 3E.

Figure 4.   The exon-F/EFBD and the exon-G/ABD interactions act synergistically. (A) Myo5a-MTD bridges between Mlph-ABD and Mlph-EFBD. Upper, diagram shows Myo5a-MTD binds to both the EFBD and the ABD of Mlph. Lower, GST pulldown assays of GST-Mlph-ABD and Flag-Mlph-EFBD with or without Flag-Myo5a-MTD. (B) Both exon-F and exon-G of Myo5a-MTD are required for the strong interaction with Mlph-ΔRBD. GST pulldown of GST-Mlph-ΔRBD with Flag-Myo5a-MTD variants. (C) ABD is essential for the strong interaction between Mlph-ΔRBD and Myo5a-MTD. The input samples were analyzed by SDS-PAGE and visualized by CBB staining. The pulled down samples were analyzed by Western blot using anti-Flag antibody (A and B) or by SDS-PAGE with CBB staining (C).
Figure 4-Source data 1 Original and uncropped gels and blots for Figure 4A.
Figure 4-Source data 2 Original and uncropped gels and blots for Figure 4B.
Figure 4-Source data 3 Original and uncropped gels for Figure 4C.

Figure 5.   Myo5a-MTD antagonizes the interaction between Mlph-ABD and actin. (A) GST-Mlph-ABD and/or Flag-Myo5a-MTD were incubated with actin and then subjected to ultracentrifugation. The supernatants (S) and the pellets (P) were analyzed by SDS-PAGE (10%) with CBB staining. (B) GST-Mlph-ABD was incubated with actin in the presence of different concentrations of Flag-Myo5a-MTD and then subjected to to ultracentrifugation. Upper panel, the supernatants (S) and the pellets (P) were analyzed by SDS-PAGE (10%) with CBB staining. Lower panel, the amounts of GST-Mlph-ABD co-sedimentated with actin in the presence of different concentration of Flag-Myo5a-MTD were quantified based on the density in the SDS-PAGE. Data are the mean ± SD of three independent experiments with one-way ANOVA with post-hoc Bonferroni test. **$p<0.01$, ****$p<0.0001$. Note: a Sf9 cell-expressed Myo5a-MTD was used in the actin cosedimentation assays shown in this figure.
Figure 5-Source data 1 Original and uncropped gels for Figure 5A.
Figure 5-Source data 2 Original and uncropped gels and statistical data for Figure 5 B.

Figure 6.   Both E449 and E452 of Mlph are essential for the perinuclear distribution of melanosomes induced by Mlph-ΔRBD overexpression. Melan-a melanocytes were transfected to express EGFP-Mlph-ΔRBD or EGFP. The distribution of melanosomes in the transfected melanocytes was imaged and the number of melanocytes with perinuclear distribution of melanosomes was counted. (A) Typical images of melanocytes expressing EGFP-Mlph-ΔRBD, its mutants, or EGFP. Cells are outlined with a dashed line. Scale bars = 10 μm. (B) The percentage of melanocytes exhibiting perinuclear melanosome aggregation among the transfected melanocytes. Results are presented as the mean ± SD of three independent experiments with one-way ANOVA with post-hoc Bonferroni test. ****$p<0.0001$.
Figure 6-Source data 1 Original and statistical data for Figure 6 B.

Figure 7.   Exon-G region of Myo5a is essential for the perinuclear distribution of melanosomes induced by Myo5a-Tail overexpression. (A) Diagram of Myo5a-Tail. (B) Melan-a melanocytes were transfected to express EGFP-Myo5a-Tail and the mutant. The distribution of melanosomes in the transfected melanocytes was imaged and the number of melanocytes with perinuclear distribution of melanosomes was counted. (B) Typical images of melanocytes expressing EGFP-Myo5a-Tail, its mutants, or EGFP. Cells are outlined with a dashed line. Scale bars = 10 μm. (C) The percentages of melanocytes exhibiting perinuclear melanosome aggregation. Data are the mean ± SD of three independent experiments with one-way ANOVA with post-hoc Bonferroni test. ****$p<0.0001$.
Figure 7-Source data 1 Original and statistical data for Figure 7 C.

Figure 8.   Effects of deletion of exon-F or exon-G on the localization of Myo5a-MTD in melan-a cells.
Melan-a melanocytes were transfected to express EGFP-Myo5a-MTD (A), EGFP-Myo5a-MTDΔF (B), or
EGFP-Myo5a-MTDΔG (C) and stained for endogenous Mlph. Scale bars = 10 μm.

Figure 9.   A model for the Mlph-mediated Myo5a transportation of melanosome. The Mlph-mediated
Myo5a transportation of melanosome is comprised of four stages. At stage 1, Mlph associates with
melanosome via its interaction with Rab27a, which directly binds to the membrane of melanosome; the
unattached Myo5a is in a folded conformation, in which the GTD binds to and inhibits the motor domain.
At stage 2, Mlph interacts with the folded Myo5a via the interactions of EFBD/exon-F and ABD/exon-G;
the attached Myo5a is still in folded conformation, because the GTBM-binding surface in the GTD is buried
at the GTD-GTD interface. At stage 3, the buried GTBM-binding surface between the GTD-GTD interface
is exposed and thus facilitate the binding of GTBM, causing the dissociation of the GTD from the motor
domain and inducing the extended conformation of Myo5a (This step is probably regulated by the binding
of Rab36/RilpL2 to the GTD). At stage 4, Mlph-ABD dissociates from exon-G and then binds to actin
filament, thus enhancing the processive movement of Myo5a (This step might be regulated by the
phosphorylation of Mlph-ABD).

Figure 2-figure supplement.   Identification of the Mlph-ABD-binding site in the middle tail of
Myo5a. GST pulldown assays were performed using GST-Mlph-ABD and FLAG-Myo5a-MTD variants.
The GSH-Sepharose-bound proteins were eluted by GSH and analyzed by Western blot using anti-FLAG
antibody; the inputs were analyzed with SDS-PAGE and visualized by Coomassie brilliant blue (CBB)
staining. (A) GST pulldown assays of GST-Mlph-ABD with Flag-Myo5a-MTD expressed in E. coli or Sf9
cells. (B and C) GST pulldown assay of GST-Mlph-ABD with Flag-tagged, truncated Myo5a tail.
Figure 2-figure supplement Source data 1 Original and uncropped gels and blots for Figure 2-figure
supplement A.
Figure 2-figure supplement Source data 2 Original and uncropped gels and blots for Figure 2-figure
supplement B.
Figure 2-figure supplement Source data 3 Original and uncropped gels and blots for Figure 2-figure
supplement C.

Figure 1-Source data 1 Original and uncropped gels and blots.

Figure 1-Source data 2 Original and uncropped gels and blots.

Figure 2-Source data 1 Original and uncropped gels and blots.

Figure 2-Source data 2 Original and uncropped gels and blots.

Figure 3-Source data 1 Original and uncropped gels and blots.

Figure 3-Source data 2 Original and uncropped gels and blots.

Figure 3-Source data 3 Original and uncropped gels and blots.

Figure 4-Source data 1 Original and uncropped gels and blots.

Figure 4-Source data 2 Original and uncropped gels and blots.

Figure 4-Source data 3 Original and uncropped gels.

Figure 5-Source data 1 Original and uncropped gels.

Figure 5-Source data 2 Original and uncropped gels and data.

Figure 6-Source data 1 Original and statistical data.

Figure 7-Source data 1 Original and statistical data.

Figure 2-figure supplement Source data 1 Original and uncropped gels and blots.

Figure 2-figure supplement Source data 2 Original and uncropped gels and blots.

Figure 2-figure supplement Source data 3 Original and uncropped gels and blots.

Figure 1

Figure 2

Figure 3

A

B

D

C

E

Figure 4

Figure 5

A

Flag-Myo5a-MTD +   –   +
GST-Mlph-ABD –   +   +

CBB staining

B

CBB staining

Figure 6

Figure 7

A

Myo5a-Tail   1106- [A | C | D | E | F | G | GTD] -1877

Myo5a-Tail-ΔG   1106- [A | C | D | E | F |⋯| GTD] -1877
                                      1453 1467

B

| | DAPI | EGFP | Bright Field |

EGFP

EGFP-Myo5a-Tail

EGFP-Myo5a-Tail-ΔG

C

Figure 8

Figure 9

Stage 1: Mlph associates with melanosome via Rab27, and the unattached Myo5a is in the folded conformation.

Stage 2: Myo5a in folded conformation binds to Mlph via exon-F/EFBD and exon-G/ABD.

Stage 3: Mlph-GTBM binds to the GTD and induces Myo5a to form the extended conformation.

Stage 4: Mlph-ABD dissociates from exon-G, and then binds to actin filament, enhancing the processibility of Myo5a.

Figure 2-figure supplement