Loss of CTRP10 results in female obesity with preserved metabolic health

Fangluo Chen[1,2], Dylan C. Sarver[1,2], Muzna Saqib[1,2], Leandro M Velez[3,4], Susan Aja[2,5], Marcus M. Seldin[3,4],

G. William Wong[1,2]

Affiliations:

[1]Department of Physiology, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA

[2]Center for Metabolism and Obesity Research, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA

[3]Department of Biological Chemistry, University of California, Irvine, Irvine, USA

[4]Center for Epigenetics and Metabolism, University of California Irvine, Irvine, USA

[5]Department of Neuroscience, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA

Running head: CTRP10 deficiency uncouples obesity from metabolic dysfunction

Corresponding author: G. William Wong, E-mail: gwwong@jhmi.edu, Department of Physiology, Johns Hopkins

University School of Medicine, Baltimore, MD 21205

Tel: 410-502-4862

Fax: 410-614-8033

ABBREVIATIONS

31	CTRP	C1q/TNF-related protein
32	DEG	Differentially expressed gene
33	EE	energy expenditure
34	gWAT	gonadal white adipose tissue
35	iWAT	inguinal white adipose tissue
36	HDL	High density lipoprotein
37	HFD	high-fat diet
38	i.p.	intraperitoneally
39	KO	knockout
40	LFD	low-fat diet
41	MHO	Metabolically healthy obese
42	NEFA	non-esterified free fatty acids
43	RER	respiratory exchange ratio
44	TG	triglyceride
45	VLDL	Very low density lipoprotein
46	WT	wildtype

ABSTRACT

Obesity is a major risk factor for type 2 diabetes, dyslipidemia, cardiovascular disease, and hypertension. Intriguingly, there is a subset of metabolically healthy obese (MHO) individuals who are seemingly able to maintain a healthy metabolic profile free of metabolic syndrome. The molecular underpinnings of MHO, however, are not well understood. Here, we report that CTRP10/C1QL2-deficient mice represent a unique female model of MHO. CTRP10 modulates weight gain in a striking and sexually dimorphic manner. Female, but not male, mice lacking CTRP10 develop obesity with age on a low-fat diet while maintaining an otherwise healthy metabolic profile. When fed an obesogenic diet, female *Ctrp10* knockout (KO) mice show rapid weight gain. Despite pronounced obesity, *Ctrp10* KO female mice do not develop steatosis, dyslipidemia, glucose intolerance, insulin resistance, oxidative stress, or low-grade inflammation. Obesity is largely uncoupled from metabolic dysregulation in female KO mice. Multi-tissue transcriptomic analyses highlighted gene expression changes and pathways associated with insulin-sensitive obesity. Transcriptional correlation of the differentially expressed gene (DEG) orthologous in humans also show sex differences in gene connectivity within and across metabolic tissues, underscoring the conserved sex-dependent function of CTRP10. Collectively, our findings suggest that CTRP10 negatively regulates body weight in females, and that loss of CTRP10 results in benign obesity with largely preserved insulin sensitivity and metabolic health. This female MHO mouse model is valuable for understanding sex-biased mechanisms that uncouple obesity from metabolic dysfunction.

Key words: Metabolism, Obesity, Diabetes, Metabolically healthy obese (MHO)

INTRODUCTION

The prevalence of obesity has nearly tripled in the past four decades and the underlying cause is complex and multifactorial (1, 2). Genetics, environmental and social factors, and demographics all play a contributing role in contributing to excessive weight gain in the setting of overnutrition (3, 4). Although obesity is a major risk factor for type 2 diabetes, dyslipidemia, cardiovascular disease, and hypertension, not all obese individuals develop the metabolic syndrome (5). There is a subset of metabolically healthy obese (MHO) individuals with an apparently healthy metabolic profile free of some or most components of the metabolic syndrome (6, 7). The molecular and physiological underpinnings of MHO are, however, not well understood. Novel preclinical animal models that can recapitulate features of MHO will be valuable in illuminating pathways that resist the deleterious effects of obesity and provide new therapeutic avenues to mitigate obesity-linked comorbidities.

The mechanisms that normally maintain body weight and metabolic homeostasis are complex and involve both cell autonomous and non-cell autonomous mechanisms. Tissue crosstalk mediated by paracrine and endocrine factors plays an especially important role in coordinating metabolic processes across organ systems to maintain energy balance (8). Of the secretory proteins that circulate in plasma, C1q/TNF-related proteins (CTRP1-15) have emerged as important regulators of insulin sensitivity, and glucose and lipid metabolism (9). We originally identified the first seven members of the CTRP family based on shared sequence homology to the insulin-sensitizing adipokine, adiponectin (10), and subsequently characterized additional members (11-16). All fifteen CTRPs share a common C-terminal globular C1q domain and are part of the much larger C1q family (17, 18). The use of gain- and loss-of-function mouse models has helped establish CTRP's role in controlling various aspects of sugar and fat metabolism (12, 19-34). Additional diverse functions of CTRPs have also been demonstrated in the cardiovascular (35-47), renal (48, 49), immune (20, 50, 51), sensory (52, 53), gastrointestinal (54), musculoskeletal (55-57), and the nervous system (58-61).

Of the family members, CTRP10 (also known as C1QL2) is understudied and consequently only limited information is available concerning its function. The best characterized role of CTRP10 is in the central nervous system (CNS). It has been shown that CTRP10 secreted from mossy fibers is required for the proper clustering of

kainite-type glutamate receptors on postsynaptic CA3 pyramidal neurons in the hippocampus (62). It serves as a

transsynaptic organizer by directly binding to neurexin3 (Nrx3) on the presynaptic terminals, and to GluK2 and

GluK4 on the postsynaptic terminals (62). Additional putative roles of CTRP10 in the CNS have also been

suggested. Genome-wide association studies (GWAS) have implicated CTRP10/C1QL2 in cocaine use disorder

(63). In rat models of depression, *Ctrp10* expression is increased in the dentate gyrus and reduced in the nucleus

accumbens (64). In humans with a history of psychiatric disorders (e.g., schizophrenia), the expression of *CTRP10*

is elevated in the dorsolateral prefrontal cortex of both males and females (64). Whether and how CTRP10

contributes to addictive behavior and psychiatric disorders is unknown.

The potential function of CTRP10 in peripheral tissues, however, is essentially unknown and unexplored. The

present study was motivated by the well documented metabolic functions of many CTRP family members we have

characterized to date using genetic loss-of-function mouse models (19-28, 30, 31). We determined that the

expression of *Ctrp10* in peripheral tissues is modulated by diet and nutritional states, and thus may have a

metabolic role. We therefore used a genetic loss-of-function mouse model to determine if CTRP10 is required for

regulating systemic metabolism. We unexpectedly discovered a female-specific requirement of CTRP10 for body

weight control. We showed that the *Ctrp10* KO mice represent a unique female model of MHO with largely

preserved insulin sensitivity and metabolic health. This valuable mouse model can be used to inform sex-dependent

mechanisms that uncouple obesity from insulin resistance, dyslipidemia, and metabolic dysfunction.

RESULTS

Nutritional regulation of *Ctrp10* expression in the brain and peripheral tissues

CTRP10 protein is highly conserved from zebrafish to human (Fig. 1A), with amino acid identity of 67%, 71%,

77%, and 94% between the full-length human protein and the fish, frog, chicken, and mouse orthologs,

respectively. The conservation is much higher at the C-terminal globular C1q domain (93-100% identity) between

the orthologs. Among the 12 different mouse tissue examined, brain had the highest expression of *Ctrp10* (Fig. 1B),

consistent with previous findings (65). Expression of *Ctrp10* in peripheral tissues was variable and generally much

lower than in the brain (Fig. 1B). We first determined whether *Ctrp10* expression is modulated by nutrition and

metabolic state. Male mice were subjected to fasting and refeeding. In the refed period after an overnight fast, we

observed a significant downregulation of *Ctrp10* in the visceral (gonadal) white adipose tissue (gWAT), liver,

skeletal muscle, kidney, cerebellum, cortex, and hypothalamus relative to the fasted state (Fig. 1C). Next, we

examined whether an obesogenic diet alters the expression of *Ctrp10*. Male mice fed a high-fat diet for 12 weeks

had a modest increase in *Ctrp10* expression in brown adipose tissue (BAT) and heart and decreased expression in

skeletal muscle relative to mice fed a control low-fat diet (LFD) (Fig. 1D). These data indicate that *Ctrp10*

expression is dynamically regulated by acute alterations in energy balance, and perhaps to a lesser extent in

alterations to chronic nutritional state on an obesogenic diet.

Generation of *Ctrp10* knockout (KO) mice

We used mice lacking CTRP10 to address whether this secreted protein has a metabolic role in vivo. The mouse

Ctrp10 gene consists of two exons (Fig. 1E). The CRISPR-Cas9 method was used to remove the entire protein

coding region spanning exon 1 and 2, thus ensuring a complete null allele (Fig. 1E-F). The targeted allele was

confirmed by sequencing. As expected, based on the gene deletion strategy, the *Ctrp10* transcript was absent from

KO mice (Fig. 1G).

CTRP10 is largely dispensable for metabolic homeostasis in young mice fed a control low-fat diet

The body weight and body composition of male mice fed a LFD were not different between genotypes (Fig. 2A-B).

By 20 weeks of age, female KO mice fed LFD had a modestly higher body weight relative to WT controls (Fig. 2

C), though the body composition was not different between genotypes (Fig. 2D). Food intake, physical activity, and

energy expenditure as measured by indirect calorimetry were also not different between genotypes of either sex

across the circadian cycle (light and dark) and metabolic states (*ad libitum* fed, fasted, refed) (Fig. 2E-J). Because

Ctrp10 expression is regulated by nutritional states (Fig. 1C), we assessed serum metabolite levels in WT and KO

mice in response to fasting and refeeding. No significant differences in fasting and refeeding blood glucose, serum

insulin, triglyceride, cholesterol, non-esterified free fatty acids (NEFA), and β-hydroxybutyrate levels were

observed between genotypes of either sex, except the female KO mice had slightly lower fasting β-hydroxybutyrate

levels (Fig. 3A-B). We performed glucose and insulin tolerance tests to determine any potential differences in

glucose handling capacity and insulin sensitivity. No significant differences in glucose and insulin tolerance were

noted between genotypes of either sex (Fig. 3C-F). Together, these data indicate that CTRP10 is dispensable for

metabolic homeostasis when mice are young (< 20 weeks old) and fed a LFD.

CTRP10-deficient female mice on a low-fat diet develop obesity with age

Because female KO mice were slightly heavier at 20 weeks of age (Fig. 2C), we suspected the weight may diverge

further with age. Consequently, we monitored the body weight of female mice fed LFD over an extended period.

Indeed, the female KO mice gained significantly more weight and adiposity with age (Fig. 4A-C). Consistent with

greater adiposity, the adipocyte cell size (cross-sectional area) was also significantly larger in both gonadal

(visceral) white adipose tissue (gWAT) and inguinal (subcutaneous) white adipose tissue (iWAT) (Fig. 4D-E). By

the time the mice reached 40 weeks of age, female KO mice weighed ~ 6 g (20 %) heavier than the WT controls.

Increased weight gain over time was not attributed to differences in food intake, as measured manually over a 24 h

period (Fig. 4F). Fecal output, frequency, and energy content were also not different between genotypes (Fig. 4G),

suggesting that weight gain was not due to greater nutrient absorption. Deep colon temperatures in both light and

dark cycle were also not different between genotypes (Fig. 4H). Indirectly calorimetry analyses also revealed no

significant differences between genotypes in food intake, physical activity, and energy expenditure across the

circadian cycle and metabolic states (*ad libitum* fed, fasted, refed) (Fig. 4I-K). Despite significantly greater body

weight and adiposity, female KO mice had the same metabolic profile as the lean WT controls. There were no

differences in fasting blood glucose, serum insulin, triglyceride, cholesterol, NEFA, and β-hydroxybutyrate levels

between genotypes (Fig. 4L). Interestingly, VLDL-TG levels were lower in female KO mice whereas HDL-

cholesterol level was not different between genotypes (Fig. 4M). Direct assessments of glucose handling capacity

and insulin sensitivity by glucose and insulin tolerance tests, respectively, also revealed no differences between

genotypes (Fig. 4N-O). Together, these data indicate that *Ctrp10*-KO female mice fed LFD develop obesity, but

preserve a largely healthy metabolic profile similar to the much leaner WT female mice.

Rapid weight gain in CTRP10-deficient female mice fed a high-fat diet

Next, we challenged the mice with a HFD to determine if the sex-dependent effects on body weight become more

pronounced. When fed a HFD, body weight gain and body composition were not different between genotypes in

male mice (Fig. 5A-B). Food intake, physical activity, and energy expenditure were also not different between genotypes in male mice across the circadian cycle (light and dark) and metabolic states (*ad libitum* fed, fasted, refed) (Fig. 5C-E). In striking contrast, female KO mice gained weight rapidly on HFD (~9 g or 28% heavier) and had greater adiposity than the WT controls (Fig. 5F-H). Surprisingly, food intake, physical activity, and energy expenditure were not significantly different between genotypes in female mice (Fig. 5I-K). The ANCOVA analysis of energy expenditure using body weight as a covariate also did not reveal any differences between genotypes in female mice (Fig. 5 L). Interestingly, the respiratory quotient (RER) was significantly lower in female KO mice relative to WT controls, especially during fasting and refeeding (Fig. 5M), suggesting a greater reliance on lipid substrates for energy metabolism during those periods. Together, these data indicate that CTRP10 is required for female-specific body weight control in response to caloric surplus, but neither food intake, physical activity level, nor energy expenditure could account for the marked increase in body weight and adiposity.

Obesity is uncoupled from insulin resistance and dyslipidemia in CTRP10-deficient female mice fed a HFD

We again measured fasting and refeeding responses in WT and *Ctrp10*-KO mice fed a HFD. No differences in fasting and refeeding blood glucose, serum insulin, triglyceride, cholesterol, NEFA, and β-hydroxybutyrate levels were noted between genotypes in male mice (Fig. 6A). Female KO mice, however, had higher fasting blood glucose and serum insulin levels, and lower β-hydroxybutyrate levels compared to the WT controls (Fig. 6B). In the refed state, serum insulin levels continued to be significantly higher in female KO mice. Unlike the WT female mice where refeeding markedly lowered serum β-hydroxybutyrate (ketone) levels as expected, KO female mice appeared unable to suppress serum β-hydroxybutyrate levels in response to refeeding (Fig. 6B).

Consistent with the fasting blood glucose and insulin data, glucose handling capacity and insulin sensitivity assessments by glucose and insulin tolerance in HFD-fed male mice also revealed no differences between genotypes (Fig. 6C-D). Female KO mice, however, had higher fasting blood glucose and serum insulin levels suggesting the presence of mild insulin resistance (Fig. 6B). We therefore expected to see differences in either glucose and/or insulin tolerance tests. To our surprise, the rate of glucose clearance in response to glucose or insulin injection was virtually identical between WT and KO female mice (Fig. 6E-F), suggesting no difference in insulin sensitivity between genotypes. VLDL-TG and HDL-cholesterol profiles were also indistinguishable between WT and KO female mice (Fig. 6G). Altogether, these data indicate that CTRP10 is not required for metabolic

homeostasis in male mice challenged with a HFD. In female mice, however, loss of CTRP10 markedly promotes weight gain in the face of caloric surplus, but, paradoxically, the excess adiposity is largely uncoupled from obesity-linked insulin resistance and dysregulated glucose and lipid metabolism.

Obesity is uncoupled from adipose dysfunction and hepatic steatosis in *Ctrp10*-KO female mice fed a HFD

Consistent with greater fat mass in visceral (gonadal) fat depot of *Ctrp10*-KO female mice (Fig. 7A), histological analysis and quantification also indicated significantly larger adipocyte cell size (Fig. 7B). Although the subcutaneous (inguinal) fat pad weight was also significantly heavier in female KO mice (Fig. 7C), the adipocyte cell size was marginally bigger but not significant (Fig. 7D). A bigger fat pad with only marginally larger cell size suggests greater adipocyte hyperplasia in the subcutaneous depot. Increased adipogenesis in response to caloric surfeit is known to be associated with improved systemic metabolic profile (66).

Obesity is known to be associated with low-grade inflammation (67), fibrosis (68), and ER and oxidative stress (69, 70). Despite marked differences in body weight and adiposity, the expression of genes associated with inflammation (except for *Ccr2*), fibrosis, oxidative stress in gWAT and iWAT were not significantly different between female KO mice and WT controls (Fig. 7E). The expression of some genes associated with ER stress (e.g., *Ddit3*/*CHOP*, *Atf4*, *Xbp1*) in iWAT were actually lower in female KO mice (Fig. 7E). Corroborating the gene expression data, quantification of hydroxyproline (marker of fibrosis) and malondialdehyde (marker of oxidative stress) revealed no significant differences between genotypes (Fig. 7F-G).

The liver weight of female KO mice was modestly increased (Fig. 7F), but when normalized to body weight it was not significantly different from WT controls (2.76 % in WT and 2.60% in KO, $P = 0.24$). Histological analysis and quantification revealed no differences in hepatic lipid content (% lipid area) between genotypes (Fig. 7G). Interestingly, although hepatic fat content was similar between genotypes, the expression of lipogenic genes (e.g., *Fasn*) was lower and fat catabolism genes (e.g., *Cpt2*, *Ppara*, *Acadl*, *Acadm*, *Acad11*, *Acadvl*) was higher in female KO mice (Fig. 7H). The expression of genes associated with inflammation, fibrosis, ER and oxidative stress in liver were not significantly different between genotypes (Fig. 7H). Consistent with the gene expression data, quantification of hydroxyproline (marker of fibrosis) in the liver revealed no significant difference between genotypes (Fig. 7K). The *Ctrp10* KO female mice, however, had higher levels of malondialdehyde (a marker of oxidative stress) in the liver, suggesting a modest increase in oxidative stress (Fig. 7L). Altogether, these data

indicate that obesity is largely uncoupled from inflammation, fibrosis, ER and oxidative stress in *Ctrp10* KO female

mice.

Transcriptomic and pathway changes associated with the metabolically healthy obesity phenotype in *Ctrp10*

KO female mice.

To define the specific mechanisms mediating the female-specific effects of *Ctrp10* ablation on favorable metabolic

outcomes, four major metabolic tissues (gWAT, iWAT, liver, skeletal muscle) from female WT and KO mice fed a

HFD were subjected to RNA-sequencing. Comparison of differentially expressed genes (DEGs) via limma (71)

showed robust changes across tissues, with the largest changes seen in the liver (Fig. 8A-D). In liver, gWAT, and

muscle, we observed comparable numbers of DEGs that were up- and down-regulated, whereas more genes were

transcriptionally suppressed in the iWAT of *Ctrp10* KO female mice (Fig. 8E, top panel). While significant DEGs

were identified in all 4 tissues, only limited overlap was observed between the DEGs in each tissue (Fig. 8E,

bottom panel). Gene set enrichment analyses of the DEGs highlighted distinct and shared processes up- or down-

regulated across the four tissues (Fig. 8F). Pathways and processes related to lipid metabolism and estrogen

receptor were the top-ranked up-regulated enrichments across tissues (Fig. 8F, top panel), whereas processes related

to blood clotting and lipoprotein metabolism were the top-ranked down-regulated enrichments (Fig. 8F, bottom

panel).

Of the DEGs, we found significant changes across tissues in relevant classes of genes that encode proteins

involved in gene expression (e.g., transcription factors), signaling (e.g., receptors), tissue crosstalk (e.g., secreted

proteins), and metabolism (Fig. 9). Notably, the nuclear receptor, *Nr1d1* (also known as *Rev-Erbα*), is the only gene

consistently suppressed across all four tissues (liver, gWAT, iWAT, and muscle) of *Ctrp10* KO female mice (Fig.

8E lower panel and Fig. 9). Interestingly, global deletion of *Nr1d1* promotes lipogenesis, adipose tissue expansion,

and obesity (72, 73). Although the whole-body and adipose-specific *Nr1d1* KO mice fed with HFD become

markedly obese, the obesity is not accompanied by insulin resistance, adipose tissue inflammation and fibrosis (73,

74). Like the *Ctrp10* KO female mice, HFD-fed mice lacking Nr1d1 can maintain a relatively healthy metabolic

profile despite being strikingly obese. Since only male mice were used in these previous studies, we do not know

whether female mice lacking Nr1d1 would also exhibit similar insulin-sensitive obesity phenotype. In WT mice,

Nr1d1 acts as a transcriptional repressor of metabolic genes whose expression are upregulated by high-fat feeding;

loss of Nr1d1 is thought to result in the de-repression of these genes, leading to greater lipid synthesis and fat mass

accrual in response to caloric excess (74). Thus, the suppression of *Nr1d1* expression—mimicking Nr1d1

deficiency—across tissues in HFD-fed *Ctrp10* KO female mice may contribute to benign fat mass expansion

without the accompanying adipose tissue fibrosis, inflammation, and oxidative stress.

Among the upregulated genes, *Fgf21* and *Fgf1* was significantly elevated in the liver of *Ctrp10* KO female

mice (Fig. 9A). FGF21 is an hepatokine known to improve systemic insulin sensitivity and to promote a favorable

metabolic profile in diet-induced obese mice (75). Likewise, FGF1 has been shown to dampen hepatic glucose

output by suppressing adipose lipolysis (76), improve systemic insulin sensitivity by reducing adipose

inflammation (77), and alleviate hepatic steatosis, inflammation, and insulin resistance (78). Thus, upregulated

expression of *Fgf21* and *Fgf1* in *Ctrp10* KO female mice could contribute to the MHO phenotype. In addition, the

upregulated hepatic expression of IL-22 receptor (*Il22ra1*) in *Ctrp10* KO female mice (Fig. 9A) may confer

protection against obesity-associated fatty liver, inflammation and fibrosis (79-81). Further, a marked increase in

uncoupling protein 3 (*Ucp3*) and Krüppel-like factor 15 (*Klf15*) expression in the skeletal muscle (Fig. 9C) may

promote lipid utilization and help mitigate lipid-induced insulin resistance in *Ctrp10* KO female mice (82, 83).

Taken together, these combined changes—at the level of gene expression and biological pathways and processes

across tissues—acting in concert likely contribute to the apparently healthy obesity phenotype seen in the KO

female mice.

Conservation of mouse DEG co-correlation in humans highlights sex-specific gene connectivity.

Next, we asked whether the female-specific transcriptomic effects across tissues were conserved in humans. To

address this, we analyzed transcriptional co-correlation of mouse DEG (Fig. 10) orthologues in GTEx (84),

consisting of 210 males and 100 females filtered for comparison of gene expression across tissues (85, 86).

Hierarchical clustering of transcriptional correlation of the orthologous DEGs among 4 metabolic tissues—

subcutaneous and visceral white adipose tissue, liver, and skeletal muscle—showed differing patterns of gene

connectivity between females (Fig. 10A) and males (Fig. 10B). When grouped according to sex in each tissue, the

degree of sex-specific gene correlation pairs of DEGs orthologues showed the most significant differences in

subcutaneous adipose tissue (Fig. 10C). Given the whole-body metabolic effects of *Ctrp10* ablation in mice, we

further examined the degree of sex-dependent DEG co-correlation across metabolic tissues. This analysis showed

that human orthologue genes in subcutaneous adipose tissue (Fig. 10D, top row) and liver (Fig. 10D, third row) also

exhibited highly significant sex differences in their transcriptional correlation with other DEGs across key

metabolic tissues (Fig. 10D). These analyses highlight the sex-specificity of CTRP10 DEG orthologues in humans,

suggest possible sex-biased mechanisms of tissue crosstalk, and overall underscores the conservation of the sex-

dependent metabolic function of CTRP10.

DISCUSSION

Our current study has established a novel function for CTRP10 in modulating body weight in a sex-specific

manner. When mice were fed a control LFD, female *Ctrp10*-KO mice developed obesity with age; increased

adiposity, however, did not impair insulin action and glucose and lipid metabolism. When challenged with an

obesogenic diet, female *Ctrp10*-KO mice gained weight rapidly. Despite having strikingly higher adiposity and

weighing ~10-11 g (~28%) more, female KO mice fed a HFD exhibited a metabolic profile largely

indistinguishable from the much leaner WT controls. Although female KO mice had higher fasting glucose and

insulin levels, direct assessments of glucose metabolism and insulin sensitivity by glucose and insulin tolerance

tests, however, revealed no differences between genotypes. Except having lower fasting ketone (β-hydroxybutyrate)

levels, the fasting lipid profile, as well as VLDL-TG and HDL-cholesterol levels, of *Ctrp10*-KO female mice

resembled the WT controls. The hepatic fat content was also comparable between genotypes. Global transcriptomic

profiling across different fat depots and liver did not reveal gene expression signatures associated with elevated

inflammation, fibrosis, and ER and oxidative stress. Altogether, these findings suggest that CTRP10 deficiency

promotes obesity in females but it also uncouples obesity from insulin resistance, dyslipidemia, steatosis,

inflammation, and oxidative stress. Thus, *Ctrp10*-KO female mice represent a novel model of female obesity with

largely preserved insulin sensitivity and metabolic health.

Our findings help inform ongoing studies on metabolically healthy obese (MHO) humans (87-92). Because

the criteria used to define MHO differs between studies, there is an ongoing debate regarding the prevalence of

MHO and what fraction of the MHO population is insulin-sensitive and metabolically healthy (93, 94).

Nevertheless, among the obese individuals, there clearly exists a subgroup that maintains long-term normal insulin

sensitivity and does not appear to develop any component of the metabolic syndrome (93). MHO is observed in

both sexes, but more common in females (6). The underlying mechanism(s) that uncouple obesity from adverse

metabolic health in MHO is not well understood (7, 89, 93). It is currently unknown whether males and females

with MHO use similar or distinct mechanism to maintain insulin sensitivity and metabolic health. Our findings in

Ctrp10-KO female, but not male, mice suggest that there may be female-biased mechanism that prevents metabolic

deterioration in the face of obesity, thus underscoring the utility of the *Ctrp10*-KO mice as a female mouse model

of MHO.

347   Obesity is frequently associated with insulin resistance, dyslipidemia, fatty liver, oxidative stress, and

chronic low-grade inflammation (67, 95, 96). The mechanisms that link obesity to metabolic dysfunctions are

complex and multifactorial. There are limited number of mouse models described where obesity is uncoupled from

insulin resistance and metabolic health (97-100); in some studies, however, only male mice were used or that the

sex of the animals was not specified. In the case of aP2/FABP4 KO male mice, the uncoupling of obesity from

insulin resistance was attributed to a marked decrease in TNF-α expression in adipose tissue (98). In the case of

adiponectin overexpression in leptin-deficient (*ob*/*ob*) male and female mice, a dramatic expansion of the

subcutaneous fat pad is thought to promote lipid sequestration in adipose compartment, thus preventing ectopic

lipid deposition in non-adipose tissues (e.g., liver, pancreas, muscle) that would otherwise induce insulin resistance

(97). Massive obesity with preserved insulin sensitivity is also observed in leptin-deficient (*ob*/*ob*) male and female

mice overexpressing the mitochondrial membrane protein, mitoNEET (100). The benign obesity is attributed to the

inhibition of iron transport into mitochondria by mitoNEET, leading to reduced mitochondrial activity, fatty acid

oxidation, and oxidative stress (100). In the Brd2 hypomorphic mice, severe obesity with lower blood glucose and

enhanced glucose tolerance is due to a combination of hyperinsulinemia and marked reduction in macrophage

infiltration into fat depot (101). Lastly, in male mice fed a high starch diet, uncoupling of obesity from insulin

resistance is associated with lower ceramide levels in liver and skeletal muscle (99). In all these cases, the

uncoupling of obesity from metabolic dysfunction is seen in either male mice only (female mice were not included)

or both sexes. These previous studies suggest that multiple mechanisms, not mutually exclusive, can contribute to the MHO phenotype in different mouse models.

In our study, loss of CTRP10 in female mice largely uncoupled obesity from insulin resistance, dyslipidemia, steatosis, inflammation, and ER and oxidative stress. The preservation of insulin sensitivity in *Ctrp10* KO female mice is due, at least in part, to the absence of obesity-linked adipose and liver inflammation, fibrosis, and oxidative stress. These phenotypes associated with a favorable metabolic profile are also observed in MHO individuals (88, 102). A healthy adaptive remodeling of white adipose tissues in response to caloric surfeit helps preserve the storage and secretory function of adipocytes (103). The expansion of benign adipose tissues further serves to sequester circulating lipids and prevent their ectopic deposition in non-adipose tissue (e.g., liver and skeletal muscle) which can impair insulin action (104, 105). The MHO phenotype seen in *Ctrp10* KO female mice reinforce the notion that adipose tissue health, rather than abundance, is an important determinant of metabolic health in obesity.

Because lipidomic analysis was not performed—a limitation of this study—we do not know whether *Ctrp10*-KO female mice have reduced ceramide or diacylglycerol levels in liver and skeletal muscle, two lipid species known to antagonize insulin action (106, 107). However, our global transcriptomic and pathway enrichment analysis across visceral and subcutaneous fat depots, liver, and skeletal muscle highlighted the relevant up- and down-regulated pathways and processes (e.g., lipid and lipoprotein metabolism, signaling) that may contribute to the MHO phenotype in *Ctrp10*-KO female mice. How these changes across tissues help to suppress the deleterious effects of obesity and maintain an apparently healthy metabolic profile in *Ctrp10*-KO female mice remains to be fully understood. Part of the mechanism may be attributable to the suppression of *Nr1d1* and the upregulated expression of *Fgf1*, *Fgf21*, *Il22ra1*, *Ucp3*, *Klf15*. Altered expression of these genes are known to reduce obesity-linked inflammation, oxidative stress, steatosis, and insulin resistance.

In many single-gene KO mouse models where both sexes are examined, it is often the males that show a more pronounced metabolic phenotype. It is known that C57BL/6 female mice generally gain significantly less weight on HFD compared to male mice (108). Therefore, it is intriguing that *Ctrp10*-KO female mice became obese

on a control LFD and gained weight rapidly when fed an obesogenic diet. After twelve weeks on HFD, the body weight of female KO mice was approaching that of WT male mice fed the same diet. What mechanism underlies the sexually dimorphic requirement of CTRP10 for body weight control? We know that the obesity phenotype was not attributed to differences in food intake, physical activity level, body temperature, and energy expenditure between WT and KO female mice. We assume that the methods used to quantify these physiologic parameters are sensitive enough to detect small differences that can give rise to divergent body weight over time. Quantification of fecal output and fecal energy content also revealed no differences between genotypes. Thus, loss of CTRP10 did not affect macronutrient intake and absorption. Although we cannot fully rule out the CNS function of CTRP10/C1QL2 (62), our data do not support a central role for CTRP10 in modulating food intake behavior, locomotor activity, and energy expenditure that affect body weight in female mice.

It is known that reduced estrogen level by ovariectomy or blocking estrogen action in estrogen receptor (ERα) KO mice will cause obesity and metabolic dysfunction in female mice fed a HFD (109-111). Conversely, estradiol supplementation decreases HFD-induced weight gain and improves glucose tolerance and insulin sensitivity (112-114). Estrogen also has the effect of reducing food intake, and promoting physical activity and energy expenditure (115-119). In our study, loss of CTRP10 promotes obesity without altering food intake, physical activity, and energy expenditure. While estrogen's role cannot be completely ruled out, the fact that female *Ctrp10*-KO mice developed obesity with largely preserved metabolic health suggest that factors other than altered estrogen level contribute to the insulin-sensitive obesity phenotype. Future studies are warranted to uncover what factor(s) is causally contributing to obesity in female mice lacking CTRP10.

The sex-dependent effects of CTRP10 on metabolism and tissue transcriptomes appear to be conserved in humans. When the human orthologues of the mouse DEGs were used to interrogate the GTEx data, clear patterns of gene connectivity within and across metabolic tissues in females and males were observed, with the strongest sex-specific gene correlations seen in subcutaneous adipose tissue and liver. These findings provide further evidence that CTRP10 modulates tissue transcriptome in a sex-dependent manner. Further, our analyses of sex-dependent DEG co-correlation across metabolic tissues also suggest possible sex-biased mechanisms of inter-organ metabolic signaling between adipose tissue and liver.

CTRP10 has been previously shown to bind to the adhesion GPCR, brain angiogenesis inhibitor-3

(Bai3/Adgrb3) (120). Bai3 is expressed in the brain and peripheral tissues, and it is a promiscuous GPCR that can

bind to multiple ligands. In addition to CTRP10 (C1QL2), Bai3 also binds to CTRP11 (C1QL4), CTRP13

(C1QL3), CTRP14 (C1QL1), neuronal pentraxins, and reticulon 4 (RTN4) receptor (56, 58, 59, 120-122). A

constitutive, whole-body KO of Bai3 mouse models have recently been generated (123, 124). Both male and female

Bai3 KO mice fed a standard chow have significantly lower body weight, beginning at weaning (3 weeks old) and

continue into adulthood (123, 124). Lower body weight in *Bai3* KO mice of either sex is attributed to a reduction in

both lean and fat mass, and is associated with higher energy expenditure and reduced food intake in male mice

(123). The impact of Bai3 deficiency on systemic metabolism in response to a high-fat diet was not examined. The

Ctrp10 KO mice do not phenocopy the phenotypes of the *Bai3* KO mice. When fed a control low-fat diet, the body

weight, food intake, and energy expenditure of *Ctrp10* KO male mice were indistinguishable from WT controls. In

striking contrast to *Bai3* KO mice, female *Ctrp10* KO mice fed a low-fat diet began to gain more weight around 20

weeks of age, and by 40 weeks had become visibly obese. While we did not rule out CTRP10-Bai3 signaling axis

in modulating energy metabolism in peripheral tissues, our findings in *Ctrp10* KO mice suggest that future works

are needed to establish the molecular mechanisms that mediate the systemic metabolic function of CTRP10.

Several limitations of our current study are noted. We use a constitutive whole-body KO mouse model of

CTRP10 to interrogate its function. It is unknown whether CTRP10 has a role during development that may

influence sex-dependent postnatal weight gain with age or in response to a high-caloric diet. Future studies using

conditional KO of *Ctrp10* gene in adult mice can help address this issue. Although the lack of differences in food

intake, physical activity, and energy expenditure between genotypes do not support a central role of CTRP10 in

mediating the metabolic phenotypes of *Ctrp10*-KO female mice, a brain-specific KO of *Ctrp10* gene is needed to

definitively rule this out. Our phenotypic analyses in the context of metabolism are relatively comprehensive but

not exhaustive. Although the metabolic profile of obese *Ctrp10*-KO female mice was largely indistinguishable from

the much leaner WT controls, we do not know if some related aspect of metabolic health (e.g., blood pressure and

heart function) may be altered in the absence of CTRP10 which we did not examine.

In summary, we have established the physiologic role and requirement of CTRP10 in modulating body

weight in a female-specific manner. Importantly, loss of CTRP10 largely uncouples obesity from insulin resistance

and metabolic dysfunction. The CTRP10-deficient female mice represent a unique and valuable model to help

dissect female-biased mechanisms that help preserve metabolic health in the face of positive energy balance and

increased adiposity.

MATERIALS AND METHODS

Mouse models

Eight-week-old mouse tissues (gonadal and inguinal white adipose tissues, interscapular brown adipose tissue,

liver, heart, skeletal muscle, kidney, pancreas, cerebellum, cortex, hippocampus, hindbrain, and hypothalamus)

from C57BL/6J male mice (The Jackson Laboratory, Bar Harbor, ME) were collected from fasted and refed

experiments as we have previously described (125). For the fasted group, food was removed for 16 h (beginning 10

h into the light cycle), and mice were euthanized 2 h into the light cycle. For the refed group, mice were fasted for

16 h and refed with chow pellets for 2 h before being euthanized. Tissues (white and brown adipose tissues, liver,

whole brain, kidney, spleen, heart, skeletal muscle, pancreas, small intestine, and colon) from C57BL/6J male mice

fed a low-fat diet (LFD) or a high-fat diet (HFD) for 12 weeks were also collected as we have previously described

(125).

The *Ctrp10/C1ql2*-KO mice (C57BL/6NCrl-*C1ql2^em1(IMPC)Mbp*/Mmucd; stock number 050587-UCD) were

generated using the CRISPR-cas9 method at UC Davis. The two guide RNAs (gRNA) used were 5'-CCGGCGCC

GCTCCACCATTACCT-3' and 5'-TCAGGCCACCCCATCCCCATCGG-3'. The *Ctrp10* gene consists of 2 exons.

The entire protein coding region spanning exon 1 and 2 was deleted. This KO strategy ensures a complete null

allele for *Ctrp10*. The KO mice were maintained on a C57BL6/6J genetic background. Genotyping primers for WT

allele were forward (m10-Com-F) 5'-TGTCGGGCTCTTCGACTCTCCA-3' and reverse (m10-WT-R) 5'-

GCATCTCGT AGTGAGCCGCTCC-3'. The size of the WT band was 360 bp. Genotyping primers for the *Ctrp10*

KO allele were forward (m10-Com-F) 5'-TGTCGGGCTCTTCGACTCTCCA-3' and reverse (m10-Mut-R1) 5'-

GTCCAATCAGCT TTCTCAAGTCTGG-3'. The size of the KO band was 422 bp. The genotyping PCR

parameters were as follows: 94°C for 5 min, followed by 10 cycles of (94°C for 10 sec, 65°C for 15 sec, 72°C for

30 sec), then 25 cycles of (94°C for 10 sec, 55°C for 15 sec, 72°C for 30 sec), and lastly 72°C for 5 min. Due to the

presence of GC rich sequences, 7% DMSO was included in the PCR genotyping reaction. Mice were generated by

intercrossing *Ctrp10* heterozygous (+/-) mice, supplemented with intercrossing WT or KO mice. *Ctrp10* KO (-/-)

and WT (+/+) controls were housed in polycarbonate cages on a 12-h light–dark photocycle with ad libitum access

to water and food. Mice were fed either a control low-fat diet (LFD; 10% kcal derived from fat; # D12450B;

Research Diets, New Brunswick, NJ) or a high-fat diet (HFD; 60% kcal derived from fat; #D12492, Research

Diets). LFD was provided for the duration of the study, beginning at 5 weeks of age; HFD was provided for 14

weeks, beginning at 6-7 weeks of age. At termination of study, all mice were fasted for 2 h and euthanized. Tissues

were collected, snap-frozen in liquid nitrogen, and kept at -80°C until analysis. All mouse protocols (protocol #

MO22M367) were approved by the Institutional Animal Care and Use Committee of the Johns Hopkins University

School of Medicine. All animal experiments were conducted in accordance with the National Institute of Health

guidelines and followed the standards established by the Animal Welfare Acts.

Body composition analysis

Body composition analyses for total fat, lean mass, and water content were determined using a quantitative

magnetic resonance instrument (Echo-MRI-100, Echo Medical Systems, Waco, TX) at the Mouse Phenotyping

Core facility at Johns Hopkins University School of Medicine.

Indirect calorimetry

LFD- or HFD-fed WT and *Ctrp10* KO male and female mice were used for simultaneous assessments of daily body

weight change, food intake (corrected for spillage), physical activity, and whole-body metabolic profile in an open

flow indirect calorimeter (Comprehensive Laboratory Animal Monitoring System, CLAMS; Columbus

Instruments, Columbus, OH) as previously described (25). In brief, data were collected for three days to confirm

mice were acclimatized to the calorimetry chambers (indicated by stable body weights, food intakes, and diurnal

metabolic patterns), and data were analyzed from the fourth day. Rates of oxygen consumption ($\dot{V}_{O2}$; mL·kg^{-1}·h^{-1})

and carbon dioxide production ($\dot{V}_{CO2}$; mL·kg^{-1}h^{-1}) in each chamber were measured every 24 min throughout the

studies. Respiratory exchange ratio (RER = $\dot{V}_{CO2}/\dot{V}_{O2}$) was calculated by CLAMS software (version 5.66) to

estimate relative oxidation of carbohydrates (RER = 1.0) versus fats (RER = 0.7), not accounting for protein

oxidation. Energy expenditure (EE) was calculated as EE= $\dot{V}_{O2}\times$ [3.815 + (1.232 × RER)] and normalized to lean

mass. Because normalizing to lean mass can potentially lead to overestimation of EE, we also performed ANCOVA

analysis on EE using body weight as a covariate (126). Physical activities were measured by infrared beam breaks

in the metabolic chamber.

Measurements of 24 h food intake

To independently confirm the food intake data collected in the metabolic cage (CLAMS), we also performed 24 h

food intake measurements manually. All mice were singly housed, with wire mesh flooring inserts over a piece of

cage paper on the bottom of the cage. A known weight of food pellets was given to each mouse. Twenty-four hours

later, the leftover food pellets remaining on the flooring insert, along with any spilled crumbs on the cage paper,

were collected and weighed. Thus, food intake was corrected for spillage.

Glucose, insulin, pyruvate, and lipid tolerance tests

All tolerance tests were conducted as previously described (21, 24, 29). For glucose tolerance tests (GTTs), mice

were fasted for 6 h before glucose injection. Glucose (Sigma, St. Louis, MO) was reconstituted in saline (0.9 g

NaCl/L), sterile-filtered, and injected intraperitoneally (i.p.) at 1 mg/g body weight (i.e., 10 μL/g body weight).

Blood glucose was measured at 0, 15, 30, 60, and 120 min after glucose injection using a glucometer (NovaMax

Plus, Billerica, MA). For insulin tolerance tests (ITTs), food was removed 2 h before insulin injection. 6.5 μL of

insulin stock (4 mg/mL; Gibco) was diluted in 10 mL of saline, sterile-filtered, and injected i.p. at 0.75 U/kg body

weight (i.e., 10 μL/g body weight). Blood glucose was measured at 0, 15, 30, 60, and 90 min after insulin injection

using a glucometer (NovaMax Plus).

Fasting-Refeeding insulin tests

Mice were fasted overnight (~16 h) then reintroduced to food as described (26). Blood glucose was monitored at

the 16 h fast time point (time = 0 h refed) and at 1 and 2 hours into the refeeding process. Serum was collected at

the 16 h fast and 2 h refed time points for insulin ELISA, as well as for the quantification of triglyceride,

cholesterol, non-esterified free fatty acids (NEFA), and β-hydroxybutyrate levels.

Blood and tissue chemistry analysis

Tail vein blood samples were allowed to clot on ice and then centrifuged for 10 min at 10,000 x g. Serum samples

were stored at -80°C until analyzed. Serum triglycerides (TG) and cholesterol levels were measured according to

manufacturer's instructions using an Infinity kit (Thermo Fisher Scientific, Middletown, VA). Non-esterified free

fatty acids (NEFA) were measured using a Wako kit (Wako Chemicals, Richmond, VA). Serum β-hydroxybutyrate

(ketone) concentrations were measured with a StanBio Liquicolor kit (StanBio Laboratory, Boerne, TX). Serum

insulin levels were measured by ELISA according to manufacturer's instructions (Crystal Chem, Elk Grove

Village, IL; cat # 90080). Hydroxyproline assay (Sigma Aldrich, MAK008) was used to quantify total collagen

content in liver and adipose tissues according to the manufacturer's instructions. Lipid peroxidation levels (marker

of oxidative stress) in the liver and adipose tissues were assessed by the quantification of malondialdehyde (MDA)

via Thiobarbituric Acid Reactive Substances (TBARS) assay (Cayman Chemical, 700870) according to the

manufacturer's instructions.

Serum lipoprotein triglyceride and cholesterol analysis by FPLC

Food was removed for 2-4 hr (in the light cycle) prior to blood collection. Sera collected from mice were pooled (n

= 6-7/genotype) and sent to the Mouse Metabolism Core at Baylor College of Medicine for analysis. Serum

samples were first fractionated by fast protein liquid chromatography (FPLC). A total of 45 fractions were

collected, and TG and cholesterol in each fraction was quantified.

Histology and quantification

Inguinal (subcutaneous) white adipose tissue (iWAT), gonadal (visceral) white adipose tissue (gWAT), and liver

were dissected and fixed in formalin. Paraffin embedding, tissue sectioning, and staining with hematoxylin and

eosin were performed at the Pathology Core facility at Johns Hopkins University School of Medicine. Images were

captured with a Keyence BZ-X700 All-in-One fluorescence microscope (Keyence Corp., Itasca, IL). Adipocyte

(gWAT and iWAT) cross-sectional area (CSA), as well as the total area covered by lipid droplets in hepatocytes

were measured on hematoxylin and eosin-stained slides using ImageJ software (127). For CSA measurements, all

cells in one field of view at 100X magnification per tissue section per mouse were analyzed. Image capturing and

quantifications were carried out blinded to genotype.

Fecal bomb calorimetry and assessment of fecal parameters

Fecal pellet frequency and average fecal pellet weight were monitored by housing each mouse singly in clean cages with a wire mesh sitting on top of a cutout cardboard that lay at the bottom of the cage for fecal collection. The number of fecal pellets and their total weight was recorded at the end of 24 h period. Additional fecal pellets collected for 3 full days were combined and shipped to the University of Michigan Animal Phenotyping Core for fecal bomb calorimetry. Briefly, fecal samples were dried overnight at 50°C prior to weighing and grinding them to powder. Each sample was mixed with wheat flour (90% wheat flour, 10% sample) and formed into 1.0 g pellet, which was then secured into the firing platform and surrounded by 100% oxygen. The bomb was lowered into a water reservoir and ignited to release heat into the surrounding water. These data were used to calculate fecal pellet frequency (bowel movements/day), average fecal pellet weight (g/bowel movement), fecal energy (cal/g feces), and total fecal energy (kcal/day).

Tissue library preparation and RNA sequencing

Total RNA was isolated from tissues using Trizol reagent (Thermo Fisher Scientific) according to the manufacturer's instructions. Library preparation and bulk RNA sequencing of liver, skeletal muscle (gastrocnemius), gonadal white adipose tissue (gWAT), and inguinal white adipose tissue (iWAT) of HFD-fed *Ctrp10*-KO female mice and WT controls were performed by Novogene (Sacramento, California, USA) on an Illumina platform (NovaSeq 6000) and pair-end reads were generated. Sample size: 6 WT and 6 KO for each tissue. All raw sequencing files are available from the NIH Sequence Read Archive (SRA) accession PRJNA971939.

Mouse RNA-Sequencing analysis.

Transcript features were assembled from raw fastq files and aligned to the current version of mouse transcriptome (Mus_musculus.GRCm39.cdna) using kallisto -aln (128). Version-specific Ensembl transcript IDs were linked to gene symbols using biomart. Estimated counts were log normalized and filtered for a limit of sum >5 across all samples. Logistic regressions comparing WT vs KO samples across tissues were performed using limma (71). Differential expression results were visualized using available R packages in CRAN: ggplot2, ggVennDiagram and pheatmap. Gene set enrichment analyses of the DEGs were performed using Enrichr (129). Scripts for analyses and

587 visualization are available at https://github.com/Leandromvelez/CTRP10-Manuscript-DEG-Sex-specific-

588 connectivities-and-integration/

Human sex difference analysis.

All the datasets and scripts to perform analyses are available at: https://github.com/Leandromvelez/CTRP10-

Manuscript-DEG-Sex-specific-connectivities-and-integration/. Male and Female human data were obtained from

Genotype-Tissue Expression (GTEx) (84) and filtered for sufficient comparison of inter-tissue transcript correlation

as described (85, 86). CTRP10/C1QL2 and other human orthologues for mouse differentially expressed genes

(DEGs) were identified by intersecting mouse gene symbols with known human orthologues from the vertebrate

homology resource at Mouse Genome Informatics (MGI) (130). Co-correlation between all human orthologues

DEGs were calculated in either self-reported male or female subjects in GTEx using the bicorAndPvalue() function

in Weighted Genetic Coexpression Network Analysis (WGCNA) package (131). To compare sex-differences of

regression coefficients, wilcoxon t-tests were compared between coefficients using the R package ggpubr.

Quantitative real-time PCR

Total RNA was isolated from tissues using Trizol reagent (Thermo Fisher Scientific). Purified RNA was reverse

transcribed using an iScript cDNA Synthesis Kit (Bio-rad). Real-time quantitative PCR analysis was performed on

a CFX Connect Real-Time System (Bio-rad) using iTaq^{TM} Universal SYBR Green Supermix (Bio-rad) according to

manufacturer's instructions. Data were normalized to the stable housekeeping gene *β-actin* or *36B4* (encoding the

acidic ribosomal phosphoprotein P0) and expressed as relative mRNA levels using the ΔΔCt method (132). Real-

time qPCR primers used to assess *Ctrp10* expression across mouse tissues were: *Ctrp10* forward, 5'-

CGGCTTCATGAC ACTTCCTGA-3' and reverse, 5'-AGCAGGGATGTGTCTTTTCCA-3'. qPCR primers used

to confirm the absence of *Ctrp10* in KO mice were: forward (qPCR-m10-F2), 5'-

CACGTACCACATTCTCATGCG-3' and reverse (qPCR-m10-R1), 5'-TCGTAATTCTGGTCCGCGTC-3'.

Statistical analyses

613 Sample size is indicated in figure and/or figure legend. All results are expressed as mean ± standard error of the

614 mean (SEM). Statistical analysis was performed with Prism 9 software (GraphPad Software, San Diego, CA). Data

were analyzed with two-tailed Student's *t*-tests, one-way ANOVA or two-way ANOVA (with Sidak's post hoc tests). 2-way ANOVA was used for body weight over time, fasting-refeeding response, and all tolerance tests. $P < 0.05$ was considered statistically significant.

ACKNOWLEDGEMENTS

This work was supported by the National Institutes of Health (DK084171 to GWW, HL138193 and DK130640 to MMS). D.C.S. is supported by an NIH T32 training grant (HL007534). The fecal bomb calorimetry analysis was performed at the University of Michigan Animal Phenotyping Core, supported by center grants 1U2CDK135066-01 (Mi-MPMOD) and DK020572 (MDRC). The FPLC/serum analyses were performed the Mouse Metabolism and Phenotyping Core (MMPC) at the Baylor College of Medicine, supported by NIH grants (DK114356 and UM1HG006348).

AUTHOR CONTRIBUTIONS

FC, GWW contributed to the experimental design; FC, DCS, MS, SA, and GWW performed the experiments; FC, DCS, SA, LMV, MMS, and GWW analyzed and interpreted the data; GWW drafted the paper with inputs and edits from all authors.

COMPETING INTERESTS

We declare that none of the authors has a conflict of interest.

DATA AVAILABILITY

High-throughput sequencing data from this study have been submitted to the NCBI Sequence Read Archive under accession number PRJNA971939. All processed datasets used and R scripts to reproduce analyses are freely available at: https://github.com/Leandromvelez/CTRP10-Manuscript-DEG-Sex-specific-connectivities-and-integration/.

REFERENCES

1. Collaboration, N. C. D. R. F. (2017) Worldwide trends in body-mass index, underweight, overweight, and
obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128.9 million
children, adolescents, and adults. *Lancet* **390**, 2627-2642
2. Flier, J. S. (2023) Moderating "the great debate": The carbohydrate-insulin vs. the energy balance models of
obesity. *Cell Metab* **35**, 737-741
3. Bluher, M. (2019) Obesity: global epidemiology and pathogenesis. *Nat Rev Endocrinol* **15**, 288-298
4. Bouchard, C. (2021) Genetics of Obesity: What We Have Learned Over Decades of Research. *Obesity (Silver*
Spring) **29**, 802-820
5. Loos, R. J. F., and Kilpelainen, T. O. (2018) Genes that make you fat, but keep you healthy. *J Intern Med*
284, 450-463
6. van Vliet-Ostaptchouk, J. V., Nuotio, M. L., Slagter, S. N., Doiron, D., Fischer, K., Foco, L., Gaye, A.,
Gogele, M., Heier, M., Hiekkalinna, T., Joensuu, A., Newby, C., Pang, C., Partinen, E., Reischl, E.,
Schwienbacher, C., Tammesoo, M. L., Swertz, M. A., Burton, P., Ferretti, V., Fortier, I., Giepmans, L.,
Harris, J. R., Hillege, H. L., Holmen, J., Jula, A., Kootstra-Ros, J. E., Kvaloy, K., Holmen, T. L., Mannisto,
S., Metspalu, A., Midthjell, K., Murtagh, M. J., Peters, A., Pramstaller, P. P., Saaristo, T., Salomaa, V., Stolk,
R. P., Uusitupa, M., van der Harst, P., van der Klauw, M. M., Waldenberger, M., Perola, M., and
Wolffenbuttel, B. H. (2014) The prevalence of metabolic syndrome and metabolically healthy obesity in
Europe: a collaborative analysis of ten large cohort studies. *BMC Endocr Disord* **14**, 9
7. Bluher, M. (2020) Metabolically Healthy Obesity. *Endocr Rev* **41**
8. Priest, C., and Tontonoz, P. (2019) Inter-organ cross-talk in metabolic syndrome. *Nat Metab* **1**, 1177-1188
9. Seldin, M. M., Tan, S. Y., and Wong, G. W. (2014) Metabolic function of the CTRP family of hormones. *Rev*
Endocr Metab Disord **15**, 111-123
10. Wong, G. W., Wang, J., Hug, C., Tsao, T. S., and Lodish, H. F. (2004) A family of Acrp30/adiponectin
structural and functional paralogs. *Proc Natl Acad Sci U S A* **101**, 10302-10307
11. Seldin, M. M., Peterson, J. M., Byerly, M. S., Wei, Z., and Wong, G. W. (2012) Myonectin (CTRP15), a
novel myokine that links skeletal muscle to systemic lipid homeostasis. *J Biol Chem* **287**, 11968-11980
12. Wei, Z., Peterson, J. M., Lei, X., Cebotaru, L., Wolfgang, M. J., Baldeviano, G. C., and Wong, G. W. (2012)
C1q/TNF-related protein-12 (CTRP12), a novel adipokine that improves insulin sensitivity and glycemic
control in mouse models of obesity and diabetes. *J Biol Chem* **287**, 10301-10315
13. Wei, Z., Peterson, J. M., and Wong, G. W. (2011) Metabolic regulation by C1q/TNF-related protein-13
(CTRP13): activation of AMP-activated protein kinase and suppression of fatty acid-induced JNK signaling.
J Biol Chem **286**, 15652-15665
14. Wei, Z., Seldin, M. M., Natarajan, N., Djemal, D. C., Peterson, J. M., and Wong, G. W. (2013) C1q/Tumor
Necrosis Factor-related Protein 11 (CTRP11), a Novel Adipose Stroma-derived Regulator of Adipogenesis. *J*
Biol Chem **288**, 10214-10229
15. Wong, G. W., Krawczyk, S. A., Kitidis-Mitrokostas, C., Ge, G., Spooner, E., Hug, C., Gimeno, R., and
Lodish, H. F. (2009) Identification and characterization of CTRP9, a novel secreted glycoprotein, from
adipose tissue that reduces serum glucose in mice and forms heterotrimers with adiponectin. *FASEB J* **23**,
241-258
16. Wong, G. W., Krawczyk, S. A., Kitidis-Mitrokostas, C., Revett, T., Gimeno, R., and Lodish, H. F. (2008)
Molecular, biochemical and functional characterizations of C1q/TNF family members: adipose-tissue-
selective expression patterns, regulation by PPAR-gamma agonist, cysteine-mediated oligomerizations,
combinatorial associations and metabolic functions. *Biochem J* **416**, 161-177

17.   Ghai, R., Waters, P., Roumenina, L. T., Gadjeva, M., Kojouharova, M. S., Reid, K. B., Sim, R. B., and
Kishore, U. (2007) C1q and its growing family. *Immunobiology* **212**, 253-266

18.   Ressl, S., Vu, B. K., Vivona, S., Martinelli, D. C., Sudhof, T. C., and Brunger, A. T. (2015) Structures of
C1q-like proteins reveal unique features among the C1q/TNF superfamily. *Structure* **23**, 688-699

19.   Lei, X., Rodriguez, S., Petersen, P. S., Seldin, M. M., Bowman, C. E., Wolfgang, M. J., and Wong, G. W.
(2016) Loss of CTRP5 improves insulin action and hepatic steatosis. *Am J Physiol Endocrinol Metab* **310**,
E1036-1052

20.   Lei, X., Seldin, M. M., Little, H. C., Choy, N., Klonisch, T., and Wong, G. W. (2017) C1q/TNF-related
protein 6 (CTRP6) links obesity to adipose tissue inflammation and insulin resistance. *J Biol Chem* **292**,
14836-14850

21.   Lei, X., and Wong, G. W. (2019) C1q/TNF-related protein 2 (CTRP2) deletion promotes adipose tissue
lipolysis and hepatic triglyceride secretion. *J Biol Chem* **294**, 15638-15649

22.   Little, H. C., Rodriguez, S., Lei, X., Tan, S. Y., Stewart, A. N., Sahagun, A., Sarver, D. C., and Wong, G. W.
(2019) Myonectin deletion promotes adipose fat storage and reduces liver steatosis. *FASEB J* **33**, 8666-8687

23.   Petersen, P. S., Lei, X., Wolf, R. M., Rodriguez, S., Tan, S. Y., Little, H. C., Schweitzer, M. A., Magnuson,
T. H., Steele, K. E., and Wong, G. W. (2017) CTRP7 deletion attenuates obesity-linked glucose intolerance,
adipose tissue inflammation, and hepatic stress. *Am J Physiol Endocrinol Metab* **312**, E309-E325

24.   Rodriguez, S., Lei, X., Petersen, P. S., Tan, S. Y., Little, H. C., and Wong, G. W. (2016) Loss of CTRP1
disrupts glucose and lipid homeostasis. *Am J Physiol Endocrinol Metab* **311**, E678-E697

25.   Sarver, D. C., Stewart, A. N., Rodriguez, S., Little, H. C., Aja, S., and Wong, G. W. (2020) Loss of CTRP4
alters adiposity and food intake behaviors in obese mice. *Am J Physiol Endocrinol Metab* **319**, E1084-E1100

26.   Sarver, D. C., Xu, C., Aja, S., and Wong, G. W. (2022) CTRP14 inactivation alters physical activity and food
intake response to fasting and refeeding. *Am J Physiol Endocrinol Metab* **322**, E480-E493

27.   Sarver, D. C., Xu, C., Carreno, D., Arking, A., Terrillion, C. E., Aja, S., and Wong, G. W. (2022) CTRP11
contributes modestly to systemic metabolism and energy balance. *FASEB J* **36**, e22347

28.   Tan, S. Y., Lei, X., Little, H. C., Rodriguez, S., Sarver, D. C., Cao, X., and Wong, G. W. (2020) CTRP12
ablation differentially affects energy expenditure, body weight, and insulin sensitivity in male and female
mice. *Am J Physiol Endocrinol Metab* **319**, E146-E162

29.   Tan, S. Y., Little, H. C., Lei, X., Li, S., Rodriguez, S., and Wong, G. W. (2016) Partial deficiency of CTRP12
alters hepatic lipid metabolism. *Physiol Genomics* **48**, 936-949

30.   Wei, Z., Lei, X., Petersen, P. S., Aja, S., and Wong, G. W. (2014) Targeted deletion of C1q/TNF-related
protein 9 increases food intake, decreases insulin sensitivity, and promotes hepatic steatosis in mice. *Am J*
Physiol Endocrinol Metab **306**, E779-790

31.   Wolf, R. M., Lei, X., Yang, Z. C., Nyandjo, M., Tan, S. Y., and Wong, G. W. (2016) CTRP3 deficiency
reduces liver size and alters IL-6 and TGFbeta levels in obese mice. *Am J Physiol Endocrinol Metab* **310**,
E332-345

32.   Peterson, J. M., Seldin, M. M., Tan, S. Y., and Wong, G. W. (2014) CTRP2 overexpression improves insulin
and lipid tolerance in diet-induced obese mice. *PLoS One* **9**, e88535

33.   Peterson, J. M., Seldin, M. M., Wei, Z., Aja, S., and Wong, G. W. (2013) CTRP3 attenuates diet-induced
hepatic steatosis by regulating triglyceride metabolism. *Am J Physiol Gastrointest Liver Physiol* **305**, G214-
224.

34.   Peterson, J. M., Wei, Z., Seldin, M. M., Byerly, M. S., Aja, S., and Wong, G. W. (2013) CTRP9 transgenic
mice are protected from diet-induced obesity and metabolic dysfunction. *Am J Physiol Regul Integr Comp*
Physiol **305**, R522-533

35.   Appari, M., Breitbart, A., Brandes, F., Szaroszyk, M., Froese, N., Korf-Klingebiel, M., Mohammadi, M. M.,
Grund, A., Scharf, G. M., Wang, H., Zwadlo, C., Fraccarollo, D., Schrameck, U., Nemer, M., Wong, G. W.,
Katus, H. A., Wollert, K. C., Muller, O. J., Bauersachs, J., and Heineke, J. (2017) C1q-TNF-Related Protein-
9 Promotes Cardiac Hypertrophy and Failure. *Circ Res* **120**, 66-77

36.   Zheng, Q., Yuan, Y., Yi, W., Lau, W. B., Wang, Y., Wang, X., Sun, Y., Lopez, B. L., Christopher, T. A.,
Peterson, J. M., Wong, G. W., Yu, S., Yi, D., and Ma, X. L. (2011) C1q/TNF-Related Proteins, A Family of
Novel Adipokines, Induce Vascular Relaxation Through the Adiponectin Receptor-1/AMPK/eNOS/Nitric
Oxide Signaling Pathway. *Arterioscler Thromb Vasc Biol* **31**, 2616-2623

37.   Kambara, T., Ohashi, K., Shibata, R., Ogura, Y., Maruyama, S., Enomoto, T., Uemura, Y., Shimizu, Y.,
Yuasa, D., Matsuo, K., Miyabe, M., Kataoka, Y., Murohara, T., and Ouchi, N. (2012) CTRP9 protein

protects against myocardial injury following ischemia-reperfusion through AMP-activated protein kinase
(AMPK)-dependent mechanism. *J Biol Chem* **287**, 18965-18973

38. Kambara, T., Shibata, R., Ohashi, K., Matsuo, K., Hiramatsu-Ito, M., Enomoto, T., Yuasa, D., Ito, M., Hayakawa, S., Ogawa, H., Aprahamian, T., Walsh, K., Murohara, T., and Ouchi, N. (2015) C1q/Tumor Necrosis Factor-Related Protein 9 Protects against Acute Myocardial Injury through an Adiponectin Receptor I-AMPK-Dependent Mechanism. *Mol Cell Biol* **35**, 2173-2185

39. Kanemura, N., Shibata, R., Ohashi, K., Ogawa, H., Hiramatsu-Ito, M., Enomoto, T., Yuasa, D., Ito, M., Hayakawa, S., Otaka, N., Murohara, T., and Ouchi, N. (2017) C1q/TNF-related protein 1 prevents neointimal formation after arterial injury. *Atherosclerosis* **257**, 138-145

40. Ogawa, H., Ohashi, K., Ito, M., Shibata, R., Kanemura, N., Yuasa, D., Kambara, T., Matsuo, K., Hayakawa, S., Hiramatsu-Ito, M., Otaka, N., Kawanishi, H., Yamaguchi, S., Enomoto, T., Abe, T., Kaneko, M., Takefuji, M., Murohara, T., and Ouchi, N. (2019) Adipolin/CTRP12 protects against pathological vascular remodeling through suppression of smooth muscle cell growth and macrophage inflammatory response. *Cardiovasc Res*

41. Otaka, N., Shibata, R., Ohashi, K., Uemura, Y., Kambara, T., Enomoto, T., Ogawa, H., Ito, M., Kawanishi, H., Maruyama, S., Joki, Y., Fujikawa, Y., Narita, S., Unno, K., Kawamoto, Y., Murate, T., Murohara, T., and Ouchi, N. (2018) Myonectin Is an Exercise-Induced Myokine That Protects the Heart From Ischemia-Reperfusion Injury. *Circ Res* **123**, 1326-1338

42. Uemura, Y., Shibata, R., Ohashi, K., Enomoto, T., Kambara, T., Yamamoto, T., Ogura, Y., Yuasa, D., Joki, Y., Matsuo, K., Miyabe, M., Kataoka, Y., Murohara, T., and Ouchi, N. (2013) Adipose-derived factor CTRP9 attenuates vascular smooth muscle cell proliferation and neointimal formation. *FASEB J* **27**, 25-33

43. Yuasa, D., Ohashi, K., Shibata, R., Mizutani, N., Kataoka, Y., Kambara, T., Uemura, Y., Matsuo, K., Kanemura, N., Hayakawa, S., Hiramatsu-Ito, M., Ito, M., Ogawa, H., Murate, T., Murohara, T., and Ouchi, N. (2016) C1q/TNF-related protein-1 functions to protect against acute ischemic injury in the heart. *FASEB J* **30**, 1065-1075

44. Sun, Y., Yi, W., Yuan, Y., Lau, W. B., Yi, D., Wang, X., Wang, Y., Su, H., Gao, E., Koch, W. J., and Ma, X. L. (2013) C1q/Tumor Necrosis Factor-Related Protein-9, a Novel Adipocyte-Derived Cytokine, Attenuates Adverse Remodeling in the Ischemic Mouse Heart via Protein Kinase A Activation. *Circulation* **128**, S113-120

45. Yi, W., Sun, Y., Yuan, Y., Lau, W. B., Zheng, Q., Wang, X., Wang, Y., Shang, X., Gao, E., Koch, W. J., and Ma, X. L. (2012) C1q/tumor necrosis factor-related protein-3, a newly identified adipokine, is a novel antiapoptotic, proangiogenic, and cardioprotective molecule in the ischemic mouse heart. *Circulation* **125**, 3159-3169

46. Han, S., Jeong, A. L., Lee, S., Park, J. S., Buyanravjikh, S., Kang, W., Choi, S., Park, C., Han, J., Son, W. C., Yoo, K. H., Cheong, J. H., Oh, G. T., Lee, W. Y., Kim, J., Suh, S. H., Lee, S. H., Lim, J. S., Lee, M. S., and Yang, Y. (2018) C1q/TNF-alpha--Related Protein 1 (CTRP1) Maintains Blood Pressure Under Dehydration Conditions. *Circ Res* **123**, e5-e19

47. Lee, S. M., Lee, J. W., Kim, I., Woo, D. C., Pack, C. G., Sung, Y. H., Baek, I. J., Jung, C. H., Kim, Y. H., and Ha, C. H. (2022) Angiogenic adipokine C1q-TNF-related protein 9 ameliorates myocardial infarction via histone deacetylase 7-mediated MEF2 activation. *Sci Adv* **8**, eabq0898

48. Rodriguez, S., Little, H. C., Daneshpajouhnejad, P., Fenaroli, P., Tan, S. Y., Sarver, D. C., Delannoy, M., Talbot, C. C., Jr., Jandu, S., Berkowitz, D. E., Pluznick, J. L., Rosenberg, A. Z., and Wong, G. W. (2021) Aging and chronic high-fat feeding negatively affect kidney size, function, and gene expression in CTRP1-deficient mice. *Am J Physiol Regul Integr Comp Physiol* **320**, R19-R35

49. Rodriguez, S., Little, H. C., Daneshpajouhnejad, P., Shepard, B. D., Tan, S. Y., Wolfe, A., Cheema, M. U., Jandu, S., Woodward, O. M., Talbot, C. C., Jr., Berkowitz, D. E., Rosenberg, A. Z., Pluznick, J. L., and Wong, G. W. (2020) Late-onset renal hypertrophy and dysfunction in mice lacking CTRP1. *FASEB J* **34**, 2657-2676

50. Kirketerp-Moller, N., Bayarri-Olmos, R., Krogfelt, K. A., and Garred, P. (2020) C1q/TNF-Related Protein 6 Is a Pattern Recognition Molecule That Recruits Collectin-11 from the Complement System to Ligands. *J Immunol* **204**, 1598-1606

51. Lahav, R., Haim, Y., Bhandarkar, N. S., Levin, L., Chalifa-Caspi, V., Sarver, D., Sahagun, A., Maixner, N., Kovesh, B., Wong, G. W., and Rudich, A. (2021) CTRP6 rapidly responds to acute nutritional changes, regulating adipose tissue expansion and inflammation in mice. *Am J Physiol Endocrinol Metab* **321**, E702-E713

800   52.   Ayyagari, R., Mandal, M. N., Karoukis, A. J., Chen, L., McLaren, N. C., Lichter, M., Wong, D. T.,
801   Hitchcock, P. F., Caruso, R. C., Moroi, S. E., Maumenee, I. H., and Sieving, P. A. (2005) Late-onset macular
802   degeneration and long anterior lens zonules result from a CTRP5 gene mutation. *Invest Ophthalmol Vis Sci*
803   **46**, 3363-3371

804   53.   Hayward, C., Shu, X., Cideciyan, A. V., Lennon, A., Barran, P., Zareparsi, S., Sawyer, L., Hendry, G.,
805   Dhillon, B., Milam, A. H., Luthert, P. J., Swaroop, A., Hastie, N. D., Jacobson, S. G., and Wright, A. F.
806   (2003) Mutation in a short-chain collagen gene, CTRP5, results in extracellular deposit formation in late-
807   onset retinal degeneration: a genetic model for age-related macular degeneration. *Hum Mol Genet* **12**, 2657-
808   2667

809   54.   Luo, Y., Wu, X., Ma, Z., Tan, W., Wang, L., Na, D., Zhang, G., Yin, A., Huang, H., Xia, D., Zhang, Y., Shi,
810   X., and Wang, L. (2016) Expression of the novel adipokine C1qTNF-related protein 4 (CTRP4) suppresses
811   colitis and colitis-associated colorectal cancer in mice. *Cell Mol Immunol* **13**, 688-699

812   55.   Youngstrom, D. W., Zondervan, R. L., Doucet, N. R., Acevedo, P. K., Sexton, H. E., Gardner, E. A.,
813   Anderson, J. S., Kushwaha, P., Little, H. C., Rodriguez, S., Riddle, R. C., Kalajzic, I., Wong, G. W., and
814   Hankenson, K. D. (2020) CTRP3 Regulates Endochondral Ossification and Bone Remodeling During
815   Fracture Healing. *J Orthop Res* **38**, 996-1006

816   56.   Hamoud, N., Tran, V., Aimi, T., Kakegawa, W., Lahaie, S., Thibault, M. P., Pelletier, A., Wong, G. W., Kim,
817   I. S., Kania, A., Yuzaki, M., Bouvier, M., and Cote, J. F. (2018) Spatiotemporal regulation of the GPCR
818   activity of BAI3 by C1qL4 and Stabilin-2 controls myoblast fusion. *Nature communications* **9**, 4470

819   57.   Cho, Y., Kim, H. S., Kang, D., Kim, H., Lee, N., Yun, J., Kim, Y. J., Lee, K. M., Kim, J. H., Kim, H. R.,
820   Hwang, Y. I., Jo, C. H., and Kim, J. H. (2021) CTRP3 exacerbates tendinopathy by dysregulating tendon
821   stem cell differentiation and altering extracellular matrix composition. *Sci Adv* **7**, eabg6069

822   58.   Kakegawa, W., Mitakidis, N., Miura, E., Abe, M., Matsuda, K., Takeo, Y. H., Kohda, K., Motohashi, J.,
823   Takahashi, A., Nagao, S., Muramatsu, S., Watanabe, M., Sakimura, K., Aricescu, A. R., and Yuzaki, M.
824   (2015) Anterograde C1ql1 signaling is required in order to determine and maintain a single-winner climbing
825   fiber in the mouse cerebellum. *Neuron* **85**, 316-329

826   59.   Sigoillot, S. M., Iyer, K., Binda, F., Gonzalez-Calvo, I., Talleur, M., Vodjdani, G., Isope, P., and Selimi, F.
827   (2015) The Secreted Protein C1QL1 and Its Receptor BAI3 Control the Synaptic Connectivity of Excitatory
828   Inputs Converging on Cerebellar Purkinje Cells. *Cell reports* **10**, 820-832

829   60.   Martinelli, D. C., Chew, K. S., Rohlmann, A., Lum, M. Y., Ressl, S., Hattar, S., Brunger, A. T., Missler, M.,
830   and Sudhof, T. C. (2016) Expression of C1ql3 in Discrete Neuronal Populations Controls Efferent Synapse
831   Numbers and Diverse Behaviors. *Neuron* **91**, 1034-1051

832   61.   Sarver, D. C., Xu, C., Cheng, Y., Terrillion, C. E., and Wong, G. W. (2021) CTRP4 ablation impairs
833   associative learning and memory. *FASEB J* **35**, e21910

834   62.   Matsuda, K., Budisantoso, T., Mitakidis, N., Sugaya, Y., Miura, E., Kakegawa, W., Yamasaki, M., Konno,
835   K., Uchigashima, M., Abe, M., Watanabe, I., Kano, M., Watanabe, M., Sakimura, K., Aricescu, A. R., and
836   Yuzaki, M. (2016) Transsynaptic Modulation of Kainate Receptor Functions by C1q-like Proteins. *Neuron*
837   **90**, 752-767

838   63.   Huggett, S. B., and Stallings, M. C. (2020) Genetic Architecture and Molecular Neuropathology of Human
839   Cocaine Addiction. *J Neurosci* **40**, 5300-5313

840   64.   Unroe, K. A., Glover, M. E., Shupe, E. A., Feng, N., and Clinton, S. M. (2021) Perinatal SSRI Exposure
841   Disrupts G Protein-coupled Receptor BAI3 in Developing Dentate Gyrus and Adult Emotional Behavior:
842   Relevance to Psychiatric Disorders. *Neuroscience* **471**, 32-50

843   65.   Iijima, T., Miura, E., Watanabe, M., and Yuzaki, M. (2010) Distinct expression of C1q-like family mRNAs
844   in mouse brain and biochemical characterization of their encoded proteins. *Eur J Neurosci* **31**, 1606-1615

845   66.   Vishvanath, L., and Gupta, R. K. (2019) Contribution of adipogenesis to healthy adipose tissue expansion in
846   obesity. *J Clin Invest* **129**, 4022-4031

847   67.   Hotamisligil, G. S. (2006) Inflammation and metabolic disorders. *Nature* **444**, 860-867

848   68.   Sun, K., Tordjman, J., Clement, K., and Scherer, P. E. (2013) Fibrosis and adipose tissue dysfunction. *Cell*
849   *Metab* **18**, 470-477

850   69.   Masschelin, P. M., Cox, A. R., Chernis, N., and Hartig, S. M. (2019) The Impact of Oxidative Stress on
851   Adipose Tissue Energy Balance. *Front Physiol* **10**, 1638

852   70.   Hotamisligil, G. S. (2010) Endoplasmic reticulum stress and the inflammatory basis of metabolic disease.
853   *Cell* **140**, 900-917

71. Ritchie, M. E., Phipson, B., Wu, D., Hu, Y., Law, C. W., Shi, W., and Smyth, G. K. (2015) limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res* **43**, e47

72. Delezie, J., Dumont, S., Dardente, H., Oudart, H., Grechez-Cassiau, A., Klosen, P., Teboul, M., Delaunay, F., Pevet, P., and Challet, E. (2012) The nuclear receptor REV-ERBalpha is required for the daily balance of carbohydrate and lipid metabolism. *FASEB J* **26**, 3321-3335

73. Hand, L. E., Usan, P., Cooper, G. J., Xu, L. Y., Ammori, B., Cunningham, P. S., Aghamohammadzadeh, R., Soran, H., Greenstein, A., Loudon, A. S., Bechtold, D. A., and Ray, D. W. (2015) Adiponectin induces A20 expression in adipose tissue to confer metabolic benefit. *Diabetes* **64**, 128-136

74. Hunter, A. L., Pelekanou, C. E., Barron, N. J., Northeast, R. C., Grudzien, M., Adamson, A. D., Downton, P., Cornfield, T., Cunningham, P. S., Billaud, J. N., Hodson, L., Loudon, A. S., Unwin, R. D., Iqbal, M., Ray, D. W., and Bechtold, D. A. (2021) Adipocyte NR1D1 dictates adipose tissue expansion during obesity. *Elife* **10**

75. Xu, J., Lloyd, D. J., Hale, C., Stanislaus, S., Chen, M., Sivits, G., Vonderfecht, S., Hecht, R., Li, Y. S., Lindberg, R. A., Chen, J. L., Jung, D. Y., Zhang, Z., Ko, H. J., Kim, J. K., and Veniant, M. M. (2009) Fibroblast growth factor 21 reverses hepatic steatosis, increases energy expenditure, and improves insulin sensitivity in diet-induced obese mice. *Diabetes* **58**, 250-259

76. Sancar, G., Liu, S., Gasser, E., Alvarez, J. G., Moutos, C., Kim, K., van Zutphen, T., Wang, Y., Huddy, T. F., Ross, B., Dai, Y., Zepeda, D., Collins, B., Tilley, E., Kolar, M. J., Yu, R. T., Atkins, A. R., van Dijk, T. H., Saghatelian, A., Jonker, J. W., Downes, M., and Evans, R. M. (2022) FGF1 and insulin control lipolysis by convergent pathways. *Cell Metab* **34**, 171-183 e176

77. Zhao, L., Fan, M., Zhao, L., Yun, H., Yang, Y., Wang, C., and Qin, D. (2020) Fibroblast growth factor 1 ameliorates adipose tissue inflammation and systemic insulin resistance via enhancing adipocyte mTORC2/Rictor signal. *J Cell Mol Med* **24**, 12813-12825

78. Fan, L., Ding, L., Lan, J., Niu, J., He, Y., and Song, L. (2019) Fibroblast Growth Factor-1 Improves Insulin Resistance via Repression of JNK-Mediated Inflammation. *Front Pharmacol* **10**, 1478

79. Kong, X., Feng, D., Wang, H., Hong, F., Bertola, A., Wang, F. S., and Gao, B. (2012) Interleukin-22 induces hepatic stellate cell senescence and restricts liver fibrosis in mice. *Hepatology* **56**, 1150-1159

80. Yang, L., Zhang, Y., Wang, L., Fan, F., Zhu, L., Li, Z., Ruan, X., Huang, H., Wang, Z., Huang, Z., Huang, Y., Yan, X., and Chen, Y. (2010) Amelioration of high fat diet induced liver lipogenesis and hepatic steatosis by interleukin-22. *J Hepatol* **53**, 339-347

81. Wang, X., Ota, N., Manzanillo, P., Kates, L., Zavala-Solorio, J., Eidenschenk, C., Zhang, J., Lesch, J., Lee, W. P., Ross, J., Diehl, L., van Bruggen, N., Kolumam, G., and Ouyang, W. (2014) Interleukin-22 alleviates metabolic disorders and restores mucosal immunity in diabetes. *Nature* **514**, 237-241

82. Choi, C. S., Fillmore, J. J., Kim, J. K., Liu, Z. X., Kim, S., Collier, E. F., Kulkarni, A., Distefano, A., Hwang, Y. J., Kahn, M., Chen, Y., Yu, C., Moore, I. K., Reznick, R. M., Higashimori, T., and Shulman, G. I. (2007) Overexpression of uncoupling protein 3 in skeletal muscle protects against fat-induced insulin resistance. *J Clin Invest* **117**, 1995-2003

83. Fan, L., Sweet, D. R., Prosdocimo, D. A., Vinayachandran, V., Chan, E. R., Zhang, R., Ilkayeva, O., Lu, Y., Keerthy, K. S., Booth, C. E., Newgard, C. B., and Jain, M. K. (2021) Muscle Kruppel-like factor 15 regulates lipid flux and systemic metabolic homeostasis. *J Clin Invest* **131**

84. Consortium, G. T. (2013) The Genotype-Tissue Expression (GTEx) project. *Nat Genet* **45**, 580-585

85. Velez, L. M., Van, C., Moore, T., Zhou, Z., Johnson, C., Hevener, A. L., and Seldin, M. M. (2022) Genetic variation of putative myokine signaling is dominated by biological sex and sex hormones. *Elife* **11**

86. Sarver, D. C., Xu, C., Velez, L. M., Aja, S., Jaffe, A. E., Seldin, M. M., Reeves, R. H., and Wong, G. W. (2023) Dysregulated systemic metabolism in a Down syndrome mouse model. *Mol Metab* **68**, 101666

87. Kloting, N., Fasshauer, M., Dietrich, A., Kovacs, P., Schon, M. R., Kern, M., Stumvoll, M., and Bluher, M. (2010) Insulin-sensitive obesity. *Am J Physiol Endocrinol Metab* **299**, E506-515

88. Primeau, V., Coderre, L., Karelis, A. D., Brochu, M., Lavoie, M. E., Messier, V., Sladek, R., and Rabasa-Lhoret, R. (2011) Characterizing the profile of obese patients who are metabolically healthy. *Int J Obes (Lond)* **35**, 971-981

89. Samocha-Bonet, D., Chisholm, D. J., Tonks, K., Campbell, L. V., and Greenfield, J. R. (2012) Insulin-sensitive obesity in humans - a 'favorable fat' phenotype? *Trends Endocrinol Metab* **23**, 116-124

90. Stefan, N., Kantartzis, K., Machann, J., Schick, F., Thamer, C., Rittig, K., Balletshofer, B., Machicao, F., Fritsche, A., and Haring, H. U. (2008) Identification and characterization of metabolically benign obesity in humans. *Arch Intern Med* **168**, 1609-1616

91. Calori, G., Lattuada, G., Piemonti, L., Garancini, M. P., Ragogna, F., Villa, M., Mannino, S., Crosignani, P.,
Bosi, E., Luzi, L., Ruotolo, G., and Perseghin, G. (2011) Prevalence, metabolic features, and prognosis of
metabolically healthy obese Italian individuals: the Cremona Study. *Diabetes Care* **34**, 210-215
92. Aguilar-Salinas, C. A., Garcia, E. G., Robles, L., Riano, D., Ruiz-Gomez, D. G., Garcia-Ulloa, A. C.,
Melgarejo, M. A., Zamora, M., Guillen-Pineda, L. E., Mehta, R., Canizales-Quinteros, S., Tusie Luna, M. T.,
and Gomez-Perez, F. J. (2008) High adiponectin concentrations are associated with the metabolically healthy
obese phenotype. *J Clin Endocrinol Metab* **93**, 4075-4079
93. Smith, G. I., Mittendorfer, B., and Klein, S. (2019) Metabolically healthy obesity: facts and fantasies. *J Clin
Invest* **129**, 3978-3989
94. Pataky, Z., Bobbioni-Harsch, E., and Golay, A. (2010) Open questions about metabolically normal obesity.
Int J Obes (Lond) **34 Suppl 2**, S18-23
95. Samuel, V. T., Petersen, K. F., and Shulman, G. I. (2010) Lipid-induced insulin resistance: unravelling the
mechanism. *Lancet* **375**, 2267-2277
96. Cohen, J. C., Horton, J. D., and Hobbs, H. H. (2011) Human fatty liver disease: old questions and new
insights. *Science* **332**, 1519-1523
97. Kim, J. Y., van de Wall, E., Laplante, M., Azzara, A., Trujillo, M. E., Hofmann, S. M., Schraw, T., Durand,
J. L., Li, H., Li, G., Jelicks, L. A., Mehler, M. F., Hui, D. Y., Deshaies, Y., Shulman, G. I., Schwartz, G. J.,
and Scherer, P. E. (2007) Obesity-associated improvements in metabolic profile through expansion of
adipose tissue. *J Clin Invest* **117**, 2621-2637
98. Hotamisligil, G. S., Johnson, R. S., Distel, R. J., Ellis, R., Papaioannou, V. E., and Spiegelman, B. M. (1996)
Uncoupling of obesity from insulin resistance through a targeted mutation in aP2, the adipocyte fatty acid
binding protein. *Science* **274**, 1377-1379
99. Brandon, A. E., Small, L., Nguyen, T. V., Suryana, E., Gong, H., Yassmin, C., Hancock, S. E., Pulpitel, T.,
Stonehouse, S., Prescott, L., Kebede, M. A., Yau, B., Quek, L. E., Kowalski, G. M., Bruce, C. R., Turner, N.,
and Cooney, G. J. (2022) Insulin sensitivity is preserved in mice made obese by feeding a high starch diet.
Elife **11**
100. Kusminski, C. M., Holland, W. L., Sun, K., Park, J., Spurgin, S. B., Lin, Y., Askew, G. R., Simcox, J. A.,
McClain, D. A., Li, C., and Scherer, P. E. (2012) MitoNEET-driven alterations in adipocyte mitochondrial
activity reveal a crucial adaptive process that preserves insulin sensitivity in obesity. *Nat Med* **18**, 1539-1549
101. Wang, F., Liu, H., Blanton, W. P., Belkina, A., Lebrasseur, N. K., and Denis, G. V. (2009) Brd2 disruption in
mice causes severe obesity without Type 2 diabetes. *Biochem J* **425**, 71-83
102. Karelis, A. D., Faraj, M., Bastard, J. P., St-Pierre, D. H., Brochu, M., Prud'homme, D., and Rabasa-Lhoret, R.
(2005) The metabolically healthy but obese individual presents a favorable inflammation profile. *J Clin
Endocrinol Metab* **90**, 4145-4150
103. Sun, K., Kusminski, C. M., and Scherer, P. E. (2011) Adipose tissue remodeling and obesity. *J Clin Invest*
121, 2094-2101
104. Shulman, G. I. (2014) Ectopic fat in insulin resistance, dyslipidemia, and cardiometabolic disease. *N Engl J
Med* **371**, 1131-1141
105. Virtue, S., and Vidal-Puig, A. (2010) Adipose tissue expandability, lipotoxicity and the Metabolic Syndrome-
-an allostatic perspective. *Biochim Biophys Acta* **1801**, 338-349
106. Samuel, V. T., and Shulman, G. I. (2012) Mechanisms for insulin resistance: common threads and missing
links. *Cell* **148**, 852-871
107. Summers, S. A., Chaurasia, B., and Holland, W. L. (2019) Metabolic Messengers: ceramides. *Nat Metab* **1**,
1051-1058
108. Yang, Y., Smith, D. L., Jr., Keating, K. D., Allison, D. B., and Nagy, T. R. (2014) Variations in body weight,
food intake and body composition after long-term high-fat diet feeding in C57BL/6J mice. *Obesity (Silver
Spring)* **22**, 2147-2155
109. Rogers, N. H., Perfield, J. W., 2nd, Strissel, K. J., Obin, M. S., and Greenberg, A. S. (2009) Reduced energy
expenditure and increased inflammation are early events in the development of ovariectomy-induced obesity.
Endocrinology **150**, 2161-2168
110. Hong, J., Stubbins, R. E., Smith, R. R., Harvey, A. E., and Nunez, N. P. (2009) Differential susceptibility to
obesity between male, female and ovariectomized female mice. *Nutr J* **8**, 11
111. Heine, P. A., Taylor, J. A., Iwamoto, G. A., Lubahn, D. B., and Cooke, P. S. (2000) Increased adipose tissue
in male and female estrogen receptor-alpha knockout mice. *Proc Natl Acad Sci U S A* **97**, 12729-12734

112. Bryzgalova, G., Lundholm, L., Portwood, N., Gustafsson, J. A., Khan, A., Efendic, S., and Dahlman-Wright, K. (2008) Mechanisms of antidiabetogenic and body weight-lowering effects of estrogen in high-fat diet-fed mice. *Am J Physiol Endocrinol Metab* **295**, E904-912

113. Stubbins, R. E., Holcomb, V. B., Hong, J., and Nunez, N. P. (2012) Estrogen modulates abdominal adiposity and protects female mice from obesity and impaired glucose tolerance. *Eur J Nutr* **51**, 861-870

114. Yonezawa, R., Wada, T., Matsumoto, N., Morita, M., Sawakawa, K., Ishii, Y., Sasahara, M., Tsuneki, H., Saito, S., and Sasaoka, T. (2012) Central versus peripheral impact of estradiol on the impaired glucose metabolism in ovariectomized mice on a high-fat diet. *Am J Physiol Endocrinol Metab* **303**, E445-456

115. Mauvais-Jarvis, F., Clegg, D. J., and Hevener, A. L. (2013) The role of estrogens in control of energy balance and glucose homeostasis. *Endocr Rev* **34**, 309-338

116. Correa, S. M., Newstrom, D. W., Warne, J. P., Flandin, P., Cheung, C. C., Lin-Moore, A. T., Pierce, A. A., Xu, A. W., Rubenstein, J. L., and Ingraham, H. A. (2015) An estrogen-responsive module in the ventromedial hypothalamus selectively drives sex-specific activity in females. *Cell reports* **10**, 62-74

117. Butera, P. C. (2010) Estradiol and the control of food intake. *Physiol Behav* **99**, 175-180

118. Xu, Y., Nedungadi, T. P., Zhu, L., Sobhani, N., Irani, B. G., Davis, K. E., Zhang, X., Zou, F., Gent, L. M., Hahner, L. D., Khan, S. A., Elias, C. F., Elmquist, J. K., and Clegg, D. J. (2011) Distinct hypothalamic neurons mediate estrogenic effects on energy homeostasis and reproduction. *Cell Metab* **14**, 453-465

119. Krause, W. C., Rodriguez, R., Gegenhuber, B., Matharu, N., Rodriguez, A. N., Padilla-Roger, A. M., Toma, K., Herber, C. B., Correa, S. M., Duan, X., Ahituv, N., Tollkuhn, J., and Ingraham, H. A. (2021) Oestrogen engages brain MC4R signalling to drive physical activity in female mice. *Nature* **599**, 131-135

120. Bolliger, M. F., Martinelli, D. C., and Sudhof, T. C. (2011) The cell-adhesion G protein-coupled receptor BAI3 is a high-affinity receptor for C1q-like proteins. *Proc Natl Acad Sci U S A* **108**, 2534-2539

121. Sticco, M. J., Pena Palomino, P. A., Lukacsovich, D., Thompson, B. L., Foldy, C., Ressl, S., and Martinelli, D. C. (2021) C1QL3 promotes cell-cell adhesion by mediating complex formation between ADGRB3/BAI3 and neuronal pentraxins. *FASEB J* **35**, e21194

122. Wang, J., Miao, Y., Wicklein, R., Sun, Z., Wang, J., Jude, K. M., Fernandes, R. A., Merrill, S. A., Wernig, M., Garcia, K. C., and Sudhof, T. C. (2021) RTN4/NoGo-receptor binding to BAI adhesion-GPCRs regulates neuronal development. *Cell* **184**, 5869-5885 e5825

123. Alsharif, H., Latimer, M. N., Perez, K. C., Alexander, J., Rahman, M. M., Challa, A. K., Kim, J. A., Ramanadham, S., Young, M., and Bhatnagar, S. (2023) Loss of Brain Angiogenesis Inhibitor-3 (BAI3) G-Protein Coupled Receptor in Mice Regulates Adaptive Thermogenesis by Enhancing Energy Expenditure. *Metabolites* **13**

124. Shiu, F. H., Wong, J. C., Bhattacharya, D., Kuranaga, Y., Parag, R. R., Alsharif, H. A., Bhatnagar, S., Van Meir, E. G., and Escayg, A. (2023) Generation and initial characterization of mice lacking full-length BAI3 (ADGRB3) expression. *Basic Clin Pharmacol Toxicol*

125. Rodriguez, S., Stewart, A. N., Lei, X., Cao, X., Little, H. C., Fong, V., Sarver, D. C., and Wong, G. W. (2019) PRADC1: a novel metabolic-responsive secretory protein that modulates physical activity and adiposity. *FASEB J* **33**, 14748-14759

126. Tschop, M. H., Speakman, J. R., Arch, J. R., Auwerx, J., Bruning, J. C., Chan, L., Eckel, R. H., Farese, R. V., Jr., Galgani, J. E., Hambly, C., Herman, M. A., Horvath, T. L., Kahn, B. B., Kozma, S. C., Maratos-Flier, E., Muller, T. D., Munzberg, H., Pfluger, P. T., Plum, L., Reitman, M. L., Rahmouni, K., Shulman, G. I., Thomas, G., Kahn, C. R., and Ravussin, E. (2012) A guide to analysis of mouse energy metabolism. *Nat Methods* **9**, 57-63

127. Schneider, C. A., Rasband, W. S., and Eliceiri, K. W. (2012) NIH Image to ImageJ: 25 years of image analysis. *Nat Methods* **9**, 671-675

128. Bray, N. L., Pimentel, H., Melsted, P., and Pachter, L. (2016) Near-optimal probabilistic RNA-seq quantification. *Nat Biotechnol* **34**, 525-527

129. Kuleshov, M. V., Jones, M. R., Rouillard, A. D., Fernandez, N. F., Duan, Q., Wang, Z., Koplev, S., Jenkins, S. L., Jagodnik, K. M., Lachmann, A., McDermott, M. G., Monteiro, C. D., Gundersen, G. W., and Ma'ayan, A. (2016) Enrichr: a comprehensive gene set enrichment analysis web server 2016 update. *Nucleic Acids Res* **44**, W90-97

130. Alliance of Genome Resources, C. (2020) Alliance of Genome Resources Portal: unified model organism research platform. *Nucleic Acids Res* **48**, D650-D658

131. Langfelder, P., and Horvath, S. (2008) WGCNA: an R package for weighted correlation network analysis. *BMC Bioinformatics* **9**, 559

1017 132. Schmittgen, T. D., and Livak, K. J. (2008) Analyzing real-time PCR data by the comparative C(T) method.
1018   *Nat Protoc* **3**, 1101-1108
1019 133. Sievers, F., and Higgins, D. G. (2018) Clustal Omega for making accurate alignments of many protein
1020   sequences. *Protein Sci* **27**, 135-145

1027 **FIGURE LEGENDS**
1028

Figure 1. Nutritional regulation of *Ctrp10* expression. (A) Sequence alignment of full-length human (GenBank #

NP_872334), mouse (NP_997116), chicken (XP_046777733), xenopus frog (XP_031749381), and zebrafish

(XP_001920705) CTRP10/C1ql2 using Clustal-omega (133). Identical amino acids are shaded black and similar

amino acids are shaded grey. Gaps are indicated by dash lines. Signal peptide, collagen domain with characteristic

Gly-X-Y repeats, and the C-terminal globular C1q domain are indicated. **(B)** *Ctrp10* expression across different

mouse tissues (*n* = 10). **(C)** Expression of *Ctrp10* across mouse tissues in response to an overnight (16 h) fast or

fasting followed by 2 h refeeding. **(D)** Expression of *Ctrp10* across mouse tissues in response to a high-fat diet

(HFD) for 12 weeks or a control low-fat diet (LFD). **(E)** Generation of *Ctrp10* knockout (KO) mice. The entire

protein coding region in exon 1 and 2 of *Ctrp10* was deleted using CRISPR/Cas9 method and confirmed with DNA

sequencing. **(F)** Wild-type (WT) and KO alleles were confirmed by PCR genotyping. **(G)** The complete loss of

Ctrp10 transcript in KO mice was confirmed in mouse cortex, one of the tissues with high *Ctrp10* expression (WT,

n = 5; KO, *n* = 5). All expression levels were normalized to *β-actin*. All data are presented as mean ± S.E.M. * *P* <

0.05; ** *P* < 0.01; *** *P* < 0.001.

1042

Figure 2. *Ctrp10*-KO mice fed a low-fat diet have normal body weight and energy balance. (A-B) Body weight

(A) and body composition analysis **(B)** of fat mass, % fat mass (relative to body weight), lean mass, and % lean

mass of WT (*n* = 17) and KO (*n* = 14) male mice at 18 weeks of age. **(C-D)** Body weight **(C)** and body composition

analysis **(D)** of fat mass, % fat mass (relative to body weight), lean mass, and % lean mass of WT (*n* = 9) and KO

(*n* = 6) female mice at 13 weeks of age. **(E-G)** Food intake, physical activity, and energy expenditure (EE) in male

mice at 18 weeks of age across the circadian cycle (light and dark) and metabolic states (ad libitum fed, fasted,

refed) (WT, *n* = 11-12; KO, *n* = 10-12). **(H-J)** Food intake, physical activity, and energy expenditure in female

mice at 13 weeks of age (WT, *n* = 9; KO, *n* = 6). All data are presented as mean ± S.E.M.

1051

Figure 3. *Ctrp10*-KO mice fed a low-fat diet have normal fasting-refeeding response and glucose homeostasis.

(A-B) Overnight fasted and refed blood glucose, serum insulin, triglyceride, cholesterol, non-esterified free fatty

acids (NEFA), and β-hydroxybutyrate levels in male **(A)** and female **(B)** mice. **(C-D)** Blood glucose levels during

glucose tolerance tests (GTT; **C**) and insulin tolerance tests (ITT; **D**) in WT (*n* = 17) and KO (*n* = 14) male mice at

12 weeks of age. **(E-F)** Blood glucose levels during glucose tolerance tests (GTT; **E**) and insulin tolerance tests

(ITT; **F**) in WT (*n* = 9) and KO (*n* = 6) female mice at 20 and 21 weeks of age, respectively. All data are presented

as mean ± S.E.M. * *P* < 0.05 (two-way ANOVA with Sidak's post hoc tests).

1059

Figure 4. *Ctrp10*-KO female mice on a low-fat diet develop obesity with age. (A) Body weights over time of

WT and KO female mice fed a low-fat diet (LFD). **(B)** Representative image of WT and KO female on LFD for 40

weeks. **(C)** Body composition analysis of WT (*n* = 9) and KO (*n* = 6) female mice fed a LFD. **(D)** Representative

H&E stained histology of gonadal white adipose tissue (gWAT) and the quantification of adipocyte cell size (*n* = 6

per genotype). Scale bar = 100 μM. **(E)** Representative H&E stained histology of inguinal white adipose tissue

(iWAT) and the quantification of adipocyte cell size (*n* = 6 per genotype). Scale bar = 100 μM. **(F)** 24-hr food

intake data measured manually. **(G)** Fecal frequency, fecal weight, and fecal energy over a 24 hr period. **(H)** Deep

colon temperature measured at the light and dark cycle. **(I-K)** Food intake, physical activity, and energy

expenditure in female mice across the circadian cycle (light and dark) and metabolic states (ad libitum fed, fasted,

refed) (WT, *n* = 9; KO, *n* = 6). Indirect calorimetry analysis was performed after female mice were on LFD for 30

weeks. **(L)** Overnight (16-hr) fasted blood glucose, serum insulin, triglyceride, cholesterol, non-esterified free fatty

acids, and β-hydroxybutyrate levels. **(M)** Very-low density lipoprotein-triglyceride (VLDL-TG) and high-density

lipoprotein-cholesterol (HDL-cholesterol) analysis by FPLC of pooled (*n* = 6-7 per genotype) mouse sera. **(N)**

Blood glucose levels during glucose tolerance tests (GTT). **(O)** Blood glucose levels during insulin tolerance tests

(ITT). GTT and ITT were performed when the female mice reached 28 and 29 weeks of age, respectively. WT, *n* =

9; KO, *n* = 6.

1076

Figure 5. Sexually dimorphic response of *Ctrp10*-KO mice to an obesogenic diet. (A) Body weights over time of WT and KO male mice fed a high-fat diet (HFD). **(B)** Body composition analysis of WT (n = 17) and KO (n = 14) male mice fed a HFD for 9 weeks. **(C-E)** Food intake, physical activity, and energy expenditure in male mice across the circadian cycle (light and dark) and metabolic states (*ad libitum* fed, fasted, refed) (WT, n = 11; KO, n = 11). Indirect calorimetry analysis was performed after male mice were on HFD for 10 weeks. **(F)** Body weights over time of WT and KO female mice fed a high-fat diet. **(G)** Representative image of WT and KO female mice after 13 weeks of high-fat feeding. **(H)** Body composition analysis of WT (n = 17) and KO (n = 13) female mice on HFD for 6 weeks. **(I-K)** Food intake, physical activity, and energy expenditure in female mice (WT, n = 11-12; KO, n = 12) across the circadian cycle (light and dark) and metabolic states (*ad libitum fed*, fasted, refed). Indirect calorimetry analysis was performed after female mice were on HFD for 6 weeks. **(L)** ANCOVA analysis of energy expenditure using body weight as a covariate. **(M)** Respiratory exchange ratio (RER). All data are presented as mean ± S.E.M. * P < 0.05; ** P < 0.01; *** P < 0.001; **** P < 0.0001

Figure 6. *Ctrp10*-KO mice on a high-fat diet have normal glucose and insulin tolerance. (A-B) Overnight fasted and refed blood glucose, serum insulin, triglyceride, cholesterol, non-esterified free fatty acids (NEFA), and β-hydroxybutyrate levels in male **(A)** and female **(B)** mice fed a HFD for 10 weeks. **(C-D)** Blood glucose levels during glucose tolerance tests (GTT; **C**) and insulin tolerance tests (ITT; **D**) in WT (n = 17) and KO (n = 14) male mice fed a HFD for 10 weeks. **(E-F)** Blood glucose levels during glucose tolerance tests (GTT; **E**) and insulin tolerance tests (ITT; **F**) in WT (n = 16) and KO (n = 12) female mice fed a HFD for 8 weeks. **(G)** VLDL-TG and HDL-cholesterol analysis by FPLC of pooled female mouse sera. All data are presented as mean ± S.E.M. ** P < 0.01; *** P < 0.001; **** P < 0.0001 (two-way ANOVA with Sidak's post hoc tests for fasted/refed data).

Figure 7. *Ctrp10*-KO female mice fed a HFD do not develop adipose tissue dysfunction and fatty liver. (A) Representative images of dissected gonadal white adipose tissue (gWAT) and the quantification of gWAT weight in WT (n = 15) and KO (n = 12) female mice fed a HFD for 14 weeks. **(B)** Representative H&E stained histological sections of gWAT and the quantification of adipocyte cell size (n = 7 per genotype). Scale bar = 100 μM. **(C)** Representative images of dissected inguinal white adipose tissue (iWAT) and the quantification of iWAT weight in WT (n = 15) and KO (n = 12) female mice. **(D)** Representative H&E stained histological sections of iWAT and the

quantification of adipocyte cell size (*n* = 7 per genotype). Scale bar = 100 μM. **(E)** Expression of genes associated

with inflammation, fibrosis, ER and oxidative stress in gWAT and iWAT of WT (*n* = 6) and KO (*n* = 6) female

mice fed a HFD for 14 weeks. Gene expression data were obtained from RNA-seq. **(F-G)** Quantification of

hydroxyproline (marker of fibrosis) and malondialdehyde (MDA; marker of oxidative stress) in gWAT and iWAT.

WT, *n* = 15; KO, *n* = 11. **(H)** Representative images of dissected liver and the quantification of liver weight in WT

(*n* = 15) and KO (*n* = 12) female mice. **(I)** Representative H&E stained histological sections of liver and the

quantification of hepatic lipid content (% lipid area; *n* = 7 per genotype). Scale bar = 100 μM. **(J)** Hepatic

expression of genes associated with inflammation, fibrosis, ER and oxidative stress, lipid synthesis, and lipid

catabolism in WT and KO female mice. Gene expression data were obtained from RNA-seq. **(K-L)** Quantification

of hydroxyproline (marker of fibrosis) and malondialdehyde (MDA; marker of oxidative stress) in liver. WT, *n* =

15; KO, *n* = 11. All data are presented as mean ± S.E.M. * *P* < 0.05; ** *P* < 0.01.

1116

Figure 8. Transcriptomic analysis of liver, adipose tissue, and skeletal muscle of female *Ctrp10* KO mice fed a

high-fat diet. (A-D) Cropped volcano plot views of all differentially expressed genes (DEGs, Log2(Fold Change)

>1 or <-1 with a *p*-value <0.05) of the liver, gonadal white adipose tissue (gWAT), inguinal WAT (iWAT), or

skeletal muscle (gastrocnemius). **(E)** Overlap analysis of tissue DEGs showing (top panel) expression unique to

gonadal white adipose tissue (gW), inguinal white adipose tissue (iW), liver (L), or skeletal muscle (M). Percent

(%) represents percent DEGs unique to each tissue. Bottom panel show DEGs shared across multiple tissues, with

all the shared DEGs listed. **(F)** Enrichr analysis (129) of biological pathways and processes significantly (p<0.01)

affected across the CTRP10 deficient female mice. Top pathways and processes derived from Gene Ontology (GO),

Reactome (R-HAS), WikiPathway human (WP), and mammalian phenotype (MP). All up- or down-regulated

DEGs across all tissues were used for analysis. The tissues contributing to the highest ranked pathways and

processes are specified. *n* = 6 KO and 6 WT for RNA-seq experiments.

1128

1129 **Figure 9. Loss of CTRP10 induces significant and wide-spread alterations in the expression of key**

1130 **transcription factors, secreted protein, membrane receptors, and metabolism-associated genes. (A-D)**

1131 Selected genes from the DEG list of each tissue organized based on gene type (genes encoding transcription factors,

secreted proteins, receptors, and proteins involved in metabolism) and ranked from highest to lowest row z-score. n = 6 per genotype

Figure 10. GTEx genetic co-correlation of mouse differentially expressed gene (DEG) orthologues. (A-B) Heatmaps showing biweight midcorrelation (bicor) coefficient among human tissue DEG orthologues in females **(A)** and males **(B)** in GTEx. Y-axis color indicates tissue of origin, *P*-value based on students' regression *P*-value. **(C)** T-tests between correlation coefficient in males and females among all DEG orthologue gene pairs for subcutaneous (SubQ) adipose tissue, visceral (visc) adipose tissue, liver, and skeletal muscle. **(D)** the same as in C, except comparisons are shown for all gene-gene pairs between tissues. For example, the top left graph compared the connectivity of males (blue color) vs females (green color) for correlation between subcutaneous (SubQ) and visceral (Visc) adipose tissue DEG orthologues.

Human DEG Correlation (GTEx)

Females

A

B

Males

Bicor

1.0
0.5
0.0
-0.5
-1.0

* = p<0.01

Tissue
- Adipo (subQ)
- Adipo (visc)
- Liver
- Sk Muscle

Within tissue gene-gene connectivity

C

Adipose (subQ)	Adipose (visc)	Liver	Skeletal Muscle
$p < 2.2e{-}16$	$p = 3e{-}04$	$p = 3.8e{-}14$	$p = 1.8e{-}08$

abs(bicor)

Females Males   Females Males   Females Males   Females Males

Across tissue gene-gene connectivity

D

Adip. subQ - Adip. visc	Adip. subQ - Liver	Adip. subQ - Sk Muscle
Wilcoxon, p = 1e-15	p < 2.2e-16	p < 2.2e-16

Adip. visc - Adip. subQ	Adip. visc - Liver	Adip. visc - Sk Muscle
Wilcoxon, p = 1e-15	p < 2.2e-16	p = 6e-06

abs(bicor)

Females Males   Females Males   Females Males

Liver - Adip. subQ	Liver - Adip. visc	Liver - Sk Muscle
Wilcoxon, p < 2.2e-16	p < 2.2e-16	p < 2.2e-16

Sk Muscle - Adip. subQ	Sk Muscle - Adip. visc	Sk Muscle - Liver
Wilcoxon, p < 2.2e-16	p = 6e-06	p < 2.2e-16

abs(bicor)

Females Males   Females Males   Females Males