1 **Astrocytes are active: An information theoretic approach reveals differences in Ca2+ signaling**
2 **patterns among distinct astrocyte subtypes**

4 **Authors:**

5 Nicholas J. Mennona[1,2], Barbara Barile[3], Hoony Kang[1,2], Valentina Benfenati[4], Grazia P. Nicchia[3],
6 Kate M. O'Neill [1], Wolfgang Losert [1,2]

7
1   Institute for Physical Sciences & Technology, Univ. of Maryland, Col. Park, College Park, MD,
USA
2   Physics Dept., Univ. of Maryland, Col. Park, College Park, MD, USA
3   Biosciences, Biotechnology and Environment, University of Bari Aldo Moro, 70125, Bari, Italy
4   Institute of Organic Synthesis and Photoreactivity, National Research Council of Italy, 40129,
Bologna, Italy
14

Keywords: Collective Dynamics, Information Theory, Astrocytes

Abstract:

The discovery that astrocytes are an active, rather than a passive, component of the brain has ushered in a
paradigm shift in thinking about how the brain processes information. Although the mechanisms by
which astrocytes integrate information from neurons are still debated, such discourse should not distract
from the importance of more completely understanding how astrocytes communicate via signals amongst
themselves. This work aims to study how different astrocytes signal within their own networks. We
investigate group calcium (Ca^{2+}) dynamics in polygonal, stellate, and reactive astrocytes. These distinct
and important astrocyte subtypes are present in the brain to varying degrees at different physiological
states. We use an information-theoretic framework to quantify the dynamics embedded in the Ca2+ traces
within astrocyte networks; specifically, we employ the Hurst exponent, cross-correlation, mutual
information, and partitioned entropy to assess differences in the astrocyte signals across subtypes. To gain
insights into the ability of astrocyte networks to respond to changes in the extracellular environment, we
probe the networks with perturbations affecting their cytoskeletal dynamics (Latrunculin B) and energetic
levels (Adenosine triphosphate). Overall, these three classes of astrocytes behave differently and respond
idiosyncratically to their extracellular environment. We find that polygonal astrocytes are not quiescent,
stellate astrocytes respond most strongly to ATP, and reactive astrocytes are uniquely perturbed by
Latrunculin B. Interestingly, despite these distinct differences in behaviors, we find a uniform speed of
information transport regardless of subtype or perturbation; this uniformity is maintained when using both
cross-correlation and mutual information to assess this speed. We conclude that the differential ways
astrocytes signal within our measured framework yield important insights into how astrocytes
communicate and contribute to this pressing issue of understanding astrocyte information processing.

Introduction:

Astrocytes are the most numerous glial cell type in the brain. They are important for physiological
functions, including K^+ clearance[1], glutamate homeostasis[2], and extracellular volume regulation[3]
(reviewed in[4–6]). Since astrocytes carry out these known supporting functions and lack electrical

excitability, most studies have overlooked the potential active role of astrocytes in information
processing. In contrast, neuronal dynamics has become synonymous with information processing[7–12].

Nevertheless, recent work has provided strong indications that astrocytes do play an active role in
information processing. As a counterpart to neurons, astrocytes also form their own networks, linked
together via gap junctions composed of connexins, including Cx43, one of the main connexins in
astrocytes[13,14]. Within these networks, astrocytes signal to each other also via slow Ca^{2+} waves.[15].
Intracellularly, IP_3 triggers intercellular Ca^{2+} waves[15] which propagate through astrocyte networks via the
aforementioned gap junctions. Recent studies reveal that these waves can occur spontaneously, with
excitable dynamics similar to action potentials, although on a slower timescale of seconds[16]. Thus,
astrocytes can have an active role in information flow, with calcium waves functioning as non-electrical
communication signals. In single astrocytes, these calcium waves are heterogeneous[17] and flow through
astrocyte processes and microdomains[18–20]. Methods such as the machine learning tool AQUA now enable
robust intracellular analysis of astrocyte calcium[17] for *ex-vivo* and *in vivo* analysis of astrocyte activity.

Calcium signals, occurring either locally in microdomains or spreading into calcium waves also determine
the release of bioactive molecules called gliotransmitters from astrocytes. Astrocyte gliotransmission (by
e.g. ATP, Glutamate and GABA) is the known information transfer mechanism from astrocytes to
neurons, and has been modeled for its role in synaptic plasticity, memory, and learning[21,26–28]. We propose
that, to encompass such network scale information, astrocytes should also process and integrate
information among themselves.

To support our hypothesis and gain insight on the information flowing between astrocytes and the
unexplored network scale implications of that[16], in this work, we utilize physics-based information-
theoretic techniques to analyze and describe the behavior of calcium dynamics or pure primary rat culture
of astrocytes *in vitro*. The current study contrasts three physiologically relevant subtypes of astrocytes,
focusing on their collective (ensemble) dynamics within *in vitro* networks. These *in vitro* cultures remain
a powerful tool[16] for analyzing astrocyte dynamics, especially since the current study focuses on how
individual astrocytes contribute to the ensemble.

Following established protocols, we study *in vitro* models of polygonal (immature), stellate (healthy), and
reactive (immature) astrocytes. Intracellular and extracellular ionic changes are reported to occur upon
several physiological and pathophysiological events, including regulation of the activity of chemical
synapses, glial scarring[21], synapsis plasticity[22], and formation upon neurodevelopment[23]. Anisosmotic
challenges are counterbalanced by water flows, which are predominantly mediated by the water channel
aquaporin-4 (AQP4) in brain astrocytes[24]. AQP4 is reported to be polarized and enriched in astrocyte
endfeet where it plays a key homeostatic role. By modulating water influx/efflux, it ensures a fine regulation
of astrocyte cell volume and morphology, thus preserving brain integrity and functionality[24]. In this view,
it is not surprising that AQP4 is found to be co-expressed or functionally coupled in astrocyte membranes
with ion channels, including potassium (i.e. Kir4.1[25]) and calcium (i.e. TRPV4) channels[3,24,26]. On the other
hand, connexins are shown to take part in astrocytes' response to injury upon the stimulation of pro-
inflammatory cytokines, including TGF-beta[27]. An increase in the levels of Cx43 in reactive astrocytes is
proposed to act as a supporting mechanism that helps maintain astrocytes' gliotic phenotype and maximize
the coordinated intervention of these cells and their communication for the injury resolution[28,29].

The astrocyte networks – polygonal, stellate, and reactive – are broadly classified as immature, healthy,
and injured, respectively, and have distinct physiological behavioral profiles[19,30–33]. A key readout of these
distinct subtypes is their unique and stark morphologies. The differing structures and morphologies of
astrocytes, in general, have been implicated in brain functionality[30,34,35] and the cell-cell signaling of

astrocyte networks[36]. Notably, previous works evidence that different morphologies are paralleled by
different functions in astrocytes, such as alteration in the cell volume mechanism[37–39],
electrophysiological properties linked with potassium spatial buffering[39,40], and actin dynamics[41].
However, studies are lacking the capability of correlating spatially and temporally the calcium signaling
in astrocytes, which are critically implicated in all the above-mentioned astrocytes functions.

With the goal of broadening our understanding of astrocyte-specific calcium signaling, we quantify the
collective calcium dynamics, both spontaneous and perturbed, of these astrocytes. We perturb these
dynamics *in vitro* cultures with low-dose Latrunculin B (LATB) and Adenosine triphosphate (ATP) as
these represent basic perturbations to the cellular cytoskeleton[42–44], and cellular energy[45–48], respectively.
Due to the slow and heterogeneous characteristics of astrocyte calcium signaling, astrocyte signals are
more analog than digital. Neuronal signals are often treated as digital, 1 (firing) or 0 (not firing).
Astrocyte signaling is too slow for such binarization. Thus, information-theoretic methods that extract
information without digitizing the signal are used. We employ amplitude ordering symbolization with an
embedding window large enough to encapsulate the dynamics of astrocyte calcium events, which are
mostly on the order of 1-5.5 seconds[16]. Amplitude ordering, rather than binarizing via the mean, is more
appropriate for quantifying the dynamic properties of these nonlinear signals. This nonlinearity allows us
to elucidate information characteristics and information speeds across polygonal, stellate, and reactive
astrocytes in a comprehensive manner.

Methods

Imaging

Primary astrocytes were obtained from Sprague Dawley rats housed at the University of Maryland (in
concordance with the recommendations of and approval by the University of Maryland Institutional
Animal Care and Use Committee; protocols R-FEB-21-04). Experiments were performed between 14 to
21 (DIV14-DIV21) days after dissection[49]. At this time, astrocytes were trypsinized from flasks and
plated into PDL-coated (Sigma-Aldrich) 35 mm dishes (MatTek). Dishes containing cultured astrocytes
were incubated with 5µM CalBryte 590 AM (AAT Bioquest) for 30-60 min at 37°C prior to imaging.
Dishes were then washed with PBS. Astrocytes are plated at densities such that they proliferate to roughly
500k cells/well on the day of imaging (a final density of approximately 10k cells/mm^2), and imaging was
performed on a spinning disk confocal microscope (PerkinElmer). Image acquisition occurred for
approximately 15 minutes at roughly one frame per second with 100 ms of exposure time per frame. The
University of Maryland Imaging Core maintains the PerkinElmer spinning disk confocal microscope used
for this research. The image sequences were taken by the equipped Hamamatsu ImagEM X2 EM-CCD
camera (C9100-23B), which records 16-bit images. Sequences were taken from an air 20x objective (0.75
NA; 0.717 µm per pixel) and under temperature (37°C), CO2 (5%), and humidity control.

All experiments are performed in NeuroBasal media (Gibco). We use this media to simulate a neuronal
environment, and we found that it allows for spontaneous activity in unperturbed wells (over DMEM).
For simplicity, throughout this manuscript, this environment is referred to as spontaneous (control). ATP
(Sigma-Aldrich) and Latrunculin B (Sigma-Aldrich) were applied immediately upon imaging and at low-
dose concentrations of 1.5uM for ATP and 0.5uM for Latrunculin B. Unlike Latrunculin A, the slower
depolymerization rate of Latrunculin B allows for perturbation of actin without drastically dissociating
the entire astrocyte network. The value of Latrunculin was found based on testing whether our astrocyte
networks were maintained (expressed minimal disassociation) for the 15-minute duration at the chosen

imaging setting (e.g., laser intensity). For simplicity, throughout the rest of this work, we refer to
Latrunculin B as LATB.

Cell Culture

Primary astrocytes were obtained from Sprague Dawley rats housed at the University of Maryland (in
concordance with the recommendations of and approval by the University of Maryland Institutional
Animal Care and Use Committee; protocols R-FEB-21-04). Astrocytes were prepared from 0-to-2 day
postnatal (P0-P2) Sprague Dawley rat brains as described previously[41]. Briefly, cortical tissue from each
pup was separately dissociated via trituration, filtered, and plated into T25 flasks containing DMEM
(ThermoFisher), 15% fetal bovine serum (Benchmark), and 1% penicillin-streptomycin (P/S). After
roughly seven days, in order to generate stellate and reactive astrocytes, the media was changed to either
(1) stellate media, which includes Neurobasal (ThermoFisher), 2% B27+ (GIBCO), 5 ng/mL HB-EGF
(TOCRIS BioSciences)[19] or (2) reactive media, which is serum-free DMEM to which 10 ug/mL of TGF-
B1 was added after five additional days *in vitro*[50].

Cell lysis and Western Blot

Protein samples were obtained as previously described, with some slight changes[3]. Briefly, astrocytes
grown for seven days in control, stellate, and reactive media were washed in ice-cold phosphate-buffered
saline (PBS) and scraped with 300 µL of RIPA lysis buffer (10 mM Tris-HCl, pH 7.4, 140 mM NaCl, 1%
Triton X-100, 1% Na deoxycholate, 0.1% SDS, 1mM Na_3VO_4, 1 mM NaF, 1 mM EDTA and 1× Protease
Inhibitor Cocktail). Lysates were vortexed every 5 minutes for 30 minutes and centrifuged at 22 000 for 30
minutes at 4°C. Supernatants were collected and Pierce™ BCA Protein Assay Kit was used to dose the
protein content. 5 µg of proteins/lane were dissolved in Laemmli Sample Buffer (Bio-Rad, Hercules,
California, USA) added with 50 mM dithiothreitol, heated to 37°C for 10 min and resolved by SDS-PAGE
in 10% (for Cx43 detection) and 12% (for AQP4 and GFAP detection) gels prepared with Mini-PROTEAN
TGX Stain-Free Precast polyacrylamide solutions (Bio-Rad, Hercules, California, USA). gels were then
activated for 5 minutes on a Chemidoc imaging system (Bio-Rad, Hercules, California, USA) and
transferred to polyvinylidene fluoride (PVDF) membranes (Merck Millipore, Burlington, Massachusetts,
USA). Stain-free signal of transferred blots was collected. Membranes were incubated with blocking
solution and incubated overnight with primary antibodies, rinsed, then incubated with the appropriate
peroxidase-conjugated IgG secondary antibodies. Protein-specific bands were revealed using enhanced
chemiluminescent Clarity Western ECL Substrate (Bio-Rad, Hercules, California, USA) and visualized on
a Chemidoc imaging system. The densitometric analysis was performed in Image Lab software 6.1.0 by
normalizing the bands to the stain-free blot signal. The Western blot analysis of Cx43, AQP4, and GFAP
for the polygonal, stellate, and reactive astrocytes confirms three distinct phenotypes: reactive upregulate
Cx43 and AQP4, stellate upregulate AQP4, and polygonal cells do not express any of these markers
significantly.

Fig 1: Astrocyte morphologies (polygonal, stellate, and reactive). The phenotypes correspond to different morphologies, as
shown in green via actin images. Images of actin are acquired with 488nm light of astrocytes that express BacMam Actin-GFP.
We record the calcium signaling channels of these dense astrocyte networks with 561 nm light of astrocyte networks stained with
CalBryte 590. It has been noted that, regardless of protocol, different media formulations result in different morphologies for
astrocytes[19]. We use time overlays to show the different manifestations of calcium propagation in the distinct morphological
conditions of astrocytes. Blue corresponds to an increase in [Ca2$^+$] during this frame, yellow a decrease, and white means that
this frame contains values similar to the previous timepoint. Western Blot analysis of Cx43, AQP4, and GFAP in polygonal,
stellate, and reactive astrocytes. The upper panels show Cx43 (band at ~43kDa), AQP4 (band at ~30-32 kDa), and GFAP
expression (band at ~50 kDa) in primary rat astrocytes. Western Blot analysis of Cx43, AQP4, and GFAP in polygonal, reactive,
and stellate astrocytes. Western Blot showing Cx43 (band at ~43 kDa), AQP4 (band at ~30-32 kDa), and GFAP expression
(band at ~50 kDa) in rat astrocytes. Using Kruskal Wallis Dunn's multiple comparison test, we find statistical significance for
polygonal compared to stellate (AQP4) and reactive (Cx43 and AQP4). *p< 0.05; **p<0.01. Scale bars are 25 μm.

In this work, we investigate astrocytes of three physiological types; a prominent distinguishing
characteristic of the differences is seen in the differences in morphology. Fig. 1 (top) provides representative
images of the three classes of astrocytes analyzed in this study. Each astrocyte network does not contain a
mix of different cell types and is cultured to maintain a singular physiological expression. For these
astrocyte types, we investigate collective Ca2+ dynamics as illustrated in Fig 1 (middle), where multiple
frames are overlaid to demonstrate the distribution of Ca^{2+} fluctuations (rising calcium in blue, diminishing
calcium in yellow) in a network. Through Western blot analysis, we evaluated the expression profiles of 1)
Cx43, one of the major isoforms forming astrocytic gap-junctions involved in calcium-mediated cell-to-
cell communication, as a molecular marker of intercellular connectivity within astrocytes networks; 2)
AQP4, the most abundantly expressed water channel in brain astrocytes, and GFAP, the predominant
intermediate filament of the astrocyte cytoskeleton in the CNS, as proxy markers of cell differentiation and
reactivity[3,51,52].

AQP4 and its supramolecular level of organization in larger-sized assemblies have been reported to
correlate with the morphological and functional maturity of astrocytes in CNS postnatal development[53] as
well as in in-vitro models of astrocyte differentiation[3]. Therefore, the overexpression of AQP4 in stellate
astrocytes is likely to refer to the morphological differentiation of this cell type in the experimental
conditions. The overexpression of Cx43 in the reactive conditions might be ascribable to the acquisition in
vitro of the in vivo-like reactive state that makes astrocytes more prompt at tissue healing and wound repair.

As for GFAP, besides the number of studies that identified a positive role for these canonical hallmarks of
astrogliosis, just as many provided opposite evidence. On one hand, it was previously found to be
downregulated in similar or even different models of astrogliosis (i.e., TNF and IL-1β[54–56] and LPS-treated
astrocytes[57]. On the other hand, its upregulation was associated with beneficial and neuroprotective effects
i.e., enhancing neuronal regeneration through protein re-localization, increasing the membrane retention of
glutamate transport GLAST in astrocytes, which protects surrounding neurons from glutamate
excitotoxicity[58]; providing scaffolds for new neurites[59]. In the present study, no changes were detected in
the three conditions, although we could detect differences in other markers in response to inflammatory
(i.e., Cx43) and differentiation stimuli (i.e., AQP4; Cx43). We, therefore, anticipate that GFAP might not
be a reliable marker to assess astrocytes' degree of astrogliosis or differentiation.

Western blot analysis shows that the three cell types have distinct expression profiles of the aforementioned
astrocyte proteins. Of note, we found that Cx43 is upregulated in the reactive state compared to control and
stellate astrocytes (Fig.1, bottom left panel), while AQP4 is overexpressed both upon stimulation with TGF-
beta and in star-shaped astrocytes, differently from controls (Fig.2, bottom middle panel). GFAP was
equally abundant in the three conditions with no statistical significance (Fig.1, bottom right panel).

Overall, while illustrating that the proteins undergo distinct changes in their profile expression in response
to the different external stimuli provided under culture conditions, these results also provide evidence that
the three cell types can be regarded as distinct "phenotypes" from the molecular perspective.

Analytical Methods

To analyze the calcium dynamics of astrocytes, we first segment individual astrocyte objects within our
imaging datasets using the morphological watershed algorithm FogBank[60]. Manually, we refined
parameters per each imaging sequence such that segmented boundaries roughly correspond to the
distinctions made by eye. FogBank preserves the unique heterogeneity of astrocyte bodies rather than
using the cell center with an arbitrary radius, as is typically done with neuron-imaging analysis[61]. See
Supplement for a representative FogBank example. All image sequences are jitter corrected, and Gaussian
smoothed in the spatial domain before further analysis. For each frame in a dataset, we take the median
fluorescence value from all the pixels identified within the astrocyte object, as shown in the five objects in

Fig 2A. We compute relative fluorescence measure ($\Delta F/F_0$), where $\Delta F = F - F_0$, for each astrocyte object. F_0
is computed using a window of approximately 30s (30 frames, 15 back, 15 forward) where F_0 is the
average of the fluorescence values below the 50th percentile of values contained within the window[62].
This is called a trace. Throughout the paper, for convenience, we address this fluorescence measure as
DF/F. As seen in representative traces, peaks in calcium dynamics are broad and not well-defined. The
sliding window is chosen based on an inspection of the data and does not meaningfully change the
dynamic properties of the traces. Please consult the Supplement for more details. We chose this window
size based on experimental observation of astrocyte calcium event duration. Examples of such traces are
shown in Fig 2B. Recall that astrocytes are not neurons, so it is inappropriate to call the activity of our
traces 'spikes.' Instead, we use the term "calcium event" to describe the smooth rises and falls exhibited
by traces in Fig 2B.

Fig 2: Astrocytes traces extracted and filtered using Hurst exponent. (A) Given a dense astrocytic network, astrocytic elements
are segmented using FogBank. Examples of astrocytic elements extracted. (B) DF/F traces of the five objects extracted in blue
(dark pixelated objects) are shown. (C) Hurst exponent measures memory in a time series, which is used to filter out inactive
cells or poorly segmented objects. Inactive (salmon, light) versus active (red, dark) cells plotted. Only time series with Hurst
exponents greater than or equal to 0.5 are used. (D) From the example points, we plot representative inactive and active traces.

Our goal is to use traces that show persistent calcium signals and to distinguish them from random
fluctuations of signals in the networks. Our goal is to define traces that exhibit smooth and meaningful

rises and falls (see red, darker curves in Fig 2D) as 'active,' regardless of the intensity values. The Hurst
Exponent (defined as *H*) allows us to carry out this analysis. H is defined for a given time series x(t) as:

249
$$< |x(t + \tau) - x(t)|^2 > \sim \tau^{2H}$$

Equation 1

We find that the Hurst exponent provides a robust and reliable way of filtering out noisy traces due to
poor segmentation or weak calcium signals. Hurst exponent values below a threshold of 0.5 represent
random activity, and the corresponding traces (e.g., the salmon, lighter colored curves in Fig 2D) are
excluded from the analysis[63,64].

The importance of correlations in neuronal encoding is well-cited in the literature[65–68].The study of
correlation in the brain relates to the collective activity of neuronal ensembles in relation to information
encoding and processing in the brain. This level of mathematical rigor has not been applied to astrocyte-
only networks, and the study of correlation within astrocyte networks has only been viewed in the context
of neuronal activation[69]. Given the success of using these methods on neurons and the absence of their use
in analyzing astrocytes, we argue that our use of these methods should yield comparable success and
insight. Correlations allow us to understand the global characteristics of an ensemble of active nodes (in
this case, astrocytes). However, we can alternatively view the dynamics in our astrocyte networks as a
dynamical system that is affected by individual astrocytes, i.e., the local dynamics of one trace affecting
the global dynamics observed. Since our Western blot analysis confirms cell-cell coupling, we argue that
these readouts provide a fairly accurate representation of the information processing performed by these
collections of astrocytes. This approach has been well-documented experimentally (e.g., with neuronal
MRI data) and theoretically[70–78] . A limitation of correlations is that the global characteristics of a network
are studied by assessing each cell's connections to others in the network. Information theory helps
overcome such limitations. An information theoretic approach can analyze a network for each astrocyte
rather than for pairs of cells. Thus, this approach provides a robust metric for assessing each individual
astrocyte's contribution to the mean field or the group characteristics of the network.

To implement this approach, from the extracted traces, we convert the traces into states using
symbolization[70,79]. This symbolization enables the interpretation of changes in calcium as an analog
signal; traces that exhibit lower entropy correspond to traces with more periodic fluctuations. We take the
time traces shown in Fig 3A, the amplitude order of each timepoint in a window, and convert this set of
timepoints into a state (an example suite of states is available in Fig 3B), and then convert the traces into
symbol sequences visualized in Fig 3C. As an illustrative example with a manageable number of states,
m=3 has been used (number of states = m!). From the resulting symbol sequence, we can find the
probability distributions of states (Fig 3D). The probability of states, in turn, allows us to compute an
information entropy for this representation of calcium traces. Amplitude ordering entails choosing two
parameters, m, and l, where m defines the embedding dimension[70,78] (the symbol set size that functions as
a coarse-graining window) and l denotes the time lag[78]. A more comprehensive review of this
symbolization method can be found in Refs[70–72,74,76–79]. We choose m to smooth out noise without losing
actual spikes in the data. Based on our experimental observation of the duration of calcium events in our
data, we chose m=5 and l=1. As m takes the role of the coarse-graining window, l can be kept at its
minimum (l=1) to prevent further information down-sampling.

Given that we have converted these time series into symbol series, we can perform time-delayed mutual
information methods. Time-delayed mutual information is the symbolized equivalent of cross-correlation.
Moreover, given that astrocyte signals are nonlinear, mutual information is a more appropriate method
than Granger causality[80]. While there are some similarities between mutual information and cross-
correlation, we emphasize that mutual information discusses the dynamics of symbols (with embedded

dynamics) from a probability distribution. Cross-correlation looks at the absolute fluctuations from the
mean for traces. In some regards, then, symbolization can be viewed as the differential expression of trace
dynamics. Lastly, from both of these measures, we can quantify the speed of information transport[81]. We
can measure speed from the measured distances between astrocytes and either the cross-correlation time
lag or the time lag associated with the mutual information measurement.

Additionally, we analyze the information content of astrocyte networks with partitioned entropy.
Partitioned entropy $h_a(\tau)$ is a useful metric for understanding the time-evolution of states over a certain

299
$$h_a(\tau) = -\sum_{i=1}^{m!} p_{ai}(\tau)\ln p_{ai}(\tau)$$

Equation 2

where $p_{ai}(\tau)$ represents the probability that the system evolves from state a to state i at τ
timesteps. Partitioned entropy differs from canonical entropy as it is a function of time. We employ this
approach to understand better how the ramps and falls of these calcium fluctuations evolve differentially
for all of the cells within the unique networks. A schematic for understanding partitioned entropy is
shown in Fig 3E. We validated this analysis on both regular and chaotic time series data.[72] See
Supplement for details. As a clarification, we note that the symbolization described enables analysis of

Shannon entropy. The rest of the analysis refers to Shannon entropy and the dynamics therein, e.g.,
partitioned (Shannon) entropy.

Fig. 3: Methods for Information Theoretic Approach to Collective Astrocyte Dynamics. (A) Astrocyte traces. Different shaded
boxes correspond to example symbol states. (B) Symbol states. Shaded boxes are those previously overlaid on traces. In this
figure, have used an m of 3 as this corresponds to 6 possible states for every time window (m!=6). We note, using grey arrows,
the physical dynamics behind each state (for states 1 and 2, the same size for the arrows is for visual purposes). (C) Symbol
sequences for the traces shown previously. (D) Visualization of Time-Delayed Mutual Information. (E) Example subset of
transition probabilities from states to future states for some given tau for any particular state 'i'. See Equation 1 for more details.
This subfigure represents partitioned entropy.

<u>Results</u>

Fig 4: Speeds of Information Transport via Cross Correlation and Mutual Information. Figures shown for all distinct subtypes of
astrocyte. Cross correlation speeds are shown for (A) spontaneous, (B) ATP, and (C) LATB affected networks. Mutual
information speeds are displayed for (D) spontaneous, (E) ATP, and (F) LATB affected networks. Due to the skewed nature of all
distributions, these distributions have been log-normalized. We find the speed of information transport is invariant across
astrocyte subtype and extracellular environment.

Astrocyte Speed of Information Transport

An important metric for assessing how astrocyte transport information is information speed (speed of
information transport). This ad-hoc (average) metric is extracted for free by any time lag calculation. We
do not track calcium flow, but can track the exchange of information as a function of time and distance
between cells. This method has been used for mutual information[81]; we extend it for cross correlation
since the time lag associated with maximal cross correlation is similarly extracted. Our measure of speed
is average ($v = DIST/\tau$), it coarsely reveals the rate at which these pairwise quantities are absolutely
maximized, reflecting the nature of the timescales associated with astrocyte calcium events. Additionally,
the maximal time lags found are rate-limited by the acquisition speed, which is 1 Hz. Due to the 1 Hz
frame rate and other coarse graining involved (e.g. distance is computed from center of mass points in a
segmented image), this measure is an average metric. Thus, we conclude that a few micron per second
difference is not significant to ascertain any meaningful discrepancy from a biological perspective.

We describe the cross-correlation speeds first. The distributions are modelled as a lognormal distribution
to obtain the mean speeds to follow as shown in Figs 4A-C. Outliers at infinity are removed (which may
happen for time lags of zero). We report the mean of these distributions modeled under lognormal

assumptions. The spontaneous speeds are: for polygonal 25 µm/s, for stellate 28 µm/s, for reactive 23
µm/s. Under LATB, for polygonal 25 µm/s, for stellate 27 µm/s, for reactive 21 µm/s. With ATP, for
polygonal, 29 µm/s, for stellate 28 µm/s, for reactive 25 µm/s. As expressed in Figs 4A-C, these
distributions are not symmetric (not entirely lognormal). Given that the order of difference between
stellate and other networks is ~5 µm/s at most, and the distributions themselves retain similar
characteristics (the distributions peak within this 20-30 µm/s range consistent with experimental studies),
we conclude that the speed of information transport is invariant across astrocyte subtypes and conditions.

Next, we report the mutual information speeds. The spontaneous speeds are: for polygonal 28 µm/s, for
stellate 32 µm/s, for reactive 30 µm/s. Under LATB, for polygonal 34 µm/s, for stellate 32 µm/s, for
reactive 29 µm/s. With ATP, for polygonal 34 µm/s, stellate 30 µm/s, and reactive 29 µm/s. These mean
values are reported from modelling the distributions as lognormal. We conclude that the apparent
differences between the reported mean values are minimized for mutual information speeds. The speeds
are slightly higher than those of cross correlation. All speeds are in the ~30-35 µm/s. This range is
roughly consistent with the cross-correlation speeds (at the higher end), and is consistent with the reported
literature.

Largely, we find a constant speed of information transport across conditions and types. Using this metric
(defined as distance between objects divided by the time lag associated with maximization of pairwise
quantity), we find that all distributions sharply peak similarly within the reported experimental range for
calcium wave speeds. The nature of these distributions across both subtype and conditions suggests an
invariance of the speed of information transport in astrocyte networks using this coarse approximation.
The similarities in the relationships between the distributions for the quantities (mutual information and
cross-correlation are similar, but differently formulated) reflects that the aggregate relationships between
calcium signals for astrocytes are independent of subtype and perturbation. This counter-intuitive result
suggests that astrocytes regulate ions' flow despite cellular state, and maintain homeostasis in stressful
environments to maintain signaling speeds.

As a final remark, we comment that the time-lags associated with either maximal cross-correlation or
mutual information may come from a symmetric peaking in a time-lag curve (i.e. bimodal peaks about the
y-axis). We argue that many of the time lags reflect a maximal value over a relevant physiological
window. All other lags reflect a mode within the time-lag curves themselves. This argument is expanded
in the Supplement.

Fig 5: Cross correlation versus mutual information, and partitioned entropy comparisons. We compare the distributions of both maximal cross correlation and maximal mutual information across time for (A) spontaneous, (B) LATB-induced, and (C) ATP-enhanced astrocyte networks. Each point represents the cross correlation (x-axis) and mutual information (y-axis) for the same pair of cells. We then look at the dynamics of individual traces using partitioned entropy for (D) spontaneous, (E) LATB-induced, and (F) ATP-enhanced networks. Partitioned entropy ratio is the average of the partitioned entropy divided by the entropy.

Information exchange of signaling amongst active astrocytes

Next, we assess how the traces in astrocyte networks can be used to infer relationships across pairs of cells. Given a time window of 31 frames (roughly 31 s), we find the maximum cross correlation and mutual information values. These relationships are shown in Figs 5A-C. First, we discuss the cross-correlation distributions. In spontaneous conditions, polygonal astrocyte pairs have higher cross correlation subtype, but stellate and reactive pairs demonstrate comparable distributions (Fig 5A). When LATB is introduced into the astrocyte networks; the distributions are separable indicating that the breakdown of the actin cytoskeleton causes idiosyncratic (i.e. each astrocyte network varies its distribution relative to its spontaneous profile) changes to astrocyte subtypes (Fig 5B). When ATP is introduced, all distributions curves aggregate (Fig 5C). The effect of ATP suggests a minimization between the global differences in astrocyte populations. This minimization is reflected in the strongly overlapping distributions.

Next, we discuss mutual information. In spontaneous conditions, we find that the distributions for the astrocyte polygonal and stellate astrocytes are similar (and lower than reactive). These results suggests that spontaneously polygonal and stellate astrocyte behave more independently than reactive astrocytes (Fig 5A). When LATB is introduced, all mutual information curves are shifted higher with reactive

astrocytes retaining larger values while polygonal and stellate astrocyte pairs are similar (Fig 5B).
Interestingly, this result indicates that a breakdown in the cytoskeleton causes uniformity in local
behavior in astrocyte interactions (i.e. higher values of mutual information reflect less independence).
When ATP is introduced, stellate astrocytes respond the most strongly (Fig 5C). Stellate astrocytes
become strongly independent in terms of their local behavior. While stellate astrocytes respond the most
strongly, all astrocyte subtypes become more independent (lower mutual information) when exposed to
ATP.

Partitioned entropy highlights temporally local dynamics

We look at the partition entropy ratios for astrocytes in networks. Unlike with cross correlation and
mutual information, partitioned entropy is a method for analyzing the dynamics of individual traces
themselves (i.e. it is not a pairwise analysis method). As a dynamical measure, it provides more ability for
understanding information dynamics than reporting entropy values alone. Partitioned entropy ratios are
calculated by finding the average partitioned entropy across some time window (in this paper we chose to
start at $\tau=10$ and end at $\tau=150$, see Supplement for details). Values of partitioned entropy closer to 1 are
classified as chaotic whereas values closer to 0.5 are deemed limit cycles (e.g. sinusoidal oscillations).
Classifying astrocytes in this way is beyond the scope of this work, but it is useful for further
distinguishing dynamical profiles of the subtypes.

We find that spontaneously, astrocytes retain similar individual characteristics, as demonstrated in Fig
5D. This result is surprising given the numerous reported physiological differences in these subtypes.
Coupled with some similarities across subtypes in the pairwise interactions results, we argue that, in
spontaneous conditions, when astrocytes signal it is similar. This result suggests a universal characteristic
to spontaneous astrocyte calcium events. LATB, comparable to the distinct effects in pairwise
interactions, causes reactive astrocyte to behave more sinusoidally (Fig 5E). Stellate astrocytes respond
with higher partitioned entropy ratios. Lastly, where LATB induces the strongest separation in cross-
correlation distributions, ATP seems to induce a comparable outsized effect in the partitioned entropy
curves shown in Fig 5F. The different perspective (focusing solely on the individual embedded dynamics)
provided by partitioned entropy complements the results mentioned above in ascertaining the nuanced
differences in astrocyte calcium signaling.

Fig 6: *Wasserstein distances for the three classes of distributions shown in Fig 5. Astrocytes types are grouped; three pairs exist as highlighted by the dark (polygonal-stellate), purple (polygonal-reactive), and pink (stellate-reactive). We contrast these groups against the natural variance within the individual subtype distributions. We find the Wasserstein distances for: spontaneous (A) cross correlation (B) mutual information, and (C) partitioned entropy distributions, LATB exposed (D) cross correlation (E) mutual information, and (F) partitioned entropy distributions, and ATP induced (G) cross correlation (H) mutual information, and (I) partitioned entropy distribution. Results are bootstrapped.*

Wasserstein distances enhance interpretability of comparisons

We use the Wasserstein distance (using its L1 norm definition, Wasserstein-1) to measure the differences in distributions shown in Fig 5. Wasserstein-1 quantifies the minimum distance needed to shift all points in a distribution to match another distribution. The Wasserstein distance considers the individual values used to generate the histograms in Fig 5., and is independent of the choice of bin. We can measure a

Wasserstein distance not only to compare distributions, but also to characterize a single distribution by
using bootstrapping. Bootstrapping was performed for all Wasserstein-1 calculations with a sample size
of 100 and 10,000 iterations. We can calculate Wasserstein distances for all metrics, cross correlation,
mutual information, and partitioned entropy with this approach.

Under spontaneous conditions, polygonal cross correlation values are the most distinct (Fig 6A). Using
symbolization, we find that the differences between stellate and reactive astrocytes (S-R) become most
apparent for mutual information (Fig 6B) and partitioned entropy (Fig 6C). Thus, we conclude that on a
global interaction perspective, polygonal dynamics behave the most uniquely; on a local level (using the
embedded symbol states to understand dynamics), we find that spontaneously stellate astrocytes have a
unique dynamical profile relative to the reactive subtype. The ability of information theory to tease out
this result is significant. Both stellate and reactive astrocytes possess branches[82]. The process formation
could have implicated such processes in the similarities seen in their cross-correlation distributions (S-R
of Fig 6A).

Relative to spontaneous conditions, LATB increases the differences in the distributions for cross-
correlation (Fig 6D), mutual information (Fig 6E), and partitioned entropy (Fig 6F). Where again, we see
that polygonal astrocytes on a global level are the most distinct relative to stellate and reactive astrocytes
(Fig 6D), but we find that in the mutual information distributions, P-R and S-R, reactive astrocytes
separate the most strongly. This suggests reactive astrocytes are more susceptible to cytoskeletal
perturbation, limiting pairwise signal propagation. Importantly, the high Wasserstein distances in Figs.
6D and 6E distinguish polygonal and reactive (P-R) astrocytes despite the similarities in AQP4
expression. The strongest separation in terms of individual trace dynamics is seen in S-R (Fig 6F),
suggesting a strong difference in effect when healthy (stellate) versus unhealthy (reactive) astrocytes are
exposed to LATB.

Lastly, it is interesting that the highest reported Wasserstein distances for mutual information (Fig 6H)
and partitioned entropy (Fig 6I) are found under exposure to ATP. The lowest differences are attributed to
the cross-correlation (Fig 6G). Outside of physiological comparisons, this highlights the importance of the
diverse set of information theoretic tools used for understanding astrocyte signal dynamics. In contrast to
the effect of LATB, ATP appears to cause the entire network to become excited (i.e. hyper-active),
consistent with its known role as a key gliotransmitter. The lowest Wasserstein differences (across all
conditions) are reported for P-R. Indeed, this result indicates that stellate astrocytes are the most
susceptible to ATP intervention (P-S and S-R in Figs. 6H and 6G) whereas polygonal astrocytes share the
unhealthy characteristics of reactive astrocytes (P-R).

<u>Discussion</u>

Collective astrocyte dynamics are rich in information distinct from neurons and previously overlooked.
Starting from filtering out inactive traces with the non-biased Hurst exponent, we employed cross
correlation, mutual information, and partitioned entropy to ascertain the dynamical properties of
collections of maturing (polygonal), healthy (stellate), and injured (reactive) astrocytes. This behavior
characterization extends to these astrocyte networks defined by distinct states (subtypes defined by
differences e.g., in molecular expression, morphology) and when gliotransmitters or biomechanics
(LATB) of the system is altered. Interestingly, this focus on information differences allowed us to
demonstrate uniformity in the speed of information transport across astrocyte subtype and extracellular
environment (ATP/LATB). Our analysis on these astrocyte networks handles the complex spatiotemporal
patterns present in these data. Not merely regarding the calcium signaling of astrocytes as an adaptive
mechanism, our work argues that the spatiotemporal patterns embedded in these networks are active. That

these patterns themselves are sources of dense information rivaling neuronal networks. The
spatiotemporal patterning of spontaneous astrocytes is shown in the Supplement.

A principal finding of this work is the corroboration of astrocyte signaling (Ca2+ waves) speed using
information theoretic methods. Using cross correlation and mutual information on astrocyte traces,
aggregating traces from independent image sequences, we find an invariance of the speed of information
transport. We discovered that for all conditions investigated, the speed of information transport is
approximately 25-40 µm/sec, consistent with the speed of Ca^{2+} waves measured directly in prior studies
[49,83,84]. These findings are independent of astrocyte subtype and chemical milieu, which strongly indicates
that Ca^{2+} waves are the primary mechanism of astrocyte-to-astrocyte information exchange, i.e. Ca^{2+}
dynamics is a dominant form of communication in astrocyte networks for all states and environmental
conditions investigated despite differences in molecular expression, information processing, and the
biochemical cell-cell coupling machinery (AQP4, Cx43). Given that we will subsequently discuss the
differences in information dynamics, this result is even more startlingly. Confirmation of the speed of
information transport with previously reported values lends confidence to using these methods for
understanding astrocyte signals.

We preface subsequent interpretations of our results with subtle distinctions between the competing
pairwise information theoretic concepts in this work. Correlation (of which cross is the most appropriate
for astrocyte signals; see Supplement for more details) deals with the specific similarities of mean field
fluctuations of the traces. One way to understand this subtle concept is that pairwise correlation values
relate to the absolute coarse-grained differences between points in a time series. Accounting for the time
lag to maximize correlation, in the context of astrocyte signals, it describes how similar the calcium
events look; the average area overlap of all calcium events. Mutual information, supported by
symbolization, is more focused and able to provide more local descriptions of how symbol state
influences across time series[85]Both are important, and our results, such as the marked separation between
cross-correlation LATB distributions, indicate that both provide a holistic ability to compare astrocyte
dynamics. Indeed, an externality of these results, including the use of partitioned entropy for comparative
analysis, is a demonstration of the varied insights that are gained by using this complete set of
information theoretic tools.

Importantly, polygonal astrocytes in control conditions (spontaneous), which have been regarded as
quiescent,[19] exhibit information processing; in other words, polygonal cells are not inactive cells and
have complex signaling patterns. The difference between these polygonal cells, and stellate and reactive
astrocytes at first lies in the global correlation structure amongst polygonal astrocytes. As shown in Figs
6A,6D,6G, polygonal cells are the most distinct with higher cross correlation values than stellate and
reactive astrocytes. As correlation is a global averaging of time values, this suggests that the mean field
behavior of polygonal cells is more coupled than those of stellate and reactive astrocytes. In other words,
polygonal astrocytes broadly have more global network synchrony. These insights are counterintuitive as
the higher expression of the astrocyte markers regulating cell-cell coupling, AQP4 and Cx43, seen in
stellate and reactive astrocytes (Fig 1) suggest that these cells would couple more closely in their
information processing. One possible explanation is that the lower expression of coupling proteins limits
the signal propagation to fewer polygonal neighbors and thus, paradoxically, these polygonal networks
are globally more aligned. Future studies will investigate this possibility.

Locally, we gain additional insights into stellate and reactive signaling dynamics using symbolization
which embeds the dynamics. We conclude that reactive astrocytes respond most strongly to cytoskeletal
perturbation (LATB) whereas stellate astrocytes react most strongly to gliotransmitter ATP. These results
are consistent with the role of the cytoskeleton and ATP in reactive and stellate astrocytes, respectively.

Reactive astrocytes must have a dynamic cytoskeleton as they migrate to form the glial scar, an adaptive function in response to injury or disease[35,86]. Stellate astrocytes communicate via more localized contacts, and ATP is suggested to be an important extracellular messenger for specifically stellate-like calcium waves[87,88].

Overall, our work shows that applying information theoretic tools to *in vitro* astrocyte networks lends insights into the collective action of astrocytes. Generally, our work has analyzed the complex spatiotemporal characteristics of polygonal, stellate, and reactive networks. We find that polygonal astrocytes are not quiescent, stellate astrocytes respond most significantly to ATP, and reactive astrocytes are most affected by LATB. Our study also provides a systematic baseline of three complementary astrocyte states and key perturbations supporting future studies of astrocyte network function in health and disease. The successful demonstration of our method across numerous astrocyte conditions and microenvironments suggests broad applicability and promise for these tools to study astrocyte dynamics.

Acknowledgments

The authors thank the University of Maryland Imaging Incubator Core Facility for providing and maintaining the systems used in collecting images for this work. This work was supported by US Air Force Office of Scientific Research Astro-COLL grant FA9550-21-1-0352.

We want to thank Dr Leonard Campanello for always being generous with his time and providing insightful guidance on the timescales of information. We additionally thank Prof Pratyush Tiwary for his comments and suggestions.

Software Availability

Software has been made available on GitHub at:
https://github.com/losertlab/Astrocyte_CollectiveNetworkInformation

1. Bellot-Saez, A., Kékesi, O., Morley, J. W. & Buskila, Y. Astrocytic modulation of neuronal excitability through K+ spatial buffering. *Neuroscience & Biobehavioral Reviews* **77**, 87–97 (2017).

2. Mahmoud, S., Gharagozloo, M., Simard, C. & Gris, D. Astrocytes Maintain Glutamate Homeostasis in the CNS by Controlling the Balance between Glutamate Uptake and Release. *Cells* **8**, 184 (2019).

3. Mola, M. G. *et al.* Cell Volume Regulation Mechanisms in Differentiated Astrocytes. *Cell Physiol Biochem* **55**, 196–212 (2021).

4. Bélanger, M., Allaman, I. & Magistretti, P. J. Brain Energy Metabolism: Focus on Astrocyte-Neuron Metabolic Cooperation. *Cell Metabolism* **14**, 724–738 (2011).

5. Verkhratsky, A. & Nedergaard, M. Physiology of Astroglia. *Physiological Reviews* **98**, 239–
389 (2018).

6. Benfenati, V. & Ferroni, S. Water transport between CNS compartments: functional and
molecular interactions between aquaporins and ion channels. *Neuroscience* **168**, 926–940
(2010).

7. Koch, C. & Segev, I. The role of single neurons in information processing. *Nature*
Neuroscience **3**, 1171–1177 (2000).

8. Luo, L. *et al.* Ten years of Nature Reviews Neuroscience: insights from the highly cited.
Nature Reviews Neuroscience **11**, 718–726 (2010).

9. Krogh, A. What are artificial neural networks? *Nature Biotechnology* **26**, 195–197 (2008).

10. Bassett, D. S. & Gazzaniga, M. S. Understanding complexity in the human brain. *Trends*
Cogn Sci **15**, 200–209 (2011).

11. Bassett, D. S. & Bullmore, E. Small-world brain networks. *Neuroscientist* **12**, 512–523
(2006).

12. Bassett, D. S. & Sporns, O. Network neuroscience. *Nature Neuroscience* **20**, 353–364
(2017).

13. Mederos, S., González-Arias, C. & Perea, G. Astrocyte–Neuron Networks: A Multilane
Highway of Signaling for Homeostatic Brain Function. *Frontiers in Synaptic Neuroscience*
10, (2018).

14. Giaume, C., Koulakoff, A., Roux, L., Holcman, D. & Rouach, N. Astroglial networks: a step
further in neuroglial and gliovascular interactions. *Nature Reviews Neuroscience* **11**, 87–99
(2010).

15. Scemes, E. & Giaume, C. Astrocyte calcium waves: what they are and what they do. *Glia*
54, 716–725 (2006).

16. Semyanov, A., Henneberger, C. & Agarwal, A. Making sense of astrocytic calcium signals
— from acquisition to interpretation. *Nature Reviews Neuroscience* **21**, 551–564 (2020).

17. Wang, Y. *et al.* Accurate quantification of astrocyte and neurotransmitter fluorescence

dynamics for single-cell and population-level physiology. *Nat Neurosci* **22**, 1936–1944

(2019).

18. Lia, A. *et al.* Calcium Signals in Astrocyte Microdomains, a Decade of Great Advances.

Frontiers in Cellular Neuroscience **15**, (2021).

19. Wolfes, A. C. *et al.* A novel method for culturing stellate astrocytes reveals spatially distinct

Ca2+ signaling and vesicle recycling in astrocytic processes. *J Gen Physiol* **149**, 149–170

(2017).

20. Agarwal, A. *et al.* Transient Opening of the Mitochondrial Permeability Transition Pore

Induces Microdomain Calcium Transients in Astrocyte Processes. *Neuron* **93**, 587-605.e7

(2017).

21. Lagos-Cabré, R., Burgos-Bravo, F., Avalos, A. M. & Leyton, L. Connexins in Astrocyte

Migration. *Frontiers in Pharmacology* **10**, (2020).

22. Szu, J. I. & Binder, D. K. The Role of Astrocytic Aquaporin-4 in Synaptic Plasticity and

Learning and Memory. *Frontiers in Integrative Neuroscience* **10**, (2016).

23. Bortolami, A. & Sesti, F. Ion channels in neurodevelopment: lessons from the Integrin-

KCNB1 channel complex. *Neural Regeneration Research* **18**, (2023).

24. Nagelhus, E. A. & Ottersen, O. P. Physiological Roles of Aquaporin-4 in Brain. *Physiological*

Reviews **93**, 1543–1562 (2013).

25. Ohno, Y., Kunisawa, N. & Shimizu, S. Emerging Roles of Astrocyte Kir4.1 Channels in the

Pathogenesis and Treatment of Brain Diseases. *International Journal of Molecular Sciences*

22, (2021).

26. Benfenati, V. *et al.* An aquaporin-4/transient receptor potential vanilloid 4 (AQP4/TRPV4)

complex is essential for cell-volume control in astrocytes. *Proceedings of the National*

Academy of Sciences **108**, 2563–2568 (2011).

27. Fan, Y.-Y. & Huo, J. A1/A2 astrocytes in central nervous system injuries and diseases:

Angels or devils? *Neurochemistry International* **148**, 105080 (2021).

28. Ben Haim, L. & Escartin, C. Astrocytes and neuropsychiatric symptoms in

neurodegenerative diseases: Exploring the missing links. *Current Opinion in Neurobiology*

72, 63–71 (2022).

29. Homkajorn, B., Sims, N. R. & Muyderman, H. Connexin 43 regulates astrocytic migration

and proliferation in response to injury. *Neuroscience Letters* **486**, 197–201 (2010).

30. Rodnight, R. B. & Gottfried, C. Morphological plasticity of rodent astroglia. *Journal of*

Neurochemistry **124**, 263–275 (2013).

31. Gradisnik, L. & Velnar, T. Astrocytes in the central nervous system and their functions in

health and disease: A review. *World J Clin Cases* **11**, 3385–3394 (2023).

32. Liddelow, S. A. & Barres, B. A. Reactive Astrocytes: Production, Function, and Therapeutic

Potential. *Immunity* **46**, 957–967 (2017).

33. Escartin, C. *et al.* Reactive astrocyte nomenclature, definitions, and future directions. *Nature*

Neuroscience **24**, 312–325 (2021).

34. Lawal, O., Ulloa Severino, F. P. & Eroglu, C. The role of astrocyte structural plasticity in

regulating neural circuit function and behavior. *Glia* **70**, 1467–1483 (2022).

35. Zhou, B., Zuo, Y.-X. & Jiang, R.-T. Astrocyte morphology: Diversity, plasticity, and role in

neurological diseases. *CNS Neuroscience & Therapeutics* **25**, 665–673 (2019).

36. Baldwin, K. T., Murai, K. K. & Khakh, B. S. Astrocyte morphology. *Trends Cell Biol* S0962-

8924(23)00204–0 (2023) doi:10.1016/j.tcb.2023.09.006.

37. Nicchia, G. P. *et al.* Aquaporin-4 orthogonal arrays of particles from a physiological and

pathophysiological point of view. *Wiley Interdisciplinary Reviews: Membrane Transport and*

Signaling **2**, 143–154 (2013).

38. Mola, M. G. *et al.* The speed of swelling kinetics modulates cell volume regulation and

calcium signaling in astrocytes: A different point of view on the role of aquaporins. *Glia* **64**,

139–154 (2016).

39. Posati, T. *et al.* A Nanoscale Interface Promoting Molecular and Functional Differentiation of

Neural Cells. *Scientific Reports* **6**, 31226 (2016).

40. Saracino, E. *et al.* Extracellular Recording Systems: A Glial-Silicon Nanowire Electrode

Junction Enabling Differentiation and Noninvasive Recording of Slow Oscillations from

Primary Astrocytes (Adv. Biosys. 4/2020). *Advanced Biosystems* **4**, 2070044 (2020).

41. O'Neill, K. M. *et al.* Decoding Natural Astrocyte Rhythms: Dynamic Actin Waves Result from

Environmental Sensing by Primary Rodent Astrocytes. *Adv Biol (Weinh)* e2200269 (2023)

doi:10.1002/adbi.202200269.

42. Giralt, A., Coura, R. & Girault, J.-A. Pyk2 is essential for astrocytes mobility following brain

lesion. *Glia* **64**, 620–634 (2016).

43. Mitsushima, M., Takahashi, H., Shishido, T., Ueda, K. & Kioka, N. Abl kinase interacts with

and phosphorylates vinexin. *FEBS Letters* **580**, 4288–4295 (2006).

44. Tomás, M. *et al.* Protective effects of lysophosphatidic acid (LPA) on chronic ethanol-

induced injuries to the cytoskeleton and on glucose uptake in rat astrocytes. *Journal of*

Neurochemistry **87**, 220–229 (2003).

45. Bowser, D. N. & Khakh, B. S. ATP Excites Interneurons and Astrocytes to Increase Synaptic

Inhibition in Neuronal Networks. *J. Neurosci.* **24**, 8606 (2004).

46. Guthrie, P. B. *et al.* ATP Released from Astrocytes Mediates Glial Calcium Waves. *J.*

Neurosci. **19**, 520 (1999).

47. Zimmermann, H. Signalling via ATP in the nervous system. *Trends in Neurosciences* **17**,

420–426 (1994).

48. Burnstock, G. Cotransmission. *Current Opinion in Pharmacology* **4**, 47–52 (2004).

49. Giaume, C. & Venance, L. Intercellular calcium signaling and gap junctional communication

in astrocytes. *Glia* **24**, 50–64 (1998).

50. Yu, P., Wang, H., Katagiri, Y. & Geller, H. M. An In Vitro Model of Reactive Astrogliosis and

Its Effect on Neuronal Growth. *Methods Mol Biol* **814**, 327–340 (2012).

51. Ahmed, Z. S. O., Hussein, S., Ghandour, R. A., Azouz, A. A. & El-Sakhawy, M. A.

Evaluation of the effect of methotrexate on the hippocampus, cerebellum, liver, and kidneys

of adult male albino rat: Histopathological, immunohistochemical and biochemical studies.

Acta Histochemica **123**, 151682 (2021).

52. Bramanti, V., Tomassoni, D., Avitabile, M., Amenta, F. & Avola, R. Biomarkers of glial cell

proliferation and differentiation in culture. *FBS* **2**, 558–570 (2010).

53. Cibelli, A. *et al.* Aquaporin-4 and transient receptor potential vanilloid 4 balance in early

postnatal neurodevelopment. *Glia* **72**, 938–959 (2024).

54. Berger, J. V. *et al.* Opposite regulation of metabotropic glutamate receptor 3 and

metabotropic glutamate receptor 5 by inflammatory stimuli in cultured microglia and

astrocytes. *Neuroscience* **205**, 29–38 (2012).

55. Song, C., Zhang, Y. & Dong, Y. Acute and subacute IL-1β administrations differentially

modulate neuroimmune and neurotrophic systems: possible implications for neuroprotection

and neurodegeneration. *Journal of Neuroinflammation* **10**, 826 (2013).

56. Tilleux, S. & Hermans, E. Neuroinflammation and regulation of glial glutamate uptake in

neurological disorders. *Journal of Neuroscience Research* **85**, 2059–2070 (2007).

57. Tilleux, S., Berger, J. & Hermans, E. Induction of astrogliosis by activated microglia is

associated with a down-regulation of metabotropic glutamate receptor 5. *Journal of*

Neuroimmunology **189**, 23–30 (2007).

58. Sullivan, S. M. *et al.* Cytoskeletal Anchoring of GLAST Determines Susceptibility to Brain

Damage: AN IDENTIFIED ROLE FOR GFAP *. *Journal of Biological Chemistry* **282**, 29414–

29423 (2007).

59. Toops, K. A., Hagemann, T. L., Messing, A. & Nickells, R. W. The effect of glial fibrillary

acidic protein expression on neurite outgrowth from retinal explants in a permissive

environment. *BMC Research Notes* **5**, 693 (2012).

60. Chalfoun, J. *et al.* FogBank: a single cell segmentation across multiple cell lines and image

modalities. *BMC Bioinformatics* **15**, 431 (2014).

61. Giovannucci, A. *et al.* CaImAn an open source tool for scalable calcium imaging data

analysis. *eLife* **8**, e38173 (2019).

62. Bowen, Z., Winkowski, D. E., Seshadri, S., Plenz, D. & Kanold, P. O. Neuronal Avalanches

in Input and Associative Layers of Auditory Cortex. *Frontiers in Systems Neuroscience* **13**,

(2019).

63. Simonsen, I., Hansen, A. & Nes, O. M. Determination of the Hurst exponent by use of

wavelet transforms. *Phys. Rev. E* **58**, 2779–2787 (1998).

64. Generalized Hurst exponent.

https://www.mathworks.com/matlabcentral/fileexchange/30076-generalized-hurst-exponent

(2023).

65. Averbeck, B. B., Latham, P. E. & Pouget, A. Neural correlations, population coding and

computation. *Nat Rev Neurosci* **7**, 358–366 (2006).

66. Panzeri, S., Schultz, S. R., Treves, A. & Rolls, E. T. Correlations and the encoding of

information in the nervous system. *Proc Biol Sci* **266**, 1001–1012 (1999).

67. Cohen, M. R. & Kohn, A. Measuring and interpreting neuronal correlations. *Nat Neurosci* **14**,

811–819 (2011).

68. Kohn, A., Coen-Cagli, R., Kanitscheider, I. & Pouget, A. Correlations and Neuronal

Population Information. *Annual Review of Neuroscience* **39**, 237–256 (2016).

69. Aguado, F., Espinosa-Parrilla, J. F., Carmona, M. A. & Soriano, E. Neuronal Activity

Regulates Correlated Network Properties of Spontaneous Calcium Transients in Astrocytes

In Situ. *J Neurosci* **22**, 9430–9444 (2002).

70. Daw, C. S., Finney, C. E. A. & Tracy, E. R. A review of symbolic analysis of experimental

data. *Review of Scientific Instruments* **74**, 915–930 (2003).

71. Cao, Y., Tung, W., Gao, J. B., Protopopescu, V. A. & Hively, L. M. Detecting dynamical

changes in time series using the permutation entropy. *Phys. Rev. E* **70**, 046217 (2004).

72. Shiozawa, K., Uemura, T. & Tokuda, I. T. Detecting the dynamical instability of complex

time series via partitioned entropy. *Phys. Rev. E* **107**, 014207 (2023).

73. Lungarella, M., Pitti, A. & Kuniyoshi, Y. Information transfer at multiple scales. *Phys. Rev. E*

76, 056117 (2007).

74. Schreiber, T. Measuring Information Transfer. *Phys. Rev. Lett.* **85**, 461–464 (2000).

75. Lehnertz, K. Ordinal methods for a characterization of evolving functional brain networks.

Chaos: An Interdisciplinary Journal of Nonlinear Science **33**, 022101 (2023).

76. Staniek, M. & Lehnertz, K. Symbolic Transfer Entropy. *Phys. Rev. Lett.* **100**, 158101 (2008).

77. Staniek, M. & Lehnertz, K. Symbolic transfer entropy: inferring directionality in biosignals.

54, 323–328 (2009).

78. STANIEK, M. & LEHNERTZ, K. PARAMETER SELECTION FOR PERMUTATION

ENTROPY MEASUREMENTS. *Int. J. Bifurcation Chaos* **17**, 3729–3733 (2007).

79. Bandt, C. & Pompe, B. Permutation Entropy: A Natural Complexity Measure for Time

Series. *Phys. Rev. Lett.* **88**, 174102 (2002).

80. Li, S., Xiao, Y., Zhou, D. & Cai, D. Causal inference in nonlinear systems: Granger causality

versus time-delayed mutual information. *Phys. Rev. E* **97**, 052216 (2018).

81. Vastano, J. A. & Swinney, H. L. Information transport in spatiotemporal systems. *Phys. Rev.*

Lett. **60**, 1773–1776 (1988).

82. Khakh, B. S. & Sofroniew, M. V. Diversity of astrocyte functions and phenotypes in neural

circuits. *Nature Neuroscience* **18**, 942–952 (2015).

83. Glaum, S. R., Holzwarth, J. A. & Miller, R. J. Glutamate receptors activate Ca2+

mobilization and Ca2+ influx into astrocytes. *Proceedings of the National Academy of*

Sciences **87**, 3454–3458 (1990).

84. Charles, A. C., Merrill, J. E., Dirksen, E. R. & Sandersont, M. J. Intercellular signaling in glial

cells: Calcium waves and oscillations in response to mechanical stimulation and glutamate.

Neuron **6**, 983–992 (1991).

85. Fraser, A. M. & Swinney, H. L. Independent coordinates for strange attractors from mutual

information. *Phys. Rev. A* **33**, 1134–1140 (1986).

86. Schiweck, J., Eickholt, B. J. & Murk, K. Important Shapeshifter: Mechanisms Allowing

Astrocytes to Respond to the Changing Nervous System During Development, Injury and

Disease. *Frontiers in Cellular Neuroscience* **12**, (2018).

87. Li, S., Graham, E. S. & Unsworth, C. P. Extracellular ATP release predominantly mediates

Ca2+ communication locally in highly organised, stellate-Like patterned networks of adult

human astrocytes. *PLOS ONE* **18**, e0289350 (2023).

88. Takano, H. *et al.* Micropatterned Substrates: Approach to Probing Intercellular

Communication Pathways. *Anal. Chem.* **74**, 4640–4646 (2002).

749