1 **Title:** Molecular subtypes of high-grade serous ovarian cancer across racial groups and gene
2 expression platforms
3

4 **Authors:** Natalie R. Davidson*[1], Mollie E. Barnard*[2,3], Ariel A. Hippen[4], Amy Campbell[4],
5 Courtney E. Johnson[5], Gregory P. Way[1,4], Brian K. Dalley[6], Andrew Berchuck[7], Lucas A. Salas[8],
6 Lauren C. Peres[9], Jeffrey R. Marks[10], Joellen M. Schildkraut[†5], Casey S. Greene[†1,4], Jennifer A.
7 Doherty[†2,8]
8
9 *Denotes equal contributions
10 [†]Denotes equal contributions
11
Author affiliations:
[1] Department of Biomedical Informatics, School of Medicine, University of Colorado Anschutz
Medical Campus, Aurora, CO, USA
[2] Huntsman Cancer Institute and the Department of Population Health Sciences at the Spencer
Fox Eccles School of Medicine, University of Utah, Salt Lake City, UT, USA
[3] Slone Epidemiology Center, Boston University Chobanian & Avedisian School of Medicine,
Boston, MA, USA
[4] Department of Systems Pharmacology and Translational Therapeutics, Perelman School of
Medicine, University of Pennsylvania, Philadelphia, PA, USA
[5] Department of Epidemiology, Rollins School of Public Health, Emory University, Atlanta, GA,
USA
[6] Huntsman Cancer Institute, University of Utah, Salt Lake City, UT, USA
[7] Department of Obstetrics and Gynecology, Duke University, Durham, NC
[8] Department of Epidemiology, Geisel School of Medicine, Dartmouth College, Hanover, NH,
USA
[9] Department of Cancer Epidemiology, H. Lee Moffitt Cancer Center and Research Institute,
Tampa, FL, USA
[10] Department of Surgery, Duke University School of Medicine, Durham, NC 27710, USA

30
31 **Running title:** High-grade serous ovarian cancer subtypes in racial groups
32

Funding statement: This work was supported by the National Cancer Institute (NCI) of the
National Institutes of Health (R01 CA200854 to JAD and JMS, R01 CA142081 to JMS, R01
CA076016 to JMS, R01 CA237170 to CSG and JAD, K99/R00CA218681 to LCP). Research
reported in this publication utilized the High-Throughput Genomics and Cancer Bioinformatics
Shared Resource at Huntsman Cancer Institute at the University of Utah and was supported by
the National Cancer Institute of the National Institutes of Health under Award Number
P30CA042014. Dr. Barnard received support from the National Cancer Institute (K00
CA212222) and the Karin Grunebaum Cancer Research Foundation. The content of the

manuscript is solely the responsibility of the authors and does not necessarily represent the
official views of the National Cancer Institute or the National Institutes of Health.
43

Conflict of interest declaration: LCP received grant funding from Bristol Myers Squibb for
work external to this manuscript. The remaining authors declare no potential conflicts of interest.

Corresponding authors:
1. Mollie E. Barnard, ScD
72 E. Concord St.
Boston, MA, 02118
T: (617) 206-6142
F: (617) 738-5119
barnard@bu.edu
54
2. Natalie R. Davidson, PhD
Department of Biomedical Informatics
University of Colorado
Anschutz Medical Campus
1890 N Revere Ct, Aurora, CO
natalie.davidson@cuanschutz.edu

61
62 **Statement of Significance:** A single pipeline was used to subtype ovarian high-grade serous
63 carcinoma (HGSC) with array-based or RNA-Seq gene expression data. Subtype distributions
64 differed by race, but subtype-specific survival was similar across racial groups.
65
66 Abstract word count: 238
67 Text word count: 4,734
68 Number of tables: 2
69 Number of figures: 4
70 Supplemental material: 11 supplemental figures, 7 supplemental tables, 1 supplemental note
71
72

Abstract

Introduction: High-grade serous carcinoma (HGSC) gene expression subtypes are associated

with differential survival. We characterized HGSC gene expression in Black individuals and

considered whether gene expression differences by race may contribute to poorer HGSC survival

among Black versus non-Hispanic White individuals.

Methods: We included newly generated RNA-Seq data from Black and White individuals, and

array-based genotyping data from four existing studies of White and Japanese individuals. We

assigned subtypes using K-means clustering. Cluster- and dataset-specific gene expression

patterns were summarized by moderated t-scores. We compared cluster-specific gene expression

patterns across datasets by calculating the correlation between the summarized vectors of

moderated t-scores. Following mapping to The Cancer Genome Atlas (TCGA)-derived HGSC

subtypes, we used Cox proportional hazards models to estimate subtype-specific survival by

dataset.

Results: Cluster-specific gene expression was similar across gene expression platforms.

Comparing the Black study population to the White and Japanese study populations, the

immunoreactive subtype was more common (39% versus 23%-28%) and the differentiated

subtype less common (7% versus 22%-31%). Patterns of subtype-specific survival were similar

between the Black and White populations with RNA-Seq data; compared to mesenchymal cases,

the risk of death was similar for proliferative and differentiated cases and suggestively lower for

immunoreactive cases (Black population HR=0.79 [0.55, 1.13], White population HR=0.86

[0.62, 1.19]).

Conclusions: A single, platform-agnostic pipeline can be used to assign HGSC gene expression

subtypes. While the observed prevalence of HGSC subtypes varied by race, subtype-specific

survival was similar.

 Introduction

Ovarian cancer is a highly fatal malignancy comprised of multiple histologically-defined

subtypes (i.e., "histotypes"). High-grade serous carcinoma (HGSC) is the most common

histotype (1), and an important contributor to ovarian cancer mortality (2–4). HGSC is also

molecularly heterogeneous, which is relevant to both prognosis and treatment. Prior studies have

described between three and five molecularly distinct subtypes (5–10), and, while no gold

standard exists for defining these subtypes, they are commonly mapped to the four TCGA-

derived subtypes, which are similar to those reported in Tothill et al., 2008: mesenchymal

(Tothill C1.MES), proliferative (Tothill C5.PRO), immunoreactive (Tothill C2.IMM), and

differentiated (Tothill C4.DIF) (5,11). Key characteristics of the mesenchymal subtype include

high expression of HOX genes, increased stromal components, and poor survival (5,9,11). The

proliferative subtype has been characterized by low expression of ovarian tumor markers (e.g.,

MUC1, MUC16), high expression of transcription factors, and intermediate survival (5,9,11).

Defining characteristics of the immunoreactive subtype include the enrichment of genes and

pathways associated with an immune response, including CD3+/CD8+ T-cell markers and genes

in the CXCL9, CXCL10, CXCL11/CXCR3 axis for immune activation, and more favorable

survival (5,9,11). The differentiated subtype has been the most difficult to characterize and

reproduce (7,12), yet multiple studies have described it as having high expression of ovarian

tumor markers (e.g., MUC1, MUC16, SLPI) and intermediate to good survival (5,9).

Since HGSC gene expression-based subtypes have potential clinical utility, it is of

interest to develop a clinical-grade, gene expression-based subtype classifier that is applicable

across diverse populations. One recently developed gene-set assay, the Predictor of high-grade-

serous Ovarian carcinoma molecular subTYPE (PrOTYPE) assay, was derived using array-based

gene expression data from mostly non-Hispanic White individuals and migrated to a NanoString

platform (11). The PrOTYPE assay has been reported to classify HGSC into the four TCGA-

derived subtypes with >95% accuracy (11). Here, we leverage RNA-sequencing (RNA-seq) data

to characterize HGSC gene expression subtypes among a more racially diverse study population,

including a population of >300 self-identified Black individuals. Prior studies have noted that

Black individuals have poorer overall ovarian cancer survival (13), and poorer three-year and

six-year HGSC survival when compared to White individuals (14,15), so inclusion of Black

individuals in the derivation of HGSC molecular subtypes is especially important.

In the present study, we used K-means clustering to assign HGSC tumors from Black,

White, and Japanese individuals to molecular subtypes (7). Following subtype assignment, we

compared subtype-specific gene expression, subtype frequency, and subtype-specific survival

across all possible pairs of studies with different racial distributions (e.g., Black, White,

Japanese), and different data types (array-based gene expression data, and RNA-seq).

Methods

Primary study population

We included epithelial ovarian cancer cases enrolled in one of two population-based

case-control studies, the North Carolina Ovarian Cancer Study (NCOCS, diagnosis dates 1999-

2005) (16), and the African American Cancer Epidemiology Study (AACES, diagnosis dates

2010-2015) (15,17). Both studies enrolled epithelial ovarian cancer cases covering a range of

histotypes, grades, and stages, though some of the most aggressive cases were missed because

they were feeling very ill or were already deceased by the time they were invited to participate in

research (15). Written informed consent was obtained for NCOCS participants, while AACES

participants provided verbal consent and signed medical record and pathology release forms to

allow for access to tumor tissue. All cases were confirmed via centralized pathology review.

Both the NCOCS and AACES studies were approved by the Duke Medical Center Institutional

Review Board (IRB) and the IRBs of participating enrollment sites.

Together, the AACES and NCOCS included 747 Black ovarian cancer cases, 464 of

which were HGSC. Of these, 325 provided consent to participate in biospecimen-based research

and had adequate tissue available to pursue RNA extraction (**Supplemental Figure 1A**). Fifty-

three of these cases were excluded from gene expression analyses due to a history of neoadjuvant

chemotherapy, which can influence observed gene expression (**Supplemental Figure 1B**).

Following these exclusions, there were 272 self-identified Black or African American cases (3

Hispanic; 269 non-Hispanic) who we subsequently refer to as "SchildkrautB". The NCOCS

study included 1,014 White cases, 484 of which were HGSC. Of these, 316 provided consent to

participate in biospecimen-based research and had sufficient tissue available to pursue RNA

extraction (**Supplemental Figure 2A**). None of these cases had neoadjuvant chemotherapy prior

to tissue collection, so all were considered eligible for gene expression analyses. We

subsequently refer to this set of 316 non-Hispanic White cases as "SchildkrautW".

Demographic characteristics, disease characteristics, and vital status were available for all

individuals included in SchildkrautB and SchildkrautW. Information on age at diagnosis was

obtained from questionnaires and pathology reports. Tumor stage, debulking status, and use of

neoadjuvant chemotherapy were abstracted from medical records and pathology reports. The

proportions of intratumoral CD3+ T cells and CD3+/CD8+ suppressor T cells were obtained

from multiplex immunofluorescence staining of formalin-fixed paraffin-embedded (FFPE) tissue

(18). Vital status was assessed using data from population-based cancer registries, obituaries,

LexisNexis, and the National Death Index.

Acquisition of gene expression data

RNA was extracted from FFPE tumor tissue and stored at -80°C. An initial quality

control (QC) evaluation revealed substantial RNA degradation, so a re-purification step

consisting of DNAase treatment and purification on a Zymo research spin column was

completed before library preparation to reduce the bulk of degraded RNA product (i.e., RNA

product <200 nucleotides in length). Following re-purification, RNA libraries were prepared

from total RNA samples (5-100 ng) using reagents from the Illumina Stranded mRNA Prep (cat#

20020189) and the Illumina RNA UD Indexes Set (20091657) for reverse transcription, adapter

ligation, and PCR amplification. Amplified libraries were hybridized to biotin-labeled probes

from the Illumina Exome Panel (cat# 20020183) using the Illumina RNA Fast Hyb Enrichment

kit (20040540) to generate strand-specific libraries enriched for coding regions of the

transcriptome. The quality of exon-enriched libraries was assessed on an Agilent Technologies

2200 TapeStation using a D1000 ScreenTape assay (cat# 5067-5582 and 5067-5583). The

molarity of adapter-modified molecules was defined by quantitative PCR using the Kapa

Biosystems Kapa Library Quant Kit (cat#KK4824). Individual libraries were normalized to 0.95

nM in preparation for Illumina sequence analysis. Sequencing libraries were chemically

denatured and applied to an Illumina NovaSeq flow cell using the NovaSeq XP workflow

(20043131). Following the transfer of the flow cell to an Illumina NovaSeq 6000 instrument, a

150 x 150 cycle paired-end sequence run was performed using a NovaSeq 6000 S4 reagent Kit

v1.5 (20028312).

Quantification of gene expression data

We trimmed adapters and filtered read quality using fastp (19). We filtered to reads with

a PHRED score of at least 15 and a length of at least 20 base pairs. While a minimum PHRED

score of 15 may include some reads with low quality, we found that most bases across all

samples in SchildkrautB and SchildkrautW had a quality score greater than 30 (**Supplemental**

Figure 3). We quantified paired-end reads with Salmon (version 1.4.0) (20) using GRCh38

release 95. We used the seqBias and gcBias flags to correct for sequence-specific biases. We also

used the recommended rangeFactorizationBins parameter value 4, which improves quantification

accuracy on difficult-to-quantify transcripts. We then filtered out low-expression genes by

excluding genes with a median expression of 0 within a dataset (SchildkrautB: 10,620 genes

removed, SchildkrautW: 10,410 genes removed). We library-size normalized samples using

upper quantile normalization. This normalization matches the 85^{th} percentile across samples to

correct for library size differences across samples.

Overview of comparator study populations

Consistent with the data processing pipeline used by Way *et al.* (7), we included data

from the TCGA (platform: AffymetrixHT_HG-U133A), Yoshihara (platform: AgilentG4112F),

and Tothill (platform: AffymetrixHG-U133Plus2) datasets included in the R package

curatedOvarianData (21), and additional data from the dataset "Mayo" (GEO Accession:

GSE74357, platform: AgilentG4112F). All gene expression data generated from these studies

was derived from fresh frozen tumor tissue. Using the R package doppelgangR (22), we

identified four more duplicate pairs across studies than found previously (7) and one pair of

duplicate samples in SchildkrautW.

Clustering Methods

We performed clustering analyses by modifying the pipeline from Way *et al.* (7) for

inclusion of RNA-Seq data. Briefly, for K-means clustering, we used the R package cluster

(version 2.0.6) (23), and for non-negative matrix factorization (NMF), we used the R package

NMF (version 0.20.6) (24). We identified the top 1,500 genes with the largest Median Absolute

Deviance (MAD) from each of the six datasets (SchildkrautB, SchildkrautW, TCGA, Mayo,

Yoshihara, and Tothill), then used their union set of 4,355 genes for clustering (**Supplemental**

Table 1). Consensus clusters for both K-means and NMF were identified using multiple random

starts for each value of K=2-4 (K-means starts: 100; NMF starts: 10). Each dataset was clustered

independently using the same set of genes.

We compared the similarity of clusters across datasets and clustering methods using the

same approach as Way *et al.* (7). We used significance analysis of microarray (SAM) (25,26) to

get cluster-specific vectors of moderated t-scores; this provided us with gene-specific expression

patterns of each cluster within each dataset. We restricted to genes that were assayed and

expressed across all datasets, which we denote as "common genes" (**Supplemental Table 2**).

Using these 8,360 "common genes", we compared clusters across: (1) datasets, and (2) clustering

methodologies by calculating the Pearson correlation between two vectors of moderated t-scores.

To use this approach on the RNA-Seq datasets, we log10(x+1) transformed the normalized

counts to more closely match the data distribution of the microarray datasets. We performed a

simulation study to examine SAM's performance on log10 transformed counts relative to

DESeq2's performance on raw counts and found comparable performance (**Supplemental Note**

1).

Additionally, we compared the similarity of clusters in a global and aligned principal

components analysis (PCA) projection. We filtered the datasets to only include the MAD genes

used in our clustering pipeline, then within-sample scaled the expression values to ensure

comparative ranges of expression across all datasets. We log10(x+1) transformed the RNA-Seq

expression values before scaling to better match the RNA-Seq and microarray data distributions.

After normalization, we concatenated the datasets to calculate a global PCA projection. We then

subtracted each dataset's total centroid from each dataset's cluster centroid to align each dataset

such that it was centered at the origin of the PCA projection. This allowed us to compare the

relative differences between each cluster across all datasets.

Survival analyses by cluster

We generated Kaplan-Meier survival curves to visualize overall survival and subtype-

specific survival within each dataset. We used Cox proportional hazards (PH) models to

calculate hazard ratios (HR) and 95% confidence intervals (CIs) quantifying the differences in

survival across HGSC subtypes. For five of the six datasets (SchildkrautB, SchildkrautW,

TCGA, Mayo, Tothill), Cox PH models were adjusted for three factors that independently

influence survival: age (in 5-year age groups), stage (I, II, III and IV), and debulking status

(optimal debulking, suboptimal or missing debulking status). Yoshihara *et al.* did not provide

information on age at HGSC diagnosis, so models for this dataset were adjusted for stage and

debulking status only.

Cluster comparison between pipeline runs and consensusOV

To ensure that our run of the updated Way *et al.* (7) pipeline was producing similar

results to those from previously published subtype predictors, we compared the assigned cluster

labels for each case in the TCGA, Mayo, Yoshihara, and Tothill datasets. To quantify the

similarity between pipeline results, we rephrased the cluster comparison problem as a prediction

problem. The ground-truth labels were the previously published cluster IDs (7) or consensusOV

(27) predictions for each case, and the predicted labels were our newly assigned cluster labels.

consensusOV version 1.16.0 was run using default parameters and no significance filtering. We

used balanced accuracy as the metric for comparing current and published labels due to

variations in cluster sizes. Balanced accuracy is defined in the following equations.

$Balanced\ Accuracy\ =\ (Sensitivity\ +\ Specificity\ )\ /\ 2$

$Sensitivity = \dfrac{Number\ of\ True\ Positives}{Number\ of\ True\ Positives\ +\ Number\ of\ False\ Negatives}$

$Specificity = \dfrac{Number\ of\ True\ Negatives}{Number\ of\ True\ Negatives\ +\ Number\ of\ False\ Positives}$

Sensitivity analysis of K-means cluster assignments

K-means is sensitive to outliers and local minima; therefore, we quantified the stability of

the cluster labels when using a subsample within each dataset. First, we removed any samples

that were more than 1.5x the interquartile range from the first or third quartiles in the first five

PCs. Using the remaining samples, we subsampled 80% of the samples within each dataset and

re-ran K-means clustering. We then matched the new cluster labels to the original cluster labels

using a greedy approach. We matched labels by finding the cluster IDs with the highest sample

overlap. We did this iteratively, removing each cluster from consideration once its matched

cluster ID was assigned.

Data Availability Statement

We provide all software under the BSD 3-Clause License. We include scripts for the

analyses presented in this manuscript, including RNA-Seq quantification, quality control,

clustering, and figure generation. To support reproducibility, we provide code to recreate the

environments and re-run both the RNA-Seq data analysis

(https://github.com/greenelab/hgsc_rnaseq_cluster/) and our updated clustering pipeline

(https://github.com/greenelab/hgsc_rnaseq_clustering_pipeline/). All derived expression data are

publicly available and included in Supplemental Tables 6 and 7.

Results

Data quality

For both SchildkrautB and SchildkrautW, we further excluded samples due to possible

technical artifacts (**Supplemental Tables 3 and 4)**. In SchildkrautB and SchildkrautW, 14 and 4

cases were re-sequenced, respectively, due to insufficient read depth, and the transcriptomic

profiles from the first sequencing attempt were excluded. After resequencing, all18 cases

attained a read depth comparable to the other sequenced samples and progressed to the next step

of the quality control pipeline (**Supplemental Figures 1B and 2B**). All other samples in both

datasets were considered high quality, with over 83.3% of reads having a base quality of at least

30 (**Supplemental Figure 3**). After normalization, read count distributions were similar across

sequencing batches, including samples previously identified by the sequencing core as low-

quality (**Supplemental Figure 4C-H)**. However, a small subset of samples had lower-than-

expected read counts. To account for this, we removed all samples where the median normalized

read count was below 925 or where the bottom 25th quantile read count for a sample was below

30. This additional read count filter removed 10 SchildkrautB cases and 5 SchildkrautW cases

(**Supplemental Figure 4A-B**). The final technical exclusion for SchildkrautB and SchildkrautW

removed samples flagged by doppelgangR (22) as having overly similar expression patterns,

suggesting they originated from the same tumor. Only SchildkrautW had a pair of samples

identified as too similar (NCO0557 and NCO0625).

Following the SchildkrautB and SchildkrautW technical exclusions, we applied the Way

pipeline exclusions to the TCGA, Mayo, Yoshihara, and Tothill datasets, and we applied

doppelgangR exclusions to all datasets. The sizes of our final analytic data sets were as follows:

SchildkrautB (n=262); SchildkrautW (n=309); TCGA (n = 499; phs000178) (5); Mayo (n = 377;

GSE74357) (9); Yoshihara (n = 255; GSE32062.GPL6480) (28), and Tothill (n = 241;

GSE9891) (6).

Validation of updated clustering pipeline

We updated an HGSC subtyping pipeline (7) that previously identified clusters across

four microarray studies from the United States, Japan, and Australia. While most of our pipeline

was the same, we used balanced accuracy to quantify the similarity in output between pipelines.

Balanced accuracy for our K-means clusters was consistently ≥0.91 across all datasets for K=2-4

(**Supplemental Figure 5**). The limited differences that arose were primarily due to changes in

stochastic elements of the clustering methods and software packages, such as differences in

computing environments, random seeds, and package versions. Another source of variation

between the output of the pipelines was the addition of the Schildkraut datasets since the genes

used for clustering must be shared across all datasets and have a high MAD in at least one

dataset (**Methods**). After adding the Schildkraut datasets, 893 genes that were previously

denoted as common were reclassified as MAD genes, 609 MAD genes were re-classified as

common genes, and 911 genes were removed from consideration. A Venn diagram of the

removed genes is provided in **Supplemental Figure 6**.

To ensure that clustering outcomes were robust across methods, we compared the clusters

from two different clustering methods: K-means and NMF. We derived a gene expression pattern

for each cluster using the significance analysis of microarray (SAM) moderated t-score

(**Methods**). Using this cluster-specific gene expression pattern, we calculated the Pearson

correlation between moderated t-scores for clusters identified by each method. We found

extremely high cluster concordance in all datasets between the clusters identified by NMF and

K-means when we selected two and three clusters. We saw diminished concordance between

methods when we used four clusters. Clusters in all datasets (except the Mayo dataset for K=4)

had the highest concordance with the matched cluster along the diagonal (**Supplemental Figure**

7). Since K-means clustering is sensitive to outliers, we additionally ran K-means clustering in

subsets of 80% of the data and found that, in all datasets, the majority of samples maintained

their same cluster label in more than 50% of the subsampled reclusterings (**Supplemental**

Figure 8).

Our final clustering validation was to compare our output to output from another external

HGSC subtype classifier, consensusOV (27). consensusOV combines four methods (5,6,8,9,29)

into a consensus classifier. Since consensusOV can only assign samples to the four Verhaak-

defined clusters, we only compare our K-means results for K=4. We found that our subtype calls

were very similar to the consensusOV calls across all microarray datasets, with a minimum per-

class balanced accuracy of 0.72 (**Supplemental Table 5**). We saw more discordance between

our calls and consensusOV when we compared RNA-Seq samples, with per-class balanced

accuracy ranging from 0.589-0.818 (**Supplemental Table 5**).

Gene expression patterns in the self-identified Black study population compared with other study

populations

Expression patterns for each cluster were summarized in two ways, first by plotting the

principal components of normalized and aligned centroids in each dataset, and second, using

SAM-moderated t-scores (7). In both approaches, we sought to determine if the relative cluster

distances for each individual dataset were consistent across datasets. In our principal components

(PC)-based approach, we normalized and projected all datasets into a shared PC space, then

centered them at the origin of the space. For K=3, PC2 and PC3 could separate each cluster

(**Figure 1a**). For K=4, PC2 and PC3 could not independently separate each cluster, but, together,

could differentiate them (**Figure 1b**). Furthermore, in the first three PCs, the SchildkrautB

dataset did not cluster away from any other datasets, and clustered closely with the

SchildkrautW, TCGA, and Tothill datasets.

In our second, complementary approach, we compared differential gene expression

between clusters by calculating the Pearson correlation for each cluster's SAM-moderated t-

score across pairs of datasets. **Figure 2** compares the cluster-specific gene expression pattern

between the SchildkrautB cases and cases from the other five datasets. For K=3, SchildkrautB

cluster-specific gene expression was highly correlated with cluster-specific gene expression

across all datasets, and showed the strongest correlations with SchildkrautW cluster-specific

gene expression. This provides strong evidence that the derived clusters from Black cases are the

same as the derived clusters from White and Japanese cases. A high correlation between clusters

was also found when performing pairwise comparisons among all six datasets (**Supplemental**

Figure 9). Similar to Way *et al.* (7), we found that the correlation of cluster-specific gene

expression patterns was diminished when using four, as opposed to three, clusters to describe the

data. This was evident when comparing SchildkrautB to each of the five other populations

(**Figure 2B vs. Figure 2A**) and when comparing gene expression patterns across all possible

pairs of datasets (**Supplemental Figure 10 vs. Supplemental Figure 9**).

Subtype distributions and characteristics by study population

We mapped the K-means clusters with K=4 to the four TCGA-derived HGSC subtypes to

compare the frequency of subtypes across datasets and evaluate how cancer characteristics vary

by subtype. The immunoreactive subtype was more common (39% vs 23-28%), and the

differentiated subtype less common (7% vs 22-31%) in the self-identified Black study population

when compared to the White and Japanese study populations (**Figure 3**). In analyses restricted to

SchildkrautB and SchildkrautW, FIGO stage differed by HGSC subtype among both Black

(*p*=0.040) and non-Hispanic White individuals (*p*=0.009); those with differentiated HGSC were

more likely to have a lower FIGO stage at diagnosis (**Table 1**). Immune infiltration also differed

by HGSC subtype, particularly among Black individuals (*p*=0.001 for CD3+ T cells; *p*=0.045 for

CD3+/CD8+ suppressor T cells; **Table 1**). In both Black and White individuals, immune

infiltration was higher for tumors categorized as mesenchymal or immunoreactive and lower for

tumors categorized as proliferative.

Survival patterns for clusters

Overall HGSC survival varied by study (**Supplemental Figure 11,** p-value for test of

heterogeneity in a stage-adjusted model <0.001), though patterns of subtype-specific survival

were generally similar across studies (**Figure 4**). Multivariable-adjusted hazard ratios and 95%

CIs indicated that, when compared to individuals with mesenchymal tumors, those with

immunoreactive tumors had better survival in most (SchildkrautB HR=0.79 [0.55, 1.13];

SchildkrautW HR=0.86 [0.62, 1.19]; Mayo HR=0.54 [0.39, 0.75]; Yoshihara HR=0.65 [0.39,

1.09]; Tothill HR=0.54 [0.31, 0.95]), but not all (TCGA HR=1.01 [0.69, 1.49]) study populations

(**Table 2**). Meanwhile, the risk of death among those with proliferative HGSC was not

statistically significantly different from the risk of death among those with mesenchymal tumors

in any study population (**Table 2**), and it was close to the null value of 1.00 in both SchildkrautB

(HR=0.98 [0.66, 1.46]) and SchildkrautW (HR=1.05 [0.70, 1.57]). The risk of death among those

with differentiated HGSC was not statistically significantly different from the risk of death

among those with mesenchymal tumors in five of the six study populations, and was lower in the

Mayo study population (HR=0.61 [0.44, 0.84]; **Table 2**).

Discussion

We performed a cross-platform and cross-population analysis of HGSC, including four

existing datasets, plus newly generated RNA-Seq gene expression data from 262 Black and 309

White HGSC cases. When comparing RNA-Seq data from the 309 White cases to microarray

gene expression data from the predominantly White populations that comprise the Tothill and

TCGA study populations, we observed high cluster stability within each dataset when advancing

from K=2 to K=4 (**Supplemental Figure 8**) and consistent cluster-specific gene expression

profiles across datasets (**Supplemental Figures 9 and 10**). This indicates that similar HGSC

gene expression clusters can be defined using combined data from array-based technologies in

fresh frozen tissue and RNA-seq technologies in FFPE tissue. We also observed consistent

cluster composition (**Figure 1**) and gene expression profiles when comparing Black HGSC cases

to White and Japanese HGSC cases (**Figure 2**). This indicates that HGSC gene expression

clusters are consistent across Black, White, and Japanese individuals, so it is unlikely that racial

differences in HGSC gene expression patterns are a key driver behind poorer HGSC survival in

Black populations.

Our interest in determining how an existing HGSC subtype clustering pipeline performs

with RNA-Seq data was motivated by the increasing use of RNA-Seq technology to interrogate

cancer gene expression (30). Previously published HGSC subtype clustering approaches were

designed around array-based gene expression data (5–7,9,28,31), and while array-based and

RNA-Seq technologies observe the same underlying biological processes, the data they produce

follow different data distributions (32,33). This is most clear in the methodological differences

between array-based and RNA-Seq differential gene expression methods (34–37). Here, we

demonstrated that, despite the difference in data distributions generated by array-based and

RNA-Seq technologies, the Way *et al.* (7) subtype pipeline built for array-based gene expression

data can be applied to normalized and log-transformed RNA-Seq data. The high cluster

correlations across all four array-based datasets using fresh frozen tissue and both RNA-Seq

datasets using FFPE tissue indicate that the delineation of HGSC subtypes is agnostic to

sequencing technology and fresh frozen versus FFPE tissue.

As done in previous studies, we assigned all cases to one of the four TCGA-derived

HGSC subtypes; however, as was previously reported in Way *et al.* (7), we observed that cluster-

specific gene expression was more concordant across datasets when clusters were assigned using

K=2 or K=3 compared to K=4. This was most evident when comparing each dataset's cluster-

specific expression profiles to the cluster-specific expression profiles across all of the other

datasets (**Figure 2, Supplemental Figures 9 and 10**). When considering K=3, the largest off-

diagonal correlation we observed over all datasets was 0.07 (**Supplemental Figure 9**). In

contrast, for K=4 we found much larger positive off-diagonal correlations for the clusters

observed in the array-based TCGA, Mayo, Yoshihara, and Tothill datasets, with the largest per-

dataset correlations ranging from 0.19 to 0.33 (**Supplemental Figure 10**). We also observed

large positive off-diagonal correlations for the RNA-Seq-based SchildkrautB and SchildkrautW

datasets, with the largest per-dataset correlations ranging from 0.17 to 0.34 (**Supplemental**

Figure 10). Since cluster-specific gene expression was more concordant for K=2 and K=3 versus

K=4 for all datasets, we encourage future studies to test whether a number other than four best

represents HGSC subtypes. Our results are consistent with a model where either two HGSC gene

expression axes (e.g., mesenchymal-like and immune) or a set of three HGSC subtypes (e.g., as

derived using K=3) may more effectively describe the biological variation in HGSC gene

expression than the four subtypes most commonly seen in the literature.

Beyond methodological advances, our work also provides strong evidence that HGSC

subtypes can be reproduced among Black HGSC cases and that Black, White, and Japanese

HGSC cases share similar subtype-specific gene expression profiles. At increasing values of K,

we observed patterns in cluster composition among Black HGSC cases that were consistent with

those observed among White and Japanese HGSC cases (**Figure 1, Supplemental Figure 8**).

Further, we observed strong correlations between cluster-specific gene expression in Black cases

and cluster-specific gene expression in White and Japanese cases, especially when comparing

clusters defined using K=2 and K=3 (**Figure 2**). Patterns of subtype-specific survival were also

generally consistent across populations. When compared to mesenchymal HGSC, the risk of

death was similar for proliferative and differentiated cases, and lower, but not statistically

significantly lower, for immunoreactive cases both in SchildkrautB (HR=0.79 [0.55, 1.13]) and

SchildkrautW (HR=0.86 [0.62, 1.19], **Table 2, Figure 4**).

The primary difference we observed when comparing HGSC subtypes in Black

individuals (i.e., cases in SchildkrautB) to HGSC subtypes in all other study populations was that

more Black HGSC cases had gene expression profiles consistent with the TCGA

immunoreactive subtype (39% compared to 23%-28%) and fewer Black HGSC cases had gene

expression profiles consistent with the TCGA differentiated subtype (7% compared to 22%-31%;

Figure 3). Differences in the Tothill, C1-C6, gene expression subtype signatures for Black

(n=29) versus White (n=156) ovarian cancers were observed previously, though in different

proportions (17), so it is possible that there exists true variation in the proportion of HGSC

subtypes for Black versus non-Black HGSC cases. However, it is also possible that variations in

study design across the six study populations contributed to the different subtype distributions

that we observed. For example, case-control studies like the AACES and NCOCS are unable to

enroll cases with rapidly fatal HGSC (15). This could have skewed the observed subtype

distribution for Black cases toward a greater proportion of less aggressive, immunoreactive

tumors, and artificially inflated estimates of overall survival in the SchildkrautB study population

(**Supplemental Figure 11**), as has been posited previously (15).

A key contribution of this study was that we were able to update a previously published

subtype clustering pipeline to accept either array-based gene expression data or RNA-Seq data

and validate our modifications. We also created the first large RNA-Seq dataset of HGSC in self-

identified Black cases, consisting of 262 high-quality expression profiles. This dataset allowed us

to compare the expression profiles of HGSC subtypes in Black cases against other study

populations, and it provided an opportunity to evaluate differences in subtype frequency and

survival in Black HGSC cases compared to non-Black HGSC cases. An important limitation of

this study was that we lacked adequate data to explore whether the observed racial variation in

the proportions of gene expression subtypes and survival outcomes was due to biological,

sociodemographic, or access-to-care differences.

In summary, we have updated an existing HGSC gene expression subtype classifier to be

compatible with both array-based gene expression data and RNA-Seq data. This advancement

will facilitate reproducible HGSC subtyping for research purposes and is available for use in

future studies. We have also demonstrated that the HGSC subtypes generated by our classifier

generalize to racially diverse populations, and we have indicated that HGSC subtype-specific

gene expression and subtype-specific survival are consistent across Black, White and Asian

study populations. Given our findings, we expect that a clinical HGSC gene expression assay

would benefit prognostication and treatment strategies similarly for women from multiple racial

and ethnic backgrounds.

Acknowledgements

We would like to acknowledge the AACES interviewers, Christine Bard, LaTonda Briggs,

Whitney Franz (North Carolina) and Robin Gold (Detroit). We also acknowledge the individuals

responsible for facilitating case ascertainment across the ten sites including: Jennifer Burczyk-

Brown (Alabama); Rana Bayakly and Vicki Bennett (Georgia); the Louisiana Tumor Registry;

Lisa Paddock and Manisha Narang (New Jersey); Diana Slone, Yingli Wolinsky, Steven

Waggoner, Anne Heugel, Nancy Fusco, Kelly Ferguson, Peter Rose, Deb Strater, Taryn Ferber,

Donna White, Lynn Borzi, Eric Jenison, Nairmeen Haller, Debbie Thomas, Vivian von

Gruenigen, Michele McCarroll, Joyce Neading, John Geisler, Stephanie Smiddy, David Cohn,

Michele Vaughan, Luis Vaccarello, Elayna Freese, James Pavelka, Pam Plummer, William

Nahhas, Ellen Cato, John Moroney, Mark Wysong, Tonia Combs, Marci Bowling, Brandon

Fletcher (Ohio); Martin Whiteside (Tennessee) and Georgina Armstrong and the Texas Registry,

Cancer Epidemiology and Surveillance Branch, Department of State Health Services. We would

also like to acknowledge the AACES investigators, Anthony J Alberg, Elisa V Bandera, Jill

Barnholtz-Sloan, Melissa Bondy, Michele L Cote, Ellen Funkhouser, Edward Peters, Ann G

Schwartz, Paul Terry, and Patricia G Moorman. This study would not have been possible without

the efforts of the North Carolina Central Tumor Registry and all of the staff of the NCOCS. We

also thank Christine Lankevich for her management of the data collection for the North Carolina

Study and Rex C. Bentley for review of the pathology in the NCOCS. We would like to thank

Rex C. Bentley and Ann M. Mills for the review of the pathology in the AACES.

References

1. Phung MT, Pearce CL, Meza R, Jeon J. Trends of Ovarian Cancer Incidence by Histotype and Race/Ethnicity in the United States 1992-2019. Cancer Res Commun [Internet]. 2023;3:1–8. Available from: http://dx.doi.org/10.1158/2767-9764.CRC-22-0410

2. Gaitskell K, Hermon C, Barnes I, Pirie K, Floud S, Green J, et al. Ovarian cancer survival by stage, histotype, and pre-diagnostic lifestyle factors, in the prospective UK Million Women Study. Cancer Epidemiol [Internet]. 2022;76:102074. Available from: http://dx.doi.org/10.1016/j.canep.2021.102074

3. Fortner RT, Trewin-Nybråten CB, Paulsen T, Langseth H. Characterization of ovarian cancer survival by histotype and stage: A nationwide study in Norway. Int J Cancer [Internet]. 2023;153:969–78. Available from: http://dx.doi.org/10.1002/ijc.34576

4. Peres LC, Cushing-Haugen KL, Köbel M, Harris HR, Berchuck A, Rossing MA, et al. Invasive Epithelial Ovarian Cancer Survival by Histotype and Disease Stage. J Natl Cancer Inst [Internet]. 2019;111:60–8. Available from: http://dx.doi.org/10.1093/jnci/djy071

5. Cancer Genome Atlas Research Network. Integrated genomic analyses of ovarian carcinoma. Nature [Internet]. 2011;474:609–15. Available from: http://dx.doi.org/10.1038/nature10166

6. Tothill RW, Tinker AV, George J, Brown R, Fox SB, Lade S, et al. Novel molecular subtypes of serous and endometrioid ovarian cancer linked to clinical outcome. Clin Cancer Res [Internet]. 2008;14:5198–208. Available from: http://dx.doi.org/10.1158/1078-0432.CCR-08-0196

7. Way GP, Rudd J, Wang C, Hamidi H, Fridley BL, Konecny GE, et al. Comprehensive Cross-Population Analysis of High-Grade Serous Ovarian Cancer Supports No More Than Three Subtypes [Internet]. G3 Genes|Genomes|Genetics. 2016. page 4097–103. Available from: http://dx.doi.org/10.1534/g3.116.033514

8. Verhaak RGW, Tamayo P, Yang J-Y, Hubbard D, Zhang H, Creighton CJ, et al. Prognostically relevant gene signatures of high-grade serous ovarian carcinoma. J Clin Invest [Internet]. 2013;123:517–25. Available from: http://dx.doi.org/10.1172/JCI65833

9. Konecny GE, Wang C, Hamidi H, Winterhoff B, Kalli KR, Dering J, et al. Prognostic and therapeutic relevance of molecular subtypes in high-grade serous ovarian cancer. J Natl Cancer Inst [Internet]. 2014;106. Available from: http://dx.doi.org/10.1093/jnci/dju249

10. Tan TZ, Miow QH, Huang RY-J, Wong MK, Ye J, Lau JA, et al. Functional genomics identifies five distinct molecular subtypes with clinical relevance and pathways for growth control in epithelial ovarian cancer. EMBO Mol Med [Internet]. 2013;5:1051–66. Available from: http://dx.doi.org/10.1002/emmm.201201823

11. Talhouk A, George J, Wang C, Budden T, Tan TZ, Chiu DS, et al. Development and Validation of the Gene Expression Predictor of High-grade Serous Ovarian Carcinoma Molecular SubTYPE (PrOTYPE). Clin Cancer Res [Internet]. 2020;26:5411–23. Available

from: http://dx.doi.org/10.1158/1078-0432.CCR-20-0103

12. Wang C, Armasu SM, Kalli KR, Maurer MJ, Heinzen EP, Keeney GL, et al. Pooled
Clustering of High-Grade Serous Ovarian Cancer Gene Expression Leads to Novel
Consensus Subtypes Associated with Survival and Surgical Outcomes. Clin Cancer Res
[Internet]. 2017;23:4077–85. Available from: http://dx.doi.org/10.1158/1078-0432.CCR-17-
0246

13. Siegel RL, Miller KD, Fuchs HE, Jemal A. Cancer statistics, 2022. CA Cancer J Clin
[Internet]. 2022;72:7–33. Available from: http://dx.doi.org/10.3322/caac.21708

14. Peres LC, Sinha S, Townsend MK, Fridley BL, Karlan BY, Lutgendorf SK, et al. Predictors
of survival trajectories among women with epithelial ovarian cancer. Gynecol Oncol
[Internet]. 2020;156:459–66. Available from: http://dx.doi.org/10.1016/j.ygyno.2019.12.011

15. Schildkraut JM, Johnson C, Dempsey LF, Qin B, Terry P, Akonde M, et al. Survival of
epithelial ovarian cancer in Black women: a society to cell approach in the African
American cancer epidemiology study (AACES). Cancer Causes Control [Internet].
2023;34:251–65. Available from: http://dx.doi.org/10.1007/s10552-022-01660-0

16. Moorman PG, Schildkraut JM, Calingaert B, Halabi S, Vine MF, Berchuck A. Ovulation
and ovarian cancer: a comparison of two methods for calculating lifetime ovulatory cycles
(United States). Cancer Causes Control [Internet]. 2002;13:807–11. Available from:
http://dx.doi.org/10.1023/a:1020678100977

17. Schildkraut JM, Iversen ES, Akushevich L, Whitaker R, Bentley RC, Berchuck A, et al.
Molecular signatures of epithelial ovarian cancer: analysis of associations with tumor
characteristics and epidemiologic risk factors. Cancer Epidemiol Biomarkers Prev [Internet].
2013;22:1709–21. Available from: http://dx.doi.org/10.1158/1055-9965.EPI-13-0192

18. Peres LC, Colin-Leitzinger C, Sinha S, Marks JR, Conejo-Garcia JR, Alberg AJ, et al.
Racial Differences in the Tumor Immune Landscape and Survival of Women with High-
Grade Serous Ovarian Carcinoma. Cancer Epidemiol Biomarkers Prev [Internet].
2022;31:1006–16. Available from: http://dx.doi.org/10.1158/1055-9965.EPI-21-1334

19. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics [Internet]. 2018;34:i884–90. Available from:
http://dx.doi.org/10.1093/bioinformatics/bty560

20. Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast and bias-
aware quantification of transcript expression. Nat Methods [Internet]. 2017;14:417–9.
Available from: http://dx.doi.org/10.1038/nmeth.4197

21. Ganzfried BF, Riester M, Haibe-Kains B, Risch T, Tyekucheva S, Jazic I, et al.
curatedOvarianData: clinically annotated data for the ovarian cancer transcriptome.
Database [Internet]. 2013;2013:bat013. Available from:
http://dx.doi.org/10.1093/database/bat013

22. Waldron L, Riester M, Ramos M, Parmigiani G, Birrer M. The Doppelgänger Effect:

Hidden Duplicates in Databases of Transcriptome Profiles. J Natl Cancer Inst [Internet].
2016;108. Available from: http://dx.doi.org/10.1093/jnci/djw146

23. Maechler M. cluster: Cluster Analysis Basics and Extensions. R package version 207–1
[Internet]. 2018 [cited 2023 Aug 14]; Available from:
https://cir.nii.ac.jp/crid/1370290617548437512

24. Gaujoux R, Seoighe C. A flexible R package for nonnegative matrix factorization. BMC
Bioinformatics [Internet]. 2010;11:367. Available from: http://dx.doi.org/10.1186/1471-
2105-11-367

25. Tusher VG, Tibshirani R, Chu G. Significance analysis of microarrays applied to the
ionizing radiation response [Internet]. Proceedings of the National Academy of Sciences.
2001. page 5116–21. Available from: http://dx.doi.org/10.1073/pnas.091062498

26. Schwender H, Krause A, Ickstadt K. Identifying interesting genes with siggenes. The
Newsletter of the R Project Volume 6/5, December 2006 [Internet]. Citeseer; 2006;6:45.
Available from: http://mirror.psu.ac.th/pub/cran/doc/Rnews/Rnews_2006-5.pdf#page=45

27. Chen GM, Kannan L, Geistlinger L, Kofia V, Safikhani Z, Gendoo DMA, et al. Consensus
on Molecular Subtypes of High-Grade Serous Ovarian Carcinoma. Clin Cancer Res
[Internet]. 2018;24:5037–47. Available from: http://dx.doi.org/10.1158/1078-0432.CCR-18-
0784

28. Yoshihara K, Tsunoda T, Shigemizu D, Fujiwara H, Hatae M, Fujiwara H, et al. High-risk
ovarian cancer based on 126-gene expression signature is uniquely characterized by
downregulation of antigen presentation pathway. Clin Cancer Res [Internet]. 2012;18:1374–
85. Available from: http://dx.doi.org/10.1158/1078-0432.CCR-11-2725

29. Helland Å, Anglesio MS, George J, Cowin PA, Johnstone CN, House CM, et al.
Deregulation of MYCN, LIN28B and LET7 in a molecular subtype of aggressive high-grade
serous ovarian cancers. PLoS One [Internet]. 2011;6:e18064. Available from:
http://dx.doi.org/10.1371/journal.pone.0018064

30. Peng L, Bian XW, Li DK, Xu C, Wang GM, Xia QY, et al. Large-scale RNA-Seq
Transcriptome Analysis of 4043 Cancers and 548 Normal Tissue Controls across 12 TCGA
Cancer Types. Sci Rep [Internet]. 2015;5:13413. Available from:
http://dx.doi.org/10.1038/srep13413

31. Bonome T, Levine DA, Shih J, Randonovich M, Pise-Masison CA, Bogomolniy F, et al. A
gene signature predicting for survival in suboptimally debulked patients with ovarian cancer.
Cancer Res [Internet]. 2008;68:5478–86. Available from: http://dx.doi.org/10.1158/0008-
5472.CAN-07-6595

32. Marioni JC, Mason CE, Mane SM, Stephens M, Gilad Y. RNA-seq: an assessment of
technical reproducibility and comparison with gene expression arrays. Genome Res
[Internet]. 2008;18:1509–17. Available from: http://dx.doi.org/10.1101/gr.079558.108

33. Robinson MD, Smyth GK. Moderated statistical tests for assessing differences in tag

621  abundance. Bioinformatics [Internet]. Oxford Academic; 2007 [cited 2023 Aug
622  14];23:2881–7. Available from: https://academic.oup.com/bioinformatics/article-
623  abstract/23/21/2881/372869

624 34. Smyth GK. Linear models and empirical bayes methods for assessing differential expression
625  in microarray experiments. Stat Appl Genet Mol Biol [Internet]. 2004;3:Article3. Available
626  from: http://dx.doi.org/10.2202/1544-6115.1027

627 35. Lönnstedt I, Speed T. REPLICATED MICROARRAY DATA. Stat Sin [Internet]. Institute
628  of Statistical Science, Academia Sinica; 2002;12:31–46. Available from:
629  http://www.jstor.org/stable/24307034

630 36. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential
631  expression analysis of digital gene expression data. Bioinformatics [Internet]. 2010;26:139–
632  40. Available from: http://dx.doi.org/10.1093/bioinformatics/btp616

633 37. Anders S, Huber W. Differential expression analysis for sequence count data. Genome Biol
634  [Internet]. 2010;11:R106. Available from: http://dx.doi.org/10.1186/gb-2010-11-10-r106

Table 1. Characteristics of the Schildkraut study populations, by subtype

	Mesenchymal	Proliferative	Immunoreactive	Differentiated	p-value	Mesenchymal	Proliferative	Immunoreactive	Differentiated	p-value
			Schildkraut Black					Schildkraut White		
Case count	79	63	101	19		106	40	84	79	
Age (%)										
Under 40	1 (1.3)	2 (3.2)	2 (2.0)	1 (5.3)		4 (3.8)	0 (0.0)	1 (1.2)	2 (2.5)	
40-44	8 (10.1)	2 (3.2)	6 (5.9)	3 (15.8)		2 (1.9)	1 (2.5)	7 (8.3)	6 (7.6)	
45-49	9 (11.4)	5 (7.9)	14 (13.9)	2 (10.5)		6 (5.7)	5 (12.5)	10 (11.9)	9 (11.4)	
50-54	17 (21.5)	8 (12.7)	13 (12.9)	3 (15.8)		17 (16.0)	2 (5.0)	17 (20.2)	13 (16.5)	
55-59	8 (10.1)	16 (25.4)	25 (24.8)	4 (21.1)		16 (15.1)	9 (22.5)	17 (20.2)	16 (20.3)	
60-64	16 (20.3)	9 (14.3)	16 (15.8)	1 (5.3)		20 (18.9)	13 (32.5)	17 (20.2)	13 (16.5)	
65-69	6 (7.6)	9 (14.3)	15 (14.9)	2 (10.5)		24 (22.6)	1 (2.5)	9 (10.7)	8 (10.1)	
70-74	12 (15.2)	8 (12.7)	7 (6.9)	1 (5.3)		17 (16.0)	9 (22.5)	6 (7.1)	12 (15.2)	
75+	2 (2.5)	4 (6.3)	3 (3.0)	2 (10.5)	0.30	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0.024
FIGO Stage (%)										
I	5 (6.3)	4 (6.3)	13 (12.9)	7 (36.8)		1 (0.9)	1 (2.5)	6 (7.1)	10 (12.7)	
II	8 (10.1)	6 (9.5)	7 (6.9)	3 (15.8)		5 (4.7)	1 (2.5)	2 (2.4)	4 (5.1)	
III	60 (75.9)	49 (77.8)	74 (73.3)	8 (42.1)		97 (91.5)	36 (90.0)	73 (86.9)	59 (74.7)	
IV	4 (5.1)	3 (4.8)	6 (5.9)	0 (0.0)		3 (2.8)	2 (5.0)	3 (3.6)	2 (2.5)	
Missing	2 (2.5)	1 (1.6)	1 (1.0)	1 (5.3)	0.040	0 (0.0)	0 (0.0)	0 (0.0)	4 (5.1)	0.009
Debulking (%)										
Optimal	33 (41.8)	24 (38.1)	45 (44.6)	8 (42.1)		28 (26.4)	14 (35.0)	23 (27.4)	26 (32.9)	
Suboptimal	25 (31.6)	15 (23.8)	21 (20.8)	2 (10.5)		8 (7.5)	3 (7.5)	2 (2.4)	4 (5.1)	
Missing	21 (26.6)	24 (38.1)	35 (34.7)	9 (47.4)	0.34	70 (66.0)	23 (57.5)	59 (70.2)	49 (62.0)	0.59
Neoadjuvant chemotherapy (%)										
No	68 (86.1)	54 (85.7)	85 (84.2)	17 (89.5)		106 (100.0)	40 (100.0)	84 (100.0)	79 (100.0)	NA
Missing	11 (13.9)	9 (14.3)	16 (15.8)	2 (10.5)	0.94					
Proportion CD3+ T cells (%)	4.24 (4.68)	1.51 (3.18)	3.88 (4.54)	1.04 (1.24)	0.001	2.46 (2.36)	0.72 (1.25)	4.35 (6.55)	2.24 (2.20)	0.053
Proportion CD3+/CD8+ Suppressor T cells (%)	1.38 (2.30)	0.52 (1.80)	1.42 (2.33)	0.32 (0.53)	0.045	1.15 (1.18)	0.22 (0.30)	2.39 (4.48)	1.16 (1.21)	0.084

Table 2. Subtype distribution and risk of death by study population

Study and molecular subtype	Case count (%)	Model 1 HR[a] (95% CI)	Model 2 HR[b] (95% CI)
Schildkraut Black			
Mesenchymal	79 (30.2)	(ref)	(ref)
Proliferative	63 (24.0)	1.13 (0.76, 1.67)	0.98 (0.66, 1.46)
Immunoreactive	101 (38.5)	0.83 (0.59, 1.19)	0.79 (0.55, 1.13)
Differentiated	19 (7.3)	0.87 (0.42, 1.83)	0.83 (0.40, 1.72)
Schildkraut White			
Mesenchymal	106 (34.3)	(ref)	(ref)
Proliferative	40 (12.9)	1.00 (0.67, 1.49)	1.05 (0.70, 1.57)
Immunoreactive	84 (27.2)	0.85 (0.61, 1.18)	0.86 (0.62, 1.19)
Differentiated	79 (25.6)	1.01 (0.73, 1.40)	0.98 (0.70, 1.36)
TCGA			
Mesenchymal	125 (25.1)	(ref)	(ref)
Proliferative	102 (20.4)	0.82 (0.56, 1.20)	0.84 (0.57, 1.23)
Immunoreactive	116 (23.2)	0.92 (0.63, 1.35)	1.01 (0.69, 1.49)
Differentiated	156 (31.3)	0.93 (0.67, 1.28)	1.03 (0.74, 1.44)
Mayo			
Mesenchymal	105 (27.9)	(ref)	(ref)
Proliferative	79 (21.0)	0.81 (0.59, 1.12)	0.92 (0.66, 1.28)
Immunoreactive	93 (24.7)	0.54 (0.39, 0.75)	0.54 (0.39, 0.75)
Differentiated	100 (26.5)	0.63 (0.46, 0.87)	0.61 (0.44, 0.84)
Yoshihara[c]			
Mesenchymal	89 (34.9)	(ref)	(ref)
Proliferative	30 (11.8)	1.35 (0.77, 2.35)	1.48 (0.86, 2.57)
Immunoreactive	71 (27.8)	0.62 (0.37, 1.04)	0.65 (0.39, 1.09)
Differentiated	65 (25.5)	1.16 (0.73, 1.85)	1.36 (0.85, 2.19)
Tothill			
Mesenchymal	75 (31.1)	(ref)	(ref)
Proliferative	45 (18.7)	0.77 (0.43, 1.37)	0.79 (0.44, 1.42)
Immunoreactive	67 (27.8)	0.55 (0.32, 0.95)	0.54 (0.31, 0.95)
Differentiated	54 (22.4)	0.64 (0.37, 1.12)	0.65 (0.37, 1.14)

[a] Multivariable model adjusted for age (in 5-year age groups) and stage (I, II, III, IV)

[b] Multivariable model adjusted for age (in 5-year age groups), stage (I, II, III, IV), and debulking status (optimal, not optimal)

[c] Yoshihara dataset does not include information on age, so multivariable models are not adjusted for age in this dataset

641
Figure 1. Principal Components Analysis (PCA) plots of cluster centroids for each dataset, after
dataset alignment. **Panel a** compares K-means cluster centroids (K=3), and **Panel b** compares K-
means cluster centroids (K=4) for each dataset considered in this study. We find that the
principal components separate each cluster centroid in a consistent way across almost all
datasets. For K=3, PC2 and PC3 are both able to separate each cluster independently, but for
K=4 the combination of PC2 and PC3 are needed to separate each cluster. Furthermore, for K=3,
we see that the Yoshihara and Mayo datasets have centroids that are much higher in PC1 than the
other datasets. This trend continues for the Mayo dataset when K=4, in all PCs.
650

Figure 2. Significance analysis of microarray (SAM) moderated t-score Pearson correlation heatmaps of clusters across datasets. **Panel a** compares K-means clusters (K=3) between SchildkrautB and every other dataset considered in this study. Across each dataset we find a strong positive correlation with the clusters in SchildkrautB, with matched cluster correlations ranging from 0.37-0.86, and mismatched cluster correlations ranging from -0.65-0.03. **Panel b** performs the same comparison, but for K=4. In this comparison, we see much more inconsistency between matched clusters, with some mismatched clusters having a higher correlation than some matched clusters.

661
662   **Figure 3.** Distribution of subtypes across datasets.
663

664
665 **Figure 4.** Kaplan Meier survival curves comparing subtype-specific survival by dataset.
666