OVO Positively Regulates Essential Maternal Pathways by Binding Near the Transcriptional Start Sites in the *Drosophila* Female Germline

Leif Benner[1,2], Savannah Muron[1], Jillian G. Gomez[1], Brian Oliver[1]

1. Section of Developmental Genomics, Laboratory of Biochemistry and Genetics, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, MD, USA.

2. Department of Biology, Johns Hopkins University, Baltimore, MD, USA.

Abstract

Differentiation of female germline stem cells into a mature oocyte includes the expression of RNAs and proteins that drive early embryonic development in *Drosophila*. We have little insight into what activates the expression of these maternal factors. One candidate is the zinc-finger protein OVO. OVO is required for female germline viability and has been shown to positively regulate its own expression, as well as a downstream target, *ovarian tumor*, by binding to the transcriptional start site (TSS). To find additional OVO targets in the female germline and further elucidate OVO's role in oocyte development, we performed ChIP-seq to determine genome-wide OVO occupancy, as well as RNA-seq comparing hypomorphic and wild type rescue *ovo* alleles. OVO preferentially binds in close proximity to target TSSs genome-wide, is associated with open chromatin, transcriptionally active histone marks, and OVO-dependent expression. Motif enrichment analysis on OVO ChIP peaks identified a 5`-TAACNGT-3` OVO DNA binding motif spatially enriched near TSSs. However, the OVO DNA binding motif does not exhibit precise motif spacing relative to the TSS characteristic of RNA Polymerase II complex binding core promoter elements. Integrated genomics analysis showed that 525 genes that are bound and increase in expression downstream of OVO are known to be essential maternally expressed genes. These include genes involved in anterior/posterior/germ plasm specification (*bcd, exu, swa, osk, nos, aub, pgc, gcl*), egg activation (*png, plu, gnu, wisp, C(3)g, mtrm*), translational regulation (*cup*, *orb*, *bru1, me31B*), and vitelline membrane formation (*fs(1)N*, *fs(1)M3*, *clos*). This suggests that OVO is a master transcriptional regulator of oocyte development and is responsible for the expression of structural components of the egg as well as maternally provided RNAs that are required for early embryonic development.

Introduction

Drosophila early embryonic development is directed by events that take place during oogenesis. Germline stem cells (GSCs) asymmetrically divide to renew the stem cell population and send one daughter cell towards oogenesis. In the germarium, oogenesis is preceded by four rounds of incomplete mitotic divisions resulting in a 16-cell egg chamber. One cell is specified as the oocyte which is arrested in prophase of meiosis I, while the rest of the 15 cells enter endoreplication cycles and become nurse cells (NCs). Once the 16 cell egg chamber buds from the germarium, the NCs begin to transcribe and translate a vast array of RNAs and proteins that serve diverse functional roles (Bastock and St Johnston 2008; Spradling et al. 2022). These roles include positioning maternal mRNAs and proteins in the correct spatial orientation to support anterior/posterior, and dorsal ventral axis specification, as well negative regulators of translation to ensure that the maternal mRNAs are not translated before fertilization (Lasko 2012). The oocyte also contains a number of proteins and mRNAs that are needed for egg activation, completion of meiosis and initiation of embryonic development after fertilization (Avilés-Pagán and Orr-Weaver 2018). To repeat the process of oogenesis from generation to generation, germ cells in the developing embryo need to be specified and maintained separately from the rest of the developing somatic cell population. This requires maternally controlled localization of the germ plasm and early pole cell formation in the embryo (Mahowald 2001). While the complex interactions between maternally supplied mRNAs and proteins have been well studied, transcriptional regulation driving the expression of these pathways are less well understood.

Few positive regulators of female specific germ cell transcription have been identified. Genes such as *grauzone* (*grau*) and *maternal gene required for meiosis* (*mamo*) have been shown to activate the transcription of *cortex* and *vasa*, respectively, in the female germline (Harms et al. 2000; S. Nakamura et al. 2019). Active repression of male specific transcription through the activity of *egg*, *wde*, and *Su(var)205* (Smolko, Shapiro-Kulnane, and Salz 2018), or global repression of non-ovarian transcriptional networks

through the function of *sov* (Benner et al. 2019), have shown the importance of heterochromatin formation in the female germline for cellular identity and oocyte development. In fact, recent work has shown the importance of transcriptional repression, mediated through changes in histone modifications, as a key regulator of egg chamber differentiation. GSCs have been shown to exist in a sort of 'ground state' of histone modifications. Characterized with modest non-canonical repressive H3K9me3 and H3K27me3 histone marks at many genes, as well as transcriptionally active H3K27ac histone marks and open chromatin at others (Pang et al. 2023; DeLuca et al. 2020). As oocyte development continues, repressive histone marks associated with heterochromatin begin to increase in abundance resulting in fewer histone marks associated with transcription and open chromatin. This suggests that gene expression becomes more restricted throughout the differentiation process. However, it is unlikely that the female germline directs oocyte development solely through a repressive transcriptional model. Whether the female germline expresses paralogs of the RNA polymerase II complex, like the male germline (M. Hiller et al. 2004; M. A. Hiller et al. 2001; Lu et al. 2020), or if there are pioneering transcription factors involved in determining the open chromatin status for female germ cell-specific expression, or something else entirely, has yet to be determined.

Although few female specific germline transcription factors have been identified, the conserved zinc-finger transcription factor OVO has long been known to be required for female germ cell viability. Female germ cells that are *ovo⁻* do not survive into adulthood (Oliver, Perrimon, and Mahowald 1987; Oliver, Pauli, and Mahowald 1990; Benner et al. 2023). Hypomorphic *ovo* alleles, specifically ones that disrupt the transcriptional activator OVO-B, show an arrested egg chamber phenotype, indicating that wild-type OVO-B activity is required for oocyte maturation (Salles et al. 2002; Benner et al. 2023). Germline OVO is expressed at all stages of oogenesis, where it is eventually maternally loaded into the egg. Maternal OVO becomes specifically localized to the developing germline and persists throughout embryogenesis until zygotic OVO is expressed (Hayashi et al. 2017; Benner et al. 2023). Thus, OVO is eternally expressed in the female germline, suggesting it may be a key regulator of female specific germline transcription. However, only two downstream targets have previously been identified for OVO. OVO has been shown to positively regulate the expression of its own transcription, therefore executing an autoregulatory loop, as well as positively regulating the transcription of the gene *ovarian tumor* (*otu*)(Lü et al. 1998; Bielinska et al. 2005; Lü and Oliver 2001; Andrews et al. 2000). *otu* is also required in the female germline, where *otu⁻* germ cells show viability and germline tumor phenotypes (Bishop and King 1984). The *ovo* phenotype is epistatic to that of *otu*, and ectopic *otu* expression cannot rescue female germ cell death due to loss of *ovo* (Hinson, Pettus, and Nagoshi 1999; Pauli, Oliver, and Mahowald 1993). Therefore, *ovo* must be responsible for activating the transcription of genes in addition to *otu* for female germ cell survival and differentiation.

We expanded our knowledge of OVO's role in the female germline by determining genome-wide OVO occupancy and global transcriptional changes downstream of OVO. This allowed us to determine which genes OVO binds, and which genes transcriptionally respond to OVO *in vivo*. We show that OVO is directly regulating essential maternal pathways such as axis specification, primordial germ cell formation, egg activation, and maternal mRNA translation regulation. Together, we show that OVO plays a pivotal role in the positive transcriptional regulation of oocyte and early embryonic development. We show that OVO likely carries out this regulation by binding at or in close proximity to the promoters of the genes it regulates and that OVO DNA binding motifs are enriched at or near the transcriptional start site (TSS) of OVO responsive genes, although the spacing of OVO binding sites suggests that it is not a component of the RNA Polymerase complex. OVO binding is also a signature of open chromatin status and active transcription throughout oocyte differentiation. Altogether, we suggest that OVO is required for the activity of a large number of female germline promoters and is likely a key regulator of oocyte maturation and RNAs and proteins that are required for early embryonic development.

Results

OVO Binds Promoters Genome-Wide

OVO-B, the predominant protein isoform expressed from the *ovo* locus in the female germline (Benner et al. 2023), is a positive regulator of transcription (Andrews et al. 2000) at both the *otu* and *ovo* locus. Transgenic reporter constructs of *otu* and *ovo* require OVO binding sites both at and upstream of the TSS in order to recapitulate full reporter expression (Bielinska et al. 2005; Lü et al. 1998; Lü and Oliver 2001).

Females hemizygous for antimorphic dominant gain-of-function (*ovo^D*) or homozygous recessive (*ovo^D1rv*)
ovo alleles lack germ cells in the adult ovary (Oliver, Perrimon, and Mahowald 1987; Benner et al. 2023).
True OVO-B null alleles created by deletion of the *ovo-B* promoter have the same germ cell loss
phenotype (Benner et al. 2023). The phenotypes of *otu* females range from germ cell death to ovarian
tumors depending on the allele and undefined stochastic factors (Bishop and King 1984). It is possible
that the germ cell death phenotype in *ovo^−* female germ cells can solely be explained by failure of OVO to
activate *otu* expression, however, this is highly unlikely. The *ovo^D1rv* phenotype is epistatic to *otu^−*, and
ectopic *otu^+* expression in *ovo^−* germ cells does not rescue the germ cell death phenotype (Hinson,
Pettus, and Nagoshi 1999). This suggests that OVO regulates the expression of additional genes in the
female germline.
123
We wanted to identify OVO target genes in the female germline by using two complementary genome-
wide approaches to test for OVO presence and function. Specifically, we determined OVO occupancy
genome-wide with ChIP-seq, and determined *ovo* function by comparing the RNA expression profiles
between *ovo^+* and *ovo* hypomorphs in the female germline. In order to determine OVO occupancy
genome-wide, we performed ChIP-seq on adult ovaries in triplicate, using two C-terminally tagged alleles
as affinity purification tools (*ovo^Cterm-3xFHA* i.e. OVO-HA and *ovo^Cterm-GFP* i.e. OVO-GFP, Figure 1A-C, Figure
S1A) (Benner et al. 2023) and called significantly enriched peaks from OVO-HA and OVO-GFP compared
to their respective input controls.
132
We first compared the pulldown results with the OVO-HA versus OVO-GFP ChIP reagents. The GFP pull
down appeared to be more efficient, but never-the-less we found excellent agreement, as most OVO-HA
peaks were also found in the OVO-GFP dataset. The OVO-GFP ChIP dataset had 7,235 significant ChIP
peaks according to peak enrichment analysis genome-wide, while OVO-HA ChIP dataset had 3,393
significant peaks genome-wide (Table S1). To determine the similarity in significant peak calling between
the two datasets, we calculated a Jaccard index (intersection/union) between the significantly enriched
peaks from the tagged *ovo* allele bearing ovaries. The Jaccard index between OVO-HA and OVO-GFP
ChIP peaks was 0.64 (where 0 = no overlap and 1 = full overlap) with a total of 3,094 ChIP peaks
overlapping. Thus, almost all significant OVO-HA ChIP peaks were also found within the OVO-GFP ChIP
dataset (91% of OVO-HA peaks overlapped OVO-GFP peaks). OVO-GFP pulldown was either more
effective, or less likely, promiscuous. Regardless, we decided to use the conservative intersection of the
two datasets (3,094 peaks) for downstream OVO occupancy informatics (Table S2).
145
OVO is a sequence-specific DNA-binding protein, but many transcription factors also have a more
general affinity for DNA. Additionally, immunoprecipitation can capture indirect interactions due to nuclear
topology in addition to direct sequence-specific binding. For example, in the particular case of OVO TSS
binding, this could be direct, as shown in the case of *ovo* and *otu* loci (Lü et al. 1998; Lü and Oliver 2001;
Bielinska et al. 2005), or could be due to looping of an OVO-bound enhancer to the core promoter.
Determining if there are canonical OVO binding sites at peaks can help to distinguish direct and indirect
binding. If OVO is directly binding to the TSS, we would expect to find OVO binding site enrichment at
that location. To examine the sequences enriched in peaks, we looked directly for the known OVO
binding sites previously defined by footprinting (Lü et al. 1998) and SELEX-seq (Bielinska et al. 2005; Lee
and Garfinkel 2000). We also did *de novo* motif finding on this substantial dataset to refine the sequence
specific motifs bound by OVO and perhaps other motifs associated with other transcription factors
preferentially bound by OVO enhancers, or OVO proximal sites in 3D nuclear space. We performed novel
motif enrichment analysis using STREME (Bailey 2021) with our overlapping ChIP peaks and found a
number of significant motifs within our dataset. The most significant motif, 5`-TAACGGTAAA-3`, was
found in 49% of significant OVO ChIP peaks (Figure 1D, 'Motif One'). This motif is highly similar to the
OVO DNA binding motif that has been reported twice before, 5`-AGTAACNGT-3` (SELEX method,
'Garfinkel OVO Motif') and 5`-TGTAACNGT-3` (Footprinting method, 'Oliver OVO Motif'). The only
differences between motif one in our dataset and the literature, is that the *de novo* motif is two
nucleotides shorter than the previously described motifs at the 5` end, extends 3 nucleotides
downstream, and includes a second G near the 3'-end (Figure S1B). Collectively, this is strong evidence
that the core OVO binding sequence is 5`-TAACNGT-3`. Some binding sites can be recognized by
multiple transcription factors. To determine if other characterized transcription factors recognize this
sequence, we searched for significant matches to motif one in the Jaspar database (Castro-Mondragon et

al. 2022) of known *Drosophila* motifs using Tomtom (Gupta et al. 2007). The OVO DNA binding motif
('Garfinkel OVO Motif') was scored as a significant match (Figure S1B, $p<0.05$). Along with the OVO DNA
binding motif, other motifs were also significantly enriched in OVO ChIP peaks. The motif 5'-
GWGMGAGMGAGABRG-3' (Figure S1C) was found in 18% of OVO ChIP peaks and is a significant
match to the DNA binding motifs of the transcription factors GAF (*Trl*) (Omelina et al. 2011) and CLAMP
(Soruco et al. 2013). *Trl* germline clones are not viable, indicating that GAF activity is required in the
germline during oogenesis (Chen et al. 2009). The possibility that OVO binds with and regulates genes
alongside of GAF given the enrichment of both transcription factors DNA binding motifs is intriguing.
Other significantly enriched motifs 5'-ACACACACACACACA-3' (29% of peaks, Figure S1D), 5'-
RCAACAACAACAACA-3' (26% of peaks, Figure S1E), and 5'-GAAGAAGAAGAAGAR-3' (17% of peaks,
Figure S1F) were present in OVO ChIP peaks, however, these motifs did not significantly match known
DNA binding motifs of other transcription factors. Determining the factors that bind to these sequences
will certainly help elucidate our understanding of transcriptional control with relationship to OVO in the
female germline.
183
Not every peak region had one of the consensus OVO DNA binding motifs. This does not mean *a priori*
that they bound OVO indirectly. Motif enrichment can be driven by a few strongly enriched sequences, so
that more minor enrichments of variants are missed. Therefore, we carefully examined the 51% of our
overlapping OVO ChIP peaks where OVO DNA binding motif one was not found. This second round of *de*
novo analysis revealed enrichment of OVO DNA binding motif derivatives. For example, the third most
significant motif (found in 37% of peaks) was 5`-RWMTAACGGV-3` (Figure 1E, motif two). This motif had
the core 5`-TAACNGT-3` sequence found in all three aforementioned methods, however, the last
nucleotide in the core sequence is unspecific and lacks the three 3` nucleotides found in motif one. Two
other derivative motifs, 5'-TAACTGTTTT-3' (found in 17% of sequences, Figure 1F, motif three), and 5'-
TTACSGTAA-3' (found in 5% of sequences, Figure 1G, motif four), vary within the central core motif (at
positions 5 and 2 of the core sequence, respectively) and upstream and downstream ends. Searching for
all variations of the OVO DNA binding motif (Table S3) within our significant overlapping ChIP peaks
indicated that 72% of peaks contained at least one variation of these four binding motifs. It is a
reasonable hypothesis that OVO peaks are most often due to direct, rather than indirect OVO binding.
198
A prediction for direct OVO binding to motifs, is that the motif should be centered within the peak of
fragmented input DNA sequences. Therefore, we plotted the significant ChIP (minus input) read density
centered on the location of the motif. We found that the read density for all ChIP peaks that contain each
one of the *de novo* OVO motifs, as well as the *in vitro* OVO motifs, are centered over the motif location
(Figure 1H). This suggests that all of these motifs from our analysis are bonafide OVO DNA binding sites
in vivo. While it is possible that OVO comes into contact with regions of DNA in three-dimensional nuclear
space non-specifically, the presence of OVO motifs within a large percentage of significant ChIP peaks *in*
vivo and enrichment of OVO ChIP read density at the location of the motifs, strongly reinforces the idea
that our OVO ChIP dataset contains regions centered on sequences specifically bound by OVO in the
ovary.
209
Given the clear function of OVO occupancy near the TSS of its two known targets, *ovo* and *otu* (Lü et al.
1998; Bielinska et al. 2005; Lü and Oliver 2001), we were interested in determining if OVO peaks are
generally near the TSS of other target genes as well. In addition to informing the biology of OVO function,
this simplifies the problem of associating peaks to potential functional target genes. As a preliminary test
of this idea, we determined if the fully overlapping OVO-HA and OVO-GFP peaks were spatially enriched
with respect to the currently annotated gene model elements such as TSS, open reading frames (ORFs),
or transcription termination sequences (TTS). If the TSS association of OVO at the two known targets
reflects a general propensity, then we expect OVO ChIP peaks to be more closely associated with TSS
than other gene elements. To carry out this analysis, we normalized the genome for these three gene
elements, such that the distance between adjacent loci was 1. If there is no enrichment for OVO peaks to
a specific gene element, then the peak location would have an equal frequency from 0.0 to 0.5 relative
distance. Measuring the relative distance of our OVO ChIP peaks to TSS, ORFs, and TTS, showed that
the OVO binding was highly enriched near TSS/promoter locations and was not correlated with ORF and
TTS locations (Figure 1I). These results confirmed that OVO is characterized by core promoter proximal
binding. Since OVO ChIP peaks as a class are associated with TSS, we plotted the ChIP minus input

read density of genes that overlap significant ChIP peaks to examine the full distribution. We found that
the OVO ChIP read density was highly enriched over the TSS and was not due to a few highly enriched
examples (Figure 1J). This builds on the idea that OVO is binding directly over, or in close proximity to the
TSS of its target genes genome-wide. In other words, the previous work showing OVO binding the *ovo-B*
and *otu* TSS (Bielinska et al. 2005; Lü et al. 1998; Lü and Oliver 2001) is typical. This very specific
binding of OVO to the TSS is intriguing and unusual, as this region associates with the basal
transcriptional machinery. It raises the possibility that OVO is not a typical transcription factor that acts
primarily via enhancer binding but might be part of the core promoter binding complex or acts to
precondition the core promoter region for example.
234
Although OVO ChIP peaks overlapping genes showed a strong read density enrichment over the TSS,
we found that only 45% (1,394/3,094) of OVO ChIP peaks directly overlapped a TSS. 43% (1,339/3,094)
of OVO ChIP peaks were found to overlap the gene body downstream of the TSS (intronic and exonic
sequences) and 12% (366/3,094) did not overlap any gene elements, indicating that they were intergenic.
We were interested in the differences between OVO binding directly over the TSS or at more distal
upstream and downstream sites. We decided to plot the OVO ChIP read density of these different classes
of OVO binding patterns and found that OVO bound over the TSS produced a sharp read density
enrichment over the TSS which was consistent with what was found for all OVO bound genes (Figure
1K). OVO binding along the gene body surprisingly also showed a read density enrichment over the TSS,
although the magnitude of read density enrichment was notably less than TSS OVO binding. Intergenic
OVO binding also showed these same characteristics with a notable upstream read density enrichment
possibly indicative of enhancer binding. This indicates that although the significantly called OVO ChIP
peaks did not overlap the TSS, there was still a propensity for TSS sequences to be enriched with OVO
ChIP over the input control. This could be due to weaker direct *in vivo* binding of OVO to these TSSs or
indirect interactions between the upstream/downstream OVO bound sequences and the TSS, possibly
through a looping enhancer-promoter interaction. However, regardless of the location of the OVO ChIP
peak, OVO seemed to always be enriched at or in close proximity to TSSs.
252
The OVO ChIP read density was highly enriched over the annotated TSS of target genes, but TSS
annotation is challenging and can be tissue specific. We were interested in empirically determining if the
same enrichment was present in TSSs utilized specifically in ovarian tissue. The 5' ends of mRNA are
capped. In order to determine where these caps mapped to the genome, we analyzed Cap Analysis of
Gene Expression (CAGE-seq) data from adult *Drosophila* ovaries (Chen et al. 2014) and extracted the
dominant significant TSSs in the ovary. CAGEr predicted 6,856 significant TSSs in the ovary dataset, of
which 1,047 overlapped with OVO ChIP peaks. We plotted the OVO ChIP minus input read density
centered on the significant ovary CAGE-seq TSSs for TSSs that overlapped or did not overlap OVO ChIP
peaks (Figure 1L). We found that OVO ChIP read density was highly enriched over the location of TSSs
from ovary CAGE-seq that overlapped OVO ChIP peaks when compared to TSSs that did not overlap
OVO ChIP peaks. Thus, OVO TSS binding is not due to poor annotation of ovarian TTSs. Furthermore,
OVO is binding at or near TSSs of genes actively being transcribed in the ovary.
265
<u>OVO Binding is Associated with Open Chromatin and Transcriptionally Active Histone Marks</u>
Our OVO ChIP data indicated that OVO was binding at or in close proximity to promoters genome-wide.
OVO could have a positive and/or negative effect on transcription at these locations. For example, OVO
could help recruit or sterically hinder RNA Polymerase binding to TSSs. However, previous *ovo* reporter
constructs show positive effects of OVO binding near TSS (Lü and Oliver 2001; Lü et al. 1998; Bielinska
et al. 2005). If OVO binding is generally promoting transcription, then we hypothesize that it would be
more closely associated with histone marks associated with active transcription, such as H3K27ac and
H3K4me3, as well as lower nucleosome density that can be measured through ATAC-seq. In contrast,
OVO binding would be expected to negatively correlate with repressive H3K9me3 and H3K27me3
histone marks and higher nucleosome density. It is technically difficult to determine changes in chromatin
status and transcription in germ cells that lack OVO, as the phenotype is cell death (although we will
return to this later for transcription profiling) but analyzing OVO binding in the context of ovarian
chromatin was highly informative.
279

Recent work profiling nucleosome density and histone marks have shown that female GSCs have a
'ground state' chromatin profile (DeLuca et al. 2020), similar to the histone mark profiles that are found in
early embryos (Li et al. 2014). This has been characterized to contain non-canonical H3K27me3 profiles
and low H3K9me3 histone levels (Pang et al. 2023; DeLuca et al. 2020). As egg follicles differentiate,
nurse cells begin to accumulate H3K9me3 marks, and H3K27me3 histone marks begin to accumulate
over more traditional polycomb domains. This in turn leads to a decrease in the number of open
chromatin peaks as well as H3K27ac histone marks, which are generally associated with active
transcription (Pang et al. 2023; DeLuca et al. 2020). Essentially, these data support the idea that egg
chambers restrict gene expression competency as they differentiate.
289
In order to determine the relationship in our OVO ChIP data and other chromatin marks, we analyzed
GSC H3K27ac, H3K27me3, H3K9me3, H3K4me3, and ATAC-seq data (Pang et al. 2023; DeLuca et al.
2020) with the same parameters used to establish significant OVO peaks in our OVO ChIP dataset. Our
OVO ChIP data was from one day old ovaries, and we did not profile specific follicle stages. So we also
analyzed 32c (roughly stage 5 egg chambers) H3K27ac, H3K27me3, ATAC-seq, and 8c H3K9me3 (32c
was not available) histone marks (Pang et al. 2023; DeLuca et al. 2020) to see if there were any stage
specific differences in comparison to OVO DNA binding. We focused specifically on GSC and 32c egg
follicles for these chromatin marks since that is when the *ovo* hypomorphic egg chambers arrest (Benner
et al. 2023). We first plotted the read density of each respective chromatin mark minus their input control
centered on either the OVO ChIP peak local maximum (Figure 2A) or OVO DNA binding motifs contained
within significant OVO ChIP peaks (Figure 2B). GSC ATAC and H3K27ac read density showed a high
degree of enrichment over OVO ChIP peak maximums (Figure 2A) and OVO DNA binding motifs (Figure
2B), consistent with positive transcriptional activity. GSC H3K4me3 read density was, to a lesser extent,
also enriched with OVO ChIP peak maximums and OVO DNA binding motifs. However, there was no
read density enrichment for repressive GSC H3K27me3 and H3K9me3 histone marks. The same findings
generally held true when looking at the overlap of OVO ChIP peaks and chromatin marks in differentiating
egg chambers. Notably, there was an even higher read density enrichment over OVO ChIP peak
maximum and DNA binding motifs for 32c ATAC-seq data, while read density enrichment decreased for
32C H3K27ac histone mark (Figure 2A, B).
309
Since there was a high degree of read density enrichment between OVO ChIP and other chromatin
marks/low nucleosome density we wanted to determine the extent of the overlap between significant
OVO ChIP peaks and significantly called peaks from the different types and stages of histone marks and
ATAC-seq data. To do this, we measured the relative distance of OVO ChIP peaks to the same datasets
described above. We found that OVO ChIP peaks had a lower relative distance, and thus were spatially
overlapping/closer in the genome, to 32c NC ATAC, GSC ATAC, GSC H3K27ac, GSC H3K4me3, and
32c NC H3K27ac peaks, in that order (Figure 2C). While the relative distance between OVO ChIP peaks
and H3K9me3 and H3K27me3, regardless of stage, showed no spatial association. There was also
further support for this association with transcriptionally active histone marks/open chromatin when
measuring the overlap between significant OVO ChIP peaks and the respective significant histone ChIP
and ATAC-seq peaks (Figure 2D). A Fisher's exact test indicated a significant enrichment in overlapping
peaks genome-wide between OVO and GSC ATAC ($p < 0.001$, odds ratio = 75.9), 32c NC ATAC ($p <
0.001$, odds ratio = Infinite), GSC H3K27ac ($p < 0.001$, odds ratio = 31.7), GSC H3K4me3 ($p < 0.001$,
odds ratio =12.0), and 32c NC H3K27ac ($p < 0.001$, odds ratio = 7.9) peaks. While there was a significant
depletion in overlapping peaks genome-wide between OVO and 32c NC H3K27me3 ($p < 0.001$, odds
ratio = 0.6), GSC H3K9me3 ($p < 0.001$, odds ratio = 0.7), and 8c NC H3K9me3 ($p < 0.001$, odds ratio =
0.5).
327
The association of OVO binding with active histone marks and open chromatin was striking, but open
chromatin is likely a general phenomenon of promoters (Haines and Eisen, 2008). Indeed, when
measuring the read density for GSC and 32C ATAC-seq for OVO bound and OVO non-bound promoters,
there is an enrichment for open chromatin at the TSS regardless of OVO binding. However, we did notice
an increase in enrichment for OVO bound promoters compared to OVO non-bound promoters (Figure
S1G), possibly suggesting that OVO bound promoters are more open or have an increase in accessibility
when compared to non-OVO bound promoters. This same relationship held true for the transcriptionally
active histone mark H3K27ac in GSCs (Figure S1H). Since only 45% of OVO ChIP peaks overlapped

TSSs, we plotted the read density of the above chromatin marks over OVO ChIP peak maximums for
OVO bound over the TSS, gene body, or intergenic regions (Figure S2A-D). We found that OVO bound
regions that were not overlapping the TSS still showed the same propensity for enrichment of open
chromatin and active histone marks. Intergenic regions were especially enriched for open chromatin
measured through ATAC-seq. Altogether suggesting that OVO binding genome-wide is tightly associated
with open chromatin regardless of germ cell stage, and active transcription in GSCs. In other words,
chromatin state data suggests OVO is acting positively on its target genes and raises the possibility that
OVO-binding and open chromatin are related.
344
OVO DNA Binding Motifs are Evenly Distributed Around Promoters and are Enriched for INR, DPE, and
MTE Elements
Our data thus far clearly indicates that OVO binding occurs at or very near the core promoter, a region
recognized by an enormous collection of factors that associate with RNA polymerase to initiate
transcription (Aoyagi and Wassarman 2000; Vo Ngoc, Kassavetis, and Kadonaga 2019). The highly
organized polymerase complex has sequence-specific DNA recognition sites with incredibly precise
spacing between them, with an overall DNA footprint of a little less than 100bp (Rice, Chamberlin, and
Kane 1993; FitzGerald et al. 2006; Ohler et al. 2002). There are upstream binding sites such as TATA,
sites at transcription start, such as the initiator (INR), and downstream promoter elements (DPE) (Vo
Ngoc, Kassavetis, and Kadonaga 2019). The combinations of these DNA motifs is not random in
mammals and Drosophila (FitzGerald et al. 2006), and distinct combinations of different motifs at the TSS
of genes expressed in *Drosophila* are conserved over tens of millions of years of evolution (Chen et al.
2014). The male germline expresses a number of TATA-associated factors that have been implicated in
male-specific promoter usage for gene expression (M. Hiller et al. 2004; M. A. Hiller et al. 2001; Lu et al.
2020; V. C. Li et al. 2009). It is possible that OVO is a female germline specific TATA-associated factor,
and if so, OVO binding sites at core promoters should share precise spacing with other core promoter
elements, suggesting it is likely part of the complex. If not, then OVO is more likely to facilitate binding of
the basal transcriptional machinery. Because of the extended footprint of engaged RNA polymerase,
OVO and the basal machinery would not be likely to occupy the same region at the same time.
364
Like OVO chip peaks, OVO DNA binding motifs were highly enriched at or near the TSS (Figure 3A). We
carefully analyzed the spacing of these sites relative to core promoter elements to see if spacing was
precise at the nucleotide level. We first searched for the presence of previously defined DNA motifs that
are enriched at promoters (FitzGerald et al. 2006; Ohler et al. 2002; Lim et al. 2004) using FIMO (Grant,
Bailey, and Noble 2011). We defined promoters by using the DNA sequences 150 nucleotides upstream
and downstream of the significant dominant TSSs in our previously analyzed ovary CAGE-seq datasets
(Chen et al. 2014). After extracting these sequences and searching for significant scoring motifs, we
plotted the density of each motif in relation to the empirically mapped TSSs (Figure 3B). We also
searched for all OVO motifs found in our significant ChIP peaks within these promoter sequences. When
plotting the density of DNA motifs found in ovary CAGE-seq promoters, we found that there were
prominent peaks for INR and M1BP (M1BP (J. Li and Gilmour 2013) = Ohler 1 (Ohler et al. 2002) = DMv4
(FitzGerald et al. 2006)) near the TSS, and MTE (Lim et al. 2004) and DPE elements downstream of the
TSS. This distribution and frequency are consistent with the constrained location of these DNA motifs
(FitzGerald et al. 2006; Ohler et al. 2002; Chen et al. 2014). Significantly, the OVO DNA binding motifs
showed a broad distribution upstream and downstream of the TSS.
380
The precise core promoter architecture of OVO bound TSSs is revealed in the CAGE-seq dataset.
Plotting the distribution of classical core promoter sequence elements in OVO bound promoters showed a
similar, but exaggerated, profile compared to all core promoters of the ovary CAGE-seq dataset. We
found a significant enrichment for INR (p < 0.01, odds ratio = 1.70), DPE (p < 0.01, odds ratio = 1.81),
MTE (p < 0.01, odds ratio = 1.65), and most importantly, OVO DNA binding motifs (p < 0.01, odds ratio =
4.83), in ovary promoters that overlapped an OVO ChIP peak in comparison to the subset of ovary
promoters that did not overlap an OVO ChIP peak (Figure 3C, 3D). This indicates that OVO bound
promoters are more likely to contain these specific promoter elements than non-OVO bound promoters.
As has been described before, promoters containing INR and DPE, but lacking TATA-box elements, are
common among *Drosophila* gonad promoters compared to promoters of other tissue types (Chen et al.
2014). The presence of TATA-box elements is negatively associated with germline-specific gene

expression (FitzGerald et al. 2006). We found that TATA-box elements were significantly depleted in
ovary CAGE-seq promoters when compared to testes ($p < 0.01$, odds ratio = 0.78) (Figure 3E) or
digestive system ($p < 0.01$, odds ratio = 0.50) (Figure 3F) CAGE-seq promoters. Indeed, both *ovo* and *otu*
have TATA-less promoters. Briefly, OVO bound promoters are characterized by the presence of INR,
DPE, MTE, and, of course, OVO DNA binding motifs. This could represent a functional class of promoters
utilized for gene expression in the *Drosophila* ovary. Importantly, the distribution of OVO DNA binding
motifs in ovary promoters is not fixed relative to TSSs or other core promoter elements. Thus, it is highly
unlikely that OVO acts as a female germline RNA Polymerase complex member that anchors the complex
to the core promoter and helps determine the +1 mRNA nucleotide. Rather, the imprecise location of
OVO binding sites might suggest that OVO is more likely to facilitate the binding of other basal
transcriptional factors.
403
<u>OVO Activates Gene Expression in the Female Germline</u>
Occupancy is a requirement for activity, but occupancy does not equal activity. Understanding the
transcriptional consequences of OVO occupancy genome-wide would allow us to investigate
mechanisms. However, as we mentioned earlier, the fact that *ovo* is absolutely required for female
germline viability greatly complicates this analysis. Measuring gene expression in dead or dying germ
cells was unlikely to be informative. Previous work from our lab has identified a transheterozygous *ovo*
allelic combination ($ovo^{ovo-GAL4}/ovo^{\Delta BP}$) that greatly reduces OVO activity resulting in sterility, however,
female germ cells are able to survive up until at least stage 5 of oogenesis (Benner et al. 2023). ovo^{ovo-}
^{GAL4} is a CRISPR/Cas9 derived T2A-GAL4-3xSTOP insertion upstream of the splice junction of exon 3 in
the *ovo-RA* transcript (Figure S1A). Importantly, this insertion in the extended exon 3 would disrupt
roughly 90% of the *ovo-B* transcripts. However, since about 10% of *ovo-B* transcripts utilize an upstream
splice junction in exon 3, these transcripts would not be disrupted with the T2A-GAL4-3xSTOP insertion
and thus allow for enough OVO activity for germ cell survival (Benner et al. 2023). Since $ovo^{ovo-GAL4}$
expresses GAL4 in place of full length OVO due to the T2A sequences, we can drive expression of a
rescuing OVO-B construct downstream of *UASp* to generate OVO⁺ female germ cells, which in fact does
rescue the arrested germ cell phenotype of $ovo^{ovo-GAL4}/ovo^{\Delta BP}$ ovaries. Therefore, in order to determine
genes that are transcriptionally responsive to OVO, we compared the gene expression profiles in sets of
ovaries that had the *ovo* hypomorphic phenotype with a negative control rescue construct (ovo^{ovo-}
$^{GAL4}/ovo^{\Delta BP}$; *UASp-GFP*)(Figure 4A) versus those that drive expression of the rescue construct expressing
OVO-B ($ovo^{ovo-GAL4}/ovo^{\Delta BP}$; *UASp-3xFHA-OVO-B*)(Figure 4B).
424
Since $ovo^{ovo-GAL4}/ovo^{\Delta BP}$; *UASp-3xFHA-OVO-B* females have full rescue of the arrested germ cell
phenotype seen in $ovo^{ovo-GAL4}/ovo^{\Delta BP}$; *UASp-GFP* females, we needed to take further measures to ensure
our analysis of gene expression was stage comparable between the two sets of ovaries. The adult female
ovary contains somatic cells, germline stem cells, and germline derived nurse cells that would be profiled
in a bulk ovary tissue RNA-seq experiment. Although OVO is only required and expressed in germline
derived cell types, we chose to dissect one day old post-eclosion $ovo^{ovo-GAL4}/ovo^{\Delta BP}$; *UASp-3xFHA-OVO-B*
female ovaries to enrich for early stages of oogenesis and collected only ovarioles containing the
germarium through previtellogenic egg chambers. $ovo^{ovo-GAL4}/ovo^{\Delta BP}$; *UASp-GFP* ovaries were collected
at the same age post-eclosion and we specifically collected ovaries that contained a visible ovariole
structure (and therefore contained germ cells) to minimize comparing germ cells to somatic ovary
structures, but rather germ cells to germ cells. We then performed RNA-seq in quadruplicate and
measured the changes in gene expression between ectopic rescue OVO and hypomorphic OVO ovaries.
We used a significance level of p-adj < 0.05 and a log2 fold change cutoff of >|0.5| to call differential
expression between these two sets of ovaries. We utilized these log2 fold change cutoffs for two reasons.
Our control ovary genotype ($ovo^{ovo-GAL4}/ovo^{\Delta BP}$; *UASp-GFP*) has hypomorphic OVO activity, hence germ
cells can survive but are arrested. With the addition of ectopic rescue OVO in $ovo^{ovo-GAL4}/ovo^{\Delta BP}$; *UASp-*
3xFHA-OVO-B ovaries, we predicted that genes that were directly regulated by OVO would
transcriptionally respond, however, we were unsure as to what degree the response would be in
comparison to hypomorphic OVO. We reasoned that if the changes were not significant between
genotypes, then minor changes in gene expression would not matter. Our second reason for using these
cutoffs is we had an internal control between the two genotypes. We knew through immunostaining that
Vas protein was present in the germline of both genotypes (Figure 4A, 4B) and therefore was likely
expressed at similar levels in the germline of both genotypes. Both genotypes also expressed *GAL4*

under the control of *ovo* in the germline. We examined the expression levels of *vas* and *GAL4* and found
that *vas* had a log2 fold change of 0.15 (p-adj = 0.03) and *GAL4* had a log2 fold change of 0.33 (p-adj =
0.18) (Figure 4C). These data suggest a slight underrepresentation of germline expression in *ovo*
hypomorphic ovaries. Therefore, by using the greater than 0.5 and less than -0.5 log2 fold change cutoffs,
and a less than 0.05 p-adj value cutoff, we would be conservative to not call genes differentially
expressed due to differences in the relative abundance of germ cells and somatic cells.
454
We were able to reliably detect the expression of 10,804 genes in these early ovarioles (Table S4). The
differential expression analysis indicated that 1,994 genes primarily expressed in the germline (see next
paragraph) significantly increased in expression with ectopic rescue OVO (Figure 4C, cyan/purple dots)
and 2,924 genes primarily expressed in the soma (see next paragraph) significantly decreased in
expression with ectopic rescue OVO expression (Figure 4C, yellow/blue dots). 5,886 genes were not
considered to be differentially expressed in our analysis (Figure 4C, gray dots). OVO is expressed in the
germline, not the soma, and previous work has shown that OVO-B is a transcriptional activator (Andrews
et al. 2000), so we hypothesized that many of the genes increasing in expression in the presence of
rescuing OVO were direct downstream targets. We found that 2,298 genes that were expressed in our
RNA-seq data overlapped an OVO ChIP peak. 666 genes significantly increased in expression and were
bound by OVO, which is a significant enrichment according to a Fisher's exact test (Figure 4C, cyan dots,
p < 0.01, odds ratio = 2.21). While conversely, 564 genes decreased in expression and were bound by
OVO, indicating a significant depletion according to a Fisher's exact test (Figure 4C, blue dots, p < 0.01,
odds ratio = 0.85). This strongly suggests that genes that are bound by OVO, transcriptionally respond in
a positive manner. This finding is fully consistent with our meta-analysis comparing OVO ChIP-seq and
histone ChIP/ATAC-seq data (Figure 4E). OVO binding was highly associated with transcriptionally active
histone marks such as H3K27ac and H3K4me3, open chromatin, and increased expression.
472
There are genes that showed decreased expression in the OVO rescued ovaries, but we believe this is
technical rather than biological. OVO is expressed only in the germline, but ovarioles contain germ cells
and somatic cells. The presence of empty ovarioles, containing leftover strings of somatic cells, are
evident even in *ovo*^{*ovo-GAL4*}/*ovo*^{ΔBP}; *UASp-GFP* ovaries that contain germ cells. Conversely, *ovo*<sup>*ovo-*
GAL4</sup>/*ovo*^{ΔBP}; *UASp-3xFHA-OVO-B* ovaries are fully rescued, and therefore possess more germ cell
containing ovarioles than *ovo*^{*ovo-GAL4*}/*ovo*^{ΔBP}; *UASp-GFP* ovaries (Benner et al. 2023). Despite our best
efforts to dissect individual ovarioles with a full complement of germ cells and egg chambers, we
wondered if there might be fewer germ cells and egg chambers in the *ovo*^{*ovo-GAL4*}/*ovo*^{ΔBP}; *UASp-GFP*
ovaries. To confirm that genes increasing in expression in ectopic rescue OVO were germline derived, we
cross-referenced the significantly expressed genes in our RNA-seq datasets with the modENCODE
developmental RNA-seq datasets (Graveley 2010). We extracted the gene names of all genes that were
considered to be 'moderately expressed' in 0-2 hour old embryos, which are produced during oogenesis
and are deposited into the early embryo. We found that 71% of genes (1,409/1,994) that had a significant
increase in expression in the presence of ectopic rescue OVO were found to be expressed in 0-2 hour old
embryos (Figure 4D, green dots), while only 21% of genes (625/2,924) that had a significant decrease in
expression were found in the same embryo dataset (Figure 4D, yellow dots). 3,448 genes from the 0-2
hour old embryo dataset were not differentially expressed in our RNA-seq dataset (Figure 4D, purple
dots). A Fisher's exact test confirmed that there was a significant enrichment for genes that significantly
increased in expression and were present in 0-2 hour old embryos (p < 0.01, odds ratio = 2.8). In
comparison, there was a significant depletion for genes that significantly decreased in expression and
were present in 0-2 hour old embryos (p < 0.01, odds ratio = 0.17). This result indicated that genes that
significantly increased in expression were more likely to be expressed in the germline and that the
presence of ectopic rescue OVO significantly increased the expression of genes that were maternally
deposited in the early embryo. While the set of genes that significantly decrease in expression are not
enriched in the embryo and are more likely specific to somatic cell gene expression. These genes are
unlikely to be direct OVO targets due to the absence of OVO in those cells, although we certainly cannot
rule out the possibility of a non-autonomous effect of OVO on somatic gene expression. In terms of the
germline proper, OVO appears to be a positively acting transcription factor.
501
OVO Positively Regulates Essential Oogenesis Genes

We wanted to examine a subset of the OVO target genes in detail, and began with the known OVO
targets, *ovo* itself and *otu* (Bielinska et al. 2005; Lü et al. 1998; Lü and Oliver 2001). Since the
relationship between OVO binding to these two genes has been well-characterized, we validated the
OVO ChIP, histone ChIP/ATAC-seq, and RNA-seq datasets by examining these two genes first. Since
OVO positively regulates the expression of both these genes, then we would expect OVO to be physically
bound at OVO motifs required for high transcription, the presence of transcriptionally active histone marks
and open chromatin at these loci, as well as a positive transcriptional response in the presence of
rescuing OVO-B. This is exactly what we observed. A significant OVO ChIP peak was found overlapping
the TSS of *ovo-B*, with four significant OVO DNA binding motifs present (Figure 5A). ATAC-seq,
H3K27ac, and H3K4me3 peaks overlapped the *ovo* promoter. Transcriptionally, *ovo* RNA-seq reads are
likely derived from the *UASp-3xFHA-OVO-B* cDNA rescue or are indistinguishable between the genomic
locus and rescuing cDNA transgene. We found a non-significant increase in exon 3 to exon 4 intronic *ovo*
reads with the expression of ectopic rescue OVO (log2 fold change = 0.76, p-adj = 0.26). These intronic
reads would be derived from the endogenous *ovo* locus, but it is difficult to conclusively determine if the
endogenous *ovo* locus would respond transcriptionally to ectopic OVO downstream of *UASp* (for
example, the pathway for *ovo* is no longer autoregulatory in *ovo$^{ovo-GAL4}$/ovo$^{\Delta BP}$*; *UASp-3xFHA-OVO-B*
germ cells, there is an additional GAL4>*UASp* activation step). So, we could not confidently assess
whether *ovo* responded transcriptionally to ectopic rescue OVO. However, when looking at the *otu* locus
(Figure S3), we found OVO occupancy over the TSS of both annotated *otu* promoters, with significant
OVO DNA binding motifs overlapping and in close proximity to the TSSs. The *otu* locus contained similar
ATAC-seq and activating histone mark peaks overlapping the TSS found at the *ovo* locus. It was also
evident that *otu* had a positive transcriptional response to the presence of OVO rescue (log2 fold change
= 2.41; p-adj < 0.001). These results confirm that OVO binds and positively regulates both itself and *otu in*
vivo, as previous work has indicated.
527
Since our overlapping OVO ChIP-seq and RNA-seq data suggests that hundreds of genes that are bound
by OVO increase in expression in the presence of ectopic rescue OVO, we wanted to know more about
the functions of those genes. To do this, we performed Gene Ontology enrichment analysis with gProfiler
software (Raudvere et al. 2019). To be especially stringent, we focused on the genes that contained an
OVO ChIP peak overlapping the transcriptional start site and significantly increased in expression in the
presence of rescue OVO. 525 genes met these criteria. Biological process GO term enrichment analysis
on these 525 genes showed a significant enrichment for 45 GO terms (Table S5). The significant GO
terms were almost exclusively related to female reproduction and maternal control of early embryonic
development (Figure 6A). Genes that are required for processes such embryonic axis specification,
mRNA localization, egg activation, and translational regulation were found within these significantly
enriched GO terms. These GO terms are well understood in the context of oogenesis and broadly
suggest that OVO expression in adult gonads is essential for constructing an egg and depositing maternal
RNAs to support zygotic embryonic development.
541
GO term enrichment analysis of genes that are bound by OVO and increase in expression in the
presence of ectopic rescue OVO suggested that OVO is likely a main transcriptional regulator of
oogenesis. These genes are the subject of decades of work on *Drosophila* oogenesis, but essentially all
the work on them has focused on what they do, not on how they are transcriptionally regulated. For
example, *bicoid* (Figure 6B), and *bicoid* mRNA binding proteins *exuperantia* (*exu*) and *swallow* (*swa*), are
essential for anterior specification of the embryo (Lasko 2012). All these genes were occupied by OVO *in*
vivo, significantly upregulated by OVO-B, and had OVO motifs in close proximity to the TSS. Genes
involved in posterior patterning (*oskar* and *nanos*) (Figure 6C), as well as pole cell specification genes
(*polar granule component*, *germ cell-less*, and *aubergine*)(Lasko 2012; L. Benner, Deshpande, and Lerit
2018), also showed similar RNA-seq, ChIP-seq, and OVO DNA binding motif profiles as *ovo* and *otu*.
Genes that are involved in translational silencing and regulation of maternally provided mRNAs, such as
cup (Figure 6D), *maternal expression at 31B* (*me31B*), *oo18 RNA-binding protein* (*orb*), and *bruno 1*
(*bru1*), as well as essential genes involved in meiosis completion and egg activation after fertilization
(*giant nuclei* (*gnu*), *pan gu* (*png*), *plutonium* (*plu*), *wispy* (*wisp*), *C(3)G*, and *matrimony* (*mtrm*)) (Figure
6E)(Lasko 2012; Avilés-Pagán and Orr-Weaver 2018) all show this stereotypic pattern of promoter
proximal OVO occupancy and DNA binding motifs, and OVO-dependent transcription. These data

indicate that the OVO is a central transcription factor activating the expression of essential maternal and
early embryonic development pathways in the female germline.
560
We also found that the genes *fs(1)N*, *fs(1)M3*, and *closca*, were all bound by OVO and responded
transcriptionally to the presence of ectopic rescue OVO. These genes are significant because they
constitute a set of genes that are expressed in the germline and the encoded proteins are eventually
incorporated into the vitelline membrane providing the structural integrity and impermeability of the egg
(Mineo, Furriols, and Casanova 2017; Ventura et al. 2010). Loss-of-function of these three genes results
in flaccid eggs that are permeable to dye and fail to develop. The loss-of-function phenotype of *fs(1)N*,
fs(1)M3, and *closca* closely resembles the dominant antimorph ovo^{D3} phenotype. The ovo^{D3} allele is the
weakest of the original dominant-negative *ovo* alleles and produces defective eggs allowing us to explore
the role of OVO in late stages (Busson et al. 1983; Komitopoulou et al. 1983). ovo^{D3}/ovo$^+$
transheterozygous females express a repressive form of OVO that results in dominant sterility, and
importantly, these females lay flaccid eggs with compromised vitelline membranes that are permeable to
the dye neutral red (Oliver, Pauli, and Mahowald 1990). Since OVO$^+$ is bound at the TSS of *fs(1)N*,
fs(1)M3, and *closca*, and these three genes respond transcriptionally to OVO$^+$, then it is plausible that the
repressive OVO^{D3} is negatively regulating these three genes that are required for vitelline membrane
formation. This is evidence that OVO is not only involved in regulating the expression of numerous
essential maternal pathways for embryonic development, but it is also essential for regulating genes that
are required for egg integrity and maturation.
578
Discussion
Since its original isolation as a dominant female sterile locus in Drosophila (Busson et al. 1983;
Komitopoulou et al. 1983), o*vo* has long been known as an essential gene in oogenesis. Female germ
cells require *ovo* for survival and differentiation, while it has no described roles or functions in the male
germline (Oliver, Perrimon, and Mahowald 1987; Oliver, Pauli, and Mahowald 1990). OVO has also been
found to be eternally present in the female germline, attesting to its likely continual requirement for female
germ cell viability and identity (Benner et al. 2023). Our work here significantly expands our knowledge on
OVO function in the female germline, showing that OVO binds and positively regulates a large array of
genes required to build an egg and pattern the resulting embryo after fertilization. OVO accomplishes this
by directly binding to the promoters of its targets, as well as more distant sites that could represent
enhancers. Altogether, we suggest that OVO is a master transcriptional regulator coordinating a number
of essential maternal pathways involved in oocyte and early embryonic development. Hints of these
functions can be found in the hypomorphic and antimorphic *ovo* alleles which show egg chamber arrest,
ventralized eggs, and permeable vitelline membranes. It is clear that OVO is required to activate multiple
pathways involved in oocyte and early embryonic development.
594
The GO term enrichment analysis on genes that were bound by OVO and transcriptionally responded to
OVO surprisingly indicated a large degree of overlap in oocyte and early embryonic developmental
pathways. Also, OVO seemed to reinforce these pathways at multiple key genes within each pathway.
For example, OVO bound to the promoters and increased the expression of *bcd*, and the *bcd* mRNA
binding proteins *exu* and *swa*, all involved in ensuring correct anterior specification of the embryo (Lasko
2012). Genes that are essential for egg activation were coordinately regulated by OVO as well. OVO
downstream target genes *gnu*, *png*, *plu*, and *wisp* all belong in the same interconnected pathway
ensuring egg activation (Avilés-Pagán and Orr-Weaver 2018). A similar story was found for genes such
as *cup*, *Me31b*, *bru1*, and *orb*, indicating that OVO controls a battery of genes involved in the positive
regulation of RNA binding proteins that negatively regulate translation (Lasko 2012). OVO also bound and
positively regulated a number of posterior and germ plasm specification genes such as *osk*, *nos*, *aub*, *gcl*,
and *pgc* (Lasko 2012; Mahowald 2001). Given this plethora of famous maternal effect loci, it might be
tempting to suggest that OVO is sufficient for egg production, but there are important exceptions. For
example, other important germ plasm factors such as *staufen* and *tudor* were not bound by or
transcriptionally responsive to OVO. This observation suggest that other transcription factors are
responsible for regulating these genes.
611
OVO binds in close proximity to the TSS of genes it positively regulates, however, it is still unclear
precisely how it regulates gene expression. The possibilities include integration into the RNA Polymerase

complex itself, a short distance sigma factor like function, a core promoter conditioning function
(pioneering), and/or garden variety transcription factor. Although active core promoters specifically in the
ovary are enriched for OVO DNA binding motifs, we did not find a strict spatial orientation for these motifs
in relation to the TSS, such as is found with other DNA elements such as INR, DPE, and MTE (Ohler et
al. 2002; FitzGerald et al. 2006; Lim et al. 2004). It is therefore unlikely that OVO is a core component of
the RNA Polymerase complex in the female germline. This suggests that it is unlikely to be analogous to
male specific TATA-associated factors that have been shown to activate gene expression in the male
germline (M. Hiller et al. 2004; M. A. Hiller et al. 2001; Lu et al. 2020; V. C. Li et al. 2009). It is therefore
possible, and previously well supported, that OVO is an activator of transcription (Lü et al. 1998; Lü and
Oliver 2001; Bielinska et al. 2005). One aspect of OVO DNA binding that showed differences with stage-
specific histone ChIP and ATAC-seq, was OVO's strong association with open chromatin. The role of
repressors of transcription such as the polycomb complex, *egg*, *wde*, and *Su(var)205* in restricting gene
expression through promoting heterochromatin formation in differentiating egg chambers is well
established (Smolko, Shapiro-Kulnane, and Salz 2018; DeLuca et al. 2020). OVO might ensure that the
chromatin status of maternally expressed genes remains open. Evidence from our work points to OVO
fulfilling that role.
630
In GSCs OVO ChIP peaks largely overlap open chromatin and transcriptionally active histone marks.
However, in stage 5 egg chambers, there was an even higher degree of association with open chromatin
(almost all OVO ChIP peaks overlapped ATAC-seq peaks), while the significant association with
H3K27ac marks was greatly reduced. This difference is likely significant. As GSCs differentiate they
accumulate repressive chromatin marks while the number of ATAC-seq and H3K27ac peaks are reduced.
This increase in association with open chromatin and OVO binding, even as the amount of open
chromatin is reduced throughout egg chamber differentiation, might indicate that OVO binding helps to
maintain chromatin accessibility, even when the locus in question is no longer actively transcribed. The
loss of histone marks of active transcription at OVO-bound open chromatin in later differentiating egg
chambers might mean that OVO does not influence the transcriptional potential of target genes as
strongly as it influences the chromatin status in this second phase. Therefore, OVO might be more similar
in function to pioneer factors/chromatin remodelers than it is to a transcription factor that is only involved
in activating transcription.
644
The requirement for OVO at the TSS of target genes has been well characterized at its own locus as well
as its downstream target *otu*. Our OVO ChIP and expression data confirm findings from previous work
that OVO is binding to these target promoters, and in the case of *otu*, strongly responds transcriptionally
to the presence of OVO. Although we did not test the requirement for OVO DNA binding motifs at other
OVO bound genes in this work, this has been extensively explored before, showing that removal of OVO
DNA binding sites overlapping the TSS results in a strong decrease in reporter expression (Lü et al. 1998;
Bielinska et al. 2005; Lü and Oliver 2001). Removal of more distal upstream OVO DNA binding sites also
reduces reporter expression to a lesser degree. However, for most cases tested, removal of OVO DNA
binding sites while leaving the rest of the enhancer regions intact, never totally abolished reporter
expression. These dynamics are highly similar to work that has been completed on the pioneer factor
zelda (*zld*). Adding *zld* DNA binding motifs to a stochastically expressed transcriptional reporter increases
the activity and response of the reporter (Dufourt et al. 2018). Distally located *zld* DNA binding motifs
influenced reporter expression to a lesser degree than proximal sites. A single *zld* DNA binding site
adjacent to the TSS produced the strongest reporter activity. Importantly, just like the activity of OVO
transgenic reporters, there is not an absolute requirement for *zld* DNA binding to activate reporter
expression, however, the addition of TSS adjacent *zld* DNA binding motifs does strongly influence
reporter response. We know that *zld* achieves this reporter response through its pioneering activity (Xu et
al. 2014; Harrison et al. 2011), whether OVO achieves this similar effect on gene expression through a
shared mechanism, or in cooperation with other transcription factors needs to be further explored.
664
665   **Methods**
666   All reagents used in this study can be found in the FlyBase recommended supplementary ART table
667   (Table S6).
668
669   Fly Husbandry

670 All fly crosses were conducted at 25°C with 65% relative humidity and constant light unless otherwise
671 noted. Flyfood consisted of premade flyfood (D20302Y) from Archon Scientific (Durham, NC).
672
673 <u>Immunofluorescence and Image Analysis</u>
674 All immunostaining procedures were done as previously described (Benner et al. 2023). Antibodies and
675 their respective dilutions are indicated in Table S6. Imaging was also completed as previously described
676 (Benner et al. 2023).
677
<u>RNA-seq library preparation and sequencing</u>
Twenty, one day old post-eclosion $ovo^{\Delta BP}/ovo^{ovo-GAL4}$; *UASp-GFP* and $ovo^{\Delta BP}/ovo^{ovo-GAL4}$; *UASp-3xFHA-*
OVO-B ovaries were dissected and germariums through previtellogenic egg chambers were removed with
microdissection scissors and placed in ice cold PBS making up one biological replicate. RNA was then
extracted from four biological replicates with a Qiagen RNeasy Plus Kit (Qiagen) according to the
manufacturer's protocol, eluted in dH_2O, and RNA concentrations were measured with Quant-iT
RiboGreen RNA Assay Kit (ThermoFisher Scientific). 500 ng of total RNA was then used to make RNA-
seq libraries with an Illumina Stranded mRNA Prep Kit according to the manufacturer's protocol (Illumina).
IDT for Illumina RNA UD Indexes Set A were used. Library concentrations were measured with Quant-iT
PicoGreen dsDNA Assay Kit (ThermoFisher Scientific), pooled, and then 50 nucleotide paired-end DNA
sequencing was completed on an Illumina NovaSeq 6000 system using a S1 flow cell (Illumina). Raw
RNA-seq reads are available at the SRA (SAMN38284748-SAMN38284755).
690
<u>ChIP-seq library preparation and sequencing</u>
Adult $ovo^{Cterm-3xFHA}$ and $ovo^{Cterm-GFP}$ females were collected and fed for 24 hours before ovaries were
dissected. 50 dissected ovaries were placed in ice cold phosphate buffered saline (PBS, Gibco,
ThermoFisher Scientific) and then incubated in 1 mL crosslinking solution containing 2% formaldehyde
(Pierce, ThermoFisher Scientific) (50mM HEPES Buffer, 1mM EDTA, 0.5 mM EGTA, 100mM NaCl), and
rotated at 37°C for 20 minutes. Ovaries were then incubated in 1 mL stop solution (125mM Glycine,
0.01% Triton X-100 (Millipore Sigma), diluted in PBS) and rotated for 5 minutes at room temperature.
Ovaries were then washed twice with 1 mL ice cold wash buffer (0.01% Triton X-100 in PBS) for 5
minutes. The last wash was removed, and ovaries were stored at -80°C until future processing. Once all
samples were collected, 4x50 ovaries were then homogenized in 250 µL RIPA lysis buffer (Pierce,
ThermoFisher Scientific) containing 1x protease inhibitor cocktail (cOmplete Mini Protease Inhibitor
Cocktail, Roche, Millipore Sigma) and 1 mM PMSF (Roche, Millipore Sigma) and kept on ice for 10
minutes. 40mg of 212-300 µm acid-washed glass beads (Millipore Sigma) were then added to
homogenized ovary lysate. Samples were then sonicated with a Bioruptor Pico sonication device
(Diagenode) at 4C for 15 cycles of 30 seconds on and 30 seconds off. Sonicated lysate was then
transferred to a new tube and centrifuged at 13,300 rpm for 10 minutes at 4°C. Three supernatants were
then combined to form one biological replicate. 100 µL for each biological replicate was removed and
stored at -80°C for input control. To pull down C-terminally tagged OVO, 100 µL of monoclonal anti-HA-
agarose (Millipore Sigma) or 50 µL of ChromoTek GFP-Trap agarose (Proteintech) were washed three
times with RIPA lysis buffer and spun down at 1,200 RPMs for one minute at 4°C. 550 µL of $ovo^{Cterm-3xFHA}$
supernatant was added to monoclonal anti-HA-agarose and 550 µL of $ovo^{Cterm-GFP}$ supernatant was added
to ChromoTek GFP-Trap agarose. Samples were supplemented with 1x protease inhibitor cocktail and 1
mM PMSF and incubated on a rotator at 4°C overnight.
714
The next day, agarose was washed in a stepwise fashion with solutions from a Chromatin
Immunoprecipitation Assay Kit (Millipore Sigma), beginning with 1 mL of a low salt wash buffer, high salt
wash buffer, LiCl buffer, and ending with 2 washes in 0.1x TE buffer. 300 µL of freshly prepared ChIP
elution/decrosslinking solution (1% SDS, 100mM $NaHCO_3$, 250 mM NaCl, 10mM EDTA, 50 mM Tris-HCl,
200 µg/mL Proteinase K) was added to the pelleted agarose, or 200 µL of chip elution/decrosslinking
solution was added to 100 µL input control and incubated at 65°C overnight. DNA was extracted by
adding 300 µL phenol:chloroform:iso-amyl alcohol (125:24:1) (Millipore Sigma). The samples were
vortexed for 30 seconds then centrifuged at 13,300 RPMs for 5 minutes at 4°C. The aqueous layer was
extracted, and this process was repeated once more. 1 µL glycogen (20 mg/mL), 30 µL 1M sodium
acetate, and 750 µL 100% EtOH was added to the extracted aqueous layer, vortexed, and incubated at -
20°C for 30 minutes. Solution was spun at 13,300 RPMs for 20 min at 4°C. Supernatant was removed

and the pellet was washed with 500 μL 70% EtOH and spun down at 13,300 RPMs for 20 minutes at 4°C.
This step was repeated but with 100% EtOH. The resulting pellet was briefly speedvacced and
resuspended in 50 μL dH$_2$O.
729
To make ChIP-seq libraries, DNA concentration for immunoprecipitated and input control samples were
measured with a Quant-iT PicoGreen dsDNA Assay Kit (ThermoFisher Scientific). 5 ng of DNA for each
sample was then used with the NEBNext Ultra II DNA Library Prep Kit for Illumina (New England Biolabs)
and completed according to the manufacturer's protocol. ChIP-seq library concentrations were then
measured with a Quant-iT PicoGreen dsDNA Assay Kit, pooled, and then 50 nucleotide paired-end DNA
sequencing was performed on an Illumina NovaSeq 6000 system using the XP workflow (Illumina). Raw
ChIP-seq reads are available at the SRA (SAMN38284736-SAMN38284747).
737
<u>RNA-seq, ChIP-seq, CAGE-seq and Gene Ontology Analysis</u>
For RNA-seq analysis of *ovo^{ΔBP}/ovo$^{ovo-GAL4}$*; *UASp-GFP* and *ovo^{ΔBP}/ovo$^{ovo-GAL4}$*; *UASp-3xFHA-OVO-B*
ovaries, 50 nucleotide paired-end reads were mapped to the FlyBase r6.46 genome (Gramates et al.
2022) for differential expression analysis and the BDGP Release 6 Drosophila Genome (dos Santos et al.
2015) for read level genome browser tracks using Hisat2 (-k 1 --rna-strandness RF --dta)(Kim et al.
2019). DNA sequences for *GAL4* and *GFP* were added to the FlyBase r6.46 genome as separate
chromosomes. Mapped reads were then sorted and indexed with Samtools (samtools sort and samtools
index)(Danecek et al. 2021). Gene level readcounts were then derived with htseq-count (-s reverse -r
pos)(Anders, Pyl, and Huber 2015) and used for differential expression analysis with DeSeq2 (Love,
Huber, and Anders 2014). Genes with 0 mapped reads were removed from the DESeq2 analysis.
748
For ChIP-seq analysis of OVO-HA, OVO-GFP, OVO-HA input, and OVO-GFP input samples, 50
nucleotide paired-end reads were mapped to the FlyBase r6.46 genome for peak calling analysis and the
BDGP Release 6 Drosophila Genome for read level genome browser tracks using Hisat2 (-k 1 --no-
spliced-alignment -X 900). Mapped reads were sorted using Samtools (samtools sort and samtools index)
and duplicate reads were removed with Picard (REMOVE_DUPLICATES=true)(broadinstitute, n.d.).
Significant ChIP peaks were called for OVO-HA and OVO-GFP versus their respective input controls
separately using Macs3 callpeak software (-g 1.2e8 -q 0.0001)(Zhang et al. 2008). Overlapping ChIP
peaks for OVO-HA and OVO-GFP were then determined with bedtools intersect software (Quinlan and
Hall 2010). Peak calling for GSC ATAC-seq (SRR24203655), 32c ATAC-seq (SRR24203650), GSC
H3K27ac (SRR11084657), H3K4me3 (SRR11084658), H3K27me3 (SRR11084656), H3K9me3
(SRR24203631), 8c NC H3K4me3 (SRR24203629), 32c NC H3K27ac (SRR24203635), and H3K27me3
(SRR11084652) ChIP-seq versus their respective input controls (SRR11084655, SRR11084651,
SRR24203634, SRR24203637) was conducted in the same manner as OVO ChIP-seq.
762
763 In order to generate gene-level read coverage tracks, deepTools' bamCompare software was used to
764 generate a single bigWig file comparing all replicates versus input controls (-bs 5 --effectiveGenomeSize
765 142573017 --normalizeUsing BPM --exactScaling --scaleFactorsMethod None)(Ramírez et al. 2016). The
766 bigWig file was then uploaded to UCSC genome browser for visualization (Kent et al. 2002).
767
To generate read coverage plots centered on the motif location or OVO peak maximums, genomic
locations of significant scoring motifs or peak maximums within overlapping OVO ChIP peaks were
determined and used as input for deepTools' computeMatrix reference-point (-a 2000 -b 2000 -bs 25 --
missingDataAsZero). Read density profiles for each motif or OVO peak maximum were then visualized
with deeptools plotProfile. In order to generate read coverage plots centered on the TSS, the same
methods as above were conducted except genes overlapping OVO ChIP peaks containing the respective
significant OVO DNA binding motifs were used as input instead.
775
776 In order to generate gene level read coverage heatmaps, deepTools' computeMatrix scale-regions
777 software was used to generate a single matrix for genes that were bound by OVO (-bs 25 --
778 missingDataAsZero -m 4000 --metagene). This matrix was then used as input for deepTools'
779 plotHeatmap software to generate heatmaps of ChIP-seq and RNA-seq for the given OVO binding
780 profiles centered on the TSS (--sortUsing max).
781

For CAGE-seq analysis, CAGE-seq libraries for ovary (SRR488283, SRR488282), testes (SRR488284,
SRR488308, SRR488285, SRR488309), and male and female digestive system (SRR488289,
SRR488288) tissues were downloaded and combined for each tissue type from the SRA. Reads were
mapped to the BDGP Release 6 Drosophila Genome with Hisat2 (-k 1). Mapped reads were sorted with
Samtools. Significant dominant TSSs were then determined with CAGEr software(Haberle et al. 2015)
from sorted BAM files with getCTSS and annotated with annotateCTSS using the dm6.ensGene.gtf
file(Hubbard et al. 2002) downloaded from UCSC(Kent et al. 2002). CAGE-seq reads were normalized
with normalizeTagCount (ce, method = "simpleTpm", fitInRange = c(5, 40000), alpha = 1.15, T = $1*10^6$)
and then TSS clusters were determined with clusterCTSS (ce, threshold = 1, thresholdIsTpm = TRUE,
nrPassThreshold = 1, method = "paraclu", maxDist = 20, removeSingletons = TRUE,
keepSingletonsAbove = 5) in order to determine the dominant significant TSS for each respective tissue.
793
Gene ontology enrichment analysis was completed with g:Profiler's g:GOSt software (Raudvere et al.
2019) on the set of genes overlapping OVO ChIP peaks over the TSS and significantly upregulated in the
presence of ectopic OVO (525 genes in total). All genes that were considered to be expressed in our
RNA-seq datasets were used as a background control (10,801 genes in total). Default parameters were
used for the enrichment analysis except for 'statistical domain scope' was set to 'custom' (our control
background genes were uploaded here), 'significance threshold' was set to 'Bonferroni correction', and
only GO biological process terms were searched for enrichment with the gene list. The GO terms listed in
Figure 6 represent the 24 smallest GO term sizes according to Table S5.
802
803   Fisher's exact test was conducted for each respective analysis with the fisher.test() command in R (R
804   Core Team 2021).
805
de novo Motif Enrichment and Promoter Motif Analysis
DNA sequences from significant overlapping OVO ChIP peaks were extracted from the Drosophila r6.46
genome and submitted to STREME software (Bailey 2021) of the MEME suite (Bailey et al. 2015). The
default parameters were used for *de novo* motif enrichment analysis, including the use of shuffled input
sequences as a control. After identifying 'OVO Motif One', OVO ChIP peaks that contained that sequence
were removed and the resulting ChIP peaks were resubmitted for STREME analysis deriving derivative
OVO DNA binding motifs like above. Significant OVO DNA binding motifs and *in vitro* OVO DNA binding
motifs were searched in the BDGP Release 6 Drosophila Genome using FIMO (Grant, Bailey, and Noble
2011). In order to find significant DNA binding motif matches for 'OVO Motif One', this motif from
STREME was submitted to Tomtom software (Gupta et al. 2007) of the MEME suite and searched within
the JASPAR Core Insect database (2022)(Castro-Mondragon et al. 2022). XSTREME software was used
with the default parameters to identify previously characterized and published DNA binding motifs within
OVO ChIP peaks (Grant and Bailey 2021).
819
Promoter motif analysis was conducted by extracting the DNA sequences 200 nucleotides upstream and
downstream of the significant dominant TSSs from CAGE-seq analysis for each respective tissue type. All
common core promoter motifs (FitzGerald et al. 2006; Ohler et al. 2002) were then searched in these
sequences depending on their strand specificity with the use of FIMO from the MEME suite using a p-
value of < 0.003 for all non-OVO promoter motifs. All OVO motifs found in this study and through *in vitro*
methods were also searched with the same method, except a p-value of < 0.0002 was used.
826
827   **Data Availability**
828   Drosophila strains used for this study are available upon request. All sequence information and datasets
829   used in this study are in Table S6.
830
Acknowledgements

We would like to thank previous and current members of the Oliver lab, Laboratory of Biochemistry and
Genetics at NIH, and L.B. committee members Mark Van Doren and Allan Spradling for insightful
discussion and comments on this work throughout. Monoclonal antibodies were obtained from the
Developmental Studies Hybridoma Bank, created by the Eunice Kennedy Shriver National Institute of
Child Health and Human Development (NICHD) of the NIH and maintained at the Department of Biology,
University of Iowa, Iowa City, IA 52242. Genetic and genomic information was obtained from FlyBase

(U41 HG-000739). This work utilized the computational resources of the NIH High-Performance
Computing Biowulf cluster (http://hpc.nih.gov). Sequencing was completed by The National Heart, Lung,
and Blood Institute (NHLBI) DNA Sequencing and Genomics Core.
841
842 **Funding**
843 This research was supported in part by the Intramural Research Program of the NIH, The National
844 Institute of Diabetes and Digestive and Kidney Diseases (NIDDK) (awarded to B.O.). L.B. was supported
845 by the NIH Graduate Partnership Program.
846
References
Anders, Simon, Paul Theodor Pyl, and Wolfgang Huber. 2015. "HTSeq--a Python Framework to Work
with High-Throughput Sequencing Data." *Bioinformatics* 31 (2): 166–69.
Andrews, J., D. Garcia-Estefania, I. Delon, J. Lü, M. Mével-Ninio, A. Spierer, F. Payre, D. Pauli, and B.
Oliver. 2000. "OVO Transcription Factors Function Antagonistically in the Drosophila Female
Germline." *Development* 127 (4): 881–92.
Aoyagi, N., and D. A. Wassarman. 2000. "Genes Encoding Drosophila Melanogaster RNA Polymerase II
General Transcription Factors: Diversity in TFIIA and TFIID Components Contributes to Gene-
Specific Transcriptional Regulation." *The Journal of Cell Biology* 150 (2): F45–50.
Avilés-Pagán, Emir E., and Terry L. Orr-Weaver. 2018. "Activating Embryonic Development in
Drosophila." *Seminars in Cell & Developmental Biology* 84 (December): 100–110.
Bailey, Timothy L. 2021. "STREME: Accurate and Versatile Sequence Motif Discovery."
Bioinformatics 37 (18): 2834–40.
Bailey, Timothy L., James Johnson, Charles E. Grant, and William S. Noble. 2015. "The MEME Suite."
Nucleic Acids Research 43 (W1): W39–49.
Bastock, Rebecca, and Daniel St Johnston. 2008. "Drosophila Oogenesis." *Current Biology: CB* 18 (23):
R1082–87.
Benner, L., G. Deshpande, and D. A. Lerit. 2018. "Primordial Germ Cells of Drosophila Melanogaster."
https://scholar.google.ca/scholar?cluster=7409236235465525669&hl=en&as_sdt=0,5&sciodt=0,5.
Benner, Leif, Elias A. Castro, Cale Whitworth, Koen J. T. Venken, Haiwang Yang, Junnan Fang, Brian
Oliver, Kevin R. Cook, and Dorothy A. Lerit. 2019. "Drosophila Heterochromatin Stabilization
Requires the Zinc-Finger Protein Small Ovary." *Genetics* 213 (3): 877–95.
Benner, Leif, Savannah Muron, and Brian Oliver. 2023. "Female Germline Expression of OVO
Transcription Factor Bridges Drosophila Generations." *bioRxiv*.
https://doi.org/10.1101/2023.08.25.554887.
Bielinska, Beata, Jining Lü, David Sturgill, and Brian Oliver. 2005. "Core Promoter Sequences Contribute
to Ovo-B Regulation in the Drosophila Melanogaster Germline." *Genetics* 169 (1): 161–72.
Bishop, D. L., and R. C. King. 1984. "An Ultrastructural Study of Ovarian Development in the otu7 Mutant
of Drosophila Melanogaster." *Journal of Cell Science* 67 (April): 87–119.
broadinstitute. n.d. "Picard." Picard. http://broadinstitute.github.io/picard/.
Busson, D., M. Gans, K. Komitopoulou, and M. Masson. 1983. "Genetic Analysis of Three Dominant
Female-Sterile Mutations Located on the X Chromosome of DROSOPHILA MELANOGASTER."
Genetics 105 (2): 309–25.
Castro-Mondragon, Jaime A., Rafael Riudavets-Puig, Ieva Rauluseviciute, Roza Berhanu Lemma, Laura
Turchi, Romain Blanc-Mathieu, Jeremy Lucas, et al. 2022. "JASPAR 2022: The 9th Release of the
Open-Access Database of Transcription Factor Binding Profiles." *Nucleic Acids Research* 50 (D1):
D165–73.
Chen, Yu-Chien, Suewei I. Lin, Ying-Kuan Chen, Chuen-Sheue Chiang, and Gwo-Jen Liaw. 2009. "The
Torso Signaling Pathway Modulates a Dual Transcriptional Switch to Regulate Tailless Expression."
Nucleic Acids Research 37 (4): 1061–72.
Chen, Zhen-Xia, David Sturgill, Jiaxin Qu, Huaiyang Jiang, Soo Park, Nathan Boley, Ana Maria Suzuki, et
al. 2014. "Comparative Validation of the D. Melanogaster modENCODE Transcriptome Annotation."
Genome Research 24 (7): 1209–23.
Danecek, Petr, James K. Bonfield, Jennifer Liddle, John Marshall, Valeriu Ohan, Martin O. Pollard,
Andrew Whitwham, et al. 2021. "Twelve Years of SAMtools and BCFtools." *GigaScience* 10 (2).
https://doi.org/10.1093/gigascience/giab008.

DeLuca, Steven Z., Megha Ghildiyal, Liang-Yu Pang, and Allan C. Spradling. 2020. "Differentiating
Drosophila Female Germ Cells Initiate Polycomb Silencing by Regulating PRC2-Interacting Proteins."
eLife 9 (August). https://doi.org/10.7554/eLife.56922.
Dufourt, Jeremy, Antonio Trullo, Jennifer Hunter, Carola Fernandez, Jorge Lazaro, Matthieu Dejean,
Lucas Morales, et al. 2018. "Temporal Control of Gene Expression by the Pioneer Factor Zelda
through Transient Interactions in Hubs." *Nature Communications* 9 (1): 1–13.
FitzGerald, Peter C., David Sturgill, Andrey Shyakhtenko, Brian Oliver, and Charles Vinson. 2006.
"Comparative Genomics of Drosophila and Human Core Promoters." *Genome Biology* 7 (7): R53.
Gramates, L. Sian, Julie Agapite, Helen Attrill, Brian R. Calvi, Madeline A. Crosby, Gilberto Dos Santos,
Joshua L. Goodman, et al. 2022. "FlyBase: A Guided Tour of Highlighted Features." *Genetics* 220
(4). https://doi.org/10.1093/genetics/iyac035.
Grant, Charles E., and Timothy L. Bailey. 2021. "XSTREME: Comprehensive Motif Analysis of Biological
Sequence Datasets." *bioRxiv*. https://doi.org/10.1101/2021.09.02.458722.
Grant, Charles E., Timothy L. Bailey, and William Stafford Noble. 2011. "FIMO: Scanning for Occurrences
of a given Motif." *Bioinformatics* 27 (7): 1017–18.
Graveley, Brenton. 2010. "The Developmental Transcriptome of Drosophila Melanogaster." *Genome
Biology* 11 (Suppl 1): I11.
Gupta, Shobhit, John A. Stamatoyannopoulos, Timothy L. Bailey, and William Stafford Noble. 2007.
"Quantifying Similarity between Motifs." *Genome Biology* 8 (2): R24.
Haberle, Vanja, Alistair R. R. Forrest, Yoshihide Hayashizaki, Piero Carninci, and Boris Lenhard. 2015.
"CAGEr: Precise TSS Data Retrieval and High-Resolution Promoterome Mining for Integrative
Analyses." *Nucleic Acids Research* 43 (8): e51.
Haines, Jenna E., and Michael B. Eisen. 2018. "Patterns of Chromatin Accessibility along the Anterior-
Posterior Axis in the Early Drosophila Embryo." *PLoS Genetics* 14 (5): e1007367.
Harms, E., T. Chu, G. Henrion, and S. Strickland. 2000. "The Only Function of Grauzone Required for
Drosophila Oocyte Meiosis Is Transcriptional Activation of the Cortex Gene." *Genetics* 155 (4): 1831–
39.
Harrison, Melissa M., Xiao-Yong Li, Tommy Kaplan, Michael R. Botchan, and Michael B. Eisen. 2011.
"Zelda Binding in the Early Drosophila Melanogaster Embryo Marks Regions Subsequently Activated
at the Maternal-to-Zygotic Transition." *PLoS Genetics* 7 (10): e1002266.
Hayashi, Makoto, Yuko Shinozuka, Shuji Shigenobu, Masanao Sato, Michihiko Sugimoto, Seiji Ito, Kuniya
Abe, and Satoru Kobayashi. 2017. "Conserved Role of Ovo in Germline Development in Mouse and
Drosophila." *Scientific Reports* 7 (1): 1–10.
Hiller, M. A., T. Y. Lin, C. Wood, and M. T. Fuller. 2001. "Developmental Regulation of Transcription by a
Tissue-Specific TAF Homolog." *Genes & Development* 15 (8): 1021–30.
Hiller, Mark, Xin Chen, M. Jodeane Pringle, Martin Suchorolski, Yasemin Sancak, Sridhar Viswanathan,
Benjamin Bolival, Ting-Yi Lin, Susan Marino, and Margaret T. Fuller. 2004. "Testis-Specific TAF
Homologs Collaborate to Control a Tissue-Specific Transcription Program." *Development* 131 (21):
5297–5308.
Hinson, S., J. Pettus, and R. N. Nagoshi. 1999. "Regulatory and Functional Interactions between Ovarian
Tumor and Ovo during Drosophila Oogenesis." *Mechanisms of Development* 88 (1): 3–14.
Hubbard, T., D. Barker, E. Birney, G. Cameron, Y. Chen, L. Clark, T. Cox, et al. 2002. "The Ensembl
Genome Database Project." *Nucleic Acids Research* 30 (1): 38–41.
Kent, W. James, Charles W. Sugnet, Terrence S. Furey, Krishna M. Roskin, Tom H. Pringle, Alan M.
Zahler, and David Haussler. 2002. "The Human Genome Browser at UCSC." *Genome Research* 12
(6): 996–1006.
Kim, Daehwan, Joseph M. Paggi, Chanhee Park, Christopher Bennett, and Steven L. Salzberg. 2019.
"Graph-Based Genome Alignment and Genotyping with HISAT2 and HISAT-Genotype." *Nature
Biotechnology* 37 (8): 907–15.
Komitopoulou, K., M. Gans, L. H. Margaritis, F. C. Kafatos, and M. Masson. 1983. "Isolation and
Characterization of Sex-Linked Female-Sterile Mutants in DROSOPHILA MELANOGASTER with
Special Attention to Eggshell Mutants." *Genetics* 105 (4): 897–920.
Lasko, Paul. 2012. "mRNA Localization and Translational Control in Drosophila Oogenesis." *Cold Spring
Harbor Perspectives in Biology* 4 (10). https://doi.org/10.1101/cshperspect.a012294.

Lee, S., and M. D. Garfinkel. 2000. "Characterization of Drosophila OVO Protein DNA Binding Specificity Using Random DNA Oligomer Selection Suggests Zinc Finger Degeneration." *Nucleic Acids Research* 28 (3): 826–34.

Li, Jian, and David S. Gilmour. 2013. "Distinct Mechanisms of Transcriptional Pausing Orchestrated by GAGA Factor and M1BP, a Novel Transcription Factor." *The EMBO Journal* 32 (13): 1829–41.

Lim, Chin Yan, Buyung Santoso, Thomas Boulay, Emily Dong, Uwe Ohler, and James T. Kadonaga. 2004. "The MTE, a New Core Promoter Element for Transcription by RNA Polymerase II." *Genes & Development* 18 (13): 1606–17.

Li, Victor C., Jerel C. Davis, Kapa Lenkov, Benjamin Bolival, Margaret T. Fuller, and Dmitri A. Petrov. 2009. "Molecular Evolution of the Testis TAFs of Drosophila." *Molecular Biology and Evolution* 26 (5): 1103–16.

Li, Xiao-Yong, Melissa M. Harrison, Jacqueline E. Villalta, Tommy Kaplan, and Michael B. Eisen. 2014. "Establishment of Regions of Genomic Activity during the Drosophila Maternal to Zygotic Transition." *eLife* 3 (October): e03737.

Love, Michael I., Wolfgang Huber, and Simon Anders. 2014. "Moderated Estimation of Fold Change and Dispersion for RNA-Seq Data with DESeq2." *Genome Biology* 15 (12): 550.

Lu, Dan, Ho-Su Sin, Chenggang Lu, and Margaret T. Fuller. 2020. "Developmental Regulation of Cell Type-Specific Transcription by Novel Promoter-Proximal Sequence Elements." *Genes & Development* 34 (9-10): 663–77.

Lü, J., J. Andrews, D. Pauli, and B. Oliver. 1998. "Drosophila OVO Zinc-Finger Protein Regulates Ovo and Ovarian Tumor Target Promoters." *Development Genes and Evolution* 208 (4): 213–22.

Lü, J., and B. Oliver. 2001. "Drosophila OVO Regulates Ovarian Tumor Transcription by Binding Unusually near the Transcription Start Site." *Development* 128 (9): 1671–86.

Mahowald, A. P. 2001. "Assembly of the Drosophila Germ Plasm." *International Review of Cytology* 203: 187–213.

Mineo, Alessandro, Marc Furriols, and Jordi Casanova. 2017. "Transfer of Dorsoventral and Terminal Information from the Ovary to the Embryo by a Common Group of Eggshell Proteins in Drosophila." *Genetics* 205 (4): 1529–36.

Nakamura, Shoichi, Seiji Hira, Masato Fujiwara, Nasa Miyagata, Takuma Tsuji, Akane Kondo, Hiroshi Kimura, et al. 2019. "A Truncated Form of a Transcription Factor Mamo Activates Vasa in Drosophila Embryos." *Communications Biology* 2 (1): 1–13.

Ohler, Uwe, Guo-Chun Liao, Heinrich Niemann, and Gerald M. Rubin. 2002. "Computational Analysis of Core Promoters in the Drosophila Genome." *Genome Biology* 3 (12): RESEARCH0087.

Oliver, B., D. Pauli, and A. P. Mahowald. 1990. "Genetic Evidence That the Ovo Locus Is Involved in Drosophila Germ Line Sex Determination." *Genetics* 125 (3): 535–50.

Oliver, B., N. Perrimon, and A. P. Mahowald. 1987. "The Ovo Locus Is Required for Sex-Specific Germ Line Maintenance in Drosophila." *Genes & Development* 1 (9): 913–23.

Omelina, E. S., E. M. Baricheva, D. Yu Oshchepkov, and T. I. Merkulova. 2011. "Analysis and Recognition of the GAGA Transcription Factor Binding Sites in Drosophila Genes." *Computational Biology and Chemistry* 35 (6): 363–70.

Pang, Liang-Yu, Steven DeLuca, Haolong Zhu, John M. Urban, and Allan C. Spradling. 2023. "Chromatin and Gene Expression Changes during Female Drosophila Germline Stem Cell Development Illuminate the Biology of Highly Potent Stem Cells." *eLife* 12 (October): RP90509.

Pauli, D., B. Oliver, and A. P. Mahowald. 1993. "The Role of the Ovarian Tumor Locus in Drosophila Melanogaster Germ Line Sex Determination." *Development* 119 (1): 123–34.

Quinlan, Aaron R., and Ira M. Hall. 2010. "BEDTools: A Flexible Suite of Utilities for Comparing Genomic Features." *Bioinformatics* 26 (6): 841–42.

Ramírez, Fidel, Devon P. Ryan, Björn Grüning, Vivek Bhardwaj, Fabian Kilpert, Andreas S. Richter, Steffen Heyne, Friederike Dündar, and Thomas Manke. 2016. "deepTools2: A next Generation Web Server for Deep-Sequencing Data Analysis." *Nucleic Acids Research* 44 (W1): W160–65.

Raudvere, Uku, Liis Kolberg, Ivan Kuzmin, Tambet Arak, Priit Adler, Hedi Peterson, and Jaak Vilo. 2019. "g:Profiler: A Web Server for Functional Enrichment Analysis and Conversions of Gene Lists (2019 Update)." *Nucleic Acids Research* 47 (W1): W191–98.

R Core Team. 2021. "R: A Language and Environment for Statistical Computing." *R Foundation for Statistical Computing*. https://www.R-project.org/.

Rice, G. A., M. J. Chamberlin, and C. M. Kane. 1993. "Contacts between Mammalian RNA Polymerase II
and the Template DNA in a Ternary Elongation Complex." *Nucleic Acids Research* 21 (1): 113–18.
Salles, Cathy, Maryvonne Mével-Ninio, Alain Vincent, and François Payre. 2002. "A Germline-Specific
Splicing Generates an Extended Ovo Protein Isoform Required for Drosophila Oogenesis."
Developmental Biology 246 (2): 366–76.
Santos, Gilberto dos, Andrew J. Schroeder, Joshua L. Goodman, Victor B. Strelets, Madeline A. Crosby,
Jim Thurmond, David B. Emmert, William M. Gelbart, and FlyBase Consortium. 2015. "FlyBase:
Introduction of the Drosophila Melanogaster Release 6 Reference Genome Assembly and Large-
Scale Migration of Genome Annotations." *Nucleic Acids Research* 43 (Database issue): D690–97.
Smolko, Anne E., Laura Shapiro-Kulnane, and Helen K. Salz. 2018. "The H3K9 Methyltransferase
SETDB1 Maintains Female Identity in Drosophila Germ Cells." *Nature Communications* 9 (1): 1–11.
Soruco, Marcela M. L., Jessica Chery, Eric P. Bishop, Trevor Siggers, Michael Y. Tolstorukov, Alexander
R. Leydon, Arthur U. Sugden, et al. 2013. "The CLAMP Protein Links the MSL Complex to the X
Chromosome during Drosophila Dosage Compensation." *Genes & Development* 27 (14): 1551–56.
Spradling, Allan C., Wanbao Niu, Qi Yin, Madhulika Pathak, and Bhawana Maurya. 2022. "Conservation
of Oocyte Development in Germline Cysts from Drosophila to Mouse." *eLife* 11 (November): e83230.
Ventura, Gemma, Marc Furriols, Nicolás Martín, Vitor Barbosa, and Jordi Casanova. 2010. "Closca, a
New Gene Required for Both Torso RTK Activation and Vitelline Membrane Integrity. Germline
Proteins Contribute to Drosophila Eggshell Composition." *Developmental Biology* 344 (1): 224–32.
Vo Ngoc, Long, George A. Kassavetis, and James T. Kadonaga. 2019. "The RNA Polymerase II Core
Promoter in Drosophila." *Genetics* 212 (1): 13–24.
Xu, Zhe, Hongtao Chen, Jia Ling, Danyang Yu, Paolo Struffi, and Stephen Small. 2014. "Impacts of the
Ubiquitous Factor Zelda on Bicoid-Dependent DNA Binding and Transcription in Drosophila." *Genes
& Development* 28 (6): 608–21.
Zhang, Yong, Tao Liu, Clifford A. Meyer, Jérôme Eeckhoute, David S. Johnson, Bradley E. Bernstein,
Chad Nusbaum, et al. 2008. "Model-Based Analysis of ChIP-Seq (MACS)." *Genome Biology* 9 (9):
R137.
1029

1030

Figure 1: Significantly Enriched OVO DNA Binding Motifs and OVO ChIP Attributes Genome-Wide.
A-C) Immunofluorescent staining of adult ovarioles of $ovo^{Cterm-3xFHA}/ovo^{Cterm-GFP}$ females (20x, scale bar = 20 μm). Ovarioles were stained for Vas (purple) to label the germline, HA (cyan) to label OVO-HA, and GFP (yellow) to label OVO-GFP. The homozygous version of these alleles were used to ChIP OVO. D-G) Significantly enriched motifs found within overlapping OVO ChIP peaks. The percentage of OVO ChIP peaks containing each motif and their corresponding p-value are indicated to the right. H) OVO ChIP minus input control ChIP-seq read coverage density centered on the location of the four *de novo* OVO DNA binding motifs and previously defined *in vitro* OVO DNA binding motifs (Lü et al. 1998, Bielinska et al. 2005, Lee and Garfinkel 2000). I) Relative distance of OVO ChIP peaks to gene level promoters, terminations sequences, and open reading frames genome-wide. J) OVO ChIP minus input control ChIP-seq read coverage density for genes containing significant OVO ChIP peaks and the corresponding OVO DNA binding motif. Genes are centered on the transcriptional start site. K) OVO ChIP minus input control ChIP-seq read coverage density for genes bound by OVO over the TSS, gene body, closest TSS in intergenic space, closest TSS for all, or not bound. Genes are centered on the transcriptional start site. L) OVO ChIP minus input control ChIP-seq heatmaps centered on the dominant significant ovary CAGE-seq TSSs overlapping or not overlapping OVO ChIP peaks.

Figure 2: OVO DNA Binding is Associated with Open Chromatin and Transcriptionally Active
Histone Marks. A, B) OVO ChIP minus input control, GSC and 32c ATAC-seq, GSC H3K27ac,
H3K4me3, H3K27me3, H3K9me3, 8c NC H3K9me3, 32c NC H3K27ac, and H3K27me3 ChIP-seq read
coverage density centered on each OVO peak maximum or OVO DNA binding motif located within a
significant OVO ChIP peak. C) Relative distance of OVO ChIP peaks to significantly called peaks for GSC
and 32c ATAC-seq, GSC H3K27ac, H3K4me3, H3K27me3, H3K9me3, 8c NC H3K9me3, 32c NC
H3K27ac, and H3K27me3 ChIP-seq genome-wide. D) Total number of significant peaks (left) and the
total number of overlapping peaks (top) between OVO ChIP and GSC and 32c ATAC-seq, GSC
H3K27ac, H3K4me3, H3K27me3, H3K9me3, 8c NC H3K9me3, 32c NC H3K27ac, and H3K27me3 ChIP-
seq. Lines connecting solid dots indicates the amount of overlapping peaks between those two
corresponding datasets. Asterisk indicates significantly enriched overlap while hashtag indicates
significantly depleted overlap between datasets.

Figure 3: OVO Bound Promoters are Enriched for INR, DPE, and MTE elements. A) Histogram of the
distance of *in vivo* and *in vitro* OVO DNA binding motifs within significant overlapping OVO ChIP peaks
from the closest genes TSS. B-F) Histogram of the percent of promoters from tissue specific CAGE-seq
analysis of common promoter motif elements centered on the dominant significant TSS.

OVO Ovary RNAseq Samples

A *ovo^{ΔBP}/ovo^{ovo-GAL4}; UASp-GFP*

B *ovo^{ΔBP}/ovo^{ovo-GAL4}; UASp-3xFHA-OVO-B*

20 µm

Vas α-Spec DAPI

C OVO RNA-seq/OVO ChIP MA Plot

log2 Fold Change

log2 Mean Expression

1,328
666

ovo
otu
vas
GAL4
GFP

D OVO RNA-seq/0-2 Hour Old Embryo

log2 Fold Change

log2 Mean Expression

1,409

0.5
-0.5

3,448
625

● Increase expression, OVO not bound
● Increase expression, OVO bound
● Decrease expression, OVO not bound
● Decrease expression, OVO bound

● Increase expression, expressed in embryo
● Decrease expression, expressed in embryo
● Expressed in embryo

E

OVO ChIP | GSC ATAC | GSC H3K27ac | GSC H3K4me3 | GSC H3K27me3 | GSC H3K9me3 | 32c NC ATAC | 32c NC H3K27ac | 32c NC H3K27me3 | 8C NC H3K9me3 | OVO RNAseq

TSS -2Kb | TSS TTS-2Kb | TSS TTS-2Kb | TSS TTS-2Kb | TSS TTS-2Kb | TSS TTS-2Kb | TSS TTS-2Kb | TSS TTS-2Kb | TSS TTS-2Kb | TSS TTS-2Kb | TSS TTS

-1   0   1   2
Normalized Control Subtracted Read Count

1065

Figure 4: Genes Bound by OVO Increase in Expression in the Presence of OVO Genome-Wide. A-B) Immunofluorescent staining of adult ovarioles of the indicated genotypes (20x, scale bar = 20 µm). Ovarioles were stained for Vas (purple) to label the germline, α-Spectrin (yellow) to label dot spectrosome and fusomes, and DAPI (cyan) to label nuclei. Line indicates the dissection point for germarium through previtellogenic RNA-seq samples. C) MA plot of $ovo^{\Delta BP}/ovo^{ovo-GAL4}$; *UASp-3xFHA-OVO-B* versus $ovo^{\Delta BP}/ovo^{ovo-GAL4}$; *UASp-GFP* RNA-seq differential expression results. Purple dots indicate genes that significantly increased in expression and were not bound by OVO, cyan dots indicate genes that significantly increased in expression and were bound by OVO, yellow dots indicate genes that significantly decreased in gene expression and were not bound by OVO, blue dots indicate genes that significantly decreased in gene expression and were bound by OVO, and gray dots indicate genes that were not differentially expressed from our analysis. D) MA plot of $ovo^{\Delta BP}/ovo^{ovo-GAL4}$; *UASp-3xFHA-OVO-B* versus $ovo^{\Delta BP}/ovo^{ovo-GAL4}$; *UASp-GFP* RNA-seq differential expression results. Cyan dots indicate genes that significantly increased in expression and were found to be moderately expressed in 0-2 hour old embryos, yellow dots indicate genes that significantly decreased in gene expression and were found to be moderately expressed in 0-2 hour old embryos, purple dots indicate genes that were not differentially expressed and were found to be moderately expressed in 0-2 hour old embryos, and gray dots indicate genes that were not differentially expressed and were not found to be moderately expressed in 0-2 hour old embryos. E) Gene level read coverage heatmaps of OVO ChIP minus input, GSC and 32c ATAC-seq, GSC H3K27ac, H3K4me3, H3K27me3, H3K9me3, 8c NC H3K9me3, 32c NC H3K27ac, and H3K27me3 ChIP-seq, and $ovo^{\Delta BP}/ovo^{ovo-GAL4}$; *UASp-3xFHA-OVO-B* minus $ovo^{\Delta BP}/ovo^{ovo-GAL4}$; *UASp-GFP* RNA-seq for genes bound by OVO. The order of the heatmap is genes with the highest to lowest amount of OVO ChIP read density.

ovo Locus

Figure 5: OVO ChIP-seq, ATAC/Histone ChIP-seq, RNA-seq, and DNA Binding Motifs at the *ovo*
Locus. *ovo* gene level read coverage tracks for OVO ChIP minus input (black), GSC and 32c ATAC-seq
(light blue), GSC and 32C H3K27ac (green), H3K4me3 (dark blue), GSC and 32c H3K27me3 (orange),
and GSC and 8c H3K9me3 (pink) ChIP-seq, and *ovo^{ΔBP}/ovo^{ovo-GAL4}; UASp-3xFHA-OVO-B* minus
ovo^{ΔBP}/ovo^{ovo-GAL4}; UASp-GFP RNA-seq (red). Red rectangles and black rectangles represent significant
OVO DNA binding motifs and OVO ChIP peaks, respectively. Gene models are represented at bottom.
Small rectangles represent untranslated regions, large rectangles represent translated regions. Arrows
indicate transcriptional start sites.

GO Term Analysis and Representative Gene Tracks

1097

Figure 6: OVO Binds and Significantly Increases the Expression of a Number of Genes Involved in
Essential Maternal Processes. A) Significantly enriched GO biological process terms for genes bound
by OVO and significantly increase in expression in the presence of ectopic rescue OVO. GO terms are
restricted to the 24 smallest term sizes. B-E) Example GO term gene level read coverage tracks for OVO
ChIP minus input and *ovo$^{\Delta BP}$/ovo$^{ovo\text{-}GAL4}$; UASp-3xFHA-OVO-B* minus *ovo$^{\Delta BP}$/ovo$^{ovo\text{-}GAL4}$; UASp-GFP*. Red
rectangles and black rectangles represent significant OVO DNA binding motifs and OVO ChIP peaks,
respectively. Gene models are represented at bottom. Small rectangles represent untranslated regions,
large rectangles represent translated regions. Arrows indicate transcriptional start sites.
1106
Table S1: OVO ChIP-seq Results by Chromosome.
Table S2: Genomic Locations of Significant Overlapping OVO ChIP Peaks
Table S3: Significant OVO DNA Binding Motifs in MEME Format
Table S4: Differential Expression Analysis Results for All Genes
Table S5: Significantly Enriched GO Biological Process Terms
Table S6: Flybase ART Table.

1113

Figure S1: Significant OVO DNA Binding Motif. A) Cartoon of the transcripts expressed from the *ovo* locus based on Benner et al. 2023. Gray box represents endogenously generated deletions, orange triangle represents the location of the *T2A-GAL4-3xSTOP* insertion, purple triangles represent the location of small peptide insertions and yellow boxes represent the location of the shared C_2H_2 zinc-fingers. Small rectangles represent untranslated regions, large rectangles represent translated regions and arrows indicate TSSs. B) Significant alignment of the *in vivo* OVO DNA binding 'motif one' and *in vitro* OVO DNA binding motif (Lee and Garfinkel 2000). C-F) Significantly enriched DNA motifs within OVO ChIP peaks. C is a signficiant match for the DNA binding motif of GAF and CLAMP (Omelina et al. 2011; Soruco et al. 2013). G, H. GSC and 32c ATAC-seq and H3K27ac ChIP-seq minus input control read coverage density for genes bound by OVO over the TSS, gene body, closest TSS in intergenic space, closest TSS for all, or not bound. Plots are centered on the TSS.

1125

Figure S2: OVO DNA Binding is Associated with Open Chromatin and Transcriptionally Active
Histone Marks Across Variations of Gene Binding Patterns. A-D) OVO ChIP minus input control,
GSC and 32c ATAC-seq, GSC H3K27ac, H3K4me3, H3K27me3, H3K9me3, 8c NC H3K9me3, 32c NC
H3K27ac, and H3K27me3 ChIP-seq read coverage density centered on OVO peak maximums located
within a significant OVO ChIP peak for genes bound by OVO over the TSS, gene body, closest TSS in
intergenic space, or closest TSS for all.

A

otu Locus

chr X: 8,486,000 | 8,488,000 | 8,490,000 |

OVO ChIP Peak & DNA Binding Motif

OVO ChIP Minus Input

GSC ATAC

32c NC ATAC

GSC H3K27ac

32c NC H3K27ac

GSC H3K4me3

GSC H3K9me3

8c NC H3K9me3

GSC H3K27me3

32c H3K27me3

UASp-3xFHA-OVO-B Minus *UASp-GFP*

Transcripts

otu-RA

otu-RB

otu-RE

Normalized Read Coverage

1132

Figure S3: OVO ChIP-seq, ATAC/Histone ChIP-seq, RNA-seq, and DNA Binding Motifs at the *otu*
Locus. A) *otu* gene level read coverage tracks for OVO ChIP minus input (black), GSC and 32c ATAC-
seq (light blue), GSC and 32C H3K27ac (green), H3K4me3 (dark blue), GSC and 32c H3K27me3
(orange), and GSC and 8c H3K9me3 (pink) ChIP-seq, and *ovo^{ΔBP}/ovo$^{ovo-GAL4}$*; *UASp-3xFHA-OVO-B*
minus *ovo^{ΔBP}/ovo$^{ovo-GAL4}$*; *UASp-GFP* RNA-seq (red). Red rectangles and black rectangles represent
significant OVO DNA binding motifs and OVO ChIP peaks, respectively. Gene models are represented at
bottom. Small rectangles represent untranslated regions, large rectangles represent translated regions.
Arrows indicate transcriptional start sites.