1 **OVO Positively Regulates Essential Maternal Pathways by Binding Near the Transcriptional Start**
2 **Sites in the *Drosophila* Female Germline**
3

4 Leif Benner[1,2], Savannah Muron[1], Jillian G. Gomez[1], Brian Oliver[1]
5
6 1. Section of Developmental Genomics, Laboratory of Biochemistry and Genetics, National Institute of
7 Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, MD, USA. 2.
8
2. Department of Biology, Johns Hopkins University, Baltimore, MD, USA.

Abstract

Differentiation of female germline stem cells into a mature oocyte includes the expression of a number of
mRNAs and proteins that drive early embryonic development in *Drosophila*. We have little insight into
what activates the expression of these maternal factors. One candidate is the zinc-finger protein OVO.
OVO is required for female germline viability, and has been shown to positively regulate its own
expression, as well as a downstream target, *ovarian tumor* (*otu*), by binding to the transcriptional start site
(TSS). To find additional OVO targets in the female germline and further elucidate OVO's role in oocyte
development, we performed ChIP-seq to determine genome-wide OVO occupancy, as well as RNA-seq
to where OVO is required. OVO preferentially binds in close proximity to target TSSs genome-wide, is
associated with open chromatin, transcriptionally active histone marks, and OVO-dependent expression.
Motif enrichment analysis on OVO ChIP peaks identified a 5`-TAACNGT-3` OVO DNA binding motif near
TSS, but without the precise motif spacing relative to TSS characteristic of RNA Polymerase II complex
binding core promoter elements. Integrated genomics analysis showed that 525 genes that are bound
and increase in expression downstream of OVO are known to be maternally loaded into eggs and early
embryos. These include genes involved in anterior/posterior/germ plasm specification (*bcd, exu, swa,*
osk, nos, pgc, gcl), egg activation (*png, plu, gnu, wisp, C(3)g, mtrm*), translational regulation (*cup*, *orb*,
bru1, me31B), and vitelline membrane formation (*fs(1)N*, *fs(1)M3*, *clos*). This suggests that OVO is a
master transcriptional regulator of oocyte development and is responsible for the expression of structural
components of the egg as well as maternally provided RNAs that are required for early embryonic pattern
formation.

Introduction
Drosophila early embryonic development is directed by events that take place during oogenesis.
Germline stems cells asymmetrically divide to renew the stem cell population and send one daughter cell
towards oogenesis. In the germarium, oogenesis is preceded by four rounds of incomplete mitotic
divisions resulting in a 16 cell egg chamber. One cell is specified as the oocyte which is arrested in
prophase of meiosis I, while the rest of the 15 cells enter endoreplication cycles and become nurse cells
(NCs). Once the 16 cell egg chamber buds from the germarium, the NCs begin to transcribe and translate
a vast array of RNAs and proteins that serve diverse functional roles (Bastock and St Johnston 2008;
Spradling et al. 2022). These roles include positioning maternal mRNAs and proteins in the correct spatial
orientation to support anterior/posterior, and dorsal ventral axis specification, as well negative regulators
of translation to ensure that the maternal mRNAs are not translated before fertilization (Lasko 2012). The
oocyte also contains a number of proteins and mRNAs that are needed for egg activation, completion of
meiosis and initiation of embryonic development after fertilization (Avilés-Pagán and Orr-Weaver 2018). In
order to repeat the process of oogenesis from generation to generation, germ cells in the developing
embryo need to be specified and maintained separately from the rest of the developing somatic cell
population. This requires maternal localization of the germ plasm and early pole cell formation in the
embryo (Mahowald 2001). While thematic elements of this complex orchestration have been well studied,
coordinate regulation of the symphony has not.
50

In particular, maternal control of stepwise embryonic pattern formation during development at the RNA
and protein level has been extensively studied. However, the transcriptional control of these pathways
during oogenesis are less well understood. Few positive regulators of female specific germ cell
transcription have been identified. Genes such as *grauzone* (*grau*) and *maternal gene required for*
meiosis (*mamo*) have been shown to activate the transcription of *cortex* and *vasa*, respectively, in the
female germline (Harms et al. 2000; S. Nakamura et al. 2019). Active repression of male specific

transcription through the activity of *egg*, *wde*, and *Su(var)205 (Smolko, Shapiro-Kulnane, and Salz 2018)*, or global repression of non-ovarian transcriptional networks through the function of *sov (Leif Benner et al. 2019)*, have shown the importance of heterochromatin formation in the female germline for cellular identity and oocyte development. In fact, important recent work has shown the importance of transcriptional repression, mediated through changes in histone modifications, as a key regulator of egg chamber differentiation. GSCs have been shown to exist in a sort of 'ground state' of histone modifications. Characterized with modest non-canonical repressive H3K9me3 and H3K27me3 histone marks at many genes, as well as transcriptionally active H3K27ac histone marks and open chromatin at others (Pang et al. 2023; DeLuca et al. 2020). As oocyte development continues, repressive histone marks associated with heterochromatin begin to increase in abundance resulting in fewer histone marks associated with transcription and open chromatin. This suggests that gene expression becomes more restricted throughout the differentiation process. However, it is unlikely that the female germline directs oocyte development solely through a repressive transcriptional model. Whether the female germline expresses paralogs of the RNA polymerase II complex, like the male germline (M. Hiller et al. 2004; M. A. Hiller et al. 2001; Lu et al. 2020), or if there are pioneering transcription factors involved in determining the open chromatin status for female germ cell-specific expression, or something else entirely, has yet to be determined.

Although few female specific germline transcription factors have been identified, the conserved zinc-finger transcription factor *ovo* has long been known to be required for female germ cell viability. Female germ cells that are *ovo⁻* do not survive into adulthood (Oliver, Perrimon, and Mahowald 1987; Oliver, Pauli, and Mahowald 1990; Benner et al. 2023). Hypomorphic *ovo* alleles, specifically ones that disrupt the transcriptional activator OVO-B, show an arrested egg chamber phenotype, indicating that wild-type OVO-B activity is required for oocyte maturation (Salles et al. 2002; Benner et al. 2023). Germline OVO is expressed at all stages of oogenesis, where it is eventually maternally loaded into the egg. Maternal OVO becomes specifically localized to the developing germline and persists throughout embryogenesis until zygotic OVO is expressed (Hayashi et al. 2017; Benner et al. 2023). Thus OVO is eternally expressed in the female germline, suggesting it may be a key regulator of female specific germline transcription. However, only two downstream targets have previously been identified for OVO. OVO has been shown to positively regulate the expression of its own transcription, therefore executing an autoregulatory loop, as well as positively regulating the transcription of the gene *ovarian tumor* (*otu*)(Lü et al. 1998; Bielinska et al. 2005; Lü and Oliver 2001; Andrews et al. 2000). *otu* is also required in the female germline, where *otu⁻* germ cells show viability and germline tumor phenotypes (Bishop and King 1984). The *ovo* phenotype is epistatic to that of *otu*, and ectopic *otu* expression cannot rescue female germ cell death due to loss of *ovo (Hinson, Pettus, and Nagoshi 1999; Pauli, Oliver, and Mahowald 1993)*. Therefore, *ovo* must be responsible for activating the transcription of genes in addition to *otu* for female germ cell survival and differentiation.

We expanded our knowledge of OVO's role in the female germline by determining genome-wide OVO occupancy and global transcriptional changes downstream of OVO. This allowed us to determine which genes OVO binds and which genes transcriptionally respond to OVO *in vivo*. We show that OVO is directly regulating essential maternal pathways such as axis specification, primordial germ cell formation, egg activation, and maternal mRNA translation regulation. Together, we show that OVO plays a pivotal role in the positive transcriptional regulation of oocyte and early embryonic development. We show that OVO likely carries out this regulation by binding at or in close proximity to the promoters of the genes it regulates and that OVO DNA binding motifs are enriched at or near the transcriptional start site (TSS) of OVO responsive genes, although the spacing of OVO binding sites suggests that it is not a component of the RNA Polymerase complex. OVO binding is also a signature of open chromatin status and active transcription throughout oocyte differentiation. Altogether, we suggest that OVO is required for the activity of a large number of female germline promoters and is likely a key regulator of oocyte maturation and RNAs and proteins that are required for early embryonic development.

Methods

All reagents used in this study can be found in the FlyBase recommended supplementary ART table (Table S1).

113   <u>Fly Husbandry, Transgenesis, and CRISPR/Cas9</u>
114   All fly crosses were conducted at 25°C with 65% relative humidity and constant light unless otherwise
115   noted. Flyfood consisted of premade flyfood (D20302Y) from Archon Scientific (Durham, NC).
116
<u>Immunofluorescence and Image Analysis</u>
Adult females were collected and fed for 3-5 days before ovaries were dissected and fixed in 5.14%
formaldehyde (Pierce, ThermoFisher Scientific) in phosphate buffered saline (PBS, Gibco, ThermoFisher
Scientific) containing 0.1% Triton X-100 (Millipore Sigma)(PBTx) for 15 minutes. Ovaries were then
washed 3 times for 5 min with PBTx followed by a blocking step in PBTx supplemented with 2% normal
goat serum (NGS, Invitrogen, ThermoFisher Scientific) and 0.5% bovine serum albumin (BSA, Millipore
Sigma)(BBTx) for 30 minutes. Primary antibodies were diluted to their appropriate concentrations in BBTx
and ovaries were incubated in primary overnight at 4°C (rabbit anti-Vasa, 1:10K; mouse anti-α-Spectrin,
1:200; rat anti-HA, 1:100; mouse anti-BAM, 1:25; chicken anti-GFP, 1:500; mouse anti-ARM, 1:200; rabbit
anti-MSL-2, 1:5K). The next day, ovaries were washed 3 times for 5 min with PBTx and incubated with
appropriate secondary antibodies diluted (1:500) in BBTx at room temperature for 2 hours (Alexa Fluor
goat anti-rat 488, Alexa Fluor goat anti-chicken 488, Alexa Fluor goat anti-rabbit 488, Alexa Fluor goat
anti-mouse 568, Alexa Fluor goat anti-rat 568, Alexa Fluor goat anti-rabbit 633; Invitrogen,ThermoFisher
Scientific). Ovaries were washed 3 times for 5 min with PBTx before incubation at room temperature in 1
µg/mL DAPI (Invitrogen, ThermoFisher Scientific) solution in PBS for 30 minutes. DAPI was then
removed and ovaries were stored in PBS at 4°C until mounting. To mount, ovaries were transferred to a
microscope slide before adding Ultramount Aqueous Permanent Mounting Medium (Dako, Agilent) and
then coverslipped. Male testes were treated in the same manner, the only difference being that they were
fixed in 4.5% paraformaldehyde in PBTx for 25 minutes. All steps were completed on a rotating nutator at
room temperature unless otherwise noted.
137
138   Ovaries and embryos were imaged on a Zeiss 780 LSM confocal microscope (Carl Zeiss AG) using the
139   Zen Black software (Carl Zeiss AG) . Image analysis was conducted using Fiji(Schindelin et al. 2012).
140
<u>RNA-seq library preparation and sequencing</u>
Twenty 3-5 day old $ovo^{\Delta BP}/ovo^{ovo\text{-}GAL4}$; $UASp\text{-}GFP$ and $ovo^{\Delta BP}/ovo^{ovo\text{-}GAL4}$; $UASp\text{-}3xFHA\text{-}OVO\text{-}B$ ovaries
were dissected and germariums through pre-vitellogenic egg chambers were removed with
microdissection scissors and placed in ice cold PBS making up one biological replicate. RNA was then
extracted from four biological replicates with a Qiagen RNeasy Plus Kit (Qiagen) according to the
manufacturer's protocol, eluted in dH_2O, and RNA concentrations were measured with Quant-iT
RiboGreen RNA Assay Kit (ThermoFisher Scientific). 500 ng of total RNA was then used to make
RNAseq libraries with an Illumina Stranded mRNA Prep Kit according to the manufacturer's protocol
(Illumina). IDT for Illumina RNA UD Indexes Set A were used. Library concentrations were measured with
Quant-iT PicoGreen dsDNA Assay Kit (ThermoFisher Scientific), pooled, and then 50 nucleotide paired-
end DNA sequencing was completed on an Illumina NovaSeq 6000 system using a S1 flow cell (Illumina).
Raw RNAseq reads are available at the SRA under accession (Not uploaded yet).
153
<u>ChIP-seq library preparation and sequencing</u>
Adult $ovo^{Cterm\text{-}3xFHA}$ and $ovo^{Cterm\text{-}GFP}$ females were collected and fed for 24 hours before ovaries were
dissected. 50 dissected ovaries were placed in ice cold phosphate buffered saline (PBS, Gibco,
ThermoFisher Scientific) and then incubated in 1 mL crosslinking solution containing 2% formaldehyde
(Pierce, ThermoFisher Scientific) (50mM HEPES Buffer, 1mM EDTA, 0.5 mM EGTA, 100mM NaCl), and
rotated at 37°C for 20 minutes. Ovaries were then incubated in 1 mL stop solution (125mM Glycine,
0.01% Triton X-100 (Millipore Sigma), diluted in PBS) and rotated for 5 minutes at room temperature.
Ovaries were then washed twice with 1 mL ice cold wash buffer (0.01% Triton X-100 in PBS) for 5
minutes. The last wash was removed and ovaries were stored at -80°C until future processing. Once all
samples were collected, 4x50 ovaries were then homogenized in 250 µL RIPA lysis buffer (Pierce,
ThermoFisher Scientific) containing 1x protease inhibitor cocktail (cOmplete Mini Protease Inhibitor
Cocktail, Roche, Millipore Sigma) and 1 mM PMSF (Roche, Millipore Sigma) and kept on ice for 10
minutes. 40mg of 212-300 µm acid-washed glass beads (Millipore Sigma) were then added to
homogenized ovary lysate. Samples were then sonicated with a Bioruptor Pico sonication device
(Diagenode) at 4C for 15 cycles of 30 seconds on and 30 seconds off. Sonicated lysate was then

transferred to a new tube and centrifuged at 13,300 rpm for 10 minutes at 4°C. Three supernatants were
then combined to form one biological replicate. 100 µL for each biological replicate was removed and
stored at -80°C for input control. To pull down C-terminally tagged OVO, 100 µL of monoclonal anti-HA-
agarose (Millipore Sigma) or 50 µL of ChromoTek GFP-Trap agarose (Proteintech) were washed three
times with RIPA lysis buffer and spun down at 1,200 RPMs for one minute at 4°C. 550 µL of $ovo^{Cterm-3xFHA}$
supernatant was added to monoclonal anti-HA-agarose and 550 µL of $ovo^{Cterm-GFP}$ supernatant was added
to ChromoTek GFP-Trap agarose. Samples were supplemented with 1x protease inhibitor cocktail and 1
mM PMSF and incubated on a rotator at 4°C overnight.
177
The next day, agarose was washed in a stepwise fashion with solutions from a Chromatin
Immunoprecipitation Assay Kit (Millipore Sigma), beginning with 1 mL of a low salt wash buffer, high salt
wash buffer, LiCl buffer, and ending with 2 washes in 0.1x TE buffer. 300 µL of freshly prepared ChIP
elution/decrosslinking solution (1% SDS, 100mM $NaHCO_3$, 250 mM NaCl, 10mM EDTA, 50 mM Tris-HCl,
200 µg/mL Proteinase K) was added to the pelleted agarose, or 200 µL of chip elution/decrosslinking
solution was added to 100 µL input control, and incubated at 65°C overnight. DNA was extracted by
adding 300 µL phenol:chloroform:iso-amyl alcohol (125:24:1) (Millipore Sigma). The samples were
vortexed for 30 seconds then centrifuged at 13,300 RPMs for 5 minutes at 4°C. The aqueous layer was
extracted and this process was repeated once more. 1 µL glycogen (20 mg/mL), 30 µL 1M sodium
acetate, and 750 µL 100% EtOH was added to the extracted aqueous layer, vortexed, and incubated at -
20°C for 30 minutes. Solution was spun at 13,300 RPMs for 20 min at 4°C. Supernatant was removed
and the pellet was washed with 500 µL 70% EtOH and spun down at 13,300 RPMs for 20 minutes at 4°C.
This step was repeated but with 100% EtOH. The resulting pellet was briefly speedvacced and
resuspended in 50 µL dH_2O.
192
To make ChIP-seq libraries, DNA concentration for immunoprecipitated and input control samples were
measured with a Quant-iT PicoGreen dsDNA Assay Kit (ThermoFisher Scientific). 5 ng of DNA for each
sample was then used with the NEBNext Ultra II DNA Library Prep Kit for Illumina (New England Biolabs)
and completed according to the manufacturer's protocol. ChIP-seq library concentrations were then
measured with a Quant-iT PicoGreen dsDNA Assay Kit, pooled, and then 50 nucleotide paired-end DNA
sequencing was performed on an Illumina NovaSeq 6000 system using the XP workflow (Illumina). Raw
ChIP-seq reads are available at the SRA under accession (Not uploaded yet).
200
RNA-seq, ChIP-seq, CAGE-seq and Gene Ontology Analysis
For RNA-seq analysis of $ovo^{ΔBP}/ovo^{ovo-GAL4}$; $UASp-GFP$ and $ovo^{ΔBP}/ovo^{ovo-GAL4}$; $UASp-3xFHA-OVO-B$
ovaries, 50 nucleotide paired-end reads were mapped to the FlyBase r6.46 genome (Gramates et al.
2022) for differential expression analysis and the BDGP Release 6 Drosophila Genome (dos Santos et al.
2015) for read level genome browser tracks using Hisat2 (-k 1 --rna-strandness RF --dta)(Kim et al.
2019). DNA sequences for $GAL4$ and GFP were added to the FlyBase r6.46 genome as separate
chromosomes. Mapped reads were then sorted and indexed with Samtools (samtools sort and samtools
index)(Danecek et al. 2021). Gene level readcounts were then derived with htseq-count (-s reverse -r
pos)(Anders, Pyl, and Huber 2015) and used for differential expression analysis with DeSeq2 (Love,
Huber, and Anders 2014). Genes with 0 mapped reads were removed from the DESeq2 analysis.
211
For ChIP-seq analysis of OVO-HA, OVO-GFP, OVO-HA input, and OVO-GFP input samples, 50
nucleotide paired-end reads were mapped to the FlyBase r6.46 genome for peak calling analysis and the
BDGP Release 6 Drosophila Genome for read level genome browser tracks using Hisat2 (-k 1 --no-
spliced-alignment -X 900). Mapped reads were sorted using Samtools (samtools sort and samtools index)
and duplicate reads were removed with Picard (REMOVE_DUPLICATES=true)(broadinstitute, n.d.).
Significant ChIP peaks were called for OVO-HA and OVO-GFP versus their respective input controls
separately using Macs3 callpeak software (-g 1.2e8 -q 0.0001)(Zhang et al. 2008). Overlapping ChIP
peaks for OVO-HA and OVO-GFP were then determined with bedtools intersect software (Quinlan and
Hall 2010). Peak calling for GSC ATAC-seq (SRR24203655), 32c ATAC-seq (SRR24203650), GSC
H3K27ac (SRR11084657), H3K4me3 (SRR11084658), H3K27me3 (SRR11084656), H3K9me3
(SRR24203631), 8c NC H3K4me3 (SRR24203629), 32c NC H3K27ac (SRR24203635), and H3K27me3
(SRR11084652) ChIP-seq versus their respective input controls (SRR11084655, SRR11084651,
SRR24203634, SRR24203637) was conducted in the same manner as OVO ChIP-seq.

225
226 In order to generate gene level read coverage tracks, deepTools' bamCompare software was used to
227 generate a single bigWig file comparing all replicates versus input controls (-bs 5 --effectiveGenomeSize
228 142573017 --normalizeUsing BPM --exactScaling --scaleFactorsMethod None)(Ramírez et al. 2016). The
229 bigWig file was then uploaded to UCSC genome browser for visualization (Kent et al. 2002).
230
To generate read coverage plots centered on the motif location or OVO peak maximums, genomic
locations of significant scoring motifs or peak maximums within overlapping OVO ChIP peaks were
determined and used as input for deepTools' computeMatrix reference-point (-a 2000 -b 2000 -bs 25 --
missingDataAsZero). Read density profiles for each motif or OVO peak maximum were then visualized
with deeptools plotProfile. In order to generate read coverage plots centered on the TSS, the same
methods as above were conducted except genes overlapping OVO ChIP peaks containing the respective
significant OVO DNA binding motifs were used as input instead.
238
239 In order to generate gene level read coverage heatmaps, deepTools' computeMatrix scale-regions
240 software was used to generate a single matrix for genes that were bound by OVO (-bs 25 --
241 missingDataAsZero -m 4000 --metagene). This matrix was then used as input for deepTools'
242 plotHeatmap software to generate heatmaps of ChIP-seq and RNA-seq for the given OVO binding
243 profiles centered on the TSS (--sortUsing max).
244
For CAGE-seq analysis, CAGE-seq libraries for ovary (SRR488283, SRR488282), testes (SRR488284,
SRR488308, SRR488285, SRR488309), and male and female digestive system (SRR488289,
SRR488288) tissues were downloaded and combined for each tissue type from the SRA. Reads were
mapped to the BDGP Release 6 Drosophila Genome with Hisat2 (-k 1). Mapped reads were sorted with
Samtools. Significant dominant TSSs were then determined with CAGEr software(Haberle et al. 2015)
from sorted BAM files with getCTSS and annotated with annotateCTSS using the dm6.ensGene.gtf
file(Hubbard et al. 2002) downloaded from UCSC(Kent et al. 2002). CAGE-seq reads were normalized
with normalizeTagCount (ce, method = "simpleTpm", fitInRange = c(5, 40000), alpha = 1.15, T = 1*10^6)
and then TSS clusters were determined with clusterCTSS (ce, threshold = 1, thresholdIsTpm = TRUE,
nrPassThreshold = 1, method = "paraclu", maxDist = 20, removeSingletons = TRUE,
keepSingletonsAbove = 5) in order to determine the dominant significant TSS for each respective tissue.
256
257 Gene ontology enrichment analysis was completed with g:Profiler's g:GOSt software (Raudvere et al.
258 2019) on the set of genes overlapping OVO ChIP peaks over the TSS, significantly upregulated in the
259 presence of OVO, and contained significant OVO DNA binding motifs. Default parameters were used for
260 the enrichment analysis and only GO biological process terms were searched for enrichment with the
261 gene list.
262
263 Fisher's exact test was conducted for each respective analysis with the fisher.test() command in R (R
264 Core Team 2021).
265
de novo Motif Enrichment and Promoter Motif Analysis
DNA sequences from significant overlapping OVO ChIP peaks were extracted from the Drosophila r6.46
genome and submitted to STREME software (Bailey 2021) of the MEME suite (Bailey et al. 2015). The
default parameters were used for *de novo* motif enrichment analysis. After identifying 'OVO Motif One',
OVO ChIP peaks that contained that sequence were removed and the resulting ChIP peaks were
resubmitted for STREME analysis deriving derivative OVO DNA binding motifs. Significant OVO DNA
binding motifs and *in vitro* OVO DNA binding motifs were searched in the BDGP Release 6 Drosophila
Genome using FIMO (Grant, Bailey, and Noble 2011). In order to find significant DNA binding motif
matches for 'OVO Motif One', this motif from STREME was submitted to Tomtom software (Gupta et al.
2007) of the MEME suite and searched within the JASPAR Core Insect database (2022)(Castro-
Mondragon et al. 2022).
277
Promoter motif analysis was conducted by extracting the DNA sequences 200 nucleotides upstream and
downstream of the significant dominant TSSs from CAGE-seq analysis for each respective tissue type. All
common core promoter motifs (FitzGerald et al. 2006; Ohler et al. 2002) were then searched in these

sequences depending on their strand specificity with the use of FIMO from the MEME suite using a p-
value of < 0.003 for all non-OVO promoter motifs. All OVO motifs found in this study and through *in vitro*
methods were also searched with the same method, except a p-value of < 0.0002 was used.
284
Results
<u>OVO Binds Promoters Genome Wide</u>
OVO-B, the predominant protein isoform expressed from the *ovo* locus in the female germline (Benner et
al. 2023), is a positive regulator of transcription (Andrews et al. 2000). OVO-B positively regulates the
gene expression of *otu* and *ovo* transgenic reporter constructs require OVO binding sites both at and
upstream of the TSS in order to recapitulate full reporter expression (Bielinska et al. 2005; Lü et al. 1998;
Lü and Oliver 2001). Females hemizygous for antimorphic dominant gain-of-function (*ovo^D*) or
homozygous recessive (*ovo^{D1rv}*) *ovo* alleles lack germ cells in the adult ovary (Oliver, Perrimon, and
Mahowald 1987; Benner et al. 2023). True OVO-B null alleles created by deletion of the *ovo-B* promoter
have the same germ cell loss phenotype (Benner et al. 2023). The phenotypes of *otu* females range from
germ cell death to ovarian tumors depending on the allele and undefined stochastic factors (Bishop and
King 1984). It is possible that the germ cell death phenotype in *ovo⁻* female germ cells can solely be
explained by failure of OVO to activate *otu* expression, however, this is highly unlikely. The *ovo^{D1rv}*
phenotype is epistatic to *otu⁻*, and ectopic *otu⁺* expression in *ovo⁻* germ cells does not rescue the germ
cell death phenotype (Hinson, Pettus, and Nagoshi 1999). This is not terribly surprising, as we know of no
examples of a transcription factor that regulates a single gene. This suggests that OVO regulates the
expression of additional genes in the female germline.
302
We wanted to identify the full stable of OVO target genes in the female germline by using two
complementary genome-wide approaches to test for OVO presence and function. Specifically, we
determined OVO occupancy genome-wide with ChIP-seq, and determined *ovo* function by comparing the
RNA expression profiles between *ovo⁺* and *ovo* hypomorphs in the female germline. In order to determine
OVO occupancy genome-wide, we performed ChIP-seq on newly eclosed adult ovaries in triplicate, using
two C-terminally tagged alleles as affinity purification tools (*ovo$^{Cterm-3xFHA}$* and *ovo$^{Cterm-GFP}$*, Figure 1A-
C)(Benner et al. 2023). We compared immuno-purified OVO associated DNA with input DNA as a control,
for a total of 12 ChIP-seq libraries, which we sequenced using the Illumina system. After quality control
and alignment to the *Drosophila* r6.46 genome (Gramates et al. 2022), we used MACS3 (Zhang et al.
2008) to call significantly enriched peaks from OVO-HA and OVO-GFP compared to their respective input
controls (see methods).
314
We first compared the pulldown results with the OVO-HA versus OVO-GFP ChIP reagents. The GFP pull
down appeared to be more efficient, but never-the-less we found excellent agreement, as most OVO-HA
peaks were also found in the OVO-GFP dataset. The OVO-GFP ChIP dataset had 7,235 significant ChIP
peaks according to peak enrichment analysis derived from MACS3 genome-wide, while OVO-HA ChIP
dataset had 3,393 significant peaks genome-wide (Table S2). To determine the similarity in significant
peak calling between the two datasets, we calculated a Jaccard index (intersection/union) between the
significantly enriched peaks from the tagged *ovo* allele bearing ovaries. The Jaccard index between OVO-
HA and OVO-GFP ChIP peaks was 0.64 (where 0 = no overlap and 1 = full overlap) with a total of 3,094
ChIP peaks overlapping. Thus, almost all significant OVO-HA ChIP peaks were also found within the
OVO-GFP ChIP dataset (91% of OVO-HA peaks overlapped OVO-GFP peaks). OVO-GFP pulldown was
either more effective, or less likely, promiscuous. Regardless, we decided to use the conservative
intersection of the two datasets (3,094 peaks) for downstream OVO occupancy informatics (Table S3).
327
OVO is a sequence-specific DNA-binding protein, but many transcription factors also have a more
general affinity for DNA, additionally, immunoprecipitation can capture indirect interactions due to nuclear
topology in addition to direct sequence-specific binding. For example, in the particular case of OVO TSS
binding, this could be direct, as shown in the case of *ovo-B* and *otu* loci (Lü et al. 1998; Lü and Oliver
2001; Bielinska et al. 2005), or could be due to looping of an OVO-bound enhancer to the core promoter.
Determining if there are canonical OVO binding sites at peaks can help to distinguish direct and indirect
binding. If OVO is directly binding to the TSS, we would expect to find OVO binding site enrichment at
that location. To examine the sequences enriched in peaks, we looked directly for the known OVO
binding sites previously defined by footprinting (Lü et al. 1998) and SELEX-seq ((Bielinska et al. 2005;

Lee and Garfinkel 2000). We also did *de novo* motif finding on this substantial dataset to refine the
sequence specific motifs bound by OVO and perhaps other motifs associated with other transcription
factors preferentially bound by OVO enhancers, or OVO proximal sites in 3D nuclear space. We
performed novel motif enrichment analysis using STREME (Bailey 2021) with our overlapping ChIP peaks
and found a number of significant motifs within our dataset. The most significant motif, 5`-TAACGGTAAA-
3`, was found in 50% of significant ChIP peaks (Figure 1D, 'Motif One'). This motif is highly similar to the
OVO DNA binding motif that has been reported twice before, 5`-AGTAACNGT-3` (SELEX method,
'Garfinkel OVO Motif') and 5`-TGTAACNGT-3` (Footprinting method, 'Oliver OVO Motif'). The only
differences between motif one in our dataset and the literature, is that the *de novo* motif is two
nucleotides shorter than the previously described motifs at the 5` end, extends 3 nucleotides
downstream, and includes a second G near the 3'-end (Figure S1A). Collectively, this is strong evidence
that the core OVO binding sequence is 5`-TAACGGT-3`. Some binding sites can be recognized by
multiple transcription factors. To determine if other characterized transcription factors recognize this
sequence, we searched for significant matches to motif one in the Jaspar database (Castro-Mondragon et
al. 2022) of known *Drosophila* motifs using Tomtom (Gupta et al. 2007). The OVO DNA binding motif
('Garfinkel OVO Motif') was scored as a significant match (Figure S1A, $p<0.05$). While there may well be
other TFs binding near OVO, we did not identify them based on known binding site data.
354
Not every peak region had one of these consensus OVO motifs. This does not mean *a priori* that they
bound OVO indirectly. Motif enrichment can be driven by a few strongly enriched sequences, so that
more minor enrichments of variants are missed. Therefore, we carefully examined the 50% of our
overlapping OVO ChIP peaks where motif one was not found. This second round of *de novo* analysis
revealed enrichment of OVO DNA binding motif derivatives. For example, the third most significant motif
(found in 37% of peaks) was 5`-RWMTAACGGV-3` (Figure 1E, motif two). This motif had the core 5`-
TAACGGT-3` sequence found in all three aforementioned methods, however, the last nucleotide in the
core sequence is unspecific and lacks the three 3` nucleotides found in motif one. Two other derivative
motifs, 5`-TAACTGTTTT-3' (found in 17% of sequences, Figure 1F, motif three), and 5'-TTACSGTAA-3'
(found in 5% of sequences, Figure 1G, motif four), vary within the central core motif (at positions 5 and 2
of the core sequence, respectively) and upstream and downstream ends. Searching for all variations of
the OVO DNA binding motif (Table S4) within our significant overlapping ChIP peaks indicated that 72%
of peaks contained at least one variation of these four binding motifs. It is a reasonable hypothesis that
OVO peaks are most often due to direct, rather than indirect OVO binding.
369
A prediction for direct OVO binding to motifs, is that the motif should be centered within the peak of
fragmented input DNA sequences. Therefore, we plotted the significant ChIP (minus input) read density
peaks centered on the location of the motif itself. We found that the read density for all ChIP peaks that
contain each one of the *de novo* OVO motifs, as well as the *in vitro* OVO motifs, are centered over the
motif location (Figure 1H). This suggests that all of these motifs from our analysis are bonafide OVO DNA
binding sites *in vivo*. While it is possible that OVO comes into contact with regions of DNA in three
dimensional nuclear space non-specifically, the presence of OVO motifs within a large percentage of
significant ChIP peaks *in vivo*, and the ChIP peak read densities centered over the location of the motifs,
strongly reinforces the idea that our dataset contains regions centered on sequence-specifically bound
OVO transcription factor in the ovary.
380
Given the clear function of OVO occupancy near the TSS of its two known targets, *ovo* and *otu* (Lü et al.
1998; Bielinska et al. 2005; Lü and Oliver 2001), we were interested in determining if OVO peaks are
generally near the TSS of other target genes as well. In addition to informing the biology of OVO function,
this simplifies the problem of associating peaks to potential functional target genes. As a preliminary test
of this idea, we determined if the fully overlapping OVO-HA and OVO-GFP peaks were spatially enriched
with respect to the currently annotated gene model elements such as TSS, open reading frames (ORFs),
or transcription termination sequences (TTS). If the TSS association of OVO at the two known targets
reflects a general propensity, then we expected OVO ChIP peaks to be more closely associated with TSS
than other gene elements. To carry out this analysis, we normalized the genome for these three gene
elements, such that the distance between adjacent loci was 1. If there is no enrichment for OVO peaks to
a specific gene element, then the peak location would have an equal frequency from 0.0 to 0.5 relative
distance. Measuring the relative distance of our OVO ChIP peaks to TSS, ORFs, and TTS, showed that

the OVO binding was highly enriched near TSS/promoter locations and was not correlated with ORF and
TTS locations (Figure 1I). These results confirmed that OVO is characterized by core promoter proximal
binding. Since OVO ChIP peaks as a class are associated with TSS, we plotted the ChIP minus input
read density of genes that overlap significant ChIP peaks to examine the full distribution. We found that
the OVO ChIP read density was highly enriched over the TSS and was not due to a few highly enriched
examples (Figure 1J). This builds on the idea that OVO is binding directly over, or in close proximity to the
TSS of it's target genes genome-wide. In other words, the previous work showing OVO binding the *ovo-B*
and *otu* TSS (Bielinska et al. 2005; Lü et al. 1998; Lü and Oliver 2001) is typical. This very specific
binding of OVO to the TSS is intriguing and unusual, as this region associates with the basal
transcriptional machinery. It raises the possibility that OVO is not a typical transcription factor that acts
primarily via enhancer binding, but might be part of the core promoter binding complex or acts to
precondition the core promoter region for example.

The OVO ChIP read density was highly enriched over the annotated TSS of target genes, but TSS
annotation is challenging and can be tissue specific. We were interested in empirically determining if the
same enrichment was present in TSSs utilized specifically in ovarian tissue. In other words, where exactly
does transcription start at these genes. The 5' ends of mRNA are capped. In order to determine where
these caps mapped to the genome, we analyzed used Cap Analysis of Gene Expression (CAGE-seq)
data from adult *Drosophila* ovaries (SRR488283, SRR488282)(Chen et al. 2014) with CAGEr software
(Haberle et al. 2015) and extracted the dominant significant TSSs in the ovary. CAGEr predicted 6,856
significant TSSs in the ovary dataset, of which 1,047 overlapped with OVO ChIP peaks. We plotted the
OVO ChIP minus input read density centered on the significant ovary CAGE-seq TSSs for TSSs that
overlapped or did not overlap OVO ChIP peaks (Figure 1K, 1L). We found that OVO ChIP read density
was highly enriched over the location of TSSs from ovary CAGE-seq that overlapped OVO ChIP peaks
when compared to TSSs that did not overlap OVO ChIP peaks. Thus, OVO TSS binding is not due to
poor annotation of ovarian TTSs. Furthermore, OVO is binding at or near TSS of genes actively being
transcribed in the ovary.

<u>OVO Binding is Associated with Open Chromatin and Transcriptionally Active Histone Marks</u>
Our OVO ChIP data indicated that OVO was binding at or in close proximity to promoters genome-wide.
OVO could have positive and/or negative effects on transcription at that location. For example, OVO
could help recruit or sterically hinder RNA Polymerase binding to TSSs. Previous OVO reporter
constructs show positive effects of OVO binding near TSS (Lü and Oliver 2001; Lü et al. 1998; Bielinska
et al. 2005). If OVO binding is generally promoting transcription, then we hypothesize that it would be
more closely associated with histone marks associated with active transcription, such as H3K27ac and
H3K4me3, as well as lower nucleosome density that can be measured through ATAC-seq. In contrast,
OVO binding would be expected to negatively correlate with repressive H3K9me3 and H3K27me3
histone marks and higher nucleosome density. It is technically difficult to determine changes in chromatin
status and transcription in germ cells that lack OVO, as the phenotype is cell death (although we will
return to this later for transcription profiling), but analyzing OVO binding in the context of ovarian
chromatin was highly informative.

Recent work profiling nucleosome density and histone marks have shown that female GSCs have a
'ground state' chromatin profile (DeLuca et al. 2020), similar to the histone mark profiles that are found in
early embryos (X.-Y. Li et al. 2014). This has been characterized to contain non-canonical H3K27me3
profiles and low H3K9me3 histone levels (Pang et al. 2023; DeLuca et al. 2020). As egg follicles
differentiate, nurse cells begin to accumulate H3K9me3 marks, and H3K27me3 histone marks begin to
accumulate over more traditional polycomb domains. This in turn leads to a decrease in the number of
open chromatin peaks as well as H3K27ac histone marks, which are generally associated with active
transcription (Pang et al. 2023; DeLuca et al. 2020). Essentially, these data support the idea that egg
chambers restrict gene expression competency as they differentiate.

In order to determine the relationship in our OVO ChIP data and other chromatin marks, we analyzed
GSC H3K27ac, H3K27me3, H3K9me3, H3K4me3, and ATAC-seq data (Pang et al. 2023; DeLuca et al.
2020) with the same parameters used to establish significant OVO peaks in our OVO ChIP dataset. Our
OVO ChIP data was from one day old ovaries and we did not profile specific follicle stages. So we also

analyzed 32c (roughly stage 5 egg chambers) H3K27ac, H3K27me3, ATAC-seq, and 8c H3K9me3 (32c
was not available) histone marks (Pang et al. 2023; DeLuca et al. 2020) to see if there were any stage
specific differences in comparison to OVO DNA binding. We first plotted the read density of each
respective chromatin mark minus their input control centered on either the OVO ChIP peak local
maximum (Figure 2A) or OVO DNA binding motifs (Figure 2B). GSC ATAC and H3K27ac read density
showed a high degree of enrichment over OVO ChIP peak maximums (Figure 2A) and OVO DNA binding
motifs (Figure 2B), consistent with positive transcriptional activity. GSC H3K4me3 read density was, to a
lesser extent, also enriched with OVO ChIP peak maximums and OVO DNA binding motifs. However,
there was no read density enrichment for repressive GSC H3K27me3 and H3K9me3 histone marks.
Since there was a high degree of read density enrichment between OVO ChIP and other chromatin
marks/low nucleosome density we wanted to determine the extent of the overlap between significant
OVO ChIP peaks and significantly called peaks from the different types and stages of histone marks and
ATAC-seq data. In order to do this, we measured the relative distance of OVO ChIP peaks to the same
datasets described above. We found that OVO ChIP peaks had a lower relative distance, and thus were
spatially overlapping/closer in the genome, to 32c NC ATAC, GSC ATAC, GSC H3K27ac, GSC
H3K4me3, and 32c NC H3K27ac peaks, in that order (Figure 2C). While the relative distance between
OVO ChIP peaks and H3K9me3 and H3K27me3, regardless of stage, showed no spatial association.
There was also further support for this association with transcriptionally active histone marks/open
chromatin when measuring the overlap between significant OVO ChIP peaks and the respective
significant histone ChIP and ATAC-seq peaks (Figure 2D). A Fisher's exact test indicated a significant
enrichment in overlapping peaks genome-wide between OVO and GSC ATAC ($p < 0.001$, odds ratio =
75.9), 32c NC ATAC ($p < 0.001$, odds ratio = Infinite, i.e. all OVO peaks overlapped ATAC-seq peaks),
GSC H3K27ac ($p < 0.001$, odds ratio = 31.7), GSC H3K4me3 ($p < 0.001$, odds ratio =12.0), and 32c NC
H3K27ac ($p < 0.001$, odds ratio = 7.9) peaks. While there was a significant depletion in overlapping peaks
genome-wide between OVO and 32c NC H3K27me3 ($p < 0.001$, odds ratio = 0.6), GSC H3K9me3 ($p <
0.001$, odds ratio = 0.7), and 8c NC H3K9me3 ($p < 0.001$, odds ratio = 0.5). Altogether, suggesting that
OVO binding genome-wide is tightly associated with open chromatin regardless of germ cell stage, and
active transcription in GSCs. In other words, chromatin state data suggests OVO is acting positively on
it's target genes and raises the possibility that OVO-binding and open chromatin precede transcription.
478
479
<u>OVO DNA Binding Motifs are Evenly Distributed Around Promoters and are Enriched for INR, DPE, and
MTE Elements</u>
Our data thus far clearly indicates that OVO binding occurs at or very near the core promoter, a region
recognized by an enormous collection of factors that associate with RNA polymerase to initiate
transcription (Aoyagi and Wassarman 2000; Vo Ngoc, Kassavetis, and Kadonaga 2019). If these
promoter proximal OVO sites are functional, they could be part of that complex, or could help facilitate the
binding of that complex. The highly organized polymerase complex has sequence-specific DNA
recognition sites with incredibly precise spacing between them, with an overall DNA footprint of a little
less than 100bp (Rice, Chamberlin, and Kane 1993; FitzGerald et al. 2006; Ohler et al. 2002). If OVO
binding sites at core promoters share this precision in spacing with other core promoter elements, then it
is likely to be part of the complex. If not, then OVO is more likely to facilitate binding of the basal
transcriptional machinery. Because of the extended footprint of engaged RNA polymerase, OVO and the
basal machinery would not be likely to occupy the same region at the same time.
493
Core promoters have sets of sequences with tightly constrained inter-motif distances that bind
components of the transcription initiation complex. There are upstream binding sites such as TATA, sites
at transcription start, such as the initiator (INR), and downstream promoter elements (DPE) (Vo Ngoc,
Kassavetis, and Kadonaga 2019). The combinations of these elements is not random in mammals and
Drosophila (FitzGerald et al. 2006), and the families of motif combinations at the TSS of genes expressed
in *Drosophila* are conserved over tens of millions of years of evolution (Chen et al. 2014). Like OVO chip
peaks, OVO DNA binding motifs were highly enriched at or near the TSS (Figure 3A). We carefully
analyzed the spacing of these sites relative to core promoter elements to see if spacing was precise at
the nucleotide level.
503

We first searched for the presence of previously defined DNA motifs that are enriched at promoters
(FitzGerald et al. 2006; Ohler et al. 2002; Lim et al. 2004) using FIMO (Grant, Bailey, and Noble 2011).
We defined promoters by using the DNA sequences 150 nucleotides upstream and downstream of the
significant dominant TSSs in our previously analyzed ovary CAGE-seq datasets (Chen et al. 2014). After
extracting these sequences and searching for significant scoring motifs, we plotted the density of each
motif in relation to the empirically mapped TSSs (Figure 3B). We also searched for all OVO motifs found
in our significant ChIP peaks within these promoter sequences. When plotting the density of DNA motifs
found in ovary CAGE-seq promoters, we found that there were prominent peaks for INR and M1BP
(M1BP (J. Li and Gilmour 2013) = Ohler 1 (Ohler et al. 2002) = DMv4 (FitzGerald et al. 2006)) near the
TSS, and MTE (Lim et al. 2004) and DPE elements downstream of the TSS. This distribution and
frequency is consistent with the constrained location of these DNA motifs (FitzGerald et al. 2006; Ohler et
al. 2002; Chen et al. 2014). Significantly, the OVO DNA binding motifs showed a broad distribution
upstream and downstream of the TSS.
517
The precise core promoter architecture of OVO bound TSSs is revealed in the CAGE-seq dataset.
Plotting the distribution of classical core promoter sequence elements in OVO bound promoters showed a
similar, but exaggerated, profile compared to all care promoters of the ovary CAGE-seq dataset. We
found a significant enrichment for INR ($p < 0.01$, odds ratio = 1.70), DPE ($p < 0.01$, odds ratio = 1.81),
MTE ($p < 0.01$, odds ratio = 1.65), and most importantly, OVO DNA binding motifs ($p < 0.01$, odds ratio =
4.83), in ovary promoters that overlapped an OVO ChIP peak in comparison to the subset of ovary
promoters that did not overlap an OVO ChIP peak (Figure 3C, 3D). This indicates that OVO bound
promoters are more likely to contain these specific promoter elements than non-OVO bound promoters.
As has been described before, promoters containing INR and DPE, but lacking TATA-box elements, are
common among *Drosophila* gonad promoters compared to promoters of other tissue types (Chen et al.
2014). The presence of TATA-box elements is negatively associated with germline-specific gene
expression (FitzGerald et al. 2006). We found that TATA-box elements were significantly depleted in
ovary CAGE-seq promoters when compared to testes ($p < 0.01$, odds ratio = 0.78)(Figure 3E) or digestive
system ($p < 0.01$, odds ratio = 0.50)(Figure 3F) CAGE-seq promoters. Indeed, both *ovo* and *otu* have
TATA-less promoters. Briefly, OVO bound promoters are characterized by the presence of INR, DPE,
MTE, and, of course, OVO DNA binding motifs. This could represent a functional class of promoters
utilized for gene expression in the *Drosophila* ovary. Importantly, the distribution of OVO DNA binding
motifs in ovary promoters is not fixed relative to TSSs or other core promoter elements. Thus, it is highly
unlikely that OVO acts as a female germline RNA Polymerase complex member that anchors the complex
to the core promoter and helps determine the +1 mRNA nucleotide. Rather, the imprecise location of
OVO binding sites might suggest that OVO is more likely to facilitate the binding of other basal
transcriptional factors.
540
<u>OVO Activates Gene Expression in the Female Germline</u>
Occupancy is a requirement for activity, but occupancy does not equal activity. Understanding the
transcriptional consequences of OVO occupancy genome-wide would allow us to investigate
mechanisms. However, as we mentioned earlier, the fact that *ovo* is absolutely required for female
germline viability greatly complicates this analysis. Measuring gene expression in dead or dying germ
cells was unlikely to be informative. Luckily, one *ovo* mRNA isoform with an extended exon 2 is also
required for egg chamber development beyond stage 5 (Salles et al. 2002; Benner et al. 2023). Utilizing
this finding allowed us to determine which genes in the female germline are transcriptionally responsive to
OVO during oogenesis. To do this, we used our *ovo^{ovo-GAL4}* allele. *ovo^{ovo-GAL4}* is a hypomorphic allele that
truncates the translation of *ovo-B* transcripts that include the exon 2 extension. About 90% of *ovo* mRNAs
are of this type (Benner et al. 2023). Females trans-heterozygous for *ovo^{ovo-GAL4}* and the *ovo-B* null allele
ovo^{ΔBP} show germline survival, with surviving germ cells arresting around stage 5 of oogenesis (Figure
4A)(Benner et al. 2023). Since this allele inherently expresses GAL4 in place of full length OVO due to
the T2A sequences, we can drive expression of rescuing and control sequences downstream with *UASp*
to generate excellent control samples. Therefore, we compared the gene expression occurring in sets of
ovaries that had the *ovo* hypomorphic phenotype with a sham rescue construct (*ovo^{ovo-GAL4}/ovo^{ΔBP}*; *UASp-*
GFP)(Figure 4A) versus those that drive the expression of the rescue construct expressing OVO-B
(*ovo^{ovo-GAL4}/ovo^{ΔBP}*; *UASp-3xFHA-OVO-B*)(Figure 4B).
559

Since *ovo$^{ovo\text{-}GAL4}$/ovo$^{\Delta BP}$*; *UASp-3xFHA-OVO-B* females have full rescue of the arrested germ cell
phenotype seen in *ovo$^{ovo\text{-}GAL4}$/ovo$^{\Delta BP}$*; *UASp-GFP* females, we needed to take further measures to ensure
our analysis of gene expression was stage comparable between the two sets of ovaries. We therefore
chose to dissect one day post-eclosion *ovo$^{ovo\text{-}GAL4}$/ovo$^{\Delta BP}$*; *UASp-3xFHA-OVO-B* female ovaries to enrich
for early stages of oogenesis, and collected only ovarioles containing the germarium through
previtellogenic egg chambers. *ovo$^{ovo\text{-}GAL4}$/ovo$^{\Delta BP}$*; *UASp-GFP* ovaries were collected at the same age
post-eclosion and we specifically collected ovaries that contained a visible ovariole structure (and
therefore contained germ cells) to minimize comparing germ cells to somatic ovary structures, but rather
germ cells to germ cells. We then performed RNA-seq in quadruplicate and measured the changes in
gene expression between the two genotypes. We used a significance level of p-adj < 0.05 and a log2 fold
change cutoff of >|0.5| to call differential expression. We utilized these log2 fold change cutoffs for two
reasons. Our control ovary genotype (*ovo$^{ovo\text{-}GAL4}$/ovo$^{\Delta BP}$*; *UASp-GFP*) has hypomorphic OVO activity,
hence germ cells are able to survive but are arrested. With the addition of ectopic rescue OVO in *ovo$^{ovo\text{-}}$*
^{GAL4}/ovo$^{\Delta BP}$; *UASp-3xFHA-OVO-B* ovaries, we predicted that genes that were directly regulated by OVO
would transcriptionally respond, however, we were unsure as to what degree the response would be in
comparison to hypomorphic OVO. So we therefore wanted to use a less conservative log2 fold change
cutoff in order to not miss any expression changes between hypomorphic OVO and rescue OVO. We
reasoned that if the changes were not significant between genotypes then minor changes in gene
expression would not matter. Our second reason for using these cutoffs is we had an internal control
between the two genotypes. We knew through immunostaining that Vas protein was present in the
germline of both genotypes (Figure 4A, 4B) and therefore was likely expressed at similar levels in the
germline of both genotypes. Both genotypes also expressed *GAL4* under the control of *ovo* in the
germline. We examined the expression levels of *vas* and *GAL4* and found that *vas* had a log2 fold change
of 0.15 (p-adj = 0.03) and *GAL4* had a log2 fold change of 0.33 (p-adj = 0.18) (Figure 4C). These data
suggest a slight underrepresentation of germline expression in the ovo mutant ovaries. Therefore, by
using the greater than 0.5 and less than -0.5 log2 fold change cutoffs, and a less than 0.05 p-adj value
cutoff, we would be conservative to not call genes differentially expressed due to differences in the
relative abundance of germ cells and somatic cells.
588
We were able to reliably detect the expression of 10,804 genes in these early ovarioles (Table S5). The
differential expression analysis indicated that 1,994 genes primarily expressed in the germline (see next
paragraph) significantly increased in expression with ectopic rescue OVO (Figure 4C, cyan/purple dots)
and 2,924 genes primarily expressed in the soma (see next paragraph) significantly decreased in
expression with ectopic rescue OVO expression (Figure 4C, yellow/blue dots). 5,886 genes were not
considered to be differentially expressed in our analysis (Figure 4C, gray dots). OVO is expressed in the
germline, not the soma, and previous work has shown that OVO-B is a transcriptional activator (Andrews
et al. 2000), so we hypothesized that many of the genes increasing in gene expression in the presence of
rescuing OVO were direct downstream targets. As a test, we intersected OVO-dependent gene
expression with OVO occupancy data. Among genes that showed OVO-dependent expression, we found
a significant enrichment for genes with OVO ChIP peaks (Figure 4C, cyan dots, p < 0.01, odds ratio =
2.21) while conversely, there was a significant depletion of genes that showed decreased expression in
OVO rescue ovaries and OVO ChIP peaks (Figure 4C, blue dots, p < 0.01, odds ratio = 0.85). This
strongly suggests that genes that are bound by OVO, transcriptionally respond in a positive manner. This
finding is fully consistent with our meta-analysis comparing OVO ChIP-seq and histone ChIP-seq/ATAC-
seq data (Figure 4E). OVO binding was highly associated with transcriptionally active histone marks such
as H3K27ac and H3K4me3, open chromatin, and increased expression.
606
There are genes that showed decreased expression in the OVO rescued ovaries, but we believe this is
technical rather than biological. OVO is expressed only in the germline, but ovarioles contain germ cells
and somatic cells. The presence of empty ovarioles, containing leftover strings of somatic cells, are
evident even in *ovo$^{ovo\text{-}GAL4}$/ovo$^{\Delta BP}$*; *UASp-GFP* ovaries that contain germ cells. Conversely, *ovo$^{ovo\text{-}}$*
^{GAL4}/ovo$^{\Delta BP}$; *UASp-3xFHA-OVO-B* ovaries are fully rescued, and therefore possess more germ cell
containing ovarioles than *ovo$^{ovo\text{-}GAL4}$/ovo$^{\Delta BP}$*; *UASp-GFP* ovaries (Benner et al. 2023). Despite our best
efforts to dissect individual ovarioles with a full complement of germ cells and egg chambers, we
wondered if there might be fewer germ cells and chambers in the *ovo$^{ovo\text{-}GAL4}$/ovo$^{\Delta BP}$*; *UASp-GFP* ovaries.
To address this directly, we tested the idea that genes with significantly increased expression with the

presence of ectopic rescue OVO were enriched for germ cell specific expression, and thus downstream of
OVO. We also tested the idea that genes that significantly decreased in expression were due to somatic
cell lineages, which should be enriched in the *ovo* hypomorph due to reduced numbers of germ cells in
the ovarioles (Benner et al. 2023). To confirm that genes increasing in expression in ectopic rescue OVO
were germline derived, we cross-referenced the significantly expressed genes in our RNA-seq datasets
with the modENCODE developmental RNaseq datasets (Graveley 2010). We extracted the gene names
of all genes that were considered to be 'moderately expressed' in 0-2 hour old embryos, which are
produced during oogenesis and are deposited into the early embryo. We found that 71% of genes
(1,409/1,994) that had a significant increase in expression in the presence of ectopic rescue OVO were
found to be expressed in 0-2 hour old embryos (Figure 4D, green dots), while only 21% of genes
(625/2,924) that had a significant decrease in expression were found in the same embryo dataset (Figure
4D, yellow dots). 3,448 genes from the 0-2 hour old embryo dataset were not differentially expressed in
our RNA-seq dataset (Figure 4D, purple dots). A Fisher's exact test confirmed that there was a significant
enrichment for genes that significantly increased in expression and were present in 0-2 hour old embryos
(p < 0.01, odds ratio = 2.8). In comparison, there was a significant depletion for genes that significantly
decreased in expression and were present in 0-2 hour old embryos (p < 0.01, odds ratio = 0.17). This
result indicated that genes that significantly increased in expression were more likely to be expressed in
the germline and that the presence of ectopic rescue OVO significantly increased the expression of genes
that were maternally deposited in the early embryo. While the set of genes that significantly decrease in
expression are not enriched in the embryo and are more likely specific to somatic cell gene expression.
These genes are unlikely to be direct OVO targets due to the absence of OVO in those cells, although we
certainly cannot rule out the possibility of a non-autonomous effect of OVO on somatic gene expression.
In terms of the germline proper, OVO appears to be a positively acting transcription factor.
639
<u>OVO Positively Regulates Essential Oogenesis Genes</u>
We wanted to examine a subset of the OVO target genes in detail, and began with the known OVO
targets, *ovo* itself and *otu* (Bielinska et al. 2005; Lü et al. 1998; Lü and Oliver 2001). Since the
relationship between OVO binding to these two genes has been well-characterized, we validated the
OVO ChIP, histone ChIP/ATAC-seq, and RNA-seq datasets by examining these two genes first. Since
OVO positively regulates the expression of both these genes, then we would expect OVO to be physically
bound at OVO motifs required for high transcription, the presence of transcriptionally active histone marks
and open chromatin, as well as a positive transcriptional response in the presence of rescuing OVO-B.
This is exactly what we observed. A significant OVO ChIP peak was found overlapping the TSS of *ovo-B*,
with four significant OVO DNA binding motifs present (Figure 5A). ATAC-seq, H3K27ac, and H3K4me3
peaks overlapped the *ovo* promoter. Transcriptionally, *ovo* RNA-seq reads are derived from the *UASp-*
3xFHA-OVO-B cDNA rescue, so we could not assess whether *ovo* responded transcriptionally to ectopic
rescue OVO. However, when looking at the *otu* locus (Figure S2), we found OVO occupancy over the
TSS of both annotated *otu* promoters, with significant OVO DNA binding motifs overlapping and in close
proximity to the TSSs. The *otu* locus contained similar ATAC-seq and activating histone mark peaks
overlapping the TSS found at the *ovo* locus. It was also evident that *otu* had a positive transcriptional
response to the presence of OVO rescue (log2 fold change = 2.41; p-adj < 0.001). These results confirm
that OVO binds and positively regulates both itself and *otu in-vivo*, as previous work has indicated.
658
Since our overlapping OVO ChIP-seq and RNA-seq data suggests that hundreds of genes that are bound
by OVO increase in expression in the presence of ectopic rescue OVO, we wanted to know more about
the functions of those genes. For the big picture, we performed Gene Ontology enrichment analysis with
gProfiler software (Raudvere et al. 2019). To be especially stringent, we focused on the genes that
contained an OVO ChIP peak overlapping the transcriptional start site and significantly increased in
expression in the presence of rescue OVO. 525 genes met these criteria. Biological process GO term
enrichment analysis on these 525 genes showed a significant enrichment for 109 GO terms (Table S6).
The significant GO terms were almost exclusively related to female reproduction and maternal control of
early embryonic development, including embryonic axis specification, mRNA localization, egg activation,
and negative regulation of translation (Figure 6A). These associated GO terms are well understood in the
context of oogenesis and broadly suggest that OVO expression in adult gonads is essential for
constructing an egg and depositing maternal RNAs to support zygotic embryonic development.
671

GO term enrichment analysis of genes that are bound by OVO and increase in expression in the
presence of ectopic rescue OVO, suggested that OVO is likely a main transcriptional regulator of
oogenesis. These genes are the subject of decades of work on *Drosophila* oogenesis, but essentially all
the work on them has focused on what they do, not on how they are transcriptionally regulated. For
example, *bicoid* (Figure 4B), and *bicoid* mRNA binding proteins *exuperantia* (*exu*) and *swallow* (*swa*), are
essential for anterior specification of the embryo (Lasko 2012). All of these genes were occupied by OVO
in vivo, significantly upregulated by OVO-B, and had OVO motifs in close proximity to the TSS. Genes
involved in posterior patterning (*oskar* and *nanos*) (Figure 4C), as well as pole cell specification genes
(*polar granule component*, *germ cell-less*, and *aubergine*)(Lasko 2012; L. Benner, Deshpande, and Lerit
2018), also showed similar RNA-seq, ChIP-seq, and OVO DNA binding motif profiles as *ovo* and *otu*.
Genes that are involved in translational silencing and regulation of maternally provided mRNAs, such as
cup (Figure 4D), *maternal expression at 31B* (*me31B*), *oo18 RNA-binding protein* (*orb*), and *bruno 1*
(*bru1*) as well as essential genes involved in meiosis completion and egg activation after fertilization
(*giant nuclei* (*gnu*), *pan gu* (*png*), *plutonium* (*plu*), *wispy* (*wisp*), *C(3)G*, and *matrimony* (*mtrm*)) (Figure
4E)(Lasko 2012; Avilés-Pagán and Orr-Weaver 2018) all show this stereotypic pattern of promoter
proximal OVO occupancy and DNA binding motifs, and OVO-dependent transcription. These data
indicate that the *ovo* locus is a central transcription factor activating the expression of essential maternal
and early embryonic development pathways in the female germline.
690
We also found that the genes *fs(1)N*, *fs(1)M3*, and *closca*, were all bound by OVO and responded
transcriptionally to the presence of ectopic rescue OVO. These genes are significant because they
constitute a set of genes that are expressed in the germline and are eventually incorporated into the
vitelline membrane providing the structural integrity and impermeability of the egg (Mineo, Furriols, and
Casanova 2017; Ventura et al. 2010). Loss-of-function of these three genes results in flaccid eggs that
are permeable and fail to develop. A testament to the likelihood of OVO regulating these genes is the
finding that females transheterozygous for the dominant antimorphic *ovo^{D3}* allele lay eggs that are flaccid
and contain a permeable vitelline membrane, assayed through their ability to uptake the dye neutral red
(Oliver, Pauli, and Mahowald 1990). Presumably in these females, the expression of the repressive form
of OVO leads to a reduction in the expression of these three essential vitelline membrane genes which
phenocopies loss-of-function alleles of each, respectively. Altogether, this largely indicates that not only is
OVO involved in regulating the expression of numerous essential maternal pathways for embryonic
development, it is also essential for regulating genes that are required for egg integrity and maturation.
704
Discussion
Since its original isolation as a dominant female sterile locus in *Drosophila (Busson et al. 1983;*
Komitopoulou et al. 1983), *ovo* has long been known as an essential gene in oogenesis. Female germ
cells require *ovo* for survival and differentiation, while it has no described roles or functions in the male
germline (Oliver, Perrimon, and Mahowald 1987; Oliver, Pauli, and Mahowald 1990). OVO has also been
found to be eternally present in the female germline, attesting to its likely continual requirement for female
germ cell viability and identity (Benner et al. 2023). Our work here significantly expands our knowledge on
OVO function in the female germline, showing that OVO binds and positively regulates a large array of
genes required to build an egg and pattern the resulting embryo after fertilization. OVO accomplishes this
by directly binding to the promoters of its targets, as well as more distant sites that could represent
enhancers. Altogether, we suggest that OVO is a master transcriptional regulator coordinating a number
of essential maternal pathways involved in oocyte and early embryonic development. Hints of these
functions can be found in the hypomorphic and antimorphic *ovo* alleles which show egg chamber arrest,
ventralized eggs, and permeable vitelline membranes. It is clear that OVO is required to activate multiple
pathways involved in oocyte and early embryonic development.
720
The GO term enrichment analysis on genes that were bound by OVO and transcriptionally responded to
OVO surprisingly indicated a large degree of overlap in oocyte and early embryonic developmental
pathways. Also, OVO seemed to reinforce these pathways at multiple key genes within each pathway.
For example, OVO bound to the promoters and increased the expression of *bcd*, and the *bcd* mRNA
binding proteins *exu* and *swa*, all involved in ensuring correct anterior specification of the embryo (Lasko
2012). Genes that are essential for egg activation were coordinately regulated by OVO as well. OVO
downstream target genes *gnu*, *png*, *plu*, and *wisp* all belong in the same interconnected pathway

ensuring egg activation (Avilés-Pagán and Orr-Weaver 2018). A similar story was found for genes such
as *cup*, *Me31b*, *bru1*, and *orb*, indicating that OVO controls a battery of genes involved in the positive
regulation of RNA binding proteins that negatively regulate translation (Lasko 2012). OVO also bound and
positively regulated a number of posterior and germ plasm specification genes such as *osk*, *nos*, *aub*, *gcl*,
and *pgc (Lasko 2012; Mahowald 2001).* Given this plethora of famous maternal effect loci, it might be
tempting to suggest that OVO is sufficient for egg production, but there are important exceptions. For
example, other important germ plasm factors such as *vasa*, *staufen*, and *tudor* were not bound by or
transcriptionally responsive to OVO. This observation suggest that other transcription factors are be
responsible for regulating these genes.
737
OVO binds in close proximity to the TSS of genes it positively regulates, however, it is still unclear
precisely how it regulates gene expression. The possibilities include integration into the RNA Polymerase
complex itself, a short distance sigma factor like function, a core promoter conditioning function
(pioneering), and/or garden variety transcription factor. Although core promoters active specifically in the
ovary are enriched for OVO DNA binding motifs, we did not find a strict spatial orientation for these motifs
in relation to the TSS, such as is found with other DNA elements such as INR, DPE, and MTE (Ohler et
al. 2002; FitzGerald et al. 2006; Lim et al. 2004). It is therefore unlikely that OVO is a core component of
the RNA Polymerase complex in the female germline. This suggests that it is unlikely to be analogous to
male specific TATA-associated factors that have been shown to activate gene expression in the male
germline (M. Hiller et al. 2004; M. A. Hiller et al. 2001; Lu et al. 2020; V. C. Li et al. 2009). It is therefore
possible, and previously well supported, that OVO is a strict activator of transcription (Lü et al. 1998; Lü
and Oliver 2001; Bielinska et al. 2005). Where OVO binding in close proximity, in any orientation relative
to the TSS, activates transcription. One aspect of OVO DNA binding that showed differences with stage-
specific histone ChIP and ATAC-seq, was OVO's strong association with open chromatin. The role of
repressors of transcription such as the polycomb complex, *egg*, *wde*, and *Su(var)205* in restricting gene
expression through promoting heterochromatin formation in differentiating egg chambers is well
established (Smolko, Shapiro-Kulnane, and Salz 2018; DeLuca et al. 2020). OVO might ensure that the
chromatin status of maternally expressed genes remains open. Evidence from our work points to OVO
fulfilling that role.
757
In GSCs and early egg chambers, OVO ChIP peaks largely overlap open chromatin and transcriptionally
active histone marks. However, in stage 5 egg chambers, there was an even higher degree of association
with open chromatin (all OVO ChIP peaks overlapped ATAC-seq peaks), while the significant association
with H3K27ac marks was greatly reduced. This difference is likely significant. As GSCs differentiate, they
accumulate repressive chromatin marks while the number of ATAC-seq and H3K27ac peaks are reduced.
This increase in association with open chromatin and OVO binding, even as the amount of open
chromatin is reduced throughout egg chamber differentiation, might indicate that OVO binding helps to
maintain chromatin accessibility, even when the locus in question is no longer actively transcribed. The
loss of histone marks of active transcription at OVO-bound open chromatin in later differentiating egg
chambers might mean that OVO does not influence the transcriptional potential of target genes as
strongly as it influences the chromatin status in this second phase. Therefore, OVO might be more similar
in function to pioneer factors/chromatin remodelers than it is to a transcription factor that is only involved
in activating transcription.
771
In light of OVOs stronger association with open chromatin over histone marks identified with active
transcription, evidence for OVO as a potential pioneering factor comes through its developmental protein
expression and the eventual expression of downstream target genes. OVO is maternally deposited into
the embryo where it eventually is sequestered to embryonic pole cells (Hayashi et al. 2017; Benner et al.
2023). Maternal OVO maintains a nuclear localization within the germline throughout embryonic
development until it eventually overlaps with its own zygotic gene expression as well as its downstream
target *otu* at stage 15 (Benner et al. 2023; Casper and Van Doren 2009). Pole cells are transcriptionally
quiescent (A. Nakamura and Seydoux 2008; Deshpande et al. 1999; Van Doren, Williamson, and
Lehmann 1998), with the first signs of zygotic transcription occurring at stage 8 (Van Doren, Williamson,
and Lehmann 1998). So why doesn't zygotic *ovo* and *otu* transcription begin until much later in
development even though maternal OVO is present and germ cells are transcriptionally competent much
earlier? One explanation for this is that OVO bound genes are also dependent on other transcription

factors in the developing germline, despite being remarkably compact. The OVO-dependent *ovo* core
promoter is very compact; showing expression with only 10s of bp of sequence in reporters. While it is
possible that OVO is co-dependent on the presence of other transcription factors that are expressed later,
or that OVO does not have the ability to activate transcription by itself and replaced by a more traditional
transcription factor. This role is more similar to pioneer transcription factors such as *zelda* (*zld*).
789
It has been well established that *zld* influences the maternal to zygotic transition by binding its target
genes before they become transcriptionally active (Harrison et al. 2011). ZLD binding increases
chromatin accessibility which allows for other activating transcription factors to bind and positively
influence gene expression (Xu et al. 2014). Loss of maternal *zld* does affect a target promoter's ability to
be activated, indicating that the difference between a pioneer factor and a transcriptional activator is
slight, but reducing chromatin accessibility can negate the effects of the presence of a transcriptional
activator and its ability to influence a target promoter. Although pioneer factors are expressed in the
female germline and are maternally deposited into the embryo to regulate the maternal to zygotic
transition, surprisingly, a functional role for pioneer factors has yet to be described in the developing
embryonic germline. If OVO is involved in influencing chromatin status, then maternally depleting it should
have a negative effect on embryonic germ cell chromatin status and gene expression. At this time, it is
technically challenging to deplete maternal OVO and determine the embryonic germ cell phenotype, as
attempts to deplete maternal OVO leads to egg chamber arrest/death in adult females. However, based
on OVO's relationship to open chromatin, its eternal expression in the female germline, and delayed
target gene expression, we propose a model where maternal OVO binds target promoters and influences
chromatin status in the transcriptionally quiescent embryonic germline. OVO binding a target promoter
without the presence of other transcription factors, however, does not in itself activate gene expression.
As germ cells develop, the eventual expression of activating transcription factors allow for OVO bound
genes to become expressed (Figure 7). Expression of OVO bound genes are required in the adult
germline to allow for germ cell differentiation and egg development. Ultimately resulting in the maternal
deposition of mRNAs and proteins required for early embryonic development, including *ovo* mRNA and
OVO protein, maintaining the cyclical nature of the germline.
812

813 **Data Availability**
814 Drosophila strains used for this study are available upon request. All sequence information and datasets
815 used in this study are in Table S1.
816

Acknowledgements
We would like to thank previous and current members of the Oliver lab, Laboratory of Biochemistry and
Genetics at NIH, and L.B. committee members Mark Van Doren and Allan Spradling for insightful
discussion and comments on this work throughout. Monoclonal antibodies were obtained from the
Developmental Studies Hybridoma Bank, created by the Eunice Kennedy Shriver National Institute of
Child Health and Human Development (NICHD) of the NIH and maintained at the Department of Biology,
University of Iowa, Iowa City, IA 52242. Genetic and genomic information was obtained from FlyBase
(U41 HG-000739). This work utilized the computational resources of the NIH High-Performance
Computing Biowulf cluster (http://hpc.nih.gov). Sequencing was completed by The National Heart, Lung,
and Blood Institute (NHLBI) DNA Sequencing and Genomics Core.
827

828 **Funding**
829 This research was supported in part by the Intramural Research Program of the NIH, The National
830 Institute of Diabetes and Digestive and Kidney Diseases (NIDDK) (awarded to B.O.). L.B. was supported
831 by the NIH Graduate Partnership Program.
832
833

References
Anders, Simon, Paul Theodor Pyl, and Wolfgang Huber. 2015. "HTSeq--a Python Framework to Work
with High-Throughput Sequencing Data." *Bioinformatics* 31 (2): 166–69.

Andrews, J., D. Garcia-Estefania, I. Delon, J. Lü, M. Mével-Ninio, A. Spierer, F. Payre, D. Pauli, and B.
Oliver. 2000. "OVO Transcription Factors Function Antagonistically in the Drosophila Female
Germline." *Development* 127 (4): 881–92.
Aoyagi, N., and D. A. Wassarman. 2000. "Genes Encoding Drosophila Melanogaster RNA Polymerase II
General Transcription Factors: Diversity in TFIIA and TFIID Components Contributes to Gene-
Specific Transcriptional Regulation." *The Journal of Cell Biology* 150 (2): F45–50.
Avilés-Pagán, Emir E., and Terry L. Orr-Weaver. 2018. "Activating Embryonic Development in
Drosophila." *Seminars in Cell & Developmental Biology* 84 (December): 100–110.
Bailey, Timothy L. 2021. "STREME: Accurate and Versatile Sequence Motif Discovery." *Bioinformatics*
37 (18): 2834–40.
Bailey, Timothy L., James Johnson, Charles E. Grant, and William S. Noble. 2015. "The MEME Suite."
Nucleic Acids Research 43 (W1): W39–49.
Bastock, Rebecca, and Daniel St Johnston. 2008. "Drosophila Oogenesis." *Current Biology: CB* 18 (23):
R1082–87.
Benner, L., G. Deshpande, and D. A. Lerit. 2018. "Primordial Germ Cells of Drosophila Melanogaster."
https://scholar.google.ca/scholar?cluster=7409236235465525669&hl=en&as_sdt=0,5&sciodt=0,5.
Benner, Leif, Elias A. Castro, Cale Whitworth, Koen J. T. Venken, Haiwang Yang, Junnan Fang, Brian
Oliver, Kevin R. Cook, and Dorothy A. Lerit. 2019. "Drosophila Heterochromatin Stabilization
Requires the Zinc-Finger Protein Small Ovary." *Genetics* 213 (3): 877–95.
Benner, Leif, Savannah Muron, and Brian Oliver. 2023. "Female Germline Expression of OVO
Transcription Factor Bridges Drosophila Generations." *bioRxiv*.
https://doi.org/10.1101/2023.08.25.554887.
Bielinska, Beata, Jining Lü, David Sturgill, and Brian Oliver. 2005. "Core Promoter Sequences Contribute
to Ovo-B Regulation in the Drosophila Melanogaster Germline." *Genetics* 169 (1): 161–72.
Bishop, D. L., and R. C. King. 1984. "An Ultrastructural Study of Ovarian Development in the otu7 Mutant
of Drosophila Melanogaster." *Journal of Cell Science* 67 (April): 87–119.
broadinstitute. n.d. "Picard." Picard. http://broadinstitute.github.io/picard/.
Busson, D., M. Gans, K. Komitopoulou, and M. Masson. 1983. "Genetic Analysis of Three Dominant
Female-Sterile Mutations Located on the X Chromosome of DROSOPHILA MELANOGASTER."
Genetics 105 (2): 309–25.
Casper, Abbie L., and Mark Van Doren. 2009. "The Establishment of Sexual Identity in the Drosophila
Germline." *Development* 136 (22): 3821–30.
Castro-Mondragon, Jaime A., Rafael Riudavets-Puig, Ieva Rauluseviciute, Roza Berhanu Lemma, Laura
Turchi, Romain Blanc-Mathieu, Jeremy Lucas, et al. 2022. "JASPAR 2022: The 9th Release of the
Open-Access Database of Transcription Factor Binding Profiles." *Nucleic Acids Research* 50 (D1):
D165–73.
Chen, Zhen-Xia, David Sturgill, Jiaxin Qu, Huaiyang Jiang, Soo Park, Nathan Boley, Ana Maria Suzuki, et
al. 2014. "Comparative Validation of the D. Melanogaster modENCODE Transcriptome Annotation."
Genome Research 24 (7): 1209–23.
Danecek, Petr, James K. Bonfield, Jennifer Liddle, John Marshall, Valeriu Ohan, Martin O. Pollard,
Andrew Whitwham, et al. 2021. "Twelve Years of SAMtools and BCFtools." *GigaScience* 10 (2).
https://doi.org/10.1093/gigascience/giab008.
DeLuca, Steven Z., Megha Ghildiyal, Liang-Yu Pang, and Allan C. Spradling. 2020. "Differentiating
Drosophila Female Germ Cells Initiate Polycomb Silencing by Regulating PRC2-Interacting Proteins."
eLife 9 (August). https://doi.org/10.7554/eLife.56922.
Deshpande, G., G. Calhoun, J. L. Yanowitz, and P. D. Schedl. 1999. "Novel Functions of Nanos in
Downregulating Mitosis and Transcription during the Development of the Drosophila Germline." *Cell*
99 (3): 271–81.
FitzGerald, Peter C., David Sturgill, Andrey Shyakhtenko, Brian Oliver, and Charles Vinson. 2006.
"Comparative Genomics of Drosophila and Human Core Promoters." *Genome Biology* 7 (7): R53.
Gramates, L. Sian, Julie Agapite, Helen Attrill, Brian R. Calvi, Madeline A. Crosby, Gilberto Dos Santos,
Joshua L. Goodman, et al. 2022. "FlyBase: A Guided Tour of Highlighted Features." *Genetics* 220
(4). https://doi.org/10.1093/genetics/iyac035.
Grant, Charles E., Timothy L. Bailey, and William Stafford Noble. 2011. "FIMO: Scanning for Occurrences
of a given Motif." *Bioinformatics* 27 (7): 1017–18.
Graveley, Brenton. 2010. "The Developmental Transcriptome of Drosophila Melanogaster." *Genome*

Biology 11 (Suppl 1): I11.

Gupta, Shobhit, John A. Stamatoyannopoulos, Timothy L. Bailey, and William Stafford Noble. 2007. "Quantifying Similarity between Motifs." *Genome Biology* 8 (2): R24.

Haberle, Vanja, Alistair R. R. Forrest, Yoshihide Hayashizaki, Piero Carninci, and Boris Lenhard. 2015. "CAGEr: Precise TSS Data Retrieval and High-Resolution Promoterome Mining for Integrative Analyses." *Nucleic Acids Research* 43 (8): e51.

Harms, E., T. Chu, G. Henrion, and S. Strickland. 2000. "The Only Function of Grauzone Required for Drosophila Oocyte Meiosis Is Transcriptional Activation of the Cortex Gene." *Genetics* 155 (4): 1831–39.

Harrison, Melissa M., Xiao-Yong Li, Tommy Kaplan, Michael R. Botchan, and Michael B. Eisen. 2011. "Zelda Binding in the Early Drosophila Melanogaster Embryo Marks Regions Subsequently Activated at the Maternal-to-Zygotic Transition." *PLoS Genetics* 7 (10): e1002266.

Hayashi, Makoto, Yuko Shinozuka, Shuji Shigenobu, Masanao Sato, Michihiko Sugimoto, Seiji Ito, Kuniya Abe, and Satoru Kobayashi. 2017. "Conserved Role of Ovo in Germline Development in Mouse and Drosophila." *Scientific Reports* 7 (1): 1–10.

Hiller, M. A., T. Y. Lin, C. Wood, and M. T. Fuller. 2001. "Developmental Regulation of Transcription by a Tissue-Specific TAF Homolog." *Genes & Development* 15 (8): 1021–30.

Hiller, Mark, Xin Chen, M. Jodeane Pringle, Martin Suchorolski, Yasemin Sancak, Sridhar Viswanathan, Benjamin Bolival, Ting-Yi Lin, Susan Marino, and Margaret T. Fuller. 2004. "Testis-Specific TAF Homologs Collaborate to Control a Tissue-Specific Transcription Program." *Development* 131 (21): 5297–5308.

Hinson, S., J. Pettus, and R. N. Nagoshi. 1999. "Regulatory and Functional Interactions between Ovarian Tumor and Ovo during Drosophila Oogenesis." *Mechanisms of Development* 88 (1): 3–14.

Hubbard, T., D. Barker, E. Birney, G. Cameron, Y. Chen, L. Clark, T. Cox, et al. 2002. "The Ensembl Genome Database Project." *Nucleic Acids Research* 30 (1): 38–41.

Kent, W. James, Charles W. Sugnet, Terrence S. Furey, Krishna M. Roskin, Tom H. Pringle, Alan M. Zahler, and David Haussler. 2002. "The Human Genome Browser at UCSC." *Genome Research* 12 (6): 996–1006.

Kim, Daehwan, Joseph M. Paggi, Chanhee Park, Christopher Bennett, and Steven L. Salzberg. 2019. "Graph-Based Genome Alignment and Genotyping with HISAT2 and HISAT-Genotype." *Nature Biotechnology* 37 (8): 907–15.

Komitopoulou, K., M. Gans, L. H. Margaritis, F. C. Kafatos, and M. Masson. 1983. "Isolation and Characterization of Sex-Linked Female-Sterile Mutants in DROSOPHILA MELANOGASTER with Special Attention to Eggshell Mutants." *Genetics* 105 (4): 897–920.

Lasko, Paul. 2012. "mRNA Localization and Translational Control in Drosophila Oogenesis." *Cold Spring Harbor Perspectives in Biology* 4 (10). https://doi.org/10.1101/cshperspect.a012294.

Lee, S., and M. D. Garfinkel. 2000. "Characterization of Drosophila OVO Protein DNA Binding Specificity Using Random DNA Oligomer Selection Suggests Zinc Finger Degeneration." *Nucleic Acids Research* 28 (3): 826–34.

Li, Jian, and David S. Gilmour. 2013. "Distinct Mechanisms of Transcriptional Pausing Orchestrated by GAGA Factor and M1BP, a Novel Transcription Factor." *The EMBO Journal* 32 (13): 1829–41.

Lim, Chin Yan, Buyung Santoso, Thomas Boulay, Emily Dong, Uwe Ohler, and James T. Kadonaga. 2004. "The MTE, a New Core Promoter Element for Transcription by RNA Polymerase II." *Genes & Development* 18 (13): 1606–17.

Li, Victor C., Jerel C. Davis, Kapa Lenkov, Benjamin Bolival, Margaret T. Fuller, and Dmitri A. Petrov. 2009. "Molecular Evolution of the Testis TAFs of Drosophila." *Molecular Biology and Evolution* 26 (5): 1103–16.

Li, Xiao-Yong, Melissa M. Harrison, Jacqueline E. Villalta, Tommy Kaplan, and Michael B. Eisen. 2014. "Establishment of Regions of Genomic Activity during the Drosophila Maternal to Zygotic Transition." *eLife* 3 (October): e03737.

Love, Michael I., Wolfgang Huber, and Simon Anders. 2014. "Moderated Estimation of Fold Change and Dispersion for RNA-Seq Data with DESeq2." *Genome Biology* 15 (12): 550.

Lu, Dan, Ho-Su Sin, Chenggang Lu, and Margaret T. Fuller. 2020. "Developmental Regulation of Cell Type-Specific Transcription by Novel Promoter-Proximal Sequence Elements." *Genes & Development* 34 (9-10): 663–77.

Lü, J., J. Andrews, D. Pauli, and B. Oliver. 1998. "Drosophila OVO Zinc-Finger Protein Regulates Ovo

and Ovarian Tumor Target Promoters." *Development Genes and Evolution* 208 (4): 213–22.

Lü, J., and B. Oliver. 2001. "Drosophila OVO Regulates Ovarian Tumor Transcription by Binding Unusually near the Transcription Start Site." *Development* 128 (9): 1671–86.

Mahowald, A. P. 2001. "Assembly of the Drosophila Germ Plasm." *International Review of Cytology* 203: 187–213.

Mineo, Alessandro, Marc Furriols, and Jordi Casanova. 2017. "Transfer of Dorsoventral and Terminal Information from the Ovary to the Embryo by a Common Group of Eggshell Proteins in Drosophila." *Genetics* 205 (4): 1529–36.

Nakamura, Akira, and Geraldine Seydoux. 2008. "Less Is More: Specification of the Germline by Transcriptional Repression." *Development* 135 (23): 3817–27.

Nakamura, Shoichi, Seiji Hira, Masato Fujiwara, Nasa Miyagata, Takuma Tsuji, Akane Kondo, Hiroshi Kimura, et al. 2019. "A Truncated Form of a Transcription Factor Mamo Activates Vasa in Drosophila Embryos." *Communications Biology* 2 (1): 1–13.

Ohler, Uwe, Guo-Chun Liao, Heinrich Niemann, and Gerald M. Rubin. 2002. "Computational Analysis of Core Promoters in the Drosophila Genome." *Genome Biology* 3 (12): RESEARCH0087.

Oliver, B., D. Pauli, and A. P. Mahowald. 1990. "Genetic Evidence That the Ovo Locus Is Involved in Drosophila Germ Line Sex Determination." *Genetics* 125 (3): 535–50.

Oliver, B., N. Perrimon, and A. P. Mahowald. 1987. "The Ovo Locus Is Required for Sex-Specific Germ Line Maintenance in Drosophila." *Genes & Development* 1 (9): 913–23.

Pang, Liang-Yu, Steven DeLuca, Haolong Zhu, John M. Urban, and Allan C. Spradling. 2023. "Chromatin and Gene Expression Changes during Female Drosophila Germline Stem Cell Development Illuminate the Biology of Highly Potent Stem Cells." *eLife* 12 (October): RP90509.

Pauli, D., B. Oliver, and A. P. Mahowald. 1993. "The Role of the Ovarian Tumor Locus in Drosophila Melanogaster Germ Line Sex Determination." *Development* 119 (1): 123–34.

Quinlan, Aaron R., and Ira M. Hall. 2010. "BEDTools: A Flexible Suite of Utilities for Comparing Genomic Features." *Bioinformatics* 26 (6): 841–42.

Ramírez, Fidel, Devon P. Ryan, Björn Grüning, Vivek Bhardwaj, Fabian Kilpert, Andreas S. Richter, Steffen Heyne, Friederike Dündar, and Thomas Manke. 2016. "deepTools2: A next Generation Web Server for Deep-Sequencing Data Analysis." *Nucleic Acids Research* 44 (W1): W160–65.

Raudvere, Uku, Liis Kolberg, Ivan Kuzmin, Tambet Arak, Priit Adler, Hedi Peterson, and Jaak Vilo. 2019. "g:Profiler: A Web Server for Functional Enrichment Analysis and Conversions of Gene Lists (2019 Update)." *Nucleic Acids Research* 47 (W1): W191–98.

R Core Team. 2021. "R: A Language and Environment for Statistical Computing." *R Foundation for Statistical Computing*. https://www.R-project.org/.

Rice, G. A., M. J. Chamberlin, and C. M. Kane. 1993. "Contacts between Mammalian RNA Polymerase II and the Template DNA in a Ternary Elongation Complex." *Nucleic Acids Research* 21 (1): 113–18.

Salles, Cathy, Maryvonne Mével-Ninio, Alain Vincent, and François Payre. 2002. "A Germline-Specific Splicing Generates an Extended Ovo Protein Isoform Required for Drosophila Oogenesis." *Developmental Biology* 246 (2): 366–76.

Santos, Gilberto dos, Andrew J. Schroeder, Joshua L. Goodman, Victor B. Strelets, Madeline A. Crosby, Jim Thurmond, David B. Emmert, William M. Gelbart, and FlyBase Consortium. 2015. "FlyBase: Introduction of the Drosophila Melanogaster Release 6 Reference Genome Assembly and Large-Scale Migration of Genome Annotations." *Nucleic Acids Research* 43 (Database issue): D690–97.

Schindelin, Johannes, Ignacio Arganda-Carreras, Erwin Frise, Verena Kaynig, Mark Longair, Tobias Pietzsch, Stephan Preibisch, et al. 2012. "Fiji: An Open-Source Platform for Biological-Image Analysis." *Nature Methods* 9 (7): 676–82.

Smolko, Anne E., Laura Shapiro-Kulnane, and Helen K. Salz. 2018. "The H3K9 Methyltransferase SETDB1 Maintains Female Identity in Drosophila Germ Cells." *Nature Communications* 9 (1): 1–11.

Spradling, Allan C., Wanbao Niu, Qi Yin, Madhulika Pathak, and Bhawana Maurya. 2022. "Conservation of Oocyte Development in Germline Cysts from Drosophila to Mouse." *eLife* 11 (November): e83230.

Van Doren, Mark, Anne L. Williamson, and Ruth Lehmann. 1998. "Regulation of Zygotic Gene Expression in Drosophila Primordial Germ Cells." *Current Biology: CB* 8 (4): 243–46.

Ventura, Gemma, Marc Furriols, Nicolás Martín, Vitor Barbosa, and Jordi Casanova. 2010. "Closca, a New Gene Required for Both Torso RTK Activation and Vitelline Membrane Integrity. Germline Proteins Contribute to Drosophila Eggshell Composition." *Developmental Biology* 344 (1): 224–32.

Vo Ngoc, Long, George A. Kassavetis, and James T. Kadonaga. 2019. "The RNA Polymerase II Core

Promoter in Drosophila." *Genetics* 212 (1): 13–24.
Xu, Zhe, Hongtao Chen, Jia Ling, Danyang Yu, Paolo Struffi, and Stephen Small. 2014. "Impacts of the
Ubiquitous Factor Zelda on Bicoid-Dependent DNA Binding and Transcription in Drosophila." *Genes*
& Development 28 (6): 608–21.
Zhang, Yong, Tao Liu, Clifford A. Meyer, Jérôme Eeckhoute, David S. Johnson, Bradley E. Bernstein,
Chad Nusbaum, et al. 2008. "Model-Based Analysis of ChIP-Seq (MACS)." *Genome Biology* 9 (9):
R137.
1012

ovo^{Cterm-3xFHA}/ovo^{Cterm-GFP}

OVO ChIP DNA Motif

D Motif One — p<0.001, Contained in 50% of ChIP Peaks

E Motif Two — p<0.001, Contained in 37% of ChIP Peaks

F Motif Three — p<0.05, Contained in 17% of ChIP Peaks

G Motif Four — p<0.01, Contained in 5% of ChIP Peaks

H ChIP Minus Input Motif Centered Read Density

- OVO Motif One
- OVO Motif Two
- OVO Motif Three
- OVO Motif Four
- OVO Motif Garfinkel
- OVO Motif Oliver

I Distance of OVO ChIP Peaks To Genome Attribute

Genome Attribute
- Promoter
- Transcription Termination
- Open Reading Frames

J ChIP Minus Input TSS Centered Read Density

- OVO Motif One
- OVO Motif Two
- OVO Motif Three
- OVO Motif Four
- OVO Motif Garfinkel
- OVO Motif Oliver
- Non-OVO Bound

K ChIP Minus Input Ovary Cage TSS Centered Read Density

- OVO Bound
- Non-OVO Bound

L ChIP Minus Input Ovary CAGE-seq
OVO Bound   Non-OVO Bound

Normalized Control Subtracted Read Count

1013

Figure 1: Significantly Enriched OVO DNA Binding Motifs and OVO ChIP Attributes Genome-Wide.
A-C) Immunofluorescent staining of adult ovarioles of $ovo^{Cterm-3xFHA}/ovo^{Cterm-GFP}$ females (20x, scale bar =
20 μm). Ovarioles were stained for Vas (purple) to label the germline, HA (cyan) to label OVO-HA, and
GFP (yellow) to label OVO-GFP. The homozygous version of these alleles were used to ChIP OVO. D-G)
Significantly enriched motifs found within overlapping OVO ChIP peaks. The percentage of OVO ChIP
peaks each motif was found and their corresponding p-value are indicated to the right. H) OVO ChIP
minus input control ChIP-seq read coverage density centered on each individual motif's location. I)
Relative distance of OVO ChIP peaks to gene level promoters, terminations sequences, and open
reading frames genome-wide. J) OVO ChIP minus input control ChIP-seq read coverage density for
genes containing significant OVO ChIP peaks and the corresponding OVO DNA binding motif. Genes are
centered on the transcriptional start site. K, L) OVO ChIP minus input control ChIP-seq read coverage
density and heatmap plots centered on the dominant significant ovary CAGE-seq TSSs overlapping or
not overlapping OVO ChIP peaks.

Figure 2: OVO DNA Binding is Associated with Open Chromatin and Transcriptionally Active
Histone Marks. A, B) OVO ChIP minus input control, GSC and 32c ATAC-seq, GSC H3K27ac,
H3K4me3, H3K27me3, H3K9me3, 8c NC H3K9me3, 32c NC H3K27ac, and H3K27me3 ChIP-seq read
coverage density centered on each OVO peak maximum or individual motif's location. C) Relative
distance of OVO ChIP peaks to significantly called peaks for GSC and 32c ATAC-seq, GSC H3K27ac,
H3K4me3, H3K27me3, H3K9me3, 8c NC H3K9me3, 32c NC H3K27ac, and H3K27me3 ChIP-seq
genome-wide. D) Total number of significant peaks and the total number of overlapping peaks between
OVO ChIP and GSC and 32c ATAC-seq, GSC H3K27ac, H3K4me3, H3K27me3, H3K9me3, 8c NC
H3K9me3, 32c NC H3K27ac, and H3K27me3 ChIP-seq. Asterisk indicates significantly enriched overlap
while hashtag indicates significantly depleted overlap between datasets.

1038
1039 **Figure 3: OVO Bound Promoters are Enriched for INR, DPE, and MTE elements.** A) Histogram of the
1040 distance of *in vivo* and *in vitro* OVO DNA binding motifs within significant overlapping OVO ChIP peaks
1041 from the closest genes TSS. B-F) Histogram of the percent of promoters from tissue specific CAGE-seq
1042 analysis of common promoter motif elements centered on the dominant significant TSS.

OVO Ovary RNAseq Samples

A *ovo^{ΔBP}/ovo^{ovo-GAL4}; UASp-GFP*

B *ovo^{ΔBP}/ovo^{ovo-GAL4}; UASp-3xFHA-OVO-B*

20 µm

Vas α-Spec DAPI

C OVO RNA-seq/OVO ChIP MA Plot

1,328
666
ovo
otu
vas
564
2,360
GAL4
GFP

D OVO RNA-seq/0-2 Hour Old Embryo

1,409
0.5
-0.5
3,448
625

● Increase expression, OVO not bound
● Increase expression, OVO bound
● Decrease expression, OVO not bound
● Decrease expression, OVO bound

● Increase expression, expressed in embryo
● Decrease expression, expressed in embryo
● Expressed in embryo

E

OVO ChIP | GSC ATAC | GSC H3K27ac | GSC H3K4me3 | GSC H3K27me3 | GSC H3K9me3 | 32c NC ATAC | 32c NC H3K27ac | 32c NC H3K27me3 | 8C NC H3K9me3 | OVO RNAseq

TSS -2Kb TTS

-1 0 1 2
Normalized Control Subtracted Read Count

1043

Figure 4: Genes Bound by OVO Increase in Expression in the Presence of OVO Genome-Wide. A-B) Immunofluorescent staining of adult ovarioles of the indicated genotypes (20x, scale bar = 20 μm). Ovarioles were stained for Vas (purple) to label the germline, α-Spectrin (yellow) to label dot spectrosome and fusomes, and DAPI (cyan) to label nuclei. Line indicates the dissection point for germarium through previtellogenic RNA-seq samples. C) MA plot of $ovo^{ΔBP}/ovo^{ovo-GAL4}$; *UASp-3xFHA-OVO-B* versus $ovo^{ΔBP}/ovo^{ovo-GAL4}$; *UASp-GFP* RNA-seq differential expression results. Purple dots indicate genes that significantly increased in expression and were not bound by OVO, cyan dots indicate genes that significantly increased in expression and were bound by OVO, yellow dots indicate genes that significantly decreased in gene expression and were not bound by OVO, blue dots indicate genes that significantly decreased in gene expression and were bound by OVO, and gray dots indicate genes that were not differentially expressed from our analysis. D) MA plot of $ovo^{ΔBP}/ovo^{ovo-GAL4}$; *UASp-3xFHA-OVO-B* versus $ovo^{ΔBP}/ovo^{ovo-GAL4}$; *UASp-GFP* RNA-seq differential expression results. Cyan dots indicate genes that significantly increased in expression and were found to be moderately expressed in 0-2 hour old embryos, yellow dots indicate genes that significantly decreased in gene expression and were found to be moderately expressed in 0-2 hour old embryos, purple dots indicate genes that were not differentially expressed and were found to be moderately expressed in 0-2 hour old embryos, and gray dots indicate genes that were not differentially expressed and were not found to be moderately expressed in 0-2 hour old embryos. E) Gene level read coverage heatmaps of OVO ChIP minus input, GSC and 32c ATAC-seq, GSC H3K27ac, H3K4me3, H3K27me3, H3K9me3, 8c NC H3K9me3, 32c NC H3K27ac, and H3K27me3 ChIP-seq, and $ovo^{ΔBP}/ovo^{ovo-GAL4}$; *UASp-3xFHA-OVO-B* minus $ovo^{ΔBP}/ovo^{ovo-GAL4}$; *UASp-GFP* RNA-seq for genes bound by OVO.

Figure 5: OVO ChIP-seq, ATAC/Histone ChIP-seq, RNA-seq, and DNA Binding Motifs at the *ovo* Locus. *ovo* gene level read coverage tracks for OVO ChIP minus input, GSC and 32c ATAC-seq, GSC H3K27ac, H3K4me3, H3K27me3, H3K9me3, 8c NC H3K9me3, 32c NC H3K27ac, and H3K27me3 ChIP-seq, and *ovo*^{ΔBP}/*ovo*$^{ovo-GAL4}$; *UASp-3xFHA-OVO-B* minus *ovo*^{ΔBP}/*ovo*$^{ovo-GAL4}$; *UASp-GFP* RNA-seq. Red rectangles and black rectangles represent significant OVO DNA binding motifs and OVO ChIP peaks, respectively. Gene models are represented at bottom. Small rectangles represent untranslated regions, large rectangles represent translated regions. Arrows indicate transcriptional start sites.

GO Term Analysis and Representative Gene Tracks

1073

Figure 6: OVO Binds and Significantly Increases the Expression of a Number of Genes Involved in
Essential Maternal Processes. A) Significantly enriched GO biological process terms for genes bound
by OVO and significantly increase in expression in the presence of ectopic rescue OVO. GO terms are
restricted to terms containing less than 125 associated genes. B-E) Example GO term gene level read
coverage tracks for OVO ChIP minus input and $ovo^{ΔBP}/ovo^{ovo-GAL4}$; *UASp-3xFHA-OVO-B* minus
$ovo^{ΔBP}/ovo^{ovo-GAL4}$; *UASp-GFP*. Red rectangles and black rectangles represent significant OVO DNA
binding motifs and OVO ChIP peaks, respectively. Gene models are represented at bottom. Small
rectangles represent untranslated regions, large rectangles represent translated regions. Arrows indicate
transcriptional start sites.

1083
Figure 7: Proposed Model for OVO Regulation in the Female Germline Throughout Development.
Embryonic pole cells, which are transcriptionally quiescent, contain maternally deposited nuclear OVO.
As development ensues and embryonic germ cells become transcriptionally competent, maternal OVO
binds to its target promoters but does not positively influence gene expression at this stage. Later in
embryonic development, in the presence of other activating transcription factors, maternal OVO works in
concert to positively influence gene expression, activating a number of essential germ cell-specific genes.
1090
Table S1: Flybase ART Table.
Table S2: OVO ChIP-seq Statistics by Chromosome.
Table S3: Genomic Locations of Significant Overlapping OVO ChIP Peaks
Table S4: Significant OVO DNA Binding Motifs in MEME Format
Table S5: Differential Expression Analysis Results for All Genes
Table S6: Significantly Enriched GO Biological Process Terms

A OVO Motif DNA Alignement

OVO Motif Garfinkel

OVO Motif One

p<0.05

1097
1098 **Figure S1: Significant OVO DNA Binding Motif Matches and *da-GAL4* X UAS Transgene Viability.**
1099 A) Significant alignment of the *in vivo* OVO DNA binding 'motif one' and *in vitro* OVO DNA binding motif
1100 (OVO Garfinkel).

1101
Figure S2: OVO ChIP-seq, ATAC/Histone ChIP-seq, RNA-seq, and DNA Binding Motifs at the *otu*
Locus. A) *otu* gene level read coverage tracks for OVO ChIP minus input, GSC and 32c ATAC-seq, GSC
H3K27ac, H3K4me3, H3K27me3, H3K9me3, 8c NC H3K9me3, 32c NC H3K27ac, and H3K27me3 ChIP-
seq, and *ovo^{ΔBP}/ovo^{ovo-GAL4}*; *UASp-3xFHA-OVO-B* minus *ovo^{ΔBP}/ovo^{ovo-GAL4}*; *UASp-GFP* RNA-seq. Red
rectangles and black rectangles represent significant OVO DNA binding motifs and OVO ChIP peaks,
respectively. Gene models are represented at bottom. Small rectangles represent untranslated regions,
large rectangles represent translated regions. Arrows indicate transcriptional start sites.